



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





800043368U

8.68.4.7

Prijs 7-8

10.6



E. BIBL. RADCL.

~~7.~~
~~10.2.~~
~~10.20.~~

19113

e . 300
5



PROCEEDINGS

OF THE

ANNUAL MEETING



PRODROMUS
SYSTEMATIS NATURALIS
REGNI VEGETABILIS.

PARS V.

On trouve aux mêmes adresses :

- DE CANDOLLE.** — *Prodromus systematis naturalis regni vegetabilis*; in-8. Tomi I, II, III, IV. 1824 à 1830..... 63 fr.
- Collection de Mémoires pour servir à l'Histoire du règne végétal.
- Premier Mémoire : sur la famille des *Mélastomactes*; in-4. avec 10 Planches. 1828..... 10 fr.
- Second Mémoire : sur la famille des *Crassulacées*; in-4. avec 13 Pl. 1828. 10 fr.
- Troisième et quatrième Mémoires : sur la famille des *Onagracées* et des *Paronichidées*; in-4. avec 9 Planches. 1829..... 8 fr.
- Cinquième Mémoire : sur la famille des *Ombellifères*; in-4. avec 19 Pl. 1829. 15 fr.
- Sixième Mémoire : sur la famille des *Loranthacées*; in-4. avec 12 Pl. 1830. 10 fr.
- Septième Mémoire : sur la famille des *Valérianiées*; in-4. avec 4 Pl. 1832. 5 fr.
- Huitième Mémoire : sur la famille des *Cactées*; in-4. avec 12 Pl. 1834. 10 fr.
- L'auteur n'ayant pu, ni dans le *Prodromus*, ni dans le *Cours de Botanique* qu'il publie, donner sur certains points de la science les développemens qui lui paraissent nécessaires pour fixer l'attention des botanistes, a cru devoir publier séparément une série de Mémoires servant d'éclaircissement aux deux ouvrages ci-dessus mentionnés, et dont dix formeront un volume. Cet ouvrage, accompagné de planches, et de même format que les *Mémoires du Muséum d'Histoire naturelle*, sera donc, à proprement parler, le *Commentaire du Prodromus*.
- Chaque Mémoire se vend séparément.
- *Regni vegetabilis Systema naturale; sive Ordines, Genera et Species Plantarum secundum methodi naturalis normas digestarum et descriptarum*. Volumen primum : *Sistens Prolegomena et Ordines quinque, nempe Rhamnales, Dilleniaceas, Magnoliaceas, Anonaceas et Menispermaceas*, in-8. de 368 p. 1818. 12 fr.
- Volumen secundum : *Sistens Ordines sex, nempe Barberidas, Podophyllas, Nymphaeaceas, Papaveraceas, Fumariaceas et Cruciferas*; in-8. de 750 pages. 1821..... 15 fr.
- *Astragologia, nempe Astragali, Biserrulae et Oxytropidis Historia*; cum 50 tab. aeneis; 1 vol. petit in-folio. 1802..... 54 fr.
- Le même ouvrage, grand in-folio, papier vélin superfine double..... 96 fr.
- JAUME SAINT-HILAIRE.** — *Exposition des Familles naturelles et de la Germination des Plantes*; 4 vol. gr. in-8. avec 117 planches en noir. 1805..... 36 fr.
- Le même ouvrage, 2 vol. in-4. Planches en couleur, papier ordinaire. 96 fr.
- Le même ouvrage, 2 vol. in-4. Planches en couleur, papier vélin..... 192 fr.
- HUMBOLDT (A.) et BONPLAND.** — *Plantes équinoxiales recueillies au Mexique, aux Andes de la Nouvelle-Grenade et au Pérou*; par MM. de Humboldt et Bonpland, avec leur description; 17 livraisons formant 1 vol. in-folio, avec 144 Pl. (Ouvrage complet)..... 500 fr.
- HOOKE (J.) et GREVILLE (R. K.).** — *Icones filicum : ad eas potissimum Species illustrandas delineatae, quae hactenus vel in herbariis delituerunt prorsus incognitae, vel saltem nondum per icones botanicas innotuerunt; cum tabulis aeneis*. Le texte en latin, avec 240 planches; 2 vol. in-fol. Londres, 1731. (Ouvrage complet. Les planches en noir..... 360 fr.
- Le même ouvrage, les planches en couleur..... 600 fr.
- WALLICH (N.),** surintendant du Jardin botanique de Calcutta. — *Plantae Asiae rariores*, ou Descriptions et figures d'un choix de Plantes inédites de l'Inde orientale, 3 vol. in-fol. avec 300 planches très soigneusement coloriées; le texte en langue latine. Londres, 1830 à 1832. (Ouvrage complet)..... 900 fr.

TYPIS CRAPELET,
vis à dicté de Vaugirard, 9.

PRODROMUS SYSTEMATIS NATURALIS

REGNI VEGETABILIS,

SIVE

ENUMERATIO CONTRACTA

ORDINUM, GENERUM, SPECIERUMQUE PLANTARUM

HUC USQUE COGNITARUM, JUXTA METHODI NATURALIS NORMAS
DIGESTA;

AUCTORE AUG. PYRAMO DE CANDOLLE.

PARS QUINTA:

Sistens Calycereas et Compositarum tribus priores.

PARISIIS,


Sumptibus Sociorum TREUTTEL ET WÜRTZ,
viâ dictâ de Lille, n° 17.

VENITQUE IN EORUMDEM BIBLIOPOLIO,

ARGENTORATI.

~~~~~  
M. DCCC. XXXVI.





---

# PRODROMUS

## SYSTEMATIS UNIVERSALIS

### REGNI VEGETABILIS.

---

Ordo CI. CALYCEREÆ *R. Brown, diss. publ. lect. Londini, febr. 1816. trans. soc. lin. Lond. v. xii (1817). p. 132-142. DC. theor. elem. ed. 2. p. 246. L. C. Rich. mem. mus. h. nat. 6. p. 77. Less. in linnæa 1831. p. 258. — Boopideæ Cass. diss. publ. lect. Paris. aug. 1816. journ. phys. oct. 1816. opusc. phyt. 2. p. 344.*

Flores densè capitati ut in Compositis super receptaculum (ramuli apicem) sessiles, cincti involucro polyphylo, squamis basi inter se concretis, paleis bracteolisve inter flores è receptaculo ortis. Calycis tubus ovario adnatus, limbus in lobos 5 sæpiùs dispares partitus. Corolla regularis infundibuliformis, tubo gracili longo, lobis 5 trinerviis, nervis lateralibus submarginalibus, areolis 5 infra stamina sitis et iis alternis notata, æstivatione loborum valvatâ. Stam. 5 monadelphæ et synanthereæ, antheris supernè distinctis appendice apicilari destitutis. Stylus lævis filiformis apice subclavatus indivisus. Stigma capitellatum subhispidulum. Achænium lobis calycinis sæpè induratis et spinescentibus coronatum, 1-spermum. Semen albumine carnosio donatum. Embryo axillis rectus inversus radiculâ nempè superâ, cotyledonibus 2 planiusculis. — Ordo omninò medius inter Dipsaceas et Compositas, prioribus affinis ob seminis embryonem inversum intra albumen carnosum, posterioribus ob staminâ synanthereâ, nervationem loborum corollæ ferè similem, et habitum simillimum. — Circa characteres præcipuos et locum ordinis in serie naturali, omninò consentiunt cl. Brown et Cassini, qui eodem ferè tempore ordinem egregiè admiserunt.

I. CAMOCARPHA — *Boopidis* sp. Poepp. Less.

*Involucrum* dentatum, constans squamis circ. 12 longe concretis. *Fimbriellæ* recept. acutæ in alveolas concretæ. Flores omnes fertiles. Calycis lobi 5 elliptici obtusiusculi. Cor. 5-fidæ 10-nerviæ. Anthera incluse exaltatæ ecaudatæ, filamentis apice distinctis. Stylus longè exsertus tenuis apice vix capitatus. Fruct. ign. — Herba Chilensis Valerianam Saliuncam aut globulariæfoliam habitu referens.

1. *G. POEPPIGII*  $\mathcal{L}$  in Chili summis montibus. Boopis? Alpina Poepp! pl. exs. n. 823. Less. in Linzæ 1831 p. 258. Herba digitalis glaberrima. Folia ad col- lum adgregata linearis-subspathulata integerrima scapi fere dimidiam longitudi- nem aquantia. Rhizoma cylindraceum subrepens. (v. s.)

- II. *BOOPIS* Juss. ann. mus. 2. p. 350. t. 58. f. 2. Cass. journ. phys. aug. 1818 p. 114. opusc. phyt. 2. p. 355. L. C. Rich. diss. cal. p. 51. t. 2. mem. mus. h. n. 6. p. 78. t. 11.

*Involucrum* constans squamis 7-8 ad medium concretis, dentibus sæpè interjectis. Receptaculum parvum convexum paleolis filiformibus apice subdilatis inter flores onustum. Flores omnes fertiles subconsimiles. Calycis lobi ovario breviores membranacei integri aut inciso-dentati. Cor. tubus gracilis, limbus campanulatus ultra medium 5-fidus, longè exsertus simplex. Achænium hialatum? — Herbae erectæ perennes. Folia alterna pinnatipartita. Capitula hemisphaerica terminalia.

1. *B. ANTHEMOIDES* (Juss! ann. mus. 2. p. 350. t. 58. f. 2), tota glabra, foliis pinnato-partitis, lobis capillaceis.  $\mathcal{L}$  in Bonariâ, ad Rio de la Plata et Rio Grande. L. C. Rich. l. c. cum. ic. Cass. 5. dict. suppl. 28 op. phyt. 2. p. 356. Herba 6-8 pollic. (v. s.)

2. *B. LEUCANTHEMA* (Poepp! pl. exs. n. 774), caule villis laxis longisque sublanato, foliis pinnatipartitis, lobis linearibus acutiusculis.  $\mathcal{L}$  in Chili legit cl. Poepp. Less. in Linzæ 1831 p. 259. Herba 2-3 poll. longa. Capitula dimidio minora quam in priore cui similis. Cor. albæ 4-fidæ. (v. s.)

- III. *CALYCERA* Rich. diss. cal. in mem. mus. 6. p. 77. t. 10 et segr. p. 50. t. 1. Cass. dict. 6. suppl. 36. journ. phys. aug. 1818 p. 113. op. phyt. 2. p. 353. — Calicera Cav. icon. 4. p. 34. t. 358.

*Involucrum* squamis 5-7 vix basi concretis inæqualibus constans. Receptaculum subglobosum paleis tenuissimis setaceo-subulatis inter flores onustum. Flores fertiles dissimiles et promiscui. Cal. lobi in aliis indurati valdè elongati corniformes, in alteris squamacei ovales obtusi. Corollæ tubus gracilis, limbus obconico-elongatus infra incisuras coarctatus vix semi-5-fidus. — Herbae perennes Chilenses. Folia integra aut grossè dentata. Capitula terminalia.

1. *C. CAVANILLESII* (Rich. l. c.), glaberrima, foliis dentibus grossis distantibus fere semi-pinnatifidis, paleis superne angusto-subulatis.  $\mathcal{L}$  in Chili. C. her- bacca Cav. l. c. Cass. l. c. *Involucrum* dicitur polyphyllum. Folia radicalia longissima petiolata.

2. *C. BALSAMITAFOLIA* (Rich. l. c.), caule subpiloso, foliis dentibus grossis distantibus fere pinnatifidis ad margines ciliato-puberulis, bracteolis superne linearis-lanceolatis.  $\mathcal{L}$  in Chili. Boopis balsamitæfolia Juss. ann. mus. 2. p. 350. Cass. dict. 5. suppl. 28. Scabiosa sympaganthæ R. et Pav. fl. per. 1. p. 40. — f. a. Caulis bipedalis. (v. s. ex h. Dombey.)

- IV. ACICARPHA Juss. ann. mus. 4. p. 347. Cass. dict. 1. suppl. 32. L. C. Rich. diss. cal. in mem. mus. 6. p. 77. t. 12. et segr. p. 31. L. C. — *Cryptocarpa* Brown. trans. soc. lin. 12. p. 131. non Raddi. — *Sommea Bory* ann. gen. sc. phys. (1820) 6. p. 92. t. 87. — *Echinolema* Jacq. f. — *Acanthosperma* Arrab. fl. flum. 8. t. 152.

Involucri squamæ 5 basi concretæ aut liberæ. Receptaculum oblongum. Paleolæ parvæ vix conspicuæ. Flores numerosi, centrales styli abortu steriles, laterales hermaphroditi ovarii densè concretis Cal. lobi 5, post anthesin excrecentes indurati conici spiniformes. Cor. limbo infundibuliformi. — Herbæ. Folia indivisa. Capitula oppositifolia. — Lobi calycini olim pro paleis receptaculi malè habitî, unde nomen minus aptum.

1. *A. TRIBULOIDES* (Juss. ann. mus. 2. p. 348. t. 58. f. 1), glabra erecta, foliis oblongo-cuneatis, sinnato-dentatis obtusis, inferioribus spathulatis, superioribus basi latiore amplexicaulibus, stylo longissimè exserto. ①? in Bonariâ. L. C. Rich. diss. cal. p. 18. t. 2. a. mem. mus. 6. p. 78. t. 11. *Cryptocarpa tribuloides* Cass. dict. 12. p. 85. atl. dict. sc. nat. ic. opusc. phyt. 12. Pedunculî axillares dicuntur, potius ramuli apice capituliferi. (v. s.)
2. *A. SPATHULATA* (Brown. comp. 129), glabra decumbens, foliis spathulatis basi in petiolum attenuatis mucronatis integris aut pauci-dentatis glaucescentibus, stylo brevissimè exserto. 3 in arenosis maritimis Brasiliæ, ad Rio-de-Janeiro, Sanctam-Catharinam, Bahiâ. L. C. Rich. diss. cal. p. 51. t. 3. mem. mus. 6. p. 78. t. 12. *Cryptocarpa spathulata* Cass. l. c. *Sommea Calcitrapa* Bory ann. gen. sc. ph. 6. p. 92. t. 87. *Echinolema arenarium* Jacq. f! in litt. *Acanthosperma littorale* Arrab. fl. flum. 8. t. 152. *Euphthalmum Bonariense* Pera. ench. 2. p. 474? Capitula duplò quàm in priore majora. (v. s.)

β. *glauca*, erecta glaberrima glauca, foliis obovato-cuneatis grossè dentato-incisis attenuatis in petiolum limbo duplò fere longiorem, stylo brevissimè exserto. ①? in Brasiliâ. Simillima *A. spathulata* sed erecta nec procumbens, foliorum petioli duplò triplòve longiores et involucri squamæ 5 nullo modo basi concretæ sed omninò liberæ! Radix in horto exilis perpendicularis herbam annuam indicat. An mera varietas, aut herba primo anno florens? (v. v. c. in h. Genev. et s. sp.)

3. *A. PROCUMBENS* (Less. in linnæâ 1831. p. 527), procumbens multicaulis glaberrima, foliis sessilibus glaucis membranaceis obtusis mucronulatis integerrimis, invol. squamis 6 patentibus. 7? in Brasiliâ. Calycis lobi indurato-corniculati minores quàm *A. spathulata*.
4. *A. SUPPLEVROIDES* (Less. l. c.), erecta glabra dichotoma, foliis membranaceis reticulato-venosis obtusis subdentatis, capitulis terminalibus, involucri corollæque 4-lobis. 7? in Brasiliæ uliginosis.
5. *A. LANATA* (Lag. ex Pera. ench. 2. p. 438) caule lanato, foliis glabris linearibus dente uno alterove incisis, summis integerrimis. — in Novâ Hispaniâ. Species vix palmaris, non satis nota.



**Ordo CII. COMPOSITÆ** *Vaill. act. acad. Par. 1718-1721. Linn. ord. nat. 1737. n. 21. 1764. n. 49. Berckh. exp. car. fl. comp. 1761. 4°. Adans. fam. (1763). 2. p. 103. DC. fl. fr. (1805). 4. p. 1. ann. mus. h. nat. Par. 16. p. 190. bot. gall. 1. p. 258. Brown. obs. comp. in trans. linn. soc. (1817). 12. p. 76. Kunth nov. gen. et sp. am. (1820). 4. p. 1. — Syngenesia polygamia Linn. syst. sex. 1737. cl. XIX. Meese class. syng. 1768. 8°. — Compositifloræ Gaertn. fruct. (1791). 2. p. 353. — Actinophytum, Glossariphytum, et Siphoniphytum Neck. elem. vol. 1. n. I. II. et III. — Synanthereæ L. C. Rich. in Marth. cat. h. bot. 1801. p. 85. Cass. opusc. phyt. 1826. v. 1. p. 1. dict. sc. nat. (1827). v. 51. p. 443. Bartl. ord. nat. p. 120. — Compositæ seu Synanthereæ Cass. dict. sc. nat. (1818). 10. p. 131. Less. syn. 1832. in-8°. — Flosculosæ, Semiflosculosæ et Radiatæ Tourn. inst. (1694). cl. 12. 13. et 14. — Cynarocephalæ, Cichoraceæ et Corymbiferæ *Vaill. l. c. Juss. gen. pl. (1789). p. 168.**

**DIAGN.** Flores capitati seu glomerati. Corollæ nevramphipetalæ, nempe nervis in tubo ad sinus tendentibus, in lobis (per æstivationem valvatis) nervis marginalibus donatæ. Anthere in tubum concretæ. Semen exalbuminosum. Embryo erectus.

**CAR. FRUCT.** Calyx gamosepalus ovario adhærens, tubo nunc ovario æquali (1) nunc ovarium superante (2), limbo seu pappo nunc omnino nullo aut ad marginem minimum reducto, rarissimè foliaceo, nunc sæpius scarioso, integro, dentato, lobato aut sæpius evadente in squamellas paleaceas seu setas piliformes, simplices, ramosas, denticulatas aut plumosas, uni-bi- aut multi-seriales. — Corolla summo calycis tubo inserta, gamopetala, nevramphipetala nempe petalis nervis duobus margini proximis donatis undè sæpius in tubo nervi prima fronte 5 (10 geminatim concreti) à basi ad sinus loborum tendentes et in lobis 10 nempe in quoque lobo marginales, adjectis hinc indè nervis accessoriis medium petalis occupantibus, in paucissimis solis ferè obviis. Tubus longitudine varius, à basi ad filamentorum liberationem productus. Faux sæpius ampliata à filamentorum liberatione ad separationem loborum : lobi sæpius 5 rariùs 4 aut 3 aut 2, per

(1) Tunc pappus seu limbus calycis abortivus incongruè dicitur *sessilis* seu potius *nactus crassatus*.

(2) Tunc pappus dicitur incongruè *stipulatus* seu potius *fiuctus rostratus*.

æstivationem valvati, nunc omnes æquales, nunc subæquales aut irregulares inæquales seu palmati, nunc bilabiati (aut 3 in labio exter. 2 in interiore, aut 4 in exter. 1 in inter.), nunc fissurâ ad latus interius productâ in ligulam 5-dentatam planam concreti. — Stamina 5, rarius 4, in floribus fœmineis neutrisve nulla aut rudimentaria; filamenta lobis corollæ alterna et ideò cum nervis tubi congruentia, tubo corollæ adnata, supernè libera, aut inter se omninò distincta aut in tubum concreta et ideò monadelpha, versus apicem articulata, articulo superiore antherifero connectivi locum tenente. Antheræ erectæ uniseriales in tubum stylo perforatum concretæ (*syngenesæ* seu *synantheræ*) lineares, biloculares, rimâ longitudinali intùs debiscentes, 4-valves, valvis extroflexis, apice in appendicem seu alam formâ et magnitudine in diversis variam, sæpiùs à parte polliniferâ consistentiâ distinctam, desinentes: basi sæpè in caudas duas longitudine et formâ subvarias productæ. Pollen globosum aut ellipticum echinulatum aut læve. — Ovarium 1-carpellare calyci adhærens, 1-ovulatum. Stylus teres aut rarius basi tumidus (tunc bulbosus dictus) in floribus masculis sæpiùs (ubi adest) simplicissimus, in fœmineis et hermaphroditis supernè bifidus; rami (seu vulgo stigmata) supernè plani, subtùs convexi, nunc omninò liberi nunc inter se plus minùs concreti. Glandulæ stigmaticæ (stigmata proprio sensu) biseriales, ad paginam superam ramorum styli sitæ, seriebus continuis marginalibus scabriusculis magis minùsve prominulis distinctisve. Pili pollinis collectores rigidi in florum hermaphroditorum parte styli superâ, aut iu ejus ramis ad latus inferius aut ad apicem siti, in floribus fœmineis neutrisve subnulli.

Fructus seu achænium constat calycis tubo, pericarpio, seminisque spermodermio inter se plus minùs arcetè concretis et embryonem includentibus. Achænium ideò 1-loculare 1-sperмум, supra receptaculum articulatum sæpiùs sessile rarius stipitatum, basi areolâ basilari aut laterali instructum, apice erostre aut rostratum, pappo suprâ descripto superatum, disco epigyno nectarium centrale cum ovario continuum styliforme aut alveolare cingente terminatum. Semen funiculo brevissimo inæ fructus parti adfixum erectum basi latere funiculo opposito affixum. Endoplevra seu spermodermii lamina interior (albumen Less.) crassiuscula tenera diaphana funiculo bifido percursa. Embryo erectus, rectus exorhizus,

dicotyledoneus, radiculâ brevi rectâ, plumulâ plerumque inconspicuâ, cotyledonibus planis dorso sæpè subconvexis, rarissimè curvatis, casu interdum ternis.

**INFL.** Flores collecti in *capitulum* (1) aut in *glomerulum* (2). Capitulum constat floribus pluribus super receptaculum (3) seu rami apicem nunc planum nunc conicum elongatumve, exterioribus seu inferioribus præcocioribus, intimis sensim serioribus. Flores aut omnes hermaphroditi et tunc capitula *homogama*, aut exteriores neutri seu fœminei, interiores hermaphroditi seu masculi et tunc capitula *heterogama*, aut omnes in eadem plantâ aut masculi aut fœminei et tunc capitula nunc *monoïca*, nempe floribus masculis et fœmineis, nunc *heterocephala* nempe in eadem plantâ alia mascula alia fœminea, nunc *dioïca* nempe in eadem plantâ omnia mascula seu omnia fœminea. Capitula aut corollis omnibus tubulosis donantur et tunc *discoïdea* seu *flosculosa* dicuntur, aut corollis omnibus ligulatis et tunc *ligulata* seu olim *semiflosculosa*, aut corollis in margine seu radio ligulatis in centro seu disco tubulosis et tunc *radiata* nuncupantur, aut omnibus bilabiatis et tunc *falso-discoïdea*, aut margine ligulatis centro bilabiatis et tunc *falso-radiata* seu *radiatifor-mia*. Capitula discoïdea et falso-discoïdea exhibent interdum flores marginales cæteris similes sed majores, et tunc *coronata* dicuntur.

**Involucrum** (4) constat *squamis* (5) plurimis, uni- aut pluri-serialibus, liberis aut (ubi uniseriales) margine coalitis, siccis scariosis coriaceis carnosis spinosis foliosisve, sæpè in squamam propriam et appendicem distinctis, seriebus æqualibus aut sæpius inæqualibus imbricatis aut calyculatis aut variè et subinordinatè elongatis.

**Glomerulus** constat capitulis uni- aut paucifloris involucro proprio instructis, densè et variè aggregatis, involucro gene-

(1) *Flos compositus* Veterum. *Calathis* Mirb. *Anthodium* Ehr. *Cephalanthium* Rich.

(2) *Capitulum* Cass.

(3) *Phoranthium* Rich. *Clinanthium* Cass. *Rachis* Less.

(4) *Calyx communis* Veterum. *Perigynanda communis* Neck. *Periclinium* Mirb. *Periphoranthium* Rich.

(5) Plurimi squamas vocant foliola, sed squamarum terminus aptior mihi videtur 1°. Quia vox foliolium sensu diversissimo jam gaudet in foliis compositis. 2°. Quia squamæ involucris omnino repræsentant squamas gemmarum et præcipuè gemmarum floriferarum 3°. Quia jam in pluribus aliis ordinibus partes involucris sunt dictæ

rali cinctis, super receptaculum generale insidentibus, sæpius centralibus præcocioribus, exterioribus serioribus aut inordinatè florentibus.

*Receptaculum* capituli nunc *paleaceum*, nempè *paleis* (1) involucris squamis similibus ad latus cujusque floris externum sitis, super basin articulatis et in folia vera hinc indè monstròse expansis donatum, nunc *semipaleaceum* nempè paleis in parte exteriore tantum donatum, nunc *epaleaceum* seu *nudum* nempè paleis omninò orbatum. In aliis capitulis (seu fortè glomerulis arctissimè cinctis) receptaculum dicitur nunc *fimbrilliferum*, nempè ubi flos singulus ex omni parte cinctus est alveolâ basi squamosâ apice sæpè in laminas setasve irregulares desinente: nunc *alveolatum* nempè ubi alveolæ breves in setas non productæ sed sæpè denticulatæ: nunc tandem *areolatum* ubi nempè areola pentagona (alveolæ rudimentum) singuli floris basin circumdat. Fortè receptacula plùs minùs paleacea flores verè spicatos et capitula proprio sensu designant, et receptacula fimbrillifera aut alveolata indicant capitula arctissimè cincta proprio sensu flores umbellatos et quorum alveolæ involucelli rudimenta repræsentant. Clariùs dicam capitula aut more spicæ contractæ aut more umbellæ contractæ disposita.

Væc. Herbæ sæpiùs perennes, aut frutices aut rariùs (præsertim in insulis) arbores, per totum orbem terrarum obviæ et decimam ferè partem regni vegetabilis comprehendentes. Folia alterna aut opposita formâ incisuris aut aspectu diversissima, semper simplicia nec composita. Caulis (ubi simplex) aut rami capitulo glomerulove terminati. Rami sæpiùs corymbosi, centrales præcociores. Folia sub capitulis glomerulisve sita à cæteris sæpè diversa et floralia seu bracteantia dicta, nunc gradatim in squamas mutata nunc abruptè squamis multò majora. Corollæ nunc xanthicæ nunc cyanicæ (2) in eodem capitulo aut *homochromæ* seu concolores aut *heterochromæ* seu discolores; ubi discolores ligulæ semper cyanicæ et tubuli xanthici! imò tubuli xanthici cum ligulati monstròse evadunt, cyanici fiunt! Succus proprius varius, interdum lacteus. Sapor sæpiùs amarus, adstringens et aromaticus.

---

(1) Aliis *Bracteola* dictis. Vox paleæ mihi servanda videtur quia notissima et quia squamæ extimæ receptaculi seu accessoris à plurimis et quidem commodè bracteolæ dicuntur.

(2) Vid. Physiol. veget. 2. p. 907.



**Obs.** Ordo omnium naturalissimus et vastissimus et ideo difficillimus, à Tournefortio facile sed artificialiter, à Vaillantio naturalius sed difficilius, à Linnæo eleganter sed arbitrariò, ab Adansonio nimis vagè, à Gærtnero nimis carpologicè, à Cassinio strenuè et philosophicè sed sæpè obscurè, à Lessingio lucidè et simpliciter dispositus, summam adhuc botanicorum attentionem requirit. An capitula alveolis (nec paleis) donata, ritè (Albertiniâ præsertim suadente) cum glomerulis veris consocianda? An pappus qui evidenter in plerisque generibus à limbo calycis variè abortivo provenit, fortè in quibusdam (v. gr. Dipterocome) etiam ex pericarpio proprio elongato provenire posset? An pars subcarnosa embryonem circumdans albuminis veri aut endoplevræ locum tenet? An fila è basi styli orta et per partem istam carnosam descendentia basin embryonis attingunt et fœcundationis principium conducunt? Quare in capitulis heterogamis flores masculi si adsint semper centrales et fœminei semper marginales? ~~Cap~~ corollæ tubulosæ xanthicæ in seriem cyanicam transeunt ~~cum~~ ligulatæ aut ampliatae evadunt? An achæniî rostrum ubi adest, à calycis tubo tantum aut etiam è pericarpio vero producto formatur?

## CONSPECTUS ORDINIS.

\* *Tubulifloræ*. Nempè floribus hermaphroditis tubulosis regulariter 5-(rarius 4-) dentatis.

**Trib. I. VERNONIACEÆ.** Stylus fl. herm. cylindræus, ramis sæpius subulatis elongatis, rarius obtusis brevibus, semper æqualiter et longiusculè hispidis. Stigmati series antè medios styli ramos desinentes.

**Trib. II. EUPATORIACEÆ.** Stylus fl. herm. cylindræus, ramis longis subclavellatis extus supernè puberulo-papillosis. Stigmati series parum prominulæ antè medios styli ramos sæpius desinentes.

**Trib. III. ASTEROIDEÆ.** Stylus fl. herm. cylindræus, ramis linearibus extus subplanis et supernè æqualiter et minutè puberulis. Stigmati series prominulæ usque ad originem pubis exterioris circiter extensæ.

**Trib. IV. SERNECIONIDEÆ.** Stylus fl. herm. cylindræus, ramis linearibus apice penicillatis nunc truncatis nunc ultra penicillum, in conum brevem aut appendiculum elongatum angustum et hispidulum productis. Stigmati series latiusculæ et prominulæ usque ad penicillum extensæ.

Trib. V. CYNARÆ. Stylus fl. herm. supernè nodoso-incrassatus et sæpè ad nodum penicillatus, ramis nunc concretis nunc liberis extùs puberulis. Stigmatis series haud prominulæ apicem ramorum attingentes et ibi confluentes.

\*\* *Labiatifloræ*. Nempè floribus hermaphroditis sæpissimè bilabiatis.

Trib. VI. MUTISACEÆ. Stylus fl. herm. supernè cylindraceus aut subnodosus, ramis sæpiùs obtusis truncatisve extùs valdè convexis et in parte superiore pube minutâ æquali (rariùs nullâ) vestiti.

Trib. VII. NASSAUVIACEÆ. Stylus fl. herm. nunquam nodoso-incrassatus, ramis linearibus longiusculis truncatis apiceque solo penicillatis.

\*\*\* *Ligulifloræ*. Nempè floribus omnibus hermaphroditis ligulatis.

Trib. VIII. CICHORACEÆ. Stylus supernè cylindraceus, ramis longiusculis obtusiusculis æqualiter pubescenti-hirtellis. Stigmatis series antè medios styli ramos desinentes.

*N. B.* Confer ad frontem tribùs cujusque subtribuum divisionumque conspectum methodicum et ad calcem totius ordinis 1°. indicem generum dichotomum, et ad tyronum faciliorem usum, artificialem : 2°. indicem nominum alphabeticum.

Trib. I. VERNONIACEÆ *Less. in linnæd* 1829. p. 241. 1830. p. 134. 1831. p. 625. *syn.* p. 145. — Vernoniæ (excl. sect. 2 et 3) et Tagetinæ Pectideæ *Cass. tabl.* II. XX *opusc.* 1. p. 333. 2. p. 206. *dict.* 20. p. 384. 36. p. 19. 57. p. 359.

Stylus supernè cylindraceus, ramis æqualiter hispidis in plurimis longis subulatis in paucissimis brevibus obtusisve; stigmatis glandulæ dispositæ in series prominulas angustas antè medios styli ramos desinentes. Corolla 5-fida sæpiùs regularis interdum obscurè bilabiata aut palmata. Pollen globosum echinulatum. — Capitula uni-pauci- aut multiflora, sæpiùs homogama discoidea, floribus omnibus hermaphroditis aut in unico genere dioicis, rariùs heterogama radiatâ floribus radii ligulatis fœmineis. Recept. alveolatum fimbrilliferumve aut nudum, in paucissimis (dubiis?) paleaceum. Flores cyanici (purpurascents, cœrulescentes aut albi) nec ferè unquam xanthici nisi in Pectidearum subtribu. — Tribus, jam rectè suadentibus cl. Cassini et Lessing, media inter Cichoraceas et Eupatorineas, sed etiam mediantibus Pectideis ad Senecioneas-Tagetineas affinis.

## CONSPECTUS TRIBUS.

Subtrib. I. VERNONIÆ. Capitula discoidea homogama. (Gen. I-XLVIII.)

Div. I. EUVERNONIÆ. Antheræ cœaudatæ. Involucrum non compressum polyphyllum. (Gen. I-XXXV.)

Subdiv. 1. *ETHULIÆ*. Pappo nullo aut simplici coroniformi. (Gen. I-V.)

Subdiv. 2. *HETEROCOMEÆ*. Pappo uni-bi- aut pluriseriali, omnibus aut saltem interiore setiformi aut angustissimè ligulato. Capitulis non fasciculatis. (Gen. VI-XXVI.)

Subdiv. 3. *ALBERTINIÆ*. Pappo prioris subdiv. Capitulis sæpius oligocephalis in glomerulum fasciculatis. (Gen. XXVII-XXXV.)

Div. II. *ELEPHANTOPÆ*. Antheræ ecaudatæ. Involucrum compressum, squamis alternis conduplicatis. (Gen. XXXVI-XXXVIII.)

Div. III. *ROLANDRÆ*. Capitula 1-flora. Invol. oligophyllum (1-5), squamis sæpius concretis. Antheræ ecaudatæ. (Gen. XXXIX-XLIIV.)

Div. IV. *BOJERIÆ*. Antheræ bicaudatæ. (Gen. XLV-XLVIII.)

Subtrib. II. *PECTIDEÆ*. Capitula radiata heterogama. (Gen. XLIX-LVIII.)

Div. 1. *LIABÆ*. Cor. disci regulariter 5-fidæ. Styli rami elongati. Folia eglandulosa eciliata. (Gen. XLIX-LIV.)

Div. II. *EUPECTIDEÆ*. Cor. disci sæpè irregulares. Styli rami abbreviati. Folia glandulosa basi sæpius ciliata. (Gen. LV-LVIII.)

Subtrib. I. *VERNONIÆ*. *Cass. trib. XX. gen. paucis excl. aut add.*

Capitula discoidea nec radiata. Styli rami elongati acuminati. — Folia alterna rariùs opposita. Flores albi, cœrulei, aut purpurei.

Div. 1. *EUVERNONIÆ*. — *Vernoniæ* prototypæ *Cass. tabl. xx. sect. iv.* — *Vernoniæ* pleræque *Less. in linnæd 1830. p. 136. syn. 146. excl. Liabeis et Pectideis.*

Capitula discoidea multiflora (in paucissimis pauciflora imò 1-flora), Antheræ ecaudatæ. Involucrum teres multiseriale imbricatum. Recept. nudum aut alveolatum imò fimbriiferum nec forsan unquam vere paleaceum (excl. fortè *Heterocoma*?) Cor. tubulosa regularis aut subpalmata. Styli rami semisubulati. — Folia alterna (in paucissimis opposita).

Subdiv. 1. *ETHULIÆ*.

Pappo nullo aut coroniformi uniseriali. Capitulis uni- aut multifloris, distinctis nec in glomerulum concretis.

I. *ADENOCYCLUS* *Less. in linnæd 1829. p. 337. f. 39-41. syn. p. 151.*

Capitulum 1-florum. Involucrum oblongo-cylindraceum imbricatum, squamis coriaceis siccis 1-nerviis interioribus multò longioribus. Recept. punctiforme. Cor. regularis, limbo à tubo haud distincto profundè 5-fido, lobis acuminatis parte integrâ glabrâ brevioribus. Anth. incluse. Filam. glabra lœvia. Achænium obconicum breve sulcatum glaberrimum, nectario alveolari, disco epigyno orbe glandularum non.

Pappus 0. — Frutex *Odontoloma* habitu. Rami juniores petiolique anescentes. Folia alterna petiolata ovato-oblonga acuminata sub-

## COMPOSITÆ-VERNONIACEÆ.

11

integerrima glabra. Corymbi terminales pluriès dichotomi condensati. Cor. rosæ.

1. *A. CONDENSATUS* (Less. l. c.). 5 in insulâ Trinitatis. Eupatorium condensatum Reich. in Sieb! fl. exs. trin. n° 73. (v. s.)

II. *ODONTOLOMA* H. B. et Kunth. nov. gen. 4. p. 43. t. 319. Juss. dict. 35. p. 395. Cass. tabl. xx. n. 46. Less. in linnæâ 1829. p. 337. syn. p. 147. non Pers.

Capitulum 1-florum. Invol. cylindraceum imbricatum, squamis siccis 1-nerviis interioribus longioribus. Recept. punctiforme. Cor. regularis glabra 5-fida, lobis acuminatis parte integra brevioribus. Fil. lævia. Styli rami semi-subulati. Achæmium breve glaberrimum, nectario alveolari. Pappus 1-serialis coroniformis coriaceus brevis apice inæqualiter et acutè multifido-dentatus. — Arbuscula, ramis fuscescenti-tomentosis, foliis sparsis petiolatis ovatis integerrimis acuminatis suprà glabris, subtus pulverulento-tomentosis. Corymbi terminales subsessiles ramosissimi. Capitula fasciculato-congesta. Cor. albidæ?

1. *O. ACUMINATUM* (H. B. et Kunth. l. c.). 5 in convalli Caracasana. Eupatorium cornifolium Willd. herb. ex Less.

III. *OIOSPERMUM* Less. in linnæâ 1829. p. 339. f. 69. 71. 72. 78. syn. p. 148. — Ethuliæ sp. Nees. Calydermi sp. Spreng.

Capitulum multiflorum æqualiflorum. Involucrum imbricatum patens teres, squamis scariosis glabris acutis, basi foliis inæqualibus capitulo longioribus cinctum. Recept. latum nudum. Cor. regularis, limbo 5-fido à tubo vix distincto, lobis acuminatis parte integrâ brevioribus. Anth. inclusæ. Filam. lævia. Achæmium apice rotundatum glandulosum multicosatum, disco epigyno parvo. Pappus o. — Herba ramosa glabriuscula ascendens basi repens. Folia alterna petiolata ovato-lanceolata utrinquè attenuata serrata. Capitula solitaria longè pedunculata oppositifolia. Cor. pallida.

1. *O. INVOLUCRATUM* (Less. l. c.), caule glabrescens, foliis petiolatis ovatis membranaceis glabris, corollâ hispidiusculâ, achænio turbinato-ovato pubescenti-hispidulo longiore. (1) in Brasiliæ arenosis circa San Pedro de Alcantara. Ethulia involucrata Nees et Mart. nov. act. nat. cur. 12. p. 3. Amphirephis pilocarpa Nees et Mart. ibid. p. 4. Calydermos repens Spreng. nov. entd. 1. p. 278. obs. ex ipso (syst. 3. p. 458) manca. (v. s. comm. à Mus. reg. Berol.)

2. *O. WIGHTIANUM*, caule villosissimo, foliis sessilibus ovatis acuminatis basi attenuatis utrinquè (præcipuè in nervis) villosissimis, corollâ glabrâ, achænio gracillimo brevioris. — In Indiâ orientali legit cl. Wight. Amphirephis Wightiana Wall! cat. comp. n. 68. Stylus ex antherarum tubo non exsertus simplex glaber fortè abortivus. Flores in capitulo circiter 100 gracillimi. An genus proprium? (v. s. com. ab ampl. cœtu mero. angl. Ind. or.)

IV. *SPARGANOPHORUS* Vaillant. act. soc. par. 1719. p. 368. Gærtn. fr. 2. p. 396. t. 165. Cass. dict. 50. p. 71. Less. in linn. 1829. p. 335. syn. p. 147. non Lag. nec Michx. — Strachium P. Brown. jam. 312. t. 34. f. 2. Juss. gen. 184. — Athenæa Adans. fam. 2. p. 121. non Schreb. — Ethuliæ sp. Lin.

Capitulum multiflorum æqualiflorum. Involucrum teres imbricatum squarrosum, squamis siccis. Recept. nudum. Cor. regularis 3-fida rarisimè 4-fida, lobis acuminatis, tubo et fauce subcontinuis brevioribus. Antheræ inclusæ. Filam. lævia. Styli rami semi-subulati. Achæmium obliquè obpyramidatum 3-5-costatum glabrum glandulosum, collo apicali magno. Pappus 1-serialis brevis coroniformis crenulatus cartilagi-

neus. — Herbarum annuarum glabrae ramosae. — Folia alterna elliptica utrinque acuminata breviter petiolata serrata. Capitula ad axillas sessilia.

1. *S. VAILLANTII* (Gært. l. c.), foliis distanter serratis. (1) in subhumidia, fossis et ripis graminosis, Jamaicae, Sancti-Thomasii, Porto-Ricci, Guadalupae, Guayaquil, Brasiliae; nec ut videtur in India orient. ut dixit Vaill. l. c. *Strachium* Brown. l. c. Sparg. *Vaillantii* et Sparg. *Strachium* Pers. ench. 2. p. 398. Stend. nomen. phan. n. 801. excl. syn. Adans. et R. Br. *Ethulia sparganophora* Lin. sp. 1171. Arrab. fl. flum. 8. t. 90. *Ethulia Strachium* Swartz. ! flor. ind. occ. 1297. *Strachium Americanum* Poir. dict. 7. p. 475 (v. a.)
2. *S. AFRICANUS* (Stend. nom. phan. 801) foliis obtusè repandis dentatisve. (1) in Africa Ovariensi et Senegalensi. *Strachium Africanum* Beauv ! fl. Ow. et Ben. 1. p. 80. t. 48. Herba priori simillima nec forsitan specificè distinguenda. Capitula in utraque variant solitaria gemina aut in axillis 3-4. (v. a.)

† *Species exclusæ.*

- S. AGERATOIDES* Lag. = *Ageratum coelestinum*.  
*S. OBTUSIFOLIUS* Lag. = *Ageratum conyzoides*.  
*S. VERTICILLATUS* Mich. = *Sclerolepis verticillatus*.  
*S. SONCHIFOLIUS* Spr. = *Cephalophorus sonchifolius*.

V. *ETHULIA* Cass. dict. 15. p. 7. et 437. Less. in linn. 1829. p. 336. f. 26. 27. 35. et 43. syn. p. 148. non Gært. — *Kahiria* Forsk. pl. æg. 153. — *Ethulia* spec. Lin. — *Pirarda* Adans. fam. 2. p. 499 non Cass. — *Leighia* Scop. non Cass.

Capitulum multiflorum æqualiflorum. Invol. teres, pluriseriale, squamis subfoliaceis acutis. Recept. nudum. Cor. regularis, limbo campanulato 5-fido, laciniis acuminatis parte integrâ brevioribus. Anth. inclusæ. Styli rami semisubulati. Achænium obpyramidatum 4-quetrum 4-costatum glabrum glandulosum apice truncatum 4-5-gonum. Pappus minimus coroniformis carnosus integerrimus. — Herbarum ramosæ erectæ. Folia alterna oblonga utrinque attenuata serrata aut integra pellucido punctata. Capitula Pisi magnitudine in corymbum disposita. Cor. roseæ aut purpureæ.

\* *Genninæ, foliis non decurrentibus.*

1. *E. CONYZOIDES* (Linn. sp. 1171 Lin. f. decad. 1. t. 1), ramis corymbosis subdivergentibus. Foliis elliptico-lanceolatis sæpius grossè serratis. (1) Cor. purpureæ. Variat foliis grossè serratis rarissimè subintegerrimis plus minus distinctè pellucido-glandulosis.
  - a. *Indica*, caulibus folisque glabriusculis. — in India orientali ad Boirugaya, Barsi, etc. (Roxb ! Wall ! ) Eth. ramosa Roxb ! herb. (v. a. sp. et v. c.)
  - β. *Africana*, caulibus folisque adpressè villosis. — In Ægypto ad ripas Nili (Forsk ! Delil ! ) in Senegaliâ (Rous ! ) in paludosis Albredæ ad ripas flum. Gambie (Perrot ! ). *Kahiria* Forsk. pl. æg. descr. 153. *Eupatorium areolatum* Perrottet ! in litt. Herba major et magis villosa quàm var. α : fortè specificè distinguenda ? Specimina glabriora et subintegrifolia. In Africa australi et orient. ad Omsamwubo et Omsanculo legit cl. Drege. n. 5065 ! (v. s. sp. )
2. *E. GRACILIS* (Delil. in Caill. voy. t. 64 et in descr. pl. Caill. p. 44. t. 3. f. 5), ramis strictis paniculatis, foliis lanceolatis integerrimis aut parcè serratis. (1) In Sennar ad El-Zerebyn (Cailland et Acerbi ! ) Cor. roseæ. Herba piloso-asperula. Folia distinctè pellucido-punctata. (v. a. comm. à cl. Acerbi. )
3. *E. ANGUSTIFOLIA* (Bojer ! in litt. 1835) ramis corymbosis apice adpressè pubescenti-villosis, foliis elongato-linearilanceolatis subintegerrimis, invol. squa-

mis obtusiusculis. ① in ins. Madagascar ad agros cultos legit cl. Bojer. Cor. purpureæ. Duabus prioribus similis, sed capitula, etiam numero florum simili 20-25, composita, duplò minora et invol. squamæ subobtusæ nec acuminatæ. (v. s. comm. à cl. inv.)

\* \* Spuriæ, foliis decurrentibus.

4. F? GARIKINA, suffruticulosa ramosissima glabra, foliis oblongo-linearibus integris aut remotè serratis in alas angustas integerrimas decurrentibus, capitulis ad apices ramorum glomerato-capitatis. 5 in Africâ australi ad flumen Gariep legit cl. Drege n. 2721! Habitus E. gracilis, sed caulis suffrutescens. Invol. squamæ acutæ. Capitula fortè dioica. Styli fl. masc? simplices, fl. form. bifidi, ramis hispidulis. Achænia minima ex statu immaturo calva. An genericè separanda? Ex capitulis junioribus non asserere queo. (v. s.)
5. E? BRASILIENSIS (Link. enum. hort. Berol. a. p. 310) caule alato, foliis lanceolatis acutis serratis pubescentibus decurrentibus, floribus subcorymbosis. 7 in Brasiliâ!†.

†† Species exclusæ.

- E. AGERATOIDES Spreng. = *Alomia ageratoides*.
- E. BIDENTIS Linn. = *Flaveria contrayerba*.
- E. AURICULATA Thunb. = *Dichrocephala Capensis*.
- E. DIVARICATA Linn. = *Epaltes divaricata*.
- E. INVOLUCRATA Nees et Mart. = *Oiospermum involucreatum*.
- E. INTEGRIFOLIA Don. = *Cotula*.
- E. PANICULATA Houtt. = *Hippia integrifolia*.
- E. REPANDA Nec. = *Cephalophora glanca*.
- E. SPARGANOPHORA Linn. = *Sparganophorus Vaillantii*.
- E. STRUCHIUM Swartz. = *Sparganophorus Vaillantii*.
- E. TOMENTOSA Linn. = *Artemisia lavandulæfolia*.
- E. UNIFLORA Michx. = *Sclerolepis verticillatus*.

#### VI. HERDERIA Cass. dict. 6o. p. 586 et 599.

Capitulum multiflorum discoideum homogamum. Recept. planum nudum. Invol. campanulatum biseriale, squamis oblongis aequalibus foliaceis. Cor. 5-fida, lobis extus glandulosis. Stigmata ut in *Ethuliâ*. Achænia tri-rariùs-tetraquetra prismatica in invol. densè conferta, glabra, angulis nerviformibus, annulo cupulari apice coronata. Pappus biformis uniserialis, paleis 8-10, crassis oblongis obtusis, imò interdum emarginatis ciliatis, demum stellato-patentibus constans, et setis 2-5 elongatis rigidis scabris ex angulis achæniæ ortis constans. — *Herderia* Senegalensis ripariæ, ramosæ sæpè ramosissimæ. Folia alterna obovata obtusa apice subdenticulata. Capitula terminalia solitaria. Flores purpureo-violacei, Genus affine *Ethuliæ* sed pappo faciliè diversum.

1. H. TRUNCATA (Cass. l. c.). ①? 7? secus flumen Senegal legit cl. Perrottet et quatuor varietates aut species observavit.
  - a. *leptorhiza*, radice exili simplici caule erectiusculo foliis obovato oblongis, nascentibus ramulisque subvillosis adultis glabris. ①? in demissis et argillosis secus flumen Senegal in regione Walo post fluv. inundationem mense decembris. Fortè sequentis ætas junior (v. s.)
  - β. *ascendens*, radice obliquâ fibrillosâ caule ascendente ramoso foliis obovato oblongis, nascentibus ramulisque subvillosis, adultis glabris. 7? in argillosis cum priore. (v. s.)
  - γ. *multicaulis*, radice crassâ subliguosâ tereti verticali, caulibus è collo

- innumeris diffusis ramosis obovatis, foliis ramulique junioribus subvillosis, aduktis subglabris 2? in inundatis argillosis cum prioribus. (v. a.)  
 1. *villosa*, tota etiam adulta villosissima ferè lanata. 2? cum priori (v. s. omnes comm. à cl. Perrottet.)

## Subdiv. 2. HETEROCOMEÆ.

Pappo uni- bi- aut rariùs pluri-seriali, interiore semper setiformi, exter. setiformi squamellato aut rarissimè coroniformi; capitulis uni- bi- aut multiflosculis distinctis nec in glomerulum concretis.

VII. PACOURINA *Aubl. guian.* 2. p. 800. t. 316. *Kunth nov. gen.* 4. p. 23. — *Pacourinopsis* *Cass. bull. philom.* 1817 septemb. p. 151. *dict. sc. nat.* 37. p. 212. *Less. in linnæa* 1829. p. 318. — *Pacourina* sect. *Vernonia* *Less. in linnæa* 1831. p. 680. *Meisteria* *Scop. introd. non Gmel. nec Ræusch.* — *Haynea Willd. spec.* 3. p. 1787. non *Schum.*

Capitulum globosum multi- et æqualiflorum. Involucrum floribus subæquale imbricatum, squamis latis ovali-oblongis parallelinerviis margine membranaceis, exterioribus foliaceis spinâ terminatis, inter. siccis obtusis. Recept. planum nudum carnosum. Cor. regularis, tubo brevi, fauce cylindricâ, 5-lobis subacuminatis parte integrâ brevioribus. Anth. inclusæ. Styli rami glabriusculi. Achænia cylindracea longè glabra, cortice crasso enervio. Pappus brevis pluriserialis, pilis rigidis crassiusculis denticulatis, interioribus basi subconcretis. — Herbæ glabræ. Caulis teres. Folia alterna ovalia aut oblonga in petiolum marginatum subamplexicaulem attenuata. Capitula oppositifolia aut lateralialia? sessilia solitaria. Cor. purpureæ. — Genus à Vernoniâ diversissimum habitu toto, achæniî formâ et pappo.

1. *P. edulis* (*Aubl. l. c.*), foliis ovali-lanceolatis dentibus setiformibus distantibus hinc indè serratis, capitulis oppositifoliis, involucri squamis exterioribus obtusis spinulâ mucronatis. — In Cayennâ. Folia utrinquè glandulis innumeris opacis notata. *Pacourinopsis integrifolia* *Cass! Dict.* 37. p. 212. *Less. in linnæa* 1829. p. 318. *Vernonia* n° 142. *Less. in linnæa* 1831. p. 681. *Haynea edulis* *Willd. l. c.* (v. s. comm. à mus. Par. et sem. mat. in herb. Desf.)
2. *P. cirsioifolia* (*H. B. et Kunth nov. gen. am.* 4. p. 30.) foliis oblongis acuminatis grossè dentato-spinosis, capitulis lateralibus? involucri squamis exterioribus spinâ terminatis 2? in humidis prope Gnayaquil Quitenarum. *Pacourinopsis dentata* *Cass. l. c.* *Vernonia* n° 143. *Less. in linnæa* 1831. p. 681.

† *Species exclusa.*

~~*Haynea pedunculata* Spr. = *Vernonia pedunculata*.~~

VIII. HETEROCOMA *DC. ann. mus.* 16. p. 191. t. 7. *choix mém.* 2. p. 34. t. 3. *Less. in linnæa* 1829. p. 339 *syn.* p. 150. *Serratulæ spec. Pers.*

Capitulum multiflorum æqualiflorum. Involucrum campanulatum subteres imbricatum, squamis coriaceis siccis plurinerviis, interioribus sublongioribus subpungentibus. Recept. paleis involucri squamas referentibus caducis onustum, et alveolis 1-floris ad palearum axillis notatum! Cor. regularis, tubo tereti, laciniis 5 linearibus tubo longioribus. Anth. subexsertæ. Styli rami semi-subulati. Achænium oblongo-cylindraceum glabrum, disco epigyno magno, nectario alveolari. Pappus biserialis, serie exteriori brevi coronæformi cartilagineâ crenatâ, interiori setis filiformibus apice serratis longis rigidis caducis. — Suffrutescens Brasiliani. Folia alterna. Capitula ad axillas sessilia. Fortè capitulum

potius dicendum esset glomerulus capitulis 1-floris ad bractearum axillas ortis constans: paleæ recept. potius bracteæ: alveolæ involucri proprii rudimentum?

1. H. ALBIDA (DC. l. c. descr. et mon.), foliis semi-amplexicaulibus oblongo-lanceolatis utrinque tomento adpresso brevissimo albidis, corollis glabris. 5 in Brasiliâ Less. in linn. 1829. p. 340. f. 50. 51. 52. Cnicus foliis semi-amplexicaulibus, etc. Vell. in Roëm. script. lus. et br. 141. Serratula albida DC. in Pers. ench. 2. p. 391. (v. s.)

† *Species exclusæ.*

H. BIFRONS DC. = Chronopappus bifrons.

H. PEDUNCULATA Steud. = Hololepis pedunculata.

- IX. VERNONIA Schreb. gen. 2. p. 541. Swartz fl. Ind. occ. 3 p. 1317. t. 16. Lessing in linnæa 1829. p. 244 syn. p. 146. (excl. spec. plur.) Vernonia, Ascaricida, Gymnanthemum, Oliganthes, Lepidaploa et verosimiliter Achyrocoma et Centrapalus Cass. dict. — Vernonia et Pollalesta H. B. et Kunth.

Capitulum pluri- seu multiflorum æqualiflorum. Involucrum imbricatum floribus brevius, squamis interioribus longioribus. Recept. nudum aut rarius alveolato-subfimbriolosum. Cor. regularis 5-fida, lobis parte integrâ subæqualibus. Filam. lævia. Achænium callo basilari cartilagineo, disco epigyno magno. Pappus sæpius biserialis, serie interiore setiformi externam sæpè paleaceam longè superante, rarius biserialis æqualis. — Herbæ erectæ aut frutescens imò ferè arbores habitu vario. Folia alterna (in paucissimis opposita) sæpè glandulosa. Inflorescentia varia sæpè scorpioidea. Flores in capitulis numero varii, in unicâ solitarii, in plurimis 3-4, in plerisque ∞. Cor. purpureæ, roseæ aut albæ, nunquam lævis, ferè in omnibus glabræ. Genus vastissimum aut fortè dividendum, aut plerisque sequentibus augendum! Habitum præcipuè secutus sum in distributione genericâ admissâ forsan ulteriori examini subiciendâ.

Sect. 1. ACILEPIS Don prod. fl. nep. 169.

Capitula multiflora sessilia. Invol. squamæ multiseriales imbricatæ subpungentes. Recept. alveolatum. Achænia angulato-villosa. Pappus biserialis, serie externâ brevissimâ. Herbæ Indicæ.

1. V. BIGIOPHYLLA, caule tereti pubero, foliis obovato-oblongis basi attenuatis apice acutis serrulatis utrinque glandulosis subtus reticulatis secus nervos raso et brevè velutinis, corymbi capitulis 1-3 sessilibus, involucri squamis sublineari-ribus valdè acuminatis glabris, exterioribus confertis brevissimis, achænio densè villosa. 2l in Indiâ orient. Nepauliâ ad Bechiaco et Hethaura legit Jcl. Wallich. 1. rigida Wall! cat. et herb. n. 2925. comp. 35. non Swartz. Folia ferè Decasemini asperi sed invol. et fruct. diversus. Pappi series externa interiori quintuplò brevior. (v. s. comm. angl. hon. coct. merc. angl. Ind. orient.)
2. V. THUNBERGII (Wall! cat. et herb. n. 2926. comp. 36), caule tereti simplici glandulosis pilis scabrido, foliis rigidis sessilibus obovatis oblongisve brevè acuminatis grosse serratis utrinque glandulosis subtus reticulatis secus nervos scabridis, capitulis axillaribus sessilibus solitariis ultimis subconfertis, involucri squamis oblongo-linearibus acutis subpungentibus multiserialibus, externis brevissimis, achænio pubescente. 2l in Indiâ orient. Acilepis squarrosa Don prod. 169. Vern. squarrosa Less. in linnæa 1831. p. 627. non p. 678. Pappi series int. interiori 4<sup>ta</sup> brevior paucisetosa.
3. angustior, foliis oblongis subovatis. Eupatorium pauciflorum Wall! ad Taogdong et Promé. (v. s.)



β. *latior*, foliis latius obovatis. Onopordon teres Ham. ad Sakanaghur.  
(v. s. comm. ab hou. cœt. merc. angl. Ind. or.)

Sect. II. HOLOLEPIS. — *Hololepis* DC. *ann. mus.* 16 p. 190. *rec. mem.* p. 34. *Cass. dict.* 21. p. 308. Hayneæ sp. *Spreng.* — *Vernoniæ* sp. *Less.*

Capitulum multiflorum. Involucri squamæ siccæ coriaceæ adpressæ imbricatæ; exteriores breves ovatae, interiores seorsim elongatae. Recept. latum planum alveolatum fimbriiferum. Achænium glaberrimum. Pappus paleis piliformibus rigidis æqualibus barbellulatis numerosis. — Herbae (forsan basi anfruticosae) Americanae erectae crassae ramosae. Folia alterna petiolata elliptica penninervia integerrima supra glaberrima, subtus sublepidoto-canescens. Rami axillares foliis longiores apice 1-cephali erecti.

§. 1. *Pedunculis 1-cephalis nudis aut immediate sub involucro folia floralia 4 bracteantia gerentibus.*

3. V. PEDUNCULATA (DC. *ann. mus.* 16. p. 190. t. 6. *rec. mem.* p. 35. t. 2. sub *Hololepis*), involucri squamis glaberrimis, corollæ lobis glabris, receptaculi fimbriis hinc inde elongatis et achænium superantibus 2? in Brasiliâ. *Serrata pedunculata* DC. in *Pers. ench.* 2. p. 391. Hayneæ *pedunculata* *Spreng.* *syst.* 3. p. 391. H. *pedunculata* *Cass.* l. c. Pappus rufescens. (v. s.)

4. V. INVOLUCRATA (Less. in *linnæa* 1829. p. 246. et 1831. p. 627 excl. syn.), involucri squamis extus margine ciliatis, corollæ lobis apice barbatis, receptaculi fimbriis brevibus 2? in Brasiliâ tropicâ. Pappus flavescens. Invol. squamm 3-nerviæ dicuntur nec ut in priore enerviæ aut multinerviæ dicendæ.

§. 2. *Capitulis ad apicem plurimis aggregatis inter folia sessilibus, invol. squamis in acumen subspinosum aut molle desinentibus.*

5. V? PROTEOPSIS, tota capitulis exceptis adpressæ sericeo-argentea, caule herbaceo tereti striato usque ad apicem foliato, foliis semisimplexicaulibus elongato-lanceolatis acuminatis integerrimis annuis basi multinerviis, capitulis ad apicem aggregatis breviter pedunculatis multifloris, invol. campanulati squamis in acumen longum subspinosum desinentibus. 2? in serrâ de Piedade prov. Min. Ger. legit. cl. Martius. *Proteopsis* argentea Mart. et Zucc! herb. Achænia glaberrima angulata. Pappi setæ 2-seriales lineari-acuminatæ ciliatæ caducæ inter se similes. An *Decaneuro* an *Hololepidi* affinis? (v. s. in h. acad. Reg. Monac.)

6. V. ANDRIEUXII, caule frutescente tereti arancoso et piloso-glanduloso, foliis oppositis petiolatis elliptico-lanceolatis utrinque acuminatis argutè et remotè serratis hinc inde sublobatis supra breviter scabridis subtus densè cano-tomentosis, capitulis ternis pedunculatis, invol. squamis inermibus multiserialibus elongatis squaroso-subtortis, achænio adpressè pubescente. 3 in Mexico inter Tehuantepec et flum. Guaracoalcos septemb. flor. legit. cl. Andrieux pl. exs. n. 269! Pappus rufescens, ext. squamellatus brevis. Petioli 1; poll. longi. Folia 4-5 poll. longa, aequipollicem lata. Species pulchra priori habitu affinis sed fortè caract. nimis aliena. (v. s. comm. cl. Andrieux.)

§. 3. *Ericoideæ, nempe fruticosaæ, ramis usque ad flores folia conferta gerentibus, apice capitula cylindracea gerentibus.*

7. V. BUXTORFII (Less. in *linnæa* 1829. p. 247. 1831. p. 629), fruticulosa, ramulis apice angulatis tomentosis, foliis sessilibus confertis obovatis integerrimis coriaceis supra glabris lucidis, subtus densè tomentosis, ramis floridis ad apicem usque foliosis, capitulis cylindraceis 7-floris, invol. pauciserialis squamis siccis 3-ner-

vūs obtusis, achenio multicostrato. ♀ in Brasiliā tropicā. Cor. glabrā purpureæ? Pappus flavescens. Foliā 3-4 lin. longa, a lin. lata.

8. *V. ERICOIDES* (Less. l. c. 247. excl. syn. et Less. in Linnæ 1831. p. 629), fruticosa, ramis confertis teretibus tomentosis, foliis sessilibus 1-nerviis coriaceis confertis linearibus integerrimis suprà glabris subtus tomentosis, ramulis ad capitula usque foliosis, capitulis cylindraceis multifloris, invol. squamis oblongis acutis basi trinerviis, achenio multicostrato, pappo biseriali, paleis aequalibus ciliatis. ♀ in Brasiliā tropicā (Less.), in monte Itambe da Villa do Principe prov. Minarum (Mart!) *Vernonia dipicola* Mart! herb. Cor. purpureæ? glabræ. Pappus flavescens. (v. s. in h. Acad. reg. Monac.)

9. *V. OLAEASTER*, caule fruticoso ramoso tereti, ramis cano-tomentosis, foliis oblongis subsessilibus integerrimis 1-nerviis subobtusis suprà virescentibus densè velutinis subhis cano-tomentosis, capitulis ad apices ramorum circ. 3 sessilibus congestis 20-floris, invol. multiserialis squamis lineari-lanceolatis acuminatis subtomentosis longè ciliato-barbatis. ♀ in Serra de Piedade prov. Minas-Geraes Brasiliæ legit. cl. Martins. *Albertinia Oleaster* Mart! herb. Folia pollicem longa, 2-3 lin. lata. Achenium glaberrimum. Pappus biserialis setis caducissimis albidis longitudine aequalibus! Receptaculum punctatum. (v. s. in herb. Reg. Monac.)

10. *V. PARUDO-MYRTUS* (S. Hil. voy. Diam. 1. p. 94 et 367), caule fruticuloso erecto ramoso tereti, ramis apice angulatis tomentosis, foliis confertis parvis petiolatis lanceolatis 1-nerviis obtusis suprà nigro-punctulatis, subtus albo-tomentosis, capitulo terminali solitario inter folia sessili, invol. cylindrici squamis linearibus obtusis ciliatis, achenio glabro 10-costato, pappi seriebus aequalibus, paleaceis barbelatis supernè capillaceis. ♀ in Brasiliæ Serra de S. Antonio. Cor. pallidè ceruleæ aut. purpureæ. Fol. 4-5 lin. longa, æquilineam lata, petiolo 1-2 lin. Pappus rufescens.

### Sect. III. LEPTOSPERMOIDES.

Capitula solitaria terminalia multiflora. Achenia sericeo-villosa. Pappi series inæquales, externā paleacè brevi. — Species Americanæ et Madagascarienses.

11. *V. AENEOPHYLLA* (Mart! herb.), caule fruticuloso ramoso tereti basi folioso inter folia villosa cæter. glabro, foliis sessilibus obovato-oblongis integerrimis obtusis vix mucronulatis penninerviis nervis utrinquè prominulis utrinquè glabris subtus crebrè glandulosis basi sparsè piligeris, ramis subaphyllis subtilissimè striatis 1-cephalis, capitulo 20-38-floro, invol. hemisphærici squamis oblongis obtusis dorso nigro-puberulis pappum æquantibus, cor. glabrā glanduliferā, achenio striato sericeo-villoso pappi ser. ext. angustè paleacè acutè brevi. ♀ in altis Serro Frio prov. Minar. Gener. Brasiliæ legit. cl. Martins. Species distinctissima vix semipedalis, habitu ferè Vernoniæ fruticulose. Folia 8-10 lin. longa, 3-4 lata. Pedunc. 2 pollic. Cor. purpureæ. (v. s. in h. Acad. reg. Monac.)

12. *V. LEPTOSPERMOIDES* (Mart! herb.), fruticosa, ramis teretibus suboppositis, junioribus tomentosis, adultis glabris, foliis confertis sessilibus coriaceis integerrimis, junioribus utrinquè molliter hirsutis, adultis suprà glabris subtus velutinis, nervo medio prominente, capitulis 9-floris ad apices ramulorum solitariis inter folia sessilibus, invol. squamis paucis oblongis obtusiusculis enerviis ciliatis dorso apice puberis. ♀ in campis editis prov. Minasum generalium Brasiliæ ad Tejuco, Serro Frio, legit cl. Martins. Achenia obovata substriata rufa glabriuscula. Pappus biserialis stramineus, exter. brevis, interior quadruplò longior scaber. Affinis ex descr. videtur *V. ericoidi* et *buxoidi*, sed pappi ext. brevi à sectione differt. (v. s. in h. Acad. reg. Monac.)

13. *V. POLYGALAEFOLIA* (Less. in Linnæ 1831. p. 628. non Licht.), caule fruticoso diffuso, ramis teretibus pube brevissimā stellato-lepidotā supernè pubescentibus, foliis brevissimè petiolatis coriaceis oblongo-obovatis obtusis integerrimis sparsè lepidotis, capitulis 1-3 terminalibus sessilibus 9-floris, invol. cylindracei squamis imbricatis acuminatis dorso albo-pilosis longè ciliatis, achenio piloso, pappi ser. ext. brevi capillari. ♀ in montibus altis prov. Emirau insule Ma-

dagascar. *Vernonia procumbens* Bojer! in litt. 1835. Folia 4-7 lin. longa, 1  $\frac{1}{2}$ -4 lin. lata; cor. lobi revoluti; limbus à tubo non distinctus. Stylus bulbosus. Pappus ferrugineus. (v. s. comm. à cl. Bojer.)

14. *V. GLUTINOSA*, fruticosa, ramis teretibus junioribus brevissimè velutinis foliisque supernè glutinosis, foliis obovatis ovalibusve basi cuneatis subpetiolatis tenèè quintuplinerviis reticulatis rigidulis integerrimis, capitulis 1-3 sessilibus terminalibus, invol. squamis lanceolato-linearibus subviscosis, achenio sericeo-villoso, pappo multiseriali, setis exterioribus paucis paulò brevioribus.  $\S$  in insulâ Madagascar propè urbem Tannanarivou, legit. cl. Bojer. *Vernonia pauciflora* Bojer! in litt. non Less. Frutex 4-5-pedalis. Folia juniora rufescentia. Pappus rotundus invol. paulò longior. Flores in spec. meo inexplicati. (v. s. comm. à cl. Bojer.)
15. *V. ARGYROPHYLLA* (Less. in *linnæa* 1831. p. 627), ramis ad apicem usque foliatis, foliis approximatis coriaceis brevissimè petiolatis obovatis acutis suprâ denum nitidis subtus argenteo-tomentosis, capitulis terminalibus solitariis, invol. campanulati squamis siccis muticis acuminatis extus pilosis, achenio villosis, pappo plariseriali, ser. ext. breviori latiori.  $\S$  ? in Brasiliâ. Cor. glabra purpureo-violacea.
16. *V. MONTICOLA* (Mart! herb.), fruticosa, ramis teretibus glabris, foliis sessilibus erectis coriaceis integerrimis oblongis margine latè revolutis suprâ junioribus sparsè puberulis demum glabris punctulatis subtus adpressè tomentosis, capitulis 5-floris oblongo-cylindricis 2-5 ad apices ramulorum aggregatis in summis axillis aut apice ipso sessilibus, invol. squamis siccis rufis glabris acuminatis.  $\S$  in monte Itambe prov. Minarum Brasiliæ, legit. cl. Martius. Habitus *Lychnophoræ*, sed capitula distincta nec verè glomerata. Achenium pubescens oblongum. Pappi exter. paleæ latè acuminatæ, inter. duplò triplòve longiores setiformes, omnes albæ. An *Chrysocoma adpressa* Arrab. fl. flum. 8. t. 35? (v. a. in h. Acad. reg. Monac.)

Sect. IV. *VANILLOSUM* Less. in *linnæa* 1831. p. 630. — *Vern. axillifloræ* Less. in *linnæa* 1829. p. 251.

Capitula pauciflora aggregata fasciculata, fasciculis axillaribus folio suo brevioribus. — Frutices Americani. Folia petiolata penninervia æspissimè subtus incana.

### §. 1. *Acheniis glabris.*

#### \* *Capitulis sessilibus.*

17. *V. SPLENDENS* (Less. in *linnæa* 1829. p. 252. et 1831. p. 630), fruticosa, caule tereti glabro, ramis flexuosis vix apice petiolisque subtomentosis, foliis petiolatis coriaceis oblongis integerrimis suprâ lucidis subtus tomento albo adpresso, capitulis spicato-congestis sessilibus 3-floris, invol. squamis coriaceis obtusis 1-nerviis ciliatis extus pubescentibus, achenio glaberrimo multicostato, pappi serie exter. brevi.  $\S$  in Brasiliâ tropicâ, ad sepes circa Rio-Janeiro. Pappus argenteus. *Conyza lucida* Spreng. neu entl. 2. p. 142. *Conyza splendens* Spreng. syst. 3. p. 509. *Carpholobus sessiliflorus* Schott. in Spreng. cur. post. 409. *Vanillosum argyrophyllum* Mart! herb. Variat foliis ovatis oblongis lanceolatisve, paginâ super. venis subprominulis reticulatâ. Pappi ser. ext. multi-setosâ, inter. triplò brevior. (v. s.)
18. *V. ISOTRICHIA*, caule fruticoso tereti glabro, ramis flexuosis junioribus petiolisque velutino-tomentosis, foliis petiolatis ellipticis breviter acuminatis integerrimis suprâ glabris lucidis subtus velutino-tomentosis, capitulis 3-floris ad axillas foliorum sessilibus 7-8 glomeratis, invol. squamis obtusis intimis glaberrimis, achenio glaberrimo pappi setis omnibus æqualibus.  $\S$  in fruticetis circa Rio de Janeiro legit. cl. Lhotsky. *V. splendens* similis, sed pappus isocharis, achenium dimidio brevius, invol. glabriora paulò longiora, folia sæpè junctura subreticulata adulta levia, petioli 5 lin. longi. Folia 2-3 poll. pæ, pollicem œquipollicemve lata. (v. s.)

19. *V. LEPROSA* (Less! in *linnæa* 1829. p. 252), fruticosa, ramis angulatis tomento adpresso fulvo petiolisque 4-gonis obtectis, foliis coriaceis oblongis acuminatis integerrimis pellucido-glandulosis glabris subtus leprosis, capitulis paucis apicato-glomerulatis 2-3-floris, invol. squamis subrotundis siccis 1-nerviis, achenio glaberrimo glanduloso multicostato, pappi serie externâ caducâ brevi.  $\frac{5}{3}$  in Brasiliâ tropicâ. Pappus argenteus.
20. *V. ROTUNDIFOLIA* (Less! in *linnæa* 1829. p. 254. 1831. p. 631), fruticosa, ramis teretibus sulcato-striatis petiolisque cano-tomentosis, foliis petiolatis coriaceis elliptico-rotundis retusis, supra glabris nervis venisque impressis, subtus densè hirsuto-velutinis nervis prominulis, capitulis 3-4 petiolo brevioribus glomeratis 3-floris, invol. squamis ovatis acutis plurinerviis glabris ciliatis, achenio glaberrimo multicostato, pappi paleis exterioribus paucis et interioribus æqualiter angustis.  $\frac{5}{3}$  in Brasiliâ tropicâ. Cor. purpureæ glabræ. Pappus argenteus. *Vanillosuma firmum* Mart! herb. (v. s.)
21. *V. PYRIFOLIA*, fruticosa glaberrima, ramulis apice angulatis, foliis breviter petiolatis ovalibus oblongive obtusis rigidis supra nitidis, subtus reticulato-venosis, capitulis in axillis sessilibus congestis 3-floris, invol. squamis oblongis obtusis, achenio glaberrimo, pappi 3-4-seriali, serie exter. interiore multò brevior.  $\frac{5}{3}$  in Brasiliâ prov. Rio de Janeiro, leger. cl. Martins et Vauthier (coll. n° 3421) *Vanillosuma pyrifolium* et *V. obtusum* Mart! herb. Folia variant.  $\frac{4}{3}$  poll. longa, vix pollicem lata, et 2-3 poll. longa,  $\frac{1}{2}$  lata, petiolo 3-6 lin. sub lente vix subpuberulo. (v. s.)
22. *V. OXYPHYLLA*, fruticosa, ramis teretibus junioribus petiolisque pube ferrugineâ velutinis, foliis longè petiolatis lanceolatis longè acuminatis integerrimis, supra glabris, subtus in nervis subhirsutis, utrinquè nervulis elevatis reticulatis, capitulis ad axillas plurimis sessilibus confertis 3-7-floris, invol. squamis exter. dorso velutinis intimis longioribus glabrisculis subobtusis, achenio glaberrimo pappo longiore, pappi setis ext. brevibus.  $\frac{5}{3}$  in sylvis propè Rio-de-Janeiro legit cl. Martins. *Vanillosuma acuminatum* Mart! herb. ex parte. Petiolus 9 lin. longus, folia 5 poll. longa, sesquipoll. lata. Affinis *V. pyrifolies*. (v. s. in h. Acad. Monac.)
23. *V. NOTATA* (Less. in *linnæa* 1829. p. 256), fruticosa subscandens, ramis angulatis tomento adpresso cinereis, foliis petiolatis rigidis lanceolatis integerrimis supra glaberrimis subtus lepidoto-tomentosis, capitulis fasciculatis petiolo subæqualibus 3-floris, involucris squamis siccis 1-nerviis apice maculâ tomentosa notatis, achenio glaberrimo, pappi serie ext. paulò, int. brevior.  $\frac{5}{3}$  in Brasiliâ. Cor. glabræ purpureæ. Pappus argenteus. Inflorescentiæ rami 1-3-cephali. (v. s. comm. à Mus. reg. Berol.)
24. *V. AXILLARIS* (Less! in *linn.* 1829. p. 256), fruticosa, ramis teretibus petiolisque fulvo-tomentosis, foliis oblongo-ellipticis subintegerrimis acutis coriaceis adultis supra glabris, junioribus supra et omnibus subtus lanato-tomentosis, capitulis cylindraceis circ. 6-floris ad axillas foliorum glomeratis sessilibus, invol. foliolis coriaceis ovato-oblongis glabris, achenio oblongo subtrigono glaberrimo.  $\frac{5}{3}$  in Brasiliâ tropicâ. Pappi straminei series externa caduca brevis, multò magis quam interior lata. (v. s. comm. à Mus. reg. Bav.)
25. *V. MACROPODA*, fruticosa, ramulis petiolisque striato-angulatis densè velutinis, foliis ovalibus utrinquè subattenuatis rariter subdenticulatis supra adultis glabris, subtus cano-velutinis nervis elevatis reticulatis, capitulis ad axillas 3-6 sessilibus petiolo brevioribus circ. 20-floris; invol. obovati squamis siccis dorso velutinis imbricatis, ext. latè ovatis, int. elongatis, achenio glaberrimo, pappi serie ext. brevi paulò quam inter. lata.  $\frac{5}{3}$  in Brasiliâ, sed locus proprius mihi ign. Petioli 12-15 lin. longi. Folia 5-6 poll. longa, 2-3 lata. (v. s.)

\* \* *Capitulis plurimis pedunculo ramoso insidentibus.*

26. *V. RAMIFLORA* (Less! in *linnæa* 1829. p. 255), fruticosa scandens, ramis teretibus junioribus cinereo-velutinis, foliis petiolatis coriaceis ovalibus subacutis subintegerrimis supra glabris lucidis, subtus lepidoto-cinereis albidisve, fasciculis

subumbellatis axillaribus, capitulis cylindraceis pedunculatis 2-3-floris invol. squamis ovatis acutis subciliatis adpressis, achænio oblongo glabro, pappi serie externâ paucisetosâ interiore triplâ brevior.  $\frac{3}{4}$  in Brasiliâ tropicâ. Cor. purpureæ glabræ. Pappus argenteus. (v. s.)

27. *V. CAPITELLATA*, fruticosa, ramis subangulatis tenuissimè velutinis, foliis petiolatis ellipticis utrinquè attenuatis suprà glabris, subtus petioloque tenuissimè subvelutinis utrinquè nervulis elevatis reticulatis, pedunculis ad axillas fol. brevissimis ramoso-corymbosis polycephalis subvelutinis, capitulis ovatis 3-floris, invol. squamis ext. brevibus obtusis dorso velutinis, int. oblongis glabrisculis, achænio glaberrimo, pappo brevior, pappi setis ext. brevibus.  $\frac{3}{4}$  in Brasiliâ (prope Rio-de-Janeiro?) legit cl. Martins. Vanillosum acuminatum Mart! herb. ex parte. Certissimè ab oxyphyllâ diversa! Proximè ad *V. tereticanlem* accedit. Achænia capituli 3 sæpè conglutinata. Petioli 6 lin. longi. Limbi 3 poll. longi 15 lin. lati. An *Chrysocoma verticillata* Arab. fl. sum. 8. t. 26? (v. s. in h. Acad. reg. Monac.)

28. *V. TERETICAULIS*, caule fruticoso, ramis junioribus teretibus pubescenti-velutinis, foliis petiolatis ellipticis subacutis basi sæpius inæquilateris integerrimis suprà glabris, subtus petiolisque velutino-tomentosis, pedunculis in axillis plurimis umbellatis petiolo brevioribus apice ramosis, capitulis ovatis 4-5-floris subvelutinis, squamis ovatis acutis, achænio gracili glaberrimo, pappi serie ext. paucisetosâ brevi.  $\frac{3}{4}$  in Americâ calidiore? (Nee!), ad Guayaquil (Haenk!) Affinis *V. Poeppigianæ*, sed differt achænio longiore non costato vix striato, pappo albissimo nec subrufescente, foliis paulò minoribus et magis coriaceis, etc. Petioli infer. interdum ovata. Petioli 5-6 lin. longi. (v. s. in herb. Haenk.)

29. *V. BLANCHETIANA*, fruticosa, ramis subacandentibus teretibus lepidoto-velutinis cinereis, foliis petiolatis ellipticis subacutis integerrimis suprà glabris, subtus petioloque lepidoto-velutinis, pedunculis in axillis confertis ramosis polycephalis, capitulis ovatis 5-floris, squamis glabris subciliatis ovatis obtusis ext. brevissimis, achænio glabro multicostato, pappi serie ext. multisetosâ brevi.  $\frac{3}{4}$  in Brasiliâ circa Bahia legit cl. Blanchet. Petioli 9 lin. longi. Folia 5-6 poll. longa, 2  $\frac{1}{2}$ -3-lata, basi sæpè inæqualia. Affinis *V. Poeppigianæ*, sed rami teretes nec apice angulati, folia minus acuminata, invol. minora. Flores 5, nec 7-8, etc. (v. s. comm. à cl. Blanchet.)

30. *V. POEPPIGIANA*, fruticosa, ramis tenniter velutinis apice angulatis, foliis petiolatis ellipticis acuminatis suprà glabris, subtus petiolo et in basi nervi medii paginæ super. subadpressè lepidoto-velutinis subrufescentibus, pedunculis ramosis axillaribus petiolo brevioribus fasciculato-polycephalis, invol. ovatis glabris 7-8-floris, squamis siccis ovatis, intimis subacutis, achænio glaberrimo, pappi setis brevioribus paucis.  $\frac{3}{4}$  in Peruvâ legit cl. Poeppig. Pl. ign. n° 23 et 1425. Petioli 9 lin. longi, Folia 5-6 poll. longa, 2-2  $\frac{1}{2}$  lata, basi sæpè inæqualia. Species *V. ramulifloræ* accedens, sed capit. ovata (nec cylindracea) pluriflora. (v. s. comm. à cl. Poeppig.)

31. *V. UMBELLATA* (Less! in linnæâ 1829. p. 255. 1831. p. 631), fruticosa scandens, ramis teretibus glabris vix apice pubescentibus, foliis petiolatis lanceolatis utrinquè acuminatis integerrimis suprà glabris, subtus dense albo-tomentosis, fasciculis umbellatis axillaribus, capitulis pedicellatis cylindraceis 7-9-floris invol. squamis subscariosis subrotundis glabris internis oblongo-linearibus, achænio glabro, pappi serie externâ brevi caducâ.  $\frac{3}{4}$  in Brasiliâ tropicâ. *Conyza ramiflora* Spreng. syst. 3. p. 509. Cor. purpureæ glabræ. Pappus argenteus. (v. s.)

32. *V. LUNDIANA* (Less! in linnæâ 1831. p. 631), caule arboreo ramis deinde atriatis tomentoso-lepidotis, foliis petiolatis oblongo-ellipticis utrinquè obtusissimis integerrimis suprà laevibus, subtus densissimè tomentoso-lepidotis, capitulorum fasciculis axillaribus petiolo brevioribus, capitulis plus quàm tria, invol. cylindraceis squamis obtusis apicè subpubescentibus, achænio immo.  $\frac{3}{4}$  in Brasiliâ ad Serra de Mar, prope novum Friburgum, colousam

Helvetiorum. Arbor 20 ped. octobri florens, odorem Philadelphi coronati spirans. (Lund!) Cor. albida. Petioli 12-15 lin. longi. Folia 8-10 poll. longa, 5 et parum ultra lata. (v. s. comm. à cl. Lund.)

§. 2. *Achæniis pilosis. — Species à Vanillosmâ subalienæ et inter se fortè vix cognatæ!*

33. V. VAUTHIERIANA, caule fruticoso ramosissimo tereti, ramulis angulatis pulverulento-incanis foliis breviter petiolatis lanceolatis acuminatis basi cuneatis integerrimis suprà adultis glabris, subtùs pube adpressissimâ canescentibus, capitulis ad axillas fol. 3-5 breviter pedicellatis confertis 10-floris, invol. squamis oblongis subacutis dorso præsertim ad apicem pubescentibus, cor. lobis apice villosis, achænio turbinato brevi sericeo-villosissimo, pappi serie ext. paleis ovatis acutis inter. multò brevioribus. § in Brasilia prov. Minasrum gener. legit. cl. Vauthier. coll. n° 336. Folia cum petiolo ferè 3 poll. longa, pollicem lata.

§. *angustifolia*, foliis lineari-lanceolatis. § cum priore. Vauth! coll. exs. n° 337. (v. s. comm. à cl. Vauthier.)

34. V. ELIASHOIDES (H. B. et Kunth nov. gen. 4. p. 42), fruticosa, ramis teretibus tenuissimè sericeo-pubescentibus, foliis brevissimè petiolatis coriaceis elongato-lanceolatis obtusis integerrimis suprà glaberrimis lucidis, subtùs sericeo-tomentosis, capitulis ad axillas glomerulatis campanulatis 15-20-floris, invol. squamis coriaceis 3-nerviis ovatis acutis extùs pilosis, achænio oblongo piloso, pappi serie ext. brevi latiusculâ. § in montosis prope flavium Caellar Novo-Granatensium. Lem. in linnæâ 1829. p. 254. 1831. p. 631. Cor. glabra, Pappus stramineus.

35. V. CANDOLLEI (Mart! herb. sub Vanillosma), caule fruticoso tereti, ramisque subangulatis cano-tomentosis foliis breviter petiolatis lanceolatis oblongiève subacutis penninerviis suprà glabris reticulato-venulosis, subtùs petioloque cano-tomentosis, pedunculis axillaribus petiolo longioribus ramoso-corymbosis incanis, capitulis oblongis 7-floris, invol. squamis acutis subpuberulis, cor. lobis tenuissimis glabris, stylo breviter bifido lobis subobtusis, achænio pilosculo, pappi setis ext. paucis brevioribus. § in campis editis ad Rio de Contas prov. Bahiensis, legit. ill. Martins, Folia majora 3 poll. longa, 12-15 lin. lata, petiolo 3-lin. Species distinctissima fortè propriæ sectionis? (v. s. in h. Acad. reg. Monac.)

36. V? GAUDICHAUDIANA, fruticosa, ramis teretibus adultis glabris junioribus vix subcanescentibus, foliis petiolatis ovato-lanceolatis acuminatis integerrimis membranaceis suprà glabris, subtùs adpressè cano-tomentosis, capitulis axillaribus 8-10 umbellulatis pedicellatis, pedicellis vix petiolo longioribus parè ramosis, invol. ovali squamis glabris subobtusis § circa Rio de Janeiro legit cl. Gaudichaud. Flores fructusque ignoti, undè species dubiâ. Accedit hinc ad V. reticulatam illinc ad V. Candollei, sed diversa videtur. Petiol. 4 lin. longus; limb. fol. 3 poll. longus, pollicem latus. (v. s. comm. à cl. inv.)

Sect. V. STROBILYX, Blum. in litt.

Invol. squamæ imbricatæ ovatæ obtusæ brevissimæ et floribus multò superatæ. Involucrum 3-5-florum teretiusculum. Pappus 1-serialis aut setis ext. paucis calyculatus. — Arborea Indica aut Africana. Folia petiolata plana penninervia.

37. V. CELEBICA, arborea, foliis longè petiolatis oblongo-ellipticove lanceolatis basi subinaequaliter attenuatis cuspidatis integerrimis glabris cum nervis subtùs petiolo ramulisque pubescenti-velutinis, corymbo terminali divaricato, capitulis 3-4-floris, invol. squamis imbricatis obtusis ciliolatis, achænio teretiusculo substriato pubero invol. circ. longitudine. § in sylvis insulæ Celebæ. Eupatorium Celebicum Blum! bijdr. 903. Leonconeris celebica Blum! in litt. Affinis V. arboreæ, sed pappus duplex, ext. brevis. Folia 5-poll. longa sesqui-poll. lata, petiolo pollicari. (v. s. comm. à cl. Blum.)

38. *V. JAVANICA*, arborea, foliis longè petiolatis ellipticis acuminatis basi rotundatis integerrimis suprà glabris, subtus reticulatis ramulique fusco-tomentosis. corymbo terminali paniculato divaricato, ramis lateralibus elongatis, invol. squamis ovali-oblongis obtusis imbricatis dorso villosis.  $\zeta$  in sylvis altioribus insulæ Java ubi dicitur *Hambirung*. *Eupatorium Javanicum* Blum! bijd. 903. *Leucomeris Javanica* Blum! in litt. Arbor 50-60 ped. Folia petiolo sesqui-poll., limbo 6 poll. longo, 2  $\frac{1}{2}$  3 poll. lato. Paniculæ rami laterales 3 poll. et ultrà longæ. Pappi series ext. paucisetosa vix ulla (v. s. comm. à cl. Blum.)
- $\beta$ ? *conferta*, foliis oblongo-lanceolatis acuminatis basi attenuatis et subtus ad marginem subreplicatis suprà glabris, subtus junioribus cum ramis et pedunculis densè rufo-velutinis, adultis glabriusculis, paniculâ confertâ, ramis nempe ferè à basi ramulosis, invol. squamis extus densè velutinis.  $\zeta$  in Javâ. Specimen habeo à cl. Blume cum var.  $\alpha$  sub nomine *Eup. Javanici* mixtum, sed meo sensu diversum. (v. s.)
39. *BLUMIANA*, arborea, ramis teretibus glabris in paniculâ pulverulento- et rufo-velutinis, foliis petiolatis oblongo-lanceolatis basi breviter apice longè acuminatis integerrimis utrinquè glabris, paniculâ corymbosâ compositâ, invol. squamis ovali-oblongis obtusis dorso subvillosis.  $\zeta$  in montanis Karang Javæ occidentalis. *Leucomeris glabra* Blum! in litt. Pappus sordidus, serie ext. brevi pauciseta. Folia petiolo 4-5 lin. longo, limbo 3-4 poll. longo, 10-14 lin. lato. (v. s. comm. à cl. Blume.)
40. *V. ARBOREA* (Ham! in trans. lin. soc. lond. 14. p. 218), arborea, ramis teretibus substriatis glabriusculis, foliis petiolatis ovato-lanceolatis acuminatis integerrimis paucinerviis glabris exceptis nervo pag. infer. et petiolo subvelutino-furfuraceis, cymæ terminalis ramosissimæ paniculæformis rachi angulatæ ramisque furfuraceis, invol. squamis paucis ovato-oblongis demum patentibus, æchenio turbinato glabro, pappi setis 1-serialibus aequalibus.  $\zeta$  in sylvis Nepauliæ inferioris et Camrupæ (Ham. tr.) et ad Gaspura Beng. (Ham. herb.) *Conyza arborea* Wall! cat. n° 2. Cor. purpurea 6-7 in capitulo unico, sed vidi æchenia 3 in unicum concreta! (v. s. comm. ab hon. cœt. merc. Anglorum Ind. or.) *Mogol* Bengalensium.
41. *V. ELÆAGNIFOLIA*, caule ramoso tereti pubescenti-albido, foliis brevè petiolatis lanceolatis acuminatis integerrimis suprà glabris, subtus villis adpressis sericeis argenteis, paniculâ ramosissimâ, ramulis ultimis capitula 3-5 gerentibus, involucri pauciflori squamis ovatis paucis marginè membranaceis subciliatis, æchenio glanduloso glabro, pappi serie externâ paucisetosâ brevi.  $\mathcal{L}$   $\zeta$  in Indiâ orient. ad Moalmyne. Wallich. *Conyza elæagnifolia* Wall! cat. n° 3041. comp. 151. *Eupatorium elæagnifolium* hort. calc. 1797. Pappus argenteus. (v. s. comm. ab hon. cœtu merc. angl. Ind. or.)
42. *V. ELLIPTICA* (DC. in Wight, contrib. bot. ind. p. 1), caule scandente ramoso tereti subpubescente, foliis breviter petiolatis exactè ellipticis integerrimis suprà glabris, subtus villis adpressis sericeis argenteis, paniculâ ramosissimâ divaricatæ ramis ultimis capitula 3-5 pauciflora gerentibus, invol. ovatis squamis ovatis subacutis glabris.  $\zeta$   $\mathcal{L}$  in Ind. orient. montibus Neelgerry legit. cl. Wight. *Vernonia* sp. scandens Wight! herb. Ex foliis et inflorescentiâ valdè accedit ad *V. elæagnifoliam* sed car. gener. ex statu specimin. imperfecto non vidi. (v. s. in herb. Wight.)
43. *V. SECUNDIFOLIA* (Bojer! in litt. 1835), fruticulosa glaberrima, ramis teretibus flexilibus subscandentibus, foliis petiolatis secundis ellipticis basi cuneatis apice mucronatis serratis coriaceis nitidis, corymbo composito subfastigiato, pedunculis axillaribus folio paulò longioribus apice corymbiferis, capitulis brevè pedicellatis 5-floris, invol. squamis lineari-oblongis vix acutis, æcheniis glabris, pappi serie ext. brevi paucisetâ.  $\zeta$  ad margines sylvarum nov. Emirnenis ins. Madagascar legit cl. Bojer. Cor. purpurea. Invol. apices rufescentes. Pappus albus. Folia. 1  $\frac{1}{2}$  poll. longa, 9 lin. lata. (v. s. cl. Bojer.)
- IMBROSA*, fruticulosa alte scandens, ramis angulatis, foliis petiolatis

elliptico-oblongis utrinque acutis integerrimis utrinque pube minimâ sparâ puberulis subtus glandulas parvas sessiles gerentibus, capitulis ad apices ramulorum circ. 3 breviter pedicellatis, invol. squamis ovatis obtusiusculis ciliatis, acheniis angulatis glabris, pappi serie externâ paucisetâ brevi.  $\zeta$  in sylvis Réfourou dictis prov. Ré-tani-mena Madagascar insulis super arbores scandens. Vern. scandens Bojer! in litt. 1835, non Wall. Flores flavidî in invol. 7. Folia 4 poll. longa, pollicem lata. Pappus flavescens rigidulus. (v. s. comm. à cl. inv.)

45. *V. DOWNIA*, fruticosa, ramulis teretibus pube ferrugineâ pedunculis petiolis nervoque medio foliorum subtomentosis, foliis petiolatis ellipticis utrinque acutis penninerviis nervo excepto glabris, paniculâ elongatâ terminali laxâ aphyllâ, invol. squamis 3-4-seriatis siccis glabris ext. brevibus ovatis, int. oblongis, achenio glabro striato-angulato, pappi albisimi serie ext. brevi setiformi.  $\zeta$  in Sierra Leona legit cl. Don. Eupatorium paniculatum herb. Lindl. Folia superiore (cum petiolis 4. lin.) 4 poll. longa, 15-18 lin. lata. Cor. non vidi (v. s. in h. Lindl.)

#### Sect. VI. TRIANTHÆA.

Capitula 3-flora rariùs 4-flora oblonga in corymbum dichotomum disposita.

46. *V. TRIMFLORA* (Less. in linnæâ 1831 p. 633), caule fruticoso, ramulis angulatis brevissimè tomentosis, foliis petiolatis membranaceis lanceolatis integerrimis suprà nitidis, subtus lepidoto-tomentosis, paniculâ terminali dichotomâ, capitulis breviter pedicellatis oblongo-cylindraceis 3-floris, invol. squamis arcetè imbricatis glabris ciliatis ovatis acutiusculis, achenio costato glabro, pappi ser. ext. paleacea brevissimâ, int. lingulatâ.  $\zeta$  in ins. Madagascar. Folia 3-3½ poll. longa, pollicem lata, petiolo 3 lin.
47. *V. TRIFLOSCULOSA* (H. B. et Kunth nov. gen. 4. p. 40), fruticosa, ramis teretibus glabris, foliis petiolatis lanceolato-oblongis apice basiue acutiusculis integerrimis membranaceis suprà glabriusculis subtus pubescentibus, cymâ paniculato-ramosissimâ foliatâ, capitulis sessilibus fasciculatis 3-floris, invol. cylindracei squamis acutis ciliatis, ext. minimis, cor. glabrâ, achenio canescente, pappi serie ext. brevi.  $\zeta$  in montibus Mexicanis prope Acaguila et in calidis prov. Vera-Cruz. Gymnanthemum congestum Cass. dict. 20. p. 110. et 26. p. 20. Pappus albidus. Cor. alba ex Kunth, flavescens ex Cass.
48. *V. DICHOCARPHA* (Less. in linnæâ 1829. p. 269. 1831. p. 634. non Spreng.), arborea, ramis teretibus fulvo-tomentosis, foliis petiolatis coriaceis ellipticis acutis integerrimis suprà glabris, subtus petiolisque fulvo-tomentosis, cymis terminalibus subcorymbosis confertis, capitulis cylindraceis 3-floris, invol. squamis siccis 1-nerviis extimis ovatis tomentosis, intimis linearibus glabris, achenio multicostato subpilosiusculo, pappi serie externâ paleacè brevi, interiori paleis tortis.  $\zeta$  in Novâ Andalusia juxta Caripe Pollalesta vernonioides H. B. et Kunth nov. gen. 4. p. 36. t. 321. Oliganthes triflora Cass. dict. 36. p. 18. Eupatorium cydoniæfolium Willd. herb. ex Less. Cor. glabræ albæ! glandulosæ. Pappus flavescens.
49. *V. SIMBRILLIFERA* (Less. in linnæâ 1829. p. 109), fruticosa, ramis striatis tomentoso-pubescentibus, foliis ovato-lanceolatis acuminate in petiolum attenuatis subdenticulatis suprà glabris, subtus glandulosè pubescentibusque, cymâ paniculæformi terminali ramosissimâ aphyllâ, capitulis pedicellatis 8-4-floris, involucri pauciserialis squamis ovalis pubescentibus, receptaculi simbrillitis setiformibus, achenio hispido, pappi setis inæqualibus.  $\zeta$  in ins. Bourbonia. Pappus rufescens. Cor. flavæ dicuntur, undè genus esset dubium, sed cl. Camini flavos dixit flores ferè omnes decoloratos. Gymnanthemum simbrilliferum Cass. dict. 20. p. 109. Eupatorium coriaceum Vahl symb. 3. p. 95. non Spreng.
50. *V. PANICULATA*, fruticosa, ramis teretibus cano-tomentosis, foliis breviter petiolatis ovalibus integerrimis supernè puncta resinosa (glandulose?) conferta gerentibus glabris, subtus petiolisque cano-tomentosis, paniculâ maximâ



myriocephalâ, capitulis ad apices ramulorum congestis oblongis, invol. squamis oblongis macronatis extus araneosis, intus et intimis glabris, fl. 1-3?... 3 in Mexico ad San-Dionysio in ditione Oaxaca legit cl. Andrieux pl. exa. n. 270! Capitula speciminis mei floribus fructibusque orbata, et idè de genere imò de sectione dubito, sed ob habitum speciosum præterire nolui. (v. s.)

Sect. VII. TEPHAODES. — Isomeria Don in Wight. herb.

Invol. squamæ imbricatæ lanceolato-lineares acuminatæ rariis acutè subtrinerviæ disco breviores, ext. breves, int. sensim longiores demum reflexo-patentes. Achenium teres villosum. Pappus ext. brevissimus regularis, int. 1-septalis albus tenuis. — Herbæ gerontogæ pleræque annuæ cinereæ (undè nomen). — Nomen Isomeriæ optimum cum discrimen erga Conyzam designabat, non servandum cum Vernoniæ sectio indicanda. — Si quis vellet omnes subsequentes species in 1 aut 2 coalescere, non valdè repugnarem.

51. V. *PHYSALEIFOLIA* (DC. in Wight contrib. bot. ind. p. 2), caule herbaceo erecto ramoso striato, foliis petiolatis latè ovatis repando-dentatis obtusis suprâ pubescentibus subtus cauleque cinereo-tomentosis, corymbo laxo, capitulis pedicellatis, invol. squamis lanceolato-linearibus dorso villosis vix acutis. ④? in Indiâ orient. Madagaspatanâ. Conyza cinerea C. Wall! cat. n. 3008. comp. 118. Conyza incana Heyn. herb. Affinis cinereæ, sed distincta videtur ex invol. squamis non acuminatis. (v. s. in herb. Wight. et h. Cur. merc. Ind.)

β. *glabrescens*, foliis tenuioribus et magis glabratâ. — Cum priori. Transit ad sequentem. (v. s.)

52. V. *CINEREA* (Less. in linnæâ 1829. p. 291. 1831. p. 673), caule herbaceo erecto ramoso pilis brevibus confertis subcanescente, foliis petiolatis, infimis subrotundis, cæteris obovato-oblongis subdentatis subtus præsertim cinereo-canescens, corymbo laxo dichotomo, capitulis pedunculatis, invol. squamis lanceolato-linearibus acuminatis subtrinerviis extus pilosiusculis. ① in Indiâ orient. continentali et insulari partibus ferè omnibus, et in Africâ equinoctiali, in ruderatis secus vias etc. Conyza cinerea Linn. 1208. Wall! cat. n. 3903 et 7137. Serratula cinerea Roxb! cat. calc. 60. Isomeria cinerea Wight! herb. Chrysocoma violacea Schum. guin. 158. Chrysocoma purpurea Forst. in h. Ups. ex Less. Conyza mollis Willd. spec. 3. p. 1924 ex Less. Rumph. samb. 6. t. 14. f. 1. Burm. Zeyl. t. 96. f. 1. Pluk. alm. t. 243. f. 3. ic. omnes mal. Herba valdè varians, cujus formæ præcipue hic subjungendæ:

β. *rotundifolia* (DC. in Wight contrib. bot. ind. p. 2), foliis etiam superioribus subrotundis obtusè repando-dentatis. ① in regione Madagaspatanâ frequens ad littora maris. Calalia rotundifolia Willd. phyt. 11. p. 6. t. 1. Wight! herb. Conyza cinerea Wall! cat. n. 1118. D. (v. s.)

γ. *scabrâ*, pube caulis et foliorum non molli et cinereâ, sed magis sparsâ subcabrâ. — in Senegaliâ (Perr.! Sierrâ Leonâ (Smeathm!) Indiâ or. (Wall.!) Conyza cinerea Wall! cat. n. 1118. H. (v. s.)

δ. *glabriuscula* (DC. l. c.), tota ferè glabra, caule striato, ramis elongatis, foliis ferè omnibus oblongis corymbis confertis. — in Indiâ aprica ad Madras et Megapatam legit cl. Wight. (v. s.) An species propria?

ε. *undulata* (DC. l. c.), tota cinereo-pubescentis, caulibus virgatis, foliis lanceolatis dentato-crispis seu undulatis acuminatis. — in Indiâ orient. Conyza cinerascens Wall! cat. n. 3062. comp. 172. Isomeria chinensis Wight. herb. Eandem ut videtur in Senegaliâ legit cl. Perrottet, sed corymbo conferto nec laxo. (v. s.)

ζ. *parviflora* (Reinw. in Blum! hijdr. ned. ind. p. 893), herbacea erecta, ramis fastigiato-paniculatis scabriusculis, foliis caulinis obovatis in petiolum angustatis, floralibus lanceolatis obtusis repandis parum pubescentibus, corymbo terminali dichotomo, squamis invol. apice cuspidatis. ① in Java graminosis et ruderatis, circa Buitenzorg (Blum!), et in insulâ Mauritiâ (Bojer! Boston!) Variat fl. violaceis et roseis et foliis infer. ovatis. Valde affinis V. cinereâ, differt tantum glabritiæ majore et capitulis dimidio brevioribus.

ribus. Pappus ext. brevissimus. Senecio indica foliis ternis Burm. fl. zeyl. p. 211. t. 96. f. 1. optimè, monente cl. L. Bontou, hæc pertinet, sed folia non ternata; lobbi laterales in icone design. sunt foliola novella ex axillaria! (v. s.)

53. *V. abbreviata*, radice suffruticulosa, caule à basi ramosissimo subdiffuso, foliis lanceolatis utrinquè attenuatis suberratis utrinquè ramisque cinereo-pubescentibus, corymbis oligocephalis, capitulis pedicellatis, invol. squamis lanceolato-linearibus acuminatis dorso puberis vix subtrinerviis. ? in Indiâ orient. *Conyza abbreviata* Wall! cat. n. 3048. comp. 158. nec fortè ab eâ diversæ *Conyza cinerea* B. Wall. n. 3008, et imò *Conyza incana* Wall! cat. n. 3046. Species *V. cinerea* affinis, sed multò brevior, basi suffruticulosa et ut videtur perennis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
54. *V. vialis*, caule herbaceo erecto tereti ramoso minutissimè puberulo, foliis lineari-lanceolatis utrinquè acuminatis breviter petiolatis, junioribus ramulisque villosis, adultis minutè pubescentibus margine denticulis remotis acutis instructis, corymbo laxo, capitulis longè pedicellatis, bracteis subulatis paucis sub capitulo, invol. squamis dorso puberulis acuminatis, æcheniis teretibus sparè hirtellis, pappi serie externâ brevissimâ. ① in insulâ Madagascar æcus vias et circa pagum Tamatave legit cl. Bojer. Flores rosei (v. s. comm. à cl. inv.)
55. *V. leptophylla* (DC. in Wight contrib. bot. ind. p. 2), caule herbaceo erecto ramoso tereti tenuiter pubescente, foliis oblongo-linearibus utrinquè acuminatis integerrimis tenuissimè pulverulento-scabris, paniculâ ramosissimâ laxâ divaricatâ nudâ, invol. squamis circ. 2-seriatis linearibus acuminatissimis vix puberulis. ① in Indiâ orient. *Conyza linifolia* Wall! cat. n. 2063. comp. 173. non Linn. *Cacalia linifolia* Heyn! herb. Diversa videtur à Vern. linifoliâ sed non malè refert iconem Rumph. 6. t. 14. f. 2. et idèd esset Vern. fasciculata Blum. bijdr. p. 893. in adn. non Michx. Achn. villosiusculum teres (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et in h. Wight.)
56. *V. erigeroides*, glabra, caule diffuso ramoso striato, foliis lineari-lanceolatis obtusis submucronatis integris in petiolum attenuatis margine angustissimè subrevolutis, corymbo laxo subpaniculato, capitulis longè pedicellatis, invol. squamis mucrone acuminatis glabris disco brevioribus, æcheniis glaberrimis. ①? in insulâ Timor et in Novâ Hollandiâ leger. collectores circumnav. Bandinanz. *V. diffusa* Decaisn. non v. ann. mus. Par. 3 (1834) p. 408. non Less. (1829) *Cyanopsis*? *erigeroides* DC. in Wight. contr. bot. ind. p. 7. var. *Timorensis*. Folia 15-18 lin. longa, 2  $\frac{1}{2}$  lin. lata. (v. s.) Pappo ext. brevissimo ambit inter Vernoniâ et Cyanopsidem.
57. *V. laxiflora* (Less. in Linnæ 1831. p. 646), glabriuscula, caule erecto à basi ramoso, foliis infimis subrusulatis petiolatis obovatis obtusis dentatis, caulinis lanceolatis linearibusve distantibus, corymbo laxo, capitulis pedicellatis, invol. squamis vix pilosiusculis acuminatis subtrinerviis. ①? in Indiâ orient. ins. Timor (Gaudich.) et peninsulâ Malaicâ (Wight.). *Conyza helldifolia* Wall! cat. n. 3047 comp. 157. Vix ab eâ differt *Conyza elegantula* Wall! cat. n. 3049 comp. 159. Invol. subrubens. Valdè accedit ad *V. cineream*. (v. s. comm. ab hon. cur. merc. angl. Ind. or.)
58. *V. conyzoides* (DC. in Wight. bot. ind. 2. p. 6), caule herbaceo erecto striato brevissimè pubescente, foliis ovato- aut oblongo-lanceolatis acuminatis in petiolum brevem attenuatis serratis suprâ glabriusculis, subtis pubescenti-villosis, corymbo composito ramoso polycephalo, invol. squamis lanceolato-linearibus pubescenti-villosis acuminatis disco brevioribus. — in Indiâ orient. prope Negapatam detexit cl. Wight. *Conyza ovata* Wall! cat. n. 3097. comp. 207. non Less. *Cineraria glaberrima* Spreng. in litt. ad Wall. Affinis *V. cinereæ*. Fol. inf. ignota, sup. 2. poll. longa, pollicem lata. (v. s. in herb. Wight et Wall.)
- β. *Leschenaultii*, foliis angustioribus lanceolatis minus serratis. — in montibus Neelgerry legit cl. Leschenault. Vernac. dicitur *Kodo*. (v. s. comm. ab am. Leschenault.)

59. *V. ALBICANS* (DC. in Wight. bot. ind. 2. p. 6), tota tomento adpresso sericeo villosa canescens, foliis ovato-rhombéis basi cuneatis integerrimis, superne grosse dentatis acuminatis, corymbo subpaniculato nudo, invol. squamis linearibus villosis acuminatis, achenio villosa, pappo ext. quadruplò quàm int. brevior. — in Indiâ orient. verisim. agro Maderaspetano. *Conyza* seu *Chrysocoma* ap. herb. Madr. *Conyza albicans* Wall! cat. n. 3072. comp. n. 182. Folia 15. lin. longa, 7-8 lata. Cor. lobi hispidi. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
60. *V. ? RUGOSA* (Blum! bijdr. ned. ind. p. 894), herbacea, ramis divaricatis pubescentibus, foliis oblongo-lanceolatis basi attenuatis apice cuspidatis remotè denticulatis suprâ glabresculis, infrâ rugosis et puberulis, corymbo terminali dichotomo divaricato laxifloro, squamis involucri acutiusculis. — in ins. Javâ ad ortum fl. Tiburrum in montanis Gede legit cl. Blum. Folia 3-4 poll. longa, 8-12 lin. lata majora subrugosa. Capitula parva. Pappus albus, ext. fortè nullus. (v. s. spec. incompl.)
61. *V. ? CYMOsa* (Blum! bijdr. ned. ind. p. 894), herbacea, ramis paniculatis pubescentibus, foliis lineari-lanceolatis remotè serrulatis scalaris subtus puberis, corymbo terminali dichotomo, pedicellis approximatis subumbellatis, squamis involucri acutiusculis. — in montanis ins. Javæ. Pedunculis compressi. Capitula parva. Folia 4-poll. longa, 6-8 lin. lata. Pappum ext. in specimine equidem maneo non vidî et hîc ex auct. cl. Blume et affin. cum *V. divergente* refero. (v. s. specim. incompl.)
62. *V. EUPATORIODES* (Blum! bijdr. ned. ind. p. 894), suffruticosa, ramis villosis foliis oblongo-lanceolatis argutè serratis rugosis suprâ sparsi-setosa-scabris, subtus villosa hirsutis, corymbo terminali divaricato, squamis involucri ext. brevibus acuminatis, intimis longioribus vix acutiusculis. — in ins. Javâ ad cacumen montis Borangrang. Pappus albus ext. brevissimus vix ullus. (v. s. fruct.)
63. *V. DIVERSIFOLIA* (Bojer! in litt. 1835), caule herbaceo erecto tereti striatulo pubero paniculato-ramoso, foliis infer. elongatis caulinis ellipticis oblongisve integris aut sinuatis suprâ scabridis, subtus puberulo hirsutulis, paniculâ laxâ apice corymbosâ, invol. squamis ext. acutis, intimis lineari-oblongis apice subobtusis subcoloratis, achenio villosa, pappo exter. brevissimo. (1) in fossis circa agros cultos prov. Emirnesia. ins. Madagascar legit cl. Bojer. Folia infer. 15 lin. longa, latitudinè varia, caulina 6-9 lin. longa, omnia obscure viridia. Cor. purpureæ. (v. s. comm. à cl. inv.)

Sect. VIII. LEPIDAPLOA. — Lepidaploæ et Vernoniæ sp. *Cass.* — Vernoniæ propriæ pleræquæ *Less.*

Capitula plæquæ 4-flora, in paniculas dichotomas seu fasciculos seu cymas scorpioideas disposita, rariùs et verisim. abortu subsolitaria. Invol. squamæ apice non appendiculata.

- §. 1. *Involucri squamis obtusis acutisve nec acuminatis, capitulis nequè solitariis nequè in cymas scorpioideas dispositis.* *Less. in linnæ 1831. p. 635. — Species Americanæ, Asiaticæ, rariùs Africanæ.*
64. *V. ZEYLANICA* (*Less. in linnæ 1829. p. 344. 1831. p. 636*), suffruticosa, caule tereti tomentoso, foliis petiolatis membranaceis acutis crenatis suprâ glabresculis, subtus densè lanato-tomentosis, aliis ovatis basi cuneatis aliis panduratis, basi auriculato-cordatis, cymis subcorymbosis terminalibus, ramis elongatis apice ramosis, invol. ovatis squamis ovatis siccis vix acutis, achenio glabresculo, pappi serie ext. angustâ paleacèa brevi. § in Zeylonâ (Burm. Lesch!) in Madagascar (Bojer). *Calalia* Burm. *Zeyl.* 52. t. 21. *Eupatorium Zeylanicum* Lin. spec. 1172. Pappus pallidè rufescens. Cor. glabræ. *Vernonia* Wight! 416 sine loco design. (v. s. comm. à cl. Lambert et Leschenault.)

65. *V. APPENDICULATA* (Less. in linnæa 1831. p. 636), caule fruticoso ramosissimo, foliis infer. maximis basi cordatis inæqualiter serratis, caulinis lanceolatis petiolatis ad basin petioli auriculatis serrulatis supra glabris, subtus cum petiolis et ramis tomentosis, summis sessilibus sæpe cordato-auriculatis, cymbo fastigiato polycephalo, capitulis 10-floris oblongis, invol. squamis imbricatis obtusis glaberrimis. ♀ in Madagascar circa urbem Tananarivon, legit. cl. Bojer! Vernaculè dicitur *Ampiati* seu *Ampiadi*. Vern. macrophylla Bojer! in litt. 1833. Folia infer. 10 poll. longa margine plicata. Flores lilacini. Achen. 3. lin. longa, subangulata sparsa-villosa. Pappus albus pluriserialis, serie ext. breviori. Variat foliis omnibus aut plerisque exaristis. (v. s. comm. à cl. Bojer.)
66. *V. PANDURATA* (Link. enum. h. her. 2. p. 276), herbacea, ramis teretibus subvillosis, foliis sessilibus amplexicaulibus ovatis obtusis apiculatis basi auriculato-cordatis crenatis supra pilosis, subtus villosis, cymæ pluriès dichotomæ ramis 1-cephalis, capitulis multifloris campanulatis, invol. squamis siccis glabris 3-nerviis glandulosis ex rotundis oblongis, achenio sparè piloso multicoato, pappi serie externâ minimâ. ♀ ad fluvium Congo, ex Less. in linnæa 1831. p. 637. Less. in linnæa 1829. p. 261. Cor. glabræ violacæ. An *V. pandurata* Jacq. in Spreng. syst. 3. p. 436. eadem?
67. *V. HYPOLEUCA*, caule fruticoso tereti erecto cum petiolis et paginâ fol. superiore araneoso-tomentoso, foliis oppositis petiolatis ovato-lanceolatis triplinerviis subtus cano-tomentosis trilobis, lobis acuminatis lanceolatis, supremis indivisis, paniculæ ramis ramulisque oppositis apice laxè corymbosis, capitulis 10-floris, invol. squamis 3-seriatis ovali-lanceolatis acutis substriatis, pappo biseriali ext. brevi latè paleaceo, int. multisetoso scabro. ♀ in Mexico legit cl. Haenke. Cor. nec genitalia vidi. Species distinctissima. Petioli 8-10 lin. longi. Limb. fol. 2. poll. longus, sesquipollicem latus, pappus rufo-croceus, (v. s. in h. Haenke à cl. Sternb. comm.)
68. *V. NUDICAULIS* (Less. in linnæa 1831. p. 637), caule herbaceo simplici imâ basi tantum foliato pilis rigidis capitatis divergentibus villosa, ramis 1-cephalis, foliis sessilibus rosulatis obovatis rigidis subintegerrimis supra pilosis, subtus sericeo-lanatis, invol. campanulati squamis 3-seriatis linearibus acutis extus pilos dimorphos gerentibus, cor. glabris, achenio hirsuto-villoso, pappi plumoso serrato, seriebus confluentibus. ♂? in Madagascar. Habitus *V. Sellowii* oligocephalæ.
69. *V. NUDIFLORA* (Less! in linnæa 1829. p. 258), basi suffruticosa, caulibus è rhizomate plurimis erectis striatis apice subvillosis, foliis sessilibus longè linearibus 1-nerviis angustissimis marginè revolutis supra scabridis, subtus adpressè villosis, corymbi subfastigiati ramis simplicibus bifidis vix folia superantibus, capitulis 15-16-floris ebracteatis pedicellatis, invol. turbinato-campanulati squamis ovatis subsentis glabris sæpius ciliatis, achenio oblongo piloso, pappi pluriserialis setis piliformibus ext. paucis brevibus cæt. inter. subæquantibus. ♀ in Brasiliâ meridionali. Cor. glabræ violacæ. Pappus albus. Folia 3-poll. longa, lineam lata. (v. s. comm. à Mus. reg. Berol.)
70. *V. SQUARROSA* (Less! in linnæa 1831. p. 678. non p. 627), caulibus erectis teretiusculis basi suffruticulosus subvelutinis, foliis sessilibus elongato-linearibus margine subrevolutis, supra glabris rugosis, subtus albo-tomentosis, cymis scorpioides summis subverticillatis, invol. campanulati squamis oblongis obtusis ciliatis apice squarrosis intimis subscariosis, acheniis multicoatis hirsutis, pappi serie ext. brevi paleacæ. ♀ in Brasiliâ. Vern. rubricaulis, squarrosa et plantaginoides Less! in linnæa 1829. p. 300. Cor. purpureo-violacæ. Differt à rubricauli, invol. squamis squarrosis nec adpressis.
- $\alpha$ , *angustifolia* (Less. ll. cc.), foliis confertis angustè linearibus uninerviis margine revolutis.
- $\beta$ , *plantaginea* (Less. ll. cc.), foliis rarioribus flexuosis brevioribus pen-ninerviis margine planis. (v. s. comm. à Mus. reg. Berol.)
71. *V. INTERMEDIA*, caule fruticuloso erecto tereti glabro, ramulis floridis pubescentibus foliosis 1-cephalis axillaribus racemoso-corymbosis, foliis sessilibus au-

gnatè sublinearihus subacutis integerrimis margine revolutis suprà glabris, subtùs albidò-tomentosis nervo medio glabro, capitulis 30-35-floris, invol. squamis lanceolatis cillatis dorso subvillosis mucrone brevi acuminatis, cor. glabrà, achenio angulato parè villosò, pappi ser. ext. paleaceà brevi  $\frac{1}{2}$   $\frac{1}{2}$  in prov. Rio-Grande Brasiliæ. Media inter V. nudicaulem et psilophyllam. Cor. purpureæ. Capitula subbracteata. (v. s. in h. Mus. reg. Par. et Mus. imp. Bras.)

$\beta$ , *ramosior*, ramis floridis valdè elongatis scorpioideis sæpè bifidis, capitulis lateralibus subsessilibus aut breviter pedicellatis secus ramos 5-6 distantibus sæpius extrà-axillaribus, invol. squamis paulò magis acuminatis.  $\frac{1}{2}$   $\frac{1}{2}$  in prov. Rio-Grande Brasiliæ. Folia 3-6 poll. longa, lineam lata. Videtur eadem ac var.  $\alpha$  in solo uberiore lecta. (v. s. in Mus. reg. Paris. ex Mus. imp. Bras.)

72. V. *PSILOPHYLLA*, caule suffruticoso tereti erecto glabro tenni apice summo pubescente, foliis linearibus sessilibus acutis integerrimis 1-nerviis margine revolutis suprà glabris, subtùs albidò pubescentibus, ramis foliosis apice pedunculatos 3-5 subcorymbosos 1-cephalos subnudos gerentibus, pedicellis axillaribus, capitulis axillaribus 15-20-floris ebracteatis, invol. squamis ovali-lanceolatis glabrisculis ext. acuminatis int. mucronatis, corollà glabrà, achenio puberulo, pappi ser. ext. paleaceà brevi acutà.  $\frac{1}{2}$  in prov. Rio-Grande Brasiliæ. Affinis habitu et inflorescentiâ V. nudifloræ, sed pappo faciliè diversa. Folia 3 poll. longa, semilineam lata. Cor. purpureæ. Pappus griseus. (v. s. in h. reg. Par. ex Mus. imp. Bras.)

$\beta$ ? *Paulina* (Mart! herb.), caulibus è collo plurimis, foliis subtùs vix pubescentibus, cymæ extra axillaribus oligocephalis subscorpioideis pedunculatis campanulatis invol. campanulatis squamis oblongis dorso vix pubescentibus ext. subacutis, int. obtusis, cor. glabrà, achenio sericeo-villoso.  $\frac{1}{2}$   $\frac{1}{2}$  in editis campis prov. S.-Pauli Brasiliæ legit cl. Martins. Herba pedalis. Pappus griseo-nigricans. (v. s. in h. Acad. reg. Monac.) An species propria? sed specimen mancum.

$\gamma$ ? *leiolepis*, corymbi polycephali pediculis inferioribus parè ramosis, capitulis 10-15-floris, invol. squamis extimis secus pedicellum minimis acutis cæteris ovalibus obtusis lævigatis glabris purpureisculis, pappo albidore.  $\frac{1}{2}$   $\frac{1}{2}$  in Rio-Grande. Habitus var.  $\alpha$ ; invol. squamæ obtusæ ut in  $\beta$ , valdè lævigatæ nec ut in  $\alpha$  et  $\beta$  subpubescentes. (v. s. in h. Mus. reg. Par. ex Mus. imp. Bras.)

73. V. *LINEARIFOLIA*, basi suffruticulosa erecta glaberrima, caule subangulato apice ramoso, foliis linearibus utrinquè attenuatis subsessilibus rigidis penni-nerviis subtùs glandulis elevatis punctatis, capitulis solitariis ramulos oppositifolios nudos aut ramos foliosos axillares terminantibus ovatis 25-floris, invol. squamis ext. ovato-oblongis obtusis adpressis, achenio sericeo-villoso, pappi serie externà vix paleaceà interiori triplò ferè breviorè.  $\frac{1}{2}$   $\frac{1}{2}$  in Brasiliâ. Herba pedalis. Folia  $1\frac{1}{2}$  poll. longa, 2 lin. lata margine revoluta (an per siccationem tantum?). Cor. glabræ verisim. purpureæ. Pappus junior albus dein è griseo sordidè rufus. (v. s.)

74. V? *REVOLUTA* (Ham! tr. linn. soc. 14. p. 218), caule gracili ramoso striato, foliis linearibus subintegerrimis mucronulatis margine revolutis un-nerviis subtùs glandulosis utrinquè ramulisque pubescentibus, capitulis terminalibus subsolitariis, invol. squamis linearibus acuminatis, exterioribus dorso carinatis, achenio villosò, pappi pilis aequalibus.  $\frac{1}{2}$  in Indiâ orient. ad Sunnysaika et Monsurguny, ex Wall. cat. 3070. comp. 180 et in Camrupa borealis et Coasæ graminosis ex Hamilton. Folia pollicem longa, lineam lata, integerrima aut dente 1-2 medio notata. Pappus tenuis argenteus. Cor. purpureæ. (v. s. comm. ab hon. cœtu merc. angl. Ind. or.)

75. V. *LUCIDA* (Less. in linnæa 1829. p. 260. 1831. p. 640), suffruticulosa glaberrima, ramis virgatis angulatis striatis, foliis sessilibus oblongis basi obtusis muc acutis serratis glandulosis lucidis, cymâ paniculiformi, capitulis longè

pedunculatis subcylindraceis 10-12-floris. invol. squamis ovatis obtusissimis 3-nerviis, achenio oblongo piloso pappi seriebus forè longitudine aequalibus setiformibus.  $\S$  in Brasiliâ meridionali. Cor. violaceæ glabræ. Pappus argenteus. Capitula parva. (v. s. comm. à Mus. reg. Berol.)

76. *V. STYDULA* (Less. in Linnæâ 1829. p. 260. 1831. p. 649), caule fruticoso tereti ramosissimo glabro, ramulis angulatis, foliis sessilibus coriaceis oblongis univariis penninerviis margine subrevolutis integris aut denticulatis utrinque glabris glandulosis, cymis ad apices ramorum subpaniculatis, capitulis pedicellatis 7-8-floris, invol. turbinati foliolis oblongis subacutis adpressis, achenio piloso multicostrato, pappi serie ext. brevi paleacea.  $\S$  in Brasiliæ campis ad Rio das Pedras, in prov. S<sup>ta</sup>. Pauli, Rio-Grande, etc. Baccharia Montevidensis Spreng. syst. 3. p. 460. Pappus rufescens. Cor. violacea glabra. Involucra piloscula seu glabra. Caulis albescens. (v. s. comm. à Mus. reg. Ber.)

77. *V. ? ARGUTIFOLIA*, caule fruticuloso? tereti striatulo foliisque pubè minimâ subpolverulento-velutinis, foliis petiolatis basi cuneatis ceterum ovatis acuminatis grossè inciso-dentatis basi 3-nerviis, nervis lateralibus bifidis, paniculâ terminali compositâ foliis intermixtâ, invol. squamis ovatis subacutis margine submembranaceis, capitulis 6-7-floris.  $\S$  ? in Africâ austr. orient. prope Omsamwubo legit cl. Drege, n. 5033! Flores in specim. varium, masculi, ovario abortivo, stylo simplici à medio ad apicem hispidi. An Webbii spec. ? ad habitus repugnat. Pappi setæ crassæ, exter. breviores. (v. s.)

78. *V. MESPILOFOLIA* (Less. in Linnæâ 1831, p. 641), fruticosa, ramis glabrioribus teretibus, foliis petiolatis longè obovato-cuneatis apice grossè 3-7-dentatis minutissimè (sub lente) puberulis, corymbo subnudo polycephalo, pedunculis pubescentibus, invol. squamis 7-8-subtriseriis obtusissimis dorso glabris margine ciliatis, floribus in cap. 4-5, pappi serie ext. brevi vix paleacè, int. multisetâ subpliseriis.  $\S$  ad Cap. Bonæ Spei, in Uitenhagen (Burchell! n. 4500. Eckl! h. 1599. 1830. 1764. 1933), Cafferland, Swellendam, territorio cesso (Eckl! n. 1137. 199. 1375. 794), ad Omsamculo et Omtata. (Dreg! n. 5035.) Eupatorium Capense Spreng. fil. suppl. p. 22. Zeyh! fl. exs. n. 201. ex h. Moric. Achenium oblongum villosum. (v. s.)

*f. subcanescent*, caule foliisque utrinque pubè brevi molli subcanescentibus.  $\S$  in Cafferland Dreg! n. 3687. (v. s.)

79. *V. CORYMBOSEA* (Less. in Linnæâ p. 647), fruticosa, ramis striatis pubescentibus, foliis vix petiolatis oblongo-cuneatis apice paucidentatis subtus tomentosis, suprà glaberrimis, capitulis corymbosis, invol. squamis 12-15 obtusissimis dorso tomentosis, floribus in capit. 5, pappi serie ext. paucisetâ, inter. pliseriis.  $\S$  ad Cap. Bonæ Spei. in coloniæ distr. oriental. (Burchell! n. 4343.) in collibus distr. Uitenhagen (Eckl! n. 144. 759. 400. 16. 1108. 1917. 1575.) ad Van Staedens rivier. (Dreg! n. 2118.) Stæbelina corymbosa Thumb. fl. cap. 628. ex Vahl! in h. Burm. Eupatorium Burm! herb. Capitula ad apices ramulorum 3-4 subsessilia subumbellata. Folia pollicem longa, 4-lin. lata. Pedunculi compressi folio florali ferè breviores. Achenium oblongum villosum. Pappus rufescens, involucri duplò longior. (v. s.)

*f. mespiloides*, foliis majoribus acutè 7-9-dentatis subtus adpressè sericeo-canis.  $\S$  in Uitenhagen legit cl. Ecklon. n. 289. et ad Omtanda cl. Drege, n. 5036! Folia formâ et magnitudine *V. mespilifoliam* referentia, sed subius canescentia. Capitula *V. corymbosæ*. An fortè utrinque hybrida? (v. s.)

80. *V. ASTERIFLORA* (Mart! herb.), fruticosa, ramis teretibus subtiliter striatis tomentoso-incanis, foliis sessilibus elliptico-oblongis utrinque vix acutis integerrimis suprà adultis glabris, subtus tomentoso-canis penninerviis, capitulis 5-7 longè pedunculatis corymbosis ebracteatis circ. 40-floris, invol. squamis floribus multò brevioribus, ext. oblongis dorso hirsutis, intimis linearibus acuminatis glaberrimis, cor. glabrâ, achenio parè villosi, pappi ser. ext. paleacè brevissimâ.  $\S$  in editis campis prov. Minarum gener. Brasiliæ. Folia 3 poll. longa, pollicem lata, pedunculi 1½-2 poll. longi, nudi. Cor. purpureæ 5 lin. longæ, pappus sordidus. (v. s. in h. Acad. reg. Monac.)

81. *V. ARIÆFOLIA* (Mart! herb.), caule suffruticoso simplici erecto tereti adpressè sericeo-tomentoso, foliis sessilibus subrotundis obtusis subintegerrimis penninerviis rigidis suprâ vix araneosis, subtus adpressè sericeo-tomentosis, pedunculis 4-5 corymbosis folio paulò longioribus: cephalis sub capitulo 1-phyllis, capitulis circ. 70-floris, invol. campanulati squamis multiserialibus coriaceis ext. ovatis subacutis dorso araneosis intimis ferè linearibus acuminatis glabris, cor. glabrâ, achænio sericeo-villoso, pappi serie ext. angustè paleacè acuminatâ, inter. triplò breviorè.  $\zeta$  in editis campis prov. Minasum gener. Brasiliz. Caulis sesquipedalis. Folia 2 poll. longa, sesquipoll. lata. Cor. purpureæ. Pappus stramineus. Affinis videtur *V. argyrophyllæ*. (v. s. in h. Acad. reg. Monac.)
82. *V. PENTACANTHA*, caule fruticoso tereti supernè albo-lanuginoso, foliis petiolatis oblongis basi longè cuneatis apice obtusis integerrimis rigidis utrinquè reticulatis penninerviis suprâ glabrisculis, subtus araneoso-tomentosis, capitulis ad apicem ramorum subternis sessilibus aut brevè pedicellatis ebracteatis ovato-subglobosis 50-floris, inv. squamis coriaceis multiserialibus imbricatis, ext. ovatis mucrone brevi squarrosos, intimis longioribus acuminatis subscariosis, cor. glabrâ, achænio villosos, pappi ser. ext. paleacè acuminatâ int. triplò breviorè.  $\zeta$  Patr. ign. Valdè accedit ad hexacantham. Folia 3 poll. longa pollicem lata, petiolo 4-5 lin. Pappus subrufescens. (v. s. in herb. Acad. reg. Monac.)
83. *V. LAURIFOLIA*, caule fruticoso ramoso tereti ferrugineo-tomentoso, foliis brevissimè petiolulatis ovali-lanceolatis basi subobtusis apice acuminatis integerrimis rigidis penninerviis utrinquè nervis elevatis reticulatis suprâ glaberrimis, subtus secus nervum puberis, capitulis longè pedicellaribus axillaribus solitariis aut ad apicem laxè cymosis multifloris, invol. ovatis squamis rigidis glabrisculis lanceolatis acuminatis, cor...., achænio sericeo-villoso, pappi ser. ext. brevè paleacè.  $\zeta$  Patr. ign. Folia 2  $\frac{1}{2}$ -3 poll. longa, pollicem lata. Pedicelli circ. pollicares. Pappus rufo-griseus rigidulus. Recept. alveolatum. Flores in invol. circ. 25 sed in specim. vix ritè observandi. (v. s. in h. Acad. reg. Monac.)
84. *V. TOLUCCANA*, fruticosa glabra, ramis teretibus foliisque oppositis petiolatis elliptico-oblongis integerrimis nitidis viscosis, pedunculis axillaribus oppositis corymbum subpaniculatum confertum gerentibus, capitulis pedicellatis circ. 20-floris, invol. squamis subacutis biserialibus disco dimidio brevioribus, achæniis angulatis pilosiusculis, pappi serie ext. brevè paucisetâ.  $\zeta$  in Mexico circa Toluccam legit cl. Audrieux, pl. exs. n. 273! Habitus Baccharidis seu Psadiæ sed stigmata elongata hispida Vernoniæ. Fol.  $\frac{1}{2}$ -2 poll. longa, 6 lin. lata. Cor. in sicco pallida. Achænia nigrescentia. Pappus albus. (v. s.)
85. *V. DOMINGENSIS*, fruticosa glabrescens, ramulis teretiusculis apice angulatis, foliis breviter petiolatis ovalibus obovatisve obtusis hinc indè dentatis penninerviis coriaceis suprâ lucidis subtus punctulatis, capitulis paucis subcorymbosis pedicellatis 10-floris, bracteis squamiformibus multiserialibus brevibus obtusis imbricatis sub invol., involucre squamis veris longioribus siccis vix acutis dorso ad apicem subpuberis, achænio 10-costato glabro, pappi ser. ext. paleacè brevè.  $\zeta$  in Sancto Domingo. *Lepidaplos buxifolia* Cass. dict. 26 (1823) p. 18 et idè Vern. *buxifolia* Less! in Linnæâ 1829. p. 313. *Promelia Domingensis* Spreng! in herb. Balb. 1821. syst. 3. (1826) p. 502. Folia summa 6-7 lin. longa 4 lin. lata. Recept. nudum. Nomen Cass. etiam vetustatis non admisi ne confundatur cum *V. buxoidi*. (v. s. ex itin. Berter.)
86. *V. ZANZIBARENSIS* (Less. in Linnæâ 1831. p. 638), caule fruticoso, ramis villosis, foliis ovalibus acutis subdentatis suprâ sparsè villosis subtus petiolisque densè cano-tomentosis, corymbo laxo dichotomo subpaniculato, capitulis 12-floris, invol. squamis lanceolatis acuminatis dorso puberis, corollæ lobis intus ad apicem longè barbatis, achænio glabro, pappo biseriali ext. paleaceo brevè.  $\zeta$  in insulâ Zanzibar locis à mare parùm dimittit cl. Bojer. Pappus intense fulvus barbellatus rufescens. (v. s. contin. à cl. Bojer.)
87. *V. SOLIDAGINIFOLIA* (Bojer! in litt. 1835), fruticosa, ramis erectis apice angulatis pubero-scarbidis, foliis subsessilibus lanceolatis acutis serratis suprâ

glabris, subtus glaucis secus nervos pubescentibus, cymâ corymbosâ, pedicellis valdè bracteolatis, invol. squamis exter. brevibus ferè subulatis, inter. linearibus acutis, acheniis teretiusculis striatis confertè puberulis, pappi serie ext. setosâ brevi.  $\frac{3}{4}$  in insulâ Zanzibar legit cl. Bojer. Frutex 8-10-pedalis. Folia omninò Solidaginis Canadensis. Flores flavi (Bojer). Folia super.  $1\frac{1}{2}$  poll. longa, 4-5 lin. lata. Fl. in invol. circ. 12. Pappus pallidè cinnamomene. (v. s. comm. à cl. Bojer.)

88. *V. SUAVEOLENS* (H. B. et Kunth nov. gen. 4 p. 30), fruticosa volubilis, ramis teretibus obscurè tomentosis, foliis breviter petiolatis oblongis penninerviis coriaceis subcrenulatis suprâ subpilosius demùm glabra, subtus petiolisque obscurè tomentosis, ramis floridis subverticillatis, capitulis 7-floris, invol. squamis 1-nerviis ex ovato-acutis ciliatis, achenio glaberrimo, pappi serie externâ brevi caducâ.  $\frac{5}{8}$  in Novâ Granat. prov. Ayavaca et flum. Cutaco, Cor. albæ glabræ. Pappus ex albo flavescens. Less. in linnæâ 1829. p. 265. 1831. p. 641.
89. *V. PECTINATA*, fruticosa, ramis teretibus glabriusculis, foliis brevè petiolatis ovatis pectinato-acutè et elongato-serratis membranaceis utrinquè pilosiusculis, cymâ terminali corymbosâ nudâ, capitulis longè pedunculatis multifloris ovato-cylindraceis, inv. squamis siccis glabriusculis ex ovato-oblongis obtusissimis 3-nerviis, achenio glabro, pappi serie externâ brevissimâ.  $\frac{3}{4}$  in Indiæ or. montibus Nilghirry legit cl. Noton. Eupatorium pectinatum Wall! cat. n. 3172. comp. 282. Species distinctissima nulli verè affinis. Cor. glabrâ. Pappus pallidè rufescens. Inv. squamæ apice obscuræ. Stylus et pappus Vernoniæ. (v. s. in herb. bon. cœtu merc. angl. Ind. or.)
90. *V. PECTINIFORMIS* (DC. in Wight, contrib. bot. ind. p. 2), fruticosa? ramis teretibus glabriusculis, junioribus angulatis pubescentibus, foliis breviter petiolatis lanceolatis acuminatis pectinato-acutè et elongatè serratis membranaceis suprâ glabris, subtus pilosiusculis, cymâ terminali corymbosâ nudâ, capitulis longè pedicellatis multifloris ovatis, inv. squamis siccis glabris ciliatis ovato-oblongis subacutis.  $\frac{3}{4}$   $\frac{1}{2}$  in Indiæ orient. montibus Dendignl dictis legit cl. Wight. Valdè affinis *V. pectinatæ* et forsân mera varietas. Præter car. indicatos differt pappo nec rufo, involacro evidentè longiore nec subæquali. Folia 3 pollices longa, vix pollicem lata. (v. s. in h. Wight.)
91. *V. ASPERA* (Ham. trans. lin. soc. 14. p. 219. non Less.), caule herbaceo? ramoso tereti subvillosa, foliis elliptico-oblongis subpetiolulatis basi attenuatis apice acuminatis serratis suprâ glanduloso scabris, subtus reticulatis secus nervos rufo-villosis et inter eos glanduloso-impressis, paniculæ subcorymbosæ ramis inferioribus folio suo brevioribus, capitulis 30-floris, invol. squamis int. oblongis mucronatis, ext. minoribus valdè acuminatis, achenio oblongo-cylindrico angulato piloso.  $\frac{1}{2}$  in Indiâ orient. ad Kodoya in Anggæ montibus, Bengalo et fortè agro Burmanico legit cl. Wall. et Ham. Eupatorium asperum Roxb. calc. 61. Vern. aspera Ham. trans. lin. soc. 14. p. 219. Wall! cat. et herb. n. 2922. comp. 32. non Less. Vernonia Roxburgi Less. in linnæâ 1831. p. 674. Conyza platylepis Spreng. in litt. ad Wall. Eupatorium pyramidale Don. prod. fl. nep. 170? — Pluk. Alm. t. 395. f. 7. malè. (v. s.)
92. *V. MULTIFLORA* (Less. in linnæâ 1831. p. 642), caule herbaceo erecto ramoso tereti tenuissimè puberulo, foliis breviter petiolatis lanceolatis utrinquè acuminatis à medio serratis suprâ asperulis, subtus tenuiter pubescentibus, paniculâ subcorymbosâ polycéphalâ nudiusculâ, invol. squamis lanceolatis acutis pallidis glabriusculis, achenio glabro pappo subuniseriali.  $\frac{1}{2}$  in Indiâ orient. ad Promè et Panlang. legit cl. Wallich. Conyza multiflora Wall! cat. n. 3032. comp. 142. An Eupatorium acuminatum Don. prod. fl. nep. 170? et idèd Eupat. Nepalense Spreng. 3. p. 412? Conyza divergens E. Wall! n. 3007. et Eupatorium versicolor Wall! cat. n. 3167. Folia 3 poll. longa vix poll. lata. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
- β. latifolia*, foliis ovali-lanceolatis suprâ magis asperis.  $\frac{1}{2}$  in collibus Raymahal legit cl. Hamilton. *V. aspera* Ham! herb. ex Wall! cat. h. calc. Var. *β. affinis V. asperæ*, sed capitula dimidio minora ut in var. *δ*. (v. s. comm. cum priore.)



93. *V. NIGERANRANSIS*, herbacea? caule tereti subpuberulo, foliis brevè petiolatis ovatis acuminatis spinuloso- et acutè serratis enprà scabridis, subtus fulvo-subtomentosis, cymæ terminalis paniculæformis ramis ramosissimis polycephalis, capitulis ovatis confertis 2-5-floris, invol. squamis siccis ovali-oblongis acutis glabris apice pilosiusculis, achenio glabro, pappi serie externâ brevissimâ deciduâ. — in Indiâ orientali legit cl. Wight, et in montibus Nilgerry cl. Lechenault. *Eupatorium polyanthum* Wall! cat. 3171. comp. 281. *V. aspera* Less. in linnæa 1831. p. 643? Cor. glabra. Pappus albus. (v. s.)
94. *V. ACUMINATA*, caule fruticoso ramoso tereti, ramis junioribus villosovelutinis, foliis breviter petiolatis elliptico-obovatis longè cuneatis apice cuspidato-acuminatis subintegerrimis glabris nervis præcipuis subtus velutinis, panicula elongatæ nudæ ramis polycephalis, invol. squamis ovalibus obtusis dorso velutinis imbricatis. 3 in Indiâ orient. prov. Silhet, in montanis ad Pundnah legit cl. Wallich. *Conyza acuminata* Wall! cat. n. 3034. comp. 144. *Eupatorium acuminatum* Wall! herb. *Eupatorium cuspidatum* Spreng. in litt. ad Wall. Folia 4-6 poll. longa 1  $\frac{1}{2}$ -2 lata, petiolo 2-5 lin. Fl. inv. 7-9 ex sicco purpurascens. Pappus albus pauciseriali, setis ext. irregularibus brevibus facilè deciduis. Achen. glabrum basi attenuatum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
95. *V. PUNDNANA*, caule herbaceo? erecto tereti vix sub lente puberulo, foliis breviter petiolatis lanceolatis longissimis basi attenuatis integerrimis apice acuminatis grossè et rariter serratis glabris, nervo medio subtus tenuissimè puberulo, panicula elongatæ nudæ ramis apice confertè polycephalis, invol. squamis oblongis obtusis, intimis longioribus demùm stellato-patulis. 2½ in Indiâ orient. prov. Silhet ad Pundnah legit Fr. de Silva. *Conyza Pundnana* Wall! cat. n. 3035. comp. 145. Folia 8 poll. longa, 2. poll. lata. Fl. in invol. 8-9. Recept. angustam. Achenia glabra basi attenuata. Pappus albus pauciserialis setis ext. brevibus irregularibus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
96. *V. VAGANS*, caule herbaceo tereti ramoso sarmentoso-acandente brevissimè puberulo, foliis breviter petiolatis ovato-lanceolatis acuminatis integerrimis glabris, ramis floridis axillaribus folio longioribus foliolosis polycephalis in paniculam amplam dispositis, invol. squamis lanceolatis acutis ciliolatis imbricatis. 2½ in Indiâ orient. prov. Silhet legit cl. Wallich. *Conyza vagans* Wall! cat. n. 3040. comp. 150. *Eupatorium vagans* Wall! herb. Flores in invol. 7-8 purpurei glabri. Pappus rufus, setis ext. paucis linearibus paleaceis ciliatis, cæteris pluriserialibus scabris. Achen. glabrum teretiusculum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
97. *V. SCANDENS*, caule scandente tereti, ramis junioribus velutinis adultis glabris, foliis breviter petiolatis ovalibus acuminatis integerrimis utrinquè glaberrimis, ramis floridis axillaribus foliosis subpaniculatis, capitulis pedicellatis ovatis, invol. squamis ciliatis dorso glabris ext. ovatis acutis, int. oblongis. 3 2½ in Indiâ or. prov. Silhet legit cl. Wallich. *Conyza scandens* Wall! cat. et herb. n. 3060. comp. 170. Fl. atropurpurei in invol. 9-11. Achenium junius teretiusculum glabrum. Pappus subrufescens multisetus æqualis, setulis ext. vix ullis deciduis. An fortè à genere removenda? (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
98. *V. BLANDA*, caule terecto tereti paniculato ramoso apice pubescente, foliis breviter petiolatis ellipticis basi attenuatis apice cuspidatis integerrimis glabris, ramis axillaribus foliosis paniculatis, capitulis pedicellatis, invol. squamis imbricatis dorso puberulis oblongo-linearibus mucronatis ext. brevissimis, int. sensim longioribus. 3 2½ in Indiâ Birmanicâ propè Trogla ad Salween legit cl. Wallich. *Conyza blanda* Wall! cat. et herb. n. 3033. comp. 143. birm. n. 1926. Affinis *V. scandenti*. Achenium teres basi subattenuatum puberum. Pappus ext. vix ullus, inter. infuscens. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
99. *VOLKAMERIÆPOLIA*, fruticosa, caule tereti glabriusculo ad axillas villosæ, brevè petiolatis membranaceis amplissimis elliptico-obovatis acutis basi subciliatis enprà glabris, subtus pubescentibus, utrinquè minutè glabris, cymæ terminalis paniculæformis rachis angulatâ, ramis tomentosis pedicellatis, invol. squamis siccis extimis ovatis acutis villosis, intimis

- oblongis obtusis, achenio glanduloso, pappi serie externâ brevi.  $\frac{5}{3}$  in Nepaliâ detexit indef. Wallich. *Conyza volkameriifolia* Wall. ! cat. n. 3001. comp. 111. *Eupatorium volkameriifolium* Wall. ! herb. Folia 9-10 poll. longa, 3 poll. lata. Cor. glabra. Pappus albus. (v. s. comm. ab hon. cœtu merc. angl. Ind. or.)
100. *V. EXTENSA*, suffruticosa glabra, ramis teretibus, junioribus subangulatis subpubescentibus, foliis brevè petiolatis oblongis basi attenuatis apice acuminatis hinc indè serratis membranaceis, cymis terminalibus paniculiformibus ramosissimis aphyllis, capitulis pedicellatis 8-10-floris, involucri squamis oblongis obtusis, achenio pubescenti-glanduloso, pappi serie ext. interiori triplo brevior.  $\frac{5}{3}$  in Nepaliâ. *Conyza extensa* Wall. ! cat. n. 3016. comp. 126. *Eupatorium rufum* Spreng. in litt. ad Wall. Pappus pallidè aut intense rufus. Folia 6-7 poll. longa, 1-2 poll. lata. Habitus ferè *V. Senegalensis* sed pappi series valdè inæquales. (v. s. comm. ab ampl. cœtu merc. angl. Ind. or.)
101. *V. MADAGASCARIENSIS* (Less. in Linnæ 1831. p. 644), caule fruticoso scandente, ramis striato-angulatis superne adpressè pilosis, foliis petiolatis membranaceis oblongo-ovatis acuminatis basi acutis punctatis albo pilosculis, paniculis terminalibus dichotomis oligocephalis, capitulis pedicellatis multifloris, invol. campanulati squamis coriaceis glabris ciliatis ovatis, cor. et achenio glabris, pappi serie ext. setacea brevi.  $\frac{5}{3}$  in insulâ Madagascar prope Maron-Voad. Cor. in sicco flavida? Recept. nudum. Pappus flavescens. Folia 2-3 poll. longa, 12-15 lin. lata.
102. *V. ATTENUATA*, caule herbaceo erecto striato infrà hirsuto apice puberulo, foliis infrà originem ramorum floralium confertis ellipticis basi cuneatis apice acuminatis grossè dentatis reticulato-nervosis suprâ glaberrimis, subtus in nervis puberulis, super. secus ramos florales distantibus linearibus acutis, dentatis, dentibus omnium mucronatis, corymbo laxo oligocephalo, capitulis longè pedunculatis. — in India orient. ad Moalmyne legit cl. Wallich. *Conyza attenuata* Wall. ! cat. n. 3020. comp. n. 130. Invol. squamæ glabriusculæ, ext. breves cuspidatæ, int. longiores acutæ. Achen. teres villosum. Pappus ext. circ. lin. longus paucisetus. Folia 4 poll. longa, sesquipoll. lata. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
103. *V. LONGICAULIS*, herbacea, caule simplici tereti striato hirtello-pubescente, foliis brevissimè petiolatis oblongis ovalibusve basi cuneatis apice acuminatis serratis membranaceis glabris reticulatim venosis in medio arcolarum suprâ elevato-glandulosis, cymæ terminalis paniculæformis ramis multi-capitatis subfoliiferis, invol. ovatis squamis siccis ovatis pungenti-mucronatis, achenio glabro, pappi serie externâ brevissimâ.  $\frac{2}{3}$  in India orient. ad Pundua, Silhet, Nepaliâ legit cl. Wallich. *Eupatorium longicaule* Wall. cat. et herb. n. 2169. comp. 279, nom 3073. *Eupatorium acuminatum* Spreng. in litt. ad Wall. 1830. Variet. foliis latioribus et angustioribus. Habitu accedit ad *V. serratuloidem*. Cor. glabræ. Pappus argenteus. (v. s. comm. ab hon. cœtu merc. angl. Ind. or.)
104. *V. SALIGNA*, caule herbaceo erecto tereti striato minutè puberulo paniculato-ramoso, foliis oblongo-lanceolatis utrinquè acuminatis hinc indè serratis, suprâ asperiusculis, subtus subpubescentibus, paniculæ polycephalæ ramis parè foliosis elongatis 7-9-cephalis, invol. squamis submembranaceis oblongis acutis. — in India orient. prov. Silhet, legit. cl. Wallich. *Conyza saligna* Wall. cat. n. 3061, comp. 171. Folia 3-4 poll. longa, 10-12 lin. lata. Flores ex sicco vid. albi. Pappus albiissimus, ext. circ. lineam longus. (v. s. comm. ab ill. Car. merc. angl. Ind. or.)
105. *V. SERRATULOIDES* (H. B. et Kunth. nov. gen. 4. p. 26, t. 316), herbacea, caule simplici glabro tereti striato, foliis brevissimè petiolatis oblongo-lanceolatis acuminatis hinc indè serratis glabris, suprâ impresso-glandulosis, cymæ terminalis paniculæformis ramis apice 5-6-cephalis, capitulis 10-floris cylindraceis, invol. squamis ovatis pungenti-mucronatis adpressis ciliatis glabris, achenio velutino pubescente, pappi serie externâ vix paleacea brevissimâ.  $\frac{2}{3}$  in Mexico temperato circa Valladolid de Mechoacan (H. et B.) et Leon (Mendez ! ) Less. in Linnæ 1829. p. 268. 1831. p. 645. *Serratula semiserrata* Willd. herb. ex

Less. Herba 2-3 ped. habitu serratula tinctoria. Caulis et rami purpureo-erecti. Cor. purpurea glabræ. Pappus argenteus. (v. s. comm. à cl. Mendes et Alaman.)

106. *V. LIATROIDES*, caule herbaceo erecto tereti striato tenuissimè puberulo apice sulcato-angulato, foliis brevissimè petiolatis elliptico-oblongis basi acuminatis apice subacutis serratis penninerviis suprà glabris, subtus puberulis, ramis foliosis apice corymbiferis in paniculam dispositis, capitalis pedicellatis ebracteatis 10-12-floris, invol. squamis oblongis glabrisculis, ext. acuminatis, int. vix mucronatis, cor. glabrâ, recept. alveolato, achenio glanduloso subpubescente, pappi serie ext. brevissimâ angustè paleacea. 2' in Mexico inter Tula et Tampico (Berl.!) Valdè affinis *V. serratuloides* sed capitula dimidio minora. Folia 2 poll. longa, 7-8 lin. lata. Pappus albidus. Invol. pallida 2. lin. longa. Variat nerviis subtus paulò prominulis aut reticulatis elevatis. (v. s.)

107. *V. MUCRONULATA* (Less. in linnæâ 1829. p. 266. 1831. p. 647), fruticosa, ramis teretibus striatis ferrugineo-tomentosis demùm glabris, foliis brevissimè petiolatis coriaceis ellipticis subcordatis retosis mucronatis integerrimis suprà glabris impresso-glandulosis, subtus fulvo-tomentosis, cymis terminalibus, capitalis brevè pedicellatis, invol. squamis coriaceis 1-nerviis ovatis ciliatis, cor. glabræ laciniis glandulosis, achenio piloso multicostato, pappi serie externâ paleacea brevi. 5 in Brasiliâ tropicâ. Folia 11-15 lin. longa, 6 lin. lata. Pappus argenteus.

108. *V. ULIGACTROIDES* (Less. in linnæâ 1831. p. 648), caule fruticoso striato ramisque fulvo-tomentosis, foliis subsessilibus coriaceis oblongis basi obtusis (nec cordatis) penninerviis suprà lucidis scabrisculis, subtus fulvo-tomentosis, margine reflexis subserratis, panicula terminalis oblonge ramis foliis parvis stipatis, capitalis 12-floris basi subequamosis, invol. squamis siccis dorso fulvo-tomentosis, ext. subrotundis mucronatis, int. oblongis obtusis, cor. glabrâ, achenio hirsutiusculo, pappi ser. ext. paleacea brevi. 5 in Brasiliâ. Folia 2 ½ poll. longa, 1 ½ lata. Cor. purpureæ. Pappus albidus. Facies *Liabi sessiliflori*.

109. *V. CORDATA* (H. B. et Kunth nov. gen. 4. p. 31), fruticosa, ramis subangulatis obscure tomentosis, foliis petiolatis coriaceis ellipticis obtusis basi cordatis subdentulatis utrinquè obscure tomentosis, cymæ terminalis ramis dichotomis, capitalis multifloris, invol. squamis ovato-oblongis, platinerviis puberulis, achenio pluricostato, costis hispidis, pappi serie externâ brevi. 5 in montibus Mexicanis prope urbem Tasco. Less. in linnæâ 1829. p. 267. 1831. p. 648. *V. cordifolia* Willd. herb. ex Less. Pappus flavescens. Cor. ign.

110. *V. LEOCARPA*, caule fruticoso tereti apice velutino demùm glabro, foliis petiolatis ovato-lanceolatis integerrimis basi subobtusis apice acuminatis penninerviis suprà pubescentibus demùm glabris, subtus lanato-tomentosis, petiolo nervoque velutino subrufo, panicula terminalis composita ramis subcorymbiferis polycephalis, pedicellis aphyllis capitula 3-5 ovata congesta 5-flora gerentibus, invol. squamis ovalibus obtusis ciliatis sublanatis, cor. et achenio glabris, pappi ser. ext. serie paucis dimidiam int. longitud. æquantibus. 5 in Mexico legit cl. Karwinaki. Petioli 8-10 lin. longi. Limbus fol. 4 poll. longus, 1 ½ poll. latus. (v. s. in h. Acad. reg. Monac.)

§. 2. *Involucris squamis obtusis aut acutis nec acuminatis, capitulis in cymis scorpioides dispositis aut rarò solitariis, folia perfecta basi non gerentibus.* Less. in linnæâ 1831. p. 648. — *Species omnes Americanæ.*

111. *V. PERRUCINÆA* (Less. in linnæâ 1829. p. 271. 1831. p. 649), fruticosa, ramis subangulatis fulvo-tomentosis, foliis petiolatis coriaceis ellipticis obtusis integerrimis margine undulatis suprà scaberrimis puberulis, subtus fulvo-tomentosis, cymis scorpioides later. et termin., capitalis campanulatis sessilibus multifloris, invol. squamula coriaceis ciliatis apice pubescentibus obtusis mucronatis, achenio glanduloso costis hispidis, pappi serie ext. brevi. 5 in Brasiliâ tropicâ. Cor. alba? glabræ. Pappus flavescens seu ferrugineus.

112. *V. DISCOLOR* (Less! in *linnæâ* 1829. p. 274. 1831. p. 649), fruticosa, ramis teretibus apice angulatis cum pedunculis petiolis et facio fol. inferâ pube brevissimâ adpressâ densâ canis, foliis petiolatis oblongo-lanceolatis utrinquè acuminatis integerrimis suprà glabris, cymis scorpioideis divaricatis, capitulis secundis brevè pedicellatis nudis, invol. turbinati squamis ovatis acutis, achenio oblongo multicostato pilosiusculo, pappi serie ext. subpaleaceâ brevissimâ.  $\beta$  in *Brasiliâ tropicâ* (Sellow!) ad Cercovado prov. Rio-Janeiro (Vauth. n. 346). *Eupatorium discolor* Spreng. syst. 3. p. 412. Pappus substramineus. Cor. glabra purpurea. (v. s.)
113. *V. DIFFUSA* (Less. in *linnæâ* 1829. p. 272. 1831. p. 649), herbacea erecta ramosa, foliis longè petiolatis membranaceis angustè elliptico-lanceolatis integerrimis suprà puberulis, subtus ramisque obscurè tomentosis, cymæ scorpioideæ ramis divaricatis, capitulis pedicellatis secundis, invol. subturbinatis squamis coriaceis 1-nerviis acutis extus lanato-tomentosis, achenio piloso, pappi serie externâ paleaceâ brevi. — in *Brasiliâ tropicâ*. *Conyza divaricata* Spreng. syst. 3. p. 508. Pappus argenteus. Petioli 1-2 poll. longi. Limbus fol. 4-10 poll. longus,  $1\frac{1}{2}$ -5 lin. latus.
114. *V. OPPOSITIFOLIA* (Less. in *linnæâ* 1829. p. 273. 1831. p. 649), herbacea, ramis oppositis teretibus subtomentosis suprà folior. orig. subincrassatis, foliis oppositis petiolatis membranaceis ellipticis acuminatis utrinquè glandulas sessiles gerentibus, suprà pubescentibus, cymæ scorpioideæ terminalis ramis aphyllis divaricatis, capitulis multifloris pedicellatis, invol. turbinati squamis 1-nerviis ovatis acutis extus pilosis demum stellato-patentibus, achenio piloso multicostato, pappi serie ext. paleaceâ brevi.  $\beta$  in *Brasiliâ tropicâ*, circa Rio-Janeiro (Less.). *Conyza stellata* Spreng. neu. entd. 2. p. 141. syst. 3. p. 508. Cor. glabra purpurea. Pappus niveus.
115. *V. VERNONIACA*, herbacea ad collum fruticosa, caule erecto tereti apice nudo nudiquè pube brevi conferta hirtello, foliis caulinis alternis ramis aspe oppositis linear-oblongis obtusis scabulis subdenticulatis suprà scabridis, subtus hirsutis, corymbo paniculato aphylo, invol. squamis lanceolatis acutis glabris, achenio hirtello striatulo, pappi serie ext. brevi.  $\beta$  in campis editis prov. Sancti Pauli *Brasiliæ*, legit cl. Lund. Flor. novemb. — Herba 1-1  $\frac{1}{2}$  ped. Fiores albi. Fol. 2-3 poll. longa, 3-4 lin. lata. (v. s.)  
 $\beta$ . *lactor*, foliis oblongo-lanceolatis basi attenuatis magis dentatis, invol. subpuberulo.  $\beta$  Cum var. et legit cl. Lund. Fl. albi in inv. 10-12. Achenium longius forsan magis etate provectum? Folia 3 poll. longa, pollicem lata. (v. s. comm. cum var. à cl. inv.)
116. *V. SERRATA* (Less. in *linnæâ* 1829. p. 275. 1831. p. 649), herbacea, caule tereti striato glabro ramoso, ramis pubescentibus, foliis petiolatis membranaceis ellipticis utrinquè brevè acuminatis argutè duplò serratis suprà glabris, subtus petiolisque pilosiusculis, cymæ terminalis scorpioideæ ramis divaricatis, capitulis multifloris lateralibus subsessilibus, invol. campanulati squamis scariosis 1-nerviis ovato-oblongis mucronato-acutis, achenio piloso multicostato, pappi serie ext. paleaceâ brevi. — in *Brasiliæ nemorosis montibus Mandiocæ*. Cor. violaceæ? glabra. Pappus argenteus subplumosus. Petioli 6-18 lin. longi. Folia  $2\frac{1}{2}$ -8 poll. longa, 1-4 poll. lata.
117. *V. PUBERULA* (Less. in *linnæâ* 1831. p. 649), caule fruticoso, ramis teretibus sordidè puberulis apice striatis, foliis petiolatis oblongis basi acuminatis apice acutiusculis subserrulatis, suprà glabris, subtus petiolisque puberulis, cymæ terminali decompositâ subaphyllâ, capitulis 8-9-floris, invol. turbinati squamis imbricatis obtusis, ext. brevissimis, cæteris gradatim longioribus, achenio oblongo hirsuto-canescente, pappi ser. ext. paleaceâ brevi.  $\beta$  in *Brasiliâ* legit cl. Sellow. Inv. squamæ sicce adpressæ pubescentes. Pappus albus. Fol. 2 poll. longa, 6 lin. lata, petiolo 2-lin. (v. s. comm. à Mus. reg. Berol.)
118. *V. WRIGHTIANA* (Less. in *linnæâ* 1831. p. 650), caule herbaceo simpliciter hispidiusculo pubescente tereti, foliis petiolatis firmis oblongis obovatis subintegerrimis penninerviis, suprà scabris hirsuto-pubescentibus, subtus tomen-

toxis, panicula terminalis ramis subverticillatis aphyllis, capitulis subsessilibus 12-floris, inv. campanulati squamis siccis ciliatis ellipticis obtusis, cor. glabra, achenio hirsuto, pappi ser. ext. brevi setosâ. — in montosis Brasiliæ prope Villa-Ricca. Cor purpureo-violaceo. Pappus rufescens. Affinis V. Beyrichii dicitur.

a. *angustifolia*, foliis oblongo-ellipticis basi obtusis semipollicem latis.

ß. *cuneata*, foliis obovatis ultrâ pollicem latis basi cuneatis.

119. V. *BEYRICHII* (Less. in Linnæ 1829. p. 275. 1831. p. 651), herbacea, caule tereti striato glabro, foliis confertis petiolatis lanceolatis rugulosis subdenticulatis, suprâ glaberrimis, infrâ minutè glandulosis ad nervos pilosiusculis, cymis terminalibus scorpioides divaricatis, capitulis subsessilibus multifloris, invol. turbinati squamis siccis multinerviis ovatis acutis pubescentibus et ciliatis, achenio oblongo multicostato, pappi serie ext. brevi. — in humidis prov. Sanctæ-Annæ ad Rio-Janeiro. Cor. rosæ glabræ. Pappus flavescens. Folia 3-3½ poll. longa, 10-11 lin. lata. (v. s. comm. à cl. Lund.)
120. V. *INCANA* (Less. in Linnæ 1829. p. 277), caulibus herbaceis teretibus simplicibus pube adpressâ subincanis apice angulatis ramoso-corymbois, foliis sessilibus lineari-lanceolatis acuminatis subdenticulatis, suprâ glabriusculis, subtis pilis adpressis incanis, cymis scorpioides in corymbum dispositis subaphyllis, capitulis multifloris sessilibus et pedunculatis, invol. subturbinati squamis adpressis oblongis acutis extis piloso-subincanis, achenio oblongo multicostato piloso et glanduloso, pappi serie externâ internam subsequantem. ½ in Brasiliâ meridionali. Rhizoma multiceps lignosum. Cor. violaceæ glabræ. Pappus argenteus. Herba fructifera refert Hieracium cymosum. (v. s. comm. à Mæa. reg. Berol.)
121. V. *ECHIOIDES* (Less. in Linnæ 1829. p. 278. 1831. p. 651), caulibus è rhizomate lignoso pluribus teretibus simplicibus striato-sulcatis strigoso-pubescentibus, foliis sessilibus membranaceis utrinquè asperis elongato-lanceolatis undulato-crenatis integerrimis margine revolutis, cyma divaricata scorpioides ramis dichotomis, capitulis multifloris subsessilibus, invol. campanulati squamis oblongis 1-nerviis glabris, achenio multicostato piloso, pappi serie ext. brevi. ¾ in Brasiliâ meridionali. Habitus Hieraci echioidis. Cor. glabræ violaceæ. Pappus niveus.
122. V. *DENTICULATA*, caule herbaceo erecto sulcato-angulato apice tenuissimè puberulo, foliis petiolatis lanceolatis utrinquè acuminatis serratulis pascia distantibus denticulatis, suprâ glabris, subtis pube tenuissimâ subpuberis, paniculâ laxâ polycephalâ aphyllâ, ramis infer. axillaribus hinc inde suboppositis folio longioribus, capitulis secus ramos sessilibus lateralibus 10-12-floris, inv. basi obtusi squamis glabris ext. brevioribus acutis, int. oblongis obtusis macronulatis, cor. glabrâ, achenio brevissimo pubero striato, pappi ser. ext. setosâ brevi. — in Brasiliâ circa Rio-Janeiro. Folia 2-2½ poll. longa, 6 lin. lata, petiolo 2-3 lin. Pappus stramineo-sabrescens. Affinis videtur V. Beyrichii ex descr. Folia summa integra. (v. s. comm. à cl. Lund. J. Gandieh. Blanch.)
123. V. *PAITACORUX*, arborea, ramis glabris, ramulis subhirtellis angulatis, foliis subsessilibus lineari-lanceolatis utrinquè acuminatis integerrimis, suprâ minutè scabridis, subtis hirsutis, cymâ ramodensâ paniculaformi polycephalâ, invol. squamis ovatis glabresculis, ext. mucronatis, inter. obtusiusculis, achenio pubero, pappi serie ext. setosâ brevi. 5 in montosis circa Rio de Janeiro. Vulgatissima arbor, vulgo *aspeiro* dicta. Herba tota et praesertim flores suaviter olentes. Flor. jul. aug. Trochili paniculas coramvolant. Paitacox vittatus (Kuhl.) semina avidè devorat (Lund.). Folia 3-4 poll. longa, 6-8 lin. lata. Cortex lenticellis verruculosus. (v. s. comm. à cl. Lund.)
124. V. *HERACLEADA*, fruticosa, ramis teretibus striatis pube brevissimâ rufâ puberula, foliis petiolatis oblongo lanceolatis utrinquè acuminatis integerrimis, suprâ aspero-scabridis, subtis hirsuto-tomentosis, cyma polycephala ramis subscorpioides, ramis inferioribus ex axillis fol. ortis, sup. nudis, invol. squamis

ovali-oblongis mucronatis, caput. aliis sessilibus aliis pedicellatis circ. 10-floris, acheniis striato-sulcatis puberis, pappi serie ext. Brevi. 5 in Brasilia prov. S<sup>t</sup>. Pauli frequens obs. cl. Lund. Fl. decemb. Valdè affinis *V. eriodada*, sed rami vix puberi, folia dimidio angustiora, panicula basi foliosa, et inv. squamæ mucronata nec acuminata. Pappus sordide rufus. Cor. purpurea. Folia 3 poll. longa, 5-6 lin. lata. (v. s. comm. à cl. Lund.)

125. *V. PETIOLARIS*, fruticosa, ramis teretibus velutino-tomentosis, foliis alternis longè petiolatis elliptico-oblongis utrinquè acuminatis integerrimis, suprâ sparè puber. subtus molliter petiolisque hirsuto-tomentosis, panicula amplis ramis scorpiolideis aphyllis, capitulis subsessilibus unilateralibus 30-floris, invol. turbinati squamis multiseriatis semipatentibus dorso velutinis oblongis acutis, ext. brevibus secus pedicellum imbricatis, cor. glabrâ, achenio villosa, pappi ser. ext. brevi vix paleacea, 5 in Brasilia sylvis primævis leg. cl. Lhotsky. Petioli 9-10 lin. longi, limbi fol. 5-6 poll. longi, sesquipoll. lati. Inv. squamæ nec adpressæ nec squarrosæ, sed semipatentes! (v. s. comm. à cl. Lhotsky.)

β *pasciflora*, foliis basi magis obtusis, capitulis paulò minoribus 10-12 floris. 5 ad margines sylvarum circa Rio-Janeiro legit cl. Lhotsky. Arbor 25-30-pedalis. (v. s. comm. à cl. invent.)

126. *V. EUPATORIIFOLIA*, arborea, ramis teretibus densè pedunculis petiolisque hirsuto-tomentosis, foliis oppositis petiolatis ellipticis acuminatis basi obtusis integerrimis glandulis sessilibus utrinquè gerentibus, suprâ pubescentibus nervo medio basi hirsuto, subtus hirsutis, paniculæ ramis ex axillis supr. aut ad apicem ortis subscorpiolideis polycephalis, capitulis pedicellatis 30-floris, invol. turbinati squamis ovali-oblongis extus villosis demum semipatulis, ext. brevioribus, cor. glandulosâ glabrâ, achenio puberulo, pappi serie ext. paleacea brevi. 5 in sylvis collium circa Rio-Janeiro legit cl. Lund. Affinis *V. petiolaris* sed oppositifolia! Differt à *V. oppositifolia* caule arboreo 8-12-pedali nec herbaceo suprâ fol. paria non dilatato, ramis folisque magis hirsutis. (v. s. comm. à cl. Lund.)

β *pasciflora*, foliis basi subacutis parcius glandulosis, capitulis 10-12-floris, corollis subglandulosis, 5 in fruticetis circa Rio-Janeiro legit cl. Lhotsky. Frutex 5-7-pedalis. (v. s. comm. à cl. invent.)

127. *V. HAVANENSIS*, caule fruticoso tereti ramoso glabro, ramulis striatis puberulis, foliis ovalibus utrinquè acuminatis basi cuneatis brevissimè petiolatis integerrimis penninerviis, suprâ muriculatis, subtus glandulosis et tenuiter pubescentibus, cymis axill. terminalibus pedunculatis laxis oligocephalis, capitulis pedicellatis aphyllis distantibus 25-floris, invol. turbinati squamis dorso glandulosis glabriusculis, ext. brevibus acutis, int. oblongis subcaneatis obtusiusculis, cor. et achenio glabris parè glandulosis, pappi ser. ext. brevi setiformi. 5 in ins. Cubâ juxta Havanam legit cl. Ramon de la Sagra. Invol. subpurpurascens. Cor. purpurea. Pappus albus. Folia 4 poll. longa, sesquipollicem lata, juniora marginibus subrevoluta. (v. s. comm. à cl. R. de la Sagra.)

128. *V. LANCEOLARIS*, caule tereti striato pube ferrugineâ parè velutino, foliis petiolatis lanceolatis basi subobtusis apice acuminatis serrulatis, suprâ glabris, subtus petiolisque parè velutinis, paniculæ ramis axillaribus ebracteatis ramosis velutinis, capitulis sessilibus ovatis 20-floris lateralibus et terminalibus, invol. squamis omnibus mucronatis dorso subpuberis, cor. glabrâ, receptaculo alveolato, achenio puberulo, pappi ser. ext. brevi vix paleacea. 2? in Mexico ex h. Haenke. Affinis *V. Haenkeana*. Pappus croceo-rufus rigidulus. Folia 5 poll. longa, pollicem lata, petiolo 4 lin. (v. s. in h. Haenke à cl. Sternberg missum.)

129. *V. HAENKEANA*, caule teretiusculo erecto ramis pedunculisque velutino-subhirsutis, foliis petiolatis ellipticis basi acutis apice acuminatis integerrimis penninerviis, suprâ demum glabris, subtus velutino-subhirsutis, ramis axillaribus foliosis apice paniculatis, ramulis bi-trifidis, capitulis axillaribus latera-

*Mucosae sessilibus aut pedicellatis ebracteatis 20-floris, invol. squamis oblongis glabrisculis ext. mucronatis inter. obtusis, cor. glabrâ, achenio pilosiusculo, pappi ser. ext. brevissimâ vix paleacea. ? inter Peruanas Orinocenas in herb. habet cl. Haenke. Folia 5-6 poll. longa, 2 poll. lata, petiolo 7-9 lin. longo velutino. Pappus croceo-rubens rigidulus. Pubes ramorum rufescens. (v. s. in h. Haenke à cl. Stenberg missum.)*

130. *V. MICRADENIA*, fruticosa, ramis teretibus densè et breviter sericeo-villosis foliis petiolatis ellipticis basi attenuatis apice acuminatis integerrimis penninerviis utrinquè secus nervos et in petiolo sericeo-pubescentibus suprâ sparè pilosis subtus sparè glandulosis, cymis scorpioides, aliis axillaribus oligocephalis folio brevioribus aliis terminalibus stellato-divaricatis polycephalis, capitulis unilateralibus ebracteatis 10-floris, invol. squamis dorso subvillosis 1-nerviis, ext. acuminatissimis, int. linearibus vix acutis, achenio subvillosis, pappi ser. ext. paleacea brevi. ? in Amer. calid. fortè Peruvia legit cl. Poeppig. (pl. exs. n. 1215.) Affinis ex descr. *V. frangulifolia*. Folia caul. 6 poll. longa, 2-2 ½ poll. lata, petiolo 8-9 lin. longa. (v. s. comm. à cl. Poeppig.)

131. *V. MENTHIFOLIA* (Less. in linnæ 1829. p. 258. 1831. p. 645), fruticosa, ramis teretibus divaricatis apice obscure pubescentibus demum glabris, foliis petiolatis ovato-oblongis acutis basi attenuatis membranaceis utrinquè scabridis subtus pilosiusculis, cymis subfastigiatis dichotomis divaricatis, capitulis 15-floris, invol. campanulatis squamis 1-nerviis coriaceis ovatis acutis ciliatis, intimis obtusis glabris, cor. glabrâ, achenio glabrisculo, pappi serie ext. setiformi breviori. ? in fruticeis insula Cuba. Eupatorium menthaefolium Poeppig. nunc. ex Less. Cor. glabrâ. Pappus flavescens. Folia 1-3 poll. longa, 4-12 lin. lata, integra aut serrata. (v. s. comm. à cl. Ramon de la Sagra.)

§. 3. *Involucri squamis plerisque acuminatis, capitulis in cymas scorpioides dispositis rarò subsolitariis, basi folia perfecta non gerentibus.* Less. in linnæ 1831. p. 651. — *Species omnes Americanae.*

132. *V. ODORATISSIMA* (H. B. et Kunth nov. gen. 4. p. 32), caule fruticoso tereti, ramis striatis velutino-pubescentibus, foliis brevè petiolatis rigidis obovatis basi obtusis apice subdentatis, suprâ scabridis, subtus reticulatis hirsuto-pubescentibus, cymæ ramis plurimis ramosis in paniculam dispositis, capitulis innumeris pedicellatis later. et terminal. folio florali destitutis, invol. squamis pluriariatis ovatis acuminatis, achenio vix pilosiusculo, pappi serie ext. setacea brevi. ? ad Caracas, Ibaque et Honda. Less. in linnæ 1829. p. 271. 1831. p. 631. *Conyza clethraefolia* et *Eupatorium obovatum*, Willd. herb. Cor. alba glabrâ. Pappus stramineus. Variat foliis serratis ex Kunth, oreatis in specim. meo Caracano, et subintegerrimis ex Less. (v. s.)

133. *V. RUBRIRAMEA* (Mart. ! herb.), caule herbaceo quinquangulato glabrisculo erecto, foliis subpetiolatis oblongo-lanceolatis basi acuminatis apice subacutis subintegerrimis, suprâ glabris glabrisculisve, subtus pubero-subhirsutis, panicula ramis axillaribus polycephalis dichotomo-ramosis folio longioribus subaphyllis, capitulis lateralibus sessilibus terminalibus ebracteatis 20-floris, invol. squamis linearis-oblongis mucronatis glabris, cor. glabrâ, achenio striato pubero, pappi ser. ext. paleacea acuminatâ vix inter. triplò brevior. ? in campis prope Gongonhas de Sabara prov. Minarum gener. Brasiliæ legit cl. Martins. Folia 4 poll. longa, 7-8 lin. lata. Rami purpurei. Corollæ albae. Capit. vix 3-lin. longa. (v. s. in h. Acad. reg. Monac.)

134. *V. ASSANA* (Mart. ! herb.), caule herbaceo ? erecto striato-angulato breviter velutino, foliis brevè petiolatis elliptico-oblongis utrinquè obtusis rigidis suprâ scabris, subtus velutino-puberis penninerviis nervis lateralibus reticulatis, ramis floridis bifidis subscorpioides oligocephalis folio longioribus aphyllis, capitulis ovatis 30-floris sessilibus, invol. squamis imbricatis glabrisculis acuminatis apice non adpressis, cor. glabrâ, achenio pubero, pappi ser. ext. vix paleacea

- brevissima panicetâ.  $\frac{1}{2}$   $\frac{3}{4}$  in campis desertis Minarum gener. Brasilia collegit cl. Martins. Petiolus 3 lin. Limbus fol.  $2\frac{1}{2}$ -3 poll. longus, pollicem latus. Pappus rufus. Recept. areolatum. (v. s. in h. Acad. reg. Monac.)
135. *V. POLYANTHES* (Less. in linnæâ 1831. p. 631), subarbores, ramis angulatis puberulis, foliis petiolatis lanceolatis utrinque acuminatis subintegerrimis suprâ scabris, subtus puberis nervis elevatis subhirsutis, panicula ramosissima puberâ ramis scorpioides subverticillatis divergentibus aphyllis, invol. 15-20-flori squamis oblongis subacuminatis glabrescens, achenio puberulo punctato, pappi seriebus setiformibus subæqualibus.  $\frac{5}{8}$  in montosis Brasiliæ circa Rio-Janeiro alt. 2-3000 ped., et in prov. Minar. gener. Eupatorium polysanthes Spreng. syst. 3. p. 435. Vern. patens Less. in linnæâ 1829. p. 276. ex parte et excl. syn. Vern. Brasiliensis Mart. ! herb. Cor. glabrâ albâ. Folia trita odorata (Lund.) 5-6 poll. longa, pollicem lata. Pappus flavescens. (v. s. comm. à cl. Lund.)
136. *V. GRANDIFOLIA*, arborea, ramis teretibus striatis apice subpubescentibus, foliis petiolatis ellipticis utrinque acuminatis simpliciter et grosse serratis membranaceis glabrescens secus nervos et venas minutè puberulis, paniculâ corymbosâ amplâ divaricatâ polycephalâ aphyllâ, capitulis pedicellatis 20-25-floris, invol. squamis glabris acuminatis margine membranaceis, cor. lobis puberulis, achenio tereti turbinate striato hispido-villoso, pappi ser. ext. paleaceâ brevi.  $\frac{5}{8}$  in monte Cercovado Brasiliæ legit cl. Lund. Arbor 8-10 ped. Folia inf. 16 poll. longa, 8 poll. lata, petiolo 2 pollic. Pappus albus caducissimus. Recept. alveolare. Inv. demum patens. (v. s. comm. à cl. Lund.)
137. *V. PATENS* (H. B. et Kunth nov. gen. 4. p. 32.), arborea, ramulis angulatis obscure pubescentibus, foliis breviter petiolatis angustè lanceolatis acuminatis basi angustatis subrenato-undulatis suprâ rugosis scabrescens, subtus hirtopubescentibus, cymâ scorpioides terminalis ramis patentibus, capitulis multifloris distantibus subsessilibus, invol. campanulati squamis siccis 1-nerviis ovatis acutis adpressis pilosisculis, achenio multicostrato piloso glanduloso, pappi ser. ext. paleaceâ brevi.  $\frac{5}{8}$  in Amer. merid. (Humb.) *V. patens* Less. in linnæâ 1831. p. 653. Baccharis rugosa Willd. herb. ex Less. Cor. glabrâ albâ.
138. *V. SCABRA* (Pers. ench. 2. p. 404. Less. in linnæâ 1831. p. 653), fruticosa, ramis angulatis divaricatis apice hirtello-puberulis, foliis brevissimè petiolatis ellipticis aut obovatis utrinque obtusis subintegerrimis suprâ scabris, subtus hirsuto-tomentosis, paniculis terminalibus ramosissimis aphyllis, capitulis subsessilibus aut pedicellatis 30-40 floris, inv. turbinati squamis glabrescens oblongo-linearibus acuminatis, cor. glabrâ, achenio pubescente, pappi ser. ext. brevi setosâ.  $\frac{3}{8}$  in Brasiliâ (ubi ex Margr. dicitur *Tremate*), in collibus juxta Bahiam (Salzm. ! Blanch. !), in montibus prov. Minarum gener. Mart. ! Baccharis Brasiliensis Linn. spec. 1205. — Margr. bres. 2. p. 81. fig. sup. Pappus subrufescens. Cor. purpurascens vanillam (Salzm. !) styracem (Margr.) olentem. Folia 12-15 lin. longa, 7-8 lin. lata. (v. s.)
139. *V. POLYCEPHALA*, caule fruticoso, ramis teretibus velutino-canescens, foliis brevè petiolatis ovalibus utrinque obtusis margine subcrenulatis et revolutis suprâ subrugosis pubescentibus, subtus tomentoso-subincanis, cymâ ramis plerimè ramosis in paniculam dispositis, capitulis 20-floris axillaribus et lateralibus subsessilibus folio floralis destitutis, involucri obovati squamis pluriseriatis ovatis subacutis, achenio vix pubero, pappi serie ext. vix paleaceâ brevi.  $\frac{5}{8}$  in collibus aridis circa Bahiam Brasiliæ legit cl. Salzm. Caud. 4-pedalia. Rami cani. Cor. pallida. Pappus stramineo-subrufescens. (v. s.)
140. *V. MICROCEPHALA*, suffruticosa, ramis teretibus pube brevi subhirsutis, foliis brevissimè petiolatis oblongis obtusis basi subcordatis integerrimis suprâ scabridis glabris subtus ferrugineo-villosis, cymis terminalibus subcorymbosis, ramis nudis lateralibus scorpioides, capitulis 20-floris lateralibus subsessilibus obovatis, invol. squamis pluriseriatis imbricatis ext. brevissimis acutis, internodiis longioribus, achenio glabrescens, pappi serie ext. paleaceâ brevi.  $\frac{3}{8}$  in Amer. calidior. legit cl. Née, sed locus proprius ignotus. Affinis *V. odora*-



tissima et scabra. Capitula subsora. Folia 2-2½ poll. longa, 6-9 lin. lata. (v. a. ex herb. Thibaud.)

141. *V. SONORIA*, suffruticosa, ramis teretibus hirsutulis, foliis breviter petiolatis ovatis acuminatis basi obtusis integerrimis supra villosis, subtus dense villosco-hirsutis, ramis floridis paniculatis elongatis bifidis hirsutulis, ramulis capitula ebracteata sessilibus remota unilateralia gerentibus, invol. ovati hirsuti squamis omnibus acuminatis subpungentibus. ♀ in Brasiliâ circa Rio de Janeiro legit cl. Gaudichaud. Valde affinis *V. microcephala*, sed differt ramis verè hirsutis, foliis ovatis supra villosis, ramis cymis magis elongatis, invol. hirsuto. Flores in specim. juniores et non satis vidi. (v. a. comm. à cl. inv.)
142. *V. HYPOCEPHALA*, caule ramoso dense et adpressè velutino, foliis petiolatis ovali-lanceolatis utrinque (apice præcipuè) acuminatis integerrimis penninerviis, subtus venoso-reticulatis supra glabris, subtus glandulosis et breviter pubescentibus, panicula ample ramosissima ramis axillaribus folio longioribus bifidis scorpioides, capitulis lateralibus et terminalibus sessilibus solitarie ebracteatis 20-floris, invol. squamis ext. acuminatis ciliatis int. glaberrimis mucronatis, cor. glabrâ, achenio turbinate sericeo-villoso, pappi ser. ext. paleacea brevi. 2½ inter Peruanas Orinocenses herb. Haenkeani odet. Affinis *V. polyanthi*. Rubes caulis et pedunculorum subflavescentes velutina. Folia 4 poll. longa, 1½-2 poll. lata, petiolo semipoll. Invol. purpurascens. Pappus albus. Cor. purpureus. (v. a. in h. Haenk. à cl. de Sternberg missum.)
143. *V. DEFFREANA* (Less. in Linnæ 1831. p. 398 et 655), fruticosa, ramis teretibus obscurè pubescentibus, foliis petiolatis membranaceis penninerviis ellipticis utrinque subattenuatis apice mucronatis integerrimis supra scabris, subtus tomentoso-pubescentibus, cymis scorpioides compositis in paniculam myriocephalam dispositis, capitulis secus ramos lateralibus sessilibus turbinate 20-floris, invol. squamis rigidis mucronato-acuminatis, cor. glabrâ, achenio piloso, pappi ser. ext. paleacea brevi. ♀ in Mexico circa Misantla. Cor. alba odoratissima. Folia 3-3½ poll. longa, 12-15 lin. lata, petiolo 2-5 lin. (v. a. in h. Acad. reg. Monac.)
144. *V. APPROXIMATA* (Less. in Linnæ 1831. p. 655), fruticosa, ramis teretibus striatis sordidè tomentosis, foliis petiolatis ellipticis basi cordatis suberratis supra scabriusculis, subtus molliter pubescenti-tomentosis, cymis scorpioides curvatis aplyllis, capitula 11-floris lateralibus pedicellatis, inv. turbinate squamis 8-seriatis siccis acuminatis muticis dorso tomentoso-incanis, ext. rotundis, cor. glabrâ, achenio villoso, pappi ser. ext. paleacea brevi. ♀ in Brasiliâ. Cor. purpureo-vioacea. Fol. circ. semipoll. 3-4 poll. lata. Similis *V. oppositifolia* et quinqueflora dicitur.
145. *V. QUIQUERFOIA* (Less. in Linnæ 1831. p. 656), caule herbaceo tereti cum pedunculis petiolis folisque subtus hirsuto-pubescentibus, foliis petiolatis elliptico-oblongis utrinque acuminatis rariter subserratis, supra scabris, cymâ terminali compositâ parè ad basin pedunculorum bracteâ, capitulis nudis pedicellatis ovali-oblongis 5-floris, invol. squamis siccis dorso villosco-pubescentibus imbricatis subacuminatis, achenio hirsuto, pappi ser. ext. brevi subpaleacea. 2½ in Brasiliâ legit cl. Sellow. Folia 5-6 poll. longa, 15 lin. lata. (v. a. comm. à Mus. reg. Berol.)
146. *V. FRANGULIFOLIA* (H. B. et Kunth nov. gen. 4. p. 37), fruticosa, ramis teretibus villosis, foliis brevè petiolatis oblongis acutis basi angustatis integerrimis rigidè pubescentibus nitidis, subtus glandulosis pubescenti-pilosis canescentibus, cyma scorpioides terminalis ramis congestis fastigiatis, capitulis multifloris approximatis sessilibus, invol. hemisphærici squamis lanceolatis acutis 3-nerviis externis tomentoso-pubescentibus, achenio piloso, pappi ser. ext. paleacea brevi. ♀ in calidis Novo-Granatensium prope Honda. Less. in Linnæ 1829. p. 276. 1831. p. 657. Cor. glabra violacea. Pappus argenteus. Fol. 2-2½ poll. longa, 4-9 lin. lata, petiolo 2 lin.
147. *V. CANESCENS* (H. B. et Kunth nov. gen. 4. p. 35. t. 317. Less. in Linnæ 1829. p. 284. 1831. p. 657), caule basi suffruticosa, ramis teretibus pilis ad-

premiis canescentibus, foliis brevè petiolatis rigidis membranaceis oblongo-lanceolatis acuminatis basi angustatis integerrimis subrugosis suprà pubescentibus, subtus piloso-canescentibus glandulosis, cymæ terminalis scorpioideæ ramis oligocephalis, capitulis distantibus sessilibus, involucri campanulati squamis siccis 1-nerviis acuminatis pilosis glandulosisque, achenio villosa, pappi serie ext. brevissima.  $\frac{1}{2}$   $\frac{3}{4}$  in Peruvia temperatâ prope Guancabamba. *Lepidaploa canescens* Cass. dict. 26. p. 18. Vern. reticulata Willd. herb. ex Less. *V. canescens* Spreng. syst. 3. p. 437. excl. syn. Pers. et Sellow? *V. geminata* Kunth nov. gen. 4. p. 35? Cor. pallidè violaceæ. Pappus argenteus. (v. s.)

*V. micrantha* (Less. in Kunth 1831. p. 637), suffruticosa? ramis subangulatis sericeo-pilosis, foliis petiolatis oblongis acuminatis basi rotundatis integerrimis suprà pubescentibus, subtus argenteo-piloso-sericeis, cymis terminalibus paniculaformibus ramosis, ramulis geminis dichotomis, capitulis solitariis sessilibus distantibus multifloris.  $\frac{3}{4}$ ? in prov. Cumanæ? Vern. *micrantha* H. B. et Kunth nov. gen. 4. p. 37. *V. acuminata* Willd. herb. ex Less.

148. *V. subequarrosa*, ramis teretibus superne villosa-hirsutis, foliis brevissimè petiolatis ovato-oblongis, basi obtusis apice acutis integerrimis suprà glabris rugosis, subtus villosa-hirsutis subcanescentibus, cymis axillaribus terminalibus bifidis divaricatis scorpioideis, capitulis ovatis 20-floris sessilibus, involucri squamis villosis lanceolatis externis subpungenti-acuminatis, acumine patente squarrosis, int. minutis acuminatis erectis, cor. glabris, achenio turbinate villosa, pappi serie ext. paleacea brevi.  $\frac{1}{2}$ ? in Brasiliâ ad margines sylvarum et aqueductus circa Rio-Janeiro (Mart.) et Bahiam (Blanch.). Cor. verisim. purpurea. (v. s.)

149. *V. mollis* (H. B. et Kunth nov. gen. 4. p. 28), caule herbaceo tereti sericeo-viloso, foliis brevè petiolatis ovato-lanceolatis acuminatis basi angustatis remotè demumatis integerrimis suprà puberulis, subtus argenteo-lanuginosis, cymæ terminalis scorpioideæ ramis sæpè geminis elongatis, capitulis sessilibus ebracteis 20-floris, inv. campanulati squamis 1-nerviis sericeo-pilosis acuminatis demum patentibus, cor. parè pilosis, achenio parè sericeo-piloso, pappi serie ext. paleacea brevi.  $\frac{1}{2}$  in Peruvia? (herb. Dombey!) ad ripam fl. Catamayo prope Gonzansua Novo-Granatensium ex Hamb. itin. et ad Orinocum ex herb. Hænk! Fol. 3-5 poll. longa, 1-2 poll. lata. Pappus argenteus. *Eupatorium polystachyum* Willd. herb. ex Less. Vern. mollis Less. in Linneæ 1823. p. 283. 1831. p. 657. (v. s. in h. Hænk. à cl. Sternb. missum.)

150. *V. flavescens* (Less. in Linneæ 1831. p. 657), fruticosa, ramis teretibus striatis rufulo-tomentosis, foliis petiolatis ovatis subrotundisve suprà levibus subtus glandulosis flavido-villosis subintegerrimis, cymæ ramis scorpioideis patentibus, capitulis sessilibus lateralibus multifloris, recept. nudo, cor. glabris, achenio piloso, pappi ser. ext. palea latissimis ovario brevioribus.  $\frac{3}{4}$  in Brasiliâ, ad Serra d'Estrella. Vern. scorpioideus Less. in Linneæ 1829. p. 282. excl. syn. Folia basi obtusa aut acuta. Capitula approximata.

151. *V. tournefortoides* (H. B. et Kunth nov. gen. 4. p. 34), herbacea, ramis teretibus striatis petiolisque flavido-pubescentibus, foliis breviter petiolatis ovato-lanceolatis subintegerrimis suprà scabris hirsutiusculis, subtus pubescentibus punctatis, spicis scorpioideis geminatis, capitulis sessilibus 25-floris, recept. levissimè alveolato, corollis glabris, achenio glabrisculo, pappi serie ext. paleacea ovario subæquali. — in Americâ merid., sed locus propr. ignotus. *V. Tournefortoides* Less. in Linneæ 1831. p. 657 non 1829. *Stanhelia solidaginoides* Willd. herb. ex Less.

152. *V. scorpioides* (Pers. ench. 2. p. 404), fruticosa, ramis teretibus puberulis, foliis breviter petiolatis ovali- aut ovato-lanceolatis acuminatis subdentatis aut subrepandis suprà glabriusculis, subtus pubescentibus hirsutisve, cymæ ramis scorpioideis, capitulis secus ramos sessilibus 1-lateralibus ebracteis circ. 20-floris, corollæ lobis pilosis, achenio turbinate piloso, recept. paleaceo fimbriatissimo, pappi ser. ext. vix paleacea achenii longitudine.  $\frac{3}{4}$  in-

Brasiliâ tropicâ, Guianâ. *V. scorpioides* Less. in linnæâ 1831. p. 658. non alibi. *V. Tournefortioides* Less. in linnæâ 1829. p. 281. non Kunth *V. centriflora* Link et Otto abb. t. 55. *Lepidaploa scorpioides* Cass. dict. 26. p. 18. *Coryza scorpioides* Lam. dict. 2. p. 48.

*a. centriflora*, foliis ovalibus utrinque acuminatis suprà glabris, subtus pubescentibus. In Brasiliâ, Guianâ. Satis refert icon. Link. (v. s.)

*ß. subrepanda*, foliis latè ovatis subrepandis apice acuminatis suprà glabris, subtus pubescentibus. — in Guianâ gallicâ et Brasiliâ. Vern. *subrepanda* Pers. ench. 2. p. 404. (v. s.)

*γ? subtomentosa*, foliis latè ovatis subdentatis apice acuminatis suprà puberis, subtus hirsutis. — ad Caracas legit cl. Vargas. (v. s.)

*δ? longifolia*, foliis elongato-lanceolatis acuminatis suprà glabris secus nervos puberis, subtus præsertim in nervis puberis. — circa Bahiam in umbrosis legit cl. Salmann. (v. s.)

*ε? longeracemosa*, foliis elongato-lanceolatis acuminatis suprà subecabris subtus hirsuto-subtomentosis, apicis longissimis. — in Brasiliâ, Peruvâ. Vern. *longeracemosa* Mart. herb. — Vauth. pl. exa. n. 262. Poepp. n. 33. (v. s.)

153. *V. IGNEOLILIA* (Less. in linnæâ 1831. p. 658), fruticosa?, ramis teretibus hispidinaculo-pubescentibus, foliis petiolatis rigidis adproximatis oblongo-obovatis obtusis suprà asperimis, subtus hirsutinaculo-pubescentibus, cymis scorpioides aphyllis, pedunculis 2-3-cephalis, capitulis 10-floris, invol. globosi squamis siccis glabrisculis & nervis acuminatis ext. rotundis, cor. glabrâ, achænio villosâ, pappi seriebus confluentibus setaceis. *γ?* in Brasiliâ. Cor. violacea. Fol. 3. poll. longa, pollicem lata.

154. *V. OBOVATA* (Less. in linnæâ 1829. p. 279. 1831. p. 659), caule herbaceo simplici tereti striato villosâ, foliis sessilibus membranaceis obovatis undulato-crenatis suprà scabris pilosis subtus villosis, cymis terminalis scorpioides ramis divaricatis dichotomis, capitulis adproximatis multifloris sessilibus, invol. campanulati squamis paucis siccis 3-nerviis adpressis acuminatis extus pilosis, achænio subsericeo, pappi serie ext. palacæâ brevi. — in Brasiliâ tropicâ. Cor. glabrâ violaceâ. Pappus argenteus. Folia variant oblonga aut elliptica, obtusa aut acuta, 2  $\frac{1}{4}$ -4 poll. longa, 10-20 lin. lata.

155. *V. PELLITA* (H. B. et Kunth nov. gen. 4. p. 28. Less. in linnæâ 1829. p. 279), fruticosa?, ramis teretibus sordidè lanatis, foliis sessilibus lanceolatis coriaceis margine revolutis suprà rugosis scabro-pilosis, subtus obscure lanatis, cymis terminalis scorpioides ramis dichotomis, capitulis adproximatis sessilibus circ. 20-floris, invol. campanulati squamis siccis adpressis acuminatis pungentibus extus et apice lanatis, achænio oblongo densè sericeo-villosâ, pappi serie ext. brevî vix palacæâ. *γ* in prov. Caracasanâ. Cor. ex Kunth glabrâ, ex Less. pilosâ, in nostro specim. lobis apice parcissimè pilosis, Pappus sordidas, ex Less. albus, in spec. Varg. rigidulus faciliè deciduus. (v. s. comm. à cl. Vargas.)

156. *V. SEPTARTHA*, caule tereti erecto densè et obscure velutino-hirsuto, foliis petiolatis elliptico-oblongis basi subacuminatis apice cuspidato-acuminatis subintegerrimis suprà brevissimè puberulis, apertis petiolisque densè hirsutis, panicula ramis axillaribus folio brevioribus simplicibus bifidis ebracteatis, capitulis sessilibus confertis spicatis 7-floris, invol. squamis lineari-lanceolatis subacuminatis dorso villosis, cor. lobis apice puberulis, achænio turbinato sericeo-villosâ, pappi ser. ext. angustè palacæâ acuminatâ int. quadruplò breviorè. *℥? γ?* inter Peruanas montanas Orinocoenses habet herb. Hænk. Fol. 3 poll. longa, 9 lin. lata, petiolo 3-4 lin. marginibus subrecurvatis. Cor. purpureâ. Tota planta in sicco obscura. (v. s. in h. Hænke à cl. Sternb. missum.)

157. *V. VIRGULATA* (Mart. herb.), caule herbaceo? erecto tenui virgato tereti striato glabrâ, foliis infer. . . . super. distantibus linearibus minimis integerrimis subtus pubescentibus, capitulis 10-12 floris secus ramos spicatis sessilibus aphyllis, invol. squamis oblongis dorso adpressè pubescentibus, ext. brevibus mucronatis, intimis longioribus subacuminatis, cor. puberâ, achænio

sericeo-villoso, pappi ser. ext. angustè palacèà acuminatè brevè.— $\chi$  ? in editis prov. Minarum gener. Brasiliæ. Folia sup. vix 2-3 lin. longa, inf. ignota. Pappus substramineus. Recept. nudum. (v. s. in h. Acad. reg. Monac.)

158. *V. DECAURONTUM* (Mart. herb.) glabrituscula, collo crasso, caulibus pluribus subsimplicibus sparsè foliosis apice hirsuto-villosis nudis 1-cephalis, foliis sessilibus linearibus acuminatis integerrimis planis nervoso-subreticulatis sparsè pilosis glabratissè, invol. 1-2-serialis squamis lanceolatis extus villosis.— $\chi$  in deserto trans flumen Sancti Francisci prov. Minarum gener. Brasiliæ legit cl. Martins. Herba 6 poll. longa. Affinis *V. campestris*. Folia 2. poll. longa vix 2 lin. lata. Cap. dimidio minus quàm *V. campestris*. Achænium sericeo-villosissimum. Pappi pallidè rufescentis paleæ ext. breves lineares acuminatæ, int. elongatæ setiformes. Ad *V. simplicem* Less. valdè accedit. Flores in invol. circ. 3o. An hujus sectionis ? (v. s. in h. Acad. reg. Monac.)

159. *V. SIMPLEX* (Less. in Linnaë 1829. p. 280. 1831. p. 659), caulibus è rhizomate lignoso-tuberoso pluribus simplicibus teretibus striatis pilosis, foliis sessilibus oblongis acutis subintegerrimis suprà pilosis, subtus villosis-lanatis, cymis scorpioideis contractis oligocephalis ebracteatis, capitulis 25-floris, involucri campanulati squamis acuminatis subpungentibus 3-nerviis sericeo-pilosis, achænio brevi densè sericeo-piloso, pappi serie ext. internam subsequante.  $\chi$  in Brasiliâ tropicâ (Less.), in campis et pedunculosis prov. Minar. gener. (Mart.!) et prov. Sancti-Pauli (Lund.). Vern. erigerontis Mart.! herb. Collum superne tuberculorum villosum. Variat caulibus adpressè aut patentim pilosis, foliis margine planis aut revolutis uninerviis angustis aut penninerviis latioribus. Cor. ex siccò pallidè glabræ. Pappus flavescens seu rufescens. (v. s.)

160. *V. CAMPESTRIS*, caule herbaceo erecto simplici confertè hispido, foliis sessilibus lineari-lanceolatis planis subintegerrimis utrinquè confertè villosis, capitulis 1-3 terminalibus, lateralibus breviter pedicellatis, terminali sessili, invol. squamis biserialibus villosis acuminatis, achænio hirtello, pappi serie ext. angustè palacèà brevè.— $\chi$  in campis editis prov. Sancti-Pauli frequens, nov. flor. legit cl. Lund. Affinis *V. simplici*. Pubes confertissima in parte superiore, parca in inferiore. Cor. purpureæ glabræ. Pappus albus vix invol. longior. Folia 1  $\frac{1}{2}$  poll. longa, 3-3  $\frac{1}{2}$  lin. lata. (v. s. comm. à cl. Lund.)

161. *V. PYZOETACHYA*, caule herbaceo erecto simplici villosa, foliis oblongo-lanceolatis utrinquè attenuatis integerrimis villosis, capitulis secus caulis apicem remotè spicatis subsessilibus ad axillas foliorum super. ortis et idè cymam gracilem erectam simplicem constituentibus, invol. 2-3-serialis squamis dorso pubescentibus acuminatis, achænio hirtello, pappi serie ext. subpalacèà brevissima acutà.  $\chi$  in campis editis prov. Sancti-Pauli sat. frequens, nov. fl. legit cl. Lund. Affinis *V. simplici* et *campestris* sed dispos. capitulorum faciliè diversa. Pappus albus, invo. vix superans. (v. s. comm. à cl. Lund.)

162. *V. DENSIVILLOSA* (Mart.! herb.), caule herbaceo erecto simplici densè hirsuto tereti apice compresso, basi patentim folioso, apice ferè aphylo, foliis sessilibus obovatis basi cuneatis apice subobtusis acutivè vix crenatis suprà adpressè sericeo-villosis, subtus et margine densè sericeo-sublanatis, cymis corymbosis confertis aphyllis subscorpioideis, capitulis 20-floris sessilibus, invol. squamis parè villosis acuminatis, cor. glabrâ, recep. nudo, achænio brevi villosa, pappi ser. ext. brevissima paleacèà acuminatâ.— $\chi$  in campis altis ad Sero Frio prov. Minar. general. legit cl. Martins, et in campis editis prov. Sancti-Pauli cl. Lund! Folia 3-3  $\frac{1}{2}$  poll. longa, 18-20 lin. lata. Cor. violaceæ. Pappus albusissimus. Hirsuties tota sericeo-flavicans. (v. s. in h. Acad. reg. Monac.)

*f. angustior*, caule minùs hirsuto, foliis oblongis glanduloso-crenatis magis distantibus.  $\chi$  cum var. *a.* Folia sesquipollicem longa, 6. lin. lata, magis crenata. (v. s. in h. Acad. reg. Monac.)

163. *V. MOLOSERICA* (Mart.! herb.), caule basi suffruticoso erecto simplici tereti pedunculisque villosis-velutinis, foliis sessilibus lanceolato-linearibus subacuminatis integerrimis utrinquè subtus præsertim sericeo-villosis, panicula

terminalis ramis bi-trifidis aphyllis, capitulis pedicellatis 25-30-floris, invol. squamis glabris lanceolato-linearibus longè acuminatis, ext. brevioribus, cor. glabrà, recept. subnudo, achenio densè sericeo-villoso, pappi ser. ext. angustà paleacè acuminatà brevi.  $\frac{5}{8}$  in campis Sancti-Philippi versus Contendos prov. Minas. gener. Brasiliæ legit cl. Martius. Similis V. schwenkianifolia, sed folia diversa 3 poll. longa, 4-lin. lata. Pappus uterque et achenii pili croci. (v. s. in h. Acad. reg. Monac.)

165. V. SCHWENKIANIFOLIA (Mart. herb.) caule basi suffruticoso erecto simpliciteretè densè pedunculisque velutino-villosis, foliis sessilibus ovatis basi obtusis apice mucrone calloso acutis utrinquè densè velutino-villosis subsericosis penninerviis, paniculæ terminalis ramis lateralibus simplicibus aut bifidis aphyllis, capitulis pedicellatis 20-floris, invol. squamis glabrisculis longè acuminatis ext. brevioribus, cor. glabrà, recept. nudo, achenio densè sericeo-villoso, pappi ser. ext. paleacè brevi.  $\frac{5}{8}$  in Brasiliæ campis editis prov. Minas. gener. (Mart.), circa Tejuco (Vanth. pl. exs. n. 258). Pappus int. in sicco croceus, ext. albus. Folia 10-12 lin. longa, 5-6 lin. lata. Caulis sesquipedalis. (v. s.)

166. V. BRUCIOLATA, fruticosa, ramis teretibus hirsutis fuscè densè imbricatè petalis lanatis, foliis petiolatis elliptico-oblongis utrinquè acutis supra subsuperis subrugulosis puberis, subtus reticulatis petiolisque fulvo-tomentosis, paniculæ terminalis ramis patentibus polycephalis bifidis subcorpioides, capitulis sessilibus ovatis 10-12-floris, involucri squamis ovato-lanceolatis obtusiusculis aut mucronato-acuminatis, achenio villosis, pappi serie externà setosa, interior. multò brevior.  $\frac{5}{8}$  in Brasiliæ prov. Sancti-Pauli. Pappus rufescens. Cor. glabrà purpurea. Affinis videtur Vern. ferruginea, sed inflorescentia diversa. Folia subtus non verè tomentosa, 2-2  $\frac{1}{2}$  poll. longa, pollicem lata. Inv. squame ciliatæ glabrescunt. (v. s.)

167. V. SEMILIFOLIA (Less. in linnæ 1831. p. 659), glaberrima, caule herbaceo erecto striato angulato apice nudiusculo, foliis lanceolato-linearibus elongatis sessilibus integerrimis penninerviis nervis utrinquè elevatis, subtus glanduloso-punctatis, capitulis circ. 40-floris longè pedunculatis aphyllis distanter racemosis, invol. campanulati squamis dorso pulverulento-puberulis et ciliatis lanceolatis acuminatis, cor. glabrà, achenio teretiusculo sericeo-villosissimo, pappi ser. ext. paleacè acuminatà achenio sextuplò ferè brevior.  $\frac{7}{8}$  in Brasiliâ. Simillima V. grandiflora, sed invol. et achenia diversa. Projectura 3 è folii cujusque basi secus caulem decurrunt. (v. s. in h. Mus. reg. Paris sec. à Mus. imp. Bras. sub n. 441.)

168. V. RARIFOLIA (Less. in linnæ 1829. p. 285. 1831. p. 659), glabrescens, caulibus imò basi suffruticulosis erectis simplicibus aut à basi ramosis adpressè puberulis apice summo aphyllis sub-1-cephalis, foliis lineari-filiformibus sessilibus acutis sparè puberulis, invol. campanulati ebracteati squamis ovatis apiculato-cuspidatis extus adpressè villosis imbricatis.  $\frac{7}{8}$  in altis campis Brasiliæ (Sellow), inter Santo João d'Elkey et V<sup>na</sup> Thitcam, prov. Minas. gener. legit cl. Martius. V. filifolia Mart. herb. Rhizoma liguosum. Folia 7-9 lin. longa. Herba in sicco nigricans. Cor. 15-20 purpurea. Achenia turbinata sericeo-villosissima. Pappus albidus pluriserialis, setis ext. triplò ferè brevioribus paniculatis. An certè hujus sectionis? (v. s.)

§. 4. *Involucri squamis intimis obtusis, cymis scorpioides, pedunculis ad basin folia perfecta gerentibus.* Less. in linnæ 1831. p. 660. — *Species omnes Americanæ.*

169. V. GRANDIFLORA (Less. in linnæ 1831. p. 660), glaberrima, caule herbaceo erecto simplici aut parè ramoso teretiusculo, foliis linearibus elongatis utrinquè acuminatis sessilibus integerrimis penninerviis utrinquè nervoso-reticulatis, subtus glanduloso-punctatis, capitulis multifloris ad apicem caulis pedunculorumve solitariis aphyllis, invol. campanulati squamis siccis imbricatis, ext. brevissimis acuminatis, medietate latè oblongis mucronatis, int. longioribus apice crocis obtusis, cor. glabrà, achenio striato serièbus 10 pilosis notato, pappi ser.

- ext. paleacea acuminatâ, achenio paulò breviorè.  $\frac{1}{2}$  in campis Brasiliæ (Sellow!), campis editis prov. Sancti Pauli (Lund.!), ad Lorena prov. Sancti Pauli (Mart.!). Vern. campestris Mart.! herb. *Chrysocoma pumila* Arrab. fl. flum. 8. t. 3a. Folia 5-6 poll. longa, 1-5 lin. lata. Capit. diam. pollicari. Cor. violaceo. Achenium sesquilineare. Pappus albidus. Variat caule monocephalo aut racemoso, pedunculis longis distantibus 1-cephalis. (v. s.)
169. *V. SELLOWII* (Less. in *linnæa* 1829. p. 301. 1831. p. 661), caule tereti herbaceo simplici subaphyllo pubescente, foliis ad collum subsessilibus obovato-oblongis subcuneatis subserratis utrinquè subcaberis penninerviis, pedunculis paucis ex axillis solitariis 1-cephalis, capitulis multifloris, invol. latè campanulati squamis oblongis acuminatis internis obtusis mucrone subpungentibus basi 3-nerviis, achenio oblongo multicostato sericeo-piloso, pappi serie ext. paleacea brevi.  $\frac{1}{2}$  in Brasiliâ meridionali. Cor. alba? glabra. Pappus sordidè flavescens. Variat pygmaea et bipedalis, involucrio ferrugineo-villoso aut subglabro. (v. s. comm. à Mus. reg. Berol.)
170. *V. OVATA* (Less. in *linnæa* 1829. p. 294. 1831. p. 661), caule herbaceo tereti striato simplici piloso, pilis sæpè reflexis, foliis brevè petiolatis membranaceis obovatis mucronulatis utrinquè pilosis, cymis scorpioides spicaformibus terminalibus, capitulis multifloris sessilibus, invol. campanulati squamis siccis 1-nerviis ciliatis, externis pungenti-acuminatis, internis obtusis, achenio piloso, pappi serie ext. brevi. (1?) in Brasiliâ ad Paraíba. Caulis pedalis. Cor. glabra violacea. Pappus argenteus. Folia 15-18 lin. longa, 6-10 lata.
171. *V. HYPOCISTIS*, caule herbaceo à collo fruticoso erecto ferè aphylo tereti pubescente, foliis radicalibus petalis sessilibus obovatis obtusis margine subcrenatis penninerviis suprà glabris subtus puberulis, caulinis oblongo-linearibus strictè adpressis crenulatis paucis, capitulis ad apicem geminis inferiore longius pedunculato ex axillâ folii supremi orto, invol. campanulati squamis ext. lanceolatis acuminatis pubescentibus, intimis oblongis subobtusis glabrisculis subsaricosis, achenio sericeo-villoso, pappi ser. ext. paleacea brevi.  $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Affinis *V. Sellowii*. Caulis pedalis. Fol. radic. 3 poll. longa, 2 poll. lata. Caulina pollicem longa, 1-2 lin. lata. Cor. purpurea. Invol. et caulis purpurascens. (v. s. in h. Mus. reg. Paris. ex h. Mus. imp. Bras. n. 1025.)
172. *V. GLABRATA* (Less. in *linnæa* 1829. p. 294. 1831. p. 661), caule herbaceo simplici erecto supernè ramoso tereti striato subglabro, foliis brevissimè petiolatis rigidè lanceolatis elongatis basi subobtusis subdenticulatis reticulatis utrinquè glabris, cymis scorpioides terminalibus longis, capitulis sessilibus approximatis 20-25-floris, invol. campanulati squamis siccis 1-nerviis, exter. mucronato-acuminatis, intern. apice scariosis latis obtusis, achenio glabro, pappi serie ext. angustè paleacea brevi.  $\frac{1}{2}$  in Brasiliâ tropicâ. Caulis 2-4-pedalis. Cor. glabra purpurea. Pappus pallescens. (v. s. comm. à Mus. reg. Berol.)
173. *V. SUBULNIFOLIA* (Mart.! herb.), fruticosa, ramis teretibus lanato-tomentosis, foliis brevissimè petiolatis elliptico-oblongis utrinquè subacuminatis crenatis, suprà glabris venis impressis, subtus cano-tomentosis penninerviis, capitulis secus ramorum apicem sessilibus aut pedunculatis paucis extrà-axillaribus ferè 60-floris, invol. subglobosè squamis coriaceis imbricatis adpressis obtusissimis dorso aranaceis, ext. ovatis, int. oblongis, cor. glabrâ, achenio tereti villoso, pappi ser. ext. paleacea brevi.  $\frac{1}{2}$  in campis ad flum. Rio-Verde prov. Minarum general. Brasiliæ legit cl. Martius. Flores rosei. Folia 5-6 poll. longa, sesquipollicem lata, superiora, 9-10 lin. lata. (v. s. in h. Acad. reg. Monac.)
174. *V. LIQUIDAFOLIA* (Mart.! herb.), fruticosa, ramis teretibus striatis glabresculis, foliis sessilibus angustè ligulatis utrinquè subobtusis integerrimis suprà glabris venoso-reticulatis, subtus cano-tomentosis penninerviis, capitulis ad apicem ramorum spicatis in axillis foliorum sessilibus sæpius geminis folio triplo brevioribus 25-30-floris, invol. ovatis squamis adpressis coriaceis glabris

- obtusis, ext. subtriangularibus brevibus, int. longis lato-linearibus, cor. glabrâ glandulosâ, achenio 10-costato basi sparsè piloso apice glabro, pappi ser. ext. paleaceâ brevi.  $\zeta$  in editis campis prov. Minarum general. Cor. purpurea. Folia 5-6 poll. longa, 9 lin. lata. (v. s. in h. Acad. reg. Monac.)
175. *V. CORIACEA* (Less. in linnæâ 1831. p. 661), caulibus è collo lignoso pluribus erectis teretibus simplicibus supernè albo-tomentosis, foliis brevissimè petiolatis coriaceis penninerviis suprâ nitidis, subtus tomentoso-incanis linearibus integerrimis, cymâ scorpioidèa spiciformi, capitulis multifloris folio flor. multò brevioribus, invol. semiglobosi squamis siccis vix subtomentosis ext. rotundis acutis, intum. linearibus obtusis, cor. glabrâ, achenio nitido sulcato, pappi ser. ext. paleaceâ brevi. — Patr. ignot. an Brasiliâ? Cor. purpureo-violaceo. Capitula *V. macrocephala*. Folia 3-6 poll. longa, 4-6 lin. lata.
176. *V. RUFO-GRISCA* (S. Hil. voy. diam. 2. p. 305 et 453), caule suffruticoso tereti, ramis supernè striato-angulatis rufo-griseove tomentosis, foliis petiolatis subulato-lanceolatis brevissimè mucronulatis basi acutis suprâ pubescenti-tomentosis subtus sericeis, cymis terminalibus bipartitis folio basilari brevioribus patentibus, capitulis secundis subsessilibus fol. flor. brevi brevioribus, invol. squamis pubescentibus acutis mucronulatis, achenio hirsuto villosa, pappi ser. ext. angustè paleaceâ brevi.  $\zeta$  in Brasiliâ inter Almeida et Rio-Doce prope Quartal do Riacho. Cor. dilutè purpurea. Pappus subrufescens.
177. *V. RUBRICAULIS* (H. et Bonpl. pl. equinox. 2. p. 66. t. 99), caule herbaceo sulcato-angulato pubescente simplici apice corymboso-paniculato, foliis sessilibus linearibus integerrimis marginè revolutis suprâ scabris, subtus sericeo-tomentosis, capitulis secus ramos ad axillas sessilibus solitariis unilateralibus 20-25-floris, invol. campanulati squamis imbricatis ovatis acutis dorso pubescentibus, acheniis glabris, pappi ser. ext. paleis lineari-subulatis brevibus.  $\zeta$  in siccis montosis Novo-Granatensium prope Ybague H. B. et Kunth nov. gen. am. 4. p. 33. Less. in linnæâ 1831. p. 66a. *V. rubricaulis* Bonplandiana, Less. in linnæâ 1829. p. 299, excl. var.  $\gamma$  et  $\delta$ . Cor. purpurea glabra. Caulis ruber. — Radem fortè ex Brasiliæ prov. Rio-Grande adest in h. Mus. reg. Par. ex Mus. imp. Bras. (ubi v. a.)
178. *V. OBTUSATA* (Less. in linnæâ 1831. p. 662), herbacea glaberrima erecta simplex, caule tereti striato, foliis petiolulatis rigidulis subintegerrimis oblongo-oboovatis basi acuminatis levibus penninerviis reticulatis, cymis scorpioides, capitulis sessilibus ad folii valdè obtusati axillam sæpiùs geminis 10-floris, invol. cylindracei squamis siccis extus subpilosis nigro-ciliatis ovatis apiculatis, intimis obtusis, cor. glabrâ, achenio hirsuto, pappi ser. ext. paleaceâ brevi. — In Brasiliâ. Cor. purpureo-violaceo. Pappus sordidus. Fol. 4 poll. longa, 15 lin. lata.
179. *V. SERICEA* (Rich. act. soc. h. nat. Par. 105), caule suffruticoso tereti striato tenuissimè puberulo, foliis brevissimè petiolatis lanceolatis acuminatis basi subobtusis subintegerrimis suprâ glabrisculis subtus adpressè pubescentibus, capitulis ad axillas sessilibus solitariis geminisve circ. 50-floris, invol. campanulati squamis glabrisculis imbricatis, ext. acumine recurvo mucronatis, intimis longioribus obtusis, achenio sericeo-villosa, pappi ser. ext. angustè paleaceâ brevi.  $\zeta$  in Brasiliâ, circa Rio-Janeiro (Gaudich. ! Lund. !), in monte Corcovado et in Serra de Estrella (Mart. !), et in Guiana Pers. ench. 2. p. 404. Kar bot. reg. t. 522. Less. in linnæâ 1829. p. 295. 1831. p. 663. Verna. imponens Mart. ! herb. Pappus uterque albusimus. Cor. glabriscula pallidè purpurea aut ex Rich. alba. Folia 3 poll. longa, 9-10 lin. lata. Petioli sesquilineares! (v. a.)
180. *V. FRASCFOLIA* (Desf. ! cat. h. Par. 1829. p. 400), caule fruticoso tereti striato flexuoso, foliis sessilibus lanceolatis longè acuminatis margine retrorsam asperis subintegerrimis deflexis nitidis, subtus subpubescentibus, capitulis axillaribus sessilibus solitariis circ. 30-floris, invol. teretis obovati squamis multiseriatis ovatis adpressis ciliatis, ext. brevibus mucrone rigido non adpresso recurvo superatis, intimis lineari-oblongis obtusis, pappi serie externâ paleaceâ brevi.  $\zeta$  in Indiâ (verosim. occidentali). Cor. dilutè rosea. Fol. 3-5 poll.

longa, 5-6 lin. basi lata. Pappus ext. fuscescens, int. setiformis albus. (v. s. in h. Deaf.)

181. *V. ACUTIFOLIA* (Hook. bot. mag. t. 3062), caule herbaceo erecto glabro simplici tereti apice in ramos angulatos pubescentes diviso, foliis sessilibus linearibus acuminatis nervoso-reticulatis margine hinc indè subserratis suprâ glabris subtus obscure pilosis, capitulis 40-floris secus ramos subsessilibus lateralibus axillaribus, invol. hemisphærici squamis imbricatis glabrisculis, ext. brevibus acutis-subsquarrosis, int. subscariosis obtusis, achenio pilos pluriseriales gerente, pappi ser. ext. brevi paleaceâ.  $\frac{1}{2}$  in Brasiliâ ad Rio-Janeiro. — Differt à Vern. grandiflorâ, capitulis plurimis dimidio minoribus ebracteatis, foliis nec integerrimis nec omninò glabris; à V. rubricauli, capitulis 40 nec 20-25-floris, achenio pubigero, foliis subtus glabrisculis, etc.

*ß. ambigua*, foliis suprâ junioribus scabridis demùm glabris subtus glabrisculis punctatis, capitulis extrâ-axillaribus.  $\frac{1}{2}$  in Brasiliâ. An Vern. rubricaulis latifolia Less. in linnæâ 1829. p. 299? (v. s. comm. sine nom. à Mus. reg. Berol.)

182. *V. SCHIEDIANA* (Less. in linnæâ 1831. p. 399 et 663), fruticosa scandens, ramis teretibus pubescenti-canescens, foliis petiolulatis sublancoolato-ellipticis membranaceis penninerviis basi obtusis subintegerrimis suprâ glabris subtus pilosis, cymâ scorpioideâ divergente, capitulis sessilibus 25-floris folio stipatis, invol. campanulati squamis erectis ciliatis extus pilosis, ext. lancoolatis mucronatis, int. oblongis obtusis, cor. glabrâ, achenio sericeo-villoso, pappi ser. ext. paleaceâ brevi.  $\frac{3}{4}$  in dumetis Mexici ad Papantla et Misantla. Folia 4-4 $\frac{1}{2}$  poll. longa, 1 $\frac{1}{2}$ -2 $\frac{1}{2}$  poll. lata; petiolo 3-4 lin. Affinis dicitur V. sericea.
183. *V. OBOVATA* (Less. in linnæâ 1829. p. 296. 1831. p. 663), caule fruticoso, ramis teretibus striatis fusco-villosis, foliis brevissimè petiolulatis lanceolatis coriaceis subdenticulatis suprâ rugosis scaberrimis subtus flavescens villosis tomentosis, cymis scorpioideis longis terminalibus, capitulis multifloris sessilibus approximatis folio florali multò brevioribus, invol. subcylindracei squamis 1-nerviis ciliatis, intimis obtusissimis, achenio multicostato inter costas hirsuto, pappi serie ext. paleaceâ brevi.  $\frac{3}{4}$  in Brasiliâ tropicâ. Cor. albæ? glabræ. Pappus flavescens.
184. *V. AXILLIFLORA* (Mart. l. herb. et hort. Mon. ex Less. in linnæâ 1829. p. 297. 1831. p. 663), caule fruticoso, ramis teretibus villosis, foliis brevè petiolulatis oblongo-ellipticis utrinquè acutis subintegerrimis suprâ scabris pilosis subtus villosis tomentosis, cymis scorpioideis terminalibus longis, capitulis sessilibus folio florali multò brevioribus, invol. campanulati squamis 1-nerviis obtusissimis subpungenti-mucronatis ciliatis, cor. lobis hirtis, achenio sericeo-piloso, pappi serie ext. brevi subpaleaceâ.  $\frac{3}{4}$  in Brasiliâ ad Rio de S. Francisco prope Salgado prov. Minar. general. legit cl. Martius. Lodd. bot. cab. t. 1690. Cor. purpureo-violaceæ glabræ. Pappus argenteus. (v. s. in h. Acad. reg. Monac.)
185. *V. COTONEASTER* (Less. in linnæâ 1829. p. 298. 1831. p. 663), caule basi suffruticuloso tereti striato apice laxè subtomentoso, foliis brevissimè petiolulatis ovatis integerrimis suprâ scabrisculis demùm glabris subtus villosis tomentosis albis, floralibus ovato-cordatis, cymis longis simplicibus, capitulis ad axillas foliorum sessilibus solitariis rariis geminis, invol. squamis oblongis obtusis mucronatis laxè pilosis, achenio oblongo-villoso multicostato, pappi serie ext. paleis brevibus latioribus.  $\frac{3}{4}$  ad Bahiam Brasiliæ. (Salzm. l. Blanch. l.) Conyza cotoneaster Willd. in Spreng. syst. 3. p. 509. Invol. squamæ membran. sicce purpureæ. Cor. purpurea glabra. Pappus argenteus. (v. s. comm. à cl. Salzm. et Lhotsky.)
186. *V. ACUMINATA* (Less. in linnæâ 1831. p. 663), fruticosa, ramis divaricatis teretibus pubescentibus, foliis petiolulatis ovato-lanceolatis utrinquè acuminatis integerrimis scabris subtus punctatis, cymis scorpioideis brevibus polycephalis, capitulis 15-floris sessilibus approximatis folio flor. brevioribus, invol. squamis extus subpilosis, ext. minutissimis acuminatis, int. oblongis vix acutis, achenio multicostato piloso, pappi ser. ext. paleaceâ brevi.  $\frac{3}{4}$  in insulis Cari-



- basis? Pappus flavescens. Recept. glabrum. Folia 1-2  $\frac{1}{2}$  poll. longa, 4-15 lin. lata. Petiolo 2 lin. Invol. in descr. nunc campanulatum nunc turbinatum dicitur.
187. *V. CHAMÆDRYS* (Less. in linnæ 1829. p. 259. 1831. p. 641), caule fruticoso inferne glabro, ramis teretibus cano-tomentosis, foliis sessilibus oblongis subrotundisve margine subreflexo parè serratis, suprà glabris luddis impressè glandulosis, subtis albo-tomentosis, cymis ramis simplicibus ramosisve, capitulis sessilibus axillaribus et extrà-axillaribus solitariis fasciculatisve cylindraceis 8-floris, invol. squamis 3-nerviis ovato-oblongis vix subtomentosis, achenio hirsuto glanduloso, pappi serie externà paleacea brevi.  $\frac{3}{4}$  in Brasiliâ tropicâ (Less.), ad Bahiam (Blanch.). Cor. purpureo-violacea glabra. (v. s.)
188. *V. AURIFODINARUM* (Mart. herb.), suffruticosa?, caule erecto tereti striato pubescenti-villoso, foliis petiolatis elliptico-oblongis utrinquè acuminatis parè dentatis, suprà muriculatis secus nervos subvelutinis, subtis pubescenti-villosis, capitulis ad axillas super. subsessilibus folio multò brevioribus solitariis 15-20-floris, invol. ovatis squamis adpressis ciliatis dorso glabrescentibus, ext. mucronatis, int. obtusis, corollâ glabrâ, achenio piloso striato, pappi ser. ext. brevi paleacè.  $\frac{3}{4}$  in campis prope Congonharde Sabara prov. Minar. gener. legit. cl. Martins. Aff. *V. lilacina*. Petioli 8-9 lin. longi; fol. limbus 4-6 poll. longus, 18-20. lin. latus. Herba obscurè virens. (v. s. in h. Acad. reg. Monac.)
189. *V. LILACINA* (Mart. herb.), fruticosa, ramis teretibus demum glabris, ramulis subangulatis pubescenti-villosis, foliis breviter petiolatis ovalibus utrinquè subobtusis integerrimis suprà tenuissimè pubescentibus, subtis pubescenti-villosis, ramis simplicibus bifidisve, capitulis ad dichotomias ram. aut axillas foliorum sessilibus solitariis rarè geminis 10-12-floris, inv. ovatis squamis oblongis adpressis dorso apice puberis obtusis aut brevissimè mucronatis, achenio piloso turbinato, pappi ser. ext. brevi angustissimè paleacè.  $\frac{3}{4}$  in campis prope Tejucco prov. Minar. gener. Brasiliæ. Cor. lilacina apice parè pubera. Pappus rufescens vix invol. longior. Recept. alveolatum. Folia 2 poll. longa, 10 lin. lata, petiolo 1-2 lin. (v. s. in h. Acad. reg. Monac.)
- §. 5. *Involucris squamis intimis acuminatis, cymis scorpioidis folia perfecta ad basin pedunculorum gerentibus.* Less. in linnæ 1831. p. 664. — *Species omnes Americana.*
190. *V. ARBUSCULA* (Less. in linnæ 1831. p. 664), caule sarbarthoreo, ramis teretibus villo brevissimo incanis, foliis petiolulatis obovatis penninerviis suprà pilosis, subtis villoso-incanis, cymis scorpioidis, capitulis sessilibus foliis ensis superatis, invol. turbinatis squamis acuminatis inermibus effatis dorso villosis, achenio villoso, pappi ser. ext. paleacè brevi.  $\frac{3}{4}$  in insulâ Mauritiæ ex h. cl. Kanth; an fortè in horto bot. collecta? Affinis videtur *V. arborescens*.
191. *V. ARBORESCENS* (Swartz fl. ind. occ. 2. p. 130), fruticosa, ramis teretibus villosis, foliis brevè petiolatis oblongo-ovatis acuminatis sæpè deflexis subintegerrimis, suprà pilosis subtis piloso-canescensibus, cymis scorpioides terminalis ramis divaricatis, capitulis sessilibus axillaribus folio florali brevioribus, invol. campanulatis squamis acuminatis, achenio sericeo-piloso, pappi serie ext. brevi.  $\frac{3}{4}$  in montibus ina. Caribæarum Portorici, Sanctæ-Crucis. Less. in linnæ 1829. p. 302. 1831. 665. *Lepidaploa phyllostachya* Cass. dict. 26. p. 16. *Conyza arborescens* Linn. sp. 1209. — Plum. am. ed. Barm. t. 130. f. 2. Cor. glabra purpureo-carulea nec flavæ ut vult Cass. Recept. subalveolatum. (v. s.)
- A. ovatifolia*, foliis ovatis acuminatis, suprà subrugosis, subtis pubescentibus paulò longius petiolatis.  $\frac{3}{4}$  in ina. Canâ juxta Havanæ et in Guadalupâ. *Eupatorium secundiflorum* Bert. herb. non Lag. An fortè *V. fruticosa* Lin.? sed folia acuminata nec obtusa. (v. s.)
192. *V. DIVARICATA* (Swartz fl. ind. occ. 3. p. 1319. non Less.), fruticosa,

ramis teretibus pubescentibus, foliis breviter petiolatis membranaceis ovali-oblongis utrinque acuminatis integerrimis supra et imprimis infra minutissimè pubescentibus, cymis axill. et terminalibus bifidis subscorpioideis, capitulis sessilibus nunc nodis nunc bracteis suis brevioribus 11-12-floris, invol. ovati squamis oblongis obtusis dorso parèe puberis, achenio turbinato sericeo-villoso, pappi serie ext. brevi vix paleacea.  $\frac{3}{4}$  in ins. Caribeis. *Conyza arborescens* Swartz! in h. L'Hér., sed antè fl. Ind. occ. script. *V. acuminata* Less. in LINNÆ 1831. p. 663? *V. divaricata* Spreng. syst. 3. p. 435 excl. syn. Cor. purpurea glabra. Pappus sordidè flavescenti-griseus. (v. s.)

193. *V. ICOSANTHA*, fruticosa, foliis petiolulatis oblongo-ovatis basi obtusis aut acutis membranaceis, subius petiolique villosis pellucido-glandulosis, cymis terminalibus scorpioideis oligocephalis, capitulis sessilibus alaribus aut extrà axillaribus folia sua superantibus 20-floris, invol. squamis oblongis adpressis macronatis interioribus acuminatis, achenio turbinato densè piloso, pappi serie ext. paleacea brevi.  $\frac{3}{4}$  in Martinicâ et fortè in Brasiliâ. *V. divaricata* Less. in LINNÆ 1829. p. 306. 1831. p. 665. excl. syn. Pappus argenteus. Invol. à cl. Less. nunc cylindraceum nunc campanulatum dictum.

194. *V. AZARUA* (Swartz! fl. Ind. occ. 1322), suffruticosa, ramis teretibus densè petiolique pubescenti-velutinis, foliis petiolatis ovato-subrotundis utrinquè obtusis integerrimis supra glabris subluccidis subtus glandulosis secus nervos velutinis, cymâ terminali subflexuosa oligocephalâ, capitulis brevè pedunculatis folia floralia ovata subaequantibus, invol. squamis ext. carinatis subvillosis, intimis linearibus subacuminatis, achenio obconico biapido, pappi serie latè paleacea brevi.  $\frac{3}{4}$  in montibus calcareis petrosis Jamaicæ borealis. *Conyza rigida* Sw. prod. 113. Cor. glabra purpurea. Pappus sordidè griseo-stramineus. An ipso medente Swartzio à *V. fruticosa* satis differt? Folia rotundiora et cyma multò magis contracta. (v. s.)

195. *V. VASELIANA* (Less. l. c. p. 306. 1831. p. 306), fruticosa, ramis teretibus striatis pube adpressâ sericeâ albidis, foliis brevè petiolatis membranaceis integerrimis ovato-oblongis utrinquè minutè glandulosis et pubescentibus demum glabris, cymis scorpioideis brevibus oligocephalis, capitulis subsessilibus approximatis, invol. cylindracei squamis acuminatis 1-nerviis demum patentibus, achenio oblongo densè sericeo-piloso, pappi serie ext. paleacea brevi.  $\frac{3}{4}$  in insulâ Sanctæ-Graciæ. *Conyza glabra* Willd. sp. 3. p. 1940. et *Eupatorium obtusifolium* Willd. sp. 3. p. 1768. ex Less. *V. albicaulis* Pers. syn. 2. p. 404. Cor. glabræ violaceæ. Pappus argenteus. (v. s. comm. à cl. Puerari, qui à cl. West recuperat.)

196. *V. LONGIFOLIA* (Pers. ench. 2. p. 404), caule suffruticoso? tereti petiolis pedunculisque pubescentibus, foliis brevè petiolatis oblongo-lanceolatis basi brevè apice longè acuminatis membranaceis glabris integerrimis utrinquè minutè glandulosis, cymis axillaribus et terminalibus folio brevioribus bifidis oligocephalis, capitulis folio oblongo-lineari brevioribus brevè pedicellatis, invol. turbinati squamis acuminatis, achenio sericeo-villoso, pappi serie ext. paleacea brevi.  $\frac{3}{4}$  in Guadalupâ. *V. punctata* Swartz in Wikstr. guad. fl. 72. an ex ipso auctore *Lepidaplos lanceolata* Cass. dict. 26. p. 18, sed cor. certè in nostrâ non sunt flavæ! Folia 4-5 poll. longa, 9-14 lin. lata. Pappus argenteus. Cor. glabræ sicco purpureæ. (v. s. comm. à cl. Krauss.)

197. *V. OSTRIFOLIA* (Less. l. c. p. 308), fruticosa, ramis teretibus glabris, junioribus pilosisculis, foliis petiolatis carnosio-coriaceis ellipticis obtusis basi acutis integerrimis supra impresso-glandulosis utrinquè pilosisculis, cymis scorpioideis brevibus terminalibus oligocephalis, capitulis sessilibus multifloris subapproximatis folii floralis longitudine, invol. campanulati squamis oblongo-ovatis subacuminatis 3-nerviis, achenio sericeo-piloso, pappi serie ext. paleacea brevi.  $\frac{3}{4}$  in Brasiliâ tropicâ. Recept. paleaceo-fimbriiferum. Pappus argenteus. Fol. 1-1  $\frac{1}{2}$  poll. longa, 4-6 lin. lata.

198. *V. ENACLES* (H. B. et Kunth nov. gen. am. 4. p. 27), herbacea, ramis teretibus glabris aut subtomentosis, foliis brevissimè petiolatis oblongis acu-

minatis subintegerrimis puberulo-scabriusculis sæpè reflexis, cymis scorpioideis terminalibus longis divaricatis, capitulis sessilibus remotis in axillis foliorum solitariis conglomeratisve, invol. subtrunati squamis acuminatis pilosis demum patentibus, achenio inter costas piloso, pappi ser. ext. brevi.  $\frac{1}{2}$  in umbrosis prope Turbaco Novo-Granatensium et in Brasiliâ tropicâ. Less. in linnæâ 1829. p. 302. 1831. p. 667. V. elongata Willd. herb. ex Less. Cor. violaceæ glabræ.

*l. villosa* (Less. l. c.) foliis subtus subtomentoso-villosis, capitulis magis conglomeratis.

199. V. STENOPHYLLA (Less. in linnæâ 1831. p. 667), caule è rhizomate lignoso erecto herbaceo tereti supernè albo-villosissimo simplici, foliis petiolulatis angustè linearibus aveniis acutis margine revolutis suprâ pilosis, subtus sericeo-villosis, capitulis secus ramos paniculæ in axillis foliorum sessilibus et iis brevioribus, invol. campanulati squamis linearibus mucronatis extus villosis, cor. glabrâ, achenio villosa, pappi ser. ext. paleaceâ brevi.  $\frac{1}{2}$  in Sancto-Domingo. Folia  $1\frac{1}{2}$ -2  $\frac{1}{2}$  poll. longa, lineam lata.

200. V. MACROPHYLLA (Less. in linnæâ 1831. p. 668), caule herbaceo altissimo tereti striato villosa-pubescente apice ramoso et foliato, foliis petiolatis membranaceis basi obtusissimis inf. ovatis acuminatis ext. lanceolato-ellipticis penninerviis suprâ hispidiusculis subtus molliter pubescentibus et punctatis, cymis scorpioideis divaricatis, capitulis sessilibus remotis solitariis plerisque folio suffultis, invol. latè campanulati squamis multinerviis, ext. ovatis acutis, med. obtusis, intimis acuminatis, cor. glabrâ, achenio villosa, pappi ser. ext. paleaceâ brevi.  $\frac{1}{2}$  in Brasiliâ prov. Rio-Janeiro in Serra do Mar. Folia 17-18 poll. longa, 9-11 lata. Herba dendromorpha 20-pedalis. Cor. violaceæ. Pappus sordidè flavescens.

201. V. MULTIFLOSCULOSA (Mart. l. herb.) suffruticosa, ramis teretibus pedunculisque pube brevissimâ velutinis, foliis breviter petiolatis ovatis basi subcuneatis apice acuminatis callosa-serratis suprâ glabresculis subtus canescenti-subtomentosis, paniculæ ramis inf. axillaribus folio aequalibus, sup. brevioribus subscorpioideis, capitulis maximis 50-floris subsessilibus aliis ebracteatis aliis bracteâ oblongâ donatis, invol. squamis ovato-oblongis mucronatis adpressis ciliatis, intimis acuminatis, achenio villosa-sericeo, pappi ser. ext. brevi paleaceâ.  $\frac{1}{2}$   $\frac{3}{4}$  in Brasiliâ umbrosis et sylvæ caduis prope Rio-Janeiro (Blanch.!), ad montem Cercovado (Mart.!), in siccis prope Rage (Lhostky!), Chrysocoma pedunculata Arrab. fl. flum. 8. t. 37. Affinis V. macrocephale. Invol. squamæ dorso vix pubescentes. Pappus sordidè rufescens. Folia 8 poll. longa, 4-5 poll. lata. Variat foliis subtus subalbidis aut virescenti-pubescentibus. An eadem ac V. macrophylla? (v. s.)

202. V. HELOPHYLLA (Mart. l. herb.), caulibus basi suffruticulosis erectis simplicibus teretibus villosa-hirsutis apice glabresculis, foliis brevissimè petiolatis ovatis basi obtusis apice acutis mucrone calloso integerrimis suprâ sparè pilosis demum scabridis subtus glandulosis villosis, ramis axillaribus distantibus simplicibus bifidiæve, capitulis sessilibus ad axillam folii ovato-lanceolati distantibus solitariis 20-floris, invol. squamis omnibus acuminatis, ext. brevioribus dorso pubescentibus, cor. glabris, achenio villosa, pappi ser. ext. paleaceâ brevi.  $\frac{1}{2}$  in paludosis montanis prov. Minarum General. prope Villa-Ricca leg. cl. Martius. Caules rubentes. Cor. albæ. Folia poll. longa, 6 lin. lata, petiolo lineam longo. (v. s.)

203. V. GERMIFATA (Less. in linnæâ 1829. p. 303. excl. syn. ex linnæâ 1831. p. 669), herbacea volubilis, ramis teretibus obscure villosis, foliis brevè petiolatis oblongo-ovatis acutis undulato-subrenulatis suprâ scabris rugosis subpilosis subtus obscure villosis, cymæ scorpioideæ terminalis ramis longis, summis geminis, capitulis sæpius solitariis 20-floris approximatis sessilibus folia floralia subtus cordata ferè superantibus, invol. campanulati squamis in acumen setiforme patulum productis intimis linearibus rectis, achenio sericeo-piloso; pappi serie ext. brevi paleaceâ. in Brasiliâ prov. Rio-Janeiro et in Serra d'Ka-

trella. Recept. alveolare. Cor. albæ glabræ. Pappus albus. Spreng. syst. 3. p. 437. excl. syn. (v. s. comm. à Mus. reg. Berol.)

β. *aprica*, caule erecto ad apicem magis villosa, foliis oblongis magis acuminatis subtus densius et mollius villosis, cymæ ramis brevioribus, invol. squamis pallidis dorso omnibus molliter villosis. ♀ in locis apricis vulgarissima circa Rio-Janeiro ex cl. Lund. Herba 1-2-pedalis. Cor. albæ. Capitula paulò minora quàm var. α. An species propria? (v. s. comm. à cl. Lund.)

204. V. *BANDANOIDES* (Less. in linnæa 1831. p. 669), caule herbaceo-erecto tereti striato apice hirsuto-viloso, foliis subsessilibus subrotundis ellipticisve basi subcordatis mucronato-acuminatis reticulato-venosis asperimis suprà lucidis subtus pilosis, cymæ scorpioideæ pauciradiatâ polycephalâ, capitulis sessilibus foliis binis stipatis, invol. campanulatis squamis acuminatis rigidis-extrâ pilosis, corollâ glabrâ, achenio . . . pappi ser. ext. paleaceâ brevi. — in Brasiliâ. Folia 2-3 poll. longa,  $\frac{1}{2}$ -2 lata. Cor. purpureo-violaceæ. Pappus argenteus. Affinis dicitur V. geminata.

205. V. *CHAMISONIS* (Less. in linn. 1829. p. 304. 1831. p. 670), herbacea, ramis teretibus albo-villosa-tomentosis, foliis breviter petiolatis rigidis ovato-lanceolatis acuminatis basi acutis integerrimis suprà rugosis pilosis subtus sericeo-incanis, cymis scorpioideis terminalibus longis, capitulis distantibus sessilibus folio florali sublongioribus multifloris, invol. campanulatis squamis ciliatis 1-nerviis ciliatis pilosis longè acuminatis ext. subpatentibus subpungentibus, cor. glabrâ, achenio densè piloso, pappi seriebus inæqualibus. ♀ in Brasiliâ ad fretum: Senæ-Catharinæ. Cor. purpureo-violaceæ. Folia 2-3 poll. longa, 9-14 lin. lata.

206. V. *TOMENTOSA* (Nees amoen. 2. p. 11. t. 3. non Kl.), caule basi frutescente simplici virgato supernè villosa-tomentoso, foliis oblongo-lanceolatis brevè petiolatis integerrimis utrinquè molliter tomentoso-villosis, capitulis 12-floris corymboso-paniculatis, cymis subscorpioideis, invol. squamis linear-lanceolatis acutis, interioribus apice coloratis, achenio obconico viloso, pappi serie ext. brevissima paleaceâ. ♀ in Indiâ or. ex sem. hort. Less. in linn. 1829. p. 305. 1831. p. 670. Cor. glabræ purpurascens, in invol. teretiusculo. Pappus argenteus.

207. V. *MEGAPOTAMICA* (Spreng. syst. 3. p. 437), caule herbaceo simplici striato tereti glaberrimo, foliis sessilibus suprà glabris nitidis subtus albo-tomentosis oblongis inferioribus basi attenuatis superioribus basi dilatatis summis cordato-lanceolatis subangulatis, cymis scorpioideis in paniculam dispositis, capitulis cylindricis 6-floris, invol. foliolis ovato-oblongis mucronato-acutis erectis albido-sericeo-pilosis, achenio obconico multicoctato piloso, pappi serie ext. ternâ paleaceâ brevi. ♀ in Brasiliâ meridionali ad Rio Grande. Less. in linnæa 1829. p. 308. 1831. p. 670. Cor. violaceæ glabræ. Pappus argenteus. Rhizoma repens. (v. s. comm. à Mus. reg. Berol.)

β. *brevifolia*, caule ad apicem foliis subtus et involucri albo-lanatis, foliis ellipticis semi-pollicem circ. longis. — Cum var. α.

γ. *melanotrichia*, caule lanâ nigricante hirsuto, fol. nervo medio villis fuscis insignito, invol. squamis ciliis nigris hispidis, pappi interioris setis nigris. ♀ in Brasiliæ prov. Sancti-Pauli in campis ad Lorena. Vern. Candolliana Mart. herb. Var. α simillima, sed colore nigro pilorum ferè omnium etiam pappi inter. distinctæ. (v. s. in h. Acad. reg. Monac.)

208. V. *APICULATA* (Mart. l. herb.), caule herbaceo erecto tereti parè ramo pubescente subhirsuto, foliis sessilibus erectis lanceolatis margine revolutis basi cordato-obtusis apice apiculatis suprà pilosis demum scabris subtus tomentosis, capitulis secus ramos sessilibus spicatis axillaribus folio florali paulò longioribus 4-floris, invol. cylindrici squamis dorso pubescentibus apice longè acuminatis, achenio adpressè pubero, pappi serie ext. setiformi paucisetâ int. triplò brevior. ☉ in campis Brasiliæ prope Barreiros Minarum Novarum legit cl. Martius. Herba sesquipedalis. Pili paginæ fol. sup. decidui basi tuberculiformes. Accedere videtur ad V. Megapotamicam. (v. s. h. Acad. reg. Monac.)

209. V. *REMOTIFLORA* (Rich. act. soc. h. n. Par. 1792. p. 117), herbacea, caule

tereti adpressè pubescente, foliis brevissimè petiolatis ovali-lanceolatis membranaceis serratis basi cuneatis apice acutis suprà pubescentibus, subtus sparsè villosis, cymis subpaniculatis ramis subscorpioideis hinc indè foliiferis, capitulis 12-15-floris sessilibus distantibus, invol. ovati squamis sublinearibus acuminatis fructus superantibus extus villosis, achenio sericeo-villoso, pappi serie ext. paleacea brevissimâ. ☉ in Cayennâ et Brasiliâ. Less. in linnæâ 1831. p. 670. Pers. ench. 2. p. 404. V. sessiliflora Willd. herb. ex Less. in linnæâ 1829. p. 309. Habitus ferè V. canescens sed pubescens nec canescens, capitula angustiora longiora hinc indè è foliorum axillis orta. Pappus argenteus. (v. s.)

210. V. *BARTZIANA*, caule herbaceo erecto ramoso tereti villosa, foliis sessilibus oblongis utrinquè subattenuatis apice macronatis integerrimis suprà setis brevissimis scabris, subtus villis adpressis subsericeis canescentibus, ramis scorpioideis, capitulis ad axillas foliorum sessilibus solitariis folio brevioribus 10-floris, invol. squamis linear-lanceolatis acuminatis glabris aculis subnitidis fructus æquantibus, cor. glabrâ, achenio sericeo-subvillosa, pappi ser. ext. paleacea brevissimâ. ☉ in insulis Porto-Ricci et Sancti-Thomasi locis aridis legit cl. Bertero. Conyza Portoricensis Bert.! herb. Pappo rufescente involucentis æquante et paginâ fol. breviter scabrâ differt à V. remotiflorâ. An varietas? (v. s.)

211. V. *VERBASCIFOLIA* (Less. in linnæâ 1829. p. 310. 1831. p. 670), caule herbaceo simplici tereti sulcato tomentoso-villoso, foliis sessilibus confertis rigidis utrinquè scabris et villosis obovatis subintegerrimis, cymis scorpioideis terminalibus subfastigiatis oligocephalis, capitulis multifloris sessilibus remotis folio florali ovato acuto longioribus, invol. campanulati pauciserialis squamis acuminatis 3-nerviis, cor. glabrâ, achenio sericeo-villoso, pappi ser. ext. paleacea interiori parum brevior. ♀ in Brasiliâ tropicâ. Cor violaceæ. Pappus argenteus. Folia 2-5 poll. longa, 1-2 lata.

212. V. *FLEXUOSA* (Sims bot. mag. t. 2477), caulibus herbaceis è rhizomate lignoso teretibus striatis albo-pilosis, foliis sessilibus oblongis seu linear-lanceolatis subintegerrimis utrinquè scabro-pilosis, cymis scorpioideis paucis, capitulis bracteis oblongis longioribus secus cymas ramos ad dichotomias sessilibus multifloris, invol. campanulati foliolis 1-nerviis oblongis mucronato-acuminatis extus subvillosis, achenio turbinato undiquè villosa, pappi serie externâ paleacea acuminatâ interiore dimidio ferè brevior. ♀ in Brasiliâ meridionali. Lessing in linnæâ 1829. p. 311. 1831. p. 670. Lodd. bot. cat. t. 1880. Variat cor. purpureis et albis. Pappus argenteus valdè inequalis aut subæqualis. Rhizoma tuberosum. (v. s. comm. à Mus. reg. Berol.)

213. V. *COGNATA* (Less. in linnæâ 1831. p. 670), caule erecto striato villosa, foliis sessilibus penninerviis basi obtusiusculis, inf. obovatis acutis, sup. linearibus acuminatis suprà scabris hirsutiusculis subtus villosa-sericeis, cymis ramis longatis erecto-patentibus villosissimis polycephalis, capitulis 10-12-floris remotis folia sua superantibus sessilibus, cor. glabrâ, achenio piloso, pappi ser. ext. paleacea brevi. — in Brasiliâ. Affinis dicitur V. Platensis et V. obovata. Caulis totus foliatus. Villi in sicco flavicantes. Fol. infer. 4 poll. longa, sesquipollicem lata.

214. V. *PLATENSIS* (Less. in linnæâ 1829. p. 312. 1831. p. 671), caule herbaceo tereti striato apice subhirsuto, foliis sessilibus confertis elongato-lanceolatis integerrimis aut undulato-crenatis suprà rugosis pubescentibus subtus villosa-canescantibus, cymis subscorpioideis axillaribus compositis in paniculam ferè dispositis, capitulis multifloris, invol. campanulati foliolis 1-nerviis oblongis mucronatis extus villosis, achenio piloso, pappi serie externâ brevissimâ. ♀ in Brasiliâ meridionali ad Rio de la Plata (Spreng.) et Rio-Grande (b. Mus. imp. Bras. n. 767). Conyza Platensis Spreng. 3. p. 509. Cor. purpureæ glabræ. Flores in invol. 35-40. Pappus argenteus. (v. s.)

215. V. *RADULA* (Mart.! herb.), caule herbaceo basi suffruticulosulo erecto suban-  
—sto glabresculo, foliis sessilibus oblongo-linearibus acutiusculis coriaceis quæ glabris asperulis venis elevatis et reticulatis subintegerrimis, capitulis

ad apicem caulis ad axillas aut extrâ axillas folio flor. multò brevioribus sessilibus aut breviter pedicellatis 70-floris, invol. subglobosâ squamis pluriserialibus ext. lanceolatis longè acuminatis semipatulis subaraneosis, intimis linearibus membranaceis subacuminatis, cor. glabrâ, achenio glabrisculo, pappi ser. ext. paleaceâ brevi, int. setis apice subclavellatis.  $\frac{3}{4}$   $\frac{1}{2}$  in editis prope Sanctam Barbaram ad fl. Sabucaby prov. Minarum General. legit cl. Martins. Fol. 3-4 poll. longa, 4-6 lin. lata. Cor. purpureæ. Rhizoma lignosum. (v. s. in h. Acad. reg. Monac.)

$\beta$ . *latifolia*, caule apice paulò magis villosiusculo, foliis oblongis vix acutis subtus puberulis perspicuè denticulatis, achenio longiore magis piloso.  $\frac{3}{4}$  in Brasiliâ legit cl. Martins et cum var.  $\alpha$  in herb. conjunxit. Folia 2-3 poll. longa, 9-12 lin. lata. An potiùs species propria? (v. s. in h. Acad. reg. Monac.)

216. *V. RECATANTHA*, caule suffruticuloso erecto ramoso tereti glabrisculo apice araneoso, foliis sessilibus linearibus utrinquè subattenuatis rigidis integerrimis suprâ glabrâ subtus subtomentoso-canescenscentibus, capitulis subcorymbosis terminalibus et paucis lateralibus folio florali vix brevioribus ovato-globosis 100-110-floris, invol. squamis multiserialibus adpressis, ext. latè ovatis vix submucronatis, inter. lineari-oblongis obtusis, cor. glabrâ glandulosâ, achenio 10-costato glabro, pappi ser. ext. paleaceâ brevi.  $\frac{3}{4}$  patr. ign. Folia et invol. *V. ligulæfoliæ*, sed folia angustiora, ferè *V. radulæ* var. *angustifoliæ* similis, dispos. capit. diversa, flor. numerus multò major, achenis glabra, etc. (v. s. in h. Acad. reg. Monac.)

217. *V. ERISYFOLIA* (Mart.! herb.), caule suffruticuloso tereti ramoso apice subviloso, foliis sessilibus linearibus acutis subvilosis suprâ subasperis subtus penninerviis subreticulatis margine subintegerrimis, capitulis secus ramos elongatos sessilibus axillaribus aut extraaxillaribus folio brevioribus circ. 35-40-floris, invol. ovatis squamis pluserialibus, ext. lanceolatis acuminatis subvilosis semipatulis, int. linearibus obtusis mucronatis, achenio glabro, pappi ser. ext. ....  $\frac{3}{4}$  in subsylvaticis montanis prov. Minarum General. legit cl. Martins. Folia 5 poll. longa, 5 lin. lata. Affinis *V. radulæ* var.  $\alpha$ . Flores in capit. juniores? (v. s. in h. Acad. reg. Monac.)

218. *V. SPFXIANA* (Mart.! herb.), caule herbaceo ramoso erecto striato apice adpressè puberulo demùm glabro, foliis subsessilibus linearibus utrinquè attenuatis integerrimis 1-nerviis suprâ et margine scabris subtus subhirsutis, paniculæ ramis bifidis folia ovali-lanceolata subsericea gerentibus subscorpioides oligo cephalis, capitulis confertis subsessilibus fol. flor. longioribus 7-floris, invol. oblongi squamis lanceolatis, exter. dorso villosis mucronatis, int. longioribus subglabris acuminatis, cor. subpilosâ, achenio sericeo-villoso, pappi ser. ext. paleaceâ brevi.  $\frac{1}{2}$  in deserto Minarum Gener. Brasiliæ legit cl. Martins. Folia pollicem longa, 2 lin. lata. Pappus albescens. Cor. pallidæ ex sicco. Invol. squamæ subrubentes. (v. s. in h. Acad. reg. Monac.)

219. *V. VAUTICULOSA* (Mart.! herb.), caule fruticuloso ramoso tereti erecto apice pulverulento-pubescente, foliis oblongis sessilibus utrinquè subacutis integerrimis aut pauciserratis utrinquè minutè et adpressè pubescentibus cinereis 1-nerviis, cymâ scorpioidèâ subaphyllâ patulâ, capitulis unilateralibus sessilibus oblongis 8-floris, invol. squamis oblongis, exter. submucronatis adpressis intimis subacuminatis, cor. hirtellâ, achenio sericeo-villoso, pappi ser. ext. angustè paleaceâ acuminatâ ferè triplò inter. breviorè.  $\frac{3}{4}$  in deserto ad flum. Sancti-Francisci prov. Minarum General. Brasiliæ legit cl. Martins. Folia 7-8 lin. longa, 2-3 lata. Capit. 3 lin. longum. (v. s. in h. Acad. reg. Monac.)

$\beta$ ? *penninervis*, ramulis striatis, foliis duplò majoribus integerrimis subtus penninerviis, capitulis magis approximatiss paulò majoribus 8-9-floris, invol. squamis mucrone subpatente brevi terminatis, pappi ser. ext. paleaceâ quintam serici internæ longitudinem vix superante.  $\frac{3}{4}$  cum priore juncta in herb. Mart. sed verisim. distinguenda. (v. s. in h. Acad. reg. Monac.)

220. *V. MENZIEBIA* (Mart. ! herb.), caule herbaceo erecto tereti hirsuto, foliis sessilibus lineari-lanceolatis basi obtusis apice cuspidatis subintegerrimis penninerviis suprà sparsè pilosis subtus hirsutis, paniculæ laxæ ramis scorpioideis elongatis, capitulis ad axillas foliorum geminis sessilibus folio longioribus 10-floris, invol. squamis lanceolatis acuminatis dorso subhirsutis, cor. glabrâ, achenio puberulo parvisimè glanduloso, pappi ser. ext. paleaceâ brevi. 2½ in campis editis prov. Sancti-Pauli Brasiliæ leg. cl. Martius et Lund. Herba bipedalis. Folia 2 poll. longa, 4-6 lin. lata. Pappus albidus. Cor. purpureæ. (v. s.)

β. *adenocarpa*, foliis brevius cuspidatis, invol. squamis dorso magis hirsutis, foliis pube magis sericeâ hirsutis, achenio copiosissimè glanduloso. 2½ in prov. Sancti-Pauli Bras. legit cl. Martius et cum var. α in herb. coadunavit. (v. s. in h. Acad. reg. Monac.)

221. *V. ARENARIA* (Mart. ! herb.), fruticosa tota tomento brevi cinascenti-velutina, ramis teretibus, foliis petiolatis ellipticis basi acutis apice obtusis integerrimis penninerviis, nervis subtus prominulis, capitulis axillaribus solitariis sessilibus 30-40-floris, invol. hemisphærici squamis imbricatis acuminatis dorso puberulis, ext. acumine subrecurvò, int. rectis, achenio sericeo-villosissimo, pappi serie ext. paleaceâ brevi. ½ in rupibus arenariis prov. Pianhiensis Brasiliæ legit cl. Martius. Cor. purpureo-violaceæ glandulosæ. Pappus nitens albidus. Fol. 1-2 poll. longa, 9-12 lin. lata, petiolo 2-3 lin. (v. s. in h. Acad. reg. Monac.)

222. *V. DEBILIS* (Mart. ! herb.), caule fruticoso ramosissimo, ramis teretibus debilibus apice velutino-tomentosis, foliis oblongo-lanceolatis vix petiolulatis utrinquè acuminatis integerrimis suprà pubescenti-velutinis subtus cano-tomentosis, ramis flor. scorpioideis, capitulis ad axillas foliorum flor. solitariis geminisve sessilibus folio multò brevioribus 20-25-floris, invol. squamis oblonga dorso subvelutinis mucronato-acuminatis adpressis, corollâ extus hirtellâ, achenio tereti puberulo, pappi ser. ext. paleaceâ brevi. ½ in apricis aridis deserti versus flum. Sancti-Francisci prov. Minar. General. Brasiliæ legit cl. Martius. Folia 15-18 lin. longa, 4 lin. lata. Pappus argenteus. Cor. purpurascens. (v. s. in h. Acad. reg. Monac.)

223. *V. HERMOPHILA* (Mart. ! herb.), suffrutescens tota pube brevi adpressâ subcanescente, caule erecto tereti ramoso, foliis elongatis lanceolato-linearibus utrinquè attenuatis integerrimis penninerviis, ramis scorpioideis oligocephalis sæpè bifidis, capitulis in dichotomiis ramorum et ad axillas fol. ram. sessilibus solitariis 10-floris, invol. ovati squamis lanceolatis subacuminatis adpressis, cor. glabrâ glandulosâ, achenio turbinato sericeo-villoso, pappi ser. ext. paleaceâ brevi acuminatâ. ½ in campis arenosis versus Goyaz prov. Minar. Gener. legit cl. Martius. Cor. albæ. Anthæræ rosæ. Pappus sordidè albescent. Folia 3-4 poll. longa, 3-4 lin. lata. (v. s. in h. Acad. reg. Monac.)

224. *V. CHALYBEA* (Mart. ! herb.), suffrutescens ? caule erecto tereti puberulo, foliis oblongo-linearibus basi acutis apice obtusis integerrimis subundulatis 1-nerviis utrinquè puberulis, ramis elongatis subscorpioideis, capitulis ad axillas foliorum sessilibus solitariis (rarè geminis) 20-floris, invol. squamis omnibus acuminatissimis glabris nitidis subcoloratis, cor. glabrâ glandulosâ, achenio sericeo-villoso, pappi ser. ext. paleaceâ brevi. ½ in aëcis prov. Bahiensis interioris legit cl. Martius. Herba ferè *Erigerontis canadensis*. Folia 6-9 lin. longa, æquilineam lata. Pappus pallidus nitens. (v. s. in h. Acad. reg. Monac.)

225. *V. TRICHOLEPIS*, caule herbaceo erecto, ramoso tereti glabro, ramulis subtiliter striatis et puberulis, foliis sessilibus membranaceis ovato-lanceolatis basi attenuatis apice acuminatis subintegerrimis, suprà glabrinaculis, subtus piloso-subhirsutis, capitulis secus ramos ad axillas foliorum sessilibus solitariis 15-18 folio flor. triplò brevioribus, invol. ovali-oblongi squamis glabris, ext. pillatis semipatulis, intimis lanceolato-linearibus acuminatis pappo et corollis agrioribus, achenio teretiusculo adpressè piloso, pappi ser. ext. brevi anguste

- paleacea?  $\frac{1}{2}$ ? in siccis insulæ Cayennæ (Leprieur!) et in montosis prope Rio de Contas prov. Bahiensis Brasilæ (Martius). Vern. patula Mart.! herb. Folia 3 poll. longa, pollicem lata. Pappus griseus invol. brevior. (v. s.)
226. *V. SALZMANNI*, caule basi suffruticoso ramoso tereti pubescente, ramis elongatis, foliis lineari-lanceolatis utrinque acuminate vix petiolatis integerrimis membranaceis utrinque subpubescentibus, capitulis 30-floris, secus ramos ad axillas foliorum sessilibus solitariis geminative ovatis, involucri squamis lineari-oblongis subacuminatis subvillosis, achenio piloso oblongo tereti, pappi serie externâ angustè paleacè brevissimâ.  $\frac{1}{2}$  in umbrosis circa Bahiam Bras. semel legit cl. Salzmann. Caulis 4-pedalis. Folia 2-2  $\frac{1}{2}$  poll. longa, 5-6 lin. lata. Invol. rufescentis. Pappus pallidè rufescens. Cor. purpureæ, lobis apice subpuberis. (v. s.)
227. *V. MUCRONIFOLIA*, caule basi suffruticuloso tereti hirsuto-pubescente, foliis brevissimè petiolatis ovatis ovalibusve basi obtusis apice mucrone acuto apiculatis utrinque minutissimè pubescentibus, cymæ ramis elongatis, capitulis ad axillas fol. sessilibus solitariis 10-12-floris, invol. squamis pilosiusculis acuminatis, achenio sericeo-villoso, pappi paleis externis brevibus oblongis paleis.  $\frac{1}{2}$  in sabulosis maritimis ad Bahiam Brasilæ leg. cl. Salzmann et Blanchet. Herba 4-pedalis. Aff. *V. cotoneastri*. Folia 15-18 lin. longa, 7-9 lata. Cor. purpureæ glabræ. Pappus argenteus. (v. s.)
228. *V. POEPPIGIANA*, caule herbaceo? tereti ramoso villosa, foliis sessilibus oblongo-lanceolatis utrinque attenuatis subintegerrimis suprà sparè pilosis subtus villosa-hirsutis, ramis scorpioideis, capitulis ad axillas folior. sessilibus aut subgeminis 35-floris folio triplo brevioribus, invol. squamis lineari-lanceolatis omnibus acuminatis, ext. submucronatis dorso adpressè pubentibus, cor. glabrâ, achenio striato nervulis obtusis lævibus valleculis seriatis pilosis, pappi ser. ext. vix paleacè brevi. — in Americâ calidiore vixima. in Peruvia legit cl. Poeppig. (pl. exs. n. 1204). Affinis *V. imponenti* et *salicifoliz*. Cor. purpureæ. Folia 3-4 poll. longa, 5-6 lin. lata. Pappus argenteus. An satis ab obscurâ diversa? (v. s. comm. à cl. inv.)
229. *V. MURICATA*, herbacea?, caule tereti ramoso striato subglabrato, ramis pedunculisque densè hirsutis, foliis subsessilibus ovato-lanceolatis basi obtusis apice acuminatis subcrenatis suprà junioribus puberis demum muricatis subtus densè et molliter hirsutis, paniculæ terminalis ramosissimæ ramis scorpioideis, capitulis sessilibus plerisque bractea foliacè suffultis 20-25-floris, invol. ovatis squamis ext. hirsutissimis acuminè rigido subpatente apiculatis, intimis glabrisculis acuminatis, achenio tereti villosa, pappi ser. ext. paleacè brevi.  $\frac{1}{2}$ ? in Brasiliâ legit cl. Lhostky. Folia 3-4 poll. longa, 15-18 lin. lata. Pappus subrufescens. Cor. purpureæ? hirtæ? Tubercula paginæ fol. superæ sunt pilorum deciduorum bases. (v. s.)
230. *V. SAGRAIANA*, fruticosa ramosissima, ramis teretibus villosis, foliis brevissimè petiolatis ellipticis basi obtusis apice breviter acuminatis distanter serrulatis penninerviis suprà glabris subtus in nervis pubescentibus, ramis elongatis subscorpioideis, capitulis unilateralibus ad foliorum axillas solitariis sessilibus folio multo brevioribus 30-floris, invol. ovatis squamis ext. lanceolatis ciliatis, inter. longioribus glabris linearibus subacuminatis, cor. et achenio glabris, pappi ser. ext. angustè paleacè brevi.  $\frac{1}{2}$  in insulâ Cubâ circa Havanam legit cl. Ramon de la Sagra. Cor. purpureæ. Pappus albus. Achenium teres. Fol. 3 poll. longa, æquipollicem lata, petiolo 1-1  $\frac{1}{2}$  lin. (v. s. comm. à cl. inv.)
231. *V. ZUCCASINTANA* (Mart.! herb.), suffruticosa? caule tereti substriato erecto ramoso vix pubescente, foliis subsessilibus elliptico-oblongis utrinque subobtusis hinc indè subcuneatis suprà muricatis subtus molliter hirsutis penninerviis, ramis scorpioideis, capitulis 20-25-floris sessilibus distantibus solitariis geminative nunc ebracteatis nunc ad foliorum axillas sitis, invol. ovatis squamis siccis imbricatis arcè adpressis obtusis longè ciliatis dorso glabris, achenio glaberrimo, pappi serie ext. paleacè brevi.  $\frac{1}{2}$ ? ad Minas Novas Brasilæ legit



- cl. Martius. Fol. 3 poll. longa, sesquipollicem lata. Pappi setæ albidæ, crassiusculæ. (v. s. in h. Acad. reg. Monac.)
232. *V. VARROXIEPOLIA*, caule fruticoso ramoso tereti basi glabresculo supernæ pube pulverulentâ ferrugineo, foliis subsessilibus ovalibus utrinque subobtusis subintegerrimis penninerviis reticulatis suprâ muriculatis bullulatis subtis ferrugineo-velutinis, capitulis ad axillas foliorum secus sessilibus solitariis aut geminis folio flor. triplò brevioribus oblongis 18-20-floris, invol. squamis ext. lanceolatis mucronato-acuminatis adpressis, inter. linearibus minutè acuminatis, cor. et achenio glabris, pappi serie ext. brevissimâ vix paleaceâ. 3 patr. igs. Cor. purpureæ 2 lin. longæ. Achenia striata. Fol. 2 poll. longa, 12-15 lin. lata. Capitula adproximata secus ramos circ. 20-30. Pappus stramineus. Paulò accedit ad *V. Sagreaham* et ad *V. aurifodinarum*. (v. s. in h. Acad. reg. Monac.)
233. *V. LEVIGATA* (Mart. l. herb.), glaberrima, caule herbaceo tereti, ramulis subangulatis, foliis lato-linearibus sessilibus integerrimis callosis-sabacutis coriaceis utrinque sabreticulatis, ramis scorpioideis, capitulis ad axillam foliorum solitariis geminisve folio multò brevioribus 12-13-floris, invol. ovatis squamis oblongis adpressis sabacutis, achenio villosio, pappi ser. ext. paleaceâ brevi. 2½ in alpestribus prope Gongonhas do campo prov. Minar. Gener. legit cl. Martius. Pappus stramineus scaber. Folia 4 poll. longa, 4 lin. lata. (v. s. in h. Acad. reg. Monac.)
- β. bupleurifolia*, caule striato crassiori, foliis oblongis, super. ferè ellipticis, capitulis 17-18-floris, squamis invol. magis obtusis. 2½ cum var. *α*? legit cl. Martius. Folia ferè Bupleuri fruticosi, 4 poll. longa, 8-10 lin. lata; summa poll. longa, 5 lin. lata. (v. s. in h. Acad. reg. Monac.)
234. *V. LACUNOSA* (Mart. l. herb.), caule herbaceo erecto tereti densè tomentoso, foliis subsessilibus ellipticis basi obtusis apice vix sabacutis integerrimis suprâ pubero-velutinis subrugosis subtis densè tomentosis penninerviis, nervis elevatis reticulatis, ramis floridis bifidis in paniculam ferè dispositis scorpioideis, capitulis sessilibus 10-floris ad dichotomias et in axillis foliorum ovatorum isæqualium, invol. ovato oblongi squamis erectis oblongis dorso lanatis intimis apice acuminatis, cor. lobis apice panicetosis, achenio sericeo-viloso, pappi ser. ext. paleaceâ acuminatâ. 2½ in pratis montium prope Villa-Ricca Minar. Gener. Brasilæ legit cl. Martius. Corollæ et stigm. rosæ. Herba 2-3-pedalis. Fol. 2 poll. longa, pollicem lata. Recept. nudum. (v. s. in h. Acad. reg. Monac.)
- §. 6. *Involucris squamis intimis acuminatis, capitulis neque solitariis neque in cymas scorpioideas dispositis.* Less. l. c. p. 671. — *Species omnes Americanae.*
235. *V. EXTRAAXILLARIS*, caulibus à collo lignoso pluribus simplicibus subherbaceis teretibus tomentosis erectis apice arcuatis, foliis sessilibus linearibus acuminatis margine revolutis suprâ glabris nervo impresso subtis tomentosis, cymâ terminali simplici, capitulis obrectatis inter folia extra-axillaribus sessilibus solitariis ovatis 20-floris, invol. squamis rigidè acuminatis dorso araneosis, achenio teretibus-sericeo-viloso, pappi ser. ext. angustè paleaceâ acuminatâ, inter. dimidio breviorè. 2½ in Brasilæ prov. Minar. Gener. prope Tejuco unicâ vice legit cl. Vauthier. Caulis sesquipedalis gracilis. Folia sesquipoll. longa, 1-2 lin. lata, rigidula. Pappus stramineus. Capitula infima axillaria gemina, cætera extra-axillaria solitaria. (v. s. commun. à cl. Vauthier.)
236. *V. LINEARIS* (Spreng. syst. 3. p. 437), caulibus è rhizomate lignoso pluribus herba lignosis lanato-tomentosis simplicibus, foliis sessilibus linearibus mucronatis 1-nerviis integerrimis margine revolutis suprâ glabris lanato-tomentosis, capitulis sparsis subsessilibus remotis folio stipatis, thelyndraceis squamis siccis pungenti-acuminatis arachnoideo-pilosis. 1, achenio densè piloso, pappi serie ext. paleaceâ brevi. 3 in Brasilâ

- tropicâ. Less. in linnæâ 1829. p. 285. 1831. p. 671. Cor. purpureæ. Pappus flavescens. Fol. 2 poll. longa, lineam lata.
237. *V. ROSEMARINFOLIA* (Less. in linnæâ 1829. p. 286. 1831. p. 672), caulibus lignosis ramosis teretibus lanato-tomentosis, foliis sessilibus 1-nerviis rigidis linearibus mucronato-acutis integerrimis margine revolutis suprà glabris subtus lanato-tomentosis, capitulis approximatis multifloris subcylindraceis foliis plerumque stipatis, invol. squamis uccis pungenti-acuminatis ciliatis apice albidis, achenio dense piloso, pappi serie ext. brevi.  $\frac{1}{2}$  in Brasiliâ tropicâ. Nervus folii pellucidus. Species valde affinis *V. lineari*.
238. *V. LINEARIFOLIA* (Less. in linnæâ 1829. p. 287. 1831. p. 672), caulibus è rhizomate lignoso plerumque teretibus subsimplicibus lanato-tomentosis, foliis sessilibus confertis linearibus 1-nerviis acuminatis coriaceis margine integerrimis et revolutis suprà glabrisculis subtus lanato-tomentosis, capitulis campanulatis multifloris ramosos nudos terminantibus, invol. squamis siccis 3-nerviis externis pungenti-acuminatis, achenio dense villosa, pappi serie ext. brevi.  $\frac{1}{2}$  in Brasiliâ tropicâ. Valde affinis *V. lineari*.
239. *V. XANTHOPHYLLA*, caule herbaceo erecto simplici supernè angulato sparè hispido apice nudo, foliis oblongo-linearibus sessilibus suprà sparè hirtis subtus nervo medio excepto glabris purpureis margine dentatis, dentibus acutis rigidè reflexis, corymbo 4-cephalo simplici aphylo, invol. squamis lanceolatis glabris, exter. obtusis, intimis acuminatis, achenio striato pilis rubris hirtello, pappi serie externâ squamellatâ brevi.  $\frac{1}{2}$  in campis editis prov. Sancti-Pauli Brasil. ad Mugidas Grues legit cl. Lund. Nov. flor. flor. retroflexis et pilis achenii rubris spectabilibus. Herba circ. pedalis. Folia 2 poll. longa, 4 lin. lata. (v. s. comm. à cl. inv.)
240. *V. CEPHALOTES*, caule herbaceo erecto simplici hirsuto, supernè nudo, infrâ medium folioso, foliis 4-5 approximato-subverticillatis sessilibus obovatis obtusis suprà sparè pilosisculis, subtus canescenti-tomentosis, bracteolâ lineari in parte caulis nudâ, capitulis circ. 5-6 breviter pedicellatis ebracteatis in umbellam capitatam dispositis, invol. squamis lineari-lanceolatis pubescentibus apice acuminatis, acheniis villosis, pappi serie externâ paleaceâ brevi.  $\frac{1}{2}$  in campis editis prov. Sancti-Pauli ad Mugl. Nov. flor. legit cl. Lund. An *Chrysocoma oligophylla* Arrab. fl. flum. 8. t. 2? Cor. purpureæ glabræ. Herba vix pedalis. Folia  $\frac{3}{4}$  poll. longa, sesquipollicem lata. (v. s. comm. à cl. Lund.)
241. *V. BARBATA* (Less. in linnæâ 1829. p. 287. 1831. p. 672), fruticosa, ramis teretibus tomentosis, foliis sessilibus imbricatis cordato-ovatis acutis coriaceis integerrimis suprà subpilis invol. villosis, cymæ terminalis ramis dichotomis, capitulis sessilibus folio stipatis, involucri turbinati pauciserialis squamis siccis 1-nerviis oblongo-acuminatis apice glabrisculis, achenio longè denseque sericeo, pappi serie ext. paleaceâ longiusculâ.  $\frac{3}{4}$  in Brasiliâ tropicâ. Cor. lobi extus longè barbati. Pappus argenteus.
242. *V. VISCIDULA* (Less. in linnæâ 1829. p. 289. 1831. p. 672), fruticosa, ramis teretibusculis viscido-pubescentibus, foliis sessilibus coriaceis ovato-oblongis acutis subdenticulatis suprà lucidis subtus viscido-pubescentibus, cymis terminalibus ramosis, capitulis brevè pedicellatis multifloris, invol. turbinati squamis siccis glabris acuminatis, cor. glabrâ, achenio piloso, pappi serie paleaceâ brevi.  $\frac{3}{4}$  in Brasiliâ tropicâ. Pappus argenteus.
243. *V. RUBROIDES* (Less. in linnæâ 1829. p. 290. 1831. p. 672), caule suffruticoso simplici tereti supernè obscure pubescente, foliis brevè petiolatis confertis rigidis obovatis apice denticulatis suprà glabris scabris subtus pubescentibus, cymis terminalibus oligocephalis, capitulis approximatis subsessilibus multifloris, invol. cylindraceis squamis siccis 3-nerviis acuminatis glabris demum patentibus, achenio oblongo glabro basi et apice subpiloso, pappi serie ext. paleaceâ brevi.  $\frac{3}{4}$  in Brasiliâ tropicâ. Cor. ign. Pappus argenteus.
244. *V. SPECIOSA* (Less. in linnæâ 1829. p. 290. 1831. p. 672), subarborea, ramis teretibus villosis-lanatis, foliis brevissimè petiolatis ovatis acutis coriaceis in-

tegerimis suprà glabris glandulosis subtùs nervis villosis-lanatis, cymà terminali spiciformi, capitulis sparsis in axillis fol. sup. sessilibus, invol. pauciserialis squamis siccis acuminatis subtomentosis, achenio dense piloso, pappi lati paleacei serie ext. longà. 5 in Brasilià tropicà. Folia  $1\frac{1}{2}$ -2  $\frac{1}{2}$  poll. longa, 12-14 lin. lata.

245. *V. ARCTIOMES* (Less. in Linnæa 1831. p. 400 et 672), fruticosa, ramis sulcato-angulatis tenerè tomentosis, foliis petiolulatis firmis obovato-lanceolatis cuneatis acuminatis inæqualiter serratis suprà lucidis glabris subtùs arachnoides, paniculà terminali contractà, capitulis multifloris, invol. squamis multinerviis ext. triangularibus rigidis, int. angustis acuminatis, corollà et achenio glaberrimis, pappi seriebus latitudinè non distinctis, ext. breviorè. 3 in Mexico. *Diazexis serrata* Don trans. lin. soc. 16. p. 254. Folia 5-8 poll. longa, 2-3  $\frac{1}{2}$  poll. lata, petiolo 2-3 lin. Cor. purpureæ. Pappus argenteus. Achenia 4-gona.

246. *V. AROINTA* (Less. in Linnæa 1831. p. 672), caule herbaceo simplici sulcato-argenteo-tomentoso, foliis subsessilibus obovatis mucronatis coriaceis penninerviis inf. serratis suprà nitidis punctatis subtùs argenteo-tomentosis, racemo terminali angusto, capitulis folio orbiculari basilari superatis, invol. campanulati squamis laxè imbricatis linearibus pungentibus, cor. glabri, achenio hirsuto, pappi ser. ext. paleaceà brevi. — in Brasilià. Fol. 4 poll. longa,  $1\frac{1}{2}$ -2 lata. Racemus 2-8 poll. Cor. purpureæ.

247. *V. XANTHOPHYLLA* (Mart. ! herb.), caule fruticoso ramoso tereti tomento denso flavicante cum pedunculis et fol. paginà infer. vestito, foliis subsessilibus ovalibus utrinquè obtusis subcrenatis coriaceis penninerviis suprà glabris, nervo medio basi tomentoso, capitulis 50-floris breviter pedunculatis in cymas contractas subpaniculatim dispositis, foliis floralibus ovatis sub ramulis, invol. squamis multiserialibus dorso tomentosis oblongis obtusis imbricatis, recept. nudo, achenio sericeo-villosissimo, pappi serie ext. ferè setiformi, int. triplo breviorè. 3 in campis arenosis in Chapada do Parana in confin. prov. Goyasæ legit cl. Martius. Cor. ign. Pappus rufo-flavescens. Folia sesquipollicem longa, pollicem lata. (v. s. in h. Acad. reg. Monac.)

248. *V. AURA* (Mart. ! herb.), herbacea tota hirsuta, caule tereti subsimplici, foliis vix petiolulatis ovalibus utrinquè obtusis integerrimis, capitulis ad apicem caulis et ramorum in axillis foliorum sessilibus solitariis folio paulò brevioribus circ. 40-floris, invol. campanulati squamis lanceolatis, ext. in acumen gracile patulum productis, inter. acuminatis erectis, achenio sericeo-villoso, pappi ser. ext. angustè paleaceà brevi. ☉ in campis editis prov. Minas Generalium Brasilie legit cl. Martius. Hirsuties partium juniorum flavescens. Fol. 15-16 lin. longa, 7-8 lata. Pappus albescens. Cor. glabræ ex siccis purpurascens. Radix descendens lateraliter fibrosa. (v. s. in h. Acad. reg. Monac.)

249. *V. CONDIGNA* (Mart. ! herb.), caulibus è collo suffruticoso pluribus herbaceis teretibus apice hirsutis, foliis sessilibus basi cordatis ovatis acutis rigidulis imbricatis integerrimis penninerviis reticulatis suprà glabris subtùs parè villosiusculis, ramulis foliosis simplicibus 1-cephalis aut bifidis oligocephalis, capitulis subcorymbosis bracteatis 20-floris, invol. squamis oblongis parè acuminatis basi glabris apice hirsutissimis, corollis acheniisque glabris, pappi ser. ext. angustè paleaceà acuminatà, inter. quadruplo breviorè. 2½ in editis prov. Minas General. legit cl. Martius. Herba vix pedalis. Folia 15 lin. longa, 6-8 lata. Capit. 7-8 lin. longa. Cor. purpureæ. Pappus albidus. Achenia juniora hinc inde subpilosa. (v. s. in h. Acad. reg. Monac.)

250. *V. PYCNOSTACHYA*, fruticosa tota molliter hirsuto-tomentosa, caulibus teretibus ad apicem usque confertè foliosis, foliis ovali-aut obovato-oblongis acutis sessilibus planis integerrimis penninerviis, capitulis ad axillas supremas sessilibus folia sua non superantibus in spicam densam confertis circ. 40-floris, invol. ovato-campanulati squamis coriaceis imbricatis pluri-serialibus lanceolatis subacutis, junior. subvelutinis adultis glabris. 3 in Brasiliæ prov. Minas. Gener. ad Serro do Frio (Vauth. n. 267) in editioribus Minas (Mart. !). Vera decussata Mart. ! herb. sed folia non decussata, alterna. Flor. purpurei. Achen.

sericeo-villosum. Pappi ser. ext. paleis oblongis apice incis, int. setis filiformibus 2-3-serialibus crassiusculis. Recept. areolatum. (v. s.)

251. *V. TOMENTELLA* (Mart. ! herb.), fruticosa, ramis teretibus junioribus tomentosis, foliis confertis sessilibus ovalibus subacutis integerrimis nervis utrinque prominulis suprà glaberrimis subtùs dense tomentosis, capitula ad apices ramorum sessilibus apicatis extrà-axillaribus folia sua non superantibus 20-floris, invol. ovati squamis imbricatis adpressis ciliatis acuminatis, cor. glabrà, achenio sericeo-piloso, pappi ser. ext. paleacea brevi.  $\frac{3}{4}$  in campis prov. Sancti-Pauli Brasiliæ legit cl. Martins. Simillima *V. pycnostachyæ* sed capitula 20 nec 40-flora, dimidio minora, folia paulò minora suprà glaberrima. (v. s. in h. Acad. reg. Monac.)

252. *V. ROSA* (Mart. ! herb.), fruticosa, ramis teretibus tomentosis, foliis sessilibus lanceolatis basi subacutis apice subacuminatis integerrimis suprà glaberrimis subtùs tomentosis penninerviis, ramis erectis, capitulis paucis sessilibus lateralibus et terminalibus folio flor. aequalibus 80-90-floris, invol. subglobosi squamis adpressis lanceolatis acuminatis dorso aranoso-sublanatis, cor. glabrà, achenio sericeo-villoso, pappi ser. ext. angustè paleacea interiore circ. dimidio brevior.  $\frac{3}{4}$  in campis prope Villa do Principe prov. Minarum General. Brasiliæ legit cl. Martins. Flores rosei. Folia 12-15 lin. longa, 3-5 lata. Capit. 7-8 lin. diam. Recept. nudum. (v. s. in h. Acad. reg. Monac.)

253. *V. VEFARTORUM* (Mart. ! herb.), caule fruticoso ramoso, ramis teretibus sericeo-villosis ad apicem foliosis, foliis sessilibus ovalibus lanceolatisve planis subintegerrimis suprà glabris subtùs adpressè sericeo-villosis, capitulis ex axillis supremis pedunculatis subcorymbosis 20-25-floris, pedunculis nudis folio suo brevioribus, invol. ovati squamis imbricatis lanceolatis siccis dorso pubescentibus subacutis.  $\frac{3}{4}$  in deserto trans Serram de Sant-Antonio et prope Caiete prov. Min. Gen. et Bahiam Brasiliæ legit cl. Martins. Recept. areolatum. Ach. sericeo-villosum. Pappi ser. ext. paleis oblongo-linearibus acutis, int. setis quadruplò longioribus scabris crassiusculis. Cor. purpureæ. (v. s. in h. Acad. reg. Monac.)

Variat  $\alpha$  : foliis latioribus capitulis distinctè pedunculatis, et  $\beta$  : foliis angustioribus capitulis subsessilibus.

254. *V. SENESCENTES* (Mart. ! herb.), caule fruticoso tereti hirsuto-tomentoso, foliis sessilibus basi cordatis ovatis aut suborbiculatis apice calloso-subacutis integerrimis utrinque (subtùs præcipuè) villosa-tomentosis, paniculæ ramis axillaribus brevibus oligocephalis, capitulis bractea foliaceà longioribus 10-floris ad apices ramorum confertis subsessilibus, invol. squamis lanceolatis acuminatis extis velutino-villosis, cor. lobis extùs ad apicem barbatis, achenio sericeo-villosissimo, pappi ser. ext. paleacea acuminatà inter. triplò brevior.  $\frac{3}{4}$  in campis editis prov. Minarum gener. et inter. Bahiensis legit cl. Martins. Variat foliis suborbiculatis 8-9 lin. longis latisque, et ovatis 2 poll. longis, pollicem latis. Invol. 4-6 lin. longum. Pappus albidus. (v. s. in h. Acad. reg. Monac.)

255. *V. FIRMULA* (Mart. ! herb.), caule fruticoso tereti erecto ramulisque hirsutis, foliis sessilibus oblongo-lanceolatis rigidis acuminatis integerrimis suprà glaberrimis subtùs villis longis sparsis hirsutis, paniculæ elongatæ ramis oligocephalis foliosis brevibus, capitulis 40-floris subbracteatis, invol. squamis linearis-lanceolatis acuminatis ext. hirsutis intus ferè glabris, cor. lobis sub apice barbatis, achenio sericeo-villoso, pappi ser. ext. paleacea brevi.  $\frac{3}{4}$  in campis editis apricis deserti Minarum General. legit cl. Martins. Valdè affinis *V. dura* sed hirsutis faciliè distinctà. Fol. 2 poll. longa, semi-poll. lata. (v. s. in h. Acad. reg. Monac.)

256. *V. DURA* (Mart. ! herb.), suffrutescens? caule erecto tereti glabro, foliis sessilibus oblongo-lanceolatis rigidis subacuminatis integerrimis utrinque glaberrimis, ramis subscorpioidis parè hirsutulis, capitulis 40-floris pedicellatis bractea foliaceà æquius suffultis, invol. campanulatis squamis acuminatis ext. dorso subhirsutis, corollæ lobis subbarbatis, achenio elongato sericeo-villo-

- assimo, pappi ser. ext. angustè paleacea, int. ciliatè externam duplò tantùm superante.  $\frac{1}{2}$ ? in campis Brasiliæ ad Contendas prov. Minar. General. legit cl. Martius. Folia 3 poll. longa, pollicem lata; axillæ subbarbatæ. Pappus rigidus stramineus. Recept. omninò nudum. (v. s. in h. Acad. reg. Monac.)
257. *V. LAMINGIANA* (Mart. herb.), caule suffruticoso tereti tomentoso-hirsuto, foliis sessilibus cordiformibus vix subacuminatis coriaceis integerrimis suprà pube deciduà subvillosis subtùs subhirsutis, paniculæ ramis brevibus axillaribus bifidis oligocephalis, foliis subcapitulis ovatis utrinquè tomentosis capitolo brevioribus, capitulis confertis 3-5 sessilibus 10-floris, invol. ovalis squamis lanceolatis acuminatis dorso subvillosis, cor. lobis dorso hirsutissimis, achenio sericeo-villosissimo, pappi ser. ext. paleacea acuminatà int. triplò brevior.  $\frac{1}{2}$  in campis prope Formigas prov. Minar. Gener. Brasiliæ legit. cl. Martius. Fol. 8-11 lin. longa, 4-6 lata. Pappus albus. (v. s. in h. Acad. reg. Monac.)
258. *V. CHITTIFOLIA* (Mart. herb.), caule herbaceo? erecto simplici tereti striato subhirsuto apice subtomentoso, foliis breviter petiolatis ovatis ovalibusve utrinquè subacutis apice distanter et callosò denticulatis suprà crebrè glandulosis puberulis subtùs parèe glandulosis subhirsutis, paniculæ ramis brevibus subscorpioides oligocephalis, capitulis confertis 10-12 sessilibus folio florali longioribus, invol. ovalis squamis lanceolatis dorso subtomentosis adpressis ext. mucronatis int. acuminatis, cor. glandulosà apice setosà, achenio glabro glanduloso?, pappi serie ext. paleacea brevissima.  $\frac{1}{2}$ ? in montibus prov. Minar. Gener. Brasiliæ legit cl. Martius. Fol. 2 poll. longa, pollicem lata, petioli 2 lin. Nervus medius suprà ad basin hirsuto-tomentosus. Pappus albus argenteus. Recept. nudum. (v. s. in h. Acad. reg. Monac.)
259. *V. TRAGIFOLIA*, caule basi suffruticulosò herbaceo confertè hispido usquè ad cymam folioso, foliis sessilibus oblongis acutis crenato-subdentatis suprà confertè villosis subtùs cano-tomentosis, capitulis secus ramos cymar sessilibus confertis hinc inde foliis stipatis in fasciculum densum glomeratis oblongis 10-floris, invol. squamis villosis pluriserialibus acuminatis, apice uncinato-recurvatis, acheniis pubescentibus, pappi serie ext. paleacea, paleis acutis ciliatis.  $\frac{1}{2}$  in campis editis prov. Sancti-Pauli Nov. flor. legit cl. Land. Species distinctissima vix scorpioides. Cor. purpureæ glabræ. Folia pollicem longa, 4 lin. lata. (v. s. comm. à cl. Land.)

Sect. IX. *ASCARICIDA* Less. in *linnæa* 1831. p. 675. *Ascaricida* et *Centrapalus* Cass. dict. 3. suppl. p. 38 et 7. p. 383. — *Baccharoides* Lin. *f. Zeyl. Manch meth.*

Involucri campanulati squamæ foliaceæ aut apice appendiculatæ plùs miuè patentès et inter se dissimiles. Herbæ capitulis terminalibus solitariis aut parvis corymbosis.

§. 1. *Involucri squamis exterioribus latioribus.*

260. *V? CALYCINA* (Wall. cat. et herb.), caule tereti glabro apice nigutissimè pubescente, foliis latè ovalibus obtusis penninerviis subsessilibus integerrimis glabrinerviis subtùs tenuissimè glandulosis, capitulis brevè pedunculatis corymboso-aggregatis, involucri squamis ext. latè elliptico-oblongis obtusis submucronulatis pedunculatis pubescentibus internerviis acutis minoribus achenio subvillosò, pappi pilis omnibus latitudine subæqualibus ext. dimidio brevioribus.  $\frac{1}{2}$  in Ind. orient. regno burmanico ad Frouse. Cor. extus glandulosa. Flor. in invol. circ. 40-45. Habitus *Vernoniæ* sed involucro et pappo fortè genericè distinguenda. (v. s. comm. ab hon. Cor. merc. angl. Ind. or.)
261. *V. MACROCEPHALA* (Less. in *linnæa* 1829. p. 298. 1831. p. 679), caulibus herbaceis basi suffruticosis teretibus simplicibus tomentosis, foliis brevè petiolatis oblongis utrinquè acuminatis integerrimis suprà glabris subtùs tomentosis penninerviis, cymis scorpioides polycephalis, capitulis distantibus muhi-

floris, involucri latè campanulati squamis oblongo-ellipticis obtusis extus tomentosis, achenio oblongo multicostato sericeo-piloso, pappi serie externà brevi paleaceâ. ♀ in Brasiliâ meridionali. Pappus subflavidus. Cor. roseæ glabræ. Rhizoma lignosum repens. Folia 3-10 poll. longa, 8-30 lin. lata. Flores in capitulo 40-50. Capit. interdum proliferum vidi. Affinis videtur *V. ligulæfolia*. (v. s. comm. à Mus. reg. Berol.)

262. *V. MEXICANA* (Less. in linnæâ 1831. p. 680), fruticosa, ramis teretibus cinereo-subtomentosis, foliis petiolatis obovatis acutis basi cuneatis denticulatis suprà glabriusculis scabridis subtus tenuiter tomentoso-incanis, panicula corymbosa ramis aphyllis, invol. globosi squamis imbricatis glabris, ext. subrotundis acuminatis, med. ellipticis obtusis apice foliaceis, int. linearibus acutis, cor. puberis, achenio glaberrimo, pappo setaceo inæquali. ♂ in Mexico ad Coasta-Grande de Chiconquisco. Cor. cæruleæ. Fol. 8 poll. longa, 6 poll. lata, petiolo 2 lin. Pappus flavescens. Habitus *V. arctioidis*.

263. *V. ALAMANI*, caule...., ramis apice striatis subpuberis, fol. inf...., summis brevissimè petiolatis elliptico-oblongis basi attenuatis apice acuminatis subintegerrimis suprà muricato-scorbis subglandulosis puberulis penninerviis, capitulis pedunculatis ex axillis et apice ramorum paucis nudis 40-45-floris, invol. subglobosi squamis coriaceo-scariosis multiseriatis glaberrimis ovalibus obtusis mucronis longo patente superatis, cor. glabrâ, achenio glanduloso striato, pappi serie ext. brevi paleaceâ. ♀? in Mexico. Frustula tantum sed distinctissima olim misit cl. Alaman. Capit. 8-10 lin. diam. Cor. purpureæ. Pappus albus, ext. subciliatus. (v. s.)

### §. 2. *Involucri squamis exterioribus angustioribus.*

264. *V. BOJERI* (Less. in linnæâ 1831. p. 676), caule herbaceo erecto ramoso tereti glanduloso-hispido, foliis super. ovali-oblongis basi attenuatis apice subacuminatis integerrimis suprà subpuberulis subtus cano-tomentosis reticulatis, capitalis circ. 30-floris, corymbo laxo divaricato, invol. campanulati squamis dorso glanduloso-puberis imbricatis oblongis productis in appendicem setiformem patulam, acheniis villosis. ♂ in campis et ad margines sylvarum insule Zanzibar legit cl. Bojer. Species distinctissima. Frutex 8-10-pedalis. Habitu *Decaneuri* sed achenium teretiusculum. Folia ferè *V. Zeylanicæ* sed indivisa. Pappus albus subplumosus, ser. ext. brevissima. (v. s. comm. à cl. Bojer.)

265. *V. ANTHELMINTHICA* (Willd. sp. 3. p. 1634), caule tereti hispidulo-pubescente superne ramoso, foliis ovato- aut ovali-oblongis utrinque acuminatis grossè serratis puberulis, capitulis corymbosis 40-45-floris, invol. squamis lanceolato-linearibus acutis, exterioribus subpatulis foliaceis obovato-linearibus, acheniis oblongis teretiusculis hispidulis, pappi serie ext. brevissima paleaceâ. ☉ in Indiâ orient. frequens. Ham. tr. lin. soc. 14. p. 217. Less. in linnæâ 1829. p. 293. *Conyza anthelmintica* Lin. sp. 1207. *Serratula anthelmintica* Roxb. cat. calc. 60. *Baccharoides anthelmintica* Moench. 578. *Ascaricida* India Cass. dict. 3. suppl. p. 38. *Serratula elegans* herb. Maurit. — Burm. Zeyl. t. 95. Rheed. mal. 2. t. 24. Cor. purpurascens. Variat caule foliisque plus minus hirsutis glabrativè. (v. v. et a. sp. è plur. Indiæ locis.)

266. *V. PAUCIFLORA* (Less. in linnæâ 1829. p. 293. 1831. p. 676), caule herbaceo ramoso teretiusculo pube brevi hirtello, foliis elliptico-lanceolatis utrinque acuminatis rariter serratis suprà sparse hirtellis subtus pubè brevi præsertim secus nervos scabris et subglandulosis, capitulis 40-45-floris solitariis terminalibus, invol. squamis ext. subulatis laxis interiores lineares 3-nerviis subaequantibus, achenio villosis, pappi setis barbellatis, serie externâ brevissimâ vix paleaceâ. ☉ in Senegaliæ planitie regionis Walo locis subhumidis legit cl. Perrotet. *Conyza pauciflora* Willd. sp. 3. p. 1927. Spreng. syst. 3. p. 514 excl. Link. syn. *Vernonia Senegalensis* Desf. cat. h. Par. 1829. p. 400. non Less. *Centropalus Galmensis*, Cass. dict. 7. p. 383. *Vern. Galmensis* Less. in linnæâ

ex iconibus citatis facillè restituenda. Variat caule glabro aut pubero, foliis basi obtusis (Less.) aut apice acuminatis. (v. v. et s.)

*ß. hebecarpa*, acheniis secus costas puberulis.  $\gamma$ . in Kentucky. Vern. altissima Rafin. in litt. 1831. Var.  $\alpha$  simillima acheniis exceptis. (v.  $\alpha$  sp.)

† *Non satis notæ.*

278. V. *ACHYROCOMA* (Less. in Linnæa 1829. p. 313), caule herbaceo striato tomentoso ramoso, foliis subsessilibus oblongo-lanceolatis serratis supra glabris subtus tomentosis, capitulis pedunculatis multifloris, invol. squamis coriaceis tomentosis apice glandulosis, ext. lanceolatis, int. oblongis obtusis, achenio villosio, pappi paleis pluriseriatis linearibus denticulatis exterioribus brevioribus, corollis apice glandulosis. — Patr. dub. fors. Amer. bor. Dispositio florum ignota. *Achyrocoma tomentosa* Cass. dict. 26. p. 21.
279. V. *AFRICA* (H. B. et Kunth nov. gen. 4. p. 39), fruticosa, ramulis subangulatis puberulento-tomentosis, foliis petiolatis ovato-oblongis acuminatis basi rotundatis integerrimis coriaceis supra glabris subtus hirtotomentosis, cymis terminalibus paniculæformibus foliatis, capitulis subsessilibus 5-6-floris, invol. squamis oblongis glabris, achenio glabro oblongo sulcato.  $\beta$  in Peruvia? cum V. floribunda à quâ vix differt nisi foliis basi magis rotundatis et acheniis minoribus.
280. V. *ANDROMACHIOIDES* (Less. in Linnæa 1831. p. 397 et 645), fruticosa, ramis diffusis teretibus striato-sulcatis superne hispido-puberulis, foliis petiolatis decussatis membranaceis triplinerviis subrotundis mucronatis basi obtusis subintegerrimis supra glabris subtus tenuiter villosio-tomentosis, paniculâ trichotomâ terminali ramis oppositis, capitulis 14-17-floris, invol. cylindraceis squamis glabris ciliatis ext. subrotundis acutis int. linearibus obtusis, recept. angusto fimbriifero, cor. glabrâ, achenio hirsuto, pappi ser. ext. brevi vix paleacea.  $\beta$  in sylvis Misantlie Mexici. Cor. latem. An ideo Vernonia vera?
281. V. *ARISTATA* (Less. in Linnæa 1829. p. 313), fruticosa? ramis....., foliis petiolatis lanceolatis apiculatis subdentato-sinuatis, dentibus spinuliformibus, supra sparse pilosis subtus villosis glandulosis, cymis scorpioides, capitulis subaggregatis uniseriatis folio florali suffultis, invol. squamis exter. apice subulatis rigidis. — Patr. ign. *Lepidoploea aristata* Cass. dict. 26. p. 17.
282. V. *EMARGINATA* (Wikstr. quad. fl. 73), fruticosa ramosa, ramis cæcâ villosis, foliis petiolatis ovalibus integerrimis apice emarginatis impresso-punctatis supra glabris subtus pubescentibus, paniculâ multicapitatâ ramis secundifloris, invol. ovatis squamis lanceolatis acutis extus villosis.  $\beta$  in Guadalupe. Affinis dicitur V. albicauli Pers. seu Vahlia Less. †.
283. V. *BACCHAROIDES* (H. B. et Kunth nov. gen. 4. p. 40), arborescens, ramulis angulatis hirtotomentosis, foliis petiolatis oblongis subacuminatis basi rotundatis remotè denticulatis supra pubescentibus tomentoso-hirtis, cymis terminalibus paniculæformibus ramosissimis, capitulis 3-5 aggregatis multifloris, invol. hemisphæricis squamis acutis glabris ciliatis.  $\beta$  in montosis Andium Novo-Granatensium. Cor. albæ glabræ. Pappus rufescens biserialis.
284. V. *CORDILOLIA* (H. B. et Kunth nov. gen. 4. p. 38), suffruticosa?, ramulis subangulatis molliter tomentoso-pilosis, foliis petiolatis ellipticis breviter acuminatis basi rotundatis integerrimis rigidis supra glabris nitidis subtus molliter piloso-pubescentibus, cymâ terminali paniculæformi patentissimâ, capitulis sessilibus, invol. semiglobosi squamis acutis subpungentibus.  $\beta$ ? in scopulis Novo-Granatensium.
285. V. *FRAGRANS* (Lallav. et Lex. nov. veg. descr. 1. p. 24), fruticoso-sarmentosa, foliis breviter petiolatis sublanceolatis integris margine revolutis scabrescentibus, cymâ apiciformi subsimplici, capitulis sessilibus unilateralibus distantibus multifloris, involucrio ventricosio.  $\beta$  ad vias in montosis prope S. Jose del Corral Mexicanorum. Flores albi vanillam olentes. Affinis videtur V. odoratissima aut Deppeana. †.

286. *V. FLOREBUNDA* (H. B. et Kunth nov. gen. 4. p. 38), fruticosa, ramulis angulatis hirsuto-tomentosis demum striatis glabris, foliis petiolatis oblongis acuminatis basi obtusis subintegerrimis coriaceis supra glabris subtus canescentibus hirsuto-tomentosis, cymis paniculæformibus axillaribus et terminalibus folio brevioribus, capitulis 4-5-floris subsolitariis, involucri oblongi squamis obovatis apice inciso-lacinatis?, achenio sulcato glabro.  $\S$  in Amer. merid. fortè in Peruvia. Cor. ign. Pappus duplex sed proportio serierum non indicata.
287. *V. FRUTICOSA* (Swartz fl. ind. occ. 1323), fruticosa, ramis teretibus..., foliis brevissimè petiolatis ovatis obtusis integris..., cymæ terminalis ramis elongatis flexuosis, capitulis sessilibus distantibus folio florali ovato brevioribus, .....  $\S$  in insul. Caribæis. Conyza fruticosa Lin. sp. 1209. — Plam. ed. Burm. t. 95. f. 1.  $\dagger$
288. *V. LINIFOLIA* (Blum. bijdr. ned. ind. p. 893), herbacea, ramis paniculatis subtrigosis, foliis lineari-lanceolatis rariter serrulatis puberulis, corymbo terminali dichotomo divaricato laxifloro, squamis involucri cuspidatis.  $\odot$ ? in montanis humidis ins. Javæ. Affinis dicitur *V. leptophyllæ*. An fortè eadem ac *V. erigeroides*?
289. *V. STELLARIS* (Lellav. et Lex. nov. veg. descr. 1. p. 23), arborescens, ramulis tomentosis, foliis brevè petiolatis lanceolatis supra pilosiusculis subtus fuscescenti-tomentosis margine revolntis subintegerrimis, cymæ terminalis scorpioides ramis tomentosis flexuosis, capitulis sessilibus, invol. campanulati squamis acutis pilosis, corollæ tubo incurvato, lobis conniventibus.  $\S$  in pascuis ad José del Corral Mexicanorum. Flores albi vanillati.  $\dagger$
290. *V. PROLIFERA* (Decaisn. n. ann. mus. h. n. Par. 3. p. 408), caule herbaceo striato puberulo, foliis subrotando-ovatis obovatis repando-subangulatis interdum oblongis obtusis subtus pallidioribus basi in petiolum attenuatis, paniculis terminalibus ramosis, ramis proliferis, invol. floribus dimidio breviori. — in insula Timor. Cl. auctor hic citat Conyzam proliferam Lam. et Sonchum volubilem Rumph., sed ex descr. icone et patriâ potius meo sensu ad Microglossam volubilem referendæ sunt.

### ++ Species vix notæ.

*Obs. 1.* Quid *Vernonia Misurica* et *Vernonia albiflora* (Raf. herb. p. 28)?

*Obs. 2.* Præter species hinc indè citatas ad *Vernoniam* referendæ videntur stirpes sequentes iconibus Floræ fluminensis incompletè cognitæ, nempe: *Chrysocoma arborea* S. t. 12, *C. repanda* t. 13, *C. paniculata* t. 14, *C. medio-serrata* t. 15, *C. cymosa* t. 16, *C. pilosa* t. 17, *C. sessilis* t. 20, *C. quadrangularis* t. 25, *C. linearis* t. 27, *C. horizontalis* t. 28, *C. herbacea* t. 29, *C. monantha* t. 33, *C. tomentosa* t. 38, *C. Prescæa* t. 39, et *C. alterna* t. 40.

### +++ Species exclusæ.

- V. AMYGDALINA* Dehil. = *Decaneurum amygdalinum*.  
*V. BRACTEATA* Wall. = *Decaneurum Silhetense*.  
*V. BRASILIENSIS* Less. = *Lychnophora brasiliensis*.  
*V. BRUNIOIDES* Less. = *Lychnophora brunioides*.  
*V. CAPITATA* Less. = *Albertinia capitata*.  
*V. CYMENSIS* Less. = *Cyanopsis pubescens*.  
*V. ECHINOPE* Mart. = *Echinopsidium*.  
*V. EXSUCCA* Mart. = *Echinopsidium exsuccum*.  
*V. FIMBRILLIFERA* Less. = *Decaneurum fimbriiliferum*.  
*V. HALIMODENDRON* Mart. = *Echinopsidium*.  
*V. HAKENFOLIA* Less. = *Lychnophora hakenefolia*.  
*V. INCANA* Less. = *Albertinia incana*.



- V. INCANESCENTS Mart. = *Albertinia incanescens*.  
 V. PINASTER Less. = *Lychnophora pinaster*.  
 V. PINIFOLIA Less. = *Webbia pinifolia*.  
 V. PLANTAGINIFOLIA Less. = *Pycnocephalum plantaginifolium*.  
 V. POLYGALÆFOLIA Licht. = *Pegolettia polygalæfolia*.  
 V. POPULIFOLIA Less. = *Distephannus popalifolius*.  
 V. PROTEÆFORMIS Less. = *Lychnophora proteæformis*.  
 V. PYCNOCEPHALA Mart. = *Echinopsidium pycnocephalum*.  
 V. SALICIFOLIA Less. = *Lychnophora salicifolia*.  
 V. SCARIOSEA Poir. = *Liatris scariosa*.  
 V. SCAPIGERA Less. = *Pycnocephalum scapigerum*.  
 V. SENEGALENSIS Less. = *Decaneurum Senegalense*.  
 V. SMITHIANA Less. = *Webbia Smithiana*.  
 V. SPATHULÆFOLIA Mart. = *Pycnocephalum spathulæfolium*.  
 V. STAAVIOIDES Less. = *Lychnophora staavioides*.  
 V. STACHYANTHUS Mart. = *Stachyanthus Martii*.  
 V. STYCHOCARPHA Spreng. = *Piptocoma lychnophoroides*.  
 V. VILLOSISSIMA Less. = *Lychnophora villosissima*.  
 V. WIGHTIANA Wall. = *Blumea vernonioides*.  
 X. DECANEURUM DC. in ann. bot. Guill. 1833. v. 2. p. 516. et in Wight contrib. bot. ind. p. 3.

Capitulum multiflorum homogamum, flor. tubulosus hermaphroditis 5-fidis. Recept. planum alveolatum. Invol. imbricatum pluriseriale, nunc bracteis foliaceis cinctum, nunc nudum, squamis ext. brevioribus aristato-cuspidatis, intimis apice scariosis exaristatis. Stigma Vernoniæ. Achænium nervis 10 prominulis angulatum sæpius glabrum. Pappus 1-serialis, setis crassistrigidis dense barbellatis, junioribus basi in annulum connatis, demum facile et segregatim deciduis. — Herbæ Indicæ aut suffrutices. Folia alterna petiolata dentata. Capitula sæpius ad apicem ramorum solitaria, rarius corymbosa pedunculata. Flores ex sicco purpurei. — Genus affine Vernoniæ, Cyanopidi et Centrathero.

#### Sect. I. PHYLLOCEPHALUM *Blum. bijdr.* 888.

Capitula bracteis foliaceis amplis arcuè cincta. Achænia basi non sensim attenuata.

1. D. FRUTESCENS, caule suffruticoso erecto hirsuto, foliis breviter petiolatis oblongo-lanceolatis basi attenuatis apice cuspidatis serratis rugosis supra puberulo hirtis subtus pallidis pubescenti-tomentosis, capitulis paucis pedunculatis bracteis. ♀ in uliginosis ins. Javæ, inter montes Barangrang et Tankawan Prabu, prov. Krawang legit cl. Blume. *Phyllocephalum frutescens* Blum. ! *bijdr.* 888. Bractes utrinque hirsutæ subtus albidæ scællæ lanceolatis, ext. capitulo longiores. Fol. 3-4 poll. longa, pollicem lata. (v. s. comm. à cl. Blume.)
2. D. RETICULATUM (DC. in Wight contrib. bot. ind. p. 3), caule suffruticoso ? erecto ramoso setis nudiquè scabro, foliis sessilibus ovatis mucronatis et mucronato-subdentatis supra asperis subtus dense albo-tomentosis, nervis venisque scabris reticulatis, pedunculis paucis axillaribus terminalibusve, capitulis bracteis pluribus foliaceis arcuè cinctis, invol. squamis internis chartaceo-scariosis glabris bracteis superantibus. ♀ ? in Ind. orient. ad montes Nealgerry legit cl. Wight. *Ampherephis Indica* Wall. ! cat. et herb. n. 3956. comp. n. 66. *Jacea* ? vulgo Kadourou Lesch. ! herb. n. 290. Capitula multiflora. Achænia glabra 10-costata. Pappus 1-ser. deciduus. (v. s.)

3. *D. MOLLE*, caule herbaceo erecto glabrescentulo aut hinc inde setoso-scabro apice tomentoso, foliis in petiolum attenuatis ovato-lanceolatis acuminatis grossè et irregulariter serratis suprâ setoso-scabris glabrisve subtus albotomentosis, pedunculis axillaribus 1-cephalis, bracteis foliaceis capit. approximatis oblongis subtus tomentosis, invol. squamis internis chartaceo-scariosis glabris subscutis. — in Ind. orient.

*a. epileium*, foliis suprâ levibus. — Ad Kortallum distr. Carnatici leg. cl. Hayne et Wight. Decan. epileium DC. in Wight contrib. bot. ind. p. 7. Ampherephis mollis Wall. ! cat. et herb. n. 3957. comp. 67. Wightia formosa Spreng. in litt. ad Wall. Folia 3-3½ poll. longa, sesquipoll. lata. (v. s.)

*β. scabridum*, foliis suprâ scabris. — Wight cat. n. 1392. — *D. scabridum* DC. l. c. (v. s.)

Sect. II. GYMNANTHEMUM. — *Gymnanthemum spec. Cass. bull.*

*philom. 1817. dict. 20. p. 108.*

Capitula aut omninò ebracteata aut bracteolis paucissimis invol. squamas inferiores referentibus subinstructa. Achenia elongato-turbinate seu basi attenuata.

4. *D. GLABRUM*, caule suffruticuloso tereti glabro striato, foliis breviter petiolatis ovalibus basi cuneatis apice acuminatis distanter repando-serrulatis glabris cum nervis utrinquè tenuè puberulis, corymbis 2-5-cephalis, invol. squamis adpressis acutis (non aristatis) flocculis multò brevioribus. *♂* in Ind. orient. *Conyza* Wall. ! cat. n. 3283. comp. n. 393. *Eupatorium glabrum* Heyn. ! herb. Folia 2-3 poll. longa, 8-10 lin. lata. Pappus rufus. Achenium teres 10-costatum basi attenuatum glabrum 2 lin. longum. (v. s. in b. Can. Ind. or. nunc Soc. lin. Lond.)

5. *D. SILBERTENSE*, caule herbaceo erecto ramoso, foliis breviter petiolatis oblongo-lanceolatis utrinquè acuminatis remotè aristato-serratis suprâ ramisque glanduloso-scabridis subtus secus nervos scabridis, capitulis ad apices ramorum subsessilibus subcorymbosis, invol. squamis inter. oblongis mucronatis, exter. filiformi-subulatis subpatulis nonnullis secus pediculum ortis. *♀* in Ind. or. prov. Silhet ad montes Pandua legit Fr. de Sylva. *Vernonia* ? bracteata Wall. ! cat. n. 2921. comp. n. 31. Achenium 10-costatum basi attenuatum glabrescentulum 2 lin. longum. Pappus rufus 1-serialis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

6. *D. DENDIGULUM* (DC. in Wight contrib. bot. ind. p. 7), collo tuberoso suffruticuloso, caule herbaceo erecto parè ramoso tereti basi striato pubescente apice tomentoso, foliis subsessilibus latè ovatis subrotundisve coriaceis dentatis suprâ glabris asperulis subtus densè albo-tomentosis, summis ovalibus acutis parvis, corymbo fastigiato composito, invol. ovatis squamis ovalibus mucronatis imbricatis extus tomentoso-lanatis intus glaberrimis. *♀* in Indiæ Peninsula montibus Dendigul legit cl. Wight. *Conyza indica* Wall. ! cat. n. 3042. comp. 152. non Blum. Folia inf. 4 poll. longa, 3 poll. lata. Achenium nervis 10 geminatis elevatis angulato-striatum glaberrimum inter nervos glandulosum. Pappus 1-serialis rigidus (v. s. in b. Wight. et comm. ab hon. Cur. merc. angl. Ind. or.)

7. *D. ? OBOVATUM*, caule suffruticuloso scandente tereti, foliis petiolatis obovatis obtusè integerrimis suprâ glabris subtus ramulisquè pube minimâ subpubescentibus, paniculâ ovatâ, invol. squamis ovato-oblongis brevissimè mucronatis. *♂* in Rawak Molaccensium. *Gymnanthemum obovatum* Gaudich. ! voy. oran. bot. p. 471. Vern. cuneata Less. in Linnæa 1831, p. 644. Folia in sicco margine subrevoluta 2-3 poll. longa, 12-15 lin. lata. Capitula 6-8-flora. Achenium glabrum versus apicem tumidum 10-costatum basi attenuatum, fortè in nostr. spec. sterile. Pappus rufus 1-serialis scaber. An omninò priorum congener. (v. s. comm. ab. am. Gandichaud.)

8. *D. GRANDE*, fruticosum glabrum, ramis teretibus apice summo pulverulento-puberulis, foliis longè petiolatis ovalibus utrinquè acuminatis integerrimis

subtus seors nervos minutissimè puberulis, corymbo composito polycephalo laxo fastigiato, invol. ovati squamis adpressis glabris, ext. acuminatis int. obtusis fere truncatis, acheniis glabris.  $\frac{5}{3}$  in insulæ Madagascar orâ occidentali circa pagum Majangay in sinu Bombatoc legit cl. Bojer! *Vernonia grandis* Bojer! in litt. 1835. Frutex 15-18-pedalis. Flores lilacini, in sicco albidii, in invol. 9-10. Pappus rutilans 1-serialis crassiusculus. Achen. junius teretiusculum. Valdè affinis *D. Senegalensis*. (v. s. sp. comm. à cl. Bojer. et in ins. Maurit. ex sem. Bojerianis cult. comm. à cl. L. Bontou.)

9. *D. SIMBRILLIFERUM*, fruticosum, ramis tomentoso-pubescentibus, ultimis seu floralibus compressis, foliis obovato-lanceolatis breviter acuminatis in petiolum attenuatis vix apice subdenticulatis suprà glabris subtus seors nervum pubero-hirtellis, corymbo amplo composito aphylo, capitulis pedicellatis obovato-cylindraceis 3-4-floris, invol. pauciserialis squamis ovatis pubescentibus inter. longioribus, recept. breviter simbrillifero, acheniis subhispidulis, pappi seriebus satis inæqualibus.  $\frac{5}{3}$  in insulæ Bourboniæ sylvis frequens (Bontou!). *Gymnanthemum simbrilliferum* Cass. dict. 20. p. 109. *Vernonia simbrillifera* Less. in linnæa 1829. p. 313. *Eupatorium coriaceum* Vahl symb. 3. p. 95. non Spreng. Cl. Cassini dixit fl. flavos, sed sæpius dixit flores omnes decoloratos. Pappus intensè rufus. (v. s. sine cor. comm. à cl. Bontou.)

10. *D. SENEGALENSE*, fruticosum, ramis subangulosis tomento brevissimo subvelutinis, foliis petiolatis ovato- aut oblongo-lanceolatis subintegerrimis suprà glabriusculis subtus villosa-pubescentibus, corymbo fastigiato nudo, invol. campanulati squamis ovatis ciliatis, ext. brevioribus mucrone acutis, inter. longioribus obtusis demum stellatis, achenio maturo glabro 10-costato inter costas glanduloso.  $\frac{5}{3}$  in Senegaliâ (Roussillon! Sparman!) *Baccharis Senegalensis* Pers. ench. 2. p. 424. *Gymnanthemum cupulare* Cass. dict. 20 p. 109. *Eupatorium coloratum* Willd. sp. 3. p. 1768 ex Less. *Vernonia Senegalensis* Less. in linnæa 1829. p. 265. 1831. p. 641. non Desf. *Conyza rutilans* Poir. suppl. 2. p. 341? *Chrysocoma amara* Schum. in act. Dan. p. 157. ex Less. Cor. alba. Pappus rufo-aureus rigidus scaber. Huc referendæ mihi videntur: *Conyza radiata* in argillosis circa Albreda ad fl. Gambie, C. aurea in insulâ prope Saint-Louis, et C. frutescens in peninsulâ Promontorii-Viridis prope Dakar, à cl. Perrotet collectæ et mecum commun. Variat squamis invol. plus minus ciliatis, imò subciliatis, foliis subrotundis, obovatis, obovato-oblongis aut oblongo-lanceolatis. (v. s.)

11. *D. AMYGDALINUM*, fruticosa, ramis teretibus apice pube brevi scabriusculis, foliis oblongo-lanceolatis petiolatis utrinquè acuminatis, adultis glabris, corymbi paniculati ramis dichotomis, ospitalis pedunculatis, involucri semiglobosi squamis obtusis, achenio villosa-hispido 10-nervio.  $\frac{5}{3}$  in Nubiâ ad Fazogl, ubi dicitur Kering et arabicè Kikir. *Vernonia amygdalina* Delil. in Cail. voy. et cent. n. 21. forsan in Sierra-Leonâ si specim. quoad patr. incertum huc ritè refero. Ex eo affinis videtur *D. Senegalensis*. (v. s.?)

12. *D. DIVERGENS* (DC. in Wight contrib. bot. ind. p. 8), caule herbaceo erecto tenuissimo velutino scabro paniculato ramoso, foliis brevissimè petiolatis ellipticis utrinquè acuminatis dentatis suprà glabris subtus puberulis reticulatis subrugosis, paniculæ ramis elongatis divergentibus foliosis ipsis paniculatis, capitula ad apicem ramulorum confertis 7-8-floris, invol. squamis oblongis acutè mucronatis glabriusculis, achenio glabro glanduloso.  $\frac{2}{4}$  in Indiâ orient. verolim. Maderaspatanâ (h. Wall.!), in Pegu (Belanger!) *Conyza divergens* Wall.! cat. n. 3027. comp. 137. excl. litt. E. *Vernonia* Wight herb.? Affinis *D. aspero*. Pappus rufus, ext. brevior paucisetus caducissimus. Folia 3 poll. longa, pollicem lata. An potiùs *Vernoniæ* sp. V. multifloræ proxima? (v. s.)

*ß. lanceolata*, foliis planis vix subrugosis capitulis paulò minoribus. — in regione Birmanorum ad Taong-Dong legit cl. Wallich. *Conyza lanceolata* Wall.! cat. et herb. n. 3059. comp. 169. cat. Birm. n. 916: Paniculâ minus divaricatâ (v. s. comm. cum priorè ab. hon. Cur. merc. angl. Ind. or.)

XI. CYANOPIS Blum. in litt. 1831. — *Cyanthillium* Blum. (1826) *bijdr.* p. 889. — *Isoneuma* Cass. *bull. philom.* 1817. *dict.* 24 p. 25. non Brown. (1810). — *Centratheri* sect. Less. in *linnæa* 1829. p. 320. — *Vernonia* spec. Less. *syn.* 146. — *Conyza* spec. Ait. *Lam.*

Capitulum multiflorum homogamum, fl. omnibus tubulosis inæqualiter 5-fidis hermaphr. Recept. alveolato-subfimbriiferum planum. Invol. hemisphærici squamæ pluriseries imbricatæ lanceolatæ rigidularistatæ apice non adpressæ. Stigm. Vernoniæ. Achæmium prismaticopentagonum glabrum, areolâ terminali latâ subcupulatâ. Pappus 1-serialis, setis rigidis imâ basi distinctissimis deciduis subscabris. — Herbæ Indicæ erectæ ramosæ verisim. annuæ. Folia petiolata ovata aut lanceolata alterna supra glabra subtus plus minus villosa. Capitula ad apices ramorum pauca pedicellata ebracteata corymbosa. Flores cœrulescentes, undè nomen ab ipso cl. auct. paulò mutatum ut habitum melius exprimat. — Genus Vernoniæ affine.

1. *C. villosa* (Blum. *bijdr.* 889 sub *Cyanthillio*), foliis petiolatis ovato-oblongis infra canescenti-villosis margine subrepando-serrulatis, summis integerrimis, corymbis 3-4-cephalis, involucri floribus subæquali. ☉ in graminosis insula Javæ circa Batavia. Proximè fortè nimis ad sequentem accedit. (v. s. comm. à cl. Blume.)
2. *C. rotundifolia* (Blum. *bijdr.* 890 sub *Cyanthillio*), foliis petiolatis ovatis utrinquè acutis subtus pubescentibus margine repando-macronalato-denticulatis, corymbis 3-4-cephalis, involucri floribus subæquali. ☉ in graminosis insula Javæ cum priori legit cl. Blume! in Manilla et Gushan (Cham.), in China (Ait.), in Marianna (Hank.), in China et Pegu (Less.), in Pegu ex cl. Belanger, ad Tavoy et Rangoon Ind. Birmanicæ ex cl. Wallich. *Conyza punctulata* Wall! cat. et herb. n. 2995. comp. 105. birm. 1167. *Synanthera* Wall! cat. n. 7136. *Conyza patula* Ait. Kew, 3. p. 184. *Conyza Chinensis* Lam. *dict.* 2. p. 83. non Linn. nec Lour. *Isoneuma ovata* Cass. *dict.* 24. p. 25. *Serratula* Mill. ic. t. 247. *Centratherum* Chinense Less. in *linnæa* 1829. p. 320. *Vernonia Chinensis* Less. in *linnæa* 1831. p. 674. Achænia glabra obpyramidata 4-5-angulata. Variat invol. glabrato aut pubero, foliis subtus plus minus pubescenti-villosis. Specim. India contin. ferè media inter spec. hujus et prioris Blumeana et fortè in unicam colligunt. (v. s.)

β. *microcephala*, capitulis dimidio minoribus. ☉ in insula Mauritiæ introd. et quasi indigena facta ex Bojer in litt. An species propria? (v. s.)

3. *C. Moluccensis* (Blum. *bijdr.* 890 sub *Cyanthillio*), foliis breviter petiolatis lanceolatis utrinquè acuminatis subtus albedo-villosis margine repando-mucronato-dentatis, capitulis geminis confertisve, invol. squamis glaberrimis longè aristatis flosculos subæquantibus. ☉? in insulis Moluccanis. Fol. 3 poll. longa, pollicem lata. (v. s. comm. à cl. Blume.)
4. *C. Madagascarensis*, caule herbaceo erecto divaricato-ramoso tereti glabro, foliis ellipticis utrinquè acuminatis grossè serratis basi integerrimis cuneatis subpetiolatis sparsè scabridis subtus punctulatis, capitulis pedicellatis ebracteatis, invol. squamis glaberrimis acuminatis, achæniis seriato-sulcatis glabris, pappo paucisetis caducissimo. — in insula Madagascar in pratis et ad margines sylvarum legit cl. Bojer. *Ethulia conyzoides* Bojer! in litt. 1835. Flores rosei. (v. s. comm. à cl. inv.)

† Species exclusa.

C. ERIGEROIDES DC. = *Vernonia erigeroides*.

XII. CENTRATHERUM Cass. bull. philom. febr. 1817. dict. sc. nat. 7. p. 383. *journal. phys.* 89. p. 24. — *Ampherephis* H. B. et Kunth nov. gen. am. 4. (1818). p. 24. Less. syn. p. 148. — *Spixia* Schrank pl. rar. h. mon. 2. (1819). t. 80. non Leand. — *Centratheri* sect. 1. Less. — *Amphirephidis* sect. 1. Nees et Mart. act. soc. cur. nat. 12. p. 4.

Capitulum multiflorum æqualiflorum. Involucrum imbricatum campanulatum quasi duplex nempe, foliis floralibus magnis cinctum, foliolis inv. veri siccis exter. brevioribus. Receptaculum planum nudum. Cor. regularis 5-fida, tubo gracili lobis acuminatis glabris longiore. Antheræ incluse. Filam lævia. Styli rami semisubulati. Achænium oblongum subcylindraceum glandulosum glabrum, collo apicali parum prominulo. Pappus 1-serialis, setis serrulatis inæqualibus rigidis caducissimis! — Suffrutices aut herbæ. Folia alterna ovata aut oblonga serrata. Capitula ramos terminantia, pedunculata. Flores purpureo-cœrulei. — Cl. Cassini in *journal. phys.* l. c. asserit *Centratherum* idem esse ac *Ampherephim*, et in tabl. comp. 1829., n. 36 et 37, genera duo admittit ex invol. squamis muticis aut spinoso-aristatis.

1. C. INTERMEDIUM (Less. in *linnæa* 1829. p. 320), caule herbaceo, involucri veri squamis integerrimis, exterioribus apice aristatis, interioribus ovato-oblongis obtusis. ☉ in Brasiliâ (Mart.), in cultis incultisque ad Bahiam (Salzm. ! Blanch. !). *Ampherephis intermedia* Link abbild. 5. t. 29. Colla hort. rip. app. 3. p. 26. *Spixia violacea* Schrank pl. rar. h. mon. t. 80. Flores coruleo-subviolacei. Cor. tubus puberulus. (v. v. cult. in h. Genev. et a. sp. à Bahiâ.)
2. C. PUNCTATUM (Cass. dict. 7. p. 384. Less. l. c.), caule fruticoso, involucri veri squamis asperne eroso-denticulatis apice spinoso-dentatis. § in ripâ flum. Orinoci ad pagum Mâypures. *Ampherephis aristata* H. B. et Kunth nov. gen. 4. p. 25. t. 344. Cor. purpurea et ex cl. auct. glabra. Folia lanceolata. (v. s.)
3. C. MUTICUM (Less. l. c.), caule fruticoso, invol. veri squamis denticulatis apice rotundatis mucronatis. § in temperatis Novæ-Andalusiæ ad montem Cocollar. *Ampherephis mutica* H. B. et Kunth nov. gen. 4. p. 26. t. 315. Folia oblonga basi cuneato-angustata.

† *Species exclusæ.*

C. CHINESE Less. = *Cyanopsis pubescens*.

AMPHEREPHIS INDICA Wall. = *Decaneurum reticulatum*.

AMPH. MOLLE Wall. = *Decaneurum molle*.

AMPH. PSILOCARPA N. et Mart. — *Oioospermum involucreatum*.

AMPH. WIGHTIANA Wall. = *Oioospermum Wightianum*.

XIII. BECHIIUM.

Capitula multiflora homogama, floribus tubulosis 5-fidis, ext. subpalmatis. Recept. alveolare. Invol. squamæ 2-3-seriales subimbricatæ longitudine subæquales, inter. angustiores glabræ apice coloratæ. Styli rami elongati hispidi. Achænium teretiusculum villosum. Pappus uniserialis pilosus scabridus. — Herbæ Madagascarienses. Folia alterna aut radicalia ovali-oblonga subsessilia subintegerrima supra pilososcabra. Capitula terminalia corymbosa. Invol. squamæ apice subcariosæ purpureæ. Flores purpurei. — Nomen apud Plinium olim Tussilagini impositum admisi.

1. B. RUBRICAULE, caule pubescente imâ basi hirtello ad medium folioso, foliis ovali-oblongis basi cuneatis subpetiolatis utrinque puberulis, corymbo composito polycephalo, invol. squamis dorso villosis-hirtis. ♀ in insulâ Mada-

gascar prov. Emirnensi legit cl. Bojer. Herba pedalis. Caulis purpurens. Folia 3 poll. longa  $\frac{1}{2}$  poll. lata obscure virentia. Corymbi rami dense-villosi, (v. a. comm. à cl. inv.)

2. *R. SCAPIFORME*, caule nodo pube sublanatâ et pilis apice glandulosis in lanâ intermixtis hispidis, foliis radicalibus ovali-oblongis basi cuneatis subsessilibus supra hirsuto-scapris subtus canis sericeo-tomentosis, capitulis circ. 3 terminalibus subsessilibus, invol. squamis dorso glanduloso-hirtis.  $\frac{1}{2}$  in montibus nudis et siccis prov. Emirnensis Madagascariensium legit cl. Bojer. (v. a. comm. à cl. inv.)

XIV. *STOKESIA* Lher. sert. 27. DC. ann. mus. 16. p. 154. Cass. dict. sc. nat. 51. p. 64. Less. in linn. 1829. p. 321. syn. p. 148. — *Cartesia* Cass. bull. philom. 1816. p. 198. — *Carthami* sp. Hill.

Capitulum multiflorum radiatiforme. Involucrum subglobosum basi bracteatum pluriseriatum imbricatum, squamis adpressis, exterioribus ovalibus in append. foliaceam patulam ciliato-spinulosam desinentibus, interioribus oblongis ciliolatis. Recept. carnosum nudum. Cor. palmatæ glandulis conspersæ, exteriores capituli majores intus profundius fissæ. Anth. inclusæ. Styli rami semi-subulati. Achænium brevè tetra- (rarius tri-) gonum, areolâ terminali latâ 4-gonâ. Pappus 1-serialis 4-5-paleaceus, paleis deciduis longis planis scariosis integerrimis. — Herba erecta ramosa glaucescens glabriuscula, caule apice villosotomentoso. Folia alterna, infer. ovalia integerrima in petiolulum longè attenuata, super. semi-amplexicaulia lanceolata ad basin serrato-spinosa. Capitula ramos terminantia solitaria. Flores cœrulei.

1. *S. CYANEA* (Lher. sert. angl. 28).  $\frac{1}{2}$  in Carolinâ australi. Videtur rara et à cl. Elliott non reperta. *Carthamus lævis* Hill. Kew. 57. t. 5. *Cartesia centauroides* Cass. bull. 1816. p. 198. *Carthamus Carolinianus* Michx. in h. Mus. Par. (v. a. ex herb. Lher.)

XV. *PLATYCARPHA* Less. in linnæa 6. p. 688. syn. p. 148. — *Cynara* Thunb. fl. cap. 622. non Linn. — *Stobæa* sp. Spreng.

Capitulum homogamum multi- et æqualiflorum. Involucri squamæ pluriseriales lanceolatæ elongatæ submembranacæ integræ acutæ, interiores angustiores recept. paleas ferè simulant. Cor. tubo longissimo extus præsertim ad apicem hispidulo, laciniiis 5-linearibus æqualibus. Stam. filam. glabra lævia. Antheræ apice obtusæ et subæqualiter appendiculatæ basi ecaudatæ. Pollen globosum. Stigm. 2 longa apice divergentia teretiuscula undiquè ferè papillosa. Fructus glaber subcylindraceus. Pappus persistens paleis 7-9 scariosis albis linearibus acuminatis integerrimis constans. — Herba perennis. Folia radicalia longa petiolata pinnatipartita subtus albo-tomentosa supra et in petiolo arachnoidea, lobis ovatis dentatis, inferioribus dissitis parvis, extremis majoribus confluentibus. Capitula plurima ad collum sessilia congesta. Cor. violacæ. — Genus distinctissimum!

1. *P. GLOMERATA*.  $\frac{1}{2}$  ad Cap. Bonæ Spei (Th. Dreg!) in distr. Uitenhagen (Eckl. et Zeyh.) juxta Swartkopp' rivier. *Cynara glomerata* Thunb. prodr. cap. 141. flor. cap. 622. Willd. sp. 3. p. 1692. *Stobæa glomerata* Spreng. syst. 3. p. 394. (v. a.)

XVI. *ODONTOCARPHA*. — Gen. nov. Papp. n. 3.

Capitulum pluriflorum homogamum. Recept. angustum ebracteolatum. Invol. oblongo-cylindricum, squamis imbricatis adpressis. Cor. tubus pappo brevior, faux oblonga tubo latior, limbus 5-lobus acu-

tiusculus. Stam. filam. glabra? brevia; antheræ ecaudatæ. Styli rami breviter exserti subulati hispiduli. Achænium villis sericeis adpressis brevibus munitum subcylindraceum striatum. Pappus 1-serialis persistens, paleis 7-8 lanceolatis acutis apice serrulatis. — Herba suffruticulosa gracilis Chilensis erecta ramosa, ramis apice 1-cephalis. Folia alterna angustè linearia acutiuscula enervia præsertim ad basin ciliolata. Capitula foliis parvis raris substipata oblonga. Cor. vix exsertæ ex sicco purpurascens. Invol. fortè viscosa. — Car. ferè Platycarpæ, sed habitus diversissimus, cor. tubus brevissimus, pappi paleæ denticulatæ, etc.

1. O. PÖPPIGII. ♂ in Chili legit cl. Pöppig. Folia pollicem longa semilineam lata. Capitula 3-4 lin. longa. (v. s.)

## XVII. WEBBIA.

Capitula pluriflora discoidea dioica! Recept. alveolatum, alveolorum marginibus subexsertis erosis. Invol. imbricatum 3-5-seriale disco brevius. — ♂ Cor. tubulosæ breviter 5-dentatæ. Antheræ inclusæ lineares. Stylus indivisus aut apice subfissus. Ovarium extus villosum abortivum. Pappus pauciserialis, setis vix dentatis. — ♀ Cor. tubulosæ in lobos 5 elongatos angustos crassiusculos divisæ. Anth. o, aut effatæ. Stylus è tubo exsertus, lobis corol. brevior, ramis 2 elongatis crassiusculis extus et styli partè indivisâ supremâ pilosiusculis. Achænium teres seu turbatum pluricostatum erostre, costis villosissimis, sulcis glandulosis. Pappus pluriserialis, setis crassiusculis densè barbellatis exter. brevioribus. — Herba suffruticosæ Africanæ, caule erecto basi simplici apice corymboso. Folia sparsa linearia marginibus parcè revoluta suprâ glabriuscula subtus cum caule involuacroque canescentia. Flores purpurei in capitulo 8-10. Pollen flavum. Stigmata purpurea. — Dixi in honorem cl. bot. angl. P. B. Webb, Orientis, Africæ borealis et insularum Canariensium strenuo observatori.

1. W. SERRATULOIDES, caule suffruticoso ramoso pubescente, espitulis ad apicem ramorum solitariis hemisphericis, invol. squamis adpressè imbricatis 5-seriatis, ♂ in Senegambis locis arenosis ad Albreda secus flumen Gambie Martio ferè defloratam legit cl. Perrottet. Folia vix pollicem longa, semilineam lata. Capitulum 6 lin. diam. Serratula aut melius Jurinea capitula referens. (v. s. specim. form. à cl. Perrottet comm.)
2. W.? SMITHIANA, caule herbaceo striato sericeo-argenteo simplici densè foliato, foliis linearibus subtus sericeo-incanis, paniculâ subfastigiata, invol. campanulati squamis 2-3-seriatis 3-nerviis emucronatis, achénio pubescente, pappi ser. ext. dimidio interiore brevior. ♀ ad Cap. B. Spei (Krebel) in montibus Langebloof et Krantzrivier (Th. Eckl. n. 352. et Zeyh.), in Coloniæ distr. orientalibus (Burch. l. ont. geog. n. 3840 et 7434), ad Willbergen, Zw. Key et Basche (Dreg.) in Cafferland et Uitenhagen (Eckl. l. et Zeyh.). Conyza canescens Thunb. fl. cap. 665. Conyza pinifolia Lam. dict. 2. p. 86. excl. syn. Seb. (1). Erigeron Capense Houtt. l
3. W. PINIFOLIA, caule herbaceo simplici striato subcanescente, foliis semilobis confertis 1-nerviis linearibus margine revolutis suprâ glabris subtus sericeo-incanis, corymbo polycephalo fastigiato, invol. squamis oblongo-ovatis subcanescentibus, achénio sericeo-piloso, pappi serie externâ dimidio interiore brevior. ♀ ad Cap. B. Spei (Krebel) in montibus Langebloof et Krantzrivier (Th. Eckl. n. 352. et Zeyh.), in Coloniæ distr. orientalibus (Burch. l. ont. geog. n. 3840 et 7434), ad Willbergen, Zw. Key et Basche (Dreg.) in Cafferland et Uitenhagen (Eckl. l. et Zeyh.). Conyza canescens Thunb. fl. cap. 665. Conyza pinifolia Lam. dict. 2. p. 86. excl. syn. Seb. (1). Erigeron Capense Houtt. l

(1) Icon enim Sebana (mus. i. t. 23. f. 3) hic à Lamakio citata non malè habitum nostræ plantæ indicat, sed ex involucri ad Helichrysum intermedium referenda videtur.

pl. syst. x. bl. 629. t. 69. f. 2. ex herb. Deless. *Vernonia pinifolia* Less. in Linnaea 1829. p. 257. 1831. p. 639. Caules sesqui- aut bipedales. Fol. 2-3 poll. longa, 1-2 lin. lata, 1-nervia. Capitula 10-20 laxè corymbosa. (v. s.)

4. *W. ARISTATA*, caule herbaceo simplici striato subcanescente, foliis subcaulibus lanceolatis acuminatis basi attenuatis penninerviis integerrimis planis utrinquè aut subtùs tantùm pube confertà sericè adpressà canis, corymbo polycéphalo fastigiato, invol. squamis subcanescentibus lanceolatis in acumen aristæforme productis, achenio sericeo-villoso, pappi serie externà triplò interiore brevior. 2½ ad Cap. Bonæ Spei ad Katrivier (Dreg. sp. exs.), ad Zw. Key et Basche (Dreg.!), in territorio cesso et Kafferland (Eckl. et Zeyh.). Affinis *W. pinifolia* sed folia ferè *Salicis albe* nunc utrinquè nunc subtùs tantùm sericea. (v. s.)
5. *W. ELAAGHOIDES*, caule herbaceo erecto striato subcanescente, foliis breviter petiolatis ellipticis acutis planis integris suprà glabris subtùs pube sericè adpressà confertà canis, corymbo polycéphalo fastigiato, invol. squamis canescentibus oblongis mucronato-acutis, achenio villosio, pappi serie externà triplò interiore breviori. 2½ in Africà australi ad Zw. Omsamwubo et Omsanculo legit cl. Drega! Folia 18-20 lin. longa, 9-10 lata. (v. s.)
6. *W. OLIGOCEPHALA*, caule herbaceo simplici striato glabresculo, foliis breviter petiolatis ovalibus abruptè acuminatis integris planis penninerviis suprà glabris subtùs pube sericè confertà adpressà canis, corymbo 3-cephalo, invol. squamis subcanescentibus lanceolatis acutis, achenio tereti villosio, pappi serie externà triplò interiore breviori. 2½ in Africà australi ad Zw. Omsamculo et Port Natal legit cl. Drega! Priori affinis sed achenium duplò longius. Caulis multò tenuior. Folia pollicem longa, 8-10 lin. lata, etc. (v. s.)
7. *W. HIRUTA*, caule herbaceo simplici striato breviter hirsuto, foliis sessilibus oblongis basi cordatis apice mucronato-acuminatis suberratis suprà glabresculis subtùs hirsutis reticulato-nervosis, corymbo polycéphalo conferto ramulis hirsuto-tomentosis, invol. squamis subcanescentibus lanceolatis acutis, acheniis villosis, pappi serie brevissimà. 2½ in Africà australi ad Tambukiland et Cafferland (Eckl.!) n. 725 et 1318, ad Katriviersberg et Port Natal (Dreg.!). Antheræ in fl. locm. effusæ. (v. s.)
8. *W. NUDICAULIS*, caule herbaceo subnudo striato glabresculo, foliis radicalibus obovatis basi attenuatis mucronulatis glabris penninerviis planis subdenticulatis, caulinis remotis linearibus erectis margine revolutis integerrimis, corymbo laxo, invol. squamis canescentibus oblongis mucronatis, acheniis villosis, pappi serie externà brevissimà. 2½ in Africà australi ad Zw. Key et Basche legit cl. Drega! Folia radice. 1½ poll. longa, 9-10 lin. lata; caulina ferè pollicem longa, vix lineam lata. Flores ferè *W. pinifolia*. (v. s.)

#### XVIII. HOPLOPHYLLUM. — *Pteroniæ* sp. *Lin. f. Thunb.*

Capitula 5-flora homogama. Invol. ovato-oblongi squamæ latæ imbricatæ obtusæ apice spinuloso-mucronatæ. Recept. angustum fimbriatiferum. Cor. glabræ tubulosæ profundè in lobos lineares æquales 5-fissæ. Antheræ ecaudatæ. Filamenta brevina glabra. Styli à medio setis erectis hispidi, ramis longissimis per totam longitudinem hispidis. Achenia crassa subtrigona villosissima apice glabra et sub pappo paululum contracta. Pappus multiserialis persistens, squamellis rigidulis exter. lineari-filiformibus, inter. basi membranaceis latoribus sensim acuminatis. — Suffrutex Capensis glaber rigidus, ramis teretibus striatis. Folia alterna rigida spinescentia integerrima aut hinc indè ad basin spinuloso-dentata crassiuscula patula utrinquè ut rami regulariter striata. Capitula in axillis supremis sessilia solitaria sed ob folia approximata 4-5 congesta capitata. Cor. pallida.



1. *H. spinosum*.  $\zeta$  in Africâ Capensi in Carro (Thunb.), ad Olifantivier in regione Gariepina (Dreg!), ad Clan William (Eckl.! n. 1420. et Zeyh.). *Pteronia spinosa* Linn. f. suppl. 357. Thunb. fl. cap. 629. (v. s.)

**XIX. PIPTOCOMA** Cass. bull. philom. 1817. p. 10. 1818. p. 50. dict. sc. nat. 41. p. 111. Less. in linnæâ 1829. p. 315. syn. p. 147.

Capitulum pluriflorum æqualiflorum. Involucrum ovato-cylindricum imbricatum, foliolis coriaceis siccis ovato-aut obovato-oblongis. Recept. nudum angustum. Cor. regularis glabra, tubo longo, limbo 5-fido, lobis acuminatis glandulosis parte integra brevioribus. Anth. inclusæ. Achæ-nium obcompressum angulatum glabrum, disco epigyno magno, nectario styliformi. Pappus biserialis (ferè *Heterocomæ*), serie externâ brevissimâ cartilagineâ crenatâ, interiori oo-paleacâ, paleis caducis longis planiusculis subserratis sæpè tortis. — Suffrutices Americani. Rami numerosi teretes tomentosi. Folia alterna integra 1-nervia suprâ glabra subtus tomentosa. Capitula plurima ad apices ramorum subsessilia ad-gregata in glomerulum subumbellatum. Flores purpurei aut rosei.

1. *P. avfescens* (Cass. II. cc.), foliis brevè petiolatis ovatis suprâ scabris subtus subreticulato-tomentosis, capitulis 12-floris.  $\zeta$  in Sancto Domingo. *Achænia* 5-gona striata. Cor. purpureæ.  
2. *P. Lychnophoroides* (Less.! l. c. f. 7. 8. 10. 12), foliis sessilibus linearibus suprâ lucidis subtus tomentosis margine revolutis, capitulis 3-4-floris.  $\zeta$  in Brasiliâ tropicâ. *Vernonia trichocarpa* Spreng. syst. 3. p. 437. ex Less. Folia ferè Phylcæ, 6 lin. longa lineam lata, coriacea. Glomeruli foliis cincti. Cor. roseæ. *Lychnophoræ* fortè nimis affinis. (v. s. comm. à Mus. reg. Berol.)

**XX. DISTEPHANUS** Cass. bull. philom. 1817. 76. dict. 13. p. 361. — *Vernonia* sp. Less. syn. 146.

Omnia *Vernoniæ*, sed involucri squamæ apice appendiculatæ. Invol. areolatum, areolis in dentes seu fimbrillas breves exsertis. Pappi series utraque constans paleis circ. 10-lamellatis coriaceis, exterioribus nunc interiores subæquantibus nunc dimidio brevioribus latioribus denticulatis, interioribus linearibus barbellatis. — Frutices Mauritanii seu Madagascarienses, ramulis foliisque tomentosis alternis integerrimis. Capitula densè corymbosa.

§. 1. *Pappi serie externâ dimidio quam interior breviori.* — Mauritanii.

1. *D. populifolius* (Cass. l. c.), foliis petiolatis latè ovatis acuminatis, corymbo 7-8-cephalo, invol. squamis dorso præsertim ad margines lanatis, capitulis 30-40-floris.  $\zeta$  in insulâ Mauritiî (Comm. ! Bory !), locis aridis elatis ventosis (Boaton !) ferè semper florens (Bojer !). *Conyza populifolia* Lam. ! dict. 2. p. 87. Willd. sp. 3. p. 1943. *Vernonia populifolia* Spreng. syst. 3. p. 434. Less. in linnæâ 1829 p. 256, et 1831. p. 673. Rami adulti atro-fusci teretes. Ramuli angulati densè albo-tomentosi. Folia ovato-cordata utrinquè tomentosa, 3 poll. longa (excl. petiol. semipollic.), 2 poll. lata. Capitula densè corymbosa. Involucri squamæ extus in parte liberâ lanato-villoæ. Cor. glabræ. Achæ-nium sericeo-villosum. (v. s.)

§. 2. *Pappi serie externâ interiorem subæquante.* — Madagascarienses.

2. *D. capitatus* (Bojer ! in litt. 1835), foliis subsessilibus lineari-lanceolatis basi attenuatis suprâ glabriusculis subtus adpressissimè cano-tomentosis, corymbo circ. 4-cephalo congesto, capitulis subsessilibus 20-25-floris, invol. squamis

acuminatis vix tomentosis.  $\S$  ad fossarum margines circa urbem Tananarivou insulæ Madagascar legit cl. Bojer. Frutex 2-3-pedalis. Rami virgati striati. Folia  $1\frac{1}{2}$  poll. longa 3 lin. lata. Recept. eleganter areolatum, areolis inæqualiter dentatis medio punctigeris. Pappus rufus. (v. s. comm. à cl. Bojer).

3. D. *TANANARIVOU* (Bojer! in litt. 1835), foliis breviter petiolatis ellipticis et super. oblongis utrinquè tomentosis trinerviis, corymbo 5-7-cephalo conferto, capitulis pedicellatis paucifloris, invol. squamis acuminatis junioribus præsertim tomentosis.  $\S$  in insulâ Madagascar legit cl. Bojer. Frutex priori similis, sed paulò elatior. Rami angulati. Folia 9 lin. longa, 4 lin. lata. (v. s. comm. à cl. Bojer).

#### XXI. STROPHOPAPPUS. — Stilpnopappi sp. Mart.

Capitulum multiflorum. Invol. imbricatum, squamis ovali-oblongis obtusis extus apice subtomentosis. Recept. vix alveolatum. Pappus duplex, seriei utriusque ligulæ planæ acuminatæ nitentes elongatæ æquales. Achænium villosissimum. — Frutex Brasiliensis, floribus non satis notis. Rami teretiusculi juniores tomentosi. Folia sessilia oblonga obtusa coriacea subintegerrima suprà glabra subtus tomentosa. Capitula secus ramos ad axillas et apicem sessilia gemina folio breviora. †

1. S. *BI-COLOR*.  $\S$  in adscensu montis Serra do Caraca prov. Minar. Gener. Brasiliæ legit cl. Martius. Stilpnopappus bicolor Mart. ! herb. Folia 3 poll. longa, 8-9 lin. lata. Flores ign. Recept. vix alveolare ex statu imperfecto judicatum. (v. s. in h. Acad. reg. Monac.)

#### XXII. BLANCHETIA.

Capitulum pluri-(8-10) florum homogamum. Invol. ovati squamæ imbricatæ ovatæ mucrone acuminatæ. Recept. angustum epaleaceum. Cor. profundè et subæqualiter 5-fidæ. Anthæræ exsertæ basi obtusæ apice acutè appendiculatæ. Styli rami exserti hispidi acuminati. Achænium glaberrimum obovato-oblongum subpentagonum striatulum. Pappi squamellæ circ. biseriales subæquales caducissimæ lineares rigidulæ subciliatæ. — Herba suffruticulosa Brasiliensis erecta alternè ramosissima. Caulis rami et petioli tomento stellato brevi subrufescente et setis rigidis nigris patulis vestiti. Folia caulis alterna, ramorum steriliùm sæpius opposita petiolata ovali-oblonga basi longè cuneata subintegra aut hinc indè denticulata suprà pube stellatâ parcâ notata subtus eodem tomento densè canescenti-rufa. Capitula in cymam umbellatam foliosam disposita subsessilia conferta ovata subaraneosa. Cor. ex sicco purpurascens. Pappus albus. — Genus habitu Stilpnopappo affine sed pappo diversum, dicavi cl. Blanchet, botanico Helvetico, qui, civitati Bahiæ dicatus, Brasiliæ res naturales strenuè collegit et plantam ipsam generis typum cum pluribus aliis detexit.

1. B. *METASTROPHICA*.  $\S$  ? in Brasiliâ ad Caxocira ab urbe Bahiâ leucis 15 dist. legit cl. Blanchet. (v. s.)

#### XXIII. STILPNOPAPPUS Mart. herb.

Capitula multiflora homogama. Invol. campanulatum squamæ circ. biseriales æquales acuminatæ. Recept. alveolatum fimbriiferum. Achænium villosum. Pappus duplex uterquè paleaceus planus nitidus, ext. paleis brevibus, int. paleis duplò longioribus vix exterioris numerum superantibus. — Herbæ annuæ Brasilienses. Ex pappo paulò accedit hinc ad Dialestam illinc ad Lychnophoram.

1. *S. TRICHOSEPTOIDEUS* (Mart. ! herb.), caule herbaceo ramoso erecto puberulo, foliis oblongis basi attenuatis acutis subintegerrimis penninerviis planis supra glabris subtus tomentosis, pedunculis secus ramos distantibus oppositifoliis 1-cephalis natis, capitulis 15-20-floris, invol. floribus brevioribus biserialis squamis lanceolatis acuminatis foliaceis, ext. supra scabris subtus subtoementosis. — in Serra do Rio Contas prov. Bahiensis Brazilie legit cl. Martins. Achenium villosum turbinatum 10-costatum. Pappus duplex persistens lencus albidus, ext. paleis 10 ovatis cuspidatis, int. triplo fere longior setis filiformibus scabris 15-20. Recept. fimbriiferum. Cor. violaceæ. Capitulum circ. magnitud. Vern. anthelminthicæ. (v. s.)
2. *S. PRATENSIS* (Mart. ! herb.), caule herbaceo erecto tereti villosa, foliis ovali-oblongis acutis basi attenuatis integerrimis supra glabrescentibus subtus cano-tomentosis, infer. paulo latioribus, cymæ ramis scorpioidis oligocephalis capitulis 30-floris lateralibus solitariis sessilibus folio suo multo brevioribus, invol. squamis 2-3-seriatis acuminatis parce villosiusculis, recept. alveolato fimbriifero, achenio brevi crasso sericeo-villosissimo, pappi ser. utraq. paleis 10 constante, ext. ovatis brevioribus, int. lineari-elongatis acuminatis. ① in campis apricis prov. Piaulsiensis et Fernambucanæ frequentissimam legit cl. Martins. Cor. purpureo-violaceæ glabræ. Herba pedalis. Radix fibrosa. Folia 1  $\frac{1}{2}$  poll. longa, 5-6 lin. lata. (v. s. in h. Acad. reg. Monac.)
- ß. crotonifolia* (Mart. ! herb.), caule pubescente elatiore, foliis linearibus aut lanceolato-linearibus. ① in cultis apricis prov. Bahiensis Brazilie. (v. s. in h. Acad. reg. Monac.)
3. *S. TOMENTOSUS* (Mart. ! herb.), caule herbaceo erecto tereti subsimplici tomentoso basi folioso apice subaphyllo, foliis sessilibus linearibus integerrimis utraq. vix acutis 1-nerviis supra araneosis subtus tomentosis, capitulis 40-50-floris, ad apices aut secus ramos sessilibus folio flor. paulo brevioribus invol. campanulati squamis ovali-oblongis acuminatis dorso villosis adpressis, recept. alveolato longe fimbriifero, achenio sericeo-villosissimo, pappi serie utraq. paleacea, ext. acuminata. ① in aridis prope Caitete prov. Bahiensis Brazilie legit cl. Martins. Folia 1  $\frac{1}{2}$ -2 poll. longa, 2-3 lin. lata. Cor. purpureæ. (v. s. in h. Acad. reg. Monac.)
4. *S. SCAPOSUS*, herbaceus collo frutescente, foliis radicalibus confertis linearibus integerrimis coriaceis supra glabris subtus cano-tomentosis 1-nerviis, scapo nudo tereti pubescente simplici aut bifido, capitulis remotis lateralibus sessilibus ebracteatis, invol. squamis dorso pubescentibus ovatis, achenio sericeo-villoso, pappi paleis ext. ovatis, inter. linearibus. ② in Brasiliâ circa Bahiam legit cl. Blanchet (scapo simplici n. 1874! scapo bifido n. 1887!). Folia pollicem longa 1-2 lin. lata. Scapus 8-9 poll. longus. (v. s.)

† *Species exclusæ.*

5. *BICOLOR* Mart. = *Strophopappus bicolor*.

XXIV. DIALESTA *H. B. et Kunth nov. gen. am. 4. p. 35. t. 320. Cass. tabl. xx. n. 40. Less. in Linnæâ 1829. p. 315. syn. p. 146.*

Capitulum 2-florum aequaliflorum. Involucrum imbricatum cylindraceum fol. siccis 1-nerviis interioribus longioribus. Recept. minimum nudum. Cor. regularis glabra 5-fida, lobis acuminatis parte integrâ brevioribus. Fil. lævia. Achenium subcylindraceum multicostatum apice pilosiusculum, nectario styliiformi. Pappus biserialis paleaceus serratus, paleis seriei ext. minimis multis inæqualibus, int. 1-4 tortis longis. — Arbor? ramis canescenti, seu fuscescenti-tomentosis. Folia alterna petiolata oblonga integerrima supra glabra subtus albedo-ferruginea. Capitula pedicellata solitaria in corymbum terminata approximata. Cor. albidæ.

1. *D. discolor* (H. B. et Kunth l. c.) 5 in calidis prope Honda Novo-Granatensium. *Eupatorium cuspidatum* Willd. herb. ex Less. (v. s. comm. à cl. Poeppig.)

XXV. MONOSIS DC. in ann. bot. Guill. 1833. v. 2. p. 515. et in *Wight contrib.* p. 1. — *Conyza* sp. Wall.

Capitula uniflora in glomerulum non concreta. Invol. oblongi squamæ imbricatæ obtusæ flore multò breviores. Cor. tubulosa 5-fida. Antheræ basi muticæ. Stigmata longè exserta Vernoniæ. Achænium glabrum teretiusculum. Pappus 2-3-serialis, setis rigidulis scabris æqualibus aut longitudine sed non latitudine discrepantibus. — Suffrutices. Folia alterna breviter petiolata obovato-cuneata acutiuscula vix subdentata suprâ glabra subtus cum petiolis et ramis tomentosa. Panicula nuda. Rami apice capitula plurima subumbellata subsessilia conferta gerentes. — Affinis videtur Albertiniæ, sed capitula distincta, Shawiæ, sed pappus crassus basi non pubescens et ovarium sæpius glabrum. — *Mexicis* græcè solitudo, ex flore in invol. unico.

### Sect. I. EUMONOSIS.

Pappi series ext. ab interiore non discrepans. — Frutex Indicus.

1. *M. WIGHTIANA* (DC. in *Wight contrib.* p. 1), caule fruticoso?, ramis teretibus velutino-tomentosis, foliis petiolatis obovatis subacutis basi cuneatis et obtusis subintegerrimis penninerviis suprâ glabris et in nervis subvelutinis subtus velutino-hirsutis, paniculâ ramosissimâ capitulis ad apices ramulorum subcorymbosis subsessilibus, invol. squamis obtusis dorso tomentosis. 5 in Indiæ orient. montibus Neelgerry leg. cl. Wight et Noton. *Conyza Wightiana* Wall. ! cat. n. 3æ28 (non 3092) comp. 138. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)

### Sect. II. EREMOSIS.

Pappus pluriserialis setaceus, ser. ext. brevi sed ab int. latitudine non discrepante. — Frutices Mexicanæ.

2. *M. SAXICIFOLIA*, fruticosa glabriuscula, ramis teretibus, foliis petiolulatis oblongo-lanceolatis utrinquè acuminatis subintegerrimis penninerviis sparse glandulosis subtus glabris ad axillas venarum pilosis aut sparse puberulis, corymbo composito terminali, capitulis pedicellatis, squamis infimis minimis, cæteris longis lanceolatis submucronatis subciliatis. 3 in Mexici montibus ad Guchinapa et Cuernavaca legit cl. Berlandier oct. florent. Seriphium uniflorum et *Eupatorium uniflorum* Moc. fl. mex. mss. Invol. squamæ corollæque subpurpurascens. Pappi setæ ext. paucae vix inter. breviores albæ. Folia 2 poll. longa, 6 lin. lata. (v. s.)
3. *M. TARCHONANTHIFOLIA*, caule fruticoso tereti cano et adpressè tomentoso, foliis petiolatis oblongo-lanceolatis utrinquè attenuatis serraturas paucas callosas gerentibus suprâ glabris glandulosis subtus cano-tomentosis, ramis flor. ex axillis supr. et apice caulis nudis apice composito-corymbiferis, pedicellis ultimis confertis ternis brevibus, invol. oblongi 1-flori squamis glabris subcartiosis imbricatis ext. brevibus subtomentosis int. longioribus subacutis glabris, achænio striato glabriusculo, pappi setis ext. paucis brevibus et medicr. ab int. latitudine non discrep. 3 in Mexico legit cl. Karvinski. Pappus albus. Fol. 4 poll. longa, ferè pollicem lata, petiolo 2-3 lin. Invol. sordidè fuscocensilis. Cor. purpurascens. (v. s. in b. Acad. reg. Monac.)
4. *M. ? TOMENTOSA*, foliis ovatis integerrimis brevè petiolatis transversè venosis subtus ramisque tomentosis, capitulis in corymbum dispositis, invol. squamis ovatis acutis apice villosis. 3 in Mexico ad montem Guinzeo prope Vallisoleum. Turpinia? tomentosa Lallav. et Lex descr. 1 p. 24. Fulcaldea? tomentosa

Less. in linnæa 1830. p. 248. Cor. purpureæ. Pappus pilosus et stigmata revoluta ex auct. undè potius huc quàm ad Fulcaldeam referenda videtur.

XXVI. SHAWIA Forst. nov. gen. p. 95. t. 48. Cass. dict. 34. p. 40. n. vi. Less. syn. p. 156.

Capitulum 1-florum. Invol. squamæ 5-6 regulariter imbricatæ adpressæ coriaceæ extûs glandulosæ, ext. breves ovaies, int. oblongo-lanceolatæ. Recept. punctiforme. Cor. infundibuliformis 5-fida. Styli rami graciles exserti divergentes. Achænium subcylindraceum villosa-subtomentosum. Pappus fructu longior, paleolis filiformibus inæqualibus numerosis basi subnudis apice barbellulatis. — Frutices. Folia alterna. Capitula in paniculam corymbumve disposita. — Genus à Cassinio olim ad Inuleas demùm ad Vernonias relatum. È Vernoniis excludit Lessing et cum dubio ad Eupatorieas refert, sed ex capitulis 1-floris potius ad Vernoniaceas fortè referendum.

1. S. PANICULATA (Forst. et Cass. ll. cc.), foliis ovali-oblongis integerrimis petiolatis subtûs ramis pedunculisque tomentosis, capitulis in paniculam dispositis. § 5 in Novâ Zeelandiâ. Rich. et Lesson fl. nov. Zeel. p. 243.

### Subdiv. 3. ALBERTINIÆ.

Pappo uni- bi- aut pluriseriali paleaceo aut setiformi, capitalis sæpius paucifloris in glomerulum concretis.

### XXVII. HAPLOSTEPHIUM Mart. herb.

Capitula 1-flora in glomerulum globosum aggregata. Bractem paucæ foliaceæ subglomerulo. Invol. partialia oblonga, squamis imbricatis ferè linearibus acutis coriaceo-scariosis glabris. Achænium obovatum glaberrimum nervis 10 costulatum, umbone prominulo. Pappus simplex, paleis caducissimis linearibus spiraliter tortis. — Frutex Brasiliensis Lychnophoræ ferè facie. Rami teretes subverticillati adpressè sericis demùm glabriusculi. Folia conferta sessilia patula subrecurva basi subdilatata ferè adnata lanceolato-linearia acuminata integerrima suprà glabra subtûs sericeo-cana ferè ericæformia. — Genus affine Piptocomæ ex pappo caducissimo, Lychnocephalo ex pappi paleis tortis, sed præsertim Lychnophoræ ex habitu.

1. H. PASSERINA (Mart. herb.). § in summo monte Itambe de Villa prov. Minas-rum Generalium Brasiliæ legit cl. Martius. Fol. 4 lin. longa, vix lineam lata. (v. s. in h. Acad. reg. Monac.)

### XXVIII. LYCHNOPHORA Mart. in act. soc. sciab. v. 3. p. 2. p. 149 et diss. in-4°.

Capitula 1-4-flora in glomerulum subglobosum densè aggregata. Folia sæpius glomerulum bracteantia. Invol. oblongi squamæ 2-3-seriales lineari-oblongæ imbricatæ coriaceo-subscariosæ. Recept. nudus aut alveolare. Achænia glaberrima teretiuscula multicostata. Pappus biserialis, uterque latè paleaceus, serie exteriori multò breviori, interiori paleis tortis. — Fruticuli Brasilienses. Folia sessilia conferta uninnervia coriacea margine revoluta subtûs sæpè cana. Glomeruli ramos terminantes. — Genus habitu distinctissimum ex cl. Martius et Saint-Hilaire.

## Sect. I. EULYCHNOPHORA.

Capitula 1-flora. — *Lychnophora* genus Less. in *linnæa* 1829.

1. *L. ROSMARINIFOLIA* (Mart. diss. p. 10. t. 9), fruticosa, ramis teretibus crasso-tomentosis, foliis patentibus à basi dilatata cordata acuminato-subulatis acutis margine revolutis supra lineâ mediâ villosis subtus villo albo vestitis, capitulis adgregatis in glomerulos subglobosos interdum ipsos adgregatos sobecylindricis 1-floris, invol. squamis obtusis.  $\frac{3}{4}$  in summis montibus distr. Adamantini Brasiliæ ad Tejuco et Millio-Verde. *Lychnophora rosmarinifolia* Less. in *linnæa* 1829. p. 316. Ob flores in invol. solitarios ab affinibus generice separat cl. Lessing, meo sensu nimis numero concedens. (v. s. in h. Acad. reg. Monac.)

## Sect. II. Lychnophoroides.

Capitula 2-4-flora. — *Lychnophora* sect. *Vernonia* Less. in *linnæa* 1829.

2. *L. PHYLLICEFOLIA*, fruticosa demissa, ramis junioribus dense albo-tomentosis demum glabratis, foliis confertis erectis sessilibus lanceolato-linearibus mucrone cuspidatis margine latè revolutis supra glabris subtus albo-tomentosis, capitulis 2-floris in glomerulum foliis cinctum aggregatis, invol. squamis glaberrimis oblongis submembranaceis mucronato-acutis.  $\frac{3}{4}$  in altis ad Lincora prov. Bahiensis Brasiliæ legit cl. Martius. Suffrutex humilis. Pappus ext. brevissimus ferè coroniformis sed paleæ distinctæ erosæ obtusiusculæ, int. paleæ elongatis apice serrulatis. (v. s. in h. Acad. reg. Monac.)
3. *L. STAAVIOIDES* (Mart. ! diss. p. 9. t. 8), fruticosa, ramis teretibus junioribus crasso-tomentosis, foliis patentibus à basi subcordatâ oblongis obtusis submembranulatis margine latè revolutis supra glabris subtus tomentosis, capitulis cylindricis 3-floris in glomerulum subglobosum foliis interpositis maxime aggregatis, invol. squamis laxè imbricatis obtusis 3-nerviis.  $\frac{3}{4}$  in summis montibus distr. Adamantini Brasiliæ. *Vernonia staavioides* Less. in *linnæa* 1829. p. 249. 1831. p. 630. *Dialesta staavioides* Mart. ! herb. Invol. squame apice summo tomentosæ. Pappi paleæ ext. sub lente apice serratæ. Cor. roseæ. (v. s. in h. Acad. reg. Monac.)
4. *L. VILLOSISSIMA* (Mart. ! diss. p. 8. t. 7), fruticosa, ramis teretibus crasso-tesselato-tomentosis, foliis patentibus linearibus obtusis margine latè revolutis supra in basi longè lanuginosis subtus ubiqûe tomentosis, capitulis 6-8 in glomerulum globosum lanatum adgregatis 4-floris cylindricis, invol. squamis brevè imbricatis subacutis.  $\frac{3}{4}$  in Brasiliæ distr. Adamantino in campis apricis prov. Tejuco. *Vernonia villosissima* Less. in *linnæa* 1829. p. 249. 1831. p. 630. Cor. pallide lilacinæ.
5. *L. SALICIFOLIA* (Mart. ! diss. p. 12. t. 8), fruticosa, ramis teretibus junioribus pulverato-tomentosis, foliis erecto-patentibus linearibus lanceolatis mucronulatis margine vix revolutis subtus tomentosis, capitulis ovatis 6-floris in glomerulos globosos adgregatis 2-floris cylindricis cinctis, invol. squamis arcuè adpressis lanceolatis.  $\frac{3}{4}$  in Brasiliæ distr. Adamant. Brasiliæ. *Vernonia salicifolia* Less. in *linnæa* 1829. p. 249. 1831. p. 630. Cor. pallide roseæ. Invol. squame apice summo tomentosæ. Pappi paleæ sub lente serratæ. (v. s. in h. Acad. reg. Monac.)
6. *L. PIRASTEA* (Mart. ! diss. p. 7. t. 6), subherum, ramis teretibus junioribus pulverato-tomentosis, foliis erecto-patentibus linearibus lanceolatis mucronulatis margine vix revolutis subtus tomentosis, capitulis ovatis 6-floris in glomerulos globosos adgregatis 2-floris cylindricis cinctis, invol. squamis arcuè adpressis lanceolatis.  $\frac{3}{4}$  in Brasiliæ montibus prov. Minas-Geraes distr. Adamant. Brasiliæ. *Vernonia pirastea* Less. in *linnæa* 1829. p. 249. 1831. p. 630. Cor. pallide roseæ. Invol. squame apice summo tomentosæ. Pappi paleæ sub lente serratæ. (v. s. in h. Acad. reg. Monac.)
7. *L. HIRSAEFORMIS* (Mart. ! diss. p. 11. t. 5), fruticosa, ramis teretibus junioribus pulverato-tomentosis, foliis erecto-patentibus linearibus lanceolatis mucronulatis margine vix revolutis subtus tomentosis, capitulis ovatis 6-floris in glomerulos globosos adgregatis 2-floris cylindricis cinctis, invol. squamis arcuè adpressis lanceolatis.  $\frac{3}{4}$  in Brasiliæ montibus prov. Minas-Geraes distr. Adamant. Brasiliæ. *Vernonia hirsaeformis* Less. in *linnæa* 1829. p. 249. 1831. p. 630. Cor. pallide roseæ. Invol. squame apice summo tomentosæ. Pappi paleæ sub lente serratæ. (v. s. in h. Acad. reg. Monac.)

foliis floralibus instructis, invol. squamis laxè imbricatis linearibus obtusis. § in summo monte Itambe do Villa Brasiliæ. *Vernonia hakeaefolia* Less. l. c. p. 249. Cor. rosæ. (v. s. in h. Acad. reg. Monac.)

8. *L. proteziformis* (Mart.! dias. p. 6. t. 5), fruticosa, ramis divaricatis basi nudis apice teselato-tomentosis, foliis erecto-patentibus linearibus cum mucrone brevi margine revolutis suprâ glabris subtùs albo-tomentosis, capitulis terminalibus sessilibus confertis ovato-cylindricis 3-4-floris, invol. squamis lineari-lanceolatis ciliatis. § in Brasiliæ monte Adamantino Serra de Santo-Antonio prov. Minas-Geraes. *Vernonia proteziformis* Less. in *linnæa* 1829. p. 249. 831. p. 629. Cor. lilacinæ.
9. *L. zecordas* (Mart.! dias. p. 6. t. 5), subarborea, ramis teretibus teselato-tomentosis, foliis erecto-patentibus linearibus submucronulatis margine revolutis suprâ glabris subtùs tomentosis, capitulis corymbois in glomerulum globosum foliis cinctum adgregatis, invol. squamis lineari-lanceolatis obtusis corollisque glabris. § in monte Adamantino Serra de Santo-Antonio prov. Minas-Geraes Brasil. Fol. 15-18 lin. longa, lineam lata. Cor. pallidè lilacinæ. Pappi paleæ subulatæ vix apice serrulatæ, exterior. interior. duplò brevior. (v. s. in h. Acad. reg. Monac.)
10. *L. brunnioides* (Mart.! dias. t. 4. p. 4), fruticosa, ramis teretibus glabris nodulosis, foliis erecto-patentibus lineari-lanceolatis acutis basi attenuatis margine revolutis suprâ glabris subtùs tomentosis, capitulis corymbois in glomerulum terminalem congestis 3-4-floris cylindricis, invol. squamis oblongo-linearibus. § in Brasiliæ montanis siccis ad Itambe da Villa. *Vernonia brunnioides* Less. in *linnæa* 1829. p. 248. 1831. p. 629. Rami tenues cicatrati sæpè lichenibus obtusi. Folia 5-6 lin. longa. Cor. pallidè rosæ in genere parvæ. Pappus ext. paleis brevibus erosis acutis, int. paleis elongatis purpurascens. (v. s. in h. Acad. reg. Monac.)
- §. *alpicola*, ramis junioribus pubescenti-subcaescentibus, foliis paulò minus confertis. — in Serro Frio ad Tejuco alt. 5000 ped. prov. Minas General. Brasiliæ legit cl. Martius. *Vernonia alpicola* var. Mart. f. herb. Pappus ext. paulò adhuc brevior quam var. α, int. paleis linearibus purpurascens. (v. s. in h. Acad. reg. Monac.)

XXXIX. *ALBERTINIA* Spreng. *neu entd.* 2 p. 133. *syst.* 3. p. 355 et 434 (*excl. car.*), Mart. *herb.*

Capitula 1-3-flora densissimè in glomerulos subglobosos collecta et sæpiùs involucrio universali brevi polyphylo cincta. Invol. partialis squamæ erectæ adpressæ sæpè pressione ad alveolorum rudimenta reductæ. Pappus bi- aut pluriserialis, setis omnibus filiformibus subscabris sæpiùs coloratis. — Frutices Brasilienses. Folia alterna petiolata elliptica utrinquè attenuata integerrima glabriuscula aut subtùs albida.

#### Sect. I. ANISOTRICHIA.

Pappi series externa interiore duplò triplove brevior.

#### §. I. Capitulis 1-floris.

1. *A. BRASILIENSIS* (Spreng. *syst.* 3. p. 355 et 434), caule fruticoso ramoso tereti apice brevissimè hirsuto, foliis petiolatis ellipticis utrinquè acuminatis integerrimis membranaceis suprâ glabriusculis subtùs petioloque subvelutinis, capitulis in racemum subcompositum ovatum dispositis pedunculatis ebracteatis, invol. campanulati floribus multò brevioris squamis velutinis concretis, recept. profundè alveolato. § in Brasiliæ prov. Rio de Janeiro ad Serra d'Estrella et Cercovado (Mart. f. Vanth. l. n. 341) in collibus circa Bahiam (Salzm. f. Blanch. l.). *Albertinia Brasiliensis* Less. in *linnæa* 1829. p. 341. *Vern. Brasiliensis* Less. in

- 1882 1831. p. 682. Pappus rufus biserialis, exterior dimidio brevior. Flores carnei seu pallide violacei suaveolentes. (v. s.)
2. *A. RUFISERTA*, caule fruticoso ramoso, ramis teretibus velutinis, foliis subsessilibus ellipticis obtusis integerrimis supra araneoso-velutinis obscure virescentibus subtus adpressè velutino-tomentosis canescentibus, summis pedunculisque suboppositis, capitulis 1-floris densè in glomerulum confertis, invol. squamis lanceolatis, intimis valdè acuminatis. § in Brasiliæ siccis apricis montosis ad Serro Frio prov. Minarum General. legit cl. Martins. Albertinia obtusata Mart. ! herb. Habitus Albertiniæ et Eremanthi, à priori capitulis 1-floris et inv. squamis liberis, à posteriore pappi 2-3-seriali, serie extimà dimidio brevior differt. Achænium sericeo-villosum turbinato-cylindraceum. Pappus rufus. (v. s. in h. Acad. reg. Monac.)
3. *A. GALLIDIERTA*, caule fruticoso ramoso, ramis teretibus velutinis, foliis ellipticis basi cuneatis obtusis integerrimis supra glabriscolis obscure virentibus subtus adpressè velutino-tomentosis canis, pedunculis folia sua superantibus, capitulis 1-floris densè in glomerulum confertis, invol. squamis liberis subhirsutis, intimis mucronatis achænia non superantibus. § in campis editis ad Serro Frio prov. Minar. Gener. Brasiliæ legit cl. Martins. Albertinia obtusata var. ? Mart. ! herb. A præcedenti tantum differt glomerulis longius pedunculatis, inv. squamis minus acuminatis et pappo paulò magis paleaceo stramineo nec rufo. An var. ? (v. s. in h. Acad. reg. Monac.)

### §. 2. Capitulis 3-floris.

4. *A. CAOROXORDES*, ramis teretibus hirsuto-tomentosis super. subcorymbosis, foliis breviter petiolatis ellipticis obtusis basi subattenuatis integerrimis supra junioribus subvelutinis adultis glabris subtus petiolisque densè cano-tomentosis, capitulis 3-floris oblongis ad apices ramulorum conferto-umbellatis brevissimè pedicellatis, invol. squamis imbricatis lanceolatis acutis dorso tomentosis, achænio pilosiusculo. § in montium sepihus prov. Minar. Gener. Brasiliæ legit cl. Martins. Albertinia bardanoides Mart. herb. Folia 2 poll. longa pollicem lata. Pappus duplex purpurascens, ser. ext. dimidio brevior magis paleacea. (v. s. in h. Acad. reg. Monac.)
5. *A. VERBASCIFOLIA* (Mart. ! herb.), ramis teretibus petiolis folisque subtus densissimè hirsuto-tomentosis, foliis supra junioribus tomentosis adultis subvelutinis basi cordatis latè ovatis obtusis integerrimis lanugine crassis penninerviis, nervis later. utrinquè 15-16 subparallelis reticulato-ramosis, capitulis 3-floris oblongis ad apices ramulorum glomerulato-umbellatis ebracteatis, invol. squamis lanceolatis acutis dorso velutinis, achænio pilosiusculo. — in ferruginosa Serra de Ant. Pereira prov. Minar. General. legit cl. Martins. Pappus subrufescens biserialis, ext. dimidio brevior. Toto cœlo diversa à *V. verbascifolia* Less. Fol. petiolus pollicaris, limbus 6 poll. longus, 4-5 poll. latus. (v. s. in h. Acad. reg. Monac.)
6. *A. ELMAEUS* (Mart. ! herb.), caule fruticoso ramoso tereti subcanescente, ramulis ultimis angulatis, foliis petiolatis oblongis obtusis basi acuminatis integerrimis supra cinascenti-subtùs rufescenti-adpressè velutinis, capitulis 5-6 ad apices ramulorum umbellato-sessilibus 3-floris, invol. squamis subincanis lanceolatis acutis, achænio pilosiusculo. § in altis lapideis Serro Frio ad Tejuco prov. Minar. Gener. Brasiliæ legit cl. Martins. Pappi pallide rufescentis series 2-3, ext. paucissimè breviori. Folia utrinquè nervo medio saltem basè transverse ruguloso ! seu tuberculoso ; an casu ? (v. s. in h. Acad. reg. Monac.)
7. *A. STYLIS*, fruticosa, ramis teretibus densè tomentosis, foliis petiolatis coriaceis ovatis-oblongis basi attenuatis vix acutis integerrimis supra glabris subtus petiolisque cano-tomentosis, capitulis oblongis 2-3-floris in glomerulum globosum foliis paucis sessilibus sæpè unctum densè adgregatis. § in altis prov. Minarum Brasiliæ legit cl. Martins. *Lychnocephalus bicolor* Mart. ! herb. Habitus *Lychnophoræ*, capitulum *Echinopsidii*, bractæ ferè *Hololepidis*, invol. squamæ apice obtusæ tomentosæ. Achænia glaberrima subangulata. Pappus



setosus pluciserialis, ser. ext. breviori. Folia 6-8 poll. longa, 1-2 poll. lata, nervo subtus valdè prominente. (v. s. in h. Acad. reg. Monac.)

### Sect. II. ISOTRICHIA.

Pappi series externa ab interiore nec longitudine nec latitudine discrepans.

#### §. 1. Capitulis 2-3-floris.

8. *A. CAPITATA*, fruticosa, ramis adultis teretibus junioribus pedunculisque angulatis canis, foliis petiolatis ellipticis utrinquè acuminatis subintegerrimis suprà glabris subtus petiolis ramulisque densè lepidoto-tomentosis albis, capitulis cylindraceis ad apices ramorum sessilibus densè corymbosis 3-floris, invol. squamis aretè imbricatis ex ovato-oblongis, achænio teretiusculo multifido-costato glabro, pappi paleis seriebus 2 aequalibus. § in Brasiliâ tropicâ, ad Portum d'Estrellam (Less.!), in monte Cercovado (Lund! Lhotsky!) *Conyza capitata* Spreng. syst. 3. p. 507. Less. in linnæa 1829. p. 270. *Vernonia incanescens* Mart. herb. Pappus stramineo-rufescens caducissimus. Achæni callus basilaris vix perspicuus. (v. s.)

9. *A. ERYTHROPAFFA*, caule fruticoso ramoso, ramis pedunculisque striato-angulatis adpressè incanis, foliis breviter petiolatis lanceolatis basi acutè apice subobtusè acuminatis integerrimis suprà glabris subtus adpressè canis, capitulis 2-3-floris densè in globulolum pedunculatum congestis, invol. squamis ext. subtomentosis truncatis, intimis paulò longioribus acutis. § in Brasiliâ prov. Minarum Generalium legit cl. Vauthier (coll. exs. n. 334). Pappus purpureo, setis biserialibus aequalibus. Achæniolum glabriusculum. (v. s. comm. à cl. Vauth.)

#### §. 2. Capitulis unifloris. — *Eremanthus* Less. in linnæa 1829. p. 317. 1831. p. 682. syn. 147. non Cass.

10. *A. GLOMERULATA*, ramis pedunculisque teretibus, foliis obtusis, floralibus sub glomerulo globoso eo longioribus reflexis, achænio pilosiusculo. § in Brasiliâ tropicâ (Less.) prov. Minar. General. (Gaudich.!) *Eremanthus glomeratus* Less. II. cc. Folia et rami supremi suboppositi. Pappus rufus. (v. s. in h. Mus. reg. Paris.)

11. *A. INCANA* (Less. in linnæa 1829. p. 342), ramis pedunculisque striatis compressis foliis lanceolatis utrinquè acuminatis, floralibus capitulo brevioribus, achænio glabro. § in Brasiliâ tropicâ (Less.) prov. Minarum Generalium (Vauth. n. 355). *Vernonia incana* Less. in linnæa 1831. p. 682. Invol. squamis ext. breves obtusæ subtomentosæ, intimæ abruptè duplò longiores glabriusculæ. Pappus albidus. (v. s.)

12. *A. POLYCEPHALA*, caule fruticoso striato corymboso ramoso, ramis angulatis adpressè subincanis, ultimis striato-sulcatis subcompressis, foliis lanceolatis suprà glabris subtus adpressè canescentibus integerrimis basi acuminatis subsessilibus apice vix acutis, capitulis 1-cephalis umbellatis glomerulis subebracteatis pedunculatis, invol. squamis ext. brevibus obtusis, intimis elongatis glabriusculis, achænio glabro. § in planitie altâ ad Pieda de Villem prov. Minar. Gener. Brasiliæ legit cl. Martius. Albertinia multiflora Mart. herb. Pappi series æquales subpurpureascentes. (v. s. in h. Acad. reg. Monac.)

13. *A. SALIGNA* (Mart. herb.) caule fruticoso tereti ramoso, ramulis ultimis angulatis subcompressis adpressè subincanis, foliis linearilanceolatis utrinquè attenuatis subobtusis integerrimis suprà glabris subtus adpressè cano-tomentosis, capitulis 1-floris in glomerulos ebracteatos pedunculatos 8-9 umbellato-congestis, invol. squamis ext. brevibus subincanis, int. elongatis glabriusculis. § in editis siccis ropestribus montium Serro Frio prov. Minar. General. Brasiliæ legit cl. Martius. Folia 2 poll. longa, 5 lin. lata. Pappus pallidus setis ex statu juniore aequalibus. (v. s. in h. Acad. reg. Monac.)

XXX. PYCNOCEPHALUM. — *Pycnocephalum* sect. *Vernonia* Less. in *linnæa* 1831. p. 630.

Capitula plurima 2-7-flora in glomerulum terminalem subglobosum collecta. Squamæ ovales multistriatæ paucæ sub glomerulo involucri locum gerentes. Invol. partialis squamæ oblongo-lanceolatæ 3-nerviis subciliatæ. Cor. glabræ. Achænium sericeo-villosum. Pappus pluriserialis setaceus scaber, setis ext. paucis brevioribus sensim ad interiorum longitudinem accedentibus. — Herbæ perennes imâ basi foliatæ cæterum aphyllæ simplicicaules.

1. *P. SPATHULÆFOLIUM*, collo crasso subtomentoso, caulibus pluribus herbaceis basi foliatis cæterum aphyllis compressis densè sericeo-pubescentibus, foliis subradicalibus lineari-oblongis (spathulatiæ ex Mart.) basi angustatis erectis integerrimis junioribus pubescentibus adultis glabris subnerviis, capitulis oblongis 5-7-floris in glomerulum globosum ebracteatum terminalem congestis, invol. squamis lanceolatis acuminatis basi 3-nerviis subciliatis, intimis glabris.  $\frac{1}{2}$  in aridis deserti versus Chapada de Parano prov. Minar. General. Brasiliæ legit cl. Martins. *Vernonia spathulæfolia* Mart. herb. Valdè affinis *P. scapigero*. Achænium sericeo-villosum. Pappus pluriserialis, setis scabris, ext. dimidio circ. brevioribus. Cor. purpureæ. (v. s. in h. Acad. reg. Monac.)
2. *P. SCAPIGERUM*, caule herbaceo simplicissimo aphylo tereti lanato-tomentoso, foliis radicalibus coriaceis brevè petiolatis spathulato-oblongis subintegerrimis glaberrimis junioribus subtomentosis, capitulis subcylindraceis 2-3-floris in glomerulum terminalem adgregatis, invol. squamis carinatis mucronato-acuminatis, achænio densè piloso.  $\frac{1}{2}$  in Brasiliâ tropicâ. *Vernonia scapigera* Less. in *linnæa* 1829. p. 250. Cor. purpureæ glabræ.
3. *P. PLANTAGINIFOLIUM*, caule herbaceo simplicissimo tereti supernè aphylo tomentoso, foliis subradicalibus membranaceis multiplinnerviis lineari-lanceolatis integerrimis glabriusculis, junioribus tomentosulis, capitulis cylindraceis multifloris in glomerulum adgregatis, invol. squamis planiusculis 3-nerviis glabriusculis, achænio densè piloso.  $\frac{1}{2}$  in Brasiliâ tropicâ. Cor. purpureæ glabræ. *Vernonia plantaginifolia* Less. in *linnæa* 1829. p. 251.

### XXXI. LYCHNOCEPHALUS Mart. herb.

Capitula plurima 8-flora ovali-oblonga densissimè conferta in glomerulum terminalem subrotundum involucri generali destitutum. Bracteæ foliaceæ sub capitulo quoque et idè in medio glomeruli hinc indè perspicuæ. Involucrum partiale oblongo-cylindraceum in lanâ immersum, squamis plurimis glabriusculis obtusis. Achænium glabrum striato-sulcatum. Pappus duplex paleaceus, paleis ext. brevibus acutis persistentibus, int. elongatis spiraliter tortis caducissimis. — Frutex Brasiliensis. Rami teretes densè tomentosi. Folia petiolata ovali-oblonga obtusa basi cuneata coriacea integerrima suprâ glabra subtus petiolique densè tomentosa.

1. *L. TOMENTOSUS* (Mart. herb.).  $\frac{3}{4}$  in campis sterilibus altis provincie Minas-Geraes Brasiliæ legit cl. Martins. Variat tomento albicante et flavicante. Invol. squamæ obtusæ glabriusculæ. Achænium glabrum striato-costatum. Pappus duplex paleaceus, ext. brevis acutus, int. elongatus spiraliter tortus caducissimus. (v. s. in h. Acad. reg. Monac.)

† *Species exclusa.*

*L. NEOLON* Mart. = *Albertinia bicolor*.

XXXII. CHRONOPAPPUS. — *Heterocomæ* sp. DC.

Capitula pauca 8-10-flora in glomerulum axillarem conferta, bracteis foliaceis involucrium verum non constituentibus circumdata. Invol. partialis ovalis squamæ imbricatæ dorso tomentosæ acutæ lanceolato-lineares, int. angustissimæ subbreviares. Recept. nudum. Corollæ lobi dorso hirsuti. Achænia breviter angulata glaberrima. Pappus duplex uterque persistens, ext. paleis brevibus oblongis acutis, int. setis elongatis serratis. — Ab *Heterocomâ* differt bracteolis recept. nullis, corollâ hirsutâ, pappo ext. paleaceo nec coroniformi. Proximè ad *Lychnocephalum* accedit.

1. *C. styriaca*, fruticosa, ramis teretibus apice subcompressis valdè lanuginosis, foliis petiolatis ovatis basi obtusissimis apice subobtusis integerrimis suprà adultis glabris eminentiâ duris prismaticis eximie muricatis suprà petiolisque camolanuginosis, glomerulis ad axillas superiores subsessilibus spicam compositam interruptam terminalem ferè constituentibus. ♀ in Brasiliâ. *Heterocomæ bifrons* DC. ann. mus. 16. p. 191. t. 8. recueil mém. 2. p. 34. t. 4. *Serratula bifrons* Pers. ench. 2. p. 391. (v. s.)

## XXXIII. PITHECOSERIS Mart. herb.

Capitula 3-4-flora in glomerulum ovatum densè congesta sessilia subspicata ebracteata. Invol. oblongi squamæ erectæ glabriusculæ acuminatæ, ext. carinatæ, int. planæ lanceolato-lineares. Recept. nudum. Cor. tubus extus hirsutus, lobi glabri. Achænia in iisdem capitulis difformia, alia (juniora aut sterilia?) sericeo-villosa, pappo duplici ext. brevissimo paleaceo, int. pluriseriali setaceo: alia (matura aut fertilia?) glaberrima oblonga compressa, pappo setiformi caducissimo! — Herba. Folia ferè lyrata seu sinuato-pinnatifida.

2. *P. racovitzoides* (Mart. ! herb.), glabriuscula, caule tereti, ramis apice fistulosis nudis, foliis sessilibus anrito-semiamplexicaulibus irreg. pinnati-lobatis lobis ovatis triangularibus dentatis subsinuatisve, terminali majore, glomerulo ovato glabro. ♀ in campis prov. Pernambuco legit cl. Martins. Herba grandis secus caulem et in foliorum paginâ infer. subtiliter pubescens. Folia membranacea pedem longa, 6 poll. lata. Capit. oblonga 3-5-flora. Invol. squamæ glabræ acuminatæ. Achænia juniora puberula adulta glaberrima. Pappus pluriserialis setosus, ser. ext. brevis, int. longa subrufescens, facillè deciduus in achæniis adultis. (v. s. in h. Acad. reg. Monac.)

## XXXIV. STACHYANTHUS.

Capitula 12-13-flora secus rami apicem sessilia spicata subconferta, rachis ad eorum exortum scrobiculatâ, bracteis subcapitulis nullis. Invol. squamæ oblongæ adpressæ obtusæ, int. submucronatæ. Achænia sericeo-villosa. Pappus pluriserialis setaceus, serie ext. brevi basi flavidâ. — Folia sinuato-angulata. Affinis *Echinopsidio*.

1. *S. Martii*, suffruticosa tota pube brevi adpressâ sericeo-canescens, caule tereti ramoso, foliis petiolatis ovato-rhombicis basi integris cuneatis à medio ad apicem dentato-angulatis, pedunculis longissimis nodis glomerulo ovato-globoso terminatis, capitulis secus axin spicatis 12-13-floris, invol. squamis oblongis submacronulatis. ♀ in siccis sylvis æstu aphyllis prov. Bahiensis et Piahiensis Brasilie legit cl. Martins. *Vernonia Stachyanthus Mart. !* herb. *Achæniom sericeo-villosum*. Pappus pluriserialis, setis ext. brevibus, basi flavidis. Folia ferè *Atriplicis roseæ*, sed limbo 3 poll. longo, 2 poll. lato, petiolo pollicari. (v. s. in h. Acad. reg. Monac.)

XXXV. CHRESTA *Fl. flumin. ic.*

Capitula 3-4-flora in glomerulos sphaericos terminales ebracteatos densissimè collecta. Invol. squamæ siccae acuminatæ paucæ. Achænium sericeo-villosissimum. Pappus multiserialis, serie ext. cæteris brevior. — Habitus distinctissimus hinc ad Elephantopum illinc ad Lychnocephalum accedens.

1. *C. SPHÆROCEPHALA*, tota canescens velutino-tomentosa, caule erecto tereti apice ramoso, foliis petiolatis ovatis penninerviis coriaceis dentato-subsinuatis, ramis nudis capitulo globoso compacto terminatis. 2*l* in saxosis calvis aridiusculis prov. Bahiensis legit cl. Martins, et in prov. Minas Geraes prope Tejuco cl. Vauthier (coll. exs. n. 294) Chresta cordata Arrab. fl. flum. 8. t. 150? In specim. Vauthieriano pedunculi apice subglomerulo fissi sen ramosi aut monstruositate aut compressione spectabiles, in Martiano indivisi. Invol. squamæ glaberrimæ. Cor. glabræ purpureæ. *Vernonia Echinope* Mart. ! herb. (v. a.)

2. *C. EXSUCCA*, caule erecto ramoso pubescenti-tomentoso, foliis sessilibus ellipticis acutiusculis integerrimis coriaceis supra glabris, subtus nervoso-reticulatis junioribus subvelutinis adultis glabris, ramis apice nudis glomerulum globosum gerentibus, invol. squamis lanceolatis cuspidatis. 2*l* in montosis prov. Minar. Gener. Brasiliæ legit cl. Martins. *Vernonia exsucca* Mart. ! herb. Folia 12-15 lin. longa, 6-7 lata. Achænium sericeo-villosissimum. Pappus flavescens, setis ciliatis pluriserialibus ext. dimidiò triplòve brevioribus. Species affinis *C. pycnocephalæ*. (v. s. in h. Acad. reg. Monac.)

4. *C. PYNOCEPHALA*, tota adpressè sericeo-pubescenti ferè lepidota, caule tereti apice ramoso, foliis sessilibus obovato-oblongis acutis subintegerrimis penninerviis, ramis nudis glomerula globoso compacto terminatis. 2*l* in campis deserti inter. Min. Nov. et S. Franciscum prov. Minar. General. Brasiliæ legit cl. Martins. *Vernonia pycnocephala* Mart. herb. Glomeruli dimidiò minores et folia diversissima à *C. sphærocephalâ* cui cæter. valdè affinis. Capitula 4-flora. Pedunculi folia quintuplò superantes. Invol. squamæ glabriusculæ submicroneatæ. (v. s. in h. Acad. reg. Monac.)

¶ *Halimodendron*, pedunculis folio triplò tantum circ. longioribus, invol. squamis extus puberulis. 2*l* in campis siccis deserti prov. Minar. Gen. prope Reacho de Togo et ad fl. S. Francisci prope Salgado legit cl. Martins. *Vern. Halimodendron* Mart. ! herb. (v. s. in h. Acad. reg. Monac.)

Div. II. ELEPHANTOPÆ *Cass. tabl. xx. sect. 4. n. 111. excl. Dialesta Less. in linnæd 1830. p. 135. — Elephantopidæ Less. syn. 149.*

Capitula discoidea pauciflora. Invol. compressum pluriseriale, foliolis seriei cætasque 4, alternis conduplicatis, alternis planis conduplicatis distichis. Recept. nudum aut alveolatum. Cor. palmata. Anthere cæcundatæ. Styli rami semisubulati. — Folia alterna.

XXXVI. ELEPHANTOPUS *Cass. dict. 14. p. 341. Less. in linnæd 1819. p. 323. syn. 149. — Elephantopi sp. Linn. Gært.*

Capitula pluriflora (3-5 sæpius 4-flora) æqualiflora in glomerulum foliis cinctum dense adgregata. Involucrum compressum biseriale, foliolis siccis oblongis alternè planis et conduplicatis interioribus sæpius 3-nerviis. Rhachis nuda. Cor. palmata, limbo 5-fido, laciniis acuminatis, sinu uno cæteris profundiore. Stam. filam. lævia. Styli rami semisubulati. Achænium compressiusculum multicostatum oblongum pilosum. Pappus 1-serialis, paleis plurimis rectis basi dilatatis cæterum

angustissimis acuminatis æqualibus serratis. — Herbæ perennes pilosæ erectæ. Folia alterna sessilia penninervia. Glomeruli ramos terminantes subcorymbosi. Cor. violaceæ rarius albæ.

### §. 1. *Corollâ glabrâ.*

1. *E. SCABER* (Linn. sp. 1313. excl. syn. Gron. et Brown.), caule dichotomè ramoso strigoso-villoso, foliis scabris radicalibus crenatis cuneatis basi longè attenuatis, caulinis lanceolatis, floralibus latè cordato-ovatis acuminatis canescentibus.  $\frac{1}{2}$  in Ind. orient. frequens in Timor, Javâ, Luçonîâ, Silhet, Nepaliâ, agro Burmannico, etc. Less. in linnæâ 1829. p. 325. Wall. ! herb. comp. n. 89. Wight. ! herb. — Dill. elth. 126. t. 106. Breyn. ic. 32. t. 34. (v. s.)
2. *E. CAROLINIANUS* (Willd. sp. 3. p. 2390 excl. Br. syn.), caule piloso apice corymboso-ramoso, foliis scabris, radicalibus crenatis cuneatis basi attenuatis, caulinis oblongis basi angustis floralibus ovato-oblongis scabridis.  $\frac{1}{2}$  in siccis substerilibus Carolinæ, agri Kentuckensis, Floridæ, et ex Willd. in Jamaicâ, ex Sieber in Martinicâ, ex variis in Braziiliâ, Mexico, etc.; sed fortè cum sequentibus sæpè confusus. Ell. sketch, 2. p. 480. Less. in linnæâ 1829. p. 325. E. scaber Swartz obs. 316? Michx. fl. bor. am. 2. p. 148. Glomeruli bracteis latitudine duplò et ultrâ longiores neo ut in priore latitudini ferè longitudine æquales. (v. s.)
3. *E. MONICAULIS* (Ell. sketch, 2. p. 481), caule subhirsuto subnudo parcè ramoso, foliis radicalibus ovali-sublanceolatis crenato-serratis suprâ scabriusculis subtus villosis, floralibus latè ovato-cordatis villosissimis.  $\frac{1}{2}$  in siccis substerilibus Carolinæ Pofr. suppl. 2. p. 543. E. Carolinianus var. simplex Nutt. gen. 2 p. 187. E. tomentosus Linn. sp. 1314? excl. syn. auct. Carib. Pursh fl. 2. p. 582? Less. in linnæâ 1829. p. 326? an E. Caroliniani var.? (v. s.)
4. *E. MOLLE* (H. B. et Kunth nov. gen. am. 4. p. 26), caule strigoso-piloso supernè dichotomo-ramoso, foliis ovatis inferioribus basi attenuatis crenato-serratis suprâ glabriusculis subtus mollissimè canescenti-villosis, caulinis oblongis acuminatis serratis, floralibus ternis latè ovatis acuminatis serratis.  $\frac{1}{2}$  in Cumanâ et Caracas ex H. B. et K., Havanâ, Portorico, Peruvîâ (Poepp.), Braziiliæ prov. S. Pauli (Lund). Videtur E. tomentosus auct. in Caribæis indicatus. E. Carolinensis albidiflorus Meyer eseq. 256? Sloan. hist. t. 156 f. 1. 2? Cor. alba ex Kunth et Meyer. (v. s.)
5. *E. MARTII* (Graham list. pl. rar. mart. 1830. p. 2), caule ramoso piloso folioso, foliis subsessilibus undulatis rugosis supernè hispidis subtus tomentosovillosis serrato-crenatis denticulatis, inferioribus spatulato-oblongis breviter attenuatis, floralibus latè ovato-cordatis acutis villosis.  $\frac{1}{2}$ ? in Braziiliâ ad Rio-Janeiro (Lund!), Sanctam Catharinam (Bacle!), Bahiam in incultis (Salzman!) El. cervinus Arrab. fl. flum. 8. t. 148. Vulgatissimus in pratis apreis et dumetis circa habitationes usque ad altitud. 3000 ped. (herb. Acad. Monac.) Ab indigenis *Yerva grossa* dictus. Infusio theæformis sudorifica. Cor. pallida. (v. s.)

### §. 2. *Corollæ lobis strigoso-pilosis.*

6. *E. HIRTIFLORUS*, caule hirtello sulcato subdichotomo, foliis oblongis sessilibus acutis villosissimis, inferioribus serratis, cæteris integris, floralibus oblongo-lanceolatis acuminatis involucrique foliolis villosis. — in sabniosis maritimis circa Bahiam legit cl. Salzman. Cor. ex eo pallidè cærulescens, lobis ad apicem strigis hispidis. (v. s.)

### + *Species non satis notæ.*

7. *E. VIRGATUS* (Desv. in Ham. fl. ind. occ. 52.) caule virgato hirsuto-piloso, foliis subsessilibus trinerviis lanceolato-linearibus remotè serratis utrinquè strigoso-pilosis, pedunculis longissimis paniculatim dispositis. — in Guianâ. Nullam novi E. speciem foliis trinerviis.

8. *E. FLEXUOSUS* (Raf. fl. lod. 73), caule ramoso flexuoso, foliis radicalibus cauliniisque sessilibus ovatis denticulatis acutis subtus tomentosis margine revolutis. — in Louisiana ex Robin, voy. p. 448. An fortè *E. mollis*, sed cor. lilacina dicitur.

†† *Species exclusæ.*

- E. ANGUSTIFOLIUS* Sw. = *Elephantosis angustifolia*.  
*E. NUDIFLORUS* Spr. = *Elephantosis angustifolia*.  
*E. NUDIFLORUS* Willd. = *Distreptus nudiflorus*.  
*E. SPICATUS* Juss. et Lam. = *Distreptus spicatus*.  
*E. STRIGOSUS* Willd. = *Distreptus spicatus*.

XXXVII. *ELEPHANTOSIS* Less. in linn. 1829. p. 322. f. 14-17 syn. 149.

Omnia Elephantopi sed pappus biserialis, paleis angustissimis rectis. — Capitula seu glomeruli spicati secus caulis apicem dispositi.

1. *E. BIFLORA* (Less. l. c.), pappi serie exteriori brevi ab interiori longitudine distinctâ, capitulis bifloris. ½ in Brasiliâ tropicâ. Pappi series interna setis deciduis tortis constans.
2. *E. QUADRIFLORA* (Less. l. c.), pappi seriebus longitudine non distinctis, capitulis 4-floris, glomerulis inferioribus subpedunculatis, foliis utrinque villosis, remotè dentatis. — in Brasiliâ tropicâ juxta Navium Friburgum (Sell.!), in prov. Sancti Pauli (Lund!). Folia etiam in specim. herb. Mus. Bras. video remotè denticulata! An ideò *El. liatroides* cat. h. Petasp. certè distincta? (v. a.)
3. *E. ANGUSTIFOLIA*, pappi seriebus longitudine non distinctis, capitulis 4-floris, glomerulis sessilibus, foliis glabrescentibus. ½ in locis montosis Jamaicae, Sancti Domingi (Bertero!) et Brasiliâ circa Bahiam (Salzman!). *Elephantopus angustifolius* Swartz fl. ind. 1383. Less. in linnæa 1829. p. 327. *Distreptus angustifolius* Cass. *Eleph. nudiflorus* Spreng. syst. 4. p. 454. non Willd. — Sloan. hist. 5. t. 148. f. 4. (v. a.)

XXXVIII. *DISTREPTUS* Cass. bull. philom. apr. 1817. dict. sc. nat. 13. p. 366. Less. in linnæa 1829. p. 327. f. 19. 33. 34. syn. 149. — *Pseudo-Elephantopus* Boer in Skrivt. af. n. h. Selsk. Kiøb. 11. p. 2. p. 213. — *Matamoria* Lallav. et Lex. nov. veg. descr. 1. p. 8.

Omnia Elephantopi sed pappi paleæ duæ cæteris longiores apice deorsum deindè sursum plicatæ (vid. Gærtn. t. 165. ic. *Eleph. spicati*). — Capitula secus axin ad foliorum axillas sessilia et ideò spicam interruptam ferè formantia.

1. *D. SPICATUS* (Cass. et Less. ll. cc.), foliis floralibus saltem inferioribus capitula superantibus. ½ in cultis ruderalisque Jamaicae, Hispaniolæ, Porto-Ricci, Martinicæ, Cubæ, Trinitatis, Mexici ad Veracruz et Tampico, prov. Caracanzæ Mexici, etiam in Philippinis verisim. introductus. *Elephantopus spicatus* B. Juss. in Aubl. guian. 808. Swartz! fl. ind. occ. 1380. Poepp. l per. exs. n. 38. *Matamoria spicata* Lallav. et Lex. nov. veg. descr. 1. p. 8. — Sloan. hist. t. 150. f. 3. 4. Flores albi. Variat foliis latioribus et angustioribus, asperficie glabrata aut villosâ, ramis erectis aut divaricatis. Huc *E. strigosus* Willd. herb. adductus ut mera varietas à cl. Lessing. (v. a.)
2. *D. CRISTATUS* (Cass. dict. 60. (1830.) p. 601. excl. syn.), foliis hirtis, floralibus saltem inferioribus densè ciliatis capitula superantibus, spicis acuminatis, achænis glaberrimis, pappi setis spiraliter tortis. — in Jamaica. *D. spiralis* Less. in linnæa 1831. p. 690. syn. 149.
3. *D. NUDIFLORUS* (Less. l. c. p. 329), foliis floralibus capitulis dimidio brevioribus. — in Sancto Domingo. *Elephantopus nudiflorus* Willd. sp. 3. p. 2390.

non Spreng. — Sub hoc nomine habeo ex Tabago? stirpem priori similem, sed apica magis continua, capitula minora ad axillas fol. flor. geminata terna quaternave 4-flora. Pappi paleæ 4-5 apice crispato-undulatae nec regulariter bis flexæ. An eadem ac stirps Willd. et Less.?

†. *Species exclusa.*

D. ANGUSTIFOLIUS Cass. = *Elephantosis angustifolia*.

DIV. III. ROLANDRÆ Cass. *tabl. xx. sect. v. (excl. gen.) Less. syn. 150.*

Capitula 1-flora. Cor. 5-fida regularis. Invol. 1-seriale ad summum 4-5-phyllum sæpe gamophyllum aut rarissime 2-phyllum basi breviter bracteolatum. Pappus brevis apicem coroniformis. — Subtribus forsitan artificialis et typus colligens plurimos valde diversos ob numerum specierum paucissimum nondum separatos.

XXXIX. GUNDELIA *Tourn. itin. 2. p. 251. ic. cor. 51. inst. t. 486. Gaert. fr. 2. p. 386. t. 163. Cass. dict. 20. p. 93. Less. in Linnæa 1829. p. 334. f. 43. 47. 68. 74. 76. 79-83. syn. p. 151. — Hacub Vaill. act. sc. par. 17. 18. p. 150. f. 18. — Gundelsheimera Cass. tabl. xx. n. 52.*

Capitula ad bractearum axillas 5-7 uniflora inter se ferè ad apicem concreta, centrali fertili, cæteris sterilibus. Involucra 1-serialia gamophylla flore breviora apice spinosa cum adstantibus demum connata. Cor. regularis glabra, limbo tubo longiore 5-fido. Antheræ inclusæ. Styli rami semisubulati extus pilis erectis hispidi intus plani glabri. Achænium glabrum compresso-subtetragonum, nectario alveolari basi pyramidato, disco epigynuo magno. Pappus 1-serialis coroniformis, ore brevi spinuloso. — Herba lactescens Dipsaci seu Eryngii facie. Folia alterna sessilia semiamplexicaulia pinnatilobata, lobis dentibusque spinosis ferè ut in Acantho spinoso. Flores purpurascens. Capitula suprà descripta in capitulum generale ovatum basi exinvolucratum adgregata.

1. G. TOURNEFORTII (Lin. sp. 1315). 2. in Persiâ, Armeniâ, Syriâ, agro Aleppensi, locis aridis et lapidosis. *Tourn. II. cc. Silybum seu Hacub, etc. Rauw. itin. 74. ic. — Lam. dict. 3. p. 60. Mill. ic. t. 267.*

a. *glabra*, capitulis glabris. — G. glabra Mill. dict. n. 2 — *Tourn. cor. 51. n. 1. (v. v.)*

f. *araneosa*, capitulis araneosâ lanugine obsitis. — G. Tournefortii Mill. dict. n. 1. — *Tourn. cor. 51. n. 2. (v. s.)*

XL. CORYMBIUM *Linn. gen. n. 1004. Lam. ill. t. 723. Gaert. fr. 2. p. 42. t. 86. DC. recueil mém. 1. p. 4. Cass. dict. 10. p. 380. Lessing in Linnæa 1829. p. 330. f. 13. 20-25. et 1831. p. 691. syn. p. 150. — Contarena Adans. fam. 2. p. 120.*

Capitulum 1-florum. Involucrum cylindraceum 2-phyllum, squamis canaliculatis liberis, basi foliolis exiguis paucissimis calyculatum. Cor. carnulosa regularis, limbo 5-partito. Antheræ inclusæ, in sicco subliberatae. Styli rami semicylindracei. Achænium teres oblongum basi longè attenuatum extus undiquè pilis longissimis densis obsitum. Pappus 1-serialis brevis membranaceus, in lacinias piliformes fissus. — Herba Capensis, radice crassâ perenni, collo lanuginoso, foliis radicalibus linearibus parallelè plurinerviis limbo nullo ad petiolum reductis. Caulis subnudus foliis parvis semiamplexicaulibus apice ramoso-corym-

bosus. — Falsò dixerunt Gærtner semen albuminosum, Willdenow antheras verè liberas, Cassini involucri squamas cum ovario et inter se concretas.

1. *C. nervosum* (Thunb. fl. cap. 729), totum glaberrimum, caule subangulato lævi, foliis latè lanceolato-linearibus, nervis subtus valdè prominulis, marginibus crassis callosis, corymbo laxo, invol. squamis lævibus.  $\frac{1}{2}$  ad Cap. B. Spei, circa urbem (Th. Gaud. ! Burch. ! n. 6771), ad montem Tabularem et Kleindrakenstein (Dreg. !); in distr. Uitenhagen (Eckl. et Zeyh. !). *C. glabrum* Lin. f. suppl. p. 392. Lam. ill. t. 273. f. 2. Less. l. c. non Thunb. Pluk. t. 272. f. 4? Axillæ et bases foliorum nunc glabræ nunc subbarbatæ. Folia 3-6 lin. lata, nervis 5-7 subtus quasi quadratis. (v. s.)
2. *C. glabrum* (Thunb. fl. cap. 729), totum glaberrimum, caule subangulato lævi, foliis linearibus seu angustè lanceolatis, nervis subtus prominulis, marginibus non incrassatis, corymbo laxo, invol. squamis lævibus.  $\frac{1}{2}$  ad Cap. Bonæ Spei (Th.), ad Drakensteinsberg et Onder-Bokkewald (Dreg. !), in distr. Worcester (Eckl. et Zeyh. !). *C. gramineum* Lam. dict. 2. p. 129? ill. t. 723. f. 3. Affine priori sed folia longiora, 2 lin. tantum lata, nervis multò minùs prominulis, margine non incrassato. (v. s.)
3. *C. cymosum* (E. Mey. ! in Dreg. coll. n. 1525), totum glaberrimum, caule teretiusculo lævi, foliis linearibus ad marginem non incrassatis, nervis vix subprominulis, corymbo conferto, invol. squamis lævibus.  $\frac{1}{2}$  ad Cap. Bonæ Spei, in parte Colonie orientali (Burch. ! n. 6661), ad Paarlberg (Dreg. !), in Stellenbosch, Swellendam et circa urbem (Eckl. et Zeyh. !). Priori affine sed corymbus confertus nec latus. An *C. gramineum* Lam. hæc potiùs referendum? (v. s.)
4. *C. scabrum* (Linn. f. suppl. 392), caule tereti muricato, foliis elongato-linearibus glabris planiusculis, invol. squamis extus muricatis.  $\frac{1}{2}$  ad Cap. B. Spei in lateribus montium prope urbem (Th.), ad Paarlberg (Dreg. !), distr. Worcester (Eckl. et Zeyh. !). Thunb. fl. cap. 729. Lam. ill. t. 723. f. 1. Less. l. c. — Burm. afr. t. 70. f. 1. Pluk. alm. t. 272. f. 5. mal. et ideò *C. Africanum* Lin. sp. 1317. Burm. ! prod. 25. sed *C. glabrum* eodem nom. insignitum adest. Folia 3-5 lin. lata, basi canaliculata. (v. s.)
5. *C. viliforme* (Lin. f. suppl. 392), caule tereti muricato, foliis convolato-semiteretibus glabris, involucri cylindrici squamis extus muricatis.  $\frac{1}{2}$  ad Cap. B. Spei in monte Tabulari et aliis circa urbem locis (Burch. ! n. 6663. Dreg. ! Eckl. et Zeyh. !). Less. l. c. Thunb. in fl. cap. fortè non immeritò cum priorè conjunxit. *C. gramineum* Burm. ! prod. fl. cap. 25. (v. s.)
6. *C. lutrum* (E. Mey. ! in Dreg. coll. n. 1527), caule tereti muricato subsetuloso, foliis lineari-filiformibus glabris, invol. oblongi subturbinati squamis vix subscabris.  $\frac{1}{2}$  ad Cap. Bonæ Spei circa Paarlberg legit cl. Dregel! Corollæ luteæ. Habitus *C. filiformis* sed invol. squamæ latiores magis foliaceæ vix ac ne vix scabræ. (v. s.)
7. *C. villosum* (Less. in Linnæ 1831. p. 691), caule hirsuto, foliolis planis utrinquè muricatis et hirsutis, invol. squamis muricatis subvillosis.  $\frac{1}{2}$  ad Cap. Bonæ Spei. Burch. ! n. 6760. Dreg. ! Eckl. et Zeyh. ! *C. villosum* et *C. hirsutum* Thunb. fl. cap. 730 et 729. *C. hirsutum* Ecklon ! pl. exs. cap. in h. Donant. Variat plùs minùs hirsutum et cum pili pauciores magis muricatum evadit. (v. s.)
8. *C. congestum* (E. Mey. ! in Dreg. coll.), caule hispido, foliis elliptico-lanceolatis utrinquè attenuatis submembranaceis utrinquè hirtellis, corymbo congesto, involucri squamis hirtellis.  $\frac{1}{2}$  in Cap. Bonæ Spei ad Drakensteinsberg alt. 2-300 ped. legit cl. Dregel! et in distr. Worcester cl. Ecklon et Zeyher. Folia infer. pollicem, imò sesquipollicem lata. (v. s.)



**XLI. ROLANDRA** *Rotb. coll. hafn. 2. p. 256. Swartz fl. ind. occ. 1388. t. 17. Cass. dict. 46. p. 170. Less. in linn. 1829. p. 332. f. 44. 75. syn. p. 150. — Echinopsis sp. Lin.*

Capitula 1-flora super receptaculum generale piloso-fimbrilliferum globosum adgregata. Invol. 2-valve compressum, valvis membranaceis carinatis aristatis, exteriore longiori. Cor. regularis glabra 3-5-fida, sæpius 4-fida, limbo subinflato. Stam. filamenta lævia. Styli rami semisubulati. Achænium oblongo-teretiusculum 4-costatum glabrum glandulosum. Pappus 1-serialis coroniformis brevis dentatus, dentibus ciliatis. — Suffrutex. Folia alterna breve petiolata oblonga utrinque attenuata supra glabra muriculata subtus araneoso-villosa alba. Capitula generalia globosa ramulos terminantia sæpè ramulis abbreviatis falso-axillaria. Cor. alba.

1. *R. ARGENTEA* (Rotb. l. c.). ♂ in subhumidis Jamaicæ, Guadalopæ, Surinami, Cajennæ, Bahiz, etc. Sloan. hist. jam. 1. p. 43. t. 7. f. 3. Plum. ed. Burm. t. 123. f. 1. Echinops fruticosus Lin. Echinops nodiflorus Lam. dict. p. 337. Variet foliis oblongis ovalibusve semper acuminatis, subtus plus minus sericeo-argenteis. Variet adhuc cor. lobis 3, 5 sæpius 4; involucri valvulâ minore nunc longiusculè nunc brevè aristatâ, nunc muticâ, in speciminibus ceterum similibus, undè *R. monocantha* et *R. diacantha* Cass. dict. 46. p. 171. vix ut varietates habendæ. (v. s.)

† *Species exclusæ.*

*R. REPTANS* Willd. = *Trichospira menthoides*.

*R. TERMINALIS* Spreng. = *Bailliera commelinoides*.

**XLII. SPIRACANTHA** *H. B. et Kunth nov. gen. am. 4. p. 25. t. 313. Less. syn. 150. — Acosta DC. mss. 1810.*

Capitula 7-11 uniflora adgregata in glomerulum ovatum bracteatum et ad bractearum axillas solitaria sessilia. Bracteæ subdisticho-imbricatæ basi concavæ in spinam desinentes. Involucrum 4-5-phyllum, foliolis lanceolatis diaphano-membranaceis subæqualibus. Cor. regularis, tubo gracili, fauce infundibuliformi, lobis 5 lineari-lanceolatis. Stylus apice bifidus? Achænium obovato-cuneatum læve. Pappus squamellulis pili-formibus brevibus integris scabridis constans. — Suffruticulus ramosissimus. Ramuli subpubescentes filiformes. Folia alterna brevè petiolata oblonga utrinque attenuata integerrima suprâ glabra subtus pube adpressâ albâ. Glomeruli ad apices ramulorum axillarium et ad apicem ramorum. Cor. purpureæ. — Genus affine *Rolandræ*.

1. *S. CORNIPOLIA* (H. B. et Kunth l. c.). ♂ in umbrosis humidis Novæ Granatæ ad portum Sapote prope Rio Sinu. Acosta *Rolandræ* DC. mss. (v. s. comm. cl. Bonpland.)

**XLIII. TRICHOSPIRA** *H. B. et Kunth nov. gen. am. 4. p. 23. t. 312. Less. in linnæâ 1829. p. 342. syn. p. 150.*

Capitulum multiflorum seu glomerulus polycephalus, bracteis è rachide ortis margine scariosis apice fimbriatis basi 1-nerviis apice 3-nerviis, in axillâ florem unicum foventibus sed involucrum verum subnullum. Cor. regularis glabra, tubo tereti, fauce campanulatâ, lobis 4 obtusis. Anthæræ è fauce semiexsertæ. Styli rami exserti semisubulati. Achænium compressum oblongo-obovatum basi acuminatum puberu-

medio utrinquē 3-costatum. Pappus 1-serialis, aristis nunc icatis nunc insuper 4-5 prioribus adjectis multò minoribus. — humiles Americanæ Filaginis facie. Radicellæ pubescentes. — lterna aut opposita subtùs lanâ albâ tomentosa sessilia integra ervia superiora opposita. Capitula axillaria et oppositifolia sess. — Genus affine Rolandræ et Spiracanthæ, Lessingii subtribus.

\* *Pappo pluriaristato.*

*ermodis* (H. B. et Kunth l. c.), pappo pluriaristato, caule repente pro-  
ente, foliis fere omnibus auriculato-semiamplexicaulis. — in arenosis  
Apures prov. Varinensis et in Cubâ. Less. in *linnæa* 1829. p. 342. f. 63.  
sila reptans Willd. herb. ex Less.

*negum* (Mart. ! herb.), pappo pluriaristato, caule repens à collo prostratus  
is non radicanibus, foliis fere omnibus auriculato-semiamplexicaulis.  
herbidis prov. Rio-Negro Brasilis fluv. Japura confluens in hunc. — Mar-  
Valdè accedit ad T. menthoidem sed ex omni parte major, et foliis  
repens et aristæ fructus paulò longiores. Squamæ paucissimæ brevissimæ  
(v. a.)

\*\* *Pappo biaristato.*

*aristata* (Less. ! l. c. f. 57. 58. 65. 66. 70), pappo biaristato, caule erecto  
ex ascendente, foliis alternis subtùs tomentosis, superioribus tantum au-  
stis. (f) ? in Brasiliâ tropicâ. Priori paulò minor. (v. s. comm. à Mus.  
berol.)

*umbra*, pappo biaristato, caule erecto, foliis sinuato-dentatis suprâ gla-  
abrâ canescenti-tomentosis, infer. alternis obovatis in petiolum attenua-  
super. oppositis basi dilatato-auriculatis. (c) in Guianâ Gallicâ ad ripas  
Gouana, in solo argillaceo paludoso legit cl. Le Prieur, qui ex naufragio  
nimo hanc et paucas alias salvavit! Herba pedalis. (v. s. comm. à cl. inv.)

*LAGASCEA* H. B. et Kunth nov. gen. am. 4. p. 24. Less. in *linnæa* 1829.  
f. 36-38. 45. 46. 48. 78. — *Noccea* Cass. dict. 25. p. 102. non Mæench.

tula 1-flora in glomerulum generale adgregata. Involucrum  
imm) 1-seriale gamophyllum 4-5-dentatum tubulosum. Cor. re-  
extus pubescens, tubo gracillimo, fauce dilatâ cylindraccâ,  
i. acuminatis. Stam. filam. papillosa. Styli rami semisubulati.  
tum obcompressio-tetragonum pluricostatum. Pappus 1-serialis  
formis membranaceus brevissimus nunc subinteger nunc denta-  
Herbæ aut suffrutices Americanæ. Folia opposita sæpius tripli-  
Capitula generalia terminalia bracteis plurimis (exterioribus  
tis) obvallata. Flores albi, flavidi aut rubri. — *Nocceæ* nomen  
avetustius (1794), sed Kunth, qui primus genera duo conjunxit,  
om nomen admisit ex usu hortulanorum. Cur autem cl. Lessing  
vera distinxit eorum nomina transposuit, ignoro.

*LAGASCA* Cav. anal. sc. nat. 6. (1800) p. 333. t. 44. et omn.  
auct. excl. Less. — *Noccea* Jacq. fragm. (1805) p. 58. t. 85.  
Less. syn. 151. non Mæench.

coroniformis minimus subinteger ciliatus. — Herbæ annuæ. Glomerati  
um pedunculi elongati nudi solitarii. Folia petiolata. Cor. albæ.

*OLLEA* (Cav. anal. sc. nat. 6. p. 333. t. 44), herbacea, foliis petiolatis ovatis  
sinatis subcrenatis molliter sericeo-tomentosis, capitulis generalibus soli-  
longè pedunculatis. (i) in collibus insulæ Cubæ et ad Santa-Fé Mexica-

norm, Desv. Journ. bot. 1. p. 28. t. 2. *Noccea mollis* Jacq. frag. t. 13. Flores albi. Involucelli foliola concreta 3-nervia. Capitula et bractes sericeo-villosæ. (v. v.)

Sect. II. *NOCCÆA*. — *Nocca Cav. ic. 3. (1794) p. 12. t. 224. non Less.*  
— *Lagascea Less. syn. 131. non Cav.*

Pappus paleis brevibus subdentatis. — Suffrutices aut herbæ perennes à Mexico. Glomeruli adgregati foliis bracteati. Folia sessilia aut vix petiolata. Cor. albida ochroleucæ aut purpurascens.

2. *L. RUBRA* (H. B. et Kunth nov. gen. am. p. 24. t. 311), suffruticosa, foliis breviter petiolatis ellipticis ovatisve subacutis dentatis coriaceo-rigidis scabris, capitulis glomerato-corymbosis. 2 in temperatis Mexici ad montes Guchilapa, etc. *Noccea rubra* Cass. l. c. *Nocca rigida* Cav. ic. 3. p. 12. t. 224. Sweet! br. fl. gard. ser. 2. p. 26. Flores pallidè rubentes. (v. s.)
3. *L. MACROPHYLLA* (Cass. in Moench. ined. mexic. 1814), suffruticosa, foliis petiolatis ovatis ovalibusve serratis supra piloso-scabridis subtus villosis, paniculae ramis distantibus oppositis, invol. generalis squamis 4-6 lineari-lanceolatis floribus oppositis. 3 in Mexico observ. cl. Moquinno et Hænke. Huc referenda est cl. Moquinno *Cacansca* Hern. mex. 184. ic. sed foliis sessilibus et glomerulorum dispositione differt et potius *Lagascea* species hodiernis ignota in nostrâ ex ic. floræ Mexic. flores purpurei. Caulis rami et involucri apice glanduloso-villosi; invol. sericeo-villosissima. Folia 2-2½ poll. longa, pollicem lata. (v. ic. pict. Moq. et sp. sicc. in h. Hænke.)
4. *L. LATIFOLIA*, fruticosa, foliis sessilibus amplexicaulibus oblongis acuminatis serratis integerrimisve supra scabris subtus molliter pilosis reticulatis, capitulis corymboso-paniculatis. 3 in Mexico. *Nocca latifolia* Lallav. et Lex. nov. veg. descr. 1. p. 31. Sweet br. fl. gard. t. 215. Flores suaveolentes ochroleuci aut albi. Cor. fere pollicem longæ. Styli longè exserti. (v. s. ex hort. Mexic. ubi Cervantez eam descr.)
5. *L. HELIANTHIFOLIA* (H. B. et Kunth l. c.), suffruticosa, foliis sessilibus amplexicaulibus oblongis subacuminatis cordatis serratis coriaceo-rigidis supra scaberrimis subtus scabriusculo-hirtellis, capitulis subcorymbosis. 3 juxta Acapulco Mexic. *Noccea helianthifolia* Cass. l. c. *Noccea macrophylla* Zucc. in h. Acad. reg. Monac. Cor. albida? pollicem fere longæ. Folia 7-8 poll. longa, 2 lata. Involucellum externè sericeo-pilosum. (v. s. comm. à cl. Martet.)
6. *L. SUAVEOLENS* (H. B. et Kunth l. c.), suffruticosa, foliis sessilibus amplexicaulibus oblongis acuminatis supernè serratis coriaceo-rigidis supra scaberrimis subtus secus nervos valdè prominentes molliusculè puberulis inter eos sparse subscabridis, capitulis corymboso-paniculatis. 3 in declivitate orientali montium Mexicanorum. *Noccea suaveolens* Cass. l. c. Cor. alba. Folia 4 poll. longa, 15 lin. lata more *Borraginearum* demùm supra albo-punctata. (v. s. in h. Hænke. à cl. Sternb. comm.)
7. *L. ANGUSTIFOLIA*, suffruticosa, foliis brevè petiolatis oblongo-linearibus integerrimis utrinquè attenuatis villosis-scabris, capitulis villosissimis subsessilibus. 3 in Mexico circa Leon legit cl. Mendez. Cor. ex statâ immaturo 3 lin. longæ albida. Involucellum villosissimum 5-dentatum, dentibus subulatis. Folia 3-4 poll. longa semipollicem lata non triplinervia. (v. s.)

#### Div. IV. *BOJERIAE*.

Capitula plariflora homogama. Anthæræ basi bicaudatæ. Pappus setis 1-seriatis. — Arbores frutices aut herbæ Madagascarienses, omnes à cl. Bojer detectæ.

#### XLV. *SYNCHODENDRON* Bojer in litt. 1833.

Capitulum pluriflorum homogamum (ex Bojer), recept. angusto nudo, invol. obconici squamis multiseriatis imbricatis coriacis siccis gle-

brevissimis, intimis longioribus linearibus. Cor. tubulosæ. Anth. longè caudatæ exsertæ. Styli rami breves. Achænia teretiuscula subvillosa. Pappus uniserialis, setis brevibus sublaminatis cor. longitudine æquantibus. — Arbor fruticosa 50-pedalis Compositarum omnium maxima! Rami sparsi et imò folia pube brevi velutino-tomentosâ obsecta. Petiolata integra orbiculata aut obovatâ, demum supernè sessile ante folia secus ramos juniores et crassos sessilia 2 lin. longa. (Bojer.)

(Bojer! in litt. 1833 et 1835). 3 in insulâ Madagascar prov. in vallibus inter montes excelsos et circa pagos cultum. Florescentia circa Madagascariensibus tempus oryzae culturae proprium indicat v. a. comm. à cl. inv.)

#### XLVI. CENTAUROPSIS Bojer in litt. 1833.

pluri- (12-20-) florum homogamum. Recept. angustum nudum linearibus paucis deciduis onustum. Invol. oblongum nudum, squamis coriaceis obtusis, ext. brevissimis, inter. linearibus. Cor. tubulosâ, fauce vix dilatâ, limbo 5-mer. incurvæ majores. Anth. loculi basi producti in apiculiferas? Stigm. 2 longa exserta divergentia recurvato-hispida. Achænium teretiusculum gracile striatum glaberrimum. Pappus 1-serialis è margine calyculi coroniformis superantibus ortus rigidus, setis inæqualibus achænio duplò brevioribus, in margine capituli brevioribus aut subnullis. Madagascarienses sylvicolæ. Folia alterna breviter petiolata ovata obovatâ aut ovalia basi cuneata penninervia. Capitula aut solitaria ovato-oblonga. Flores ex sicco purpurascetes. (v. Stokesiæ, Platycarphæ et Distephano.)

(Bojer! in litt.), foliis latè obovatis obtusissimis suprâ adultis, subtus ramulis pedunculisque densè velutino-tomentosis, capitulo rymbois. 5? 3? in insulâ Madagascar sylvæ vastis loco Bè-Fou- detexit cl. Bojer! Folia 6 poll. longa, 4 poll. lata, basi distinctè angulata jun. furfuraceo-tomentosa, adulta punctatis elevatis subscabrida. Flores exteriores dorso velutinæ. Flores in invol. 12-13. Cor. extus (v. a. comm. à cl. inv.)

(Bojer! in litt.), ramulis junioribus vix pubescentibus, foliis pedunculisque glaberrimis, pedunculis solitariis angulatis. 3? ad marginem in prov. Emirneni inter. ins. Madagascar detex. cl. Bojer. Priori. Folia pollicem longa, 5-6 lin. lata. Pedunculi folii circ. longitud. Marshalli sub capitulo bracteâ foliaceâ ovalem sessilem gerentes. Priori sed pappus paulò longior et magis caducus. Flores vix long. circ. 20 in invol. (v. a. comm. à cl. Bojer.)

#### XLVII. TECMARSIS.

circ. 6-florum homogamum. Recept. angustum nudum. Invol. ante explicationem acutum, squamis arcè imbricatis, setis obtusis, inter. linearibus subacutis. Cor. tubulosâ filiformi-plicatâ. Anth. Stigm. Achænia gracilia subangulata glabra erostris. Pappus 1-serialis, setis basi in tubum conæqualibus erectis subrigidulis achæni longitudine æqualibus superantibus. — Frutex Madagascariensis debilis,

ramis teretibus procumbentibus apice setoso-ciliatis villosisque. Folia alterna fasciculata glabra obovato-oblonga acuta basi cuneata paucidentata, petiulis ciliato-villosis brevibus. Capitula corymbosa, pedunculis ramosis nudis. Invol. glabra. — Genus ex floribus ignotis dubium, et ideo Tecmarsis (græcè conjectura) dictum, fortè ex invol. et pappo ad Centauropsidem et Synchodendron accedens.

1. T. BOJER. ♂ ad margines sylvarum in locis umbrosis insulæ Madagascar legit cl. Bojer, qui mecum sine nomine commun. Rami et folia in sicco non male Bucidam referunt. (v. s. comm. à cl. invent. sed capitula alia juniora minima alia nimis ætate provecta!)

#### XLVIII. BOJERIA.

Capitulum multiflorum homogamum, fl. tubulosus 5-fidis hermaphroditus. Invol. campanulatum polyphyllum, squamis multiseriatis longitudine subæqualibus acutis, ext. lineari-lanceolatis foliaceis dorso villosis, int. linearibus glabriusculis. Recept. planum subalveolatum. Anth. basi bicaudatæ. Stylus basi bulbosus, ramis exsertis teretibus pube minimâ scabridis. Achænia striato-subangulata glabra. Pappus 1-serialis, setis longis subscabris basi inter se variè et brevè subconcretis. — Herba perennis elata Madagascariensis. Caulis teres ramosus præsertim ad apices densè villosus. Folia caulina alterna amplexicaulia ovato-lanceolata integra suprà sparsè puberula subtùs subtomentosa. Capitula ad apices ramorum solitaria. Flores numerosissimi. Cor. purpureæ. Calyx in achænio facilè à pericarpio separatur et pappum secum trahit! Pericarpium teres apice cupulam exiguum styli basi per anthesin foveentem gerit. — Genus spectabile Eupatoriaceis stylo subaffine, sed habitu et antheris Vernoniaceis referendum, dicavi cl. Guill. Bojer, hist. nat. in insulâ Mauritii professori, plantæ detectori et qui insulas Mauritanas, Madagascar et alias Australi-Africanas laboribus indefessis egregiè explorat.

1. B. SPECIOSA. in locis humidis insulæ Madagascar circa montes Angavon dictos detexit cl. Bojer. Herba interdum 10-pedalis. Capitula diam. pollicari. (v. s. comm. sine nomine à cl. invent.)

#### Subtrib. II. PECTIDÆ Less. in linnæa 1830. p. 134.

Capitula radiata nec discoidea. Styli flor. herm. rami apice obtusi nec acuminati. Folia sæpius opposita. Herbæ aut frutices. Species omnes Americane. Flores sæpè flavi.

- DIV. 1. LIABEÆ Cass. tabl. p. 28. Less. in linnæa 1831. p. 696. syn. p. 151. — Vernonicarum gen. Cass. Less. in linnæa 1829. p. 318.

Cor. disci regulariter 5-fidæ. Stylus fl. herm. ramis longiusculis plerumque obtusis. — Herbæ seu frutices. Folia sæpius opposita eglandulosa basi ciliis destituta, subtùs tomentosa.

#### XLIX. XANTHISMA.

Capitulum multiflorum radiatum, fl. radii ligulatis neutris? seu femineis? integris, disci hermaphroditis 5-fidis lobis erectis. Recept. fimbrilliferum, fimbrillis laceris achænio ferè longioribus. Invol. hemisphæricum, squamis imbricatis coriaceis ferè ovalibus obtusissimis

adpressis. Antheræ (disci) ecaudatæ. Stylus radii brevis simplex intra partem ligulæ tubulosam inclusus, disci apice bifidus, lobis linearibus inclusis obtusis breviter hispidulis. Achænium obovatum vix subangulatum pubescens. Pappus paleis elongatis acuminatis inæqualibus, exter. paulò brevioribus et angustioribus. — Herba Mexicana monocarpica erecta glabra simplex apice laxè ramosa subcorymbosa, ramis erectis foliosis. Folia caulina alterna sessilia oblonga aut linearia uninervia rigidula, pleraque integerrima, pauca apice acutè serrata. Flores lutei.

1. X. TEXANUM. (1) (2) in sylvis Mexicanæ provinciæ Texas legit cl. Berlandier (pl. exs. n. 263g). Herba habitu ad Centauream cerinthefoliam ferè accedens. (v. s.)

## L. HECTORÆA.

Capitulum multiflorum radiatum, ligulis fœm. 12-15 acutè 3-dentatis. Invol. pluriseriale, squamis acuminatis subimbricatis. Recept. alveolatum. Cor. disci glabræ 5-fidæ. Stylus teres nec basi bulbosus, ramis teretibus hispidulis. Achænium teres villosum. Pappus biserialis, serie exter. paleaceâ brevi, inter. elongatâ setiformi scabrâ. — Herba Mexicana basi suffruticosa simplex erecta tota cano-villosa. Caulis teres ad apicem usquè foliosus. Folia alterna conferta erecta oblongo-lanceolata integra mucrone calloso superata. Capitula ad apicem caulis breviter pedicellata subcorymbosa flava. Pappi series ext. alba, int. subrufa. — Cor. Andromachiæ sed habitus diversissimus, stylus non bulbosus, etc. Nomen ad Hectoris et Andromachiæ affinitatem alludit.

1. H. VILLOSISSIMA.  $\frac{1}{2}$   $\frac{3}{4}$  in Mexico prope Real del Monte legit cl. Hænke. Caulis pedalis. Folia 10-12 lin. longa, 2-3 lin. lata.

- LI. ANDROMACHIA H. et Bonpl. pl. eq. 2. p. 104. t. 112. Less. in *linnaea* 1829. p. 318. non Cass. — Andromachiæ sect. 2 et 3. H. B. et Kunth nov. gen. 4. p. 99. — *Oligactis* Cass. dict. 36. p. 16. — *Diplostephii* sp. Spreng. — *Viviania* Willd. herb. non Cav. nec Colla.

Capitulum radiatum, disco regulari multifloro, radio uniseriali ligulato fœmineo. Invol. squamæ imbricatæ siccæ acuminatæ. Recept. breviter paleaceo-fimbriatiferum. Cor. glabræ, disci 5-fidæ, lobis acuminatis revolutis tubo paulò brevioribus, radii ligulatæ, tubo longo, ligulâ oblongâ. Filam. lævia. Styli basi bulbosi rami semicylindracei. Achænium cylindraceum multicostatum, valleculis hirsutis, nectario styliformi. Pappus 2-serialis paleaceus serratus, paleis seriei ext. multò brevioribus et sublateralioribus quam inter. — Herbæ aut frutices. Folia opposita petiolata subtus tomentosa dentata. Capitula corymbosa.

## Sect. I. PLEIONACTIS.

Capitulum multi (20-30)-florum. Ligulæ radii crebræ. — Herbæ, caulibus foliisque subtus cano-tomentosis. Flores ligulati flavi. — Andromachiæ verè Kunth.

1. A. GRANDIFLORA (H. B. et Kunth l. c. p. 99), herbacea ramosa, foliis ovatis acutis leviter cordatis suprà glabris rugoso-bullatis subtus albo-lanatis, petiollis alatis basi auriculatis et connatis, capitulis 3is-4is.ve corymbosis.  $\frac{1}{2}$  in aridis prope urbem Alausi Quitensium. *Diplostephium grandiflorum* Spreng. *eyst.* 3. p. 544.
2. A. IGNIARIA (H. et Bonpl. pl. eq. 2. p. 104. t. 112), herbacea ramosa, foliis ovatis acutis basi obtusis suprà glabris subtus albo-lanatis, petiollis basi auricu-

latia et connatis, capitalis fasciculatis in corymbos terminales trifidos dispositis.  $\mathcal{L}$  in altâ planitie Quitensi. H. B. et Kunth nov. gen. 4. p. 100. Less. in *linnaea* 1829. p. 318. *Amellus floribundus* Willd. herb. ex Less. *Diplostephium igniarium* Spreng. l. c. *Liabum Bonplandii* Cass. dict. 26. p. 206. *Tomentum Agarici igniarii succedaneum*.

3. *A. solidaginea* (H. B. et Kunth l. c. p. 100), herbacea ramosa, foliis ovatis oblongis acuminatis basi rotundatis et in petiolum basi nudum decurrentibus supra glabris subtus albo-tomentosis, corymbis terminalibus ramosis, ramis fasciculatis.  $\mathcal{L}$  in Andibus Peruvianorum inter Ayavaca et fl. Cutaco. *Diplostephium solidagineum* Spreng. syst. 3. p. 544. (v. s. in h. reg. Acad. Mon. specimen Peruv. à descr. Kunth discrepans tantum tomento paginae inf. fo in sicco rufo nec albo.)
4. *A. verbascifolia* (H. B. et Kunth l. c. p. 100), herbacea ramosa, foliis oblongis acutis basi in petiolum nudum angustatis supra rugoso-bullatis glabris subtus lacunosis cano-lanatis, corymbis terminalibus ramosissimis, capitali fasciculatis.  $\mathcal{L}$  in montibus Quitensium cum Cinchonâ inter Malacates et Gusanama. *Diplostephium verbascifolium* Spreng. l. c.
5. *A. melastomoides* (H. B. et Kunth l. c. p. 101. t. 337), herbacea ramosissima foliis oblongo-ellipticis acuminatis basi acutis coriaceis supra scabris subtus cano-tomentosis, petiolis nudis, pedunculis axillaribus et terminalibus polycephalis.  $\mathcal{L}$  in temperatis prope Ibaguê Novo-Granatensium. Less. in *linnaea* 1829. p. 318. f. 1-3. 6. *Diplostephium melastomoides* Spreng. l. c. *Vivian bicolor* Willd. herb. ex Less.

## Sect. II. OLIGACTIS H. B. et Kunth nov. gen. 4. p. 101.

Capitulum pauciflorum. Ligulae radii 3-7. — Frutices, ramis glabris. Flores ligulati pallidi fortè albi?

6. *A. nubigena* (H. B. et Kunth nov. gen. 4. p. 102), fruticosa, foliis lanceolatis oblongis acutis basi obtusis membranaceis supra petiolisque glabris subtus nervo medio excepto albo-tomentosis, corymbis terminalibus ramosis, capitali pedicellatis.  $\mathcal{L}$  in excelsis montis Chimborazo. *Diplostephium nubigenum* Spreng. syst. 3. p. 543. *Oligactis nubigena* Cass. dict. 36. p. 17.
7. *A. sessilifolia* (H. B. et Kunth l. c.), fruticosa, foliis oblongis acutis basi rotundatis membranaceis supra glabris subtus petiolisque albo-tomentosis paniculis axillaribus et terminalibus, capitulis conglomeratis sessilibus.  $\mathcal{L}$  Peruvia? *Diplostephium sessiliflorum* Spreng. l. c. *Oligactis sessiliflora* Cass. l. c.
8. *A. volubilis* (H. B. et Kunth l. c.), volubilis, foliis lanceolato-linearibus acutis brevè petiolatis supra glabris subtus cano-tomentosis, paniculis terminalibus simplicibus, capitalis solitariis pedicellatis.  $\mathcal{L}$  in monte Antianâ Quitensium. *Diplostephium volubile* Spreng. l. c. *Oligactis volubilis* Cass. l. c.

## † Species exclusæ.

- A. acaulis* Kunth. = *Liabum acaule*.  
*A. hieracioides* Kunth. = *Liabum hieracioides*.  
*A. Jussieu* Cass. = *Liabum Jussieu*.  
*A. Portraui* Cass. = *Liabum Brownei*.

LII. *LIABUM* Adans. fam. 2. p. 131. excl. sp. Cass. dict. 26. p. 203. excl. 4 Less. in *linnaea* 1829. p. 319. — *Andromachia* sp. Kunth. — *Andromachia* Cass. bull. philom. 1817. p. 184. non *Bonpl.* — *Amelli* sp. *Linna.* — *Starb* Willd. sp. 3. p. 226. — *Starckia* Juss. dict. 50. p. 403.

Capitulum radiatum, disco regulari multifloro, radio uniseriale lig

lato fœmineo. Invol. squamæ imbricatæ subulatæ. Recept. fimbriis membranaceis subulatis hispidum. Cor. glabræ, disci 5-fidæ regulares, lobis 5 linearibus tubo brevioribus, radii ligulæformes longæ lineares apice integræ. Styli (disci) rami semiteretes graciles parçè hispidi. Achæmium cylindraceum striatum, callo basilari instr. Pappus 1-serialis setis filiformibus barbellatis subinæqualibus. — Herbæ Americanæ sæpè cano-lanatæ, caulescentes aut acaules. Folia opposita triplinervia subdentata. Flores flavi.

§. 1. *Capitulis plurimis corymbosis.* — *Starkea Willd. l. c.*

1. L. BROWNII (Cass. dict. 26. p. 203), caule erecto infernè folioso tomentoso, foliis ovatis subserratis in petiolum attenuatis suprà lanuginosis demùm glabris subtùs tomentosis, cymà umbellatà terminali, pedunculis elongatis aliis monocephalis oligocephalis. ½ in Jamaicà et Santo-Domingo. Solidago P. Brown. jam. 320. t. 83. f. 2. Amellus umbellatus Linn. sp. 1296. Swartz! obs. 310. *Starkea umbellata* Willd. sp. 3. p. 2216. *Andromachia Poiteavi* Cass. bull. philom. 1817. p. 184. Variat foliis obtusis et acutis. (v. s.)
2. L. JUSSIEI (Cass. dict. 26. p. 205), caule erecto glabriusculo subanguloso, foliis petiolatis ovalibus dentatis basi stipulato-auriculatis suprà glabris subtùs tomentosis, capitulis in cymam paniculato-corymbosam dispositis. ½ in Peruvia. *Andromachia Jussiei* Cass. bull. philom. 1817. p. 184. *Conyza stipulata* Vahl in herb. Juss.

§. 2. *Capitulis solitariis.* — *Chrysactinium H. B. et Kunth l. c.*

3. L. HIERACIOIDES, caule procumbente cano-lanato, foliis sessilibus connatis lanceolatis acutis subdentatis suprà pilosis subtùs cano-lanatis, pedunculis terminalibus 1-cephalis. ½ in siccis temperatis prope Lozam Quitensium. *Andromachia hieracioides* H. B. et Kunth nov. gen. am. 4. p. 98. *Solidago hieracioides* Spreng. syst. 3. p. 543.
4. L. ACAULE, caule subnullo, foliis radicalibus sessilibus lanceolato-oblongis basi angustatis remotè dentatis suprà glabris subtùs dense candido-lanatis, scapo erecto longissimo 1-cephalo lanato. ½ in montibus Quitensibus. *Andromachia acaulis* H. B. et Kunth nov. gen. 4. p. 98. t. 336. *Solidago acaulis* Spreng. syst. 3. p. 543.

† *Species exclusa.*

L. BONPLANDII Cass. = *Andromachia igniaria*.

LIII. ALIBUM Less. syn. 152.

Omnia Liabi sed pappus disci biserialis, serie exteriori coroniformi dentatà, interiori multisetaceà: radii coroniformis dentatus 1-serialis. — Herba trichotoma erecta Australi-Americana. Rami teretes hirsuti. Folia opposita obsoletè auriculata et connata ovato-elliptica integra seu basi pinnatifida suprà hirsutiuscula subtùs tenuiter albo-tomentosa. Capitula solitaria nutantia. Invol. squamæ biseriales æquales oblongo-ellipticæ mucronatæ plurinerviæ. Recept. planum nudum. Achænia turbinata hirsuta.

1. A. LIABOIDES (Less. l. c.). — in Pomelacte Americæ æquinoctialis.

LIV. CACOSMIA H. B. et Kunth nov. gen. 4. p. 289. t. 404. Cass. tabl. xx. n. 4. Less. in linnæa 1829. p. 338. f. 31. 53-55. syn. 152. — *Xantholepis Willd. herb.*

Capitulum radiatum, floribus disci plurimis hermaphr., radii ligulatis fœmineis. Invol. teres imbricatum, squamis siccis plurinerviis



interioribus longioribus. Recept. nudum. Cor. glabræ, disci regulares 5-fidæ, lobis acuminatis revolutis parte integrâ brevioribus, radii ligulæ latè oblongo-ellipticâ apice 3-crenulatâ tubo duplò longiore. Filam. lævia. Styli rami semicylindræci. Achænium obpyramidatum 4-quetrum truncatum glaberrimum, nectario styliformi. Pappus o. — Herba suffruticosa graveolens. Rami tenuiter lanati. Folia opposita subpetiolata connata ovato-oblonga remotè dentata trinervia suprâ rugosa glabra subtus pubescenti-cana. Corymbi ad apices ramorum axillarium. Cor. flavæ.

1. C. RUGOSA (H. B. et Kunth, l. c.). 2. in Andibus Peruviz. Xantholepis tomentosa Willd. herb. ex. Less. l. c.

**DIV. II. EUPECTIDÆ.** — Pectidæ Less. in *linnæâ* 1831. p. 706. *syn.* p. 152. — Tagetineæ Pectidæ pleræque Cass. *tabl.*

Corolla disci subbilabiata aut rarissimè regulariter 5-dentata. Styli disci rami semicylindræci breves. — Herbæ glabrescentes pleræque annuæ Americane. Folia opposita basi lineâ transversali subconnata pellucido-punctata uniuersia basi denticulis ætigeris ciliata, apice rariâ serrulata. Invol. 1-seriale. Recept. nudum. Cor. flavæ glabræ nunc regulares 5-dentatæ nunc lobo unico profundius fissæ subbilabiata, lobo exteriore indiviso, inter. 4-dentato. Anthæræ ecaudatæ. Stylus radii glaber. Achænia angulata basi attenuata, callo basilari. Pappus flavescens. — Pectidæ à cl. Cassini ad Tagetineas relatæ, à cl. Kunth inter Tagetem et Steviam mediæ, à cl. Lessing juxta Liabeas in Vernoniacearum tribu admixtæ.

#### LV. PECTIDOPSIS.

Cor. disci regulariter 5-dentatæ. Pappus paleaceus brevis 5-dentatus. — Herba borealis-americana basi ramosa diffusa lævis. Folia angustolinearia basi dentato-ciliata. Capitula solitaria brevissimè pedicellata. Invol. 8-phyllum. Radii ligulæ 7-8. — Habitus Pectidis sed pappo eodem jure ac Pectidium et Lorentea separanda.

1. P. ANGUSTIFOLIA. ① in Americ. bor. montibus dictis Rocky-Mountains. Pectis angustifolia Torr. acc. rock. mount. in ann. Lyc. N.-York. 2. p. 214.

**LVI. PECTIDIUM** Less. in *linnæâ* 1831. p. 706. *syn.* p. 152. — Pectis Cass. *dict.* 38. p. 202. non Less. — Pectidis sp. Linn.

Cor. disci regulariter 5-dentatæ. Pappi setæ subulatæ rigidæ cornæ læves nudæ demum divergentes. — Herba annua glabra erecta di- aut trichotoma. Folia linearia cilia utrinquè 1-2 basi gerentia glandulis grossis pellucidis ad utrumque nervi medii latus ornata. Capitula longè pedicellata solitaria. Invol. 5-phyllum. Ligulæ breves.

1. P. PUNCTATUM (Less. in *linnæâ* 1831. p. 707). ① in insulis Caribæis nempe Santo-Domingo (Bertero!), Santo-Thomasio (Ehr.), Sancta-Mariha (West!), Sanctæ-Crucis (Jacq.), Porto-Ricco (Ledru!). Pectis punctata Jacq. amer. 216. t. 128. Linn. sp. 1230. Swartz obs. p. 308. Linn. fil. herb. Pectis trifolia Gertn. fr. 2. t. 171 (excl. syn.). Lam. ill. t. 684. — Plum. et Barn. t. 86. f. 1? Sloan. jam. hist. 1. t. 149. f. 3? sed icon utraque pessima et fortè removenda. Conf. Pectidem linifoliam et Swartzianam. (v. s.)

**LVII. PECTIS** Less. in *linnæâ* 1831. p. 708. *syn.* 153. — Lorentea Lag. *nov. gen.* (1816). p. 28. Cass. *dict.* 27. p. 202. nec Less. nec Ort. — Chthonia Cass. *bull. philom.* 1817. p. 33. — Pectidis sp. pleræque Linn.

Cor. disci obscurè bilabiata. Pappus disci et radii uniserialis, paleis basi scariosis latiss, apice setiformibus serrulatis, sæpius inæqualibus. —

Herbæ annuæ aut rarius perennantes. Folia glabra margine augustissimè cartilaginea. Capitula pedunculata, ramis nempe floridis nudis aut bracteolatis 1-cephalis elongatis, aut subsessilia inter folia plùs minùs occultata.

\* *Capitulis pedicellatis, pedicellis nudis capitulo longioribus.*

1. *P. LINIFOLIA* (Linn. sp. 1250. amœn. 5. p. 107), caule suberecto trichotomo tetragono ad angulos puberulo, foliis linearibus mucronatis margine scabris grossè glandulosis basi cilia setosa gerentibus, capitulis pedicellatis 4-18-floris, invol. squamis 4-6 elliptico-lanceolatis conduplicatis, pappo latè paleaceo, setis disci 5 longis, radii 2 oppositis. — in Jamaicâ et Sagnet. Thomas. (Less.). — Sloan. jam. hist. 1. t. 149. f. 3. ex Less. in linnæâ 1831. p. 709, sed icon à descr. Less. in plurimis recedit nec malè Pectidis convenit.
2. *P. SWARTZIANA* (Less. in linnæâ 1831. p. 711), erecta, ramis patulis glabris subretrogonis, foliis oblongo-linearibus acutis basi cilia 2 utrinquè gerentibus grossè pellucido-glandulosis, pedicellis ex dichotomiis et apice ramorum sphyllis elongatis 1-cephalis, invol. squamis 5-6 linearibus obtusiusculis subconvolutis, floribus 10, achæniis glabriusculis, pappo 1-seriali brevi dentatoparvo radii brevi inæquali interdum uniaristato. ① in Sancti-Domingi aridis. Pectis patula Linn. fil. herb. *P. punctata* Swartz in herb. L'Hér. *P. ciliaris* Swartz obs. bot. 307. non Linn. *P. Swartziana*? *Cithonia leptocephala* Cass. dict. 27, p. 206? — Icon Plum. ed. Burm. t. 86. f. 1. ad *P. punctatam* à Linnæo et Lessingio relata huc meo sensu pertinet, sed pessima! Icon Plum. t. 151. f. 2. à Linnæo ad ciliarem suam relata toto cœlo differt, nec descr. amm. acad. convenit. (v. s.)
3. *P. CAPILLARIS*, caule erecto diffusè ramoso glaberrimo, foliis linearibus basi setigero-ciliatis subitùs glandulas grossas ad utrumquè nervi mediî latus gerentibus, pedicellis elongatis capillaribus, invol. squamis 5 oblongo-linearibus acutis margine scariosis, capitulis 12-15-floris, ligulis oblongis minimis, pappo 5-seriatis. ② in Mexici ditione Oaxacana prope las Bacas septemb. flor. Caulis pedicellique purpurascens tenues. Herba 5-6 poll. (v. s. comm. à cl. Mairet.)
4. *P. TAMELLA*, caule herbaceo erecto glabriusculo tetragono oppositè ramosissimo, foliis linearibus submucronatis marginibus subrevolutis minutè et parcè glandulosis cilia ad basin utrinquè sæpiùs solitaria gerentibus, pedicellis 1-cephalis parcè bracteolatis capitulo paulò longioribus, invol. squamis 8 linearibus obtusis convolutis, floribus 13-15, pappo radii 3-4, disci 5-6-setoso. ① in Mexico prope Laredo ad Tamaulipas legit cl. Berlandier flor. mense julio. Herba 3 pollic. glabra. Radix exilis simplex. Fol. 9-12 lin. longa. Flores flavi. (v. s.)
5. *P. ELONGATA* (H. B. et Kunth nov. gen. 4. p. 262. t. 392), caule erecto subtereti glabro virgato, foliis elongato-linearibus acutis basi dentibus aristatis utrinquè 3-8-ciliatis utrinquè subscabris, capitulis subcorymbosis ad apices ramulorum pedicellatis, involucri squamis 6 lineari-oblongis acuminatis, pappi paleis in disco et radio 10-20 setiformibus scabris inæqualibus. ① circa Papayan? (Kunth), ad Hondam (Willd.), ad Panama (h. Hænk!). Herba pedalis basi simplex. Folia 15-18 lin. longa, 1-1½ lata. (v. s. in h. Hænk. à cl. Sternb. comm.)
6. *P. BONPLANDIANA* (H. B. et Kunth nov. gen. 4. p. 263), erecta oppositè aut dichotomè ramosa, foliis lanceolato-linearibus acuto-mucronatis grossè pellucido-glandulosis basi dentibus aristatis 1-2 utrinquè ciliatis, capitulis terminalibus et alaribus longè pedunculatis solitariis, involucri sub-10-phyllo, pappo disci 5-setoso, radii pauciseti, utroque inæquali setis 3 elongatis scabris 2 brevibus membranaceis denticulato-erosis. ① in Mexico prope Gueretaro et Tampico de Tamaulipas. Less. in linnæâ 1831. p. 712. *P. linearis* Willd. herb. ex Less. Achænia pilosiuscula. Pedunculi sesquipollicares. (v. s.)
7. *P. UNIARISTATA*, caule herbaceo erecto tereti glabro oppositè ramosissimo, foliis linearibus mucronato-aristatis cilia utrinquè 3-4 ad basin et glandulas

- pellucidis ad utrumque nervi medii latius 1-serialis gerentibus, pedicellis elongatis bracteolatis 1-cephalis, invol. squamis 5 linearibus acuminate, floribus circ. 10, pappi radii et disci coroniformi hinc in setam uncam aristatam producto. (1?) in Mexico. Herba pedalis elegans tota glabra Cor. flava. Folia 6-10 lin. longa, vix ultra lineam lata. v. s. comm. a cl. Alaman.
8. *P. PYCNIA* (H. B. et Kunth nov. gen. 4. p. 281. caule erecto basi ramoso, ramis teretibus puberulis fastigiatis, foliis lanceolatis-linearibus acuminato-piliferis per totam longitudinem dentibus aristatis ciliatis, capitulis terminalibus subcorymbosis brevissimis pedicelatis inter folia suprema absconditis, capitulis 12-floris, invol. 5-phyllo tubuloso, pappi paleis 3-5 subcartaceis inaequalibus acutis. (1) in calidis prope Honda ad fl. Magdalena. Less. in linnæi 1831. p. 113. P. minuta Willd. herb. ex Less. non Lin. Proxima dicitur *P. serpyllifolia* et *ancipitia*.
9. *P. PERTUSUM*, caule herbaceo erecto di-trichotome ramoso, ramis angulatis, scens angulatis hinc inde subcaberis, foliis linearibus mucronatis subglandulosis superius et margine plus minus scabris basi cilia 2-4 utrinque gerentibus, pedicellis 1-cephalis subbracteolatis involucri paulo longioribus, invol. squamis linearibus acuminatis convolutis, pappi setis 1-serialibus vix basi dilatatis disci 5-6, radii 4-5, (1) in Mexico ad Tancoyuca legit cl. Berlandier. Radix exilis ramosa filiformis. Herba 5-6 pollicaris. Habitus fere *Lorentea Hanceana*, sed pedicellis brevior, caulis fere omnino glaber, involucri acuminate et pappus brevior (v. s.)
- \*\* Capitulis sessilibus aut subsessilibus
10. *P. VIRENS* Less. in linnæi 1831. p. 715. caule ascendente trichotomo ancipiti, hirsutum puberulo, foliis obovato-linearibus mucronatis subrugulosis, angulis scabris basi setigeris, capitulis subsessilibus circ. 12-floris, ligulis brevissimis, invol. squamis 5 obtusissimis, pappi paleis latiusculis in disco 4-5, in caulis 1 lineâ unilateralibus. (1) Patr. non satis nota. An ins. Caribæ? Achenium hirsutiuscula.
11. *P. INSTATATA* Ser. et Merc. mss. in h. DC., prostrata ramosissima, foliis oblongis acuto-mucronatis basi dentibus aristatis utrinque 2-5-ciliatis, capitulis inter folia suprema congesta sessilibus, involucri 5-angulato 5-phyllo, pappi paleis 3-5 membranaceis erectis basi latis apice aristatis subserratis inaequalibus. (1) in insula Cuba legit cl. Ramon de la Sagra. *Lorentea* prostrata Lag. nov. gen. et sp. p. 23 n. 339. v. s.
12. *P. PROSTRATA* (Cav. ic. 4. p. 12. t. 324), prostrata ramosissima, foliis oblongis acuto-mucronatis basi dentibus aristatis 5-7 utrinque ciliatis, capitulis ad ramosum dichotomias et inter folia suprema congesta sessilibus, involucri 5-phyllo ovato-cylindraceo, pappi paleis membranaceis erectis basi latis apice subaristatis in disco 5, in radio 4 unilateralibus. (1) in Mexico, in montibus Sancti Angeli (Moc. ic. ined., ad occid. urbis Guanajuato (Alam.), circa Leon et alibi. Less. in linnæi 1831. p. 714. *Chthonia* prostrata Cass. dict. 9. p. 173. *P. fasciculata* Poir. dict. 5. p. 120. Achenium hirsutiusculum. (v. s.)
13. *P. CILIATA* Lin. sp. 1050. amon. acad. 5. p. 497, erecta dichotome ramosa, caule tetragono sub foliis bifariam puberulo, foliis linearibus amplexicaulibus acutis basi ciliis utrinque 3-5-ciliatis, capitulis intra folia superiora congesta sessilibus, invol. squamis 5, pappi paleis membranaceis lanceolatis acuminatis serratis radii 3 inaequalibus, disci 5. (1) in Santo-Domingo. *P. ciliaris* Swartz: in herb. L'Hér. (sed descr. suæ non congr.) Less. in linnæi 1831. p. 708. An fortè *Chthonia glaucescens* Cass. dict. 9. p. 173? (v. s.)
14. *P. FASCICULIFLORA*, erecta glabra, caule tereti subimplici, foliis lato-linearibus subobtusis, infer. oppositis ciliatis, summis alternis ad ramos floridos congestis pauci-ciliatis, omnibus crebre et minute pellucido-punctatis, capitulis ad apicem et apicem inter folia floralia congestis, invol. oblongis squamis 5 acutiusculis, ligulis parvis oblongis, acheniis 1-2-aristatis. (1) in Mexico legit cl. Hb. Affinis *P. carthasianorum* sed duplo major. Capitula fere ut in *Campomerata* disposita. (v. s. in h. Haenk. a cl. de Sternberg misso.)

15. *P. CARTUSIANORUM* (Less. in *linnæâ* 1831. p. 712), caule erecto teretibus foliis obscure biserialibus pubescentibus, foliis lato-linearibus acutiusculis ciliis 5-9 utrinque ad medium usque instructis minutè et crebrè pelliculo-glandulosis, ramis floridis brevissimis apice corymboso-congestis, capitulis inter folia occultatis sessilibus, invol. squamis 6 oblongis obtusis conduplicatis, flor. circ. 12, acheniis hirsutiusculis, pappi paleis latis in setem acuminatis, radii sæpius 3, disci 5. (1) in Santo-Domingo legit cl. Bertero, qui fortè non immeritò sub nomine *P. ciliaris* comm. Huc enim apprimè convenit icon *Plum. ed. Burm. t. 151. f. 2. à* Linnæo et aliis ad ciliarem relata. (v. s.)
16. *P. SERPYLLIFOLIA* (Less. in *linnæâ* 1831. p. 715), caule suffruticulosò ramosisimo procumbente basi repente, foliis oblongis obovatisve obtusis cilia utrinque 3-4 ad basin conferta gerentibus glandulosis internodio longioribus, capitulis paucis ad apices ramorum sessilibus solitariis, invol. squamis 5 obtusis, floribus 15-30, pappi paleis basi dilatatis in radio 2-3, in disco 5-6, achenio hirsuto. 2/3 in insulis Caribæis (Sw.), Portorico (Wydll.), Sancti-Thomas. (Less.), Chthonia repens Cass. dict. 27. p. 205. *P. humifusa* Wydl. pl. exs. n. 208. Icon *Plum. ed. Burm. t. 95. f. 2. ad* Lorenteam *serpyllifoliam* ab auct. relata huc fortè ex capitulis sessilibus potius pertinet. Valde accedit ad *P. humifusam* Sw. (Lorenteam Less.) sed differt radice perenni, caule basi radicante capitulis sessilibus et pappo disci uniseriali. (v. s.)
17. *P. SIKEREX* (Less. in *linnæâ* 1831. p. 717), caule ascendente basi repente tetragono subdichotomo, foliis oblongo-obovatis muticis internodio sæpius brevioribus basi cilia 4-6 gerentibus, capitulis inter folia subsessilibus sparsis, invol. squamis obtusis, pappo latè palesco disci paleis circ. 8, radic. paucioribus. 2/3 in Martinicâ (Sieb.) et Guadalupâ (Perr.). Pectis prostrata Sieb. fl. Mart. exs. suppl. n. 24. Pappi radii paleæ plùs quàm 2. Cor. intez. Affinis *P. serpyllifoliz*. (v. s.)

† *Species exclusæ.*

- P. ANGUSTIFOLIA* Torr. = *Pectidopsis angustifolia*.  
*P. CANESCENS* Kunth. = *Lorentea canescens*.  
*P. CANESCENS* Less. = *Lorentea auricularis*.  
*P. DISCOIDEA* Horn. = *Leysera capillacea*.  
*P. HUMIFUSA* Sw. = *Lorentea humifusa*.  
*P. MEUFOLIA* Wall. = *Glossocardia Boswellæ*.  
*P. MINUTA* Linn. = *Bellium minutum*.  
*P. PINNATA* Lam. = *Schkuhria abrotanoides*.

VIII. *LORENTEA* Less. in *linnæâ* 1831. p. 717. syn. 153. non *Ort.* (1) nec *Lag.* (2). — *Cryptopetalum* Cass. dict. 27. p. 206. non *Hook.* — *Pectidis* sp. *Linn.*

Corollæ disci obscurè bilabiatae, Pappus disci biserialis inæqualis serrulatus. — Herbæ interdum basi suffruticulosæ. Capitula pedicellata in omnibus, unicâ (*L. sessiliflora*) exceptâ.

Sect. I. *CRYPTOPETALUM* Cass. dict. 12. p. 123. 27. p. 206. non *Hook.*

Pappus radii paleis plurimis acuminato-setiformibus pappo disci brevioribus constans.

1. *L. CRYPTOPETALA*, caule diffuso bifariam piloso, foliis lineari-lanceolatis,

(1) *LORENTEA* Ort. = *Sanvitalia*.

(2) *LORENTEA* Lag. et Cass. dict. 27. p. 202. = *Pectidis* sp. Less. seu *Chthonia* Cass.

capite punctatis dentibus aristatis utrinque 7-8-ciliatis, capitulis terminalibus plerisque pedunculatis, invol. cylindraceo 5-phylo, ligulis brevissimis, pappi paleis inaequalibus pleriseriatis rigidis scabris. (1) in Peruvia ad Guamanga legit Jos. de Jussieu. Cryptopetalum ciliare Cass. l. c. (v. s. in h. Juss.)

2. *L. HÆNKEANA*, caule herbaceo erecto tereti plus minus hirsuto di-trichotomè ramoso, foliis linearibus aut sublancoolato-linearibus mucronatis plus minus utrinque hirsuto-scabris cilia utrinque 2-3 basi gerentibus, pedicellis 1-cephalis bracteolatis capitulo quintuplò longioribus, invol. squamis 5 oblongo-linearibus subobtusis achenia vix superantibus, pappo disci biseriali setis 5 elongatis cæteris brevioribus tenuioribus, radii paucioribus brevioribus. ? in Mexico legit cl. Hænke.

a. *hirsuta*, ramis foliisque hirsuto-scabris, cillis foliorum breviusculis. (v. s. in h. Hænke à cl. Sternberg comm.)

β. *laviuscula*, ramis foliisque glabresculis, setis foliorum longioribus. — Etiam in Mexico legit cl. Hænke. Habitus omninò Pectidis Berlandieri, sed caulis basi durior et verisim. suffruticulosus, pappus et involucria valdè diversa etc. (v. s. cum var. α.)

3. *L. NUMIFUSA* (Less. in linnæ 1831. p. 719), prostrata ramosissima, foliis obovato-oblongis obtusis sparè et minutè glandulosis basi dentibus aristatis utrinque 2-4-ciliatis, capitulis ex axillis foliorum superiorum confertorum brevè pedicellatis, involucriis ovatis 5-phyllis, pappo disci biseriali, paleis majoribus 5 membranaceis basi latè apicè angustè acuminatis serratis disci brevioribus. (1) in Antillarum sterilibus apricis, Guadalupà, Sancta-Cruce. Pectis humifusa Swartz fl. ind. oec. p. 136a. — Plum. ed. Burm. t. 95. f. 2 ? (v. s.)

4. *L. MULTIFLOSCULOSA*, caule suffruticulosò diffuso tereti glabro, foliis lato-linearibus seu lingulatis obtusis mucronatis basi cilia utrinque 3-4 et in paginà inf. glandulas plurimas parvas gerentibus, pedunculis elongatis bracteolatis 1-cephalis, invol. squamis 8 oblongis obtusis, floribus 30-60, acheniis glaberrimis, pappo disci biseriali aristato serrato inaequali, radii 1-seriali breviori. ? in Peruvia montibus Guanacentibus (sic script. in herb. an Guanacacensis ex Leon de Guanaco ?) legit cl. Hænke. Folia 9-12 lin. longa, 2 lin. lata. Invol. 4 lin. longa, floribus breviora. Ligula disco longiores. Pappus etiam disci achenio brevior. Variat ramis foliis pedunculisque plus minus elongatis. Specim. longifolia optimè referunt ic. Banks. rel. honest. t. 19. à Lessingio ad Lorenteam satyroidem relatum, sed in nostrà achenia radii papposa nec calva. (v. s. in h. Hænke à cl. Sternberg comm.)

5. *L. ASSEUTIFLORA* (Less. in linnæ 1831. p. 720), caule herbaceo, ramis ad apicem usquè foliatis bifariam sub foliis puberulis, foliis linearibus mucronatis internodio brevioribus basi ciliigeris pellucido-glandulosis, capitulis inter folia absconditis, invol. 5-phylli squamis obtusis, pappo in radio et disco conformi — Patr. ign.

Sect. II. STAMMARIUM Willd. herb. — Lorentea Less. in linnæ 1830. p. 135.

Pappus radii nullus aut dimidiatus auriculæformis brevis.

6. *L. AURICULARIS*, canescenti-hirtella, caulibus erectis aut procumbentibus ramosis, foliis linearibus acutis basi cillis utrinque 2-3 instructis, capitulis pedicellatis circ. 40-floris, pappo radii auriculæformi. (1) in temperata prope Chilpancingo Mexicanorum ex Less. Pectis canescens Less. in linnæ 1830. p. 135. t. 2. f. 7-12. et 1831. p. 718. non Kunth. Stammarium hyssopifolium Willd. herb. ex Less. (v. s. in h. Hænke à cl. de Sternb. comm.)

β. *Acapulcensis*, caulibus diffusis, foliis latius linearibus, invol. disco dimidio breviorè. ? circa Acapulco. An species propria? (v. s. in h. Hænke.)

7. *L. CANESCENS*, canescenti-subhirtella, caulibus basi suffruticulosis diffuso-

ramosis, foliis linearibus acutis basi ciliis 2-3 utrinque instructis, capitulis pedicellatis 40-floris, pappo radii nullo aut brevi paleaceo setigero.  $\frac{1}{2}$   $\frac{5}{8}$  in Mexico ad Tehuentepec et Chiguitan. Pectis canescens H. B. et Kunth nov. gen. am. 4. p. 263. t. 393. non Less. Cinerarias sp. nov. Alam. herb. Affinis *L. auriculari*, sed pappus radii nunc omninè nullus, nunc setis paucis inaequalibus constans, nunquam nec coroniformis nec auriculiformis. (v. s. comm. à Alaman et Mairet.)

8. *L. SATUREIODES* (Less. in *linnæa* 1830. p. 135. t. 2. f. 7-12. et 1831. p. 718), caulibus suffruticulosis erectis, foliis oblongo-linearibus mucronatis glabris basi ciliis utrinque circiter 8-10 instructis, capitulis multifloris longè pedicellatis, pappo radii nullo.  $\frac{1}{2}$   $\frac{5}{8}$  in collibus arenosis prope Vera-Cruz, Aster satureis Banks in Houst. rel. t. 19. Inula satureioides Mill. dict. Inula Satureis Spreng. syst. 3. p. 520.

9. *L. TENUIFOLIA*, glabra, caule basi suffruticuloso erecto-subdiffuso ramosissimo, foliis linearibus acuminatis grossè glandulosis cilia utrinque 3-4 basi gerentibus, capitulis longè pedicellatis circ. 10-floris, pappo radii nullo.  $\frac{5}{8}$   $\frac{1}{2}$  in insula Albemarle à numero archip. Gallapagos sub æquatore non longè à littore Peruviano jacentia, legit cl. Macrae. Rami tenues teretes. Fol. 8-10 lin. longa, semilineam lata. Involucra oblongo-cylindracea. Achænia elongata, radii glabra, disci puberula. (v. s. comm. ab hon. Soc. hortie. Londin.)

Trib. II. *EUPATORIACEÆ* Less. syn. 154. — Eupatoriæ, Adenostyleæ et Tussilagineæ Cass. tabl. XIX et XX.

Stylus fl. hermaphr. supernè cylindraceus, ramis longis teretibus seu clavatis exsertis extus supernè papillosis rariùs puberulis, fl. masc. integer aut breviter bifidus clavulatus. Stigmatum series angustæ parùm prominulæ antè medios styli ramos desinentes, rariùs apicem attingentes, rarissimè confluentes. Corolla staminigera regularis 5-dentata, dentibus sæpiùs erectis. Anthæræ ecaudatæ. Pollen globosum echinulatum. — Capitula pauci- aut multiflora nunquam 1-flora, homogama fl. tubulosis, rariùs heterogama fl. radii tubulosi ligulatis. Flores (paucissimis exceptis) cyanici. Herbæ aut frutices. Folia sæpiùs opposita. — Tribus accedens hinc ad Vernoniaceas et ad Senecionideas oppositifolias.

## CONSPECTUS TRIBUS.

Subtrib. I. *EUPATORIÆ*. Capitula homogama. (Gen. LIX-LXXXIX.)

Div. I. *ALOMIÆ*. Pappus nullus. (Gen. LIX-LXIV.)

Div. II. *AGERATÆ*. Pappus paleaceus aut partim squamellatus. (Gen. LXV-LXXIII.)

Div. III. *ADENOSTYLÆ*. Pappus setosus scaber aut plumosus. (Gen. LXXIV-LXXXIX.)

Subtrib. II. *TUSSILAGINÆ*. Capitula heterogama aut dioica. (Gen. XC-XCVII.)

Div. I. *PETASITÆ*. Flores fœminei tubulosi. (Gen. XC-XCIII.)

Div. II. *EOTUSSILAGINÆ*. Flores fœminei ligulati. (Gen. XCIV-XCVII.)

Subtrib. I. *EUPATORIÆ*. — Eupatoriæ et Adenostyles Cass. —  
Alomiæ Ageratæ, Eupatoriæ Less.

Capitula homogama. — Flores nunquam stricto sensu flavi.

Div. I. *ALOMIÆ* Less. *syn.* 154.

Pappus nullus. — Divisio nimis artificialis ex affinitate Alepidis cum Agerato, Astemmatidis cum Adenostemmate et Orsinis cum Eupatorio, imò Isocarpis cum Carphophoro.

LIX. *ORSINIA* Bertol. in *lit.* 1834. non in *ann. st. nat.* — Eupatorii sp. Balb.

Capitulum 6-7-florum. Recept. nudum. Invol. ovati squamæ 6-7 ovals erectæ æquales. Cor. 3-5-fidæ, hirtellæ, lobis ciliatis. Styli rami exserti muriculati obtusi. Antheræ incluse. Achænium obcompressum apice villosum. Pappus nullus. — Suffrutex aut Herba suffruticans Brasiliensis. Caulis teres pubescenti-velutinus apice ramoso-corymbosus. Folia opposita et alterna breviter petiolata ovata rigida acutiuscula acutè serrata penninervia reticulata suprâ scabrida subhis secus nervos nervulosque densè hirta. Corymbi terminales, ramis hirsutis. Capitula subsessilia. Invol. squamæ pallidæ dorso puberæ latè ovales subconcaavæ. — Cum primum genus cl. Orsini à cl. Bertoloni dictum in Jasoniam ceciderit hoc novum consecravit cl. auctor et mecum caract. ex specim. authentico desumptos communicavit.

I. *O. EUPATORIA*. § in Brasiliæ prov. Minarum Generalium ad Mariannam legit cl. Vauthier. Eupatorium Armani Balb. pl. rar. h. taur. 1810. p. 27. t. 6. *des* loci indie. Flores albi. Folia a poll. longa, 15 lin. lata, petiolo 3 lin. (v. s. comm. à cl. Vauthier sub n. 263.)

LX. *PIQUERIA* Cav. ic. 3. p. 19. t. 235. Willd. sp. p. 1748. Cass. dict. 41. p. 115. Less. *syn.* 154. — Flaveria sp. Juss.

Capitulum 4-5-florum. Invol. 4-5-phyllum 1-seriale oblongum. Recept. nudum planum parvum. Cor. tubus brevis sub dilatatus sæpè hirsutus. Antherarum appendix terminalis subnulla. Styli rami obtusi. Achænia juniora compressa adulta 5-angulata glabra supiti brevi articulata. — Herbæ aut suffrutices Americanæ. Folia opposita sæpius trinervia. Capitula corymbosa aut paniculata parva. Corollæ albæ etiãsi ex siccis flavas dixerit Cassini, qui sæpissimè album videbat flavum.

#### Sect. I. *EUPUIQUERIA*.

Capitula 4-flora. Cor. tubus hirsuto-barbatus. Invol. squamæ obtusæ mucronatæ. — Caulis herbacei.

I. *P. TRINERVIA* (Cav. ic. 3. p. 19. t. 235), herbacea glabra, foliis ovato- aut oblongo-lanceolatis subserratis trinerviis, paniculâ corymbosâ laxâ polycéphalâ. 2/ in Mexico locis rupestribus prope S. Rosa de la Sierra (H. B.), Jalapam (Schied. et Depp.), etc. in omnibus Mexici circuitibus ubi *Xoromiztli* et *Yoloxitlic* nuncupatur (Moc. fl. mex.) Ageratum febrifugum Sess. fl. mex. ined. *Febrifuga* Moc. fl. mex. ic. ined. — Hern. mex. p. 353. fl. 1. ex Moc. et à p. 437. f. 2. ex DC. — *P. trinervia* Jacq. f. eclog. 1. p. 70. t. 48. Variat ræ omnino glabris ent sub foliorum origine bifariam pilosis. (v. v. c. et a. sp.) *pyroea* (H. B. et Kunth nov. gen. sm. 4. p. 153), herbacea piloso-glabra, foliis latè ovatis aut ovato-oblongis trinerviis grossè serratis, paniculâ laxâ. 2/ in temperatis Mexici inter Capula et Pazcuaro (H. B.) 3/ gill. longa, 1 1/2 fere lata. (v. s. comm. à cl. Alaman.)

## Sect. II. PIQUERIOIDES.

Capitula 5-flora. Invol. squamæ striatæ in appendicem subulatam desinentes. — Caulis fruticosi. Folia triloba.

3. *P. PUNESCENS* (Smith in Rees cycl. 27. n. 2), fruticosa, caule pube densâ brevi velutinâ, foliis trilobis acutis glabrescentibus, paniculâ densâ corymbosâ villosâ, capitula 5-floris. ♀ in Peruvâ ex herb. Linn. fil.
4. *P. QUINQUEFLORA* (Cass. dict. 4t. p. 116), fruticosa, caule basi glabro apice rufo-tomentoso, foliis trilobis glabrescentibus pellucido-glandulosis, lobis acutis dentatis, paniculâ densâ corymbosâ subviscosâ, capitulis 5-floris. ♀ in Peruvâ legit Dombey. Corollæ à Cassinio dicuntur flavæ, sed frequentissimè ex sicco flores albos dixit flavos. An certè à priorè diversa?

## Sect. III. ARTEMISIOIDES.

Capitula 3-4-flora. Cor. glabræ, tubo longiusculo basi suprâ ovarium subdistans. Invol. squamæ acuminatæ aut obtusæ. — Caulis fruticosi glabri apice summo pubescentes aut glutinosi. Folia glabra petiolata. — An genus proprium?

5. *P. FLORIBUNDA*, fruticosa glaberrima apice glutinosa, foliis oppositis petiolatis ovato seu rhombico-lanceolatis grossè dentatis, paniculæ ramosissimæ ramis corymbiferis, capitulis 4-5-floris, invol. squamis acuminatis. ♀ in Peruvâ montanis Oronocensibus (et fortè in Chili) legit cl. Hænke. Inv. 3 lin. longa. Folia ferè Sedi populifolii sed membranacea. Specimina Chilensia in herb. Hænkeano (fortè schedulæ transpositione) dicta Peruvianis simillima in ditissimis Chilensibus herbariis nunquàm vidi et mihi quoad patriam valdè dubia! (v. s. in herb. Hænke à cl. Sternb. comm.)
6. *P. ARTEMISIOIDES* (H. B. et Kunth nov. gen. am. 4. p. 153), glabrescens, foliis caulinis oppositis petiolatis lanceolatis acuminatis basi cuneatis grossè dentatis summis lineari-lanceolatis integerrimis sparsis, invol. squamis obtusis. ♀ in Peruvâ (Domb.?), in frondosis Quitensium (H. et B.), in vallibus Cordillieris (Hænke). Flaveria Peruviana Juss. gen. 187. Flaveria spicata Smith in Rees cycl. v. 14. n. 2. Herba sicca subflavicans. Rami et folia novella subpubescentia, cæteræ partes glabræ. (v. s.)
7. *P. CALIODES*, fruticosa glaberrima, foliis linearibus integerrimis subnerviis, paniculæ ramosissimæ ramis racemiferis, capitulis 3-floris, involucri squamis subobtusis. ♀ in Peruvâ vallibus Cordillieris legit cl. Hænke. Folia 8-10 lin. longa, vix lineam lata. Invol. 2 lin. longa. (v. s. in h. Hænke à cl. Sternb. comm.)

LXI. *ALOMIA* H. B. et Kunth nov. gen. am. 4. p. 151 et 312. t. 354.  
Less. syn. 154.

Capitulum multiflorum. Invol. campanulatum imbricatum, squamis angustis acutis. Recept. convexum nudum. Cor. tubo gracili glabro. Antheræ appendicibus ovato-oblongis obtusis terminatæ. Styli lobi elongati obtusi. Achænia 5?-angularia apice calva. — Herba Mexicana ramosissima glabriuscula albiflora. Piquerias (nec Ethulias) characteribus et habitu omninò referens.

1. *A. AGERATOIDES* (H. B. et K. l. c.), in Mexicis calidis prope Mescala. *Ethulia ageratoides* Spreng. syst. 3. p. 458.

## LXII. PHALACRÆA.

Capitulum multiflorum. Invol. squamæ subbiseriales æquales 3-nerviæ. Recept. nudum. Corol. tubus basi piloso-hispidus, faux ampliata, limbus 5-fidus. Antheræ vix apice appendiculatæ. Styli rami exserti



apice clavati. Achænium compresso-angulatum calvum. — Herba Peruviana breviter puberula fortè subviscida. Folia opposita petiolata latè ovata basi truncata et ad petiolum subcuneata grossè et interdum duplicato-dentata. Pedunculi nudiusculi apice corymbosi. Capitula pedicellata. Flores purpurascentes. — Genus ex pappo nullo affine Alomiæ sed diversum cor. tubo hispido, invol. non imbricato, etc. Aff. etiam Astemmati, sed differt cor. fauce ampliata nec cylindracea, tubo hispido nec glabro. — Nomen ex φαλλος, calvus.

1. *P. LATIFOLIA*. ♀ in Peruvia circa Limam legit cl. Néé. *Stevia* seu *Ageratum* Néé! herb. *Ageratum latifolium* Cav. ic. 4. p. 33. t. 35? Planta mea (etiam in herb. Hækeano obvia) ut Cavanillesiana à Neesano itinere proveniens apprimè, ex habitu toto, iconi et desc. respondet, sed Cav. vidit pappum 5-paleaceum et in meâ pappus nullus. An diversa? An à Cavan. malè descr.? Ex Cav. vulgò *Teatina* nuncupatur. Petioli 6 lin. longi pedunculique caulesque pube brevi velutini. Limbus 2 poll. ferè longus æsequipollis. latus. (v. s.)

♀. *glabra*, caule glabro, pedunculis vix apice puberulis, petiolis 15 lin. longis, fol. limbo 3 poll. longo, 2 poll. lato, capitulis paulò majoribus. ♀ Etiam habeo ex itin. Neesano sed locum propr. ignoro. (v. s.)

#### LXIII. GYMNOCORONIS.

Capitulum multiflorum homogamum. Invol. squamæ 1-seriales acuminatæ subnerviæ. Recept. nudum. Corolla tubulosa glabra, fauce non dilatata, lobis brevibus conniventibus. Styli rami exserti clavati. Achænium compressum dorso nervoso-subangulatum calvum. — Herbæ Brasilienses erectæ glabræ. Caules infernè sulcati. Folia opposita petiolata oblongo-lanceolata acuminata dentata. Pedunculi corymbosi. Capitula pedicellata subpuberula. Flores albi. — Genus valdè *Adenostemmati* affine sed achænia calva! Nomen ex γυμνός nudus, et κορυφή, apex, terminus. An stirpes sequentes sint species aut varietates dubius hæreo.

1. *G. ATTENUATA*, foliis oblongato-lanceolatis basi cuneatis secus petiolum decurrentibus. ♀ in Brasiliâ prov. Rio-Grande. (v. s. in b. Mus. reg. Par. à Mus. imp. Bras. sub n. 782 miss.)
2. *G. SUBCORDATA*, foliis elongato-lanceolatis basi truncatis subcordatis in petiolum non decurrentibus. ♀ in Bras. prov. Rio-Grande. (v. s. cum priori sub n. 886 miss.)

LXIV. ISOCARPHA *R. Brown. obs. comp. 110 excl. syn. Less. in Linnæâ 1830. p. 141. 1831. p. 405. syn. p. 154.*

Capitulum multiflorum. Invol. squamis circ. biseriatis conduplicatis. Recept. conicum, paleis acuminatis squamas referentibus. Corolla tubulosa 5-dentata. Styli rami elongati extus convexi hispiduli acuti. Achænium prismaticum epapposum. — Herbæ Americæ æquinoctialis erectæ oppositifoliæ. Folia integra rariùs dentata. Capitula terminalia.

1. *I. ATRIPLICIFOLIA* (BROWN. l. c.), erecta glabra, caule tereti, foliis oppositis auriculato-connatis aut alternis stipulato-auriculatis in petiolum alatum basi attenuatis supernè deltoideo-ovatis acuminatis grossè et obtusè dentatis triplinerviis membranaceis, capitulis paucis pedicellatis subpaniculatis. — in Amer. merid. (Mill.) ad flumen Magdalensæ (Bertero!), *Calidemos atriplicifolius* Spreng.! syst. 3. p. 457. *Bideus atriplicifolia* Linn. sp. 2. p. 1168. *Spilanthus atriplicifolia* Linn. syst. veg. 610. Ad aliam stirpem alludere videtur Necker cum recept. nudum dixit, achænia angulata nulla, juniora interdum setas 1-2-gerentia. (v. s.)

2. *I. OPPOSITIFOLIA* (R. Brown. l. c.), caule dichotomo, ramis longissimis erectis tripartitis simplicibusque, foliis lanceolatis, capitulis subternis sessilibus. — in *Jamaica montibus*. *Calea oppositifolia* Linn. sp. 1179. amom. acad. 5. p. 404. Swartz obs. 302. — *Santolina Brown. jam.* 315. Gærtner fr. 2. p. 391. Invol. squamis binerviis laxæ muticæ rectilineo-acuminatæ ex Less.
3. *I. BILLBERGIANA* (Less. in Linnæ 1831. p. 405. in adn.), ramis flexuosis, foliis sessilibus penninerviis basi auriculatis spathulato-rhomboidis, squamis invol. penninerviis, achæniis pubescentibus. — in *Tierra Bomba* prope *Carthagenam*.
4. *I. ECHIOMES* (Less. in Linnæ 1830. p. 141. t. 2. f. 14-16), caule infernè dense foliato superne nudo, foliis sessilibus lineari-lanceolatis triplinerviis scabris et hispidis, paniculâ corymbosâ confertâ, corollis apice pilosiusculis, invol. squamis linearibus canaliculatis ciliato-hirsutis, achæniis glaberrimis. ♀? in *graminosis Mexici* prope *Hacienda de la Laguna*. — Paniculæ rami foliis suis breviores. Invol. floribus æquale, squamis ciliatis, inter. latib. scariosis.

† *Species exclusa.*

*I. BENTHAM* Balb. = *Crinia Dalea*.

Div. II. AGERATEÆ Less. syn. 155.

Pappus paleaceus, squamellis membranaceis rigidisve distinctis aut in coronam plus minus concretis.

LXV. COELESTINA Cass. dict. 6. suppl. p. 8. et v. 26. p. 227. Kunth nov. gen. am. 4. p. 150. Less. syn. 155. — *Agerati* sp. auct.

Capitulum multiflorum. Invol. cylindraceo-hemisphæricum polyphyllum subimbricatum. Recept. convexum paleaceum aut nudum. Achænia pentagona. Pappus coroniformis membranaceus dentatus, paleis nempe concretis apice subinæqualibus. — Herbæ annuæ erectæ ramosæ. Caulis teretes. Folia opposita breviter petiolata dentata. Capitula corymbosa approximata pedicellata. Corollæ cæruleæ aut purpureæ. — Genus vix ab *Agerato* distinctum et cum eo fortè (eodem jure ac sectiones *Stevie*) coadunandum?

Sect. I. ISOCARPHOIDES.

Recept. paleis linearibus onustum.

1. *C. ISOCARPHOIDES*, caule suffruticoso tereti hispidulo scabro, foliis lineari-lanceolatis acuminatis subsessilibus triplinerviis integerrimis supra scabris subtus velutino-villosis, invol. squamis lineari-subulatis scabridis, capitulis circ. 25-floris, pappo brevi truncato. ♂ in Mexico legit cl. Henke. Proximè accedit ad *Isocarpham echinoidem* ex descr. Less., sed habitus ab *Hieracio echinoidi* diversus. Caulis per ferè totam longitudinem foliosus. Pappus brevis equidem non verò nullus. (v. s. in h. Hænk. à cl. Sternb. comm.)

♂. *dentata*, foliis majoribus hinc inde dentatis. (v. s. cum var. a.)

2. *C. PALMACHA* (Gay! in Andr. pl. mex. exs. n. 287), caule herbaceo erecto tereti molliter ac brevissimè velutino, foliis petiolulatis ovatis seu ovato-lanceolatis triplinerviis subdentatis supra scabridis subtus velutino-villosis, invol. squamis lineari-subulatis scabridis, capitulis 25-floris, pappo coroniformi dentato hinc inde in setam expanso. ♀? in Mexico circa *Oaxacam* legit cl. Andrieux mense julio. Flores violacei. (v. s.)
3. *C. ALBIDA*, caule herbaceo erecto puberulo, foliis ovalibus utrinquè attenuatis vix dentatis quintuplinerviis supra hispidulo-scabris subtus tomentoso-hirsutis, invol. squamis glabrisculis linearibus acutis, capitulis circ. 60-floris, pappo coroniformi dentato. ♀? in Mexico inter *Oaxacam* et *Mitla* julio ineunte legit cl. G. Andrieux. Cor. albidæ. (v. s. comm. à cl. inv. snb n. 548.)

## Sect. II. AGERATOIDES.

## Receptaculum epaleaceum.

4. *C. AGERATOIDES* (H. B. et Kunth nov. gen. am. 4. p. 151), caule pulverulento scabrido, foliis ovato-oblongis acuminatis utrinque scabris, corollis tubo extus pilifero, pappo cupuliformi obtusè dentato. ① in Mexici montanis ut videtur sat. frequens. *Ageratum celestinum* Sims bot. Mag. t. 1730. *Sparganophorus ageratoides* Lag. elench. h. madr. 1815. p. 25. ex Spreng. *Ageratum corymbosum* Balb. hort. taur. 1806 et 1808. Pers. ench. n. 4. (v. a. sp.)

*β. latifolia*, foliis latè ovatis. — in Mexico. (v. a.)

5. *C. CORYMBOSA*, caule glabriusculo, foliis ovatis acuminatis utrinque subscabridis, corollæ tubo extus glandulifero, pappo cupuliformi inæqualiter dentato, dentibus 2 longioribus acutis. — in Mexico? *Ageratum corymbosum* Zacc. obs. n. 85. *Celestina cærulea* Cass. dict. 6. suppl. p. 9. An prioris varietas? (v. v. c. in hort. Florent. à Zuccagnii tempore seminibus servatam.)

6. *C. PARVIFOLIA*, caule vix puberulo, foliis ovatis crenato-dentatis brevissimè petiolatis, pedunculis racemoso-corymbosis, corollæ tubo glabro, pappo coroniformi obtusè dentato. — in Amer. talid. (fortè circa Limam) legit el. Née. Petioli lineam vix longi. Limbus folii 6 lin. longus, 4. lin. latus. (v. a. ex itin. Née.)

LXVI. *AGERATUM* Linn. gen. n. 936. *Gærtn. fr. 2. p. 398. t. 165.* — *Carollæ Pont. diss. Adans. fam. 123. non Less.*

Capitulum multiflorum subglobosum. Invol. polyphyllum imbricatum, squamis linearibus acuminatis. Recept. nudum. Cor. 5-fida. Styli rami elongati cylindracei obtusiusculi. Achænium subpentagonum basi attenuatum. Pappus paleaceus, paleis 5-10 liberis aristato-acuminatis aut pectinatis obtusis. — Herbes pleræque annuæ erectæ Americane aut rarius Indicæ. Folia opposita petiolata ovata dentata. Capitula corymbosa. Flores cærulei aut albi. — Sectiones non magis inter se differre videntur quam sectio *Stevia*.

Sect. I. EUAGERATUM. — *Ageratum* Cass. dict. Less. syn. 155.

Pappi paleæ 5 ad basin distinctæ subscerratæ apice acuminatæ aristatæ.

1. *A. CORYNTOIDES* (Linn. sp. 1175), caule ramoso, foliis ovatis rhombeis cordatisve longiusculè petiolatis, pappi paleis basi dilatatis serrulatis apice longè aristatis corollam subsequantibus. ① frequens in America et Africa æquinoctiali, in Indiâ orientali, insulis Mauritanis (introd.?) et Maderæ secus vias (Low!), in ins. Sandwich Owyhee (Macrae!). — Schkuhr handb. t. 238. Hook. exot. flor. t. 15. Herba valdè varians, caule nunc hispido nunc ferè glabro, foliis nunc ovatis nunc obcordatis, acutis aut obtusis, plus minus dentatis pilosis, petiolis hispidissimis aut glabriusculis, corollis cæruleis (*Ag. cæruleum* Desf. cat. h. par.) aut albis (*A. album* Steud. nom. ph. p. 13), undè plurimæ species ab auctoribus admixtæ hic ut varietates indicatæ, vix inter se nunc satis distinctæ sed tamen recognoscendæ.

a. *obtusifolium*, caule glabriusculo, foliis ovatis obtusis, floribus pallidè cæruleis. *A. obtusifolium* Lam. dict. 1. p. 54. ill. t. 672 f. 1. — *Sparganophorus obtusifolius* Lag. elench. h. madr. 1815. p. 25? (v. a.)

β. *hirtum*, caule hirtio, foliis ovatis acuminatis. *A. hirtum* Lam. l. c. Poir. suppl. 1. p. 241. — Lam. ill. t. 672. f. 2. (v. v. c.)

γ. *Mexicanum*, caule hirtello, foliis ovatis subobtusis, floribus cæruleis. *A. Mexicanum*, bot. mag. t. 2524. Sweet br. fl. gard. t. 89. (v. a.)

δ. *cordifolium*, caule hispido, petiolis hispidissimis, foliis ovatis acuminatis. *A. cordifolium* Roxb. hort. calc. Wall. herb. et list, n. 237. (v. a.)

## Sect. II. PECTINELLUM.

Pappi paleæ circ. 10 distinctæ per totam ferè longitudinem pectinato-ciliatæ.

2. *A. MELISSA-FOLIUM*, caule ramoso glabro, foliis petiolatis glabresculis ovatis crenato-dentatis truncatis aut subcordatis apice subacutis, pedunculis corymbosis, corollâ glabrâ brevi, pappi paleis oblongis pectinato-ciliatis subacutis. — in Brasiliâ circa Bahiam leg. cl. Salzmann (n. 431) et Blanchet (n. 136 et 1974 !). An *Chrysocoma pauciflora* Arrab. fl. Rum. v. 8, t. 3? Petioli 6-7 lin. longi; limbi fol. pollicem longi, 6 lin. lati. Achænia angulata basi attenuata glabra. (v. s.)
3. *A. MATRICARIOIDES* (Less. syn. 155), caule ramoso, foliis petiolatis rhombeo-ovatis obtusis crenatis glabresculis, pedunculis corymbosis, pappi paleis lanceolatis incisio-laceris pubescentibus. — in insulâ Cubâ. *Hymenopappus matricarioides* Spreng. syst. 3. p. 450. *Anthodium patens* (Spr.) Corolla Piquerie (Less.). Cæc. ign.
4. *A. DOMINGENSIS* (Spreng. syst. 3. p. 446), caule gracillimo vix puberulo, foliis petiolatis orbiculatis cordatis puberulis crenato-dentatis, pedunculis 1-cephalis solitariis, pappi paleis oblongis serratis breviter acuminatis corollâ dimidio brevioribus. ① in Santo-Domingo legit cl. Bertero! Herbula fortè decumbens. Petioli 4 lin. longi. Folia 4 lin. diam. Invol. æquamæ paucæ. (v. s.)

† *Species dubia.*

5. *A. CULARE* (Linn. sp. 1175), caule glabro, foliis petiolatis ovatis crenatis obtusis. — in Ind. orient. prope Bisnagas. Pluk. alm. t. 81. f. 4. Spec. omninò obscura †.
6. *A. ? MARITIMUM* (H. B. et Kunth nov. gen. am. 4. p. 150), caule herbaceo ramoso glabresculo, foliis petiolatis ovatis leviter cordatis obtusiusculis crenatis glabris, capitulis corymbosis pedicellatis, circ. 70-floris, corollâ glabrâ, pappi paleis crebris subulatis. ① in litore insulæ Cubæ prope Havanam. Corollæ violacæ. An Eupatorii species? An genus proprium?

† *Species exclusæ.*

- A. ANGUSTIFOLIUM* Sp. = *Eriophyllum angustifolium*.
- A. ARBUTIFOLIUM* Kunth. = *Phania arbutifolia*.
- A. COELESTINUM* Sims. = *Cœlestina ageratoides*.
- A. CORYMBOSUM* Balb. = *Cœlestina ageratoides*.
- A. CORYMBOSUM* Zucc. = *Cœlestina corymbosa*.
- A. FERRUGINEUM* fl. mex. = *Piqueria trinervia*.
- A. GUIANENSIS* Aubl. = *Habeclinium macrophyllum*.
- A. LATIFOLIUM* Cav. = *Phalacraea latifolia*.
- A. LINEARE* Cav. = *Palasfoxia linearis*.
- A. PEDATUM* Ort. = *Stevia pedata*.
- A. PUNCTATUM* Jacq. = *Stevia serrata*.
- A. PUNCTATUM* Ort. = *Stevia Eupatoria*.
- A. VINOSUM* Cav. = *Stevia salicifolia*.

## LXVII. ANISOCHÆTA.

Capitulum multiflorum (fortè dioicum?). Invol. ovatum imbricatum, squamis lineari-lanceolatis adpressis disco brevioribus. Recept. nudum. Cor. tubulosæ profundè 5-fidæ glabræ exter. ad peripheriam dejectæ. Antheræ ecaudatæ. Styl. in plerisque inclusus subabortivus, in paucis ramis exsertis cylindraceis subclavatis vix papillois. Achænia teretiusscula vix subpuberula erostris (imatura aut abortiva?)

Pappus paleis sæpius 3 rarius 4-5 erectis squamellois angustis acuminatissimis inæqualibus. — Frutex Australi-Africanus subscandens. Caulis teres striatus pubescens, ramis patulis. Folia alterna petiolata latè ovata basi subtruncata 3-5-nervia grossè inciso-dentata glabra, petiolo nervisque pubescentibus. Paniculæ ramos terminantes subnudæ laxæ. Bracteolæ minimæ secus pedicellos. Cor. ex sicco albide.

1. A. MIKAWOIDEA.  $\S$  in Africâ austr.-orient. ad Zw. Omsamcaba et Omsamwak legit cl. Drege! Petiolus semipollic. Limbus folii a poll. longus et latus. (v. s. comm. à cl. inv.)

LXVIII. ADENOSTEMMA Forst. nov. gen. (1776) p. et t. 45. Cass. dict. 25 p. 360. Less. syn. 156. — Lavenia Swartz Prod. (1788) 112. fl. ind. occ. 139 — Verbesinæ et Cotulæ sp. Linn.

Capitulum multiflorum homogamum. Recept. planum nudum foreolatatum. Invol. campanulatum floribus ferè brevius post anthesin reflexum, squamis 1-seriatis foliaceis oblongis. Corolla tubulosa 5-dentata infra lobos subvillosa, fauce vix ampliata. Stigmata longè exserta apice dilatata colorata. Achenium obovato-oblongum subangulatum superatum, aristis 3-5 brevibus rigidis apice glanduloso-pilosæ seu glabriusculæ. Folia opposita petiolo ovata seu rhomboidea triplinervia dentata. Capitula pedunculata corymboso-paniculata. Flores albi. — Nomen Swartzii à Rajo mutuatum erat ritè conditum (planta enim erat Verbesina Lavenia Linn.), sed nomen Forteri vetustius et characterem indicat. — Species inter se valdè affines examen ulterius requirunt.

\* *Caule basi radicante adscendente.*

1. A. SWARTZII (Cass. dict. 25, p. 362), caule basi radicante adscendente simplici apice minutissimè puberulo, foliis petiolatis ovato-subcordatis basi cuneatis grosse dentato-crenatis subacutis glabris, pedunculis 3-5 corymbosis elongatis, invol. squamis obtusis dorso glabris apice ciliolatis, acheniis distinctè stipitatis scabridis. (I) in subhumidis sylvaticis umbrosisque Jamaicæ. Lavenia decumbens Swartz! fl. ind. occ. 1331. Cotula Verbesina Linn. an. ac. 5, p. 407. — Sloan. jam. 126. p. 262. t. 155. f. 2. Caulis non decumbens ut vult Swartz sed adscendens. Cor. 15-20 albide. Achenii puncta parva albis pulveris moleculam referunt. Pappi setæ 3. (v. s.)
2. A. BERTERII, caule basi adscendente glanduloso-puberulo, foliis petiolatis ovato-rhomboidis basi cuneatis apice acutis hinc inde dentato-serratis glabris, corymbo 5-7-cephalo, involucri squamis ovalibus dorso puberulis subacutis, acheniis distinctè stipitatis muricatis. (I) in Santo-Domingo legit amic. Bertr. Lavenia decumbens ex Hispaniolâ Spreng. syst. 3, p. 445. A. Swartzii similis, sed differt notis iudicatis, consist. foliorum magis papyraceâ, capitulis duplè ferè majoribus, acheniis multò magis muricatis, setis pappi ferè ubique 3 nec 3. (v. s. comm. à cl. Balbis.)
3. A. PERROTETII, caule basi radicante adscendente glanduloso-subpuberulo, foliis petiolatis ovatis crenatis obtusis basi cuneatis glabris, paniculâ polycephalâ laxâ, invol. squamis oblongis acutiusculis dorso puberulis, acheniis distinctè stipitatis muricatis. (I) ? in paludibus turfosis peninsulæ Promontorii Viridis Africanæ legit cl. Perrottet. Similis A. Swartzii, sed folia rigidiora et tusalora. Capitula 11-15. Achenia punctis glandulosis rubris muricata. Caulis rubescentes. (v. s. comm. à cl. inv.)
4. A. MAURITIANUM, caule basi radicante adscendente simplici vix apice puberulo, foliis petiolatis ovato-lanceolatis basi cuneatis grosse dentatis glabris, pedunculis elongatis 3-7 in corymbum dispositis, invol. squamis obtusis

glabris, acheniis lævibus. — in insulæ Mauritiæ sylvia montosis (Comm. , Sieb. Bontou! etc.). Ad. viscosa var. Commersonii Cass. dict. 25. p. 362. *Lavenia erecta* Sieber! fl. maur. exa. 2. n. 119 et 364. Non malè refert t. 42. Surin. Zeyl. Folia papyracea cum petiolo 3-5 pollic. longa. (v. s.)

1. *A. DARGH*, caule basi radicante ascendenti-erecto glabro vix hinc inde asperulo, foliis petiolatis ovatis basi longè cuneato-attenuatis apice acutis dentatis membranaceis suprâ pube minimâ sparsâ puberulis subtus nervos glanduloso-scabridis, corymbo laxo, pedicellis elongatis, invol. squamis elliptico-oblongis pubescentibus, acheniis valdè muricatis. — in Africa australi ad Zw. Omsamwubo et Omsameulo legit cl. Drege! Habitus *A. Mauritanii*, sed acheniis muricatis et aliis notis facilè differt. (v. s.)

6. *A. MICROCEPHALUM*, caule basi radicante ascendente minutissimè glanduloso-pubero, foliis breviter petiolatis ovatis denticulatis obtusis glabris, paniculæ laxiusculæ capitalis parvis, invol. squamis oblongis subacutis glabriusculis, acheniis vix muricatis. — in Indiâ orient. ad Gualpara et Silhet legit cl. Hamilton. *Lavenia viscida* Ham. ex Wall. ! cat. n. 332. Affine *A. Perrottetii*, sed capitulis dimidio minoribus sanè primâ fronte diversum. (v. s. comm. ab ampl. Cur. merc. angl. Ind. or.)

7. *A. TINCTORIUM* (Cass. dict. 25. p. 364), caule subprocumbente diffuso, radice repente, foliis oppositis lanceolatis inæqualiter serratis glabris euerviis, invol. squamis obtusis. — in Cochinchinâ et Chinâ ad colorem cœruleum foliis coarctatis eliciendum culta et *cham-lonla* nuncupata. *Spilanthes tinctorius* Lour. coch. ed. Willd. 2. p. 590. Species ex pappi aristis 3-capitatis ad Adenostemma ritè videtur relata, sed fruct. non descr.

\* \* *Caule erecto, acheniis muricatis.*

4. *A. GLUTINOSUM*, caule erecto ramoso villosa-glutinosa, foliis ovatis acutis remotè dentato-serratis ciliatis basi cuneatis ntrinque glutinosa-hirtellis, corymbo paniculato laxo, pedicellis capitulo triplo longioribus, acheniis papuloso-muricatis. — in insulis Sandwich ad alt. 450-500. hexap. *Lavenia glutinosa* Gaudich. ! bot. voy. Freyc. p. 471. Ach. basi attenuata sessilia glabra, sed evidentissimè tuberculis minimis muricata. (v. s. comm. à cl. invent.)

9. *A. VISCOSUM* (Forst. ! nov. gen. n. 15), caule erecto pubescente aut glabro, foliis petiolatis oblongo-lanceolatis basi cuneatis grossè serratis, capitulis laxè paniculatis, pedicellis capitulo vix duplo longioribus, acheniis papuloso-scabris. ☉ ? in ins. Moluccanis, Rawak (Cham. ! ) Timor (Gaudich. ! ), et in ins. Societatis (Forst.) *Lavenia erecta* Gaudich. in bot. voy. Freyc. 470. excl. syn. — Putumba Rheed mal. 10. t. 63 in arenosis Malabaræ crescens, huc ex forma foliorum fortè referenda sed capitulis majoribus rufescentibus differt. (v. s.)

12. *A. PARVIFLORUM* (Blum. bijdr. p. 906. sub *Laveniâ*), caule erecto simplici glabriusculo, foliis petiolatis rhombéo-ellipticis obtusiusculis basi cuneatis repando-obtusè-serratis, corymbo terminali divaricato, ramis puberulis, invol. squamis dorso puberis, stigmatibus vix exsertis, acheniis muricato-tuberculatis, ☉ in umbrosis ins. Javæ circa Buitenzorg. (v. s. in h. Blum.)

a. *divaricatum*, paniculâ laxiore valdè divaricatâ, invol. minis puberulis. — in ins. Javâ. *Lavenia erecta* Blum. ! bijdr. 906. non alior. (v. s. comm. à cl. Blum.)

11. *A. FASTIGIATUM* (Blum. ! bijdr. p. 905, sub *Laveniâ*), caule erecto subramoso glabro, foliis in petiolum attenuatis ovato-lanceolatis obtusè acuminatis grossè et inæqualiter dentato-serratis glabris, paniculâ terminali fastigiatâ, ramis involucriisque glanduloso-pilosiusculis, acheniis pube minimâ et tuberculis densè scabridis. — in ruderatis et humidis ins. Javæ (Blum. ! ), ins. Marianarum (Nanke ! ), in Indiâ orient. ad Bolojâ. *Lavenia divaricata* Ham. herb. L. erecta var. f. Wall. ! cat. n. 328. Folia papyracea 5-6 poll. longa, 1 ½-2 poll. lata. (v. s.)

12. *A. ROYLEI*, caule erecto teretiusculo ramoso, foliis (saltem superioribus) alternis ovalibus basi cuneatis subsessilibus apice acutis grossè dentatis secus nervos ramis canibusque brevissimè pulverulento-puberulis, corymbo laxo paniculato, invol. squamis ovali-oblongis obtusis, acheniis muriculatis. — in Ind. orient. regionibus ad radices Hymalayæ sitis legit cl. Royle. Flores albi. Affinis *A. elato* et hirsuto. (v. s. comm. à cl. inv.)
13. *A. KLATUM* (Don prod. fl. nep. 181), caule erecto obtusè tetragono glabro simplici, foliis ovato-oblongis subsessilibus basi attenuatis crenato-dentatis subobtusis glabris, paniculà subconfertà, invol. squamis lineari-oblongis obtusis glabris, acheniis sparsè muricatis. — in Nepalià. *Lavenia carnea* Wall. ! cat. et herb. n. 329 (exclusis specim. nonnullis casu mixtis). *Lavenia elata* Spreng. syst. 3. p. 446 ex ipso in litt. ad Wall. Folia summa interdum alterna. (v. s.)
- $\beta$ . *laxum*, caule ramoso, paniculà polycephalà divaricatà. — in Nepalià et Silhet. *Lavenia alba* (sen alta) Wall. ! cat. et herb. n. 330. exclusa var. *C. Lavenia nudiflora* hort. bot. calc. à var.  $\alpha$  nullo modo nisi car. indic. differre videtur. (v. s. comm. cum var.  $\alpha$  ab hon. aulà merc. angl. Ind. orient.)
14. *A. LATIFOLIUM* (Don prod. 181), caule erecto ramoso apice præsertim puberulo hirsuto, foliis petiolatis basi cuneato-acuminatis latè ovato-rhombicis aut subcordatis vix acutis grossè serratis secus nervulos puberulis, paniculà corymbosà hirsutà polycephalà, invol. squamis subacutis dorso scabridis, acheniis muricato-tuberculatis. ① in Nepalià. *Lavenia latifolia* Spreng. syst. 3. p. 445. Habeo ex pl. Wallichianis, sed ex confus. quadam in numer. serie locus proprius incertus. Specim. habeo à cl. Leschenault in montibus Nelligherryanis lectum. Ageratum strictum Sims bot. mag. t. 2410 huc ex icon. pertinere videtur, sed descr. desideratur. (v. s. comm. ab ampl. aulà merc. angl. Ind. or.)
15. *A. RUGOSUM* (Wight in Wall. cat. n. 331. sub *Lavenià*), caule erecto basi subtetragono ramoso puberulo seu scabrido, foliis petiolatis latè ovatis subcordatis irregulariter subtrilobis grossè dentatis nervis subtus prominulis reticulato-rugosis rigidis suprà glabresculis subtus puberulo-scabridis, paniculà laxà polycephalà, invol. squamis oblongis dorso scabridis, acheniis muricato-tuberculatis. — in Peninsulà Indicà legit cl. Wight. DC. in Wight. contrib. bot. ind. p. 8. *Aden. aquaticum* Don prod. fl. nepal. p. 81. ipso mon. Lindl. Valde affine *A. reticulat.* sed achenia muriculata nec lævia? An character varians in hoc genere? (v. s. in h. hon. Cur. Ind. or. nunc Soc. Lin. Lond.)
16. *A. CAFFRUM*, caule erecto basi glabro apice pedicellisque pube brevi molli subvelutinis, foliis petiolatis ovato-rhombicis longè acuminatis basi cuneatis grossè villosò-dentatis suprà glabris subtus secus nervos junioribus tenuissimè subvelutinis, corymbo trichotomo laxo, invol. squamis lineari-oblongis obtusis basi velutinis, acheniis glanduloso-muricatis. ① in Africæ austr. dist. Caffreland legit cl. Drege! Affine *A. elato* et latifolio, sed capitula duplò fortè majora. Cor. albæ. Folia 5 poll. longa, 2 poll. lata. (v. s.)
17. *A. NATALENSE*, caule erecto hinc indè (præsertim sub foliorum jagis) petiolisque muricato-echinulatis, foliis petiolatis ovato-subcordatis basi cuneatis apice acutis dentatis suprà puberulis subtus hirtellis, corymbo laxo subpaniculato, pedicellis involucrisque puberulis, acheniis muricatis. — in Africà australi orient. ad Port Natal legit cl. Drege! Caulis è nodis infimis subradicans, circ. bipedalis. Petiol. pollicaris. Limbus folii 3 poll. longus, 2 poll. latus. Capitula florida parva. (v. s.)
18. *A. BRASILIANUM* (Cass. dict. 25. p. 363), caule erecto scabrido, foliis petiolatis latè cordato-ovatis obtusis secus petiolum subcuneatis grossè et irregulariter dentatis glabresculis, corymbi polycephali ramis puberulis, invol. squamis glabris ovali-oblongis obtusis, corollà sub lobis hirsutà, acheniis muricatis. — in Brasilià (Vand. ex Cass.), ad Sanctam-Catharinam (Bacle). *Verbesia*

*Brasiliana* Petr. ench. s. p. 472. Petioli 12-15 lin. Folia 2 poll. longa, 2  $\frac{1}{2}$  poll. lata. (v. s. comm. à cl. Bacle.)

19. *A. TRIANGULARE*, caule herbaceo erecto petiolisque subtrigono foliis petiolatis basi truncatis aut subcordatis ideo fere triangularibus dentatis glabrisculis, corymbo laxo paniculato, capitulis longè pedunculatis, invol. campanulato truncato, squamis subconcretis corollâ sub lobis hirsutâ, acheniis subtuberculatis, pappi setis 2-3.  $\frac{1}{2}$  in montosis ad margines rivulorum circa Rio-Janeiro legit cl. Lund. Flores albi. Rhizoma obliquum. Caulis pedalis. Petiol. pollicaris. Limbus fol. 2 poll. longus, 1  $\frac{1}{4}$  latus. (v. s. comm. à Mus. reg. Ber.)

\*\*\* *Caule erecto, acheniis lævibus.*

20. *A. ERECTUM*, caule erecto simplici glabro, foliis petiolatis ovatis basi acuminate grosse dentatis glabris, paniculâ divaricatâ laxâ, ramis glanduloso-puberulis, invol. squamis glabris obtusis, acheniis lævibus. — in Zeylonâ et in Ind. orientali (Wall.) comp. n. 328 et 329, sed in n. 327 est *Myriactis* spec. (forte nom. transp.). Burm. Zeyl. p. 95. t. 42. *Lavenia erecta* Willd. sp. 3. p. 1725 ? excl. Rheed. et Forst. syn. *Verbesina Lavenia* Linn. fl. Zeyl. 310. excl. syn. Plures species sub hoc nomine in herbariis et in libris meo sensu remanent confusæ ! (v. s.)

21. *A. VIRSUTUM* (Blum. ! bijdr. p. 905. sub *Laveniâ*), caule erecto subramoso glanduloso-pilosiusculo, foliis petiolatis infer. ovato-lanceolatis utrinquè acuminatis à medio ad apicem argutè serratis, paniculâ terminali virgatâ depauperatâ, ramis glanduloso-puberis, invol. squamis subobtusis dorso puberulis, acheniis subscabridis. — in humidis insulæ Javæ ad montem Gede. (v. s. comm. à cl. Blum.)

22. *A. MADURANSE* (DC. in Wight contrib. bot. ind. p. 9), caule erecto juniore cum foliis brevissimè puberulo adulto glabrato, foliis petiolatis latè ovato-rhombicis utrinquè acuminatis à basi grossè et acutè dentatis, corymbo suprâ ultima folia breviter exserto ramis puberulis, invol. squamis glabris subacutis, acheniis lævisculis. — in Indiæ orient. peninsulâ Maderaspatanâ prope Madura legit cl. Wight. *Lavenia macrophylla* Wight ! herb. non malè refert ic. Burm. Zeyl. t. 42. sed corymbus minùs exsertus et folia acutè nec obtusè dentata. (v. s. in h. Wight.)

23. *A. RETICULATUM* (DC. in Wight contrib. bot. ind. p. 8), caule erecto subtetragono glabro secùs paniculæ divaricatissimæ ramos parè glanduloso-puberulo, foliis ovatis grossè dentatis rigidis, nervis subtùs prominulis reticulatis puberulis, invol. squamis lineari-oblongis obtusis dorso subpubescentibus, acheniis lævibus. — in Ind. orient. Maderasp. australi circa Dendigul legit cl. Wight. Folia infer. non vidi. Species distinctissima nervatione foliorum ! (v. s. in h. Wight.)

24. *A. LEOCARPUM* (DC. in Wight contrib. bot. ind. p. 9), caule erecto tetragono glabro secùs paniculæ ramos glanduloso-puberulo, foliis ovali-oblongis subobtusis basi attenuatis subpetiolatis grossè dentatis glabris, paniculâ laxâ polycephalâ, invol. squamis lineari-oblongis obtusis dorso vix puberulis, acheniis sublævibus. — in Ind. or. *Lavenia alta* Wall. ! cat. et herb. n. 3220. comp. n. 330. *Ageratum aquaticum* Roxb. ! herb. sed diversum ab Aden. aquatico Don. Herba fere à clati, fructus *A. Mauritiani*. (v. s.)

+ *Species recognoscendæ.*

25. *A. MACROPHYLLUM* (Blum. ! bijdr. 905 sub *Laveniâ*) caule erecto simplici glabro, foliis infer. longè petiolatis subrhombico-ovatis utrinquè acuminatis à medio ad apicem distanter serratis, paniculâ laxæ ramis vix puberulis, invol. squamis linearibus obtusis ciliolatis, acheniis.... — in locis humidis montis Gede in ins. Java. (v. s.)

! ? *repens*, caule subramoso infernè repente, foliis angustè serratis, ramis paniculæ subalternis — ad cataractas fervidas montis Gede. Huo relata ex auctoritate cl. Blume l. c.



26. *A. AQUATICUM* (Don fl. nep. prod. p. 81), caule glabro, foliis cordatis serratis acutis glabris. ☉ in Nepaliæ inundatis. *Ethulia aquatica* Roxb. mss.
27. *A. PLATYPHYLLUM* (Cass. dict. 25. p. 363), caule erecto, pilis glandulosis sparsis, foliis oppositis petiolatis glabris ovato-deltoides secus petiolum canento-decurrentibus grossè crenato-dentatis, paniculâ corymbiformi laxissimâ. — in Peruvîâ legit Jos. de Jussieu. Fruct. indeser.
28. *A. BIFLORUM* (Less. syn. 136). — Huc cl. Lessing. refert sine descript. Verbesinam bifloram Linn. sp. 1272. sed descr. Linn. ab *Adenostemmati* caret. differt floribus lateis, pedunculis axillaribus bifloris et acheniis muticis. Icon Rheed. mal. 10 t. 40. à Willd. sub *Verbesinâ* biflorâ citatâ ab *Adenost.* omnino diversa. Species ideò omnino dubia.

† *Species exclusæ.*

*LAVENTIA DENTATA* Wall. = *Myriactis Wallichii*.  
*LAVENTIA RIGIDA* Wall. = *Leptocoma racemosa*.  
*LAVENTIA SPHERANTHA* Wall. = *Myriactis Nepalensis*.

LXIX. *SCLEROLEPIS* Cass. dict. 25 (1816). p. 365. 26. p. 227 et 233. 48. p. 153. Less. syn. 136. non Monn. — *Sparganophorus Michx.* Nutt. Ell. non Germ. — *Ethulia* sp. Walt.

Capitulum multiflorum. Invol. squamæ biseriales lineares æquales. Recept. nudum. Corolla tubulosa glabra 5-dentata, fauce vix à tubo distinctâ. Styli rami exserti cylindraceo-clavati. Achenium 5-angulare. Pappi squamæ 5 uniseriales æquales subcorneæ breves subovales obtusiusculæ. — Herba Boreali-americana glabra simplex. Folia verticillata linearia. Capitula sæpius solitaria, rarius 2-3, ramos supernè aphyllis breves terminantia. Cor. pallidè purpureæ.

1. *S. VERTICILLATA* (Cass. dict. 48. p. 155).  $\mathcal{U}$  in aquosis humidisve pinetis Carolinæ mediæ (Walt. Michx.!), et Novæ-Cæsareæ (Greene! Torr.!). *Ethulia* uniflora Walt. car. 195. *Sparganophorus verticillatus* Michx. fl. am. 2. p. 95. Nutt. gen. 2. p. 139. Ell. sketch 2. p. 312. (v. s.)

LXX. *PHANIA*. — *Stevia* sp. Linn. — *Agerati* sp. Kunth.

Capitulum 15-25-florum. Invol. squamæ 8-12 subuni- aut biseriales non imbricatæ. Corolla tubulosa glabra 5-fida, lobis acuminatis. Achenium angulatum glabrum. Pappi paleæ 5-8 oblongo-lineares ciliatæ. — Frutices Americani glabriusculi. Rami tenues. Folia opposita ovalia integerrima aut subdentata basi subattenuata. Invol. puberula. Capitula pauca pedicellata. Corollæ albæ. — Genus à *Stevia* ferè ut *Homia* à *Piqueria* differt, habitu etiam sat distinctum, sed fortè dividendum. — Nomen à Phanio vetustissimo botanico Theophrasti comæ.

Sect. 1. *EUPHANIA*.

Corolla pappi longitudine æqualis, tubo vix nullo, fauce obovatâ, lobis brevibus subacutis. — Suffrutex multicaulis. Folia trifida aut trisecta.

1. *P. MULTICAULIS*, glaber, caulibus pluribus erectis, foliis petiolatis trilobis (nunc trifidis nunc trisectis) lobis obovatis obtusis subdentatis, capitulis 1-5 cæthyumbulosis axillaribus 20-floris, invol. squamis 10-12 acutiusculis.  $\mathcal{U}$  in Americâ calidiore, Santo-Domingo? seu Brasiliâ? Species distinctissima. Capitula parva albidâ. (v. s.)

Sect. II. OXYLOBUS. — *Oxylobus Moc. fl. mex. ic. ined.*

Corolla pappo triplò circiter longior, lobis acuminatis. — Frutices Mexicani. Folia integerrima aut subdentata.

2. *P. ARBUTIFOLIA*, ramulis hirtis, foliis apice subcrenatis, inv. 12-phyllis, capitulis 20-25-floris. ♀ in Mexici montibus Nauhcampatepetl vel Cofre del Perote. *Agrostum arbutifolium* H. B. et Kunth nov. gen. et sp. 4. p. 149. *Stevia arbutifolia* Willd. herb. ex Less. in linnæa 1830. p. 140.
3. *P. TRINERVIA*, ramulis glabris, foliis integerrimis, invol. 8-phyllis, capitulis 15-16-floris. ♀ in Mexico. *Oxylobus trinervius* Moc. ic. fl. mex. Pappus dicitur 8-paleaceus et achænium tetragonum. (v. ic. pict.)

LXXI. *STEVIA* Cav. icon. 4 (1797). p. 33. Lag. nov. gen. et sp. (1816). p. 26. H. B. et Kunth nov. gen. et sp. 4 (1820). p. 139. Cass. dict. 26. p. 227. Less. syn. 135.

Capitulum 5-florum. Invol. cylindricum squamis 5-6 acutis acuminateve subæqualibus. Recept. nudum. Achænium striato-nervosum angulatimve elongatum. Pappus paleis 1-serialibus nunc in iisdem aut sæpius in diversis individuis omnibus scariosis planis parvis, nunc aliis scariosis, aliis (pler. alternis) in aristas 1-5 rigidas margine scabras abeuntibus, nunc omnibus (pler. 5-20) in aristas mutatis. — Suffrutices aut herbæ omnes ex Americâ præsertim à Mexico, rariùs ex Amer. merid. Folia infer. opposita, super. opposita aut alterna, in unicâ ternato-verticillata, linearia lanceolata aut ovata, sæpiùs triplinervia integerrima serrata aut in unicâ trifida. Capitula in corymbum densum laxumve disposita, rariùs subsparsa longius pedicellata. Cor. albæ, rosææ aut purpureæ, interdum subochroleucæ.

\* *Exaristata, nempe pappo exaristato paleaceo scarioso, paleis 5-6 nunc liberis parvis nunc in coronulam dentatam concretis.*

## a. Herbaceæ perennes.

1. *S. TRIFLORA*, caule herbaceo erecto tereti oppositè ramoso tenuissimè præsert. ad apicem pulverulento-velutino, foliis oppositis lanceolatis in petiolum subattenuatis trinerviis apice serratis suprâ glabris subtus pulverulento-velutinis, corymbo laxo composito fastigiato myriocephalo, invol. squamis acutis glabrisculis, floribus sæpiùs in invol. 3, pappo coroniformi dentato brevi. ♀ in Mexico legit cl. Karwinski. Cor. extus subglandulosæ ochroleucæ. (v. s. in h. Acad. reg. Monac.)
2. *S. CORONIFERA*, herbaceæ glabra erecta, caule tereti oppositè ramoso, foliis oppositis petiolatis ovali-lanceolatis utrinquè acuminatis subdentatis, corymbo laxè composito, capitulis ad apices ramulorum confertis, invol. squamis acuminatis, floribus circ. 5, pappo coroniformi brevi subintegro. — à Mexico misit cl. Mairet. Cor. albæ eglandulosæ. (v. s. comm. à cl. inv.)
3. *S. TRACHELOIDES*, caule herbaceo erecto densè pubescenti-velutino, ramis oppositis alternisque subcorymbosis, foliis pl. oppositis basi cuneatis sessilibus integerrimis cæterum latè ovatis acutis triplinerviis grossè dentatis utrinquè minutè pubescentibus subtus pallidis subvillosis, corymbis confertis polyccephalis, inv. squamis puberis mucronato-acuminatis, corollâ puberulo-hirtellâ, pappo coroniformi brevissimo. ♀ in Mexici valle Tolucana legit cl. Berlandier. Species distinctissima. Pappo perexiguo ferè transit in Atomium seu Piquetiam sed evid. *Stevia* congener. Cor. purpurascens. Folia 3 poll. longa, 1½-2 lata. (v. s. comm. à cl. Berl. sub n. 1164.)
4. *S. MONARDZEOIDIA* (H. B. et Kunth nov. gen. am. 4. p. 147), caule herbaceo apice ramoso hispido-hirto, ramis fastigiatis polyccephalis foliisque suboppositis

petiolatis ovatis acutis duplicato-serratis basi angustatis triplinerviis involucri-  
que hirtis, capitulis arcuè congestis, pappo paleaceo brevi.  $\frac{1}{2}$  in Mexico versas  
Valladolid. Cor. violaceæ. (v. s.)

5. *S. CLINOPODIA*, caule herbaceo erecto puberulo, foliis oppositis in petiolum  
brevem attenuatis ovato-lanceolatis parèe dentatis utrinquè puberulis tenuiter  
nervosis, ramis oppositis paniculatis apice corymbiferis, corymbis confertis,  
invol. puberuli squamis mucronatis, pappo brevi paleaceo subacuto.  $\frac{1}{2}$  in  
Mexico leg. el. Alaman et Malret. Corollæ purpureæ involucri paulò longiores  
extus glabriusculæ. Internodia distantia. Caulis fuscus. Variat foliis ovatis et  
lanceolatis. (v. s.)

6. *S. JORULLENSIS* (H. B. et Kunth nov. gen. et sp. p. 144), caule herbaceo  
apice ramoso pubescente, ramis sparsis abbreviatis corymbosis subfastigiatis  
polycephalis, foliis oppositis petiolatis ovatis obtusis crenatis basi cuneatis  
triplinerviis involucrique puberulis, pappo paleaceo brevi.  $\frac{1}{2}$  in monte Jo-  
rullo Mexici. Cor. purpurea.

7. *S. ELONGATA* (H. B. et Kunth nov. gen. am. 4. p. 144), caule herbaceo erecto  
apice ramoso piloso-hirto, ramis alternis elongatis polycephalis, foliis oppositis  
sessilibus obovatis basi cuneatis serratis triplinerviis piloso-hirtis, super. rhom-  
boideo-ovatis acutis, capitulis longè pedunculatis, involucri glanduloso-hirto,  
pappo scarioso brevi irregulariter fissis.  $\frac{1}{2}$  prope Santa-Fé de Bogota. Affinis  
St. Jorullensi et St. ovatæ. Cor. glabra.

8. *S. CONFERTA*, caule herbaceo erecto piloso aut pubescente paniculato, foliis  
pl. alternis adproximatis sublinearibus integerrimis aut subdentatis puberulis  
subglandulosis, corymbis confertis polycephalis, invol. squamis glabriusculis  
apice acuminatis, pappi paleis 5-6 adproximatis exaristatis.  $\frac{1}{2}$  in Mexico prope  
Guanaxuato (Née) et Mexico (Berl.). Cor. purpureæ involucri apice purpu-  
rascente paulò longiores. Achænia glabra.

*α. pilosa*, caule pilis articulatis subhispido. — St. pilosa Lag. elench. h.  
madr. 1805. nov. gen. et spec. 26. n. 341. Pappi paleæ obtusæ. Invol.  
apice vix patentia. (v. s.)

*β. puberula*, caule pube brevi confertà velutino-pubescente. — Pappi paleæ  
acutiores. Invol. apice subpatula. (v. s.)

*β. Fruticosa.*

9. *S. CORIACIA* (Lag. nov. gen. et sp. p. 28. n. 355), fruticosa, foliis oppositis  
ternisve subcoriaceis latè lanceolatis argutè serratis summis linearibus acutissi-  
mis integerrimis  $\frac{1}{2}$  corymbo umbellato polycephalo compacto, pappo paleaceo  
exaristato.  $\frac{1}{2}$  in Peruvia.

10. *S. SUBPUBESCENS* (Lag. nov. gen. et sp. p. 28. n. 35a), fruticosa minutè pu-  
bescens, ramis teretibus, foliis oppositis petiolatis ovato-lanceolatis acutis hinc  
indè serratis subtiùs pubescentibus, corymbo fastigiato composito, invol. pube-  
rulis, pappi subcoroniformis paleis brevibus exaristatis.  $\frac{1}{2}$  in Mexico (Lag.)  
St. lanceolata Moc. fl. mex. ic. ined. ? Cor. ex albo carneæ. Specimen habeo  
à cl. Berlandier in montibus Guchilacensis lectum huic descr. simile, sed  
foliis suprà glandulosis. (v. s.)

11. *S. GLUTINOSA* (H. B. et Kunth nov. gen. am. 4. p. 148. t. 353), fruticosa glu-  
tinosa glabra, ramulis alternis corymboso-fastigiatis, foliis oppositis longè  
petiolatis ovato-oblongis utrinquè acutis argutè serratis, corymbis capitato-  
congestis polycephalis subsessilibus, achæniis. . . . , pappo paleaceo brevi exa-  
ristato.  $\frac{1}{2}$  in planitiis Novo-Granatensium prope Santa-Fé de Bogota et  
Suacha. Cor. albæ ? Petioli 9-10 lin. longi, quà notà differt ab omnibus et spe-  
cialiter à St. lucidà ad quam cæterum accedit. Achænia ex icone puberulo-hirto  
sed de hoc caract. descriptio silet.

*α. Oaxacana*, achæniis glaberrimis.  $\frac{1}{2}$  in Mexici ditione Oaxaca legit cl.  
Andrieux (pl. exa. mex. n. 285). Optimè refert iconem et descr. var. *α*  
sed folia longiora ( $3\frac{1}{2}$  poll.) et ferè angustiora (9-12 lin.) minùs serrata.

Cor. ferè purpurascens et certè achænia glabra. An species propria?, suadente patriâ sat distante. (v. s.)

12. *S. LUCIDA* (Lag. nov. gen. am. 28), suffruticosa glabra glutinosa, foliis oppositis petiolatis lato-lanceolatis dentatis suprâ lucidis penninerviis, ramulis oppositis, corymbo conferto polycephalo, pappo coroniformi paleaceo subintegerrimo.  $\zeta$  in Mexico ad Ixmiquilpan et Cimmapan legit cl. Née. St. hymenocoppa Moc. fl. mex. ic. ined. Achænia angulata glabra. Cor. glabræ ex sicco albidæ? Petioli 3 lin. longi, limbus 2 poll. longus 6-8 lin. latus. Affinis videtur St. glutinosæ. (v. s. ex h. Née sine nom.)

13. *S. ANGUSTIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 149. excl. var.  $\alpha$ ?), suffruticosa, caule supernè puberulo-hirto, ramis fastigiato-corymbosis oppositis rariùs alternis, foliis oppositis breviter petiolatis lineari-lanceolatis utrinquè angustatis glabris subtùs involucrique subglutinosi, capitulis densè conglomeratis, pappi subcoroniformis paleis ovatis acutis brevissimis.  $\zeta$  in Mexico. Cor. albæ. Achænia angulata vix subscabrida. Pappum in floribus 1-2 inter sexcentas vidi 1-aristatum. St. salicifoliæ affinis.

$\alpha$ . *integrifolia*, foliis integerrimis. — Folia 3 poll. longa 5-6 lin. vix lata. Videtur diversa à var.  $\alpha$  Kunthianâ, fortè ipso auct. monente à specie separandâ. (v. s.)

$\beta$ . *subserrata*, foliis remotè serratis. — St. angustifolia  $\beta$  Kunth.

14. Paleaceo-aristatæ, nempe pappo paleis paucis scariosis brevibus et aristis 1-5 rigidulis scabrido-serrulatis constants.

#### $\alpha$ . Fruticosæ.

14. *S. OLIGOCEPHALA*, caule gracili imâ basi fruticose et ramoso infernè hispido, foliis oppositis lineari-lanceolatis petiolulatis tenuiter trinerviis à medio remotè serratis glabris, paniculâ laxâ oligocephalâ subnudâ, capitulis longè pedicellatis, invol. squamis glabriusculis acuminatis, pappo brevissimè paleaceo et 3-aristato.  $\zeta$  in Brasiliæ prov. Minarum Gener. ad Serro-Frio legit cl. Vauthier (pl. exa. n. 309) Species elegans distinctissima. Cor. tubus purpureus, limbus albidus. (v. s. comm. à cl. Vauthier.)

15. *S. NERVOSA*, fruticosa, ramis breviter et densè velutinis, foliis oppositis alternis ellipticis basi attenuatis apice subacutis subserratis utrinquè tenuiter pubescentibus penninerviis, nervis subtùs prominulis, ramulis summis alternis involucriisque subhirtellis, corymbis confertis, achæniis glabris, pappo paleaceo et 1-3-aristato.  $\zeta$  in Mexico prope Villalpando ad orient. urbis Guanaxuato legit cl. Mendez. Corollæ pubescentes albæ. Achænia nigra. Folia 12-15 lin. longa 5-6 lata minutè subtùs glandulosa, nervis elevatis albidis distincta. (v. s.)

16. *S. SALICIFOLIA* (Cav. ic. 4. p. 32. t. 354), fruticosa glabra, foliis oppositis angustè lanceolatis brevissimè petiolatis et subconnatis integerrimis aut medio subserratis utrinquè attenuatis, corymbo patente, pappi paleis nunc omnibus brevibus scariosis nunc 1, 2 aut 3 in setas aristatas exsertis.  $\zeta$  in Mexici montanis (Berl. ex. n. 868. 675. 825. 1202.), juxta Ixmiquilpan (Lag.) et alibi. Lag. nov. gen. p. 28. Ageratum viscosum Ort. dec. p. 36. ex Lag. St. semper-florens Ten. in litt. Rami et folia subtùs sæpè subviscosa. Pedunculi pube brevi villosiusculi. Cor. albæ. (v. s.)

17. *S. CONNATA* (Lag. elench. h. madr. 1815. nov. gen. et sp. p. 27. n. 351), fruticosa stricta glaberrima, foliis sessilibus oblango-lanceolatis serratis acutissimis, infer. oppositis connatis, super. subsparsis, corymbo fastigiato, invol. 5-7-floris, pappo paleaceo et aristato.  $\zeta$  in Mexico. St. glabra Moc. fl. mex. ic. ined. Corollæ albæ. An fortè eadem ac St. viminea?

#### $\beta$ . Herbacæ.

18. *S. VIMINEA* (Schrad. hort. Gœtt. 1832), herbacæ glabra erecta, caule simpliciter apice corymboso, foliis oppositis alternisve basi semiamplexicaulibus elongato-linearibus acutis apice subserratis, corymbo umbellato composito

- polyccephalo laxiusculo, inv. squamis acutis, pappo paleis brevibus 2-4 et aristis 1-3 constante.  $\mathcal{L}$  in Mexico. St. longifolia Moq. fl. mex. ic. ined. St. umbellata Alam. in litt. 1831. Valde accedit ad Scouarnatam Lag. sed herbacea. Cor. alba invol. ferè duplò longiores. (v. s.)
19. *S. serrata* (Cav. ic. 4. p. 33. t. 355), caule herbaceo erecto ramoso pubescente, foliis alternis subfasciculatis lineari-lanceolatis glabrisculis serratis basi integerrimis in petiolum attenuatis, corymbis fastigiatis, pappo paleaceo et bi-sæpius triaristato.  $\mathcal{L}$  in Mexici montanis, circa urbem, in valle Toluca (Berl.) ad Temascatio et Guanaxusto (H. et B.), etc. Lag. nov. gen. et sp. n. 336. H. B. et Kunth nov. gen. am. 4. p. 141. Moq. fl. mex. ic. ined. Ageratum punctatum Jacq. h. Schoenb. 3. p. 28. t. 300 non Ort. An St. virgata H. B. et Kunth nov. gen. am. 4. p. 142. satis differt? Variat caule erecto aut demisso, foliis lanceolatis aut ferè linearibus sæpè subsecundis, corollis albis aut ex albo roseis, pubescentiâ parvâ aut subhirsutâ (v. v. c. et s. sp.)
20. *S. ivæfolia* (Willd. mag. gea. naturf. Berl. 1807. p. 137. enum. 855), caule herbaceo erecto hirsuto-villoso apice ramoso-corymboso, foliis lanceolatis in petiolum attenuatis triplinerviis apice serratis, corymbis fastigiatis, invol. et corollâ glandulosis et puberis, pappo brevi paleaceo et hinc inde 1-2 aristato.  $\mathcal{L}$  in Mexico. Affinis St. serratæ sed caulis villosco-hirsutus, folia sæpius latiora plana. (v. v. c. in hort. Par.)
- $\beta$ ? *Bogotensis*, foliis minoribus non triplinerviis supra glabris et viscosis, corymbo coerctato, pappi aristis ternis.  $\mathcal{L}$  in altâ planitie Bogotensi inter Santa Fe et Chipo. St. ivæfolia H. B. et Kunth nov. gen. am. 4. p. 142 sed à verâ ivæfoliâ patriâ et caract. videtur aliena.
21. *S. eupatoria* (Willd. sp. 3. p. 1775. enum. 854. mag. naturf. Berl. 1807. p. 138), herbacea subdiffusa glabriscula, foliis lanceolatis in petiolum subtenatis trinerviis superuè obsolete serratis, corymbis fastigiatis laxiusculis, pappo paleaceo et sæpius 2-4-aristato.  $\mathcal{L}$  in Mexico. St. byscopifolia Cav. præl. 465. Lag. nov. gen. et sp. 26. n. 340. non Kunth nec Sims. Ageratum punctatum Ort. dec. 3. p. 37. (ex Lag.) non Jacq. St. punctata Pers. ench. 2. p. 403. Mustelia eupatoria Spreng. trans. Linn. soc. Lond. 6 p. 152. t. 13. St. eupatoria Sims bot. mag. t. 1849. Corollæ involucro duplò longiora, tubo incarnato, limbo intrâ albo. Achæu. glabra. (v. v. et s.)
22. *S. purpurea* (Pers. ench. 2. p. 402 excl. syn.), caule herbaceo erecto pubescenti-velutino ramosissimo, foliis lanceolatis alternis, inferioribus obovatis canaliculatis in petiolum angustatis triplinerviis apice serratis, corymbo subcoerctato, pappo paleaceo et triaristato.  $\mathcal{L}$  in Mexico (Pers. Moq.), ad radices montis Orizabæ (Sch. et Dep.). Willd. enum. 855. Moq. fl. mex. ic. ined. Less. in linnæa 1830. p. 139. Jacq. fragm. 80. t. 127. f. 2. St. eupatoria bot. reg. t. 93. Corollæ purpureæ extus puberulæ. Invol. vix puberum pallidè virescens. An satis à St. eupatoriâ diversa? An St. stricta Horn. hort. hafn. 2. p. 799. hæc revoc. ? (v. v.)
23. *S. microphylla* (H. B. et Kunth nov. gen. am. 4. p. 140), viscoso-hirta, caule herbaceo pauculato-ramoso, ramis floriferis dichotomis polyccephalis folisque oppositis subsessilibus lineari-lanceolatis obsolete crenatis, invol. squamis acuminatis, pappo breviter paleaceo et triaristato.  $\mathcal{L}$  in uliginosis inter Tasco et Thuilotepec Mexicanorum. Cor. glabra alba aut rosea.
24. *S. origanoides* (H. B. et Kunth nov. gen. am. 4. p. 147), caule herbaceo oppositè ramosissimo hirta pubescente, foliis oppositis sessilibus lanceolatis angustato-acuminatis medio serratis triplinerviis utrinquè tenuissimè hirtis, capitulis fasciculatim congestis, involucri squamis pubescentibus vix acutis, pappo breviter paleaceo et triaristato, aristis cor. puberâ brevioribus.  $\mathcal{L}$  in Mexico. Habitus Origani. Cor. albæ. Achænia glabra. (v. s.)
25. *S. microcephala*, caule herbaceo erecto simplici hirsuto-villoso, foliis alternis subsessilibus ovalibus obovatisve basi attenuatis integerrimis apice serratis dentatis subtriplinerviis utrinquè puberulo-hirtis, summis lanceolatis

- linearibusve acutis, corymbo conferto alternè ramosissimo polycephalo, capitulis pedicello proprio multò longioribus, achenio glabro, pappo brevi paleaceo et 3-5-aristato, aristis cor. extus puberulâ paulò brevioribus.  $\mathcal{L}$  in Mexico legit cl. Hauke. Capitula 3 lin. longa. Cor. ex sicco pallidè purpureascentes. Folia 9-10 lin. longa, 4-5 lata. Au St. incanescens Lag. nov. gen. et sp. n. 348? an fortè eadem ac St. pubescens Lag. ? (v. s.)
26. *S. LANCEOLATA* (Lag. nov. gen. et sp. 26. n. 338), caule herbaceo subhirsuto supernè ramoso-corymboso, foliis lanceolatis acutis planiusculis in petiolum attenuatis subtùs scabris, summis lineari-lanceolatis integerrimis, corymbo fastigiato laxiusculo, pappo paleaceo et triaristato.  $\mathcal{L}$  in Mexico circa Pachua. Var. duas dist. cl. Lagasca.
- a. floribus albis, foliis subtùs glabris.  
 ß. floribus purpureis, foliis subtùs subhirsutis.
27. *S. CANESCENS* (H. B. et Kunth nov. gen. am. 4. p. 143), caule herbaceo apice corymboso-ramosissimo polycephalo villosio-tomentoso, ramis subfastigiatis folisque sparsis subsessilibus lanceolatis basi angustatis serratis hirtopubescentibus, involucri pilosiusculo, pappo paleaceo et triaristato.  $\mathcal{L}$  in Mexico inter Guanajuato et montem Serena. Cor. alba.
28. *S. PUBESCENS* (Lag. nov. gen. et sp. 26. n. 337), caule herbaceo simpliciterectiusculo pubescente, foliis infer. oppositis subspathulatis apice dentatis in petiolum attenuatis, super. sparsis linearibus subintegerrimis, corymbo fastigiato densiusculo, invol. pubescentibus, pappo paleaceo et triaristato.  $\mathcal{L}$  in Mexico circa Guanajuato et Chalma. Cor. purpureo-sanguinea. — St. pubescens H. B. et Kunth nov. gen. am. 4. p. 143. Videtur diversa foliis lanceolatis, corollis albis, etc.
29. *S. CARACASANA*, caule herbaceo erecto tenuiter puberulo, foliis oppositis in petiolum attenuatis ovatis acutis basi cuneatis dentato-serratis subtriplinerviis subtùs glandulosis puberulisque, ramis sup. oppositis paniculatis apice corymbos polycephalos confertos gerentibus, invol. puberuli squamis apice mucronatis, pappo breviter paleaceo et 3-aristato corollam æquante.  $\mathcal{L}$  circa Caracas detexit cl. Vargas. Glandulæ fol. in sicco nigricantes. Invol. apice purpurascentia. Cor. albidæ, tubo ochroleuco. (v. s. comm. à cl. invent.)
30. *S. TRIPLIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 146), caule herbaceo obsoletè trigono ramosissimo subpubescente, ramis fastigiatis polycephalis folisque ternis sessilibus oblongis acutis crenato-serratis triplinerviis involucrique glabris, capitulis pedicellatis fasciculatis, pappo paleaceo et subbiaristato.  $\mathcal{L}$  in provinciâ Mechoacanensi Mexici. Cor. alba. Species foliis ternato-verticillatis facillè distincta.
31. *S. NEPTEFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 145), caule herbaceo erecto paniculato-ramoso tomentoso-pubescente, ramis oppositis fastigiatis apice corymbiferis, foliis oppositis ovato-lanceolatis basi longè cuneatis et subsessilibus supernè triplinerviis grossè serrato-crenatis suprâ tenuissimè pubescentibus subtùs pallidis pubero-subtomentosis, corymbis confertis, invol. pubescenti-hirtellis, pappo paleaceo et 3-aristato.  $\mathcal{L}$  in planitie Mexicanâ (H. et B.) inter Vittoria et Tula (Berl.) circa Guanajuato (Lag.?) St. suaveolens Lag. nov. gen. et sp. 27. n. 342? St. pubescens Moc. fl. mex. ic. ined. Cor. albæ aut fortè rosæ extus puberulæ. Aristæ cor. æquales. Achen. glabra. (v. s. comm. à cl. Berlandier.)
32. *S. RHOMBEFOLIA* (H. B. et Kunth nov. gen. am. 4 p. 143), pubescens, caule herbaceo apice subramoso polycephalo, foliis oppositis petiolatis rhomboideo-ovatis serrato-crenatis triplinerviis, capitulis fasciculatis breviter pedicellatis, invol. squamis acutis, pappo triaristato tubum cor. basi pubescentem æquante.  $\mathcal{L}$  in collibus montis Sanctæ Guadalupæ prope Mexico. Cor. alba.
33. *S. FASTIGIATA* (H. B. et Kunth nov. gen. am. 4. p. 148) glutinosa, caule herbaceo ramoso, ramis superioribus corymboso-fastigiatis polycephalis folisque oppositis breviter petiolatis ovato-oblongis, super. oblongo-lanceolatis

acutis grossè serratis basi cuneatis glabris, capitulis congestis, pappo brevi paleaceo rarius 1-aristato.  $\frac{1}{2}$  locis scopulosis Mexici. Affinis *St. glutinosæ* dicitur.

34. *S. UNIARISTATA*, caule herbaceo erecto pubescente subhirtello, foliis oppositis ovato-lanceolatis acuminatis in petiolum brevem attenuatis dentatis subtriplinerviis utrinquè puberulis, ramis oppositis apice corymbiferis in paniculam dispositis, corymbis confertis, involucri puberuli squamis mucronatis, achaeni capituli circ. 4, pappo paleaceo brevi obtuso, uno aristam unicam gerente.  $\frac{1}{2}$  in montibus Leoninis ad occid. urbis Guanaxuato legit cl. Mendez. *St. incanescens* Lag. nov. gen. et sp. 27. p. 348. var.  $\beta$  ? et  $\gamma$  ? Folia subtus nervosa. Corollæ in sicco flavidæ involucri paulò longiores. Folia summa oblonga lanceolata utrinquè acuminata integerrima. (v. s.)
35. *S. FASCICULARIS* (Less. in *linnæa* 1830. p. 140), caule herbaceo, foliis oppositis rhomboideo-lanceolatis profundè et argutè serratis vix pilosiusculis, super. sessilibus, ramis inflorescentiæ fastigiatis aut subfastigiatis plerique oppositis, capitulis fasciculatim congestis, corollis glabris, pappo florum cuneisque capituli sæpius 2 brevi paleaceo, aliorum 3-aristato. — in Mexico prope Jalapam et Hacienda de la Laguna.
36. *S. TOMENTOSA* (H. B. et Kunth nov. gen. am. 4. p. 145. t. 352), caule herbaceo supernè fastigiato-ramoso pubescente, foliis alternis rariùs oppositis basi cuneatis et petiolatis ovatis dentato-serratis triplinerviis suprà tenuissimè pubescentibus subtus albo- et densè tomentosis, corymbis polycephalis confertis, invol. squamis puberulis, pappo nunc brevi paleaceo nunc 1-3-aristato.  $\frac{1}{2}$  in Mexici et Toluca vallibus (Berl. ! exs. n. 1100. et 1196), prope Valladolid (H. B.) Cor. ex sicco incarnatæ. (v. s. comm. à cl. Alam. Mairet, etc.)
37. *S. MOLLIS* (Schrad. ! in *litt.* 1832), caule herbaceo pubescente supernè fastigiato-ramoso, foliis infer. oppositis petiolatis latè ovatis grossè dentato-serratis basi subcuneatis suprà minutissimè pubescentibus subtus cano-tomentosis, super. oblongis sessilibus integerrimis, corymbis laxis oligocephalis, invol. squamis dorso puberulis, pappo nunc brevi paleaceo nunc 1-2-aristato  $\frac{1}{2}$  patr. ign. Cor. ex sicco albæ. An fortè var. *St. tomentosa*? (v. s. comm. à cl. Schrad.)
38. *S. ADENOPHORA* (Lag. nov. gen. et sp. 27. n. 347), caule herbaceo glanduloso-villoso, foliis oppositis sessilibus ovatis obtusis serratis triplinerviis, basi subdeltoideâ integerrimâ, paniculâ trichotomâ laxè ramosâ, pedicellis 1-cephalis villosis glanduliferis, pappo 3-aristato, rariùs 2 seu 4-aristato.  $\frac{1}{2}$  in Chili. Cor. purpureæ. Folia summa minora integerrima.
39. *S. PANICULATA* (Lag. *elench. h. madr.* 1805. nov. gen. et sp. 27. p. 343), caule herbaceo erecto breviter pubescente paniculato, foliis infer. oppositis ovatis triplinerviis, super. alternis ovato-oblongis serratis basi cuneatis integerrimis, summis lineari-lanceolatis, pedunculis parè ramosis 3-4-cephalis laxè corymbosis, pappo nunc paleaceo brevi obtuso, nunc in fisdem capitalis 1-2-aristato.  $\frac{1}{2}$  in Mexico (Lag.) *St. hyssopifolia* Sims bot. mag. t. 1861. non Cav. nec Kunth. Corollæ paulò involucri longiores albæ, tubo purpurascente. Achaenia glabra 2-3 exaristata, 3-2 aristam 1-2 gerentia. (v. v. c.)
40. *S. OVATA* (Lag. *elench. h. madr.* 1805. nov. gen. et sp. 27. n. 344), caule herbaceo erecto glabriusculo paniculato, foliis ovatis triplinerviis serratis basi cuneatis integerrimis, super. oblongis subintegris, corymbis fastigiatis subcompactis, pappo plerumquè brevi paleaceo obtuso hinc inde uni-rariùs bi-aristato.  $\frac{1}{2}$  in Mexico. Willd. *enum.* 855. Dum. *Cours. bot. cult.* 3. p. 93. Similis *St. paniculatæ* nec fortè satis distincta. (v. v. c.)
41. *S. ELATIOR* (H. B. et Kunth nov. gen. am. 4. p. 144), caule herbaceo erecto, ramis oppositis elongatis piloso-hirtis, foliis oppositis, infer. petiolatis subrotundo-ovatis obtusis grossè crenatis basi cuneatis pilosiusculis, super. subsessilibus oblongo-ovatis, capitulis pedicellatis, invol. glanduloso-hirto, pappo

brevis paleaceo et triaristato.  $\frac{1}{2}$  prope Santa-Fé de Bogota in umbrosis humidis. Corolla alba extus tenuissimè glanduloso-hirta. Aff. *St. elongata*.

42. *S. QUINQUEVIA* (H. B. et Kunth nov. gen. am. 4. p. 145), caule herbaceo erecto supernè ramoso viscoso-villoso, ramis subfastigiatis polyccephalis folisque oppositis petiolatis ovato-oblongis acutis grossè crenato-serratis triplinerviis subtus hispidulis, capitulis pedicellatis confertis, involuero glabriusculo, pappo paleis parvis 3-1 et aristis 3-5 alternantibus.  $\frac{1}{2}$  in apricis inter urbem Quito et montem Barro Potrero. Cor. alba glabriuscula.
43. *S. PODOCEPHALA*, caule herbaceo erecto villosa-subhirsuto in paniculam laxam divaricatam ramosissimam supernè diviso, foliis caulinis oppositis sessilibus basi cuneatis ovato-rhomboides grossè dentatis triplinerviis utrinquè pubescenti-hirtis, ramis alternis ovatis subintegris, ramis pedicellis pilosoglandulosis, invol. pedicello brevioribus puberulis, pappo 3-4-paleaceo et 3-2-aristato, aristis rigidis demum divergentibus.  $\frac{1}{2}$  in Mexico leg. cl. Née, Alamán, Berlandier, Mendez. Flores invol. duplò longiores albidè extus parè puberuli. Achænia glabra. Pili caulis articulati. An idè *St. enarthotricha* Lag. huc revocanda? sed in nostrâ invol. puberum nec glabrum, folia (saltem mihi obvia) sessilia nec petiolata, corolla albidà nec purpurea. (v. s.)
44. *S. VIRATA*, caule herbaceo erecto apice subnudo folisque utrinquè pubè articulatis longâ confertâ subdecidua hirsutis, foliis in petiolum attenuatis basi cuneatis ovato-lanceolatis dentatis, capitulis fasciculatis sessilibus, fasciculis in corymbum trichotomum dispositis, invol. glabriusculis, pappo paleaceo et triaristato, aristis divergentibus corollâ puberulâ paulò longioribus.  $\frac{1}{2}$  in Mexico ad Villalpando legit cl. Mendez. Pappus *St. podocephala*. Cor. ferè *St. tenella*. Hirsuties *St. enarthotrichæ*. Cor. pallidè purpurea ex sicco. Inv. purpureascentia. (v. s. comm. à cl. Mendez.)
45. *S. MULTIFIDA* (fl. mex. ic. ined.), caule herbaceo filiformi piloso-glanduloso, foliis imis oblongis dentatis, summis alternis linearibus integerrimis, cæteris omnibus oppositis trifidis, lobis elongatis aliis dentatis aliis pinnato-incisis, corymbo laxo paucifloro, invol. squamis acuminatis dorso puberulis, achæniis tenais glabro, pappo vix paleaceo et 3-aristato. (1) in Mexico, verisim. circa urbem. Species distinctissima. (v. s. in h. Henk. ab ill. Sternberg ruis.)  
 $\beta$ . *trifida*, foliis trifidis, lobo medio aspè dente utrinquè exciso. — in Mexico. *St. trifida* Lag. nov. gen. et sp. 27. n. 350.
46. *S. MICRANTHA* (Lag. elench. h. madr. 1805. nov. gen. et sp. 27. n. 349), caule herbaceo sparse pubero paniculato, foliis oppositis alternisque ovatis serratis, inf. subcordatis, capitulis pedicellatis laxè subcorymbulosis, pappi paleis minimis, aristis 3 corollâ pubescente tubulosâ longioribus. (1) in Mexico prope Leon ad occid. urbis Guanaxuato legit cl. Mendez. *St. tenella* Moc. fl. mex. ic. ined. Species distinctissima. Flores albidè minimi. (v. s. sp.)
47. *S. DISMITIVOLIA*, herbacea glabriuscula erecta, ramis oppositis distantibus divaricatis, ramulis alternis, foliis oppos. alternisque dissitis sessilibus ellipticis integerrimis, sup. ovali-oblongis, capitulis pedicellatis ad apices ramulorum paucis, invol. squamis acuminatis, pappo breviter paleaceo et aristis 3-6 corollam vix puberulam paulò superantibus.  $\frac{1}{2}$  in Mexico legit cl. Henke. Habitus ferè *Hyperici pulchri*. Internodia ferè 3 pollic. Folia 4-5 lin. longa 3-4 lata. Cor. parvæ albidæ. Ambigit inter secundam tertiamque divisionem. (v. s. in h. Henke à cl. Sternberg comm.)
48. *S. LEUCONURA*, caule basi fruticuloso ramoso, ramis elongatis infernè pilis articulatis demum deciduis hispidis cæterum pedicellis involucriusque brevissimè puberulis, foliis oppositis et alternis lanceolatis utrinquè acuminatis serratis et super. integerrimis, nervis albis elevatis, foliis infer. basi penninerviis, super. uninerviis, paniculâ laxâ diffusâ, capitulis pedicellatis, pappo 4-5 aristato, aristis cor. æqualibus.  $\frac{1}{2}$  in Mexico ad Leon ad occid. urbis Guanaxuato legit cl. Mendez. *St. longiseta* Moc. fl. mex. ic. ined.? Affinis *St. laxifloræ*. Invol. et cor. extus purpurascentia. Achænia glabra aristis breviora (v. s. sp.)



49. *S. LAXIFLORA* (DC. h. monsp. 1813 et in Moq. fl. mex. ic. ined.), caule herbaceo pubescente ramosissimo, foliis lineari-lanceolatis plerisque alternis canaliculatis apice serratis, corymbo laxo, capitula pedicellatis, pappo paleis parvis et aristis 5 alternantibus.  $\frac{1}{2}$  in Mexico. St. purpurea (quinquearistata) Lag. nov. gen. et sp. 26. n. 339. non Pers. St. hyssopifolia H. B. et Kunth nov. gen. am. 4 p. 140. non Cav. nec Sims. Affinis St. purpureæ et cum eâ in hortis sæpè ob flores purpureos confusa. Differt staturâ majore, caule infernè pâlè raris subhispido, ramulis pube tenuissimâ subincanis, corymbo laxiore, corollis involucri duplò longioribus et aristis pappi 5 cum paleis parvis alternantibus. (v. v. c.)
50. *S. VISCIDA* (H. B. et Kunth nov. gen. am. 4. p. 140. t. 351), caule herbaceo viscoso-hirto supernè paniculato, foliis alternis sessilibus linearibus apice subcrenulatis uninerviis pubescentibus et glandulosis, corymbis oligocephalis in paniculam subcorymbosam dispositis, involucri squamis hirtò-viscosis acuminatis, pappo 5-aristato epalaceo.  $\frac{1}{2}$  in umbrosis circa Mexico (Humb. et Bonpl., Berl.!) Ageratum purpureum Sesse et St. villosa Moq. fl. mex. ic. ined.? ex quo fortè *Piptisahnati* Mexicanorum. Affinis laxifloræ. Cor. purpureæ invol. duplò longiores extûs puberæ. An potius ad sect. 3<sup>am</sup>? (v. s.)

\*\*\* Multiaristatæ, pappo paleis nullis aut subnullis sed aristis 6-20 constante.

51. *S. HEPTACHEATA*, caule herbaceo? gracili vix sparso pilosulo, foliis oppositis linearibus obtusis integerrimis vix pilosiusculis, infer. apice sublatis et hinc indè dentatis, paniculâ laxâ nudiusculâ, capitulis longè pedicellatis, pappo 6-8 aristato.  $\frac{1}{2}$ ? in Brasilîæ prov. Minarum General. ad Sabarra legit cl. Vauthier. Caulis fortè imâ basi fruticulosus ad apicem foliis distantibus minoribusque subnudus. Invol. et pedicelli extûs puberuli. Aristæ achænio puberulo longiores corollâ æquales, hinc indè in achænio quodam capituli subnullæ. (v. s. comm. à cl. Vauthier sine nom. n. 310.)
52. *S. LUNDIANA*, subherbacea simplex erecta, caule glabresculo tereti, foliis oppositis sessilibus oblongis erectis dentatis trinerviis antrisque subreticulatis glabris, paniculâ corymbosâ laxâ, invol. squamis glandulosis pubescentibusque subacutis, pappo 12-13-aristato.  $\frac{1}{2}$  in campis editis prov. Sancti Pauli nov. flor. legit cl. Lund. Herba circ. pedalis. Cor. rubentes. Pappus subnigrescens. Habitu affinis St. heptachætæ. (v. s. comm. à cl. inv.)
53. *S. SUBOCTOARISTATA* (Lag. nov. gen. et sp. 27. n. 345), herbacea glabrescula, foliis petiolatis obovatis venosis subserratis, corymbo fastigiato (Spr. syst. 3. p. 448), pappo 7-9-aristato (Lag.).  $\frac{1}{2}$  in Peruvîâ. Habitus St. ovata. Cæ. ign. †.
54. *S. HÆNKEANA*, caule basi suffruticuloso erecto oppositè ramoso apice tenuiter velutino, foliis oppositis petiolatis ellipticis utrinquè attenuatis dentatis utriusquè punctulato-subscabris, capitulis in corymbos plurimos ramosos confertis, invol. squamis acuminatis, pappo 10-12-seto corollâ vix puberulâ evidentè breviorè achæniî longit. æquali.  $\frac{1}{2}$  in Mexico legit cl. Hænke. Habitus ferè Origani. Capitula breviter pedicellata. Achænia scabriuscula. Cor. ex alio purpurascens. Petioli 3 lin. longi. Limbus fol. 12-14 lin. longus 6 lin. latus. (v. s. in h. Hænke à cl. Sternberg comm.)
55. *S. DODECACHÆTA*, caule herbaceo adscendente supernè villosa confertè folioso, foliis oppositis sessilibus ovalibus grossè dentatis triplinerviis subpuberulis, corymbis confertissimis ad apices ramorum subsessilibus polycephalis, pappo constante aristis circ. 12 achænio longioribus corollam æquantibus.  $\frac{1}{2}$  in Peruvîâ circa Lima lectam misit cl. Abadia. Habitus ferè *Pycnanthem.* Cor. tubus purpureus, limbus albidus. Invol. squamæ vix puberulæ breviter acuminatæ. (v. s.)
56. *S. HISPIDULA*, caule herbaceo erecto apice tenuissimè glanduloso-velutino, foliis oppositis oblongo-lanceolatis utrinquè attenuatis apice serratis margine et subius in nervo sparso setoso-hispidis, corymbo terminali conferto, capitulis pedicellatis, invol. squamis lanceolatis acuminatis dorso puberulis achænio

parce longioribus, pappi setis 20-25 scabris corollæ intus ad faucem et extus subvillosas subæqualibus.  $\frac{1}{2}$  in Mexico verisim. legit cl. Née, saltem ex ejus herb. habui commixtam cum St. vigintiseta. (v. s.)

57. *S. PUBERULA* (Hook. misc. bot. 2. p. 225), herbacea erecta pubera apice corymbosa, foliis alternis sessilibus ovalibus basi apiceque acutis obtusè serratis triplinerviis, corymbo denso, invol. glanduloso, pappi aristis subnullis scabris. — in Peruvia valle Canta ad Obrajillo. Cor. albæ aut carneæ.
58. *S. DIVARICATA*, caule herbaceo erecto pubescente, ramis alternis divaricatis, foliis alternis sessilibus lanceolatis integerrimis aut subdentatis utrinque attenuatis parce pubescentibus subtriplinerviis, capitulis in fasciculos 2-3-cephalos pedicellatos adgregatis laxè paniculatis, invol. brevi subobtusò, pappo 15-seto.  $\frac{1}{2}$  in Mexico? Cor. tubus purpurascens puberulus, limbus albidus pappi longitudinem vix superans. Achenium in capitulo sæpè adest unicum exaristatum. (v. s. in h. Mart.)
59. *S. POLYPHYLLA*, caule basi suffruticuloso erecto apice corymboso et pubescente, foliis oppositis in petiolum attenuatis linearibus subintegerrimis subænnerviis pubescentibus, rameis ex axillis ortis subconfertis, fasciculis oligocephalis in corymbum dispositis, invol. squamis 5-6 pappo dimidio brevioribus vix acutis, pappo 12-16-seto, corollæ puberæ æquali, acheniis puberulis.  $\frac{1}{2}$  inter Peruvianas et Panamenses herb. Hænkei recensetur. Caulis pedalis strictus. Rami steriles frequentes foliosi breves. Cor. purpurascens. Habitus fere *Silenes polyphylla*. (v. s. in h. Hænk. comm. à cl. Sternberg.)
60. *S. VERONICA*, caule herbaceo basi suffruticuloso erecto simplici glanduloso-piloso, foliis oppositis oblongis basi connatis integris apice serratis subacutis minutè et sparè puberulis, corymbo fastigiato, capitulis pedicellatis, invol. squamis lanceolatis pubescentibus corollas fere adæquantibus, pappo 15-16-seto stramineo scabrido corollam glabram subæquante.  $\frac{1}{2}$  in Brasiliæ prov. Minas Generalium ad Serro do Frio legit cl. Vanthier (n. 325!). Cor. vid. roseæ ex sicco. Achenium glabrum. (v. s. comm. à cl. inv.)
- $\beta$ . *erythrochata*, pappi setis rubescentibus. — in prov. Rio-Grande. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 943 miss.)
- $\gamma$ . *glabrata*, invol. squamis glabris. — in prov. Rio-Grande (v. s. in eod. herb. n. 925.)
61. *S. LINARIÆFOLIA*, caule herbaceo erecto ramosissimo parce piloso, foliis alternis linearibus integerrimis vix uniuerviis utrinque pubescenti-subhirtellis, pedicellis elongatis 1-cephalis paucis ad apices ramorum, invol. squamis acutiusculis, pappo 12-14-seto corollæ puberulæ longitudinem subæquante.  $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Folia 12-15 lin. longa, sesquilineam lata. Cor. ex sicco albidæ. Achenium subscabridum. (v. s. in h. Mus. reg. Par. sub n. 852 à Mus. imp. Bras. acc.)
62. *S. MEGAPOTAMICA*, caule basi subfruticuloso ramoso, ramis sparè hispidulis teretibus, foliis oppositis linearibus glabrinusculis hinc indè subserratis, pedicellis elongatis 1-cephalis in corymbum laxum dispositis, invol. 5-6-phyllò, squamis acutis glabrinusculis, achenio glabro striato maturo invol. vix breviorè, pappo 20 seto scabro cor. æquali.  $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Herba 6-8 poll. longa. Folia pollicem longa, lineam lata. Cor. purpurascens. Pappus subfuscescens. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 928 miss.)
63. *S. OKYLENA*, caule imò basi suffruticuloso tereti glabrinusculo, foliis oppositis oblongo-lanceolatis sessilibus utrinque attenuatis dentatis hinc indè subitis et ad margines sparè setosis, capitulis ad apices ramorum paucis brevè pedicellatis, invol. squamis 5-6 lanceolatis acuminatis dorso setosis, achenio glabro, pappi setis circ. 20 scabris.  $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Herba 7-8 pollic. Cor. apice puberæ. Pappus flavescens. (v. s. in h. Mus. reg. Par. à Mus. Bras. sub n. 998 miss.)
64. *S. VIGINTISETA*, caule herbaceo glabrinusculo erecto alternè paniculato-ramoso, ramis apice corymbiferis, foliis alternis brevissimè petiolatis lanceolatis

acuminatis subintegerrimis triplinerviis parvè subcabridis, capitulis breviter pedicellatis confertiusculè corymbosis, invol. squamis subacutis achænia vix superantibus, achæniis multicoctatis piloso-hirtellis, pappi aristis 20-24 scaberrimis corollâ paulò brevioribus. — in Mexico legit cl. Karvinaki. Cor. purpureæ glabræ. Pappus sordidè purpurascens. Styli rami valdè elongati fert omnino epapulosi. (v. s. in h. Acad. reg. Monac.)

65. *S. narunculoides*, caule herbaceo glabresculo apice pubero erecto alternè paniculato, foliis alternis brevissimè petiolatis elongato-lanceolatis utrinquè attenuatis triplinerviis crenato-dentatis utrinquè scabridis, capitulis ad apices ramorum paucis pedicellatis, invol. squamis linearibus acuminatis glabris achænia superantibus, achænio angulato sulcato pilosinaculo, pappi aristis 25-30 scaberrimis achænio longioribus. ♀ in Mexico ad occid. urbis Guanaxuato legit cl. Mendez. Inv. purpurascens. Pappus flavescens. Capitula majora quàm in cæteris Steviis. (v. s. comm. à cl. inv.)

66. *S. calycina*, caule suffruticoso erecto tereti basi glabro apice pulverulento-subvelutino, foliis linearibus utrinquè attenuatis integerrimis glabresculis axillis foliiferis, corymbo laxo paniculato, invol. duplici, ext. squamis bracteivæ 5-6 linearibus, inter. squamis 5 linearibus lanceolatis margine scariosis pappum maturum subæquante, achænio nitido glabro, pappi setis circ. 50 scabris. ♂ in Brasiliæ prov. Rio-Grande. Habitus ferè *St. polyphylla*, sed invol. achæni. et pappus valdè differunt. Fruticulus sesquipedalis. Folia pollicem longa,  $\frac{1}{2}$  lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 905 miss.)

67. *S. multiaristata* (Spreng. syst. 3. p. 449), herbacea glanduloso-pubescent, foliis alternis linearibus cauliculis subintegerrimis subtus canescentibus, pedunculis 1-cephalis subracemosis, pappo multiaristato corollam longè superante. — in Monte-Video.

† *Non satis nota.*

68. *S. urticæfolia* (Thunb. ex florè 1821. p. 331), herbacea, foliis ovatis acutis serratis capitulis paniculatis. — in Brasiliâ. Cæt. ex opere Thunb. mihi ignoto invitus omitto.

†† *Species exclusæ.*

*S. callosa* Nutt. = *Florestina callosa*.

69. *FERRIFUGA* Fl. mex. = *Piqueria trinervia*.

*S. linearis* Cav. = *Palafoxia linearis*.

*S. pedata* Cav. = *Florestina pedata*.

LXXII. *PALAFoxia* Lag. nov. gen. h. madr. (1815. et april. 1816) p. 26. *Sims bot mag. t. 2132. Spreng. syst. 3. p. 449.* — *Paleolaria* Cass. bull. philom. decemb. 1816. mart. 1818. dict. suppl. 1. p. 59. 37. p. 256. Less. syn. 155. — *Polypteris* Nutt. gen. am. (1818). 2. p. 139. non Less. — *Stevia* et *Agerati* sp. auct.

Capitula pluriflora. Invol. oblongum campanulatumve vix subimbricatum, squamis 8-14 in fructu maturo stellato-patentibus. Recept. nudum. Achænia angulata, exter. squamis involuta. Styli rami elongati obtusi semiteretes dorso papillis elongatis subhispidi. Pappus paleis 8-12 scariosis 1-nerviis pinnato-striatis siccis 1-serialibus alternis brevioribus aut biserialibus exterioribus brevioribus constans. — Frutices aut herbæ Americane habitu Stevias referentes. Folia inf. opposita et sup. alterna indivisa. Capitula laxè corymbosa. Cor. albæ carnæ aut purpureæ. — Genus habitu valdè accedens ad *Florestinam Helianthis* adjunctam, et styl. habitu ferè *Vernoniaceas* referens.

1. *P. linearis* (Lag. ! nov. gen. et sp. h. madr. 1815. n. 335), fruticosa cinereo-

pubescens, foliis linearibus integerrimis obtusis, involucri oblongo 10-12-floro, pappi paleis 8 inæqualibus acuminatis longioribus achenio puberulo elongato brevioribus.  $\S$  in Mexico. Bot. mag. t. 2132. *Stevia linearis* Cav. l. præl. n. 464. *Ageratum lineare* Cav. icon. 3. p. 3. t. 205. *Paleolaria carnea* Cass. ll. cc. *Stevia linearis* et *lavandulæfolia* Willd. Corollæ carneæ. Achenia exteriora interdum pappo minimo subabortivo tantum donata. (v. v. et s.)

2. *P. FASTIGIATA* (Less. syn. 156), caule herbaceo subscabro, ramis fastigiato-corymbose, foliis lineari-lanceolatis integerrimis utrinque scabris, invol. squamis 12-15-biserialibus, ext. brevioribus, pappi paleis 12-14 lineari-lanceolatis, achenii subscabri longitudine.  $\mathcal{Q}$  in Amer. bor. ad ripas flum. Altamahah. *Polypteris integrifolia* Nutt. gen. am. 2 p. 139. ex Less. Cor. tubus pubescens. Achenium subquadrangulare obconicum. Pappi paleæ interdum lacere.
3. *P. TEXANA*, caule herbaceo puberulo, ramis glanduloso-puberis, foliis lanceolato-linearibus acuminatis trinerviis integerrimis puberulis, capitulis circ. 30-floris, invol. squamis 10-12 dorso glanduloso-puberis, pappo fl. ext. brevi vix acuto, fl. inter. aristato-acuminato achenio æquali. — in Mexici provinciâ Texas legit cl. Berlandier (pl. exs. n. 2014). Folia petiolo 4 lin., limbo bipoll. Cor. ex sicco incarnatæ. Achenia tetragona puberula basi attenuata. (v. s.)
4. *P. LATIFOLIA*, caule herbaceo erecto basi pubescente apice pubescenti-glanduloso, foliis oppositis petiolatis cordatis suborbiculatis grossè dentatis utrinque pubescenti-villosis basi trinerviis, paniculâ laxâ subnudâ, invol. squamis obovatis obtusis, pappi paleis 8 alternis elongatis acuminatis, alternis brevioribus obovatis. — in Mexico circa Tehuantepec mens. august. legit cl. G. Andrieux (pl. exs. mex. n. 286). Species distinctissima! Cor. purpureæ in invol. circ. 20. (v. a. comm. à cl. inv.)

LXXIII. *CARELIA* Less. syn. 156. non *Adans.* nec *Ponted.* nec *Rieux.*

Capitulum multiflorum. Recept. convexum pilosiusculum. Invol. campanulati squamæ ovali-oblongæ 2-3-seriales margine membranaceæ. Cor. tubulosæ 5-dentatæ. Achenia 5-gona. Pappus biserialis æqualis, paleis brevibus obtusis nerviis apice serrulatis. — Fruticulus Brasiliensis villosus-canescens. Folia opposita oblongo-ovata petiolata basi obtusa. Capitula dichotomè corymbosa.

1. *C. CISTIFOLIA* (Less. l. c.).  $\S$  in Brasiliâ (Less.), in prov. Rio-Grande. Hoc refero specimen descr. equidem brevissimæ Lessingii optimè conveniens, sed fortè specificè distinguendum. Petioli 4 lin., limbus fol. 15-16 lin. 5-6 lin. latus. Capitula pedicellata 50-flora. (v. a. in h. Mus. reg. Paris. à Mus. imp. Bras. sub n. 907 miss.)

LXXIV. *AGRIANTHUS* Mart. in herb. Acad. Monac.

Capitulum circ. 20-florum. Invol. squamæ 1-2-seriales lineares acuminatæ subæquales. Recept. nudum. Cor. cylindraceæ obtusè 5-dentatæ. Styli rami longè exserti cylindrici obtusi. Achenia acutè pentagona secus angulos scabra, areolâ basilari prominulâ. Pappus corollâ brevior 1-serialis, squamellis lineari-lanceolatis acuminatis rigidulis inæqualibus densè barbellato-ciliatis. — Frutices Brasilienses ramosi humiles glabriusculi. Rami teretes. Folia conferta subimbricata alterna rigida lineari-lanceolata aut acerosa. Capitula in axillis supremis sessilia aut breviter pedicellata in corymbum densum capitatum adgregata. Cor. ex sicco pallidè purpurascens. — Genus ambiguum inter *Agerateas* et *Adenostyleas*.

1. *A. CAMPESTRIS* (Mart. l. c.), foliis lato-linearibus planiusculis minutissimè pubescentibus subtus reticulatis, pappo corollâ triplo ferè breviori.  $\S$  in Brasiliâ prov. Minarum Generalium, in altis graminosis deserti inter Minas Novas et

flum. Sancti-Francisci, legit cl. Martius. Folia 10-12 lin. longa, 1  $\frac{1}{2}$ -2 lata. (v. s. in h. Acad. reg. Monac.)

2. *A. AMPETRIFOLIUS* (Mart. l. c.), foliis acerosis subtriquetris glaberrimis, pappo corollâ paululum brevior.  $\frac{5}{8}$  in Brasiliæ prov. Bahiensi interiori in montibus ad Rio de Contas Vales legit cl. Martius. Flor. octob. Rami divaricati breves. Folia 2-3 lin. longa, lineam lata. (v. s. in h. Acad. reg. Monac.)

**DIV. III. ADENOSTYLÆ. — Eupatoriæ Less. syn. 156. — Eupatoriæ prototypæ et Adenostylæ Cass. tabl.**

Pappus pilosus barbellatus plumosusve uni- aut pluriserialis. — *Adenostyles* non divellendæ, nec ex recept. paleaceo aut nudo, nam *Chromolaena* et *Decachæta* ad *Eupatorium* nimis accedunt; nec ex pappo piloso aut plumoso, nam *Liatrides* plurimæ transitum exhibent.

**LXXV. KUHNIA** Linn. sp. pl. ed 2 (1763). p. 1662. Vent. h. cels. n. 91. Cass. dict. 24. p. 515. Less. syn. 157. Rafin. herb. p. 29. — *Critonia Gært. fr. 2. p. 411. t. 174. non R. Brown.* — *Eupatorii* sp. Ort.

Capitulum multiflorum. Invol. squamis pauciserialibus. Recept. nudum. Cor. tubus ad faucem non ampliatus. Antheræ interdum liberæ (nunc sæpius abortivæ). Styli rami exserti cylindracei obtusi. Achænum elongatum sessile aut stipitatum. Pappus 1-serialis plumosus aut rarius barbellatus. — Herbæ aut suffrutices Americani. Folia alterna 1-nervia rarius triplinervia linearia aut lanceolata. Capitula paniculato-corymbosa pedicellata. Cor. purpurascens aut albæ.

**Sect. I. STRIGIA.**

Achænia teretiuscula multistriata. Pappus plumosus. Invol. squamæ imbricatæ, exter. breviores.

1. *K. EUPATORIODES* (Linn. sp. 1662. Linn. f. deced. 2. p. 21. t. 11), caule herbaceo, foliis lato-lanceolatis serratis, corymbis paniculatis terminalibus.  $\frac{2}{3}$  in Pensylvaniâ (Linn.), in Sibiria? (Ard. sed verisim. falso). Vent. h. Cels. n. 91. in adn. Ell. sketch 2. p. 291. *Eupatorium alternifolium* Ard. spec. 2. p. 40. t. 20. Flores albi aut albi. Folia ex Ell. subtus pubescentia glanduloso-punctata. Capitulum 10-florum.
2. *K. PANICULATA* (Cass. dict. 24. p. 516), caule herbaceo, foliis linearibus aut lineari-lanceolatis integerrimis junioribus margine revolutis, ramis apice corymbosis in paniculam laxam dispositis.  $\frac{2}{3}$  in siccis Carolinæ (Ell. Bosc!), in montanis Pensylvaniæ, Virginie (Pursh) et Alabama (Torr.!). *Critonia Kuhnia* Michx. fl. bor. am. 2. p. 101. *Kuhnia Critonia* Willd. spec. 3. p. 1773. Ell. sketch 2. p. 291. Flores albi. Folia subtus punctata. (v. s.)
3. *K. ROSMARINIFOLIA* (Vent. h. Cels. n. et t. 91), caule suffruticuloso, foliis lineari-lanceolatis integerrimis junioribus margine revolutis, pedunculis panicis subcorymbosis.  $\frac{5}{8}$  in insulâ Cubâ. *Eupatorium canescens* Ort. dec. p. 34. *K. frutescens* Horn. h. hafn. 2. p. 791. *Eupatorium Cubense* Pers. ench. 2. p. 408. Flores purpurei. Folia subtus glabra punctata. (v. v. c. in h. Cels. et s. ex h. Madr.)

**Sect. II. TRICHOGONIA.**

Achænia pentagona, angulis piloso-scabris. Pappus plumosus. Invol. squamæ æquales nec imbricatæ. — An genus proprium?

4. *K. ARGUTA* (H. B. et Kunth nov. gen. am. 4. p. 105. t. 393), caule herbaceo subviscoso-hirto, foliis angustè lanceolatis argutè serratis tenuissimè hirtis, corymbis terminalibus.  $\frac{2}{3}$  in montanis siccis prope Popsyen Novo-Granat. Flores rosei.

5. *K. villosa* (Spreng. *syat.* 3. p. 439), caule fruticuloso ramoso apice dense piloso-glanduloso, foliis linearibus integerrimis aut remotè denticulatis marginè revolutis dense piloso-glandulosis, pedicellis paucis 1-cephalis in corymbum subdigestis.  $\frac{3}{4}$  in Brasiliæ prov. Minarum General. ad Serra do Frio legit cl. Vauthier (pl. exs. n. 313). Folia secus ramos conferta, pollicem longa vix lineam lata. Invol. oligophyllum dorso pubescens. Corollæ subhispidulæ. (v. s. comm. à cl. Vauthier.)
6. *K. hirtiflora*, fruticulosa, ramulis striatis hirtis, foliis brevissimè petiolatis ovatis subcordatis acutiusculis dentatis basi trinerviis suprâ sparsè puberulis subtus pubescenti-subhirsutis, corymbo terminali subsimplici oligocephalo conferto, capitulis breviter pedicellatis circ. 40-floris, invol. squamis 1-2-seriatis aequalibus oblongis dorso puberis obtusiusculis, cor. fauce dilatata extus hirtâ.  $\frac{3}{4}$  in Brasiliæ prov. Minarum General. legit cl. Gaudichaud. Achænia longa 5-costata basi attenuata secus angulos hirta. Pappus plumosus. Folia 6 lin. longa 5 lin. lata. (v. s. in h. Mus. reg. Par.)
7. *K. glabrata*, caule erecto simplici basi suffruticoso glabro apice nudo ad summitatem pubescente, foliis glabris sessilibus obovato-cuneatis triplinerviis, corymbo composito, invol. squamis circ. biserialibus ovali-oblongis obtusiusculis.  $\frac{3}{4}$   $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Habitus *K. glutinosæ*. Flor. in invol. 10. Achænia pentagona obpyramidata parcè villosa. Pappi setæ crassæ barbellatæ.
- a. foliis subintegerrimis. (v. s. in h. Mus. Par. à Mus. imp. Bras. sub n. 1055 miss.)
- b. foliis serratis. (v. s. in eod. herb. sub n. 784.)

## Sect. III. LEIOGONIA.

Achænia pentagona omnino lævia. Pappus barbellatus. Invol. squamæ paucæ subimbricatæ. — An potiùs Eupatorii species?

8. *K. podocarpa*, fruticulosa erecta ramosa, ramis teretibus minutissimè hirtellis, foliis alternis petiolatis ovatis acutis dentatis suprâ glabresculis subtus petiolisque puberulo-velutinis, capitulis ad apices ramorum paucis pedicellatis, invol. squamis oblongis biserialibus, ext. foliaceis dorso puberulis, achæniis 5-gonis glabris longiusculè stipitatis.  $\frac{3}{4}$  in Brasiliæ arenosis maritimis Restingas dictis. Fl. nov. et dec. Tota planta odorata (Lund in litt.). Folia ferè formâ *Melissæ*. Cor. in sicco pallidæ. Habitu accedit ad sect. secundam, sed achænia lævia. (v. s.)
9. *K? glutinosa* (Ell. sketch 2. p. 292), glutinoso-pubescent, caule herbaceo, foliis lanceolatis basi attenuatis triplinerviis integerrimis apice acuminatis rariter et grossè dentatis, capitulis corymboso-paniculatis.  $\frac{1}{2}$  in pratis Alabama Amer. bor. Capitulum 8-10-florum. Cor. alba. Styli rami breviter exserti. Achænium acutè pentagonum, angulis lateribusque glaberrimis. Pappus albidus 1-serialis multisetus, setis scabris nec verè plumosis. (v. s. comm. à cl. Elliott.)
10. *K. baccharoides*, fruticosa glabra ad summitates viscosa, foliis obovatis basi cuneatis in petiolum attenuatis apice obtusis dentatis coriaceis reticulato-venosis glabris, corymbo composito nudo, invol. squamis linearibus paucis-seriatis.  $\frac{3}{4}$  in Brasiliâ circa Bahiam legit cl. Blanchet (pl. exs. n. 1826). Folia sequalipollic. longa 8-9 lin. lata. Cor. ex sicco pallidè flavæ. Achænium acutè pentagonum læve. Pappus uniserialis setis breviter et dense plumosis. (v. s.)

## LXXVI. CLAVIGERA.

Capitulum 5-20-florum. Invol. pluriseriale imbricatum, squamis longitudinaliter striatis, ext. brevissimis intimis elongatis linearibus. Recept. angustum epaleaceum. Cor. tubulosæ cylindricæ brevissimè 5-dentatæ, dentibus extus glanduliferis. Anthæræ apice appendiculatæ. Styli rami teretes nunc inclusi nunc parcè exserti. Achænium subcy-

lindraceum glabrum. Pappus 1-serialis, setis breviter barbellato-plumosis. — Suffrutices Mexicani erecti ramosissimi. Folia alterna integerrima aut dentata, linearia aut oblonga, uninervia impunctata. Capitula in paniculam spiciformem aut corymbosam disposita. Flores albi. — Genus medium inter Kuhniam et Liatridem, ob specierum in Mexico omnium habitantium, dicavi cl. Franc. Xav. Clavijero qui historiam Mexici non tantum civilem, sed etiam naturalem scripsit. (1)

1. *C. CORYMBOSA*, glabra, caule basi suffruticoso, foliis linearibus integerrimis, capitulis circ. 20-floris corymbosis. ♂ ♀ in Mexici provinciâ Leonensi legit cl. Mendez. Invol. squamæ intensè striatæ. (v. s.)
2. *C. SCOPARIA*, glabra, caule basi suffruticuloso, foliis linearibus integerrimis, capitulis circ. 10-floris in paniculam angustam spiciformem longè digestis. ♂ ♀ in Mexici prov. Leonensi legit cl. Mendez, in Cordillera Gochilapana Berlandier, et ad sinum Mulgrave ex herb. Hænke. Invol. squamæ striatæ. Rami paululum ad Liatridem mucronatam accedunt sed caract. diversissimi. (v. s.)
3. *C. DENTATA*, pubescens cinerea fruticosa, foliis ovali-oblongis dentatis hinc inde apice sublobatis, ramulis foliolis oligo- aut monocephalis in paniculam angustam dispositis, capitulis 12-floris, invol. squamis linearilanceolatis acuminatis ciliatis apice subscariosis, parce striatis. ♂ in Mexicanæ prov. Texas distr. (Commancheria) orientalibus necnon circa Bejar legit cl. Berlandier, pl. exs. n. 1897. 2079 et 2081! (v. s.)

LXXVII. LIATRIS Schreb. gen. (1791). n. 1263. Willd. sp. 3. p. 1634. Nutt. gen. am. 2. p. 131. Less. syn. 157. non Don. — Anonymos Walt. fl. car. (1788). p. 196. — Suprago Gærtn. fr. (1791). 2. p. 402. — Liatris Suprago et Trilba Cass. — Serratulæ sp. Linn. — Vernoniæ sp. Poir. — Kuhniz sp. Juss.

Capitulum multi- (5-30) florum. Invol. pauciserialè imbricatum. Recept. nudum. Cor. tubulosa ad faucem ampliata, lobis elongatis. Styli rami longè exserti cylindracei. Achænium circ. 10-costatum subcylindraceum. Pappus 1-3-serialis, setis plumosis barbellatisve. — Herbæ rarò frutices. Species omnes genuinæ ex Amer. bor. ortæ. Caulès sæpius simplices. Folia alterna integerrima aut in paucissimis dentata. Capitula spicata racemosa paniculata aut corymbosa. Cor. purpureæ rosæ rarius variantes albæ. — Interdum in plantis cultis vidi styli ramos in ligulas petaloideas eglandulosas mutatos. Radices tuberosæ resinosæ terebinthinam spirant undè à Canadensibus dicuntur *Pinetta de Prairie* et hinc inde ut diuretica imò antisiphilitica admissæ.

Sect. 1. EULIATRIS. — Liatris Cass. dict. 26. p. 235. non alior.

Pappus 1-serialis evidenter et longè plumosus. — Herbæ radice globoso-tuberosâ vix deorsum subfibrillosâ. Caulès simplices. Capitula spicata aut simpliciter racemosa.

1. *L. TENUIFOLIA* (Nutt. gen. am. 2. p. 131), radice tuberosâ, caule gracili glabro, foliis filiformi-linearibus longissimis infernè confertis summis brevibus setaceis infernè subpilosiusculis, capitulis pedicello ramoso insidentibus in racemum longissimum digestis oblongis 5-floris, invol. squamis oblongis submucronatis, achænio villosa, pappo plumoso. ♀ in pinetis arenosis Carolinae

(1) Opus anno 1780 italicè, et curante C. Cullenio anno 1787 anglicè editum (Conf. Bibl. voy.)

- utrinque. Ell. sketch 2. p. 276. *L. gysaminifolia* Willd. sp. 3. p. 1636? Folia ferè Pini palustris. Racemus 1-2-pedalis. Pedicelli circ. pollicares. Flores purpurei.
2. *L. mucronata*, radice. . . . , caule simplici glabra, foliis angustè linearibus acutiusculis calloso-mucronatis glaberrimis erectis punctatis, capitulis circ. 5-floris ad axillam bracteis sessilibus approximatis in spicam digestis, invol. squamis ciliatis obtusis abruptè mucronatis, achenio pubescente, pappo plumoso.  $\frac{1}{2}$  in Mexicanæ provinciæ Texas distr. (Cammancherias) orientalibus legit cl. Berlandier. Fructif. mens. nov. et dec. Affinis videtur *L. tenuifolia* sed capitula semilia. (v. a.)
3. *L. elegans* (Willd. sp. 3. p. 1635), radice tuberosâ, caule simplici pubescenti-villoso, foliis linearibus mucronatis scabrido-punctatis, capitulis breviter pedicellatis adproximatis 3-5-floris, racemo spiciformi elongato, invol. squamis acuminatis, interioribus coloratis, achenio villoso, pappi setis plumosis.  $\frac{1}{2}$  in Virginis (Ker), Carolinæ? (Walt. Fras. etc.), Floridæ (Michx.) et Georgiæ (Dab.) siccis. Bot. reg. t. 267. Ell. sketch 2. p. 279. *Stæbelina elegans* Walt. fl. car. 202. *Serratula elegans* Ait. Kew. ed. 1. v. 3. p. 138. *Eupatorium speciosum* Vent. h. Cals. n. et t. 79. Folia sæpè falcata patentia sæpè reflexa. (v. a.)
4. *L. squarrosa* (Willd. sp. 3. p. 1634), radice tuberosâ, caule simplici pubescente, foliis longissimè linearibus nervosis margine scabriusculis, capitulis paucis in racemum digestis ad apicem pedicelli folioli solitariis, invol. cylindrici squamis superne foliaceis lanceolatis rigidis patentibus, achenio glabriusculo, pappi setis plumosis.  $\frac{1}{2}$  in arenosis campis et sylvis Canadæ (Hook. fl. bor. 1. p. 306), Virginis, Carolinæ et agri Kentuckensis. Parah fl. 2. p. 509. Cass. dict. 26. p. 235. Sweet br. fl. gard. t. 44? malè. Ell. sketch 2. p. 282. *Serratula squarrosa* Linn. sp. 1147. *Pteronia caroliniana* Walt. fl. car. p. 292. ex Parah. — Dill. elth. 83. t. 71. f. 82. Moris. ox. 3. t. 27. f. 1. Cor. latè purpurea, exter. demùm lateraliter rejecta. Pappus 1-serialis rufo-purpureus. Ach. striatum. Var. foliis et invol. glabriusculis aut pubescentibus. (v. a.)
- fl. intermedia*, caule humiliore folisque magis pilosis, invol. squamis ext. elongatis acuminatis erectis. — in Canada. *L. intermedia* bot. reg. t. 948. (v. a.)
5. *L. punctata* (Hook. fl. bor. am. 1. p. 306. t. 105), radice. . . . , caule erecto-nuculo simplici piloso, foliis linearibus marginatis glabris ciliatis utrinquè impresso-punctatis, spicis elongatis densis non rarè foliosis, invol. subcylindrici circ. 5-flori squamis imbricatis subaquarrosis, ext. rigidioribus, omnibus impresso-punctatis pungenti-acuminatis margine ciliato-lanatis, achenio piloso, pappo valdè plumoso. — in Amer. bor. occid. planitiibus ad Saskatchewan Red-deer et Eagl. hills. Cor. roseæ.
6. *L. resinosa* (Nutt. gen. am. 2. p. 131), glabra, radice tuberosâ, caule erecto simplici aut apice vix ramoso, foliis linearibus rigidulis integerrimis utrinquè impresso-punctatis, spicis elongatis densis aspius foliosis, invol. cylindracei 5-6-flori squamis imbricatis erectis longè acuminatis punctulatis, achenio villoso, pappo densè plumoso. — in Carolinæ utriusque pinetis (Nutt.), in territorio Arkansaseno (Pourt.). Flores purpurei elegantes. (v. v. c. in h. Genev. ex sem. à cl. de Pourtales lectis.)

Sect. II. SUPRAGO. — Suprago Cass. dict. 51. p. 385. — Supraginis sp. Gartn.

Pappus 1-3-serialis, setis densè et breviter barbellatis. — Herba, radice globoso-tuberosâ vix deorsum subfibrillosâ. Caules simpliciores. Capitula spicata aut racemosæ. — Radix et inflorescentia sect. 1<sup>ma</sup>. Pappus inter 1<sup>am</sup> et 3<sup>am</sup> medius.

7. *L. scariosa* (Willd. sp. 3. p. 1635), radice tuberosâ, caule simplici pubescenti-villoso, foliis lanceolatis pubescentibus margine scabris, infer. longissimis basi attenuatis, capitulis breviter pedicellatis racemoso-spicatis 15-30-floris, invol.



- aquamis obovatis obtusis margine scariosis, exter. patentibus acutis, achænio villosio, pappo subplumoso achænio subbreuiore.  $\mathcal{L}$  à Virginîæ ad Carolinam in montanis pratis (Pursh), in siccis (Ell.! sketch 2. p. 280 cum var. plur.). bot. reg. t. 590. non Curt. Anonymos racemosus Walt. car. 198. ex Ell. *Liatris squarrosa* Michx. fl. bor. am. 2. p. 92. *Serratula scariosa* Linn. sp. 1147. excl. Pluk. syn. *Vernonia scariosa* Poir. dict. 8. p. 502. — Pluk. alm. t. 177. f. 4. Variat capitulis 15-20-25-30-floris, foliis glabrisculis pubescentibus aut fere villosis, floralibus involucri brevioribus aut longioribus, spicâ compactâ aut dissitiflorâ. (v. s. comm. à cl. Ell.)
8. *L. SPHERIOLEA* (Michx. fl. bor. am. 2. p. 92), radice tuberosâ, caule pubescente, foliis lævibus infer. lato-lanceolatis super. lanceolato-linearibus, capitulis pedunculatis majusculis distantibus subglobosis 30-40-floris, invol. squamis latè ovalibus obtusis adpressis margine scariosis, achænio multicoctato piloso scabro, pappi setis barbellatis achænio longioribus.  $\mathcal{L}$  in Carolinæ distr. superioribus. Ell. sketch 2. p. 281. Sw. br. fl. gard. t. 87. *Suprago sphaerocephala* Cass. dict. 51. p. 386. *Liatris scariosa* bot. mag. t. 1709. excl. syn. (v. s.)
9. *L. SPICATA* (Willd. sp. 3. p. 1636), radice tuberosâ, caule simpliciter glabro, foliis linearibus glabris basi ciliatis nervosis et punctatis, capitulis longè spicatis ad axillam bractearum sessilibus 8-floris, invol. squamis oblongis adpressis obtusis sæpè mucronatis, achænio piloso, pappi setis densè barbellatis.  $\mathcal{L}$  in pratis et pinetis demissis à Pennsylvania ad Georgiam. Sims bot. mag. t. 1411. *Serratula spicata* Linn. sp. 1147. And. rep. t. 401. *Suprago spicata* Gært. fr. 2. p. 402. t. 167. Cass. dict. 51. p. 385. *Liatris picta* Bart. ex Green. herb. L. gracilis Lodd. bot. cab. t. 1909. non Nutt. Anonymos ciliatus Walt. car. p. 197. An L. *linaria* Raf. fl. lud. 61 satis differt? Flores purpurei. (v. s.)
- $\beta$ . *macrostachya*, spicâ densiore, capitulis brevissimis pedicellatis. *L. macrostachya* Michx. fl. 2. p. 91. ex Ell. sketch 2. p. 273. *L. spicata* Sweet br. fl. gard. t. 1. 49. *L. pumila* Lodd. cab. t. 147? — Dill. elth. t. 72. f. 83. (v. s.)
- $\gamma$ . *racemosa*, capitulis distinctè pedicellatis. *Liatris pilosa* bot. reg. t. 395. Lodd. bot. cab. 4. t. 356. non alior. An *L. turbinata* Sweet in Loud. hort. brit. 383. sine descr.? Differt à verâ *L. pilosâ* caule glabro nec pubescente, foliis ciliatis ut in var.  $\alpha$  nec pilosis, floribus in invol. 8 nec 10-14. (v. s.)
10. *L. PYCNOSTACHYA* (Michx. fl. 2. p. 91), radice tuberosâ, caule simplici hirsuto, foliis strictis angusto-linearibus pubescentibus, capitulis in spicam longam digestis confertim sessilibus, invol. squamis adpressis superne squarrosis.  $\mathcal{L}$  in pratis Illinoensibus (Michx.), montanis Virginîæ et Carolinæ (Pursh.) Hæc cl. Pursh refert ic. Dill. t. 72. f. 83. à cæteris adspicatam pertin. — Anonymos pilosus Walt. car. 197. Capitula minora quàm *L. spicata*. Cæter. ign.
11. *L. GRAMINIFOLIA* (Pursh fl. bor. am. 2. p. 508), radice tuberosâ, caule simplici glabro, foliis linearibus longissimis nervosis eciliatis margine scabriusculis costâ mediâ hispidis, capitulis remotiusculis ad axillam bractearum sessilibus in spicam digestis, invol. squamis oblongis obtusis submucronatis ciliatis adpressis, inter. coloratis, achænio villosulo, pappi setis longiusculè barbellatis.  $\mathcal{L}$  in pinetis humidis demissis Carolinæ Ell.! sketch 2. p. 274. Anonymos graminifolius Walt. car. 197. — Pluk. alm. t. 424. f. 6? In plantâ cultâ capitula majora et magis adproximata. (v. v. c. et s. ap.)
12. *L. CYLINDRACKA* (Mich. fl. bor. am. 2. p. 93), radice tuberosâ, tota hirsuta, caule gracili, foliis gramineis, capitulis cylindraceis paucifloris sessilibus distanter spicatis, squamis apice rotundatis abruptè mucronatis.  $\mathcal{L}$  in sylvis pratisque Illinoensibus. Cor. lobi intus hirsuti (Nutt.). — Species homonyma à cl. Pursh in Carolinâ indicata et à cl. Elliot sketch 2. p. 276. descr. videtur diversa sed mihi non satis nota.
13. *L. ASPERA* (Michx. fl. bor. am. 2. p. 92), radice....., caule subramoso scabro-pubescente, foliis lineari-lanceolatis asperimis, capitulis brevibus sessilibus spicatis distinctè alternis solitariis, invol. squamis rotundato-obtusis conniventibus.  $\mathcal{L}$  in pratis Illinoensium. Cæter. car. ign. — Anonymos ramosa

Walt. fl. car. 198 huc à cl. Pursh revocata videtur diversa et species in Carolinâ non indigena ex Ell. sketch 2. p. 276.

14. *L. HETEROPHYLLA* (Ait. kew. ed. 2. v. 4. p. 502), radice . . . , caule simplici glabro, foliis lanceolatis glabris lævibus, superioribus lineari-lanceolatis multoties minoribus, capitulis brevissimè pedunculatis spicato-racemosis, invol. subquarrosi squamis lanceolatis acutis nudis.  $\frac{1}{2}$  in Americâ bor. (Ait.), in Carolinâ australi et Georgiâ (Ell. sketch 2. p. 277). *L. varia* Banks ined. Cæt. ign.

15. *L. PILOSA* (Willd. sp. 3. p. 1636), radice . . . , caule simplici pubescente subpiloso, foliis linearibus pilosis ciliatis, capitulis longè pedicellatis in racemum laxum dispositis 10-14-floris, invol. squamis oblongis obtusis adpressis villosis, achænio multicostrato puberulo, pappi setis densè barbellatis.  $\frac{1}{2}$  in Amer. bor. à Georgiâ ad Nov. Cæsaream. Pursh fl. 2. p. 508. Ell. sketch 2. p. 277. Cor. lætè purpureæ. Species ab hortulanis sub hoc nomine admixta est *L. spicata* var.  $\gamma$ .

$\beta$ . *laevicaulis*, caule glabro, foliis basi remotè ciliatis, invol. squamis oblongis, capitalis 7-8-floris.  $\frac{1}{2}$  in Novâ Cæsareâ (Torr.!) *L. pilosa* Torr. herb. 1835. *L. gracilis* Pursh fl. bor. am. 2. p. 508. Var. aut spec. recognoscenda. (v. s.)

16. *L. GRACILIS* (Ell. sketch 2. p. 278), radice . . . , caule gracili pubescente subpiloso, foliis linearibus glabris basi subciliatis punctatis, capitulis longè pedicellatis laxè racemosis sub-7-floris, invol. squamis obovatis ciliatis adpressis, achænio villosis, pappi setis barbellatis.  $\frac{1}{2}$  in pinetis siccis Carolinæ (Ell.) et Georgiæ (h. Duby!). Cl. Nuttall ut mera var. *L. pilosa* hanc fortè non immeritò habet. *L. gracilis* Pursh et Lodd. bot. cab. t. 1909. fortè diversa? (v. s.)

17. *L. SECUNDA* (Ell. sketch 2. p. 278), radice tuberosâ, caule reclinato pubescente, foliis linearibus glabris basi parcè ciliatis, capitulis breviter pedicellatis racemosis secundis, invol. squamis lanceolatis acuminatis adpressis dorso punctatis.  $\frac{1}{2}$  in siccis et arenosis collibus Carolinæ, vulgaris prope Columbia (Ell.). Cor. pallidè purpureæ in invol. 6-8. Achænium villosum. Pappi setæ 2-3-seriales breviter et densè barbellatæ. (v. s.)

### Sect. III. TRILISA. — *Trilisa* Cass. dict. 55. p. 310.

Pappi setæ 1-3-seriales breviter et distanter barbellatæ. — Herbæ aut suffrutices. Radices à trunco subtuberoso fibrosæ. Capitula in corymbum paniculamve disposita.

#### \* *Herbaceæ.*

18. *L. PAUCIFLORA* (Pursh fl. bor. am. 2. p. 510), radice . . . , caule simplici glabro, foliis linearibus, paniculâ virgatâ foliosâ, ramis brevibus oligocephalis, capitulis subsessilibus secundis 3-5-floris, invol. squamis erectis lanceolatis acutis glabris. — in Georgiâ. Capitula magnit. *L. heterophyllæ*. Cæt. ign.

19. *L. ODORATISSIMA* (Willd. sp. 3. p. 1637), radice crassâ fibrosâ, caule simplici foliisque glaberrimis ovatis lanceolatisque nervosis subdenticulatis subglaucis, canlinis semi-amplexicaulis, paniculâ polycephalâ subcorymbosâ, capitulis pedicellatis 7-8-floris, invol. squamis oblongo-obovatis obtusis, achænio multicostrato vix puberulo, pappi setis parcè et breviter barbellatis.  $\frac{1}{2}$  in paludosis apriciis et pinetis demissis Virginis (Nutt.) Carolinæ et Floridæ. Michx. fl. 2. p. 93. Pursh fl. 2. p. 510. Nutt. gen. am. 2. p. 132. Andr. bot. rep. t. 633. *Trilisa odoratissima* Cass. dict. 55. p. 310. Anonymos odoratissimus Walt. car. 198. Cor. tubulosa, lobis brevibus. Pappus biserialis. Cor. lætè purpureæ. Folia trita vanillam ferè spirant. (v. s.)

20. *L. PANICULATA* (Willd. sp. 3. p. 1637), radice fibrosâ, caule simplici piloso-viscoso, foliis lanceolatis nervosis glabrescentibus, paniculâ coarctatè ramulis brevibus oligocephalis, capitulis sub-5-floris, invol. lanceolatis puberis erectis, achænio puberulo, pappi setis densè barbellatis.  $\frac{1}{2}$  in pinetis Virginis (Pursh fl. 510), Carolinæ (Fras. Bosc.), Floridæ (Pursh), Georgiæ (Duby!), Ell.

sketch 2. p. 283. *Anonymos paniculatus* Walt. car. 198. Variat cor. purpureis et albis. Capitula generis minora. Folia radic. multò majora 3-5-nervia basi attenuata. (v. s.)

21. *L. oppositifolia* (Nutt. in Sillim. Journ. 5. 1822. p. 299), pubescens, caule ramoso, foliis lanceolatis suboppositis nervosis, ramulis paniculato-corymbosis, capitulis cylindraceo-ovatis 15-20-floris, invol. squamis oblongis obtusis, infer. coloratis, pappi setis scabriusculis. ? in Louisiana circa Novam Aureliam. Capitula parva *L. paniculatae* sed numerosiora.
22. *L. corymbosa* (Nutt. gen. am. 132), radice fibrosa, caule solitario simplici hirsuto, foliis lanceolato-oblongis obtusis laeviusculis, super. brevioribus, corymbo partim simplici circ. 20-cephalo, pedicellis hirsutis subundis 1-5-cephalis, capitulis 20-floris, invol. squamis oblongo-ovalibus obtusis laevibus. ? in Carolinae bor. (Nutt.) et Georgiae (Ell.) humidis ad margines paludum. *L. tomentosa* Michx. fl. bor. 2. p. 93 ? Ell. sketch am. 2. p. 284. Cor. pallide purpureæ.
23. *L. Walpae* (Ell. sketch 2. p. 285), radice fibrosa, caule simplici superne piloso, foliis lanceolatis acutis glabris punctatis basi attenuatis, capitula paucis corymbosis multifloris, invol. squamis acutis tomentosis, schenio parè piloso, pappi setis subplumosis. ? in Carolina ad St. John's Berkeley (Ell.). *Anonymos uniflora* Walt. car. 198. Cor. intense purpureæ.
24. *L. bellidifolia* (Nutt. gen. am. 2. p. 133), demissa laxiuscula, radice tuberosa fibrosa, caulibus aggregatis, foliis radicalibus spatulato-lanceolatis obtusis 3-nerviis, caulinis sublinearibus, corymbi ramis subfastigiatis 1-5-cephalis, capitulis circ. 20-floris, invol. squamis oblongis obtusis, infer. subpetalis, pappi setis vix plumosis. ? in collibus arenosis Carolinae borealis. Cor. purpureæ.

\* \* *Fruticosa.*

25. *L. fruticosa* (Nutt. in Sillim. am. Journ. 5. 1822. p. 299), glabra fruticosa, ramis corymbosis superne nudis, foliis cuneato-obovatis punctatis, capitulis sub-5-floris, invol. squamis lineari-lanceolatis acuminatis. ? in Florida orientali. Dumetum ramosum. Folia sèrè Portulacæe oleraceæ. Invol. basi attenuatum stomis resinosis punctatum. Cor. purpureæ. Pappus subconcolor.

† *Species exclusæ.*

*L. Baicalensis* Adams. = *Carphephorus Baicalensis*.

*L. latifolia* Don. = *Ainalæa pteropoda*.

LXXXVIII. *CARPHEPHORUS* Cass. bull. philom. 1816. dict. 7. p. 149. Less. syn. 158.

Capitulum multiflorum. Receptaculum paleaceum !, paleis inv. squamas referentibus corollâ brevioribus deciduis. Invol. squamæ 3-seriales imbricatæ adpressæ. Achænium cylindraceum seu 10-costatum basi attenuatum. Pappus uni- aut pluriserialis, setis crassis plumosis aut grossè et irregulariter barbellatis. — Herbae perennes, aut Sibiricæ aut originis dubiæ, habitu *Liatridis*. Caulis simplex. Folia alterna linearia aut lanceolata integra aut dentata. Capitula corymbosa aut racemosa. Cor. purpureæ. — Species recognoscendæ.

1. *C. pseudo-liatris* (Cass. l. c), caule pubescente, foliis linearibus acutis amplexicaulibus integerrimis? adpressis, capitulis densè corymbosis, invol. squamis oblongis, achæniis 10-costatis hirtellis. ? pair 194. Cor. et styl. purpurei.
2. *C. Baicalensis*, caule infernè glabro superne subvillosa, foliis lanceolatis in petiolum attenuatis dentatis, racemo terminali folioso, pedunculis 1-cephalis nodis, invol. squamis ovato-lanceolatis acuminatis. ? in regione alpina ad meridiem lacus Baikal. *Liatris Baicalensis* Adams act. soc. h. n. Mosc. 6. (1817)

p. 115. Cor. violaceæ. Pappus plumosus 20-setus, setis basi subconcreta. Recept. paleaceum.

3. *C. REVOLUTIFOLIUS*, caule simplici suffruticoso basi glabro apice cum pedicellis puberulo, foliis linearibus glabrescentibus margine revolutis, ramulis axillaribus confertè foliosis, capitulis pedicellatis in racemum terminalem folio-  
rum elongatum dispositis, acheniis teretibus puberulis. ♀ in Mexico legit cl. Hænke. Folia pollicem longa, explicata 2 lin. lata. (v. s. in h. Hænke ab ill. de Sternberg comm.)

## LXXIX. DEGACHETA.

Capitulum multi- (20-) florum. Invol. ovatum imbricatum, squamis 3-4-seriatis substriatis caducis ext. ovatis, inter. oblongis obtusiusculis. Recept. subconvexum, paleis linearibus caducissimis inter flores onustum. Cor. glabræ. Antheræ inclusæ. Styli rami longè exserti. Achæmium acutè pentagonum glabrum, areolâ terminali convexâ. Pappus 1-serialis, setis 10 persistentibus basi vix scabris apice subclavellatis. — Suffruticulus erectus glaber Mexicanus. Rami teretes. Folia alterna subpetiolulata oblongo-lanceolata utrinquè acuminata serrata penninervia tenuiter reticulata. Thyrsus terminalis. Rami axillares apice capitula racemoso-congesta sessilia gerentes.

1. *D. HÆNKIANA*. ♀ in Mexico legit cl. Hænke. Folia 3-6 poll. longa, 10-12 lin. lata. (v. s. in h. Hænke ab ill. com. de Sternberg miss.)

## LXXX. CHROMOLÆNA.

Capitulum multiflorum. Recept. paleis paucis linearibus apice coloratis inter flores hinc inde onustum. Invol. squamæ innumere erectæ sed laxiusculæ (Lupuli more) imbricatæ oblongæ obtusæ subscariosæ apice coloratæ, venis subelevatis parallelis substriatis. Cor. tubulosæ ad faucem non ampliata. Antheræ appendicè ovatâ subpetaloideâ coloratâ donatæ. Styli rami crassi elongati glandulosi. Achæmium 5-gonum basi attenuatum glaberrimum. Pappus 1-serialis setosus scaber. — Herba perennis Brasiliensis tota capitulis exceptis hirsuto-tomentosa. Radices fibrosæ. Caules teretes. Folia opposita lanceolata breviter petiolata basi integrâ à medio ad apicem dentato-serrata. Capitula cymosa, ad apicem caulium, medio subsessili precociori, ramis ex hujus basi ortis mono- aut oligocephalis. Invol. (et corollæ?) latè roseo-violacea, formâ Lupulina capitula, colore Salvie Hormini bracteas, referentia. — Differt ab Eupatorio ut Carphæphorus à Liatride.

1. *C. NORMINOIDES*. 2/ in Brasiliæ prov. Minarum Generalium legit cl. Vauthier. Herba sesquipedalis basi simplex ramis paucis oppositis. (v. s. comm. à cl. Vauthier sub n. 296.)

LXXXI. OOCLINIUM. — *Præxels Cass. dict.* 10. p. 261? — Eupatorii sp. *ant.*

Capitulum multiflorum. Invol. squamæ biseriales aut sæpius pluri-seriales imbricatæ deciduæ. Recept. nudum post invol. collapsum ovatum! Achæmium angulatum. Pappus 1-serialis setosus scaber. — Herba sæpius suffruticosæ Americanæ. Folia opposita dentata aut integrâ. Pedunculi sæpius elongati 1-cephali. Cor. purpureæ. — Genus affine Eupatorio, sed receptaculo oviforme et etiam habitu diversum.

§. 1. *Species fruticosa aut perennes, foliis integris aut dentatis, corollis æqualibus.*

1. *O. GRANDIFLORUM*, collo suffruticuloso, caulibus herbaceis teretibus sparè hispido-setosis, foliis oppositis sessilibus lanceolatis integerrimis acuminatis hispido-setosis, ramis elongatis subnudis 1-cephalis, capitulis obconicis circ. 40-floris, invol. squamis imbricatis acuminatis glabris.  $\frac{3}{4}$  in Brasiliæ prov. Sancti-Pauli. Herba vix pedalis. Folia pollicem longa, 2-3 lin. lata. Capitula 6 lin. longa. Flosculi 5 lin. longi duplò nempe triplòve sequentibus majores, invol. æquales, ex sicco purpurei. Achænia exter. ad angulos scabra obcompressa. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 434 missa.)
2. *O. PAUCIDENTATUM*, collo suffruticoso, caule erecto basi sparè hispido apice glabro simpliciter aut trichotomo, ramis aphyllis apice 1-cephalis, foliis oppositis sessilibus lanceolatis ciliatis utrinquè glabris dentibus paucis acutis remotis suberratis, invol. ovali-subcylindracei glabri squamis adpressis ovalibus acutè mucronatis acuminatisve, exter. striatis, recept. ovato.  $\frac{1}{2}$  in Peraviâ legit olim cl. Hænke ex ill. de Bausa (sed in herb. Hænk. non reperire quivi). Medium inter grandiflorum et pedunculare. (v. s. in h. Acad. reg. Monac.)
3. *O. PEDUNCULARE*, suffruticulosum gracile diffusum adscendensve, caule tereti hispidulo, foliis oppositis linearibus utrinquè acutis subdentatis hispidulis, ramis elongatis subnudis apice 1-cephalis, capitulis 35-40-floris obovato-cylindraceis, invol. squamis imbricatis, ext. striatis acuminatissimis, inter. submembranaceis vix acutis, recept. ovato.  $\frac{3}{4}$  in Brasiliæ prov. Minas Generalium ad Sabara legit cl. Vauthier et hab. etiâ in prov. Sancti-Pauli (h. Mus. Paris!). Folia 1-1  $\frac{1}{2}$  poll. longa, 1-2 lin. lata. Rami floridi 3-4 poll. Invol. apice purpurascens. Achænia compresso-tetragona hispidiuscula. Cor. et styli purpurei. (v. s.)
4. *O. SIDERITIS*, caule herbaceo adscendente tereti sparè hispido oppositè ramoso, foliis oppositis distantibus ovato-lanceolatis sparè hispidulis basi cuneatis integerrimis trinerviis apice acuminatis dentatis, ramis elongatis apice 1-3-cephalis nudis, capitulis obovato-cylindraceis circ. 30-35-floris, invol. squamis 2-seriatis erectis glabris, exter. acutis, int. obtusis apice subscariosis, recept. ovato.  $\frac{1}{2}$ ?  $\odot$ ? circa Bahiam semel legit cl. Saltzman. Affine *E. pedunculari*. Habitus ferè *Stachydis Sideritis*. Radix fibrosa. Cor. ex sicco purpurascens. Capit. 4 lin. longa. (v. s. comm. à cl. Saltzman.)
5. *O. RIGIDUM*, suffruticulosum, caule erecto tereti striato glabro, foliis oppositis breviter petiolatis ovatis acuminatis remotè dentato-serratis trinerviis superè glabris subtus crebrè glanduloso-punctatis secus margines et nervos villosis, corymbo oligocephalo, capitulis longè pedunculatis 20-30-floris, invol. squamis acutè imbricatis deciduis striatis ext. obtusis, recept. ovato.  $\frac{3}{4}$  in calcareis montibus Jamaicæ borealis. *Eupatorium rigidum* Swartz! prod. III. flor. 1313. Caulis subherbacens. Folia pollicem longa, 6-7 lin. lata. Flores cærulei. Odor plantæ suaveolens, sapor terebinthinaceus. (v. s. ex h. Swartz.)

§. 2. *Species herbacea annua, foliis tripartitis, coroll. marginalium lobis exter. majoribus.*

6. *O. CAPILLARE*, herbaceum erectum glabrum tenue, foliis oppositis usquè ad basin in segmenta 3 lineari-filiformia partitis et inde falso-verticillatis, pedunculis elongatis laxis 1-cephalis, invol. oblongi squamis acutis, corollis radii palmatifidis majoribus, achæniis angulatis glabris. (I) in Brasiliæ prov. Minas Generalium ad Serra do Frio legit cl. Vauthier (pl. exa. n. 311). Herba ferè Galii facie ramosissima gracillima. Cor. latè violaceæ (v. s.)

† *Species dubia.*

7. *O. VILLOSUM*, herbaceum erectum, caule pilis articulatis sublanato, foliis oppositis petiolatis ovalibus serratis utrinquè petioloque villosissimis, capitulis

paucis subpaniculatis, invol. squamis glabris deciduis, recept. ovali seu conoideo, achæniis hispidulis. — in Guianâ Gallicâ legit cl. Poiteau. Præxelis villosa Cass. dict. 10. p. 262. An Ooclinii, an Conoclinii spec.?

LXXXII. CONOCLINIUM. — Eupatorii sp. auct.

Capitulum multiflorum. Invol. campanulati squamæ 2 3-seriales lineares acutæ subæquales. Recept. nudum conicum! Achænium angulatum. Pappus 1-serialis pilosus scaber. — Herbæ Americanæ aut suffrutices. Caules teretes striati. Folia opposita petiolata dentata. Corymbi conferti terminales. Cor. cœruleæ aut purpureæ. — Genus affine Eupatorio, sed recept. conico nec plano diversum sat videtur.

\* *Species herbaceæ.*

1. *C. COELESTINUM*, herbaceum, caule tereti pubescente, foliis oppositis petiolatis ovatis basi truncatis aut subcordatis apice acutis obtusè dentatis triplinerviis subacabris, corymbis confertis, invol. squamis circ. 30, 2-3-serialibus ferè subulatis, floribus in invol. circ. 40, recept. conico, achænio angulato glabro.  $\frac{1}{2}$  in pinguibus umbrosis montanis Virginie (Pursh), Carolinæ (Fras.! Bosc.), ad ripas fl. Ohio (Raf.). Eupatorium celestinum Linn. sp. 1174. Ell. sketch 2. p. 306. Steud. nom. phan. 322. excl. Cass. syn. E. deltifolium Raf. in litt. 1831. — Dill. elth. 140. t. 114. f. 139. Pluk. t. 394. f. 1. Cor. lætè cœruleæ parvæ odoratæ. Styli lobi longè exserti. (v. s.)
2. *C. BETONICUM*, herbaceum, caule tereti striatulo glabro apice pubescenti-velutino, foliis oppositis sinu rotundato cordatis oblongis crenatis vix subacutis supra scabridis subtus petiolisque pubescenti-villosis, capitulis in corymbis terminales simplices confertis 60-70-floris, invol. squamis circ. 25 biserialibus linearibus acuminatis scabridis, recept. conico, achænio angulato glabro.  $\frac{1}{2}$  in Mexico ad Tampico de Tamaulipas legit cl. Berlandier. Affine E. celestino. Cor. purpureæ. Folia exactè Betonicæ. (v. s.)
3. *C. BETONICIFORME*, herbaceum?, caule tereti cum petiolis pedicellisq. densè breviterque velutino, foliis oppositis petiolatis sinu rotundato cordatis oblongis crenatis acutis supra pulverulento-scabris subtus velutinis, capitulis in corymbis terminales confertos digestis pedicellatis circ. 40-floris, invol. squamis circ. 16 biserialibus linearibus acutis dorso subpuberis, recept. conico, achænio angulato glabro globulos resinosos gerente. — in Brasiliâ circa Rio-Janeiro legit cl. Lund. E. betonico simillimum. Petioli 4 nec 6-8 lin. longi. Superficies tota magis velutina. (v. s. comm. à cl. Lund.)

\*\* *Species basi suffruticosæ.*

4. *C. FRASIFOLIUM*, basi suffruticosum, caulibus erectis teretibus hispidis, foliis alternis rariùs oppositis petiolatis ovatis basi cordatis truncatisve grossè crenato-dentatis supra glabriusculis subtus pallidis villosis, capitulis in corymbum terminalem dense congestis pedicellatis circ. 20-floris, invol. squamis 10-12 subbiserialibus ovali-oblongis mucronatis dorso puberis, recept. elevato subconico, achænio angulato pilosiusculo.  $\frac{3}{4}$   $\frac{1}{2}$  in Brasiliâ circa Bahiam leger. cl. Blanchet, Salzmann et Lhotsky, et circa Caracas cl. Vargas. E. nepetoides Lindl.! herb. Petioli 3-9 lin. longi; limbus fol. 9-18 lin. longus, 6-9 lin. latus. Folia subtus nunc ferè cana nunc glabriuscula, foliis E. chamædrifolii sat similia, sed fructif. valdè diversa. Flores rosei odorati. (v. s.)
5. *C. MICROPHYLLUM* (Linn. suppl. 355 sub Eupatorio), suffruticosum adscendens, ramis teretibus subhirsutis, foliis oppositis petiolatis triangulari-ovatis obtusis crenatis margine revolutis supra glabris subtus albo-lanatis, corymbo terminali conglomerato, capitulis pedicellatis multifloris, invol. squamis subimbricatis lanceolatis ciliatis, recept. conico, achænio glabro.  $\frac{3}{4}$  in Novâ Granatâ. Crenæ fol. utrinquè 3-4. Petiolus ferè limbo æqualis nempe 3-4 lin. longus. Cor. purpureæ ex icone fortè 10-12, lobis puberis.

LXXXIII. HEBECLINIUM. — *Eupatorii* sp. auct. — *Coleosanthi* sp. Cass.

Capitulum multiflorum. Invol. campanulati squamæ pluriseries laxè subimbricatæ sæpè in appendicem subcoloratam productæ. Recept. elevatum superne plano-convexum pube brevi confertè hirsutum et ideò ferè piloso-frimbilliferum. Achænia angulata. Pappus 1-serialis pilosus scaber. — Herbæ australi-americanæ pubescentes. Caules teretes. Folia opposita petiolata cordata acuminata dentata. Corymbi terminales compositi conferti. Cor. albæ aut roseæ. — Genus affine *Eupario*, sed differt recept. villosa et invol. squamis sæpius appendiculatis.

\* *Involucri squamis exappendiculatis.*

1. *H. macrophyllum*, basi suffrutescens, caule tereti pubescente, ramis patulis, foliis oppositis longè petiolatis lato-cordatis acuminatis crenato-serratis trinerviis reticulatis suprà glabris subtus molliter pubescentibus, corymbo terminali composito polycephalo, capitalis ad apices ramulorum confertis ovatis 30-floris, invol. squamis laxè imbricatis linearibus obtusis, achæniis angulato glabro.  $\zeta$   $\eta$  in rudertis montium Jamaicae (Sw.), Guadalupe (Kraus!), Porto-Ricci (Wydlor!), Cubæ (Ossa), Guianæ (Aubl.), Brazilie (Lund!), agri Guyaquilensis (Hænke!), Peruviae (Poepp.), Novæ-Andalouciæ (H. B. K.). *Eupatorium macrophyllum* Linn. sp. 1175. Vahl symb. 3. p. 96. E. molle Swartz prod. 111. *E. macrophyllum* Sw. flor. 1303. *E. molle* et *macrophyllum* Willd. *Ageratum guianense* Aubl. guian. 800. *Coleosanthus tiliaefolius* Cass. dict. 24. p. 519. — Plum. et Burm. t. 129. Herba 4-5-pedalis. Flor. albi. (v. s.)

$\beta$ . *rivale*, frutescens, caule tereti glabro ramis divaricatis, foliis oppositis longè petiolatis lato-cordatis acuminatis dentato-serratis trinerviis setosis subtus vix pubescentibus, corymbo terminali conferto parvo, capitulis pedicellatis circ. 20-floris, invol. squamis oblongis glabriscaulis ferè obtusis, achæniis angulato glabro.  $\zeta$  in montosis juxta rivos Brazilie ad Rio-Janeiro legit cl. Lund. (v. s. in h. Mus. reg. Berol.)

\* \* *Involucri squamis omnibus aut plerisque in appendicem subcoloratam productis.*  
— *Urolepis*.

2. *H. hecatanthum*, caule herbaceo erecto subsimplici tereti striato superne præsertim cum petiolis et pediculis pube glandulosâ velutinis, foliis oppositis longè petiolatis cordatis brevè acuminatis acutè dentatis suprà sparse subtus secus nervos puberulis, corymbo terminali composito conferto, capitalis ad apices ramulorum plurimis sessilibus circ. 100-floris, invol. squamis pluriserialis striatis, intimis productis in appendicem linearem villosam subcorollinam albidam, achæniis glabro. ☉ in Brazilie prov. Rio-Grande. Herba pedalis sesquipedalisve. Petioli ferè 2 poll.; limbus 2  $\frac{1}{2}$  poll. longus, 2 poll. latus. Cor. albæ. Recept. hemisphæricum villosissimum. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 787 miss.)
2. *H. Urolepis*, caule herbaceo erecto tereti, petiolis pedunculisque pube brevi glandulosâ confertè velutinis, foliis oppositis petiolatis cordatis acutis dentatis utrinque glanduloso-puberulis, corymbo terminali oligocephalo, capitulis pedicellatis circ. 60-floris, invol. squamis 2-3-seriatis ferè omnino linearibus striatis puberulis productis in appendicem longam puberam roseo-coloratam, recept. hemisphærico villosa, achæniis glabro. — in Brazilie prov. Rio-Grande. Cor. roseæ. Valdè accedit ad *H. hecatanthum*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 807 et 803 miss.)

LXXXIV. CAMPULOCLINIUM. — *Eupatorii* sp. auct.

Capitulum multiflorum. Invol. campanulati squamæ 2-5-seriales inæquales, ext. ovales, inter. angustiores. Recept. convexum hemisphæ-

paucis subpaniculatis, invol. squamis glabris deciduis, recept. ovali seu conoideo, achæniis hispidulis. — in Guianâ Gallicâ legit cl. Poiteau. Præxelis villosa Cass. dict. 10. p. 262. An Oocliniæ, an Conocliniæ spec.?

## LXXXII. CONOCLINIUM. — Eupatorii sp. auct.

Capitulum multiflorum. Invol. campanulati squamæ 2 3-seriales lineares acutæ subæquales. Recept. nudum conicum! Achænium angulatum. Pappus 1-serialis pilosus scaber. — Herbæ Americane aut suffrutices. Caulis teretes striati. Folia opposita petiolata dentata. Corymbi conferti terminales. Cor. cœruleæ aut purpureæ. — Genus affine Eupatorio, sed recept. conico nec plano diversum sat videtur.

\* *Species herbaceæ.*

1. *C. COELESTINUM*, herbaceum, caule tereti pubescente, foliis oppositis petiolatis ovatis basi truncatis aut subcordatis apice acutis obtusè dentatis triplinerviis subacabris, corymbis confertis, invol. squamis circ. 30, 2-3-serialibus ferè subulatis, floribus in invol. circ. 40, recept. conico, achænio angulato glabro.  $\frac{1}{2}$  in pinguibus umbris montanis Virginie (Pursh), Carolinæ (Fras.! Bosc.!), ad ripas fl. Ohio (Raf.!). Eupatorium coelestinum Linn. sp. 1174. Ell. sketch 2. p. 306. Steud. nom. phan. 322. excl. Cass. syn. E. deltifolium Raf. in litt. 1831. — Dill. eth. 140. t. 114. f. 139. Pluk. t. 394. f. 1. Cor. lætè cœruleæ parvæ odoratæ. Styl. lobi longè exserti. (v. s.)
2. *C. BETONICUM*, herbaceum, caule tereti striatulo glabro apice pubescenti-velutino, foliis oppositis sinu rotundato cordatis oblongis crenatis vix subacutis suprâ scabridis subtus petiolisque pubescenti-villosis, capitulis in corymbis terminales simplices confertis 60-70-floris, invol. squamis circ. 25 biserialibus linearibus acuminatis scabridis, recept. conico, achænio angulato glabro.  $\frac{1}{2}$  in Mexico ad Tampico de Tamaulipas legit cl. Berlandier. Affine E. coelestino. Cor. purpureæ. Folia exactè Betonica. (v. s.)
3. *C. BETONICIFORME*, herbaceum?, caule tereti cum petiolis pediculisque densè breviterque velutino, foliis oppositis petiolatis sinu rotundato cordatis oblongis crenatis acutis suprâ pulverulento-scabris subtus velutinis, capitulis in corymbis terminales confertos digestis pedicellatis circ. 40-floris, invol. squamis circ. 16 biserialibus linearibus acutis dorso subpuberis, recept. conico, achænio angulato glabro globulos resinosos gerente. — in Brasiliâ circa Rio-Janeiro legit cl. Lund. E. betonico simillimum. Petioli 4 nec 6-8 lin. longi. Superficies tota magis velutina. (v. s. comm. à cl. Lund.)

\*\* *Species basi suffruticosæ.*

4. *C. FRASIFOLIUM*, basi suffruticosum, caulibus erectis teretibus hispidis, foliis alternis rarius oppositis petiolatis ovatis basi cordatis truncatisve grossè crenato-dentatis suprâ glabriusculis subtus pallidis villosis, capitulis in corymbum terminalem densè congestis pedicellatis circ. 20-floris, invol. squamis 10-12 subbiserialibus ovali-oblongis mucronatis dorso puberis, recept. elevato subconico, achænio angulato pilosiusculo.  $\frac{3}{4}$  in Brasiliâ circa Bahiam legit. cl. Blanchet, Salzmann et Lhotsky, et circa Caracas cl. Vargas. E. nepetoides Lindl.! herb. Petioli 3-9 lin. longi; limbus fol. 9-18 lin. longus, 6-9 lin. latus. Folia subtus nunc ~~ferè~~ cana nunc glabriuscula, foliis E. chamædrifolii sat similia, sed fructif. valdè diversa. Flores rosei odorati. (v. s.)
5. *C. MICROPHYLLUM* (Linn. suppl. 355 sub Eupatorio), suffruticosum adscendens, ramis teretibus subhirsutis, foliis oppositis petiolatis triangulari-ovatis obtusis crenatis margine revolutis suprâ glabris subtus albo-lanatis, corymbo terminali conglomerato, capitulis pedicellatis multifloris, invol. squamis subimbricatis lanceolatis ciliatis, recept. conico, achænio glabro.  $\frac{3}{4}$  in Novâ Granatâ. Crenæ fol. utrinquè 3-4. Petiolus ferè limbo æqualis nempè 3-4 lin. longus. Cor. purpureæ ex icone fortè 10-12, lobis puberis.



## LXXXV. BULBOSTYLIS. — Coleosanthi sp. Cass. — Eupatorii sp. auct.

Capitulum 10-25-florum. Invol. squamæ lineari-lanceolatæ acuminate striatæ laxè imbricatæ 2-3-seriales. Recept. angustum epaleaceum. Cor. tubulosæ basi dilatatæ brevissimè 5-dentatæ. Stylus basi bulbosus. Achenium teres striatum sessile. Pappus 1-serialis setosus scaber. — Suffrutices Mexicani aut rarius austr.-Americani. Rami teretes. Folia opposita aut superiora alterna, petiolata, lanceolata, aut ovata serrata, pubera aut glabra, subtus sæpè glandulosa nunquam pellucido-punctata. Panicula subthyrsoidea foliosa. Capitula oblonga.

1. *B. CAVANILLESII*, suffruticosa erecta, caule tereti villosulo-velutino, foliis oppositis (summis alternis) petiolatis ovato-lanceolatis subacutis crenato-dentatis suprè sparè scabridis subtus trinerviis reticulatis glandulosus nervis petiolisque pubescentibus ramulis axillaribus sæpius 1-3-cephalis in paniculam elongatam foliosis dispositis, capitulis 20-25-floris, invol. squamis lanceolatis acuminatis striatis laxè imbricatis erectiusculis (non verè squarrosis), achenio tereti striato pubero.  $\frac{3}{4}$  in Mexico ut vid. non infrequens et in Californiâ (Douggl.). Eupatorium axillare fl. mex. ic. ined. Eupatorium squarrosum Cav. ic. 1. p. 66. t. 98. Coleosanthus Cavanillesii Cass. dict. 10. p. 37. Specim. habeo ex herb. Cavan. proveniens sed à descr. cl. auct. subrecedens foliis basi truncatis nec cordatis, floribus imò fructibus vix involucreo longioribus, etc. Cor. ochroleucæ basi tumidæ apice brevissimè 5-dentatæ. Stylus basi dilatatus et hispidus. (v. s.)
2. *B. SECUNDIFLORA*, suffruticosa erecta, caule tereti pubescenti-villosulo, foliis alternis et oppositis petiolatis subcordato-ovatis acuminatis dentatis trinerviis utrinquè parè pubescenti-hirtellis subtus glandulosis, ramis floridis racemosis in paniculam foliosam secundam dispositis, pedicellis glanduloso-villosis, capitulis nutantibus circ. 20-floris, invol. squamis imbricatis lanceolatis striatis acuminatis, achenio teretiusculo striato puberulo.  $\frac{3}{4}$  in Mexico (Lag.) inter Tula et Tampico (Berlandier!) Eup. nutans H. B. et Kunth nov. gen. sm. f. p. 105? Eup. secundiflorum Lag. nov. gen. 1814. p. 25. n. 334. Cor. ochroleucæ. Species media inter priorem et sequentem. (v. s.)
3. *B. PENDULA*, suffrutescens erecta, caule tereti glabriusculo, foliis oppositis (vix summis alternis) petiolatis ovato-lanceolatis basi cuneatis apice acuminatis dentatis triplinerviis tenuiter pubescenti-hirtellis subtus crebrè glandulosis, paniculâ racemosâ secundâ, pedicellis glanduloso-pilosis, capitulis pendulis 11-15-floris, invol. squamis imbricatis lanceolatis acuminatis striatis, achenio tereti striato puberulo.  $\frac{3}{4}$  in Mexico. Enpat. pendulum Schrad! cat. sem. h. Gott. 1830. pl. rar. descr. et ic. ined. 1832. E. scariosum Wallr. in litt. 1834. Differt à duabus præced. florum in invol. numero minore, foliis tripli- nec trinerviis subtus pilos pauciores glandulas numerosiores gerentibus et basi cuneatis. Cor. flavo-virides basi vix dilatata. An hæ tres species in unicam formam colligendæ? (v. s. cult. h. Götting. et à cl. Schrad. comm. cum ic. exim.)
4. *B. REBECARPA*, suffruticosa ramosissima, caulibus ramis pedunculis petiolisque pubescenti-hirtellis, foliis oppositis breviter petiolatis ovatis utrinquè subacutis dentato-serratis suprè scabridis subtus trinerviis reticulatis villosi-hirtellis, ramulis axillaribus 1-cephalis, capitulis circ. 10-floris, invol. squamis laxè imbricatis subacutis striatis pubescentibus, ext. ovatis, inter. oblongo-linearibus, achenio teretiusculo striato pubescenti-villosulo.  $\frac{3}{4}$  in Mexico legit cl. Hænke. Cor. tubulosa brevissimè 5-dentata. Stylus basi bulbosus et villosus. Folia (incl. petiolo 2 lin.) 15 lin. longa, 6 lin. lata. (v. s. in h. Hænk. ex Mexico et comm. à cl. Née sine loci design.)
5. *B. PEDUNCULOSA*, fruticosa ramosissima subcespitosa, ramulis teretibus pulverulento-velutinis apice longè in pedicellum 1-cephalum denudatis, foliis oppositis (et summis alternis) breviter petiolatis ovatis dentibus paucis crenatis pube minimâ utrinquè cinereis subtus glandulosis, capitulis circ. 25-floris, invol. squamis laxè imbricatis striatis mucronato-acuminatis, achenio ter-

salo striato glabro.  $\frac{1}{2}$  in Mexico circa Villapando legit cl. Mendez. Suffrux vix pedalis tortuosus. Cor. ex sicco albæ basi tumidulæ. Pedunculus  $\frac{1}{2}$  lin., quæ notâ facili ab *E. veronicæ* folio differt. Fol. cum petiolo 1 lin., limbus longa. (v. s.)

*VERONICÆFOLIA*, fruticosa ramosissima, ramulis tenuissimè pulverulentis, foliis infer. ternis, cæteris oppositis, brevissimè petiolatis ovato-subrotundis obtusis subcordato-reniformibus crenatis tenuissimè pubescenti-canescentibus, capitulis ad apices ramorum solitariis breviter supra ultimum jugum sitis circ. 15-20-floris, invol. squamis laxè imbricatis striatis mucronato-acuminatis, achanio glabro.  $\frac{1}{2}$  in collibus Mexici præpe sacellum Guadalupe. Esp. glechomæfolium Moq. fl. mex. ic. ined. Eupat. veronicæfolium H. et Kunth nov. gen. am. 4. p. 112. t. 441. Cor. alba. Suffrutex pedalis ad pedunculosam accedens, sed pedunculus vix invol. longitudine.

*NEPETÆFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 112), fruticosa, ramis nudo-striatis hispidis-hirtis, foliis alternis petiolatis triangulari-ovatis acutis serratis crenatis trinerviis rigidiusculis canescentibus supra scabris subtus glaucescentibus tomentosis, capitulis axillaribus et terminalibus subspicatis subsessilibus 20-25-floris, invol. squamis imbricatis striatis linearilanceolatis, ext.  $\frac{1}{2}$  lin.  $\frac{1}{2}$  in exsiccatis montis Sanctæ Rosæ Mexici. Eupat. nepetæfolium H. B. et Kunth nov. gen. am. 4. p. 112. Cor. alba. Petiol. 2 lin., limbus pollicem long., 3-9 lin. latus. An fortè B. Cavanillesii affine? sed de stylo et corollâ t. cl. auct.

*OLIGANTHUS*, herbacea erecta, foliis oppositis alternisve breviter petiolatis sub-striatis hispidis-hirtis pubescentibus, capitulis solitariis paucisve cymosis axillaribus multifloris, invol. turbinatis squamis locis multinerviis ext. subnullis ciliatis, intim. linearibus elongatis. — in collibus apricis Mexici prope elunda de la Laguna. Eupat. oliganthus Less. in linnæa 1830. p. 137. t. 2.  $\frac{1}{2}$  Cor. ut in *B. nepetæ* foliâ cylindricæ à typo generis recedunt, sed cor. non inflata ex icon. cit. Cæter. tamen ad *Bulbostylis* accedunt. Forma fol. vix omnes.

*GRASSA*, herbacea glabra, ramis teretibus, foliis oppositis petiolatis ovali-angulatis acuminatissimis remotè serratis triplinerviis, paniculâ laxâ subcoarctatâ, capitulis longè pedicellatis erectis 16-floris, invol. squamis laxè imbricatis lanceolatis acuminatis striatis, ext. multò brevioribus, achanio angulato, pappo puberulo. — Patr. ign. Petioli 8-9 lin. longi; fol. limbus 5-6 poll. long., 1  $\frac{1}{2}$ -2 latius. Capit. 5-6 lin. longa. (v. s. comm. à cl. Gaudichaud.)

*SPINACLEFOLIA*, fruticosa glabra, ramis teretibus, foliis oppositis longè hastato-hastato-triangularibus lobis apiceque acuminatis ad margines grosse serratis dentatis triplinerviis punctulatis, paniculis foliosis, pedunculis trifidis, foliis longitudine, capitulis pedicellatis 10-12-floris, invol. squamis à imbricatis oblongis obtusis substriatis, achanio nervis exsertis 5-costato, rima lateraliter hispidis.  $\frac{1}{2}$  in Mexico inter Victoria et Tula legit cl. Berlandier. Petioli 2 poll. longi. Limbus basi breviter cuneatus 3 poll. longus, 2 poll. am. Cor. teres albida. (v. s.)

*PAUCIFLORA*, herbacea erecta hispido-pilosa, ramis teretibus, foliis ovatis petiolatis ovatis acutis serratis basi cuneatis et integerrimis reticulatis-triplinerviis, capitulis axill. et termin. solitariis longè pedunculatis multifloris, invol. compresso-striatis squamis subimbricatis lanceolatis acuminatis striatis subnullis, achanio ad angulos scabro.  $\frac{1}{2}$  in Novâ Granatâ? Eupat. pauciflorum H. B. et Kunth nov. gen. am. 4. p. 120. Cor. violaceæ? Recept. indescr. lamina compressa quadrangulari ex cl. auct. Petioli 2 lin. longi. Limbus hispidus, 7 lin. latus.

LXXXV. BULBOSTYLIS. — Coleosanthi sp. Cass. — Eupatorii sp. auct.

Capitulum 10-25-florum. Invol. squamæ lineari-lanceolatæ acuminatæ striatæ laxè imbricatæ 2-3-seriales. Recept. angustum epaleaceum. Cor. tubulosæ basi dilatatæ brevissimè 5-dentatæ. Stylus basi bulbosus. Achænium teres striatum sessile. Pappus 1-serialis setosus scaber. — Suffrutices Mexicani aut rariùs austr.-Americani. Rami teretes. Folia opposita aut superiora alterna, petiolata, lanceolata, aut ovata serrata, pubera aut glabra, subtùs sæpè glandulosa nunquàm pellucido-punctata. Panicula subthyrsoidea foliosa. Capitula oblonga.

1. B. CAVANILLESII, suffruticosa erecta, caule tereti villosulo-velutino, foliis oppositis (summis alternis) petiolatis ovato-lanceolatis subacutis crenato-dentatis suprâ sparsè scabridis subtùs trinerviis reticulatis glandulosis nervis petiolicque pubescentibus ramulis axillaribus sæpiùs 1-3-cephalis in paniculam elongatam foliosam dispositis, capitulis 20-25-floris, invol. squamis lanceolatis acuminatis striatis laxè imbricatis erectiusculis (non verè squarrosis), achænio tereti striato pubero.  $\frac{3}{4}$  in Mexico ut vid. non infrequens et in Californiâ (Dougl.). Eupatorium axillare fl. mex. ic. ined. Eupatorium squarrosum Cav. ic. 1. p. 66. t. 98. Coleosanthus Cavanillesii Cass. dict. 10. p. 37. Specim. habeo ex herb. Cavan. proveniens sed à descr. cl. auct. subrecedens foliis basi truncatis nec cordatis, floribus imò fructibus vix involuero longioribus, etc. Cor. ochroleucæ basi tumidæ apice brevissimè 5-dentatæ. Stylus basi dilatatus et hispidus. (v. s.)
2. B. SECUNDIFLORA, suffruticosa erecta, caule tereti pubescenti-villosulo, foliis alternis et oppositis petiolatis subcordato-ovatis acuminatis dentatis trinerviis utrinquè parè pubescenti-hirtellis subtùs glandulosis, ramis floridis racemosis in paniculam foliosam secundam dispositis, pedicellis glanduloso-villosis, capitulis nutantibus circ. 20-floris, invol. squamis imbricatis lanceolatis striatis acuminatis, achænio teretiusculo striato puberulo.  $\frac{3}{4}$  in Mexico (Lag.) inter Tula et Tampico (Berlandier!) Eup. nutans H. B. et Kunth nov. gen. am. 4. p. 105? Eup. secundiflorum Lag. nov. gen. 1814. p. 25. n. 334. Cor. ochroleucæ. Species media inter priorem et sequentem. (v. s.)
3. B. PENDULA, suffrutescens erecta, caule tereti glabriusculo, foliis oppositis (vix summis alternis) petiolatis ovato-lanceolatis basi cuneatis apice acuminatis dentatis triplinerviis tenuiter pubescenti-hirtellis subtùs crebrè glandulosis, paniculâ racemosâ secundâ, pedicellis glanduloso-pilosis, capitalis pendula 11-15-floris, invol. squamis imbricatis lanceolatis acuminatis striatis, achænio tereti striato puberulo.  $\frac{3}{4}$  in Mexico. Eupat. pendulum Schrad! cat. sem. h. Gott. 1830. pl. rar. descr. et ic. ined. 1832. E. scariosum Wallr. in litt. 1834. Differt à duabus præced. florum in invol. numero minore, foliis triplinerviis subtùs pilos pauciores glandulas numerosiores gerentibus et basi cuneatis. Cor. flavo-virides basi vix dilatæ. An hæ tres species in unicam formam colligendæ? (v. s. cult. h. Götting. et à cl. Schrad. comm. cum ic. exim.)
4. B. HEBECARPA, suffruticosa ramosissima, caulibus ramis pedunculis petiolicque pubescenti-hirtellis, foliis oppositis breviter petiolatis ovatis utrinquè subacutis dentato-serratis suprâ scabridis subtùs trinerviis reticulatis villosulo-hirtellis, ramulis axillaribus 1-cephalis, capitulis circ. 10-floris, invol. squamis laxè imbricatis subacutis striatis pubescentibus, ext. ovatis, inter. oblongo-linearibus, achænio teretiusculo striato pubescenti-villosulo.  $\frac{3}{4}$  in Mexico legit cl. Hænke. Cor. tubulosa brevissimè 5-dentata. Stylus basi bulbosus et villosus. Folia (incl. petiolo 2 lin.) 15 lin. longa, 6 lin. lata. (v. s. in h. Hænke ex Mexico et comm. à cl. Née sine loci design.)
5. B. PEDUNCULOSA, fruticosa ramosissima subcæspitosa, ramulis teretibus puberulento-velutinis apice longè in pedicellum 1-cephalum denudatis, foliis oppositis (et summis alternis) breviter petiolatis ovatis dentibus paucis crenatis pube minimâ utrinquè cinereis subtùs glandulosis, capitulis circ. 25-floris, invol. squamis laxè imbricatis striatis mucronato-acuminatis, achænio ter-

Sapè cum C. Dalea mixta, sed præst. notas ind. differt odore ferè nullo, corymbo magis divaricato, foliis non nitidis, etc.

5. C. DALEOIDES, fruticosa, ramulis pedunculis foliorumque petiollis nervisque utrinque villis rufis velutino-subhirtis, foliis oppositis breviter petiolatis elliptico-oblongis utrinque acuminatis penninerviis hasi integris cæterum serratis punctis pellucidis crebris liturisque paucissimis pertusis, paniculâ divaricatâ, pedunculis apices capitula 3-5 glabra straminea 5-flora gerentibus, invol. oblongi squamis imbricatis 3-4 seriatis vix acutis.  $\zeta$  ad Tampico de Tamaulipas in Mexico legit cl. Berlandier. Affinis C. Dalea sed faciliè distincta. (v. s.)
6. C. HESPEROTATA, fruticosa, ramis subangulatis glabris, ramulis panicula velutino-villosis, foliis oppositis petiolatis ellipticis acuminatis minutè et rariter serratis penninerviis punctis pellucidis rotundis et paucis oblongis pertusis, paniculâ ramulosissimâ, capitulis ad apices ramulorum confertis breviter pedicellatis 5-floris, invol. oblongi squamis stramineis obtusiusculis striatis, achenio 5-costato inter costas hispidulo.  $\zeta$  in Mexico legit cl. Hænke. Petioli pollicares. Limbus fol. 5 poll. longus, 2-2  $\frac{1}{2}$  latus. (v. s. in h. Hænke. à cl. Sternb. comm.)

LXXXVII. EUPATORIUM Tourn. inst. 455. t. 259. Linn. gen. n. 935. Gærtn. fr. 2. p. 401. t. 166: Adans. fam. 2. p. 123. Cass. dict. 16. p. 2. 25. p. 432. 26. p. 228. Less. syn. 157. excl. hinc indè sp.

Capitulum multi (3-100)-florum. Recept. planum nudum. Invol. squamæ uni- bi- aut pluriseriales æquales aut inæquales laxè aut strictè imbricatæ. Cor. faux vix dilatata. Antheræ inclusæ. Styli rami exserti cylindracei obtusi. Achæmium angulatum striatumve. Pappus 1-serialis pilosus scaber. — Herba aut suffrutices. Species paucæ gerontogæ, innumerae, Americanae. Folia sæpius opposita, rarè alterna aut verticillata. Capitula sæpius corymbosa aut paniculata. Cor. purpureæ cœruleæ, albæ aut albidæ uno verbo cyanicæ, nunquam flavæ seu xanthinæ. — Genus vastissimum, etiam post priorum et sequentis separationem species ferè innumerosas continens, fortè quoad specierum in sectionibus seriem recognoscendum.

Ser. I. IMBRICATA. *Nempè capitula squamis pluribus serialibus imbricatis, est. brevibus, inter. sensim elongatis constantia.*

### §. 1. CYLINDROCEPHALA.

Capitula cylindrica 5-60-flora. Invol. squamæ pluriseriales arcuè imbricatæ obtusæ intensè striatæ.

\* Capitulis 20-60-floris, foliis oppositis.

1. E. MULTIFLOECULOSUM, fruticosum, ramis teretibus hirsutis, foliis oppositis brevissimè petiolatis ovatis subcrenatis basi subtrinerviis subtus nervulosis utrinque glabrescentibus subtus multiglandulosis, capitulis ad apices ramulorum inter duo ultima folia subsolitariis sessilibus ovato-cylindricis 60-65-floris, invol. squamis arcuè imbricatis obtusis ciliatis substriatis.  $\zeta$  in Brasiliâ. Affine E. aqualido sed faciliè ab eo et ab omnibus hujus sect. differt floeculis in invol. numerosissimis, capitulis solitariis, etc. (v. s.)
2. E. MACRANTHUM (Swartz fl. ind. occ. 1315), fruticosum glaberrimum, ramis angulato-sulcatis, foliis oppositis petiolatis ovatis acuminatis cartilagineo-serratis trinerviis subtus granulis pellucidis adspersis, corymbo terminali decomposito, pedicellis angulatis, capitulis cylindricis 30-40-floris, invol. squamis arcuè imbricatis obtusis.  $\zeta$  in Martinicâ. Spreng. syst. 3. p. 410. exclud. syn. Petioli semunciales. Capitula semipoll. longa. Cor. albidæ.

LXXXVI. CRITONIA *P. Brown. jam.* (1756) p. 490. non *Gærtn. nec Cass.* (1) — *Dalea P. Brown. p.* 314. non *Linn.* — *Wikstrœmia Spreng. syst.* 3. (1826) non *Schrad. nec Endl.* (2). — *Eupatorii sp. auct.*

Capitulum 3-5-florum. Invol. oblongi squamæ 2-3-seriales imbricatæ oblongæ striatæ obtusæ. Recept. angustum nudum. Stylus basi non bulbosus. Achænia acutè pentagona basi attenuato-stipitata inter costas hispidula aut glandulifera. Pappus 1-serialis setosus scaber. — Frutices Caribæi et Mexicani glabri elegantes. Rami teretes striati. Folia opposita petiolata ovalia utrinquè acuminata subserrata membranacea punctis glanduloso-pellucidis ut in *Samydâ* sæpius oblongis perforata tria suaveolentia. Corymbi compositi. Capitula straminea ad apices ramulorum corymbi congesta. — Genus characteribus levibus sed habitu distinctissimo insignitum.

1. *C. DALEA*, fruticosa glaberrima, foliis oppositis petiolatis lanceolatis utrinquè sensim acuminatis subserratis penninerviis punctis liturisque pellucidis notatis, paniculâ subcorymbosâ, capitulis ad apices ramulorum ternis fasciculatis oblongis 4- (rariùs 3-) floris, invol. squamis 7-8 oblongis imbricatis substriatis, achænio acutè 5-gono hispidulo et glandulifero. ♂ in frutetis Jamaicæ (nec alibi ritè cognitum!) *Eupatorium Dalea Linn. sp.* 171. *Swartz obs.* 298. *Jacq.!* h. *Sehcn.* 2. p. 10. t. 146. *Isocarpha Berterii Balb.!* herb. et ex eo *Calydemon Jamaicensis Spreng. syst.* 3. p. 457. excl. syn. — *Dalea sen Critonia P. Brown. jam.* p. 314 et 490. t. 34. f. 1. Odor totius fruticis gratissimus etiam in statu sicco diù permanens. Petioli 4 lin. longi. Limbus fol. 3 poll. longus, pollicem latus. (v. s.)
2. *C. PSEUDO-DALEA*, fruticosa glaberrima, foliis petiolatis lanceolatis utrinquè sensim acuminatis serratis penninerviis punctis liturisque pellucidis notatis, paniculâ subcorymbosâ, capitulis ad apices ramorum quinis fasciculatis oblongis 5-floris, invol. squamis 7-8 oblongis imbricatis substriatis, achænio acutè pentagono hispidulo eglanduloso. ♂ in frutetis insulæ Cubæ circa *Havanam*. *Eup. Dalea H. B. et Kunth nov. gen. am.* 4. p. 106. excl. syn. Valdè accessit ad *C. Daleam* sed præter notas ind. differt paniculâ magis confertâ. De odore fruticis silent autoptæ. Petiolus 3-4 lin. limbus 4 poll. longus, 12-15 lin. latus. (v. s. comm. à cl. Ramon de la Sagra.)
3. *C. MACROPODA*, fruticosa glaberrima, foliis oppositis longè petiolatis ovato-lanceolatis basi breviter attenuatis apice subabruptè et longè acuminatis penninerviis serratis punctis liturisque pellucidis notatis, paniculâ corymbosâ, capitulis ad apices ramulorum 3-5 breviter pedicellatis oblongis 5-floris, invol. squamis imbricatis oblongis acutiusculis, achænio acutè pentagono hispidulo eglanduloso. ♂ in insulâ Trinitatis *Sieb. fl. trinit. exs. n.* 184. Affine *C. Daleæ* et præsertim pseudo-daleæ, sed ab utrâque differt petiolo pollicari et idèò triplo longiori, foliis 5-7 poll. longis, 2 poll. latis magis ovatis et abruptius acuminatis, capitulis pedicellulatis, etc. (v. s. comm. à cl. *Sieb.*)
4. *C. PARVIFLORA*, fruticosum glabrum, ramulis nervisque fol. subitè subpærescentibus, foliis oppositis longè petiolatis ovato-lanceolatis acuminatis serratis penninerviis punctis liturisque notatis, paniculâ corymbosâ, capitulis ad apices ramulorum confertis sessilibus 3-floris, invol. squamis 6-8, inter. obtusis, achænio angulato. ♂ inter fruteta in montibus excoctis Jamaicæ australi *Eupator. parviflorum Swartz prod. p.* 111. fl. ind. oce. ♂. p. 1299. non *Aubl.*

(1) *Critonia* *Gærtn.* (1791) est *Kuhniz* sp. *Critonia* *Cass. dict.* v. 12 (1818) p. est *Decaneuri* sp.

(2) *Wikstrœmia* *Schrad.* (1821) est genus *Tiliaceis* affine. *Vid. prod.* 1. p. 51. *Wikstrœmia* *Endl.* (1833) est genus *Thymelæarum*.

10. *E. FLORIBUNDUM* (H. B. et Kunth nov. gen. am. 4. p. 118. t. 344), fruticosum, ramis sexangularibus sulcato-striatis glabriusculis, foliis oppositis petiolatis ovatis angustato-acuminatis leviter serratis supra glabriusculis subtus molliter tomentoso-pubescentibus, corymbis ramosissimis diffusis, capitulis longè pedicellatis circ. 25-floris, invol. cylindracei squamis arcetè imbricatis striatis obtusis glabris, achenio glabro. ♂ in temperatis versus Loxam Peruvianorum. Cor. violaceæ. Petioli 3-6 lin. Limbus fol. 3 poll. longus, 2  $\frac{1}{4}$  poll. latus.
11. *E. SERRATULOIDES* (H. B. et Kunth nov. gen. am. 4. p. 117), caule herbaceo glabriusculo oppositè ramoso, ramis apice furcatis oligocephalis, foliis oppositis petiolatis ovatis acutis serratis triplinerviis supra scabriusculis subtus glabris margine ciliato-scabris, capitulis solitariis pedunculatis multifloris, invol. subcylindracei squamis arcetè imbricatis obtusis striatis ciliatis, achenio glabro. ♀ in præruptis juxta Guancabamba Peruvianorum. Cor. violaceæ. Petioli 3 lin. Fol. limbus 13-14 lin. longus, 7 lin. latus.
12. *E. MAXIMILIANI* (Schrad. in litt. 1832), fruticosum, ramis virgatis teretibus apice tenuissimè pubescentibus, foliis oppositis petiolatis ovato-rhombeis acuminatis per totam ferè longitudinem grossè serratis trinerviis utrinquè glabriusculis, corymbo composito terminali, capitulis longè pedicellatis breviter cylindraceis 35-40-floris, invol. squamis arcetè imbricatis obtusis. ♂ in Brasiliâ circa Rio-Janeiro inter sepes primus detexit Princ. Maxim. Neovidensis. Subascandens. Rami oppositi patentes. Flores pallidissimè violacei aut nivei. Achenia glabra. (v. s. comm. à cl. Schrad. ex hort. Gœtt. et sp. à cl. Lund.)  
*β. hispidulum*, foliis utriusquè pilis sparsis hispidulis. ♂ in Brasiliâ ad Rio-Janeiro et Bahiâ (Blanchet!), in Guianâ Gallicâ (Perr.). Flores in invol. 36 reperi. (v. s.)
13. *E. ODORATUM* (Linn. sp. 1174. amœn. 5. p. 405), fruticosum, ramis angulo recto divaricatis pubescenti-villosis, foliis oppositis petiolatis triangulari-ovatis apice integris acuminatis basi grossè dentato-serratis triplinerviis subtus densè pubescentibus punctatis corymbis ramos terminantibus, capitulis pedicellatis cylindricis 20-floris, invol. squamis arcetè imbricatis plerisque acuminatis. ♂ in ruderatis Jamaicæ. (Br. Linn. Sw.), Sancti Domingi (h. Lher.), Guadalupæ ubi vulgò dicitur *langue-de-chat* (Rad. ! Krauss ! ) Martinicæ (h. Doby ! ), Guianæ (h. Lindl. ! ). Swartz obs. 300. — Pluk. alm. t. 177. f. 3? Flores albi suavè odorati ! Floret decemb. Fortè à plurimis cum sequentibus confusum (v. s.)  
*β. Cubense*, invol. squamis striatis plerisque obtusis. — in insulâ Cubâ prope Havanam leg. cl. de la Ossa et R. de la Sagra. Fl. in invol. circ. 18. (v. s.)  
*γ. brachiatum*, invol. squamis plerisque acuminatis, foliis subtus parcius pubescentibus. ♂ in Guadalupâ (Badier) E. atriplicifolium Hort. Par. Lam. dict. 2. p. 407. excl. Plum. syn. E. brachiatum Wikstr. fl. guad. p. 71. (v. s. ex hort. Par. tempore Lam. lectum et sp. sicc. à cl. Badier comm.)
14. *E. COMYZOIDES* (Vahl symb. 3. p. 96), suffruticosum, ramis angulo ferè recto divaricatis pubescentibus, foliis oppositis petiolatis latè ovatis acuminatis basi grossè dentato-serratis triplinerviis apice integris supra subpuberulis subtus densè pubescentibus et punctulatis, corymbis terminalibus compositis, capitulis pedicellatis cylindricis circ. 20-floris, invol. squamis arcetè imbricatis striatis obtusis. ♂ in umbrosis Brasiliæ circ. Bahiam et in prov. Sancti-Pauli, in ins. Cubâ, et in Mexico inter Santander et San. Fernando. Schrank pl. rar. h. mon. t. 85. Affine odorato sed certè dist. Cor. pallidè cœroleæ ex Berland. purpurascentes ex Saltzm. sed cæt. nullum discrimen video inter specim. Mexicana et Brasiliensia. Florum vix fragrantium numerus in invol. variat à 15 ad 27. (v. s.)
15. *E. DIVERGENS* (Less. in linnæâ 1830. p. 138), herbaceum, ramis divergentibus, foliis oppositis breviter petiolatis ovato-lanceolatis longè acuminatis triplinerviis subcrenato-serratis sparsè pilosiusculis, paniculis parvis oligoce-

- phalis fastigiatis, capitulis 20-25-floris, invol. cylindraceis squamis erectis obtusissimis glabris striatis apice subciliatis. — in Mexico prope Hacienda de la Laguna. Cor. lilacin. Affine *E. conyzoidi*.
16. *E. CLEMATIDIS*, fruticosum, ramis teretibus striatis glabrisculis, ramulis angulo recto divergentibus, foliis oppositis petiolatis triangularibus basi truncatis apice acuminatis subintegerrimis subtrinerviis glabris, corymbo paniculato laxo oligocephalo, capitulis longè pedicellatis cylindricis 25-floris, invol. squamis arcè imbricatis obtusis.  $\zeta$  in Peruvia? legit cl. Poeppig. Petioli 8-10 lin. longi. Limbi vix prope basin 1-dentatis. Capit. semipoll. (v. s. comm. à cl. inv. sub n. 3108.)
17. *E. AMILACINUM* (H. B. et Kunth nov. gen. am. 4. p. 110), caule herbaceo volubili sulcato-angulato, ramulis piloso-hispidis, foliis oppositis petiolatis ovatis acuminatis grossè crenato-serratis trinerviis suprà hispidulis subtus hispidis, corymbo terminali subtrifido divaricato, capitulis sessilibus congestis, invol. oblongi squamis imbricatis obtusis striatis ciliatis.  $\zeta$  in Andibus Quindensis Novæ Gran. ? Flosc. et fruct. ign.
18. *E. INTERIOFOLIUM* (Bert. ! herb. Spreng. syst. 3. p. 410), fruticosum, ramis petiolisque minutè pubescentibus, foliis oppositis petiolatis cordatis ovatis obtusis subintegerrimis suprà glabris subtus glabrisculis densè punctatis, corymbo terminali subsessili, capitulis cylindricis circ. 20-floris, invol. squamis arcè imbricatis obtusis.  $\zeta$  in Guadalupà legit cl. Bertero. Capit. 4 lin. longa. Petioli 2-4 lin. longi. Folia subtus pallidiora. (v. s. comm. à cl. amic. inv.)
19. *E. ATRAPLICI-FOLIUM* (Vahl symb. 3. p. 96. non Lam.), fruticosum, ramis petiolisque velutinis, foliis oppositis petiolatis triangulari-ovatis dentatis acutis suprà glabris subtus glabrisculis densè punctatis, corymbo terminali subsessili, capitulis cylindraceis circ. 20-floris, invol. squamis arcè imbricatis obtusis.  $\zeta$  in ins. Sanctæ Crucis. *E. triangulare* Poir. suppl. 2. p. 604. Fortè nimis ad *E. repandum* accedit? — Stirpe Guadalupensis huc à Willd. relata videtur *E. odorati* aut punctati var. (v. s. sp. sine fl. cum herb. Vahl. coll. in h. Puer.)
20. *E. OSSÆANUM*, fruticosum glabrum, ramis teretibus vix cum foliis novellis subpuberulis, foliis oppositis petiolatis ovato-lanceolatis basi obtusis aut subattenuatis apice acuminatis grossè dentatis rigidis basi 3-5-nerviis suprà laevibus subtus tenuiter multipunctatis, corymbo trichotomo, capitulis ad apices ramulorum 3 sessilibus cylindricis circ. 20-floris, invol. squamis arcè imbricatis vix striatis obtusiusculis, achænio angulato glabro.  $\zeta$  in insulâ Cubâ legit cl. de la Ossa. Petioli 6-12 lin. Limbus fol. 4 poll. longus, ferè 2 poll. latus. (v. s. comm. à cl. inv.)
21. *E. ALTERNANS*, suffruticosum glabrum, caule tereti striatulo, ramis alternis, foliis lanceolatis subpetiolatis basi cuneatis apice acuminatis grossè serratis triplinerviis subreticulatis subtus punctatis, corymbo composito, ramis ramisque alternis, capitulis pedicellatis 20-25-floris cylindraceis, invol. squamis arcè imbricatis striatis obtusis.  $\zeta$  in Brasiliæ prov. Sancti-Pauli. Valdè accedit ad *E. propinquum*. Folia 2 poll. longa, 9 lin. lata. Capitula 3 lin. longa. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 815 miss.)
22. *E. GLABERRIMUM*, fruticosum undiquè glabrum, ramis teretibus, foliis oppositis subsessilibus ovalibus acuminatis basi obtusis serratis penninerviis reticulato-nerviosis, corymbo composito fastigiato polyccephalo, capitulis pedicellatis oblongo-cylindraceis circ. 20-floris, invol. squamis adpressis tenuibus striatis.  $\zeta$  in Mexico legit cl. Hænke. Petioli semilineares primà fronte nulli. Folia 6-7 poll. longa, 2 poll. lata. Capitula 3-4 lin. longa. Styli rami refecti nudè in siccò primo intuitu non apparent. (v. s. in h. Hænke. à cl. Steud. comm.)
23. *E. PSIADI-FOLIUM*, fruticosum glaberrimum subviscosum, caule tereti, ramis subhexagonis, foliis oppositis breviter petiolatis oblongo-lanceolatis utrinquè breviter acuminatis trinerviis basi integris apice grossè et penni-serr-

tis, corymbo trichotomo laxo, capitulis pedicellatis cylindricis 15-18-floris, invol. squamis arcuè adpressis obtusis. ♀ in Brasiliâ (h. L'Hér.), in sicca ad Cepovado (Lhotsky!). (v. s.)

β. *latifolium*, ramis subhexagonis, foliis ovatis acuminatis. ♀ in Cajennâ legit cl. Patris! (v. s.)

γ. *tereticaule*, ramis teretibus, foliis ovatis acuminatis grossius dentatis, capitulis gracilioribus. ♀ in Mexico legit cl. Hænke! (v. s. in ejus herb. à cl. Sternb. comm.) An species propria?

\*\* *Capitulis 10-20-floris, foliis oppositis.*

24. *E. MACRONON*, fruticosum glaberrimum, ramis teretibus, foliis oppositis petiolatis ovatis acuminatis grossè et obtusè cartilagineo-serratis subtrinerviis subtus eglandulosis, corymbo terminali decomposito, pedicellis compresso-angulatis, capitulis 16-floris, invol. squamis arcuè imbricatis obtusis. ♀ ad vertices montium Dominicæ locis humidis. An var. *E. macranthi*?, sed differt præter notas indicatas, petiolo 3 lin., capitulis 4 lin. longis. (v. s. ex h. L'Hér.)

25. *E. EXPANDUM* (Willd. sp. 3. p. 1767. excl. Lam. syn.) fruticosum, ramis petiulis pedunculisque pubescenti-tomentosis, foliis oppositis petiolatis ovatis acuminatis sinuato-dentatis supra junioribus velutinis adultis glabris subtus velutino-tomentosis parè glandulosis, corymbi ramis vix folia superantibus, capitulis pedicellatis cylindraceis circ. 15-floris, invol. squamis arcuè adpressis obtusis. ♀ in Sancto-Domingo (Plum. Lam.) et Cuba (R. de la Sagra!). Plum. Amer. ed. Burm. t. 130. f. 1. (nec 128. f. 1 in quâ corollæ involucri duplò longiores et folia obtusa). (v. s.)

26. *E. SINVATUM* (Lam. dict. 2. p. 407. ill. t. 672. f. 5), anfruticosum, ramis teretibus apice pubescentibus gracilibus, foliis oppositis petiolatis ovatis basi truncatis subtrinerviis apice subobtusis margine dentibus utrinquè 4-5 grossis obtusis lobato-crenatis supra scabridis subtus velutino-pubescentibus et glandulosis, pedunculis axillaribus divaricatis folio æqualibus et terminalibus parè corymbosis, capitulis pedicellatis oblongis 8-10-floris, invol. squamis imbricatis ext. dorso puberis int. glabris substriatis, achenio glabro. ♀ in Sancto-Domingo legit cl. Bertero. Petioli 2 lin. Limb. fol. pollicem longus, 8-9 lin. latus. Icon Plum. t. 128. f. 1. à Lamarckio citata verisim. excludenda. (v. s. comm. à cl. Bert.)

27. *E. GRACILIFLORUM*, fruticosum glabrum, ramis teretibus angulo recto divergentibus, foliis oppositis petiolatis ovato-lanceolatis acuminatis basi trinerviis dentatis apice integerrimis, corymbo oligocephalo laxo, capitulis pedicellatis teretibus gracilibus circ. 12-floris, invol. squamis arcuè imbricatis obtusis. ♀ in Mexico leg. cl. Hænke, circa Acapulco (Karvinski). Affine *E. clematiti*. Petioli 5 lin. longi. Fol. limbi 2 poll. longi, 7-9 lin. lati. Capitula 4-5 lin. longa sesquilineam lata. Corymbi 5-15-cephali. (v. s. in h. Hænke à cl. Sternb. comm. et h. Acad. reg. Monac.)

28. *E. FUSCATUM* (Lam. dict. 2. p. 408. non Willd.), fruticosum glaberrimum, ramis obliquè divergentibus, foliis oppositis petiolatis ovatis utrinquè acuminatis grossè serratis subtus punctatis, corymbis ramos terminantibus, capitulis pedicellatis cylindricis 15-16-floris, invol. squamis arcuè imbricatis obtusis. ♀ in ins. Caribæis (Sur. Vaill. ex Lam.), Guadalupe (Bad!) Trinitatis (Sieb!) E. odorato affine Sieb. fl. trin. exs. n. 75. — Syn. Pluk. cum signo? à Lam. citatum est certè exclud. Herba ex sicco videtur subviscosa. (v. s.)

29. *E. OXYLEPIS*, fruticosum, ramis teretibus densè hirsuto-velutinis, foliis oppositis brevissimè petiolatis ellipticis subæntis integerrimis aut subundentatis supra subscabridis subtus densè velutino-hirsutis, paniculæ oligocephalæ ramulis oppositis circ. 3-cephalis, capitulis cylindricis glabris 15-16-floris, invol. squamis arcuè imbricatis vix striatulis, ext. brevibus ovatis æntis, int. linearibus acuminatis. ♀ in Brasiliæ prov. Sancti-Pauli. Fol. pollicem longa 6 lin. lata. Petioli 1-2 lin. Capitula 5 lin. longa ad *E. leptocephalum* accedentia. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 468.)



30. *E. SALICINUM* (Lam. dict. 2. p. 409), fruticosum, caule subangulato pubescente ramis teretibus substriatis glabriusculis, foliis oppositis subpetiolatis oblongo-lanceolatis acuminatis supernè rariter dentatis suprà glabris et rugosis subtùs cano-tomentosis, corymbo glomerato convexo, capitulis 10-12-floris, invol. ovalis squamis imbricatis pubescentibus, intus longioribus obtusis.  $\frac{5}{8}$  in Peruvia ex h. Juss. ad radices montis ignivomi Tunguraguz Quitensiam. Spreng. syst. 3. p. 412. excl. syn. Folia 6-7 poll. longa.
31. *E. PROPINQUUM*, fruticosum glabrum, caule tereti, ramis oppositis, foliis oppositis subpetiolatis lato-lanceolatis basi cuneato-attenuatis apice acuminatis grossè serratis triplinerviis subreticulatis subtùs subpunctatis, corymbo composito trichotomo polycephalo, capitulis pedicellatis cylindricis 16-floris, invol. squamis arcuè adpressis striatis obtusis.  $\frac{5}{8}$  in Brasiliæ prov. Rio-Grande. Folia (cum petiolo 2-3 lin.) 3 poll. longa, ferè pollicem lata. Capitula 4 lin. lata. Accedit ad *E. ivæfolium* et *E. alternans*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 1064.)
32. *E. ivæFOLIUM* (Linn. amœn. ac. 5. p. 405. sp. 1175), herbaceum, caule tereti hispidulo, foliis oppositis angusto-lanceolatis utrinquè attenuatis subpetiolatis trinerviis subserratis glabris, corymbo trichotomo laxo, capitulis oblongis pedicellatis circ. 15-20-floris, invol. squamis paucis erectis striatis obtusis.  $\frac{2}{3}$  in cultis Jamaicæ (Swartz! obs. 301), Cubæ (Spr.) et in Mexici ditione Oaxaca (Andr. ! pl. exs. n. 283). (v. s.)  
 $\frac{1}{2}$  foliis medio serratis, capitulis 20-25-floris. — in Guadalupà (Bert.!). (v. s.)
33. *E. GRAMÆDRYFOLIUM* (H. B. et Kunth nov. gen. sin. 4. p. 113), fruticosum, ramis teretibus pubescenti-velutinis, foliis oppositis breviter petiolatis ovatis obtusiusculis crenatis trinerviis suprà scabris subtùs pubescenti-hirtis subto mentosis, junioribus canescentibus, corymbis compositis confertis breviter ramosis, capitulis sessilibus 10-12-floris subcylindraceis, invol. squamis adpressis obtusis substriatis dorso subpuberis, inf. ciliatis.  $\frac{5}{8}$  in Andibus Peruvianorum frequens (H. et B.), in vallibus (Hænke!). Cor. violaceæ. (v. s. in h. Hænke à cl. Sternb. comm.)
34. *E. FURCATUM* (Lam. dict. 2. p. 407), caule dichotomo glabro tereti, foliis oppositis, infer. ovatis, sup. lineari-lanceolatis, angustis integerrimis internodio longioribus, paniculis terminalibus, capitulis 10-12 floris, invol. ovali-oblongi imbricati squamis striatis obtusis, achenio angulato. — in ins. Caribæis ex h. Juss.
35. *E. RHINANTHACEUM*, suffruticosum glabrum, caulibus à collo plurimis teretibus simplicibus, foliis oppositis (supremis subalternis) subsessilibus lanceolatis basi attenuatis trinerviis parèe dentatis impunctatis, corymbo pedunculato subgloboso conferto, capitulis oblongis pedicellatis 10-12-floris, invol. squamis imbricatis oblongo-linearibus substriatis acutiusculis.  $\frac{5}{8}$  in Brasiliæ provinciâ Sancti-Pauli. Caulis pedalis. Folia sesquipollicem longa, 6 lin. lata. Accedit ad *E. ivæfolium* sed caule glabro, corymbo coarctato et invol. squamis acutis facile differt. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 500 miss.)
36. *E. VERBENACEUM*, suffruticosum, caule tereti sparè hispido, foliis oppositis petiolatis lanceolatis basi cuneatis apice acuminatis grossè dentatis margine scabris secus nervum subtùs hispidis cæterum glabriusculis, corymbo fastigiato, ramis oppositis, capitulis longè pedicellatis cylindricis 10-12-floris, invol. squamis arcuè imbricatis striatis glabris obtusis.  $\frac{5}{8}$  in Brasiliæ prov. Rio-Grande. Folia internodiis dimidio ferè breviora, caulina bipollicaria, ramos pollicaria, Verbenæ officinalis folia referentia. Corymbi rami glabri. Capit.  $\frac{4}{5}$  lin. longa. Cor. ex sicco purpureæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 760 miss.)
37. *E. TOXIFOLIUM*, caulibus ad collum suffruticulosis simplicibus teretibus apice præsertim puberulo-velutinis, foliis oppositis brevissimè petiolatis ovatis obtusè dentatis trinerviis suprà pubescentibus subtùs tomentosis, summis etiam dentatis, corymbi ramis oppositis folio longioribus, capitulis ad apicem

- ramulorum subsessilibus confertis 10-floris, invol. squamis imbricatis 4-5-seriatis striatis obtusis, int. apice villosiusculis.  $\frac{5}{2}$  in Brasiliæ prov. Sancti-Pauli. Valdè accedit ad *E. barstæifolium*. Achænia 5-gona ad angulo subvillosa. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 504 miss.)
38. *E. SANTÆFOLIUM*, suffruticosum, caule simplici tereti supernè pubescente, foliis confertis, infer. oppositis subsessilibus ovato-lanceolatis, sup. alternis exactè sessilibus angustius lanceolatis basi tantùm dentatis apice acuminatis integris, summis integerrimis angustis, sessilibus basi trinerviis grossè incisedentatis utrinquè pubescenti-scabris subtùs glandulosis, corymbo composito fastigiato, ramis alternis erectis, capitulis pedicellatis cylindricis 11-floris, invol. squamis imbricatis striatis obtusis dorso puberulo et glanduloso, intimis mucrone subpatulo brevi apiculatis.  $\frac{5}{2}$  in Brasiliæ prov. Rio-Grande (h. Mus. imp. bras. n. 822), et in Bandâ orientali (Bacle!). Folia pollicem longa. Rami floridi foliis multò longiores. Capit. 4 lin. Cor. et styli purpurei (v. s.)
39. *E. RHODOVOLUM*, fruticosum, caule tereti glabro, ramis hispidis, foliis oppositis sessilibus angusto-lanceolatis acuminatis hinc indè subseratis glabris subtùs punctulatis, corymbo composito fastigiato polycephalo, capitulis oblongis 5-floris. invol. squamis 9-10 glabris striatis obtusis.  $\frac{5}{2}$  in Brasiliæ prov. Rio-Grande. Habitus ferè *Eup. ivæfolii* sed corymbo multò magis amplo, ramis magis hispidis et capitulis 5-floris facilè distinctum. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 799 miss.)
40. *E. SERRULATUM*, fruticosum, ramis teretibus apice sex-sulcatis puberulo-sabhispidis, foliis oppositis breviter petiolatis lingulato-lanceolatis utrinquè subattenuatis regulariter serrulatis penninerviis utrinquè sparè glandulosis subtùs secus nervos puberulis, petiolo ciliato, corymbo oligocephalo, capitulis oblongis 5-floris. invol. squamis imbricatis glabris oblongis striatis obtusiusculis.  $\frac{5}{2}$  in Brasiliæ prov. Rio-Grande. Folia (incl. petiolo 2-3 lin.)  $2\frac{1}{2}$  poll. longa, 4-6 lin. lata. Capit. 3 lin. longa pallida. Cor. ex sicco ochroleucte. Affinis *E. comptoniæfolio*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 790 miss.)
41. *E. TAICHOPELOUM*, caulibus herbaceis erectis simplicibus setoso-hispidis, foliis oppositis et sup. alternis petiolo brevissimo hispidissimo donatis lanceolatis basi subtruncatis apice acuminatis grossè et obtusè dentatis utrinquè hispidis, corymbo terminali composito conferto polycephalo, capitulis breviter pedicellatis 10-floris, invol. squamis circ. 3-seriatis oblongis basi glabris striatis apice submembranaceis obtusis villosis. — in Brasiliæ prov. Sancti-Pauli. Setæ articulatæ. Folia sesquipollicem longa, 7 lin. lata. Achænia glabriuscula. Affine *E. barstæifolio*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 497 miss.)
- \*\*\* Capitulis 10-15-floris, foliis alternis.
42. *E. PULCHELLUM* (H. B. et Kunth nov. gen. am. 4. p. 119. t. 345), fruticosum, ramis teretibus pubescentibus, foliis alternis oblongo-lanceolatis utrinquè attenuatis dentatis suprâ scabriusculis subtùs tenuissimè pubescentibus penninerviis, corymbo composito terminali conferto polycephalo, capitulis 8-13-floris cylindricis pedicellatis, invol. squamis arcè imbricatis obtusis ciliatis.  $\frac{5}{2}$  in cœvalli Mexicana prope lacus Texcusenses. Cor. supernè violacæ. Styli longè exserti. (v. s. comm. à cl. Alaman, Mendez, Mairët.)
- \*\*\*\* Capitulis 5-9-floris, foliis oppositis.
43. *E. PUNCTULATUM*, fruticosum, caule tereti glabriusculo, foliis oppositis petiolatis ellipticis basi attenuatis apice acuminatis rariter dentatis aut integerrimis triplinerviis glabriusculis subtùs punctulatis, corymbo fastigiato composito, pediculis pubescentibus apice capitula 3-5 sessilia fasciculata cylindrica 6-9-flora gerentibus, invol. squamis arcè imbricatis obtusis substriatis.  $\frac{5}{2}$  in Amer. calid. sed locus proprius ignotus. Valdè accedit ad *E. punctatum*. Petioli 6-8 lin. longi. Folia 4 poll. longa, 12-14 lin. lata. Capit. 4 lin. longa. Achænia glabra. (v. s. in h. Martius.)

44. *E. SCABRUM* (Linn. f. suppl. 354), suffruticosum? erectum, ramis striatis villoso-scabris, foliis oppositis breviter petiolatis ovatis acuminatis sub serratis rarius integris subtrinerviis rugosis subtus hirtis supra scaberrimis, paniculis coarctatis terminalibus bracteatis, capitulis pedicellatis cylindricis 7-8-floris, invol. squamis arcuè imbricatis glabris oblongis obtusis trinerviis apice callosa, achen. glabro.  $\zeta$  in Novâ-Granatâ (Mut. in Linn.) juxta Santa-Fe (H. B. et Kunth). Smith pl. ined. 3. t. 67. *E. scabrum* H. B. et Kunth gen. nov. am. 4. p. 119 à Mutisio ut specim. Linn. communic. ab icone et descr. Smith. multum recedit ex cl. auct. Planta nigrescens. Flores ex Sm. flavescentes, potius verisim. albid.
45. *E. LEPTOCEPHALUM*, fruticosum glabrinsculum, ramis teretibusculis, foliis oppositis breviter petiolatis ovali-lanceolatis acuminatis serratis triplinerviis subtus crebre glanduloso-punctatis, corymbo composito fastigiato polycephalo, pedicellis 3-cephalis, capitulis cylindricis 7-8-floris gracilibus, invol. squamis arcuè adpressis substriatis ext. obtusiusculis intimis subacutis.  $\zeta$  in Andibus Peravianorum legit indef. Hænke. Corymbus ferè Odontolomæ. Rami subrubentes. Capitula glabra 4 lin. longa. Petioli sesquilineares. Fol. 2 poll. longa, 7-9 lin. lata. (v. s. in h. Hænke à cl. Sternb. comm.)
46. *E. HEXANTHUM*, suffruticosum, caule simplici tereti pubescente subhirtello, foliis sessilibus infer. oppositis cordatis breviter acuminatis serratis grosse denticatis subtus reticulato-nervosis supra scabris subtus secus nervos puberis, super. sensim alternis vix cordatis longius acuminatis subintegris, corymbi compositi ramis alternis paulò folia superantibus capitula 3-7 fasciculata cylindracea 6-flora gerentibus, invol. squamis imbricatis dorso puberulis et subglandulosis, int. acutè mucronatis.  $\zeta$  in Brasiliæ prov. Rio-Grande. Pedale. Affine *E. bartschifolium*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 519. miss.)
47. *E. GAUDICHAUDIANUM*, caule herbaceo? erecto tereti densè et sordidè cum petiolis pedicellisque velutino-hirtis, foliis oppositis petiolatis ellipticis obtusis basi subcuneatis crenato-dentatis penninerviis supra sparse scabridis, subtus pubescenti-hirtellis, corymbo terminali composito polycephalo densè conferto, capitulis congestis subsessilibus 5-floris, invol. ovalis squamis 10 biseriatis, ext. ferè subnatis, inter. ovali-oblongis obtusis apice ciliatis, achenio angulato glabro.  $\mathcal{U}$ ? circa Rio-Janeiro Brasiliæ in apricis legit cl. Gaudichaud. Petioli 2-3 lin. longi. Limb. fol. 15-16 lin. longus, 7-8 latus. Folia suprema sæpè alterna. (v. s. comm. à cl. invent.)
48. *E. INTERMEDIUM*, fruticosum, ramis teretibus apice præsertim velutino-hirtis, foliis oppositis basi brevissimè cuneatis et petiolulatis elongato-lanceolatis serratis penninerviis supra puberis subtus velutino-hirsutis, corymbi compositi terminalis ramis oppositis polycephalis, capitulis ad apices ramulorum confertis subsessilibus 5-floris, invol. squamis laxè imbricatis, ext. minimis acutis, inter. oblongis basi striatis glabris apice submembranaceis obtusis ciliatis dorso villosulis, achenio glabro.  $\zeta$  in Brasiliæ prov. Rio-Grande. Exactè medium inter *E. Gaudichaudianum* et *E. comptoniaefolium*. Folia 2-3 poll. longa, 6-9 lin. lata. (v. s. in h. Mus. Par. à Mus. imp. Bras. sub n. 844 miss.)
49. *E. COMPTONIAEFOLIUM*, fruticosum, caule tereti glabro, ramulis brevissimè velutinis, foliis oppositis brevissimè petiolatis lineari-sublanceolatis acutis regulariter serratis 1-nerviis supra glabris subtus brevissimè velutinis, corymbo composito fastigiato densò, capitulis oblongis 4-5-floris vix pedicellatis, invol. squamis 8-9 imbricatis subacutis striatulis int. apice ad dorsum tomentosia.  $\zeta$  in Brasiliæ prov. Rio-Grande. Cor. ex sicco subochroleæ breves. Achenia glabra. Fol. 1  $\frac{1}{2}$ -2 poll. longa, 3-4 lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 76 miss.)
50. *E. ARCLEPIADEUM*, suffruticosum, caule tereti apice villoso-hispido, foliis oppositis sessilibus aut summis alternis cordatis lanceolatis acuminatis integerrimis glabris subtus crebre punctatis, corymbi compositi ramis oppositis villoso-hispidis, ramulis apice capitula 3-5 subsessilibus oblonga 4-5-flora gerentibus.

- invol. squamis imbricatis striatis achenisque glabris.  $\zeta$  in Brasiliæ prov. Rio-Grande. Folia cōnfecta more Gentianæ asclepiadæ decussata sesquipollicem longa 5 lin. lata. Invol. 2-3 lin. longa pallida subpurpurascens. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 473, 774 et 842.)
51. *E. cuneatum*, fruticosum erectum ramosum glaberrimum viscosum, ramis teretibus striatis, foliis breviter petiolatis oppositis obovatis basi cuneatis et plurinerviis apice obtusis grossè dentatis coriaceis, corymbo composito fastigiato, invol. squamis 3-4 seriatis lanceolatis acutis ciliolatis pallidis, flor. 5, acheniis obovato-teretiusculis minutissimè scabridis.  $\zeta$  in Brasiliæ prov. Sancti-Pauli. Folia sesquipollicem longa, pollicem lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 503 miss.)
52. *E. polystachium* (DC. non Willd.), fruticosum glaberrimum, ramis teretiusculis striatis, foliis oppositis petiolatis ellipticis acutis serratis trinerviis subtus punctatis suprà glutine viscoso illinitis, paniculæ oblongæ ramis subracemosis, capitulis oblongis 5-floris pedicellatis, invol. squamis 8-10 imbricatis obtusis substriatis.  $\zeta$  in Brasiliæ prov. Sancti-Pauli. Petioli 3 lin. longi. Folia sesquipollicem longa, 8-9 lin. lata. Capit. 2-3 lin. longa. Achenia glabra angulata. Styli longè exserti. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 488 miss.)
- $\beta$ . *angustius*, foliis longius petiolatis oblongis suprà vix viscosis.  $\zeta$  in prov. Rio-Grande (h. Mus. imp. Bras. n. 976!). Petioli 6 lin. longi. Fol. limb. 2 poll. longus, 8-9 lin. latus. (v. s. in h. Mus. reg. Par.)
53. *E. xylophyllodes*, fruticosum glaberrimum, foliis oppositis longissimis sublinearibus utrinquè attenuatis subpetiolatis denticulatis coriaceis reticulato-venosis, paniculæ subcorymbosæ ramis oppositis, capitulis oblongis pedicellatis 4-floris, invol. squamis 8-10 imbricatis brevissimis subrotundis int. longioribus obtusiusculis substriatis.  $\zeta$  in Brasiliæ prov. Rio-Grande et Sancti-Pauli (h. Mus. imp. Bras. n. 997 et 513), prov. Minarum General. (Gaud. in h. Mus. Par.), circa Tejuco (Vauthier! pl. exs. n. 277). Species distinctissima, foliis coriaceis subfalcatis Xylophyllas angustifolias subreferens. Folia 4-7 poll. longa, 1  $\frac{1}{2}$ -3 lin. lata. Capit. vix 3 lin. longa. Achenia angulata glabra. Receptac. elevatum minimum. (v. s.)
54. *E. pinnatifidum*, fruticosum erectum ramosum glabrum, ramulis angulatis, adultis teretibus, foliis oppositis profundè pinnatifidis, lobis 1-2-jugis linearilanceolatis integerrimis, capitulis racemosis subpaniculatis 5-floris, invol. squamis 1-3-seriatis imbricatis oblongis acutis scarioso-rufis, achenio glabro.  $\zeta$  in Americâ austr. legit cl. Née, sed locus propr. mihi ignotus. Affine *E. xylophylloidi* ex inflorescentiâ, sed foliis facilè diversum (v. s.)
- \*\*\*\*\* *Capitulis 5-floris, foliis alternis.*
55. *E. eupleurifolium*, fruticosum, caule tereti glabro, paniculæ ramis puberulis, foliis alternis sessilibus lato-linearibus acuminatis integerrimis trinerviis nervis lateralibus tenuissimis subtus minutè glandulosis utrinquè subviscosis, paniculâ corymbosâ compositâ polycephalâ, capitulis oblongis 5-floris brevè pedicellatis, invol. squamis viscosis glabris imbricatis obtusis striatis, pappo involucri non superante.  $\zeta$  in Brasiliæ prov. Sancti-Pauli. Affine *E. linifolio* sed ex omni parte majus. Folia 4 poll. longa, 3-4 lin. lata. Corymbus major. Invol. fusco-purpurascens. Folia summa basi dilatata more Eupleurii. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 464 et 465 miss.)
56. *E. linifolium*, suffruticosum, caule tereti breviter pubescenti-hispido, foliis alternis subsensilibus erectis linearibus acuminatis integerrimis 1-nerviis margine subrevolutis subtus crebrè glandulosi, corymbo composito fastigiato, capitulis oblongis brevè pedicellatis 5-floris, invol. squamis erectis, ext. brevibus puberis ciliatis, int. longioribus obtusiusculis pallidis striis tribus notatis, pappo involucri longiore.  $\zeta$  in Brasiliâ ad Sanctam-Catharinam legit cl. Gaudichaud. Folia 2 poll. longa, 2 lin. lata. Cor. ex sicco purpurascens. Styli rami longissimi. (v. s. comm. à cl. invent.)

57. *E. ERICOIDES*, fruticosum erectum, caule tereti infernè glabro superne pulverulento-velutino, foliis alternis sessilibus linearibus basi subattenuatis integerrimis apice callosis et obtusis erectis confertis, corymbo composito terminali fastigiato, capitulis 5-floris pedicellatis, invol. squamis circ. 10 dorso velutinis margine ad apicem ciliatis demum patulis.  $\frac{1}{2}$  in Bras. prov. Rio-Grande. Specimen quoad flores mancum sed evident. *E. ericoidi* affine et foliis faciliè distinctum. Folia 6-7 lin. longa, lineam lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 834 miss.)
58. *E. HALIMIFOLIUM*, fruticosum erectum, caule tereti glabro, ramis ad apicem velutino-canis, foliis alternis cuneatis obtusis basi attenuatis integerrimis erectis confertis utrinquè pube sericeâ glandulis intermixtâ cano-sericeis, corymbo composito terminali, capitulis pedicellatis 5-floris, invol. squamis laxè imbricatis dorso subvillosis ext. acutis int. obtusis ciliatis, acheniis 5-gonis glabris.  $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 766) et in prov. Minas Generalium in monte Serra do Caraca (Mart.) *Vernonia halimifolia* Mart. ! herb. Invol. subpurpurea. Cor. purpurea. Pappus sordidus. Folia 6-9 lin. longa,  $1\frac{1}{2}$ -2 lin. lata. (v. s. in h. Mus. reg. Par. et Acad. reg. Monac.)
59. *E. ERICOIDES*, fruticosum glabrum erectum, caulibus teretibus, foliis alternis sessilibus linearibus subulatis acutis erectis confertis integerrimis rigidulis, corymbis terminalibus oligocephalis, capitulis pedicellatis 5-floris, invol. squamis 10 laxè imbricatis, ext. parvis acutis, int. longioribus ovalibus obtusis margine membranaceis ciliatis flosculis dimidio ferè brevioribus, achenio pentagono glabro.  $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Fruticulus pedalis. Folia 3-4 lin. longa. Styli exserti obtusi. Cor. ex sicco albidæ. Pappus achenio longior, setis rigidis basi ferè in anulum concretis crebrè barbellatis. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 843 et 775 miss.)
60. *E. STRAMINUM*, caule suffruticoso erecto ramosissimo tereti, ramulis folisque minutè puberulis, foliis alternis linearibus sublanco-latis uninerviis integerrimis, capitulis 5-floris ad axillas supremas breviter pedicellatis in racemos foliosos subdispositis, invol. squamis imbricatis lanceolatis acuminatis stramineis glabris striatis, achenio 5-gono ad angulos subscabro.  $\frac{1}{2}$  in Peruvia legit cl. Hænke. (v. s. in h. Hænke. à cl. de Sternberg. comm.)

### §. 2. Pteropoda.

Capitula 8-10 flora. Invol. oblongi squamæ imbricatæ oblongæ pallidæ. — Caulis herbacei. Folia in petiolum alatum angustata glabra impunctata.

61. *E. QUADRANGULARE*, herbaceum? undiquè glabrum, caule acutè tetragono, foliis sessilibus subconnatis lanceolato-ovatis apice acuminatis basi attenuatis penninerviis serratis membranaceis, paniculæ amplæ ramis oppositis striato-angulatis corymboso-paniculatis polycephalis, capitulis oblongis 8-10-floris, invol. squamis adpressis oblongis vix obtusis striatis. — in collibus Mexici ad Tantoyuca legit cl. Berlandier (n. 2151), ad Acapulco (seu fortè ad montes Oronocenses) legit cl. Hænke. Folia 6 poll. longa, 2 poll. lata. Panicula pedalis. Capit. 3 lin. longa pallida. Achenia angulata puberula. (v. s. in h. Hænke.)
62. *E. AROMATISANS*, herbaceum? glaberrimum, caulibus striato-sulcatis subangulatis, foliis oppositis latè ovatis acuminatis basi longè cuneato-angustatis in parte petioliformi subintegris cæterum grossè serratis, paniculæ amplæ ramis oppositis polycephalis, capitulis oblongis 10-12-floris pedicellatis, invol. squamis striatulis subciliatis obtusiusculis adpressis. — in insulâ Cubâ juxta Havanam, ubi ex cl. Ramon de la Sagra vulgò adhibetur ad odore aromatico impregnandas præparationes tabacinas vulgò *cigarres* dictas. Affine *E. quadrangulare*! Fol. pars petioliformis 3 poll. longa, limbus 6-7 poll. longus, 5 poll. latus. Styli rami longissimè exserti! (v. s. comm. à cl. de la Sagra.)
63. *E. THYRSOIDRUM* (Moc. fl. mex. ic. ined.), herbaceum? glaberrimum, caulibus teretibus striatis, foliis oppositis cordatis acuminatis serratis, petiolo late marginatis integerrimis basi subdilatis et ferè connatis, paniculæ amplæ

thyrsoides ramis oppositis, capitulis pedicellatis oblongis 8-9-floris, invol. squamis imbricatis acutis. — in Mexico. Affine *E. aromatisantii*, sed folia cordata serrataris minoribus donata et rigidiora, petioli subconnati basi latiores nec angustiores et capitula pauciflora. (v. ic. pict. et spec. sicc. in h. Hænk. à cl. Sternberg comm.)

℞. *puberum*, foliis secus nervos velutino-puberis, caule apice subpubero. — in Mexico legit cl. Hænke. Stirps coll. Andrieux exs. mex. n. 282. foliis subtus ubique sed tenuissimè pubescentibus huic fortè etiam referenda? (v. s. in ejus herb. à cl. Sternb. comm.)

### §. 3. *Sessilifolia*.

Capitula 5-12-flora. Invol. oblongi squamæ oblongæ imbricatæ. — Folia opposita sessilia. Caulis herbacei.

64. *E. sessilifolium* (Linn. sp. 1172), caule tereti glabriusculo, foliis sessilibus semiamplexicaulibus distinctis ovato-lanceolatis acuminatis basi rotundatis serratis glabris, corymbo composito, invol. squamis paucis oblongo-linearibus obtusis imbricatis, floribus in invol. 5-12. ½ in montanis rupestribus præ. calcareis in Alleghany (Pursh) occidentilibus (Michx.), in sylvis ad Roxburg prope Bostoniam (Big. fl. ed. 2. p. 293). Flores albi. Affine *E. perfoliatum* sed glaber et folia non connata (v. v. c. et sp.)

65. *E. truncatum* (Willd. sp. 3. p. 1751), caule tereti striato villosulo-hispidulo, foliis oppositis amplexicaulibus distinctis basi obtusis lanceolatis acuminatis rugosis dentato-serratis subtus pubescenti-villosis, corymbo composito conferto, capitulis 5-10-floris, invol. squamis 12-15 imbricatis linearibus obtusis. ½ in Marylandiâ (Sims) Pennsylvania (Willd.). *Eup. salicifolium* Sims bot. mag. t. 2010. Flores albi. Omnino medium inter *E. perfoliatum* et *E. sessilifolium*. (v. s.)

66. *E. revolutum* (Linn. sp. 1174), caule tereti striato villosulo-hirsuto, foliis oppositis connato-perfoliatis lanceolatis acuminatis dentatis rugosis supra glabris pubescentibus subtus tomentosis aut subhirsutis, summis distinctis, corymbo conferto, invol. squamis 12-15 imbricatis lineari-lanceolatis, floribus in inv. 8-10, achenio glabro. ½ in pratiis humidis Amer. bor. à Canadâ ad Floridam. Pursh fl. bor. am. 2. p. 516. Ell. sketch 2. p. 302. Big. fl. bost. ed. 2. p. 297 et med. bot. 1. p. 38. t. 2. Raf. fl. med. 1. n. 36. ic. *E. connatum* Michx. fl. bor. am. 2. p. 99. Flores albi. Variat foliis super. alternis! Herbar. (vulgò *Boneset*) amaræ decoctum emeticum sudorificum febrifugum in Amer. popolare. (v. s.)

### §. 4. *Verticillata*.

Capitula 5-10-flora. Invol. oblonga squamis oblongis imbricatis. Folia 3<sup>ies</sup> 4<sup>ies</sup> verticillata. Caulis herbacei.

67. *E. ternifolium* (Ell. sketch 2. p. 306), caule tereti striato pubescente solido, foliis 3-4 verticillatis ovato-lanceolatis acuminatis dentatis supra sparse subpuberis subtus pubescentibus glandulosis, corymbo conferto, invol. squamis imbricatis, ext. brevibus valde puberis, int. lineari-oblongis striatis subpubescentibus, floribus in capitulo paucis. — in pinguibus et demissis Carolinæ (Fras. ! Ell.). Capitula parva pallida. Achenia angulata. Cor. albidæ vix purpureo tinctæ. *E. trifoliatum* Linn. sp. 1173 ex syn. Gron. virg. 119, sed species cum aliis verticillatis sæpe confusa. (v. s.)

℞. foliis majoribus basi longius cuneato-acuminatis. *E. verticillatum* var. Torr. ! in litt. 1820, ~~Ell.~~ l. c. in adn. (v. s.)

68. *E. purpuræum* (Linn. sp. 1173), caule tereti lævi glaucescente fistuloso, foliis 4-5<sup>ies</sup> verticillatis brevissimè petiolatis ovato-lanceolatis dentato-serratis subrugoso-venosis scabriusculis, capitulis corymbosis 5-floris, invol. squamis interioribus longioribus subscariosis linearibus oblongis glabris, achenio pentagono vix pubescente. ½ in humidis pinguibus Amer. bor. à Canadâ ad Caro-

- linam. Pursh fl. bor. am. 2. p. 515. Ell. sketch 2. p. 307. — Cora. cad. cap. 72. p. 191. ic. Moris. ox. 3. s. 7. t. 13. f. 4. malè. Dod. mem. t. 217. An *E. laevigatum* Torr. cat. pl. New-York. p. 92. non Lam. huc revocandum? *E. purpureum* Lour. videtur omnino diversum! Herba obscure virens sæpè purpureascent. Cor. pallidè purpureæ. (v. s.)
69. *E. MACULATUM* (Linn. amœn. ac. 4. p. 288), caule solido sulcato superne pubescente sæpè lineis purpureis maculato, foliis 4-5<sup>æ</sup> verticillatis petiolatis ovato-lanceolatis inæqualiter serratis subtus pubescentibus, corymbo polycephalo, capitulis pedicellatis 5-8-floris, invol. squamis imbricatis lævibus ext. subrotundis int. linearibus elongatis, achenio angulato glabro glanduloso.  $\mathcal{L}$  in humidis Amer. bor. à Canadà ad Carolinam. Pursh fl. 2. p. 1760. Ell. sketch 2. p. 308. — Herm. parad. 158. t. 159. Moris. ox. s. 7. t. 18. f. 3. An confirmante cl. Hooker fl. bor. 1. p. 305, ab *E. purpureo* satis differt? (v. s.)
70. *E. VERTICILLATUM* (Muhl. in Willd. sp. 3. p. 1760), caule tereti lævi solido sæpè lineis purpureis maculato, foliis 3-4<sup>æ</sup> verticillatis basi cuneato-attenuatis ovato-lanceolatis inæqualiter serratis glabriusculis, corymbo composito, capitulis pedicellatis 5-10?-floris, invol. squamis imbricatis lævibus.  $\mathcal{L}$  in pinguib. Carolinæ super. (Bosc!) Georgiæ (Ell. sketch 2. p. 308) agri Bostonienas (Big. ! fl. bost. ed. 2. p. 298). *E. purpureum* Michx. fl. 2. p. 99. non alior. *E. fusco-rubrum* Walt. car. 199? Cor. purpureæ. Affine *E. maculato* et sæpè cum eo confusum, sed foliis non verè petiolatis facilè differt. (v. s.)
71. *E. DUBIUM* (Poir. suppl. 2. p. 606), caule tereti striato solido pubescente, foliis 3<sup>æ</sup> verticillatis petiolatis ovatis vix acuminatis subserratis super sparse puberulis subtus pubescentibus glandulosisque. corymbo longè pedunculato, capitulis 3-floris, invol. squamis imbricatis lævibus, int. longioribus acuminatis lineatis, achenio angulato glanduloso.  $\mathcal{L}$  in Pensylvanià (Muhl. ! ) novà Casarsè (Pursh.) Carolinà merid. (Fraser. ! ). *E. punctatum* Willd. enum. 853. non Lam. Affine videtur *E. ternifolium* sed capitula pauciflora, invol. glabra, folia ovata, etc. (v. s.)

Ser. II. SUBIMBRICATA. *Nempè involucris aut laxè aut paucis seriebus imbricatis ovatis campanulatisve.*

\* Capitulis 3-5-floris, foliis alternis.

72. *E. TRIFLORUM* (Aubl. guian. 2. p. 795. t. 314), fruticosum, ramis teretibus sarmentosis albo-velutinis, foliis alternis breviter petiolatis ovato-lanceolatis acutis subcordatis integerrimis suprà scabris subtus albidove-lutinis, pedunculis axill. et termin. racemoso-paniculatis, capitulis glomeratis 3-floris, invol. oblongi squamis imbricatis oblongis acutis concavis.  $\mathcal{J}$  in Guianà Gallicà ad ripam fl. Aroura. Lam. dict. 2. p. 404. Willd. sp. 3. p. 1749. Cor. alba. Petiol. 4 lin. Limb. fol. 5 poll. longus, 2 poll. latus.
73. *E. CAMPESTRIS*, herbaceum erectum glabrum, caulibus simplicibus striatulis, foliis alternis oblongis basi subtrinerviis cuneato-subpetiolulatis integris, à medio ad apicem dentatis, corymbo composito fastigiato, capitulis breviter pedicellatis 5-floris, invol. squamis subimbricatis acutiusculis, corollis acheniisque puberulis.  $\mathcal{L}$  in campis editis prov. Sancti-Pauli novemb. flor. legit cl. Lund (n. 852 ! ). Herba pedalis stricta. Folia pollicem longa, 4 lin. lata. Cor. dilatè violacæ. (v. s.)
74. *E. SUBALTERNIFOLIUM*, caule herbaceo erecto gracili tereti pubescenti-velutino, foliis inf. oppositis cæteris subalternis breviter petiolatis ellipticis seu obovato-oblongis utrinquè attenuatis rigidulis ~~ambidentatis~~ <sup>puberulis</sup> suprà glabris subtus trinerviis et conspicuè reticulatis canescenti-puberulis, corymbo terminali pedunculato conferto, capitulis ad apices ramulorum plurimis sessilibus 5-floris, invol. squamis 2-seriatis subimbricatis, ext. obtusè dorso puberulis, inter glabris apice serrulatis, achenio glabro.  $\mathcal{L}$  in Brasiliæ prov. Minarum General ad Serra do Frio (Vanth. ! n. 335), in campis editis prov. Sancti-Pauli (Lund herb. n. 851 ! ). Herba circ. pedalis, rhizom. verisim. lignoso. Folia pseudo-

opposita cum petiolo 10-12 lin. longa, 4 lin. lata. Cor. ex sicco purpurascens. Styli longè exserti. Pappus longit. corollæ. (v. s.)

75. *E. DICTYOPHYLLUM*, caule suffruticoso erecto tereti pulverulento-velutino foliis alternis coriaceis ovatis basi in petiolulum cuneato-attenuatis superne remotè et callosè-serratis utrinquè glandulosis et puberulis subtus reticulo-nervulorum elevatorum eleganter areolato, corymbo paniculato polycephalo, invol. squamis omnibus linearibus acuminatis exter. dorso puberulis cæt. glabris, flor. 5, acheniis villosis. ♀ in campis editis prov. Sancti-Pauli nov. flor. legit cl. Lund (n. 853!). Flores albi. Folia 3 poll. longa,  $1\frac{1}{2}$  lata. (v. s.)

\*\* Capitulis 5-10-floris, foliis oppositis.

76. *E. INESHOIDES* (H. B. et Kunth nov. gen. am. 4. p. 106. t. 340), caule herbaceo volubili striato villosò-pubescente, ramis divaricatis, foliis oppositis petiolatis deltoideo-ovatis acutis serrato-dentatis triplinerviis suprâ pubescentibus subtus molliter canescenti-pubescentibus, paniculâ terminali trichotomâ diffusâ, capitulis pedicellatis subfastigiatis sub-5-floris, invol. squamis circ. 16 subimbricatis oblongis glabris, ext. mucronatis, int. acuminatis. ♀ ad radices Andinum Novo-Granatensium prope Ybague. Cor. alba. Habitus lissimè. Petiol. 4 lin. limbus 20-21 lin. longus, 13-14 latus.

77. *E. PYRIFOLIUM*, fruticosum glabrum, ramis teretibus, foliis oppositis petiolatis ovatis ovalibusve utrinquè vix acutis rigidulis integerrimis penninerviis, paniculis axillaribus oppositis terminalibusque polycephalis, capitulis pedicellatis 5-floris, invol. squamis 4-seriatis imbricatis ferè 5-fariis striatis ext. brevibus ovatis acutis, int. oblongis longioribus, achenio puberulo. ♀ in fruticetis ad Cercovado Brasiliæ legit cl. Lhostky. Petioli 6-9 lin. longi. Limb. fol. 3 poll. longus. 18-21 lin. latus. (v. s. comm. à cl. inv.)

78. *E. DENDROIDES* (Spreng. syst. 3. p. 415), arboreum glabrum, ramis sexangularibus, foliis oppositis petiolatis ovatis acutis basi cuneatis crenato-dentatis coriaceis reticulato-venosis, corymbis terminalibus ramosissimis, capitulis geminis ternisve subsessilibus sub-5-floris, invol. oblongis squamis 7-8 imbricatis linearibus obtusis viscosis, achenio glabro. ♀ prope Loxa Novo-Granatens. Mikania arborea H. B. et Kunth nov. gen. am. 4. p. 139. Petioli 7-8 lin. longi. Limbus fol. sub 4-pollicaris, 21-22 lin. latus.

79. *E. RACCHAROIDES* (H. B. et Kunth nov. gen. am. 4. p. 132), fruticosum glabrum, ramis sulcato-striatis, foliis oppositis petiolatis lanceolato-oblongis utrinquè acutis remotè serratis coriaceis, corymbis terminalibus, ramulis diffusis viscosis, capitulis pedicellatis 6-floris, invol. cylindraceo-campanulati squamis imbricatis oblongis acutiusculis subciliatis, achenio ad angulos scabro. ♀ in umbrosis juxta Alaguer Novo-Granatensium. Petiol. 3 lin. Limb. fol.  $2\frac{1}{2}$  poll. longus, 10-11 lin. latus. Cor. alba.

80. *E. ANGUSTIFOLIUM* (Spreng. syst. 3. p. 415), fruticosum, ramis teretibus albido-tomentosis, foliis oppositis petiolatis oblongo-lanceolatis acutiusculis basi rotundatis integerrimis reticulato-venosis coriaceis suprâ glabris subtus albido-tomentosis, corymbo paniculatis, capitulis subgeminis sessilibus 6-floris, invol. oblongis squamis subdenis imbricatis oblongo-linearibus apice fimbriatis, acheniis glabriusculis. ♀ in umbrosis Nova-Granatæ inter Ybague et Cuestada Tolima. Cor. pilosiuscula. Anthæræ subexsertæ. Mikania angustifolia H. B. et Kunth nov. gen. am. 4. p. 138. Petioli semipoll. Limb. fol.  $4\frac{1}{2}$  poll. longus, 14 lin. latus.

81. *E. ACUMINATUM* (H. B. et Kunth nov. gen. am. 4. p. 107), caule herbaceo-sexangulari glabro, foliis oppositis petiolatis subrotundo-ovatis leviter cordatis valdè acuminatis dentatè-serratis 5-nerviis glabris subtus in venis nervisque pubescentibus, corymbo terminali conglomerato, capitulis sessilibus sub 7-floris, invol. cylindracei squamis imbricatis lineari-oblongis striatis glabris, exter. minimis, achenio glabro. ♀ in Nova-Granatâ? Petiol. 15 lin. longus. Limb. fol. 4 poll. longus, 3 poll. latus. Cor. alba? glabra.



82. *E. tenuifolium* (H. B. et Kunth nov. gen. am. 4. p. 107), caule herbaceo erecto subangulati glabro, foliis oppositis petiolatis ovatis grosse dentato-serratis trinerviis glabris subtus in venis nervisque pubescentibus, corymbo terminali sessili oppositè ramosissimo, capitulis sessilibus conglomeratis sub 7-floris, invol. cylindracei squamis imbricatis, ext. ovatis brevibus ciliatis, int. oblongis obtusis glabriusculis. 2 $\frac{1}{2}$  in humidis Amer. austr. inter Cumana et Bosdones. Petiol. 10 lin. Fol. limbus 2  $\frac{1}{2}$  poll. longus, 20 lin. latus.
83. *E. micranthum* (Less. in linnæa 1830. p. 138), fruticosum glabrum, foliis oppositis petiolatis papyraceis penninerviis ovato-lanceolatis acutiusculis remotè serratis subtus densè glandulosis, paniculis trichotomis subfastigiatis, capitulis 7-8-floris, invol. turbinato-cylindracei squamis siccis erecto-patentibus apice ciliatis. 3 in sylvis Kalappæ Mexic. Cor. purpureæ. An *E. micranthum* Lag. nov. gen. et sp. 25 n. 333 omninò idem? Ex Lessingio aff. *E. fastigiato* Kunth! ex Lag. *Conoclinio cælestino*!
84. *E. pallescens*, suffruticosum?, ramis teretibus tenuissimè et molliter pubescentibus, ramulis divaricatis, foliis oppositis petiolatis ovato-lanceolatis basi breviter cuneatis integris triplinerviis apice longè acuminatis dentato-serratis suprâ glabriusculis subtus pubescentibus, corymbis ad apices ramorum confertis polycephalis, capitulis breviter pedicellatis 7-9-floris, invol. squamis ovali-oblongis obtusis striatis 2-3-seriatis laxè imbricatis glabriusculis. 3 in Brasiliæ prov. Minarum General. ad Sabara legit cl. Vauthier. Invol. straminea, corollas subæquantia. Achænio glabrum. Petiol. semipollic. Limb. fol. 4 poll. longus, pollicem circ. latus. (v. s. comm. à cl. Vauthier n. 273.)  
 $\beta$ . *hirsutum*, fruticosum, ramis velutino-hirsutis, foliis suprâ puberulis subtus velutino-hirsutis. 5 in Brasiliæ prov. Rio-Grande. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 789 miss.)
85. *E. glomeratum*, caule suffruticuloso erecto cum petiolis et pedunculis hirsutulo, foliis oppositis petiolatis basi cordatis ovato-lanceolatis acuminatis dentato-crenatis suprâ minutè pubescentibus subtus molliter subtomentosis, ramis floridis oppositis divaricatis paniculam confertissimam gerentibus, ramulis apice 3-cephalis, capitulis 7-8-floris, invol. squamis glabris striatulis parè imbricatis, achæniis pentagonis hispidulis. 5 inter Chilenses herb. Hænkiani adnumeratur. Folia ferè Betonicæ. Styli longè exserti. Capitula adeò adgregata ut primâ fronte multiflora videantur. (v. s. in h. Hæнке.)
86. *E. leucophyllum* (H. B. et Kunth nov. gen. am. 4. p. 115), fruticosum, ramis teretiusculis albo-tomentosis, foliis oppositis petiolatis ovatis apice angustatis acutis leviter cordatis subtrinerviis suprâ pubescentibus subtus albo-tomentosis, paniculis terminalibus ramosissimis foliatis diffusis, capitulis pedicellatis 9-10-floris, invol. oblongo-campanulati squamis imbricatis obtusis pubescentibus, ext. brevibus acutiusculis, achænio glabriusculo. 3 in temperatis Andium Peruvianorum. Cor. albæ? Petiol. subsemipollic. Limb. fol. 4 poll. longus, basi pollicem latus.
87. *E. lavandulæfolium*, fruticosum, ramis teretibus junioribus cano-tomentosis, foliis oppositis subsessilibus linearibus integerrimis obtusiusculis margine revolutis suprâ molliter villosis demùm glabris subtus tomentosis, paniculâ ovatâ brevi confertâ, capitulis ad apices ramorum congestis sessilibus 9-10-floris, invol. squamis oblongis obtusis laxè imbricatis, ext. dorso villosis, inter. glabriusculis subscariosis. 5 in Amer. merid. legit Dombey, sed locus prop. ign. in vallibus Andium Peruviz (h. Hænk!). Accedit ad *E. stachadifolium* sed folia revoluta nec plana. Recept. planum nec semiglobosum. Invol. squamarum obtusæ nec acutæ etc. (v. s. comm. à Mus. reg. Paris.)

\*\*\* Capitulis 10-20-floris, foliis oppositis.

88. *E. decemflorum*, suffruticosum, ramis teretibus apice præsertim pubescentibus, foliis oppositis brevissimè petiolatis elliptico-oblongis utrinquè acuminatis suprâ basin triplinerviis serrulatis suprâ sparè puberulis subtus pubescentibus et glandulosis, corymbi terminalis congesti ramis oppositis, capituli

- subsessilibus 10-floris, invol. squamis 2-3-seriatis subimbricatis obtusis, exter. brevibus pubescentibus, inter. longioribus membranaceis glabris, achænio glabro.  $\frac{5}{3}$  in Peruvia (Poeppig!), in vallibus Andium (h. Hænk!). Folia 3 poll. longa, pollicem lata. Capitula straminea. Invol. corollas æquantia. (v. s. comm. à cl. Poeppig. sub n. 30.)
89. *E. PELLUCIDUM* (H. B. et Kunth nov. gen. am. 4. p. 108), caule herbaceo erecto sexangulari glabro, foliis oppositis petiolatis subrotundo-ovatis leviter cordatis acuminatis dentato-serratis 5-nerviis glabris subtus in venis nervisque pubescentibus, venulis pellucidis, corymbo terminali conglomerato, capitulis sessilibus 10-floris, invol. cylindracei squamis imbricatis lineari-oblongis glabris nitidis albidis, achænio secus angulos scabriusculo.  $2\frac{1}{2}$  in Novâ-Granatâ. Simile *E. acuminato* sed præter notas ind. differt ex cl. auct. Invol. 24-27 nec 15-phyllis.
90. *E. HELIANTHIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 127), fruticosum, ramis teretibus hispidis, foliis oppositis petiolatis ovato-oblongis acuminatis serratis supra scabriusculis subtus glanduloso-punctulatis et pilosiusculis, corymbis terminalibus trifidis, capitulis pedicellatis 10-floris, invol. campanulato-cylindracei squamis imbricatis oblongo-linearibus obtusis glabriusculis.  $\frac{3}{2}$  prope pagum Ayayaca Peruvianorum. Petiol. 4 lin. Limb. fol. sub 5-pollicaris, 22 lin. latus.
91. *E. CANESCENS* (Vahl! symb. 3. p. 95. t. 73), fruticosum, ramis teretibus apice pulverulento-velutinis, foliis oppositis breviter petiolatis ovatis dentibus paucis obtusis sublobatis integerrimisve obtusis supra glabriusculis subtus canescenti-velutinis, corymbo simplici terminali, capitulis pedicellatis 10-floris ovatis, invol. squamis 3-serialibus laxè imbricatis obtusis, inter. striatis, achænio pube minimâ subvillosa  $\frac{3}{2}$  in Ins. Sanctæ-Crucis (Vest in h. Puer!) et Sancti-Thomas. (Willd.). Folia petiolo 1-2 lin., limbo 9-12 lin. longo, 5-6 lato. (v. s.)
- β. adenocarpum*, foliis paulò minoribus, achænio glabro globulis glandulose consperso.  $\frac{3}{2}$  in Sancto-Domingo legit cl. Bertero. (v. s.)
92. *E. VARGASIANUM*, fruticosum, ramis teretibus petiolis pedunculisque densè breviterque hirsutis, foliis oppositis petiolatis latè ovatis acutis basi subcordatis 5-nerviis dentatis supra sparse scabridis subtus pubescenti-hirsutis, corymbo amplo conferto polycephalo oppositè ramoso, capitulis pedicellatis 10-floris, invol. squamis ovali-oblongis vix acutis 2-3-seriatis ciliatis dorso glabris, achænio puberulo.  $\frac{5}{2}$  circa Caraccas legit cl. Jos. Vargas. Petiol. ferè sesquipollic. Fol. limbus 3-4 poll. longus, 2-2  $\frac{1}{2}$  poll. latus. Invol. straminea. (v. s. comm. à cl. Vargas.)
93. *E. NIVÆUM* (H. B. et Kunth nov. gen. am. 4. p. 115. t. 342), fruticosum, ramis sulcatis tenuissimè petiolis pediculisque albo-lanatis, foliis oppositis petiolatis triangulari-ovatis acutis leviter cordatis crenatis trinerviis supra glabris nitidis subtus niveo-tomentosis, paniculis subcorymbosis ramosissimis foliatis diffusis, capitulis pedicellatis 10-12-floris, invol. oblongi squamis imbricatis striatis villosa-pubescentibus, ext. ovatis, achænio glabro.  $\frac{3}{2}$  juxta Popayan Novo-Granatensium? Cor. alba? Petiol. 5-6 lin. Limb. fol. 15 lin. longus, 11 lin. latus. Species elegans!
94. *E. POPULIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 111), arborecens glabrum, ramis sulcatis, foliis oppositis petiolatis ovatis acutis obtusè grossè et inæqualiter dentatis subcoriaceis, paniculis terminalibus ramosissimis patentibus, capitulis sessilibus subquaternis approximato-congestis 10-12-floris, invol. oblongo-cylindraceo imbricato, squamis obtusiusculis trinerviis.  $\frac{5}{2}$  in aridis calidissimis juxta pagum Mazatlan Mexicanorum. Petiol. 2 poll. Limb. fol. 7 poll. longus, 5 poll. latus.
95. *E. SOLIDAGINOIDES* (H. B. et Kunth nov. gen. am. 4. p. 126), fruticosum, ramis sexangularibus glabris, foliis oppositis petiolatis ovatis acuminatis serratis subtrinerviis supra glabris subtus punctulatis et pilosiusculis, paniculis terminalibus ramosissimis foliatis divaricatis, capitulis pedicellatis 10-13-floris,

- invol. campanulati squamis imbricatis lanceolatis acuto-subulatis subciliatis, achenio ad angulos hispido.  $\bar{\gamma}$  in temperatis Quitensium inter Ticaen et Alausi. Cor. albæ? E. celtidifolio et E. cœlestino (mihi diversissimis) dicitur affine.
96. *E. ellipticum*, scandens, ramis elongatis teretibus striatis pubescentibus, foliis alternis petiolatis ellipticis acuminatis glabris integerrimis, capitulis ad apicem ramorum paniculatis pedicellatis 12-15-floris, invol. squamis 10-12 glabris subciliatis acutis, int. longioribus sed disco dimidio brevioribus.  $\bar{\gamma}$  (?) in Indiâ or. ad Penang legit cl. Wallich. Commixtum errore quodam cum Mikaniâ volubili à Wall. cat. n. 3174 sed evidenter diversissimum. Flores violaceo-purpurei, fauce campanulatâ. Pappus rufus copiosus. (v. a. comm. ab hon. Cur. merc. angl. Ind. or.)
97. *E. fastigiatum* (H. B. et Kunth nov. gen. am. 4. p. 125. t. 347), fruticosum glabrum, ramis sulcatis, foliis oppositis petiolatis oblongis acutis basi angustatis crenato-serratis coriaceis subviscosis, corymbis terminalibus ramosissimis fastigiatis, capitulis pedicellatis circ. 12-floris, invol. subcampanulati squamis imbricatis lineari-oblongis obtusis subciliatis.  $\bar{\gamma}$  in frigidis montium Novo-Granatensium. Cor. alba. Petiol. 7-8 lin. Limb. fol. sesquipollicem longus, 13-14 lin. latus.
98. *E. frunifolium* (H. B. et Kunth nov. gen. am. 4. p. 132. t. 349), arboreum, ramis teretiusculis, ramulis piloso-pubescentibus, foliis oppositis sparsive petiolatis oblongis acutis basi rotundatis subintegerrimis coriaceis suprâ glabris subtus hirtopubescentibus, corymbis terminalibus oppositis ramosissimis, capitulis pedicellatis 12-floris, invol. oblongi squamis laxè imbricatis acutis glutinoso glabris, achenio ad angulos scabriusculo.  $\bar{\gamma}$  in convalli Tarquienis juxta urbem Quito. Petiol. 2-3 lin. Limb. fol. 16 lin. longus, 8 lin. latus. Cor. alba.
99. *E. inulæfolium* (H. B. et Kunth nov. gen. am. 4. p. 109), caule herbaceo erecto sexangulari tomentoso-pubescente, foliis oppositis breviter petiolatis ovato-lanceolatis longissimè acuminatis basi in petiolum angustatis remotè serratis suprâ pubescentibus subtus tomentoso-pubescentibus et canescentibus, paniculâ corymbosâ diffusâ, capitulis subsessilibus congestis circ. 13-floris, invol. oblongi squamis laxè imbricatis lineari-oblongis obtusis striatis ciliatis, achenio glabro.  $\bar{\gamma}$  in dumetis circa urbem Mariquita Novo-Granatensium.
100. *E. molle* (H. B. et Kunth nov. gen. am. 4. p. 109), caule herbaceo erecto sexangulari cano-tomentoso, foliis oppositis petiolatis ovato-oblongis longissimè acuminatis basi attenuatis obsolete crenatis suprâ pubescentibus subtus molliter et tenuissimè tomentosis canescentibus, paniculâ subcorymbosâ patente, capitulis subsessilibus conglomerato-congestis circ. 13-floris, invol. oblongi squamis laxè imbricatis lineari-oblongis obtusis ciliatis, achenio glabro.  $\bar{\gamma}$  juxta Mariquita Novo-Granatensium cum *E. inulæfolio* à quo vix satis differt.
101. *E. suaveolens* (H. B. et Kunth nov. gen. am. 4. p. 109), caule herbaceo erecto teretiusculo molliter pubescente, foliis oppositis petiolatis ovato-oblongis angustato-acuminatis basi angustatis obtusè serratis subtriplinerviis suprâ scabris subtus hirtopubescentibus subrugosis, paniculâ corymbosâ fastigiatâ, capitulis subsessilibus conglomeratis circ. 13-floris, invol. oblongi squamis laxè imbricatis lineari-oblongis obtusis ciliatis, achenio glabro.  $\bar{\gamma}$  in temperatis montium Novæ-Granatæ. Flores suaveolentes. Valdè affine *E. inulæfolio* et *E. molli*. — *E.* circa Montevideo à Sellowio lectum et huc à Spreng relatum est verisim. diversum.
102. *E. concinnum*, fruticosum, caule tereti glabro, ramis præs. ad apicem hirsutis, foliis oppositis breviter petiolatis ovali-lanceolatis acuminatis serratis penninerviis suprâ glabriusculis subtus cano-tomentosis, corymbo laxo fastigiato, capitulis pedicellatis 15-floris, invol. squamis 3-seriatis laxè ambimbricatis oblongo-linearibus acutiusculis ciliatis dorso puberulis, achenio angulato glabro.  $\bar{\gamma}$  in Brasiliæ prov. Rio-Grande. Folia 2 poll. longa, 7 lin. lata.

- Capit. 6-7 lin. longa. Achænium bilineare. Cor. ex sicco pallidæ. Styli valdè exserti. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 827 miss.)
103. *E. PENTLANDIANUM*, fruticosum glabrum, ramis teretibus, foliis oppositis breviter petiolatis ovato-lanceolatis subacuminatis serratis basi trinerviis subtus pallidis, corymbis terminalibus fastigiatis, capitulis pedicellatis 12-15-floris, invol. squamis oblongis obtusiusculis 2-3-seriatis laxè imbricatis puberulis corollis dimidio brevioribus, acheniis glandulosis glabris. ♂ in Amer. merid. parte centrali nunc Bolivia dicta legit cl. Pentland. Frutex ramosissimus obscure virens. Folia conferta. Cor. purpureæ. Pappus sordidus. Achænia albidæ. (v. s. comm. à cl. inv.)
104. *E. MONTANUM* (Swartz) prodr. 111. flor. 1311), fruticosum, ramis teretibus hirsutiusculis, foliis oppositis breviter petiolatis cordiformibus acutis denticulatis suprâ sparè hispido-scabris subtus secus nervos hirsutis granuliferis, corymbo pedunculato, ramis trifidis polycephalis, capit. circ. 15-floris, invol. squamis imbricatis conniventibus acutis hirtis. ♂ in montibus altis Jamaicae. Petioli 2 lin. hirsuti. Limbi fol. 2  $\frac{1}{2}$  poll. longi, 1  $\frac{1}{2}$  lati, lobis bascos conniventibus. Cor. exalbidæ. Pubes ferruginea. Affine *E. cordifolium*. (v. s. ex h. Swartz.)
105. *E. GLABRATUM* (H. B. et Kunth nov. gen. am. 4. p. 127), fruticosum glabrum, ramis teretiusculis striatis, foliis oppositis petiolatis ovato-oblongis acutis dentato-serratis subcarnosis, pedunculis axillaribus terminalibusque subtricephalis, capitulis circ. 16-floris, invol. campanulatis squamis subimbricatis lanceolatis obtusiusculis externè subviscosis, achænio 5-angulato ad angulos scabro. ♂ in locis altis montium Mexicanorum inter Omitlan et Real del monte. Cor. albæ glabræ invol. duplò longiores. Petiol.  $\frac{1}{2}$  poll. Limb. fol. 2  $\frac{1}{2}$  poll. longus, pollicem et paulò ultrâ latus.
106. *E. RUBRICAULE* (H. B. et Kunth nov. gen. am. 4. p. 124), herbaceum glabrum, canle simplicî, foliis oppositis petiolatis ovato-deltoides acuminatis grossè dentato-serratis trinerviis, venulis pellucidis, corymbi terminalis ramosis ramulis pubescentibus, capitulis pedicellatis circ. 16-floris, invol. subcampanulatis squamis subimbricatis lanceolatis acutis subciliatis. ♀ in Mexici montanis inter Puerto de Varientes et Santa-Rosa de la Sierra. Cor. albæ. Caulis petiolique purpurei. Petiol. 7-8 lin. Limb. 3 poll. longus, 2 poll. latus.
107. *E. GLUTINOSUM* (Lam. dict. 2. p. 408), fruticosum, ramis teretibus sulcato-rugosis glabris apice glutinosi, foliis oppositis petiolatis cordato-lanceolatis acuminatis serratis suprâ bullato-rugosis glabris subtus tomentosis, corymbo oblongo paniculato terminali, capitulis 15-18-floris, invol. ovali-oblongi squamis imbricatis pubescentibus acutis. ♂ in Peruvia ex h. Juss. inter Mulalo et El Tambillo Quitensium (H. B. et Kunth 4. p. 131). Fol. 5-6 poll. longa, 1  $\frac{1}{2}$  lata. Cor. purpureæ. Arbor 2-3-org. cortice cinereo.
108. *E. SALVIA* (Colla pl. chil. f. 2. p. 8. n. 53 act. tur. 38. p. 8), fruticosum, ramis teretibus striatis, junioribus folisque novellis viscosis, foliis oppositis petiolatis lanceolatis acutis basi truncatis aut subattenuatis grossè crenato-dentatis rigidulis suprâ glabris bullulatis subtus nervis elevatis reticulatis pubescentibus, corymbo terminali composito conferto, capitulis pedicellatis 15-20-floris, invol. squamis 2-3-seriatis imbricatis lanceolatis acuminatis, ext. brevioribus pubescenti-viscosis, intimis lævibus coloratis. ♂ in Chili circa Valparaiso, Quillota, Rancagua, inter fruticeta et sylvas (Bert. Gay! etc.). An ex loco natali *E. reticulatum* Hook. et Arn. in Beech. bot. 1. p. 29? non Desv. Cor. violaceæ. Achæni. 5-costatum. Vulgò à Chilensibus dic. *Salvia Macho* ex Bertero. (v. s.)
109. *E. BACLEANUM*, suffruticosum pube brevi confertâ pubescenti-hirtellam, caulibus teretibus apice subaphyllis, foliis oppositis et hinc indè alternis in petiolum subattenuatis oblongis ovalibusve dentatis aut profundè incisis, paniculæ subcorymbosæ ramis alternis, capitulis pedicellatis circ. 16-20-floris, invol. squamis 2-3-seriatis laxè imbricatis lineari-oblongis obtusissimis apice membranaceis subciliatis, achænio angulato villosa. ♂ in Brasiliæ merid. ad

oram dictam Banda orientale legit cl. Bacle, et paulò post naufragium miserabile collectiones suas ferè totas amisit ! undè levis solatii modò hujus nomen huic stirpi miraculè salvatæ dedi. Etiam vidi inter Personas (h. Hænk.!) et sine loco (h. Née!). Species ex habitu cinerascens, foliis variè incisus facillè distincta. Pappus sordidus. (v. s.)

110. *E. xanthophyllum*, herbaceum ? caule erecto tereti infernè hispido folioso supernè nudo pulverulento-pubescente, foliis oppositis dorso et margine hispidis in petiolum longè attenuatis supernè pinnatipartitis, lobis pinnatifidis lobulis oblongo-linearibus subacutis, corymbi terminalis conferti ramis alternis approximatis, capitulis 18-20-floris pedicellatis, invol. squamis circ. 2-seriatis, ext. paucis parvis acutis, int. oblongis basi striatis apice submembranaceis obtusis puberis. ♀? in Brasiliæ prov. Rio-Grande. Affine *E. Bacleano* sed dist. caule foliisque setoso-hispidis, capitulis dimidio minoribus, etc. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 808 miss.)

111. *E. staviævolium*, herbaceum, caule erecto tereti hirsuto oppositè ramoso, foliis oppositis in petiolum attenuatis lanceolatis acuminatis serratis trinerviis supra sparsè puberis subtus pubescenti-hirsutis, corymbis axillaribus parvis et terminalibus fastigiatis, capitulis breviter pedicellatis 15-20-floris, invol. biserialis squamis ext. parvis paucis acutis, int. oblongis pubescentibus apice membranaceis obtusis. ♀? in Brasiliæ prov. Sancti-Pauli. Invol. apice subcolorata. Folia 15-18 lin. longa, 5 lin. lata. Cor. parvæ invol. ferè breviores. (v. s. in Mus. reg. Par. à Mus. imp. Bras. sub n. 496 miss.)

β. *angustius*, caule magis hirsuto, foliis angustioribus et longioribus (2 circ. pollices longis, 3 lin. latis). — in prov. Rio-Grande (h. Mus. imp. Bras. n. 806). An species diversa. (v. s. in h. Mus. reg. Par.)

112. *E. paulinum*, herbaceum, ramis divaricatis teretibus hispidis, foliis oppositis et alternis lanceolatis basi in petiolum attenuatis triplinerviis integris supernè acuminatis dentatis, paniculæ corymbose ramis alternis velutinis, capitulis ad apices ramulorum confertis pedicellatis 15-20-floris, invol. squamis biserialis, ext. paucis parvis acutis, int. oblongis basi striatis apice obtusis pubescentibus ciliolatisque submembranaceis, achænio glanduloso. — in Brasiliæ prov. Sancti-Pauli. Folia sesquipollicem longa, 4-5 lin. lata, summa subintegerrima. Species affinis *E. pubescenti* et *E. decemflora*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 498 miss.)

\*\*\*\* *Capitulis 20-floris.*

113. *E. Hænkianum*, fruticosum, ramis teretibus apice pubescenti-velutinis, foliis oppositis petiolatis latè ovato-rhombeis basi cuneatis trinerviis apice acuminatis integerrimis glabresculis, corymbo laxo, pedicellis longis bracteolatis, capitulis circ. 20-floris, invol. ovalis squamis imbricatis acutis vix pubescentibus, achæniis glabris. ♂ in Mexico legit cl. Hænke. Petiol. 9-12 lin. longi. Limb. fol. 3 poll. longus sesquipollicem latus. Folia referunt *Hedera helix* var. *integrifolia*. (v. s. in h. Hænke à cl. Sternb. comm.)

β? *velutinum*, ramis petiolis foliisque pubescenti-velutinis, capitulis circ. 30-floris, achæniis brevioribus glaberrimis. ♂ ad Acapulco legit cl. Hænke. An fortè species propria inter *E. Neeanum* et *E. Hænkianum* media? (v. s. in h. Hænke à cl. Sternb. comm.)

114. *E. hemisphaericum*, fruticosum erectum, ramis 3-4-angulatis petiolis pedicellisque pubescenti-hirtellis, foliis oppositis petiolatis ovatis acutis trinerviis dentatis glabresculis aut sparsè puberulis subtus crebrè glanduloso-punctatis, paniculâ laxâ subirregulari, capitulis longè pedicellatis 20-floris ferè hemisphaericis, invol. squamis 3-seriatis laxè imbricatis striatis ext. brevibus acutis int. longioribus ciliatis apice ad dorsum hirsutis corollas æquantibus, achænio glabro. ♂ verisim. in Brasiliâ legit cl. Lund! Habitus paulò *E. glechonophylli* sed capitula diversissima. Petiol. 7-8 lin. longi. Limb. fol. pollicem longus, 8-10 lin. latus. (v. s.)

β. *molliusculum*, foliis duplò triplòve majoribus vix glandulosis, ramis ter-

- retinaculis.  $\frac{3}{4}$  circa Rio-Janeiro (Lund!). Capitula exactè var.  $\alpha$ . Prioris videtur var. solo pinguiori et magis umbroso lecta. (v. s. cum.  $\alpha$ .)
115. *E. CACALIOIDES* (H. B. et Kunth nov. gen. am. 4. p. 130), fruticosum, ramis tetragonis striatis glabris viscosis, foliis oppositis petiolatis oblongo-lanceolatis acuminatis basi acutis profundè crenatis subcoriaceis suprà glabris subtùs tomentoso-pubescentibus, corymbis terminalibus trifidis ramosissimis, capitulis pedicellatis 20-floris, invol. campanulato-oblongi squamis laxè imbricatis striatis acutis ciliatis viscosis.  $\frac{3}{4}$  in agro Quitensi? Cor. violacea? Petiol.  $\frac{7}{8}$  lin. Limb. fol. 3 poll. longus, pollicem latus.
116. *E. XALAPPENSE* (H. B. et Kunth nov. gen. am. 4. p. 128), fruticosum glabrum, ramis sexangularibus striatis, foliis oppositis petiolatis ovato-ellipticis acutis serratis subtriplinerviis subcoriaceis, paniculis terminalibus ramosissimis foliatis, ramis subgeminis oppositis, capitulis pedicellatis ternis circ. 20-floris, invol. campanulatis squamis oblongo-lanceolatis acutis imbricatis subviscosis.  $\frac{3}{4}$  prope urbem Xalappam Mexicanorum. Cor. alba. Petiol.  $\frac{1}{2}$  poll. Limb. fol. 20 lin. longus, 10 lin. latus. Affine *E. glabrato*.
117. *E. RUGOSUM* (H. B. et Kunth nov. gen. am. 4. p. 114. non Schrad.), fruticosum, ramis teretibus molliter pubescentibus, foliis oppositis subsessilibus triangulari-ovatis acutis grossè crenatis coriaceis suprà bullato-rugosis scabris subtùs hispido-tomentosis et canescentibus, corymbis terminalibus ramosis densis, capitulis circ. 20-floris, invol. campanulato-cylindracei squamis imbricatis, ext. subrotundis, int. oblongis obtusis striatis ciliatis, achænio ad angulos scabriusculo.  $\frac{3}{4}$  in collibus apricis aridis inter urbem Alausi et Cerro de Sitzan prov. Quitensis. Cor. pallidè purpureæ. Folia 14-15 lin. longa, 13 lin. lata, petiol. 2 lin.
118. *E. SCHREBERIANUM* (Schrad.! ind. sem. h. Goett. 1832. p. 3), suffrutescens, caulis teretibus pubescentibus, foliis oppositis petiolatis ovato-lanceolatis acuminatis crenato-serratis basi obtusis truncatisve suprà sparè subtùs secus nervos scabriusculis, corymbis pedunculatis oligocephalis axillaribus terminalibusque, capitulis breviter pedicellatis circ. 20-floris, invol. squamis laxè imbricatis oblongo-linearibus apice scariosis glabris ciliatis obtusis mucronatisve, achænio acutè angulato secus nervos scabro.  $\frac{1}{2}$   $\frac{3}{4}$  in Mexico (Schrad.) circa urbem et ad Tampico de Tamaulipas (Berl.!). Cor. lilacinæ. Habitus ferè *E. iresinoidis*. (v. s.)
- E. virgatum*, foliis deltoideis imà basi cuneatis. — in Americà meridionali.
- E. virgatum* Schrad.! ind. sem. h. Goett. 1829 ex litt. 1832. Caulis 3-4-pedalis, cortice sanguineo-castaneo. (v. s. ex h. Goett. comm. à cl. Schrad.)
- $\gamma$ . *diversifolium*, foliis deltoideis longè acuminatis, ramorum juniorum subcordatis, summis subrotundo-ellipticis. — in Americà merid. sed ut var.  $\beta$  locus proprius incertus. *E. diversifolium* Schrad.! ind. sem. h. Goett. 1829. ex litt. 1832. Var.  $\beta$  et  $\gamma$  vix ab  $\alpha$  distinctæ ex specim. Schraderianis.
- Nomen var.  $\alpha$  selegi etiamsi paulò recentius, ob patriam certam. (v. s. ex h. Goett.)
119. *E. SALZMANNIANUM*, caule herbaceo tereti apice glabro cæterùm sparè pubero, foliis oppositis petiolatis elliptico-oblongis obtusis basi attenuatis rigidulis integerrimis penninerviis suprà scabris subtùs glabriusculis, paniculæ corymbosæ ramis oppositis elongatis subfastigiatis, capitulis breviter pedicellatis 20-floris, invol. squamis 3-4-seriatis imbricatis linearibus obtusiusculis dorso puberis, achænio minutè pubescente.  $\frac{1}{2}$  circa Bahiam Brasiliæ in collibus aridis legit cl. Salzmann. Caulis sesquipedalis. Invol. purpurascens. Cor. purpurea ex Salzmann sed ex sicco aurea videtur. Pappus albus. Folia 3 poll. longa, 10 lin. lata, petiolo 4 lin. (v. s.)
120. *E. VAUTHIERIANUM*, caule herbaceo erecto tereti oppositè ramoso tenuissimè pulverulento-velutino, foliis oppositis breviter petiolatis oblongo-lanceolatis utrinquè attenuatis subtriplinerviis distanter et acutè serratis suprà scabridis subtùs pubescentibus, paniculâ compositâ amplâ polyccephalâ, capitulis pedicellatis 20-floris, invol. squamis 3-4-seriatis laxè imbricatis striatis pube-

rulis et apice subciliatis, intimis acutis.  $\frac{1}{2}$ ? in Brasiliæ prov. Minarum Gener. prope Mariannam legit cl. Vauthier. Achænium glanduloso-puberulum. Folia 5-6 poll. longa, 1-2 poll. lata. (v. s. comm. à cl. Vauthier sub n. 274.)

\*\*\*\*\* *Capitulis 20-25-floris, foliis alternis.*

121. E. VINDEX, rhizomate crasso lignoso, caulibus pluribus herbaceis simplicibus erectis teretibus folisque puberulis, foliis paucis remotis parvis alternis ovali-lanceolatis subsessilibus integris, capitulis 7-8 subracemosis, invol. ovatis squamis subimbricatis striatis ovali-oblongis acutis disco brevioribus, achæniis pentagonis subhirsutis.  $\frac{3}{4}$   $\frac{1}{2}$  in Brasiliæ prov. Minarum Gener. in altis deserti Minar. Novarum legit cl. de Martins. E campis adustis invicta quotannis assurgit. Vernonia vindex Mart. ! Herb. Habitus omnino Vernoniæ, sed pappus et stigmata Eupatorii. Caules pedales. Folia 4-5 lin. longa. Cor. et styli purpurascens. Flores in inv. 20-25. (v. s. in h. Acad. reg. Monac.)

\*\*\*\*\* *Capitulis 25-30-floris, foliis oppositis.*

122. E. NERANUM, fruticosum, ramis teretibus apice pubescenti-velutinis, foliis oppositis petiolatis late ovatis acuminatis ferè triangularibus angulis infer. obtusis subcrenatis trinerviis anprà glabriusculis subtus pubescentibus, panicula subcorymbosæ ramis oppositis, capitulis pedicellatis obovatis circ. 25-floris, invol. squamis imbricatis pubescentibus, ext. acutis intimis obtusis, achænio angulato secus angulos scabro.  $\frac{3}{4}$  circa Acapulco legit cl. Née. Petioli 9-10 lin. longi. Limb. 2 poll. longus, 15 lin. latus. Cor. invol. longiores. (v. s. ex herb. Thibaud.)
123. E. TINIFOLIUM (H. B. et Kunth nov. gen. am. 4. p. 133), fruticosum glabrum viscosum, ramis subsulcatis, foliis oppositis petiolatis ovato-ellipticis acutis basi rotundatis crenato-serratis coriaceis nitidis, corymbis terminalibus oppositis ramosis, capitulis breviter pedicellatis 20-25-floris, invol. campanulati squamis imbricatis lanceolatis acutiusculis striatis, achænio glabriusculo.  $\frac{3}{4}$  in Novâ-Granatâ? Petiol. semipoll. Limbus fol.  $3\frac{1}{2}$  poll. longus, 2 poll. latus. Cor. alba.
124. E. STILLINGIAEFOLIUM, fruticosum glabrum, ramis teretibus, foliis oppositis petiolatis late ovato-lanceolatis acuminatis basi trinerviis obtusis serratis apice integerrimis subtus crebrè et minutè glandulosi, panicula composita corymbosâ, capitulis pedicellatis ovatis circ. 20-25-floris, invol. squamis paucis striatis imbricatis linearibus acutis pedicellatisque minutè pubescentibus, achænio angulato secus angulos scabrido.  $\frac{3}{4}$  in Mexico ad Tamaulipas legit cl. Berlandier. Cor. roseæ. Petiol. 9-12 lin. longi. Fol. limbi 3 poll. longi, 2 poll. lati. Affinis E. collino. (v. s. comm. à cl. Berland.)
125. E. MENDEZII, fruticosum, ramis teretibus apice breviter velutino-hispidulis, foliis oppositis petiolatis ovato-lanceolatis acuminatis basi èt apice integris medio serratis trinerviis anprà subpuberulis subtus petiolisque velutino-hispidulis, panicula amplâ trichotomâ ramosissimâ, capitulis pedicellatis oblongo-obovatis circ. 25-floris, invol. squamis imbricatis acuminatis dorso puberulis, achænio glabro.  $\frac{3}{4}$  in Leon Mexic. ad occid. urbem Guanaxusto legit cl. Mendez. Petiol. 6 lin. longi. Limbus 30-33 lin. longus, sesquipollicem latus. Frondescentia cinerea. (v. s.)
126. E. MALVÆFOLIUM, fruticosum?, ramis teretibus tenuiter cum pedunculis petioli nervisque foliorum pulverulento-velutinis, foliis oppositis longè petiolatis late cordiformibus breviter acuminatis grossè dentatis trinerviis hinc indè sublobis utrinquè glabriusculis, corymbis terminalibus compositis foliis intermixtis, capitulis pedicellatis circ. 25-floris, invol. squamis laxè imbricatis striatulis subacutis glabris, achænio glabriusculo, pappo tenui caducissimo.  $\frac{3}{4}$ ? in Mexico inter Vittoria et Tula legit cl. Berlandier. Petiol. 3-5 pollic. Limbi fol. 5-6 poll. longi, 4-5 lati. Cor. teretes brevissimè 5-dent. Styli longissimi. (v. s.)
- J. E. PULGOSIUM (H. B. et Kunth nov. gen. am. 4. p. 110), caule suffruti-

coeo volubili, ramis ferrugineo-tomentosis, foliis oppositis petiolatis oblongis acuminatis basi angustatis remotè denticulatis suprâ scabris subtus tenuissimè tomentoso-hirtis fuligineo-fuscescentibus, paniculis terminalibus et axillaribus pedunculatis divaricatis, capitulis subsessilibus approximatis circ. 25-floris, invol. semiglobosi squamis imbricatis oblongis obtusis ciliatis, recept. convexo, achenio ad angulos scabro.  $\frac{5}{8}$  in Andibus Quindensibus Novo-Granat. Cor. albæ? Petioli pollicares. Fol. limbus 6 poll. longus, 2-2  $\frac{1}{2}$  latus.

128. *E. ARBOREUM* (H. B. et Kunth nov. gen. am. 4. p. 131), arboreum, ramis sexangularibus glabris, foliis oppositis petiolatis oblongo-lanceolatis basi rotundatis crenato-denticulatis coriaceis suprâ glabris subtus tenuissimè pubescentibus, paniculis terminalibus subcorymbosis, capitulis pedicellatis sessilibus circ. 25-floris, invol. cylindraceo-campanulatis squamis imbricatis striatis acutis subciliatis subviscosis, achenio ad angulos scabro.  $\frac{5}{8}$  inter Pomalacta et Alami Quitensium. Petiol. 9 lin. long. Limb. fol. 3 poll. longus, 10-11 lin. latus. Cor. lilacina. Affine *E. cecalioidi*.

129. *E. DENDROXYPHOLUM*, fruticosum glaberrimum, ramis teretibus, foliis oppositis subpetiolatis basi cuneato-attenuatis oblongis obtusis integerrimis aut summis callosò-denticulatis rigidis reticulatis subtus punctulatis, paniculæ ramis oppositis apice subcorymbosis, capitulis pedicellatis circ. 25-floris, invol. squamis 3-4-seriatis imbricatis linearibus striatulis apice obtusis ciliatis dorso pubescentibus.  $\frac{5}{8}$  in Peruvia legit cl. Poeppig. Affine *E. Blanchetiano*. Folia ramea internodio multò breviora, 2-2  $\frac{1}{2}$  poll. longa, 7-9 lin. lata. Achenia minima glabra. Pappus tenuis albus. (v. s. comm. à cl. Poeppig sub n. 18.)

130. *E. DISCOLOR*, fruticosum, ramis sexangularibus glabris vix apice summo subpuberulis, foliis oppositis petiolatis oblongo-lanceolatis basi attenuatis apice acuminatis suprâ glabris subrugosis subtus densè cano-tomentosis margine junioribus angustè revolutis vix crenulatis, corymbo trichotome conferto terminati, capitulis ad apices ramulorum congestis subsessilibus 25-floris, invol. squamis laxè imbricatis pluri-seriatis acutis ciliatis, acheniis ad angulos subcabridis.  $\frac{3}{8}$  in Amer. calid. (Née!), in Peruvia montanis et in Huanacera circa urbem (Hank!). An *E. persicæ-folium* H. B. et Kunth sed folia basi non rotundata! An *E. salicinum* Lam. sed capitula non 10-12 sed 25-flora? Cæt. à foliis Persicæ et Salicis differt lanugine canâ paginæ infer. *E. discolor* Spreng. est *Vernonia*. (v. s.)

131. *E. SALICIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 131 non Sims), fruticosum volubile, ramis teretibus sulcatis glutinosis, foliis oppositis petiolatis oblongo-lanceolatis acuminatis basi rotundatis argutè serrato-dentatis coriaceis suprâ bullato-rugosis glabris subtus cano-tomentosis, corymbis terminalibus, capitulis pedicellatis et sessilibus 20-30-floris, invol. cylindraceo-campanulatis squamis imbricatis striatis acuminatis glabris acutis, achenio ad angulos scabro.  $\frac{5}{8}$  in frigidis ad declivitatem montis Chimborazo. Cor. violacea. *E. glutinoso* affine.

132. *E. GLABRIUSCULUM*, herbaceum erectum glabriusculum, ramis oppositis cum nervis fol. subtus pedicellisque minutissimè pubescentibus, foliis oppositis petiolatis ovato-lanceolatis longè acuminatis grossè et acutè serratis basi cuneatis triplinerviis membranaceis, paniculæ amplæ ramis ramisque oppositis, capitulis longè pedicellatis circ. 25-30-floris, invol. squamis 3-4-seriatis laxè imbricatis dorso minutè puberulis vix substriatis, intimis longioribus acutis, achenio glanduloso-pubero.  $\frac{7}{8}$  in Brasiliâ. Nimis affine *E. Vauthieriano* sed glabritie, petioli triplo longioribus, foliis longius cuneatis regul. serratis, etc. differre videtur. (v. s.)

133. *E. SUBROTUNDEUM*, fruticosum glabrum subviscosum, ramis teretibus, foliis oppositis breviter petiolatis oblongis obtusis basi attenuatis integerrimis rigidulis penninerviis, paniculæ ramis oppositis elongatis apice trifidis 15-20-cephalis, capitulis pedicellatis circ. 30-floris, invol. squamis 2-3-seriatis linearibus imbricatis dorso subpuberulis, intimis apice obtusis araneoso-ciliatis.  $\frac{3}{8}$  in Brasiliâ circa Bahiam legit cl. Blanchet (n. 1409!) et in sicco ad montem Sineri



prope Cayennam el. Leprieur! Cor. purpureæ. Pappus albusissimus. Styli longè exserti. Achenium secus costas scabridum. Folia 3 poll. et ultra longa, pollicem circ. lata. (v. s.)

134. *E. OXYCLÆNUM*, suffruticosum, ramis teretibus hispidis, foliis oppositis integerrimis aut apice subrepando-dentatis obtusis rigidulis supra scabris subtus pubescenti-hirtis penninerviis, inf. ellipticis in petiolum attenuatis, super. oblongis basi cuneatis subsessilibus, paniculæ ramis oppositis apice 5-6-cephalis, pedicellis glanduloso-pilosis, capitulis pedicellatis circ. 3o-floris, invol. squamis 2-3-seriatis imbricatis linearibus acuminatissimis corollas paulò superantibus dorso glanduloso-puberis.  $\frac{5}{8}$  in Bras. prov. Minarum Gener. prope Mariannam legit cl. Vauthier (n. 280!). Affine prior. Cor. purpureæ. Rami rubentes. (v. s.)
135. *E. FRUSCÆPOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 130), fruticosum, ramis sexangularibus glabrisculis, foliis oppositis petiolatis oblongo-lanceolatis acuminatis basi rotundatis crenulatis subcoriaceis supra glabris subtus cano-tomentosis, corymbis terminalibus, capitulis pedicellatis sessilibusve multi-? floris, invol. cylindraceo-campanulati squamis laxè imbricatis striatis acutis ciliatis.  $\frac{5}{8}$  ad radices montis ignivomi Cotopaxi Quitensium prope Malalo. Valdè affine *E. cecalioidi* dicitur.
136. *E. CORDIFOLIUM* (Swartz prod. III. flor. 1309), fruticosum, ramis teretibus glabris, ramulis hirsutis, foliis oppositis brevissimè petiolatis cordiformibus acuminatis serratis supra sparè hispidis subtus præsertim secus nervos majores hirsutis et granulis conspersis, corymbo terminali sessili, capitulis ad apices ramulorum confertis subsessilibus multifloris, invol. squamis imbricatis linearibus acutis semireflexis hirtis, achenio lineari.  $\frac{5}{8}$  in rupibus calcareis et præcipitiis montium Jamaicae. Hirsuties ferrugineæ. Folia 2 poll. longa,  $\frac{1}{2}$  poll. lata. Recept. convexum. (v. s. sine fl. ex h. Swartz.)
137. *E. GRACILE* (H. B. et Kunth nov. gen. am. 4. p. 124), fruticosum procumbens repens, ramis ascendentibus hirtis, foliis oppositis breviter petiolatis ovatis acutiusculis crenato-serratis glabris subtrinerviis, pedunculis axillaribus et terminal. 3-cephalis, capitulis pedicellatis multifloris, invol. campanulati squamis imbricatis lanceolatis acutis ciliatis glabris, achenio hispidulo.  $\frac{5}{8}$  in temperatis juxta Guaduas Novo-Granat. Cor. alba? Petiolus vix lin. longus. Folia semipollicaria.
138. *E. EPILOBRIDEZ* (H. B. et Kunth nov. gen. am. 4. p. 125), fruticosum erectum glabrum, ramis teretibus lævibus, foliis oppositis subsessilibus ovato-oblongis acutiusculis remotè crenulatis trinerviis, pedunculis axill. et termin. 3-cephalis, capitulis pedicellatis multifloris, invol. campanulati squamis imbricatis oblongis acutis ciliatis, achenio ad angulos scabro.  $\frac{5}{8}$  in temperatis Novæ-Granatæ? Affine *E. gracili* dicitur.
139. *E. ELEGANS* (H. B. et Kunth nov. gen. am. 4. p. 133 non Vent.), fruticosum, ramis teretibus piloso-tomentosis, foliis oppositis petiolatis elliptico-ovatis ovatis acutiusculis crenatis coriaceis utrinquè scabrisculis subtus in nervis venisque pubescentibus, corymbis terminalibus congestis, capitulis subsessilibus multifloris, invol. campanulati squamis imbricatis lanceolatis obtusiusculis pilosis, achenio glabro.  $\frac{5}{8}$  circa urbem Quito? Petiol. 2-3 lin. Limbus fol. 15 lin. longus, 8 lin. latus. Cor. alba?
140. *E. ARGUTUM* (H. B. et Kunth nov. gen. am. 4. p. 121), caule herbaceo striato hirtello, foliis suboppositis petiolatis lanceolatis acutis argutè serratis utrinquè tenuissimè glanduloso-hirtis, capitulis 1-2 longè pedunculatis terminalibus semiglobosis multifloris, invol. squamis subæqualibus 2-3-seriatis imbricatis hirtellis linearibus apice subulatis, achenio glabro.  $\frac{2}{3}$  ad radicem montis Chapoltepec Mexicanorum. Cor. violacea supernè hirta. Petiol. semipoll. Limb. fol. sesquipollicem longus, 4 lin. latus.
41. *E. SORDIDUM* (Less. in Linnaë 1831. p. 403), suffruticosum, ramis teretibus sordidè lanuginoso-tomentosis, foliis oppositis petiolatis membranaceis supra

basin quintuplinervio-penninerviis ellipticis utrinquè obtusis supra hispidiusculis subtus tomentoso-pubescentibus, paniculis subcorymbosis trichotomis, capitulis pedicellatis multifloris, invol. campanulati squamis 3-seriatis linearibus acuminatis villosa-pubescentibus.  $\frac{3}{4}$  in Mexici Cuesta-Grande de Chiconquisaco et in sylvis Jalapensis. Cor. purpureo-violaceæ. Fol. 2-5 poll. longa,  $2\frac{1}{2}$ - $3\frac{1}{2}$  lata.

\*\*\*\*\* *Capitulis 30-100-floris, foliis oppositis alternisqve.*

142. *E. KARVINIANUM*, fruticosum, ramis teretibus vix apice puberulis, pediculis et præsertim petiolis hispidis, foliis oppositis petiolatis ovato-lanceolatis acuminatis argutè et parcè serratis basi ciliatis glabriusculis, corymbis ramosis terminantibus 8-12-cephalis, capitulis longè pedicellatis 35-40-floris, invol. campanulati laxè imbricati squamis oblongo-linearibus obtusis ciliatis, achenio glabro.  $\frac{3}{4}$  in Mexico legit cl. Karvinski. Affine *E. adenophoro*. Pili petiolorum articulati. Petiol. 4 lin. longi. Limb. fol. pollicem longus, 5-6 lin. latus. Sryli longè exserti. (v. s. in h. Acad. reg. Monac.)
143. *E. AMPHYDICTYUM*, caule à collo lignoso simplici herbaceo tereti confertè hirtello à medio ad apicem aphylo, foliis alternis sessilibus obovato-oblongis basi cuneatis obtusis integris apice serratis acutis utrinquè puberulis et æqualiter nervis elevatis reticulatis, corymbo 7-8-cephalo conferto terminali, invol. squamis omnibus acuminatis ext. vix puberulis, flor. 30-35, acheniis minutissimè scabridis.  $\frac{1}{2}$  in campis editis prov. Sancti-Pauli nov. flor. legit cl. Lund. Herba pedalis. Affinis *E. subalternifolio* et *dyctiophyllo* sed florum numero (30 nec 5) evidenter diversum. (v. s.)
144. *E. BULLOTÆFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 121), caule herbaceo erecto striato pubescenti-hirto, foliis alternis petiolatis ovato-triangularibus cordatis acutis inciso-crenatis subtrinerviis supra hispidis subtus hispidopubescentibus, corymbis terminalibus ramosissimis, capitulis pedicellatis 36-40-floris, invol. cylindraceo-campanulati squamis imbricatis linearibus acuminatis glabris, ext. brevibus pubescentibus, achenio glabro.  $\frac{1}{2}$  in Novâ-Granatâ? Cor. violacea. Petiol. 7 lin. longi. Fol. subpollic.
145. *E. LAMIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 126 non Link.), caule herbaceo? volubili? striato hirto, foliis oppositis petiolatis ovatis cordatis angustato-acuminatis grossè crenatis rigidis supra bullato-rugosis et scabriusculis subtus hirtis, corymbis terminalibus, capitulis pedicellatis circ. 40-floris, invol. cylindraceo-campanulati squamis imbricatis striatis ciliato-pubescentibus, ext. acutis, int. lanceolatis, achenio glabro.  $\frac{1}{2}$ ? in agro Quitensi? Cor. violacea?
146. *E. VISCOSUM* (H. B. et Kunth nov. gen. am. 4. p. 129), fruticosum glabrum viscosum, ramis sexangularibus, foliis oppositis petiolatis obovato-oblongis acutis crenato-serratis coriaceis glanduloso-punctatis, corymbis terminalibus ramosissimis diffusis subfastigiatis, capitulis pedicellatis circ. 40-floris, invol. campanulati squamis laxè imbricatis lanceolatis acutis, achenio ad angulos hispido.  $\frac{3}{4}$  in agro Quitensi? Cor. violacea. Petiol. 5 lin. Limbus fol.  $3\frac{3}{4}$  poll. longus, 16 lin. latus.
147. *E. FRATERUM*, fruticosum glabrum, caule tereti, foliis oppositis vix petiolatis ovalibus integerrimis mucrone calloso subacutis rigidis penninerviis reticulatis, sup. internodio brevioribus, paniculæ ramis oppositis polycephalis, sup. corymbosis, capitulis pedicellatis 40-floris, invol. squamis 2-3-seriatis subimbricatis linearibus dorso et ad apicem margine subvillosis, intim. mucronatis, ext. subobtusis.  $\frac{3}{4}$  in Amer. merid. (fortè Peruvia?) legit cl. Poeppig. Proximè accedit ad *E. Blanchetianum*, sed folia magis distantia, 3 poll. longa,  $1\frac{1}{2}$  lata, capit. 40-flora, achenia tenuia puberula. (v. s. comm. à cl. inv. sub D. 2075.)
148. *E. VITALBÆ*, fruticosum scandens glabrum, ramis teretibus junioribus pubescentibus, foliis oppositis petiolatis ellipticis acutis seu plerisque ovato-lanceolatis acuminatis subdentatis quintuplinerviis, paniculæ ramis oppositis

161. *E. DELTOIDEUM* (Jacq. h. schoenbr. 3. p. 63. t. 369), herbaceum erectum, caule teretibus pulverulento velutino, foliis oppositis petiolatis hastatis triangularibus lobis apiceque acuminatis trinerviis inæqualiter serratis subtus pubescentibus, paniculâ thyrsoidæ apice subcorymbosâ, capitulis pedicellatis circ. 40-floris, invol. squamis linearibus acutis subuniseriatis, achenio puberulo.  $\frac{1}{2}$  in Mexico ad Villalpando (Mend.!), in valle Toluca (Berl.), in Yxtapalapan montibus (Moq.). Bonpl. nav. 1, p. 38. t. 14. *E. triangulare* Moq. fl. mex. ic. Mexicanis dicitur *Xolochichil* ex Moq. Cor. pallide purpureæ. (v. v. et s.)
162. *E. PETIOLARE* (Moq. fl. mex. ic. ined.), fruticosum, ramis teretibus pubescenti-velutinis, foliis oppositis longè petiolatis cordiformibus acuminatis crenatis basi trinerviis supra sparse puberulis subtus pubescenti-velutinis, panicula ramis axillaribus et terminalibus folio æqualibus polycephalis, capitulis pedicellatis circ. 35-40-floris, invol. squamis linearibus acuminatis subuniseriatis pubescentibus, achenio gracili pubero.  $\frac{1}{2}$  in Mexico. Bustamenta cordata Alam. in litt. 1831. Petiol. sesquipollic. Limbus fol. 3 poll. longus, 2-2  $\frac{1}{2}$  latus. Cor. alba basi attenuata. (v. s. comm. à cl. Alaman et Mairet et in Andr. pl. mex. ined. n. 279.)
163. *E. LOWICEROIDES* (H. B. et Kunth nov. gen. am. 4. p. 116), fruticosum, ramis subviscosis, foliis oppositis subsessilibus elliptico-oblongis obtusis subintegerrimis subcoriaceis glabris, paniculis terminalibus patulis, capitulis breviter pedicellatis 30-40-floris, invol. campanulatis squamis lanceolato-linearibus acutis subviscosis ciliato-pubescentibus, achenio ad angulos hispidulis.  $\frac{1}{2}$  in temperatis Andium Novo-Granatensium. Cor. purpurea. Fol. 2 poll. longa, pollicem lata. Affine *E. amygdalino*.
164. *E. CORSAQUINÆUM*, suffruticosum, caule erecto tereti cum petioliis pedunculisque tenuissimè pubescente, foliis oppositis petiolatis ovato-triangularibus vix obtusè acuminatis grossè crenatis basi truncatis trinerviis membranaceis supra æcus nervos subtus ubique pube minimè velutinis, ramis floridis axillaribus oligocephalis folio brevioribus, capitulis pedicellatis circ. 30-floris, invol. squamis 1-2-seriatis linearibus acutis dorso subpuberulis achenio angulato glabriusculo.  $\frac{1}{2}$  in Brasiliâ circa Rio-Janeiro legit cl. Gaudichaud. Petioli pollicares. Limb. fol. 3 poll. longus, 2 poll. latus. Affine *E. sordido* sed multò minùs velutinum, longius petiolatum, folia latiora magis deltoidea, rami floridi breviores, etc. (v. s. comm. à cl. Gaudichaud.)
165. *E. TRISTE*, fruticosum, ramis teretibus cum petioliis pedunculisque pube ferrugineâ densè hirsuto-velutinis, foliis oppositis petiolatis latè ovatis acuminatis basi obtusissimis crenato-dentatis supra scabridis subtus velutino-hirsutis, corymbi ramis primariis oppositis, capitulis longè pedicellatis 25-floris, invol. vix biserialis squamis linearibus acutis dorso velutinis, acheniis angulatis glandulos.  $\frac{1}{2}$  in Jamaicâ legit cl. Bertero. Eup. cranthifolium Spreng. syst. 3. p. 418. excl. syn. et patr. primâ. Petioli æmipollicares. Limb. fol. 2 poll. longus, 1  $\frac{1}{2}$  latus. Cor. faux magis inflata quàm in cæteris. Antheræ odore perispermæ. (v. s. comm. à cl. Bertero et Balbis.)
166. *E. ARALIEFOLIUM* (Less. in linnæa 1831. p. 403), fruticosum scandens, ramis glabris junioribus subancipitibus, foliis oppositis petiolatis oblongo-ellipticis obovatis integerrimis penninerviis carnosissimis, panicula supra decompositis trichotomis, capitulis 23-28-floris, invol. squamis pauciserialis glabris, intim. linearibus acuminatis.  $\frac{1}{2}$  in sylvis Nisanthe Mexicanorum.
167. *E. OLIGOCEPHALUM*, caulibus basi fruticulosâ à rhizomate lignoso plarimis erectis teretibus pulverulento-subvelutinis, foliis oppositis petiolatis ovato-lanceolatis acutiusculis serrulatis glabriusculis, capitulis paucis corymbosis longè pedicellatis circ. 25-floris, invol. squamis 1-2-seriatis oblongis acuminatis glabris, achenio angulato glabriusculo, pappo pauciseto caducissimo.  $\frac{1}{2}$   $\frac{1}{2}$  in Mexico legit cl. Hænke. Herba suffruticulosa vix pedalis. Radix teres lignosa simplex. Folia intermodi breviora, cum petiolo bilineari 9 lin. longa, 3 lin. lata. (v. s. in h. Hænke à cl. Sterub. comm.)

168. *E. MAIRETIANUM*, fruticosum, ramis teretibus glabrisculis, foliis oppositis petiolatis ovato-lanceolatis acuminatis minutè serrulatis triplinerviis glabris subtus albidis, paniculæ ramis oppositis apice corymbosis, capitulis pedicellatis campanulatis circ. 25-floris, invol. squamis linearibus acutis 1-2-seriatis, acheniis angulatis glabris.  $\frac{3}{4}$  in Mexico. Bustamentæ sp. Mair. ! Folia ferè Populi. Petiolus ferè pollicaris; limbus 3 poll. longus,  $1\frac{1}{2}$  latus. (v. s. comm. cum pl. aliis Mexic. à cl. Mairet.)
169. *E. BERLANDIERI*, fruticosum glabrum, ramis teretibus striatis, foliis oppositis petiolatis ovatis acuminatis basi et apice integris medio crenas paucas gerentibus utrinquè eglandulosis, paniculæ ramis oppositis apice corymbosis, capitulis pedicellatis circ. 25-floris, invol. squamis subbiseriatis linearibus acuminatis, achenio angulato ad angulos vix scabro, pappo corollæ ferè duplò brevioris.  $\frac{3}{4}$  in Mexico inter Tula et Tampico legit cl. Berlandier. Affine *E. ageratifolio* var.  $\beta$  fortè ab  $\alpha$  distincta. Petiol. 4 lin. longi; limbi fol. sesquipollic. longi, 9 lin. lati. Cor. albæ. (v. s. comm. à cl. Berland.)
170. *E. GRANDIDENTATUM*, caule herbaceo tenuissimè pulverulento-velutino, ramis oppositis erectis elongatis, foliis oppositis petiolatis ovato-lanceolatis acuminatis basi obtusis subquintuplinerviis grossè dentato-serratis glabriusculis internodio brevioribus, corymbis pluribus fastigiatis trichotomis, capitulis pedicellatis 25-floris, invol. squamis subbiseriatis linearibus acutis, achenio 5-gono vix scabrido.  $\frac{1}{2}$ ? in Mexico legit cl. Luc. Alaman. Petioli 9-12 lin. longi. Limbus fol. 3 poll. longus,  $1\frac{1}{2}$  latus. Juga foliorum distantia. Cor. ex sicco albidæ. Achenia parva nigra. Affine ex descr. *E. Pasquarensi*. (v. s.)
171. *E. STERNBERGIANUM*, caule herbaceo? erecto tereti glabro apice minutè puberulo, foliis oppositis petiolatis latè ovatis vix acuminatis grossè dentatis suprà glabriusculis subtus petiolisque subpuberulis, corymbis compositis confertis, capitulis pedicellatis 25-30-floris, invol. squamis biseriatis linearibus glabriusculis, achenio pentagono ad angulos scabro. — in Peruviam collibus Cordillerianis legit cl. Hænke. Petiol. 5-6 lin. longus, limbus 1-1  $\frac{1}{2}$  poll. longus, 10-13 lin. latus, imò basi trinervius, supernè penninervius. Affine *E. Dombeyano*. (v. s. in h. Hænk. ab ill. com. de Sternberg comm.)
172. *E. DOMBEYANUM*, fruticosum glaberrimum, ramis teretibus, foliis oppositis petiolatis ovatis acuminatis serratis longè suprà basin triplinerviis, paniculà tripartitâ, ramis divergentibus ramosis laxis, capitulis longè pedicellatis circ. 25-floris, invol. squamis biseriatis sublinearibus acuminatis vix subciliatis apice ferè scariosis, achenio glabro angulato.  $\frac{1}{2}$  in Amer. merid. legit cl. Dombey, sed locus propr. in ejus herb. non indic. Petioli 4 lin. Limb. fol. 2  $\frac{1}{2}$  poll. longus, pollicem latus. Cor. ex sicco purpureascentes. (v. s. comm. à Mus. reg. Par.)
173. *E. PANICULATUM* (Schrad. ! ind. sem. h. Gætt. 1832. p. 3. diss. Blum. p. 44). caule herbaceo tereti glabriusculo, foliis inf. et ramis oppositis super. et ramis alternis petiolatis ovato-triangularibus basi truncatis aut breviter cuneatis grossè et obtusè dentatis membranaceis glabriusculis, corymbo laxo foliis parvis intermixto, capitulis pedicellatis circ. 25-floris, invol. squamis oblongo-linearibus mucronatis glabris 1-2-seriatis subæqualibus, achenio angulato glabriusculo. ☉ in Brasiliâ ad sepes circa Bahiam legit cl. Salzmann, etiam vidi ex insulâ Trinitatis ortum. Habitus in sicco *E. Schiedeani* sed ex cl. invent. annum herbaceum. Invol. albida. Cor. minimè pallidè rubentes. *E. microste-mon* Casa. dict. 25. p. 432. Habui etiam sub nomine *E. paniculati* Bernh., sed descr. editam non reperio. (v. s.)
174. *E. SORDIDUM*, fruticulosum, caule erecto tereti cum petiolis pedunculisque velutino-hirtello, foliis oppositis petiolatis ovato-lanceolatis acuminatis dentatis trinerviis suprà sparse scabris subtus pubescenti-velutinis pallidioribus, paniculæ ramis folio paulò longioribus ultimis corymbosis, capitulis pedicellatis 25-floris, invol. squamis 1-2-seriatis linearibus acutis dorso puberis, achenio secus costas scabrido.  $\frac{3}{4}$  in Brasiliâ circa Rio-Janeiro legit cl. Lund. An *E. lamiifolium* Link. enum. h. Ber. 2. p. 306 (non Kunth) huc refer. ? Petiol.

- 9-10 lin. longi. limb. fol. 3 poll. longus, pollicem latus. (v. s. comm. à cl. Lond.)
175. *E. strumosum* (Lindl.) a herb. Mus. reg. Berol.; fruticosum, ramis teretibus glauca apice rubro-velutino-hispida, foliis oppositis petiolatis ovalibus basi sub-attenuatis apice longae et angustae acuminatis subintegerrimis penninerviis supra glabris subtus ramis velutino-hispidis. paniculae amplae ramis oppositis polycephalis. capitulis pedicellatis 20-25-floris, invol. squamis circ. biserialis linearibus acuta basi puberulis corollis multo brevioribus, achenio glabro.  $\frac{3}{4}$  in montibus prope Rio-Janeiro Brasiliae legit cl. Lindl. Petiol. 1  $\frac{1}{2}$ -2 poll. longi. Limb. fol. 1 - 2 poll. longus, 2 - 3 poll. latus. Cor. ex sicco purpurascens. Pappus annuus. v. s. in n. cit.
176. *E. maynensis*. fruticosum, ramis teretibus velutino-hispidis, foliis oppositis breviter petiolatis ovali-oblongis acuminatis tripplinerviis à medio ad apicem serratis supra scabris subtus reticulatis cinereo-pubescentibus, nervis petiolo-que villosis. corymbo terminali conferto vix pedunculato composito, capitulis 20-25-duris pedicellatis, invol. squamis linearibus 1-2-seriatis acutis denum pinnatis. achenio angulato vix pubero.  $\frac{3}{4}$  in Mexico ad Villalpando legit cl. Mendez. Fil. incl. petiolo 1-2 lin. sesquipollicem longa, 5-6 lin. lata. Cor. ex sicco pallide purpurascens. v. s. comm. à cl. Mendez.)
177. *E. lucayanum*. suffrutescens, ramis teretibus dense villosa-velutinis, foliis oppositis petiolatis ovato-lanceolatis acuminatis excepto apice grossè et inaequaliter serratis supra glabrescentibus subtus petiolisque velutinis subvillosis, corymbo terminali conferto, capitulis pedicellatis 20-25-floris, invol. squamis linearibus acuminatis, achenio costato glabrescentibus.  $\frac{3}{4}$  in Mexico ad Montemorelia Nova Letras prov. legit cl. Berlandier. Cor. corollae. Petiol. 6 lin. longi. Limbus fol. pollicem longus, 3 lin. latus. (v. s.)
178. *E. Boissierianum*. suffrutescens, ramis teretibus apice pilosulis, foliis oppositis petiolatis ovatis basi cuneatis apice acuminatis medio dentatis triner-vis supra sparse puberulis subtus secus nervos villosis, corymbo trichotomo, capitulis ad apicem ramorum confertis breviter pedicellatis circ. 20-floris, invol. squamis linearibus subnerviatis mucronulatis margine scariosis, achenio angulato puberulo — in Mexico Bustamenta ovata Mairet! in litt. Cor. alba. Petioli pollicares. Limbus 28-30 lin. longus, 14-15 lin. latus. (v. s.)
179. *E. pauciflorum* (H. B. et Kunth nov. gen. am. 4. p. 123), caule her- baceo substriato hirsuto subglutinoso, foliis oppositis petiolatis ovatis acutis serratis utrinque pilosis, capitulis longe pedunculatis terminalibus sub- corymbosis multifloris, invol. campanulatis squamis lanceolatis apice subulatis ciliatis, exter. pubescenti-hirsutis.  $\frac{2}{3}$  in Mexico juxta pagum Ario, in decliv. montium Aguacacensium et in monte Orizaba. Cor. violacea? Petiol. 4 lin. l. sub. fol. 12 lin. longus, 5-6 latus.
180. *E. Pazcuarensis* (H. B. et Kunth nov. gen. am. 4. p. 123), caule her- baceo erecto pubescenti-hirsuto, foliis oppositis petiolatis subrotundo-ovatis breviter acuminatis leviter cordatis grosse subduplicato-serratis subtripplinerviis supra scabrescentibus subtus in venis pubescentibus, corymbis terminalibus fastigiatis, capitulis longe pedicellatis multifloris, invol. subcampanulatis squamis acutis subciliatis, exter. pubescentibus, achenio ad angulos hispido.  $\frac{2}{3}$  in temperatis montibus Mexici ad lacum Pazcuarensium. Cor. alba? Affine *E. ageratoidi* di- citur.
181. *E. vallincola*, caule suffrutesco erecto tereti, ramulis petiolisque pilos- tulus articulatos gerentibus, foliis oppositis longe petiolatis subcordatis acutis triangularibus (angulis inf. obtusis) basi trinerviis grossè dentatis utraq. hispidulis, capitulis 5-13 in corymbum compositum fastigiatum dispo- sicellatis 20-25-floris, invol. squamis linearibus 1-2-seriatis disco bre- vi, acheniis pentagonis ad angulos puberulo-scabris.  $\frac{3}{4}$  in vallibus Andium legit cl. Hanke. Petioli circ. 10-12 lin. longi. Limbus fol. pollicem 10-12 lin. latus. Cor. ex sicco purpurascens. (v. s. in h. Hanke à cl. Jberg comm.)

182. *E. PAUPERTATUM* (H. B. et Kunth nov. gen. am. 4. p. 121), caule suffruticoso? tereti piloso, foliis oppositis petiolatis triangularibus acutis basi truncatis grosse crenato-serratis subtrinerviis supra pilosiusculis subtus molliter piloso-pubescentibus, corymbo terminali simplici, capitulis pedicellatis multifloris, invol. campanulati squamis lanceolatis apice subulatis striatis ciliatis, ext. paulò brevioribus, achenio glabro.  $\zeta$ ? in Novâ Granatâ? Fortè idem ac *E. microphyllum* L.

\*\* Capitulis 20-floris.

183. *E. MEAVOSUM* (Swartz! prod. III. flor. 1307), fruticosum, ramis teretibus erectis, ramulis pubescentibus, foliis oppositis petiolatis ovatis acuminatis grosse et acutè dentatis triplinerviis rugoso-nervosis glabris subtus globuliferis, corymbo composito polycephalo fastigiato, capitulis pedicellatis 20-floris, invol. squamis linearibus acuminatis 8-10 subuniseriatis.  $\zeta$  in altis montibus Jamaicæ (Sw.!), Sancti Domingi (Mack. in h. Lindl.!) Ramuli rubentes. Petioli puberi 3 lin. longi. Fol. 2 poll. longa, 10-12 lin. lata, infra nervos 2 laterales, nervulos 2 minores è nervo medio emittentia. Capitula parva. Cor. albidæ. Achenia minima. (v. s. ex h. Swartz.)
184. *E. PAPANTLENSIS* (Less. in linnæa 1831. p. 403), fruticosum, ramis teretibus junioribus pubescentibus, foliis oppositis membranaceis triplinerviis ovatis basi cordatis acuminatis serrato-crenatis non glabris, paniculis fastigiatis myriocephalis, capitulis pedicellatis 20-floris, invol. campanulati squamis biseriatis linearibus acuminatis glabris, corollis tubulosis.  $\zeta$  in Mexico prope Papantla. *E. leiophyllo* proximum. Petioli, pubes foliorum, achenia indeser.  $\dagger$ .
185. *E. AREOLARE*, fruticosum, ramis teretibus petioliis pedunculis nervisque foliorum subtus pube rufâ brevi confertâ velutinis, foliis oppositis petiolatis ovato-lanceolatis acuminatissimis rigidis basi obtusis 3-nerviis supra glabris minutissimè areolatis subtus puberis reticulatis, pedunculis axillaribus oppositis et subterminalibus trichotomis folio brevioribus, capitulis brevissimè pedicellatis 20-floris, invol. squamis 1-2-seriatis linearibus acutis, achenio pubescente.  $\zeta$  in Mexici montibus Guachilacensibus specimina pauca legit cl. Berlandier. Petioli 3 lin. longi. Fol. limb. 30 lin. longus 12-14 latus. Cor. ex sicco albidæ. (v. s.)
186. *E. SARVATUM*, fruticosum, ramis teretibus densè petiolisque velutinohispidis, foliis oppositis petiolatis ovato-lanceolatis acuminatis serratis trinerviis supra aparsè scabris subtus pubescenti-subvelutinis glandulosisque, paniculis axillaribus terminalibusque folio brevioribus oppositè ramosis, capitulis pedicellatis circ. 20-floris, invol. squamis linearibus acutis 1-seriatis dorso puberis, pappi setis corolla dimidiò brevioribus, achenio glabro.  $\zeta$  in Mexico. Genus novum Alam! in litt. 1831. Mair! exs. 1833. Cor. albæ invol. duplò longiores. Petioli 4 lin. longi; limbus 3 poll. longus, sesquipollicem latus. (v. s.)
- $\beta$ . foliis supra confertè scabris subtus velutino-tomentosis. — in Mexico. (v. s.)
187. *E. LÆVRE*, fruticosum glaberrimum, ramis teretibus substriatis, foliis oppositis petiolatis ovato-lanceolatis acuminatis apice obtusè serratis triplinerviis utrinquè lævigatis, paniculis ramosissimis ramos terminantibus, capitulis pedicellatis 20-floris, invol. squamis 1-2-seriatis linearibus acuminatis.  $\zeta$  inter frutes circa Rio-Janeiro leg. cl. Lond Gaudichaud et Lhotsky. Frutex 10-12 pedalia. Rami junior. subrubentes, adulti albid. Petioli. 3-5 lin. longi. Folia 3-5 poll. longa, 9-15 lin. lata. Pappus stramineus corollæ longit. Achen. angulatum glabrum. An fortè *E. Freyreii* Thunb. pl. bras. dec. 2. n. 19. ex florâ 1821. p. 382? sed descr. non suff. (v. s.)
188. *E. AYAPANA* (Vent. ! malm. t. 3. mag. enc. au VIII. v. 3. p. 76), caule basi suffruticoso ascendente ramoso glabro, foliis subsessilibus oppositis lanceolatis triplinerviis acuminatis subintegerrimis glabris, corymbo laxo oligocephalo, capitulis pedicellatis circ. 20-floris, invol. subuniseriatis squamis

201. *E. villosum* (Swartz! prod. III. flor. 1305), fruticosum, ramis teretibus ferrugineo-velutinis, foliis oppositis petiolatis ovatis acutis subseriatis trinerviis supra scabridis subtus ferrugineo-hirsuto-subtomentosis, corymbo trichotomo fastigiato polycephalo, capitulis pedicellatis 8-16-floris, invol. squamis 8-10 uni- seu biseriatis linearibus subpatulis pubescentibus, achenio angulato puberulo.  $\frac{3}{4}$  in frutetis montosis Jamaicae (Sw.!), Sancti-Domingi? (Willd. sp. 3. 1763? excl. syn. Sloan.). Hirsuties tota ferruginea. Petioli 2-3 lin. longi hirsuto-velutini; limb. fol. 12-15 lin. longi, 6-7 lati. Capitula parva. Cor. albidæ. (v. s. ex h. Swartz.)
202. *E. orbiculatum*, caulibus fruticulosis erectis simplicibus teretibus glabris, foliis oppositis brevissimè petiolatis orbiculatis dentatis basi trinerviis internodii longitudini plerisque subæqualibus utrinquè glabrescentibus, corymbo fastigiato ramis oppositis oligocephalis, capitulis pedicellatis 12-floris, invol. squamis 1-2-seriatis acutis oblongo-linearibus dorso subpuberulis.  $\frac{3}{4}$  in Brasiliæ prov. Sancti-Pauli. Petiolus lineam longus. Limb. fol. diam. 9-10 lin. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 507 miss.)
203. *E. fuscaulium*, suffruticosum, ramis teretibus tenuissimè puberulis, foliis oppositis petiolatis ovato-lanceolatis subacuminatis grossè dentatis apice integris basi trinerviis supra glabrescentibus subtus petiolisque tenuè velutino-puberulis, panicula ramis apice ramulosis, capitulis ad apices ramulorum confertis sessilibus circ. 15-floris, invol. squamis linearibus acutis 1-2-seriatis dorso subpuberulis, achenio breviter hispido.  $\frac{3}{4}$  in Sancto-Domingo legit cl. Bertero. *E. molle* Bert! herb. non Sw. nec Kunth. *E. salicinum* ex Hispaniâ Spreng. syst. 3. p. 412? excl. syn. Petioli 6 lin. longi, limb. fol. 2-2  $\frac{1}{2}$  poll. longus, pollicem latus. Cor. ex sicco albidæ. (v. s. comm. à cl. Bertero.)
204. *E. rhomboidum* (H. B. et Kunth nov. gen. am. 4. p. 127), fruticosum, ramis striatis glabris ramulis pubescenti-hirtis, foliis oppositis petiolatis rhomboideo-ovatis obtusiusculis crenatis subtriplinerviis supra pilosisculis subtus glabris, corymbis terminalibus, capitulis pedicellatis, invol. campanulati squamis oblongo-lanceolatis acutis hirsuto-pubescentibus.  $\frac{3}{4}$  in frigidis Mexici inter Toluca et Tianguillo. Petiol. 4 lin. Limb. fol. 15 lin. longus, 10 latus.
205. *E. triangulatum* (Alam.! in litt. 1831), caule herbaceo erecto simpliciter tereti purpureo glabro, foliis oppositis petiolatis ovato-triangularibus apice acuminate basi truncatis integris, angulis lateralibus rotundatis, lateribus dentato-serratis rigidis trinerviis reticulatis glabrescentibus, panicula subcorymbosa, pediculis pubescentibus, capitulis breviter pedicellatis 12-floris, invol. squamis subuniseriatis linearibus acutis, pappo corollâ dimidiò brevior.  $\frac{3}{4}$  à Mexico misit cl. Luc. Alaman cum plur. aliis rarissimis. Ex descr. valdè accedit ad *E. rubricaulis*, sed caulis teres nec hexagonus, fol. venæ obscuræ nec pellicidæ, folia rigidula nec membranacea, invol. non imbricata, etc. Cor. albæ invol. duplò longiores. (v. s.)
206. *E. cubense*, fruticosum, ramis teretibus apice pubescenti-velutinis, foliis oppositis petiolatis ovato-lanceolatis basi subtruncatis apice acuminate subdentatis trinerviis supra glabris subtus pubescentibus canescentibus glandulosisque, corymbo trichotomo fastigiato, capitulis 10-12-floris, invol. squamis 8-10 linearibus subpatulis pubescentibus, achenio angulato puberulo.  $\frac{3}{4}$  in ins. Cubâ prope Havanam legit cl. Ramon de la Sagra. Petioli hirsutis brevi canescentes, 2-3 lin. longi. Limb. fol. 15-16 lin. longi, 7-8 lati. Inflorescentia omnino *E. villosi*. (v. s. comm. à cl. de la Sagra.)
207. *E. cynanchioides*, fruticosum, ramis teretibus apice tenuissimè pulverulento-velutinis, foliis oppositis breviter petiolatis ovatis subcordatis obtusis trinerviis reticulato-venosis supra glabrescentibus subtus pulverulento-velutinis, corymbo trichotomo fastigiato, capitulis pedicellatis 10-12-floris, invol. squamis 8-10 ambiseriatis pubescentibus subacutis, achenio subangulato vix puberulo.  $\frac{3}{4}$  in ins. Cubâ juxta Havanam legit cl. Ramon de la Sagra. Folia cinereo-sulcanescentia petiolo 1-2 lin., limbo pollicem longo, 8 lin. lato. Infloresc. *E. cubensis*. Cor. albæ. (v. s. comm. à cl. de la Sagra.)

208. *E. LEROTHYLLUM* (Less. in *linnæa* 1831. p. 402), fruticosum, ramis teretibus junioribus pubescentibus, foliis oppositis membranaceis triplinerviis ovatis acuminatis basi obtusissimis profunde serrato-crenatis glaberrimis, paniculis subfastigiatis myriocephalis, capitulis pedicellatis 12-floris, invol. campanulati squamis biserialis linearibus acuminatis glabris.  $\frac{1}{2}$  in Mexico prope Papantla. Folia 12-21 lin. longa, 9-15 lin. lata, an sessilia? an petiolata? Affine *E. micrantho* dicitur, sed paniculae rami primaria folia sua superantes, corollae limbo amplo tubum superante.

209. *E. LUCIDUM* (Ort. dec. 35), fruticosum, ramis teretibus, junioribus breviter glanduloso-hirsutis, adultis glabris, foliis oppositis petiolatis ovatis utrinque breviter acuminatis integerrimis aut hinc inde serratis, supra sparse subcabridis, subtus præsertim secus nervos et petiolum hirtellis, corymbo terminali conferto, capitulis pedicellatis 10-12-floris, invol. squamis biserialis linearibus subobtusis extus subhirtellis, achenio angulato glabro glanduloso.  $\frac{1}{2}$  in Mexico (Ort.). Specimen habeo sub Ortegiano nomine à cl. Née lecto, sed locus non indic. Folia supra vix in sicco lucentia. Affine *E. calamintæfolio*. (v. s.)

210. *E. OSTRUSIACUM*, fruticosum, ramis teretibus hirsuto-villosis, foliis oppositis breviter petiolatis subcordatis ovato-rotundis obtusis supra minute velutinis subtus velutino-tomentosis subtriplinerviis reticulatis, corymbo terminali subcespiti conferto, capitulis breviter petiolatis 10-12-floris, invol. squamis linearibus subacutis puberis 2-seriatis subæqualibus, achenio angulato glabro.  $\frac{1}{2}$  in Sancto-Domingo legit cl. Bertero. *E. villosum* Bert. herb. non Willd. *E. domingense* Spreng. syst. 3. p. 412? (etiamsi folia alterna dicat). Petioli 2-3 lin. longi. Limb. fol. sesquipoll. longus latusque (v. s. comm. à cl. Balbis.)

211. *E. INCOMPTUM*, caule herbaceo erecto teretiusculo villosa-subtomentoso, foliis alternis fere sessilibus basi longè canecatis ovatis acutis triplinerviis supra scabridis subtus villosa-tomentosis, panicula elongata ramis alternis apice racemoso-paniculatis, capitulis ad apices ramulorum confertis brevè pedicellatis 10-12-floris, invol. squamis 1-2-seriatis linearibus obtusis pubescentibus, achenio angulato glabriusculo. — in Mexico legit cl. Hænke. Fol. infer. ignota, an opposita? media 3 poll. longa, sesquipollicem lata; super. longè acuminata. (v. s. in h. Hænke à cl. Sternberg comm.)

\*\*\*\* *Capitulis 6-10-floris.*

212. *E. AGERATIFOLIUM*, fruticosum glabrum, ramis teretibus, foliis oppositis petiolatis latè ovatis basi subtruncatis apice attenuatis obtusis grosse dentatis trinerviis eglandulosis, corymbis terminalibus trichotomis, pedicellis subviscosis vix puberulis, capitulis circ. 10-floris, invol. squamis linearibus biserialibus acutiusculis apice subciliatis, demum patulis, achenio puberulo ad angulos subcabrido.  $\frac{1}{2}$  in insulâ Cubæ circa Havanam legit cl. R. de la Sagra. *E. Havanense* H. B. et Kunth nov. gen. am. 4. p. 128? sed pedicelli non verè glabri, peduncululi trichotomi nec trifidi, corollæ pallide nec violaceæ, involucrium vix imbricatum, etc. — Petioli 9 lin. longi; limb. fol. 24-30 lin. longus, 20-22 latus. (v. s. comm. à cl. de la Sagra.)

$\beta$ ? *Mexicanum*, foliis magis ovatis basi subobtusis medio crenato-dentatis, pedicellis subpubescentibus, capitulis 12-13-floris.  $\frac{1}{2}$  in Mexico inter Victoria et Tula legit cl. Berlandier. *E. Berlandieri* formâ foliorum fere accedit, sed pappus corollam aequans.

$\gamma$ ? *Domingense*, foliis magis ovatis basi obtusis crenato-dentatis, pedicellis subpubescentibus, capitulis 18-20-floris.  $\frac{1}{2}$  in Sancto-Domingo legit Bertero. *E. ageratoides* Bert. herb. non Linn. Foliorum limbus sesquipollicem longus poll. latus. Invol. squamæ acutiores. An species propria? (v. s. comm. à cl. Balbis.)

213. *E. CRETIDIFOLIUM* (Lam. dict. 2. p. 406), fruticosum glabrum, ramis teretibus striatis, foliis oppositis petiolatis ovato-lanceolatis basi obtusis apice acu-



- minatissimis subseeratis basi subquintuplinerviis, paniculae amplae ramis oppositis polycephalis, capitulis pedicellatis 10-floris, invol. squamis 7-9 subuniseriatis linearibus acuminatis apice serratis 1-2 notatis demum patulis, achenio subpuberulo.  $\frac{5}{2}$  in ins. Caribæis (Lam.), in montibus Guadalupe (Perrottet!) et in Cubâ circa Havanam (La Osa!) Mikania verrucosa Spreng. syst. 3. p. 423? Verrucæ sunt lenticellæ omnibus Eupat. lignosis solemnes. Fol. 4-5 lin. longa,  $1\frac{1}{2}$  lata, petiolo 4 lin. longo. Cor. parvæ albidae. (v. s.)
214. E. SCHRADERI, frutescens, ramis teretibus striatis pubescenti-hirtellis, foliis infer. oppositis rameis alternis brevissimè petiolatis oblongo-lanceolatis basi cuneatis integerrimis à medio dentatis suprâ sparsè scabridis subtus pubescenti-hirtellis triplinerviis, corymbis terminalibus laxis, capitulis pedicellatis circ. 10-floris, invol. squamis subuniseriatis erectis lanceolatis acuminatis, achenio angulato puberulo.  $\frac{5}{2}$  in Americâ merid. sed locus proprius ign. E. rugosum Schrad. ! pl. rar. h. Gœtt. ined. ex litt. 1832. non Kunth nec Sprengel. Folia subrugosa 3 poll. longa, 6-9 lin. lata. Cor. albæ. (v. s. cult. ex h. Gœtt. à cl. Schrad. comm.)
215. E. POLYLOTOSUM, caule suffruticoso tereti vix tenuissimè pulverulento-velutino, foliis oppositis (alternisque?) petiolatis ellipticis crenatis subtus reticulatis utrinquè cinereo-subvelutinis, paniculae elongatæ ramis brevibus oligocephalis, capitulis pedicellatis 9-10-floris, invol. squamis linearibus obtusiusculis subuniseriatis pubescentibus, achenio angulato scabrido  $\frac{5}{2}$  in Chili circa Coquimbo leger. cl. Gandichaud et Macrae. Petioli 2-3 lin. longi. Limb. fol. 3 lin. longus, circ. 2 lin. latus. Folia ramea ad axillas conferta. Rami alterni. (v. s.)
216. E. DIFFUSUM (Vahl symb. 3. p. 94?), caule herbaceo erecto tereti vix pubescente, foliis oppositis petiolatis basi cuneatâ cordato-ovatis acuminatis obtusè et sæpè duplicato-serratis membranaceis basi trinerviis glabris subtus crebrè glandulosi, paniculae diffusæ ramosissimæ ramis inf. oppositis sup. alternis, pedicellis longis capillaribus, capitulis 7-8-floris, invol. squamis 1-2-seriatis linearibus acuminatis nitidis subscariosis demum reflexis, achenio tertiusculo pubescente. — in Americæ merid. (Vahl), prov. Santa-Marta (Bert.), ad fluv. Magdalenæ (Spreng. syst. 3. p. 419), in sylvis Jamaicæ interioris (Swartz flor. Ind. occ. 1301), in Hispaniolâ (Ham.). E. capillare Desv. in Ham. flor. Ind. occ. p. 51. E. flaccidum Spreng. in litt. ad Balb. Species habitu distinctissima sed corollæ ignotæ. An fortè hæc species duæ confusæ? (v. s. comm. à cl. Bert. et Balb.)
217. E. POLYNOTRYUM, caule fruticoso tereti ramosissimo tenuissimè cum pedunculis petiolis nervisque fol. pulverulento-velutinis, foliis oppositis rariis ternis et super. alternis petiolatis ovatis acuminatis subdentatis penninerviis glabrisculis, ramis axillaribus racemoso-paniculatis in paniculam longam foliosam dispositis, capitulis pedicellatis 6-8-floris, invol. squamis paucis 1-2-seriatis oblongo-ovalibus acutiusculis, achenio vix pubente.  $\frac{5}{2}$  in Mexico legit cl. Hænke. Petioli 4-6 lin. longi. Limb. fol. 3o lin. longus, circ. 2o lin. latus. Capitula parva innumerosa. Cor. ex sicco albidæ. Styli valdè elongati. Achenium exangulatum. (v. s. in herb. Hænke à cl. Sternb. comm.)
218. E. AURICULATUM (Vahl symb. 3. p. 95. t. 72. non Lam.), fruticosum, ramis teretibus striatis tomentoso-canis demum calvis, foliis alternis petiolatis ad basin petioli utrinquè rotundè auriculatis ovatis acutis denticulatis suprâ glabris subtus cano-tomentosis, apicis plurimis paniculatis elongatis, capitulis sessilibus distantibus 7-floris, invol. squamis 8 fusco-ferrogineis basi squamulis parvis calyculato.  $\frac{5}{2}$  in Brasiliâ.
219. E. ORGYALE, herbaceum altissimum, ramis hexagonis vix apice summo puberulis, foliis oppositis petiolatis latè ovato-lanceolatis basi cuneatis apice acuminatis grossè dentatis penninerviis reticulatis suprâ sparsè scabridis subtus pubescenti-hirtellis et glandulosi, corymbo trichotomo terminali confertissimo, capitulis ad apicem ramulorum densè adgregatis sessilibus 6-floris, invol. squamis 2-ser. oblongis obtusis ext. brevioribus subpubescentibus, int.

longioribus glabris.  $\frac{1}{2}$  circa Bahiam (Blanch. ! n. 1923), in montosis circa Rio-Janeiro frequens ex cl. Lund! Herba 6-7-pedalis. Rami oppositi. Flor. albi odoratissimi insecta odore allicientes. Invol. albida. (v. s.)

220. *E. PIQUERIOIDES*, caule herbaceo tereti glabro, ramulis oppositis apice petiolatisque puberulis, foliis oppositis petiolatis lanceolatis acuminatis trinerviis remotè serratis secus nervos subtus puberulis cæterum glabris, paniculis laxis oppositifloris ramos terminantibus, invol. 5-phyllis, floribus circ. 6-7, styli ramis exsertis apice crassè clavellatis. — in Peruviae montibus Guanacensibus legit cl. Haenke. Simillimum Piquerie trinervie sed folia subtus secus nervos hirtella et car. gener. diversi. (v. s. in h. Haenke ab ill. de Sternberg comm.)

\*\*\*\* *Capitulis 8-20-floris. — Herba Boreali-Americæ, foliis petiolatis oppositis aut in unica specie alternis.*

221. *E. CORDATUM* (Walt. car. 199), caule tereti tenui-pubescente, foliis oppositis breviter petiolatis ovatis subcordatis grossè et obtusè dentatis triplinerviis suprà sparè et subtus secus nervos scabridis, corymbo composito, capitulis 12-20-floris, invol. squamis circ. 10 lanceolatis subæqualibus vix pubescentibus, achænio subangulato glabro.  $\frac{1}{2}$  in siccis Carolinæ (Michx. ! Bosc !), Novæ-Cæsareæ (Green. !). *Eup. aromaticum* Ell. sketch 2. p. 304. non Lin. *E. cordiforme* Poir. suppl. 2. p. 600. *E. melissoides* Willd. spec. 3. p. 1754? sed capit. dicuntur 5-6-flor. Cor. albæ odoratæ. Petioli 2 lin. longi. (v. s.)

*Æ. Fraseri*, foliis ovatis paulò longioribus nec basi subcordatis.  $\frac{1}{2}$  in Carolinæ. *E. Fraseri* Poir. suppl. 2. p. 600. Lam. ill. t. 672. f. 4. (v. s. ex herb. Fraser ex quo etiam Lam. et Poir. habuerunt.)

222. *E. AROMATICUM* (Lin. sp. 1175), caule tereti pubescente, foliis oppositis petiolatis ovatis acuminatis trinerviis grossè et inæqualiter serratis scabriusculis, corymbo amplo subpaniculato, capitulis circ. 20-floris, invol. squamis 10-12 linearibus acutis æqualibus, achænio angulato glabro.  $\frac{1}{2}$  in Virginia (Lin.), Novâ-Cæsareâ (Torr. !), à Pensylviâ ad Floridam (Pursh) Willd. sp. 3. p. 1765. — Pluk. alm. t. 88. f. 3. Cor. albæ. Styli breviter exserti. (v. s.)

223. *E. CRANOTRIFOLIUM* (Muhl. in Willd. sp. 3. p. 1755), caule tereti glabresculo, foliis oppositis petiolatis ovatis acuminatis dentatis triplinerviis subglabris, corymbo subconferto, capitulis 5-10-floris, invol. squamis 10 linearibus subæqualibus, achænio angulato glabro.  $\frac{1}{2}$  in pinguibus Carolinæ ad Paris island prope Beanfort (Ell. sketch 2. p. 303), à Noveboraco ad Virginiam in rupibus (Pursh fl. 2. p. 514). An stirps Purshiana eadem ac Elliotiana? (v. s.)

224. *E. AGERATOIDES* (Lin. f. suppl. 355), caule tereti glabro, foliis oppositis longè petiolatis latè ovatis acuminatis trinerviis inæqualiter et grossè serratis glabris, corymbo composito, capitulis circ. 20-floris, invol. squamis linearibus acutis æqualibus, achænio angulato glabro.  $\frac{1}{2}$  in Amer. bor. Canadâ (Michx.) Virginiâ (Willd.) Carolinâ (Ell.) Pensylviâ et montibus Allegany (Pursh). *E. urticæfolium* Reich. syst. 3. p. 719. Michx. fl. bor. 2. p. 100. *E. altissimum* Lin. syst. veg. 614. *Ageratum altissimum*. Lin. sp. 1176. — Corn. can. t. 21. Mor. ox. s. 7. t. 18. f. 11. Cor. albæ. Styli longè exserti. Petiol. 1-2 poll. longi. Fol. interdum subcordata. (v. s.)

225. *E. SÆROTINUM* (Michx. fl. bor. am. 2. p. 100), caule tereti pulverulento-pubescente, foliis oppositis petiolatis ovato-lanceolatis supernè attenuatis acutis grossè et acutè serratis triplinerviis pubescentibus, corymbo composito, capitulis 10-12 floris, invol. squamis 10 linearibus puberis subimbricatis, achænio angulato.  $\frac{1}{2}$  in vallibus Carolinæ maritimæ (Michx.). Ell. sketch 2. p. 304. Folia non rariter serrata ut vult Michaux, sed serratoris grossis plurimis utrinquè notata. Cor. albæ. (v. s. c. ex sem. Michaux in h. L'Hér.)

226. *E. INCARNATUM* (Walt. flor. car. 200. Ell. ! sketch 2. p. 305), caule tereti pubescente, foliis oppositis longè petiolatis ovatis basi truncatis subcordatisve

acuminatis grossè et obtusè dentatis triplinerviis glabriusculis, corymbi laxi ramis subracemosis, capitulis 20-floris, invol. squamis circ. 15-20 linearibus acutis biserialibus subæqualibus, achenio angulato glabro.  $\frac{1}{2}$  in pinguibus Carolinæ. Cor. purpureæ. Petioli 1-2 poll. longi. Affin. E. ocellato sed certè distinctum. An fortè E. violaceum Raf. fl. Ind. 62? (v. s.)

227. E. ASSIROSUM (Torr. l. in litt. 1835), caule tereti erecto brevissimè velutino, foliis oppositis sessilibus elongato-linearilanceolatis acuminatis serratis basi obtusis superne glabriusculis subtus velutino-tomentosis, corymbo composito fastigiato polycephalo, invol. squamis ovalibus obtusis dorso velutino-villosis, flor. 10-12, acheniis resinoso-muricatis.  $\frac{1}{2}$  in Novâ-Cæsariâ (Torr. l. Guss.). Habitus potius sect. sequentis sed ex numero florum huc referendus. Flores albi. Styli modicè exserti. (v. s.)

228. E. OCCIDENTALE (Hook. fl. bor. am. 1. p. 305), glabriusculum, caule striato, foliis alternis brevè petiolatis ovatis acuminatis grossè inequaliter serratis apice integerrimis, summis lanceolatis integerrimis, capitulis paniculato-corymbois, invol. pluriflori squamis 1-serialibus acutis laxis, acheniis linearibus angulatis. — in collibus lapideis inter flum. Lewis et Clark Amer. bor. occid. Flores albi.

♀. *subrosum*, caule tereti, corollis apice stylisque roseo-purpureascentibus. — ad flum. Columbia (Douglas). Flor. in invol. 15-20. (v. s. comm. ab hoc Soc. hort. Londin.)

\*\*\*\*\* Capitulis 5-floris. — *Herbæ Boreali-Americanae, foliis indivisis pinnatifidum alternis aut oppositis, rarissimè verticillatis.*

229. E. LEPTOPHYLLUM, caule glaberrimo tereti erecto apice paniculato-ramoso, foliis alternis filiformibus tenuibus glabris integerrimis, racemis spicatis elongatis secundis ramos ramulosque terminantibus in paniculam dispositis, capitulis 3-5-floris breviter pedicellatis, invol. squamis lanceolatis margine albo angusto cinctis.  $\frac{1}{2}$  in Georgia circa Savannah à cl. Herbmont lectam habi. Habitus inflorescentiæ Solidaginum, cæterum E. feniculaceo affine. (v. s. comm. à cl. Duby.)

230. E. FENICULACEUM (Willd. sp. 3. p. 1750), caule striato pubescente paniculato ramosissimo, foliis alternis glabris, infer. pinnatifidis, super. fasciculatis, omnibus linearifiliformibus, capitulis innumeris parvis 3-5-floris, invol. 10-phylli squamis ext. 5 brevioribus pubescentibus, achenio cylindrico glabro.  $\frac{1}{2}$  in pascuis pinguibus præ. maritimis à Virginia ad Floridam et Georgiam. Pursh fl. 2. p. 512. Ell. sketch 2 p. 294. E. feniculoides Walt. car. 199. Chrysocoma capillacea Michx. fl. bor. am. 2. p. 101. Cor. ochroleuca interdum purpureo tinctæ (Ell.). Habitus Artemisiæ ex Pursh. (v. s.)

♀. *traganthes*, panicula stricta angustè ramulis rachi approximatis, invol. 6-phyllis 6-floris. — in Amer. bor. (ex L'Hér. l. mss. qui asserit plantam in horto Par. natam in terrâ in qua Dionæa muscipula adfecta fuerat) nec in Chinâ. Artemisia capillifolia Lam. l. dict. 1. p. 266. Art. tenuifolia Willd. sp. 3. p. 1819. non Moench nec Adans. Traganthes Wallr. sched. crit. 456. Mikania artemisioides Cass. dict. 55. p. 130. — Dill. elrh. 37. t. 33. Specimen habeo ex cl. Jan lectum Venetiis. An spontè? an in horto lectam? (v. s.)

231. E. CORONOPOLIUM (Willd. sp. 3. p. 1750), caule tereti pubescente diffusopaniculato, foliis subpubescentibus alternis, infer. pinnatifidis lobis linearibus, reliquis fasciculatis angustè linearibus integerrimis, capitulis 5-floris, invol. 10-phyllis pubescentibus basi imbricato, achenio angulato glabro.  $\frac{1}{2}$  in siccis sterilibus Carolinæ. E. compositifolium Walt. car. 199. Chrysocoma coronopifolia Michx. fl. am. 2. p. 102. Cor. albæ. Affine E. feniculaceo. (v. s.)

232. E. PINNATIFIDUM (Ell. sketch 2. p. 295), caule striato pubescente apice corymboso-ramoso, foliis pubescentibus pinnatifidis, infer. 4<sup>te</sup> verticillatis, super. alternis, capitulis 5-floris, invol. squamis 8-10 oblongo-lanceolatis pubescentibus, achenio oblongo sulcato.  $\frac{1}{2}$  in humidis Carolinæ mediæ. Cor. albæ. Species media inter. E. feniculaceum coronopifoliumve et reliquas generis stirpes.

233. *E. LINZABIFOLIUM* (Walt. car. 199), caule tereti pubescente apice ramoso-corymboso, foliis inferr. oppositis, sup. alternis, omnibus pubescentibus linearibus aut sublancoelatis 1-nerviis integerrimis aut hinc inde subdentatis, capitulis 5-floris, invol. squamis 10 oblongo-linearibus imbricatis dorso puberis et glandulosis, achenio sulcato-angulato glabro glanduloso.  $\frac{1}{2}$  in Carolinâ frequens (Fras.!), circa Charleston (Nois.!), à Novâ-Cæsareâ (Greene), ad Carolinam (Pursh fl. a. p. 513), in Novo-Eboraco (Torr.). Michx. fl. 2. p. 97. Willd. sp. 3. p. 1750. Pluk. alm. t. 88. f. 2? — An *E. hyssopifolium* Ell. sketch 2. p. 297? *E. crassifolium* Rafin. fl. lud. 62? — Styli plerumque corollâ longiores. Flores albi. (v. s.)
234. *E. HYSSOPIFOLIUM* (Lin. sp. 1171), caule tereti pubescente apice laxè corymboso-ramoso, foliis oppositis summiusve alternis lineari-lanceolatis trinerviis pubescentibus punctatisque integerrimis aut inferioribus dentatis, capitulis 5-floris, invol. squamis 10 imbricatis dorso puberis glandulosisque.  $\frac{1}{2}$  in pinetis et arenosis à Carolinâ (Fras.! Bosc!), ad Novam-Cæsaream (Pursh fl. bor. am. 2. p. 512). — Dill. elth. 141. t. 115. f. 140. Flores albi. An *E. linearifolium* Ell. sketch 2. p. 296 huc referendum? sed caulis dicitur subprocumbens, atque styli corollam subæquantes! unde suspicor speciem Elliottii diversam et fortè novam sed mihi ignotam. (v. s.)
235. *E. GLAUCESCENS* (Ell. sketch 2. f. 297), caule pubescente, foliis plerisque oppositis subsessilibus lanceolatis obtusè pauci-serratis basi integerrimis triplinerviis subglancis pubescentibus, capitulis corymbosis 5-floris, invol. squamis 8-10 lanceolatis acutis dorso glandulosis, stylo corollâ longiore  $\frac{1}{2}$ . In pinibus umbrosis Carolinæ. Cor. albæ. Serraturæ 3-4 utriusque à medio ad apicem folii.
- fl. leucolepis*. capitulis 5-7-floris, invol. squamis lanceolatis acuminatis apice albedo-scariosis.  $\frac{1}{2}$  in Novâ-Cæsareâ (Torr.! Greene!) Eup. linearifolium Michx? ex Torr. in litt. sed ab *E. linearifolio* differt foliis evidenter à medio ad apicem serratis, invol. acuminatis nec subacutis. (v. s.)
236. *E. ALTIMUM* (Lin. sp. 1171), caule tereti pubescente apice ramoso-corymboso, foliis oppositis subsessilibus lanceolatis trinerviis utrinque attenuatis pubescentibus, inf. medio aut à medio ad apicem serratis, sup. integerrimis, capitulis 5-floris, invol. squamis 10 oblongo-linearibus imbricatis obtusiusculis pubescentibus, achenio glabro glanduloso teretiusculo.  $\frac{1}{2}$  in arenosis sylvaticis demissis Pennsylvaniae, Virginiae et ad ripas fl. Mississipi, Missouri. Pursh fl. am. bor. 2. p. 514. Jacq. hort. Vind. t. 164. Willd. sp. 3. p. 1754. Fl. albi. Caulis 3-4-pedalis. (v. s. sp. et v. c.)
237. *E. SEMISERRATUM*, velutino-pubescent subvillosum, caule tereti, foliis 3<sup>is</sup> verticillatis, rameis summiusve oppositis lanceolatis utrinque longè acuminatis triplinerviis infernè integerrimis subpetiolatis, supernè serraturis obtusis 9-10-profundè notatis, corymbo composito fastigiato, capitulis 5-floris, invol. squamis 10, int. linearibus obtusis, ext. brevissimis, achenio angulato glabro.  $\frac{1}{2}$  in Georgia circa Savannah legit cl. Herbesmont. Cor. albæ parvæ. Affine *E. canefolio*. (v. s. comm. à cl. DuRoi.)
238. *E. CUMIFOLIUM* (Willd. sp. 3. p. 1753), caule tereti pubescente, foliis oppositis alternisve angusto-lanceolatis basi cuneato-atenuatis subpetiolatis et integerrimis ceteram dentatis triplinerviis utrinque velutino-pubescentibus, corymbo composito, capitulis 5-floris, invol. squamis 10 linearibus obtusis, ext. brevissimis, achenio angulato parè glanduloso.  $\frac{1}{2}$  in Carolinâ (Bosc!) ad St. Thomas (Ell.). *E. parviflorum* Ell. sketch 2. p. 299. non Sw. Cor. albæ parvum involucre longiores. *E. Marrubium* Walt. huc à Willd. revocatum videtur omninò diversum. (v. s. comm. à cl. Bosc.)
239. *E. SCABRUM* (Ell. sketch 2. p. 299), caule pubescente, foliis oppositis sessilibus ovato-lanceolatis acutè serratis basi integris pubescentibus subscabris subtus subglancis, capitulis corymbosis 5-floris, invol. squamis 10 acutissimis puberis glandulosis, achenio angulato.  $\frac{1}{2}$  in siccis Carolinæ. Cor. albæ invol. longiores. Rami inf. brachiati, sup. alterni.

240. *E. PUBESCENS* (Muhl. in Willd. sp. 3. p. 1753), caule tereti pubescente, foliis oppositis sessilibus ovatis acuminatis utrinque sparse pubescentibus et punctato-glandulosis, infer. duplicato-serratis, sup. anbeserratis, corymbo composito fastigiato, capitulis 5-floris, invol. squamis 10 linear-lanceolatis acutis puberis, achenio angulato.  $\frac{1}{2}$  in sylvis arenosis Pennsylvaniae (Muhl. Ell.) à Novâ-Caesarâ ad Carolinam? ex Pursh fl. 514. excl. syn. Michx. Pers. ench. 2. p. 400. excl. syn. Michx. Fl. albi. An *Critonia elliptica* Raf. in Kentucky orientali lecta ab hoc diversa? (v. s.)
241. *E. AMERUM* (Pursh fl. bor. am. 2. p. 514), caule solido levi, foliis oppositis ternisque brevè petiolatis lanceolato-oblongis utrinque acutis serratis glabrisculis subrugosis subtus reticulato-venosis, panicula corymboso-fasciculata confertè pedunculis tomentosis, capitulis 5-floris, invol. squamis oblongis acutis coloratis.  $\frac{1}{2}$  in montanis Novâ-Caesaræ. Fl. carol.
242. *E. FALCATUM* (Michx. fl. bor. am. 2. p. 99), glabellum, foliis 4<sup>to</sup> verticillatis ovali-lanceolatis utrinque acuminatis subfalcatis rarius serratis, panicula brachiata multicorymbosa, capitulis 5-floris, invol. gradatim imbricatis, squamis extimis brevibus ovalibus.  $\frac{1}{2}$  in Amer. bor. ad ripas fl. Ohio et Scioto. Affine dicitur *E. albo*.
243. *E. ALBUM* (Linn. mant. 111), caule tereti pubero apice ramoso-corymboso, foliis oppositis subsessilibus lato-lanceolatis basi attenuatis apice grossè pinnidentatis penninerviis utrinque scabrinusculis subtus punctatis, capitulis 5-floris, invol. squamis 10 elongato-lanceolatis acuminatis corollas ferè superantibus dorso glandulosis, achenio angulato glabro.  $\frac{1}{2}$  in Pennsylvania (Linn.), Novâ-Caesarâ (Torr.), Georgiâ prope Savannah (herb. Duby!), Willd. sp. 3. p. 1752 excl. syn. Kämpf. Icon Banks. Kämpf. t. 26. ad *E. Finlaysonianum* relata, hanc speciem refert sed in pl. Japonica corollæ involucro longior et circ. aequalis in Americâ. Cor. albæ. Invol. squamæ apice albide. (v. s.)
- $\beta$ . *glandulosum*, caule magis hispidulo, foliis basi minis attenuatis per totam ferè longitudinem dentatis subtus involucrisque multò magis punctato-glandulosis.  $\frac{1}{2}$  in aridis sylvarum Carolinæ. *E. glandulosum* Michx. fl. bor. am. 2. p. 98. *E. album* Ell. sketch 2. p. 298. (v. s.)
244. *E. TRICAPITOLITUM* (Willd. sp. 3. p. 1753. hort. ber. t. 32), caule tereti pubescente, foliis oppositis sessilibus distinctis ovatis subacuminatis scabris dentato-serratis basi profundius dentatis, corymbo composito subpaniculato, capitulis 5-6-floris, invol. squamis dorso puberis apice acutis, achenio angulato glanduloso.  $\frac{1}{2}$  in Amer. bor. à Novâ-Angliâ ad Carolinam. Fl. albi. (v. s.)
- $\beta$ . *verbenæfolium*, foliis oblongo-lanceolatis profundius incis.  $\frac{1}{2}$  à Bostoniâ ad Carolinam. Eup. pilosum Walt. car. 199. Eup. *verbenæfolium* Michx. fl. bor. am. 2. p. 98. An fortè *E. lanceolatum* Willd. sp. 3. p. 1752? (v. s.)
245. *E. ROTUNDIFOLIUM* (Linn. sp. 1173), caule tereti pubescenti-velutino, foliis oppositis sessilibus distinctis subrotundo- aut ovato-cordatis obtusis dentatis venosis utrinque pubescentibus subtus punctato-glandulosis, corymbo fastigiato, capitulis 5-floris, invol. squamis 10 acuminatis, achenio angulato glanduloso.  $\frac{1}{2}$  in pinetis et sylvis à Canadâ ad Carolinam (Pursh), in Novâ-Caesarâ (Raf.), Georgiâ (Herbm.), Pursh fl. 2. p. 514. Ell. sketch 2. p. 300. — Plukn. alm. t. 88. f. 4. — *E. Marrabium* Walt. car. 199? ex Ell. — Flor. albi. Herbar. decoctio tonica febrifuga. (v. s.)
- $\beta$ ? *ovatum* (Torr. in litt. 1835), capitulis 7-8-floris, caule magis hirsuto.  $\frac{1}{2}$  in demissis circa Bostoniam. Eup. *ovatum* Bigel. ex Torr. an spec. propria? (v. s.)

\*\*\*\*\* Capitulis 5-8-floris. — Species *Gerontogæ*, foliis indivisis.

246. *E. LONGICAULE*, caule herbaceo erecto tereti ramoso hirsuto-velutino, foliis isolatis ovato-lanceolatis acuminatis serratis basi triplinerviis superè glabris

subtus pubescenti-velutinis, corymbo composito fastigiato amplo, capitulis 8-floris, invol. squamis circ. 10 oblongis acutis glabriscentis.  $\mathcal{L}$  in Nepaliâ legit cl. Wallich, etiam recepi à cl. Royle. *Conyza longicaulis* Wall.! cat. n. 3073. comp. 183 non n. 2169 ad *Vernoniam longicaulem* referend. *Mikania*? *longicaulis* Wall.! herb. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

247. *E. WALLICHII*, caule erecto herbaceo ramoso tereti striato pubescenti-velutino, foliis vix petiolatis ovali-lanceolatis acuminatis serratis suprà glabriusculis subtus pubescentibus, capitulis in corymbum subglobosum densum terminalem congestis subsessilibus paucifloris, invol. squamis paucis obtusis, ext. brevioribus subacutis.  $\mathcal{L}$  in Nepaliâ legit cl. Wallich. *E. viscosum* Wall.! cat. n. 3287. comp. 397 non Kunth. Ex statu florum juniore nec numerum nec formam ritè vidi sed ex habitu sine dubio huc refero. (v. s. comm. ab. hon. Cur. merc. angl. Ind. or.)

248. *E. BIRMANICUM*, caule suffruticoso erecto ramoso apice subpuberulo, foliis breviter petiolatis lanceolatis utrinquè acuminatis parcè serratis glabriusculis, corymbo oligocephalo, capitulis parvis paucifloris, pedicellis canescenti-villosis, invol. glabro.  $\mathcal{L}$  in Indiâ or. regione Birmanorum ad Segain legit cl. Wallich. *E. suaveolens* Wall.! cat. n. 3290. comp. n. 400 non Kunth. Folia ferè *E. triplinervii* sed *penninervia*. Capitula in specim. obvio nimis juniora! (v. s. in h. Cur. Ind. or. nunc Soc. Lin. Lond.)

249. *E. FIRLAYSONIANUM* (Wall.! cat. n. 7133), caule herbaceo ramoso erecto? striato tereti apice puberulo, foliis lanceolatis in petiolum attenuatis acuminatis grossè et irregulariter inciso-serratis glabris, corymbo composito fastigiato, pedunculis subbracteolatis, capitulis 5-floris, invol. squamis circ. 10 inæqualibus oblongo-linearibus glabris, ext. acuminatis.  $\mathcal{L}$  in Indiâ or.? (ex herb. Finl.) et Japoniâ si ut autumo *E. album* Thunb. fl. jap. 308. Banks in Kämpf. t. 26 non Linn. huc pertinet. (v. s. comm. ab ill. Cur. merc. angl. Ind. or.)

250. *E. PUNDUANUM* (Wall.! cat. n. 3170. comp. n. 280), caule herbaceo erecto ramoso tereti pubescenti-velutino, foliis vix petiolatis lanceolatis acuminatis à medio acutè serratis utrinquè puberis subtrinerviis, ramis in paniculam subdispositis nudiusculis apice confertè corymbois, capitulis 4-floris, invol. squamis glabris 5-7 oblongo-linearibus obtusis.  $\mathcal{L}$  ad Pundua Ind. or. legit cl. Wallich. *Achnia subangulata* glabra basi attenuata subpuberula. Pappus albidus cor. aequalis. Styli rami exserti longissimi! (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

$\mu$ . *Lambertianum* (Wall.! cat. n. 3285. comp. 280), caule paulò magis hirsuto. — cum priori. (v. s. comm. cum  $\alpha$ .)

251. *E. REEVESII* (Wall.! cat. n. 3168. comp. 278), caule herbaceo erecto tereti glabriusculo, foliis subsessilibus ovatis acutis grossè et obtusè dentatis glabris nervis subtus et interdum suprà puberulis, pedunculis axill. nudis apice congestè corymbosis, supremis in corymbum compositum confertis, capitulis subsessilibus 5-floris, invol. squamis circ. 10 inæqualibus glabris oblongo-linearibus obtusis.  $\mathcal{L}$  in Chinâ legit cl. Reeves. Fl. ex sicco vid. albi. Pappus albidus. *Achnia* glabra striato-angulata. (v. s. in h. Calc. comm. ab ill. Cur. Ind. or. et in h. Lindl. à cl. Parkes circ. Canton lect.)

252. *E. MODIFLORUM* (Wall.! cat. n. 3166. comp. 276), caule herbaceo erecto subsimplici tereti hirsuto, foliis sessilibus oblongo-lanceolatis mmerone acutis grossè serratis sparè puberis subtus in nervis hispidis, corymbo composito conferto, capitulis ad apices ramorum congestis 4-floris, invol. squamis 5-6 inæqualibus dorso pubescentibus.  $\mathcal{L}$  in Indiâ or. montibus Silhetensibus. Strictum 1-2-pedale. Folia vere *penninervia* sed nervis infimis 2 validioribus. *Achnia* striato-angulata glabra glandulosa. Pappus sordidè albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

253. *E. CHINENSE* (Linn. sp. 1172), herbaceum, caule tereti striatulo tenuè

puberulo oppositè ramoso, foliis oppositis breviter petiolatis ovato-lanceolatis acutis serratis utrinque tenuissimè puberulis, corymbis terminalibus compositis subsagittatis, capitulis pedicellatis 5-floris, invol. squamis 5-7 oblongo-linearibus acutis dorso puberulis, achenio teretiusculo striato glabro.  $\frac{1}{2}$  in Chinâ (Linn. h. Lindl.!), Japoniâ? (Thunb. fl. jap. 308). An stirps Japonica de quâ Thunb. caulem et folia glabra et Willdenowiana (sp. pl. 1753) cui auctor folia obovata glabra tribuit, sunt omninò similes? (v. s. in h. Lindl.)

254. E. LINDLEYANUM, caule herbaceo erecto simplici tereti striato puberulo apice nudiusculo, foliis oppositis subsessilibus lineari-lanceolatis remotè serratis glabrescentibus basi trinerviis, corymbo composito terminali, pedicellis sub capitulo minutissimè bracteatis, invol. squamis 5 oblongo-lanceolatis acutis, floribus 5, acheniis angulatis minutè glandulosis.  $\frac{1}{2}$  in Chinâ collibus. Radices è collo fasciculatè teretes. Folia sesquipoll. longa, 2-3 lin. lata. (v. s. in h. Lindley.)

\*\*\*\*\* *Cannabina*. — Capitula 5-6-flora. Invol. squamæ 7-9 circ. biseriales oblongo-lineares non verè imbricatæ. — Folia opposita aut verticillata, indivisa aut trifida. Caules herbacei.

255. E. CANNABINUM (Linn. sp. 1173), caule herbaceo erecto striato scabro, foliis plerisque trisectis, segmentis lanceolatis acuminatis basi attenuatis regulariter serratis pubescentibus, corymbo fastigiato, capitulis 5-6-floris, invol. cylindrico glabro circ. 10-phyll. squamis ext. 5 obtusis brevissimis.  $\frac{1}{2}$  in fœmie, aquosis, inundatis totius Europæ, Persiæ (Sow.!), Tauriæ (Nieb.). OED. fl. dan. t. 745. Sow. engl. bot. t. 478. E. trifoliatum Hb. non Linn. Flores pallide purpurascetes et var. albi, odore melleo insignes. Achenia subpentagona glandulosa. (v. v.)

$\frac{1}{2}$ . *indivisum* (DC. fl. fr. n. 3107), foliis omnibus aut plerisque indivisis. Ray. syn. 180. E. Soleirolii Lois. nouv. not. 36. fl. gall. ed. 2. v. 2. p. 223 huc relat. ex Petit ann. sc. obs. 1829. p. 463 et ex specim. Solms. (v. s.)

256. E. SYRIACUM (Jacq. ic. rar. 1. t. 170. misc. 2. p. 349), caule herbaceo erecto subpubescente, foliis plerisque trisectis, segmentis ovato-lanceolatis acuminatis basi breviter attenuatis subpubescentibus grossè et inæqualiter dentatis, corymbo fastigiato, capitulis 5-6 floris, invol. cylindrico glabro circ. 10-phyll. squamis ext. 5 obtusis brevissimis.  $\frac{1}{2}$  in Syriâ (Jacq.) et Persiâ (Sow.) Willd. sp. 3. p. 1757. Fortè vera E. cannabini varietas? (v. v. c.)

$\frac{1}{2}$ . *indivisum*, foliis plerisque indivisis.  $\frac{1}{2}$  in Persiâ boreali. (Stev. l. Sow.!) (v. s.)

257. E. HETEROPHYLLUM, caule herbaceo tereti striato puberulo, ramis gracilibus, foliis oppositis petiolatis suprâ sparè subtus æcne nervos puberulis, caulinis trisectis, segmentis ovali-lanceolatis utrinque acuminatis medio grossè serratis, ramis indivisis ovali-lanceolatis subintegerrimis, corymbis terminalibus confertis, capitulis pedicellatis 5-floris, invol. glabro 7-8-phyll. squamis 3-4 ext. inæqualibus brevibus, 4 oblongis obtusis subæqualibus, achenio glabro. — Patr. ign. Speciem distinctissimam vidi in tepidario univ. Vindob. anno 1827 à cl. et amic Jacquin commun. Stylus basi filiformis. Cor. basi inflato-globosa teretiuscula intus glabra. (v. s.)

258. E. JAPONICUM (Thunb. fl. jap. 308), caule herbaceo erecto striato scabro, foliis longiusculè petiolatis, inf. trisectis, segmentis foliisque super. ovatis acutis sæpius serratis subtus pallidè scabridis, paniculis terminalibus ovatis. pedunculis brevissimis, capitulis 5-floris.  $\frac{1}{2}$  in Japoniâ. An ex Willd. ab E. cannabino (seu syriaco) satis differt? Petioli dicuntur unguicularum nec 2-4 lin. longi. †

\*\*\*\*\* Capitula 4-5-floris. — Species ex Americâ calidiore foliis indivisis.

259. E. XITIDUM, fruticosum glaberrimum, caule tereti, foliis oppositis bre-

- viximè petiolatis approximatis ellipticis basi et apice attenuatis integerrimis triplinerviis utrinque lucidis, corymbis compositi fastigiati ramis inf. oppositis, summis alternis compressis, capitulis pedicellatis 5-floris, invol. squamis 1-2-seriatis subaequalibus mucronato-acuminatis, achenio puberulo.  $\frac{3}{4}$  in sabulosis aridis circa Bahiam Brasiliæ leg. cl. Salzmänn et Blanchet. Frutex 5-7-ped. Folia 2 poll. longa, pollicem lata, petiolo bilineari. Cor. alba. Antheræ ferè cor. orem sequentes. (v. s.)
260. *E. BOGOTENSE*, fruticosum glabrum, ramis teretibus, foliis breviter petiolatis oblongo-lanceolatis utrinque subacutis serratis coriaceis penninerviis, paniculâ terminali cernuâ viscosâ, capitulis longè pedicellatis 5-floris, invol. squamis 7-8 biserialibus inæqualibus obtusiusculis, achenio angulato glabro.  $\frac{3}{4}$  circa Santa-Fe de Bogota. Species paniculâ cernuâ et quasi reflexâ (ex a. specim.) distinctissima. Petiol. a lin. Limb. fol. 2 poll. longus, 7 lin. latus. (v. s. comm. ab amic. Delessert.)
261. *E. ANRATIFOLIUM*, fruticosa erecta glabra, ramis sexangularibus, foliis petiolatis ovato-oblongis acuminatis basi cuneatis dentato-serratis reticulato-trinerviis, corymbis terminalibus ramosissimis, capitulis solitariis pedicellatis 5-floris, invol. squamis 5 oblongo-linearibus obtusis subellipticis viscosis, achenii angulis pilosisculis.  $\frac{3}{4}$  in Novâ Granatâ. Petiol. 3-4 lin. Limb. fol. 15-16 lin. longus. Mikania serratifolia H. B. et Kunth nov. gen. am. 4. p. 138.
262. *E. SINERIANUM*, herbaceum pubescens, caule tereti striatulo erecto, foliis petiolatis ovato-lanceolatis basi cuneatis apice acuminatis repando-dentatis suprâ sparse subpuberulis subtus in nervis petiolisque puberis, paniculâ laxâ subtrichotomâ, ramis apice capitula oblonga glabra 5-flora fasciculato-sessilia gerentibus, invol. squamis 6-7 subbiserialis oblongis mucronatis, achenio angulato puberulo. — in insulâ Trinitatis legit cl. Sieber. Mikania serratifolia Sieb. f. fl. trin. exa. n. 72 non Kunth. Petiolu 6 lin. longi. Limbus 5-6 poll. ferè longus, 12-15 lin. latus. Capitula pallida 2 lin. longa. (v. s.)
263. *E. ZEUSTANUM*, fruticosum, ramis teretibus glabris, ramulis subangulatis pulverulento-velutinis, foliis oppositis petiolatis oblongis aut subellipticis obtusiusculis basi attenuatis et subrecurvis ad apicem dentibus paucissimis notatis glabris subtus crebrè punctatis, corymbis trichotomis polycephalis, pedicellis glanduloso-puberulis ferrugineis, capitulis 4-5-floris, invol. squamis 5-6 linearibus obtusiusculis sub 1-seriatis, ext. panicis parvis acutis, achenio angulato subcabrido.  $\frac{3}{4}$  in Mexico ad Tamaulipas legit cl. Berlandier. Cor. albæ, tubo brevi, fauce obconicâ. Folia circ. Liguatri. (v. s.)
264. *E. TRIPIDUM* (Vahl symb. 3. p. 94), scandens, caule striato vix pubescente, foliis alternis petiolatis tripartitis, lacinii linearibus extus serratis, summis indivisis lanceolatis, corymbis longè pedunculatis compositis, capitulis 5-floris, invol. squamis glabris 10-phyllis biserialis, extimis brevissimis. — in Caribæis.
265. *E. GLUMACEUM*, caule herbaceo erecto tereti pubescenti-piloso, foliis oppositis petiolatis ovato-lanceolatis acuminatis hinc indè subserratis suprâ pubescenti velutinis subtus velutino-tomentosis trinerviis, paniculâ laxâ diffusâ trichotomâ, ramulis ad apices capitula plurima oblonga 4-flora breviter pedicellata gerentibus, invol. squamis stramineis glaberrimis acuminatis biserialis, ext. brevioribus. — in insulâ Cubâ legit cl. de la Ossa. Folia internodiis multò breviora, adjecto petiolo 4-lin., 3 poll. ferè longa, ferè pollicem lata. Invol. glumacea subnitentia. Achenium angulatum puberulum. (v. s.)
266. *E. CHILCA* (H. B. et Kunth nov. gen. am. 4. p. 126), fruticosum glabrum, ramis striatis, foliis oppositis petiolatis ovato-oblongis acuminatis serratis basi angustatis triplinerviis, paniculis terminalibus corymbosis foliatis, capitulis pedicellatis et sessilibus sub 4-floris, invol. squamis 4-6 subimbricatis oblongo-lanceolatis acutis externè glutinosas, achenio glabro  $\frac{3}{4}$  ad radices montis Sanctæ-Poloniz juxta Caxamarca Peruvianorum. Cor. alba. Petiol. lineam longus. Fol. limbus sesquipollicem longus, 4 lin. latus. Frutex aromaticus Chilca vern. dictus. An Mikaniæ spec.?



267. *E. AFFINE* (H. B. et Kunth nov. gen. am. 4. p. 126), fruticosa glabrum. ramis striatis, foliis oppositis petiolatis ovato-oblongis acuminatis serratis basi cuneatis triplinerviis, panicula terminalibus corymbosis, capitulis pedicellatis et sessilibus 3-6-floris, invol. squamis 4-6 subimbricatis oblongo-lanceolatis acutis externe glutinosi, achenio ad angulos scabriusculo. ♀ cum *E. Chilcã*? à quo ipsa docemur Kunthio vix differt, nisi foliis latioribus longius petiolatis, capitalisque majoribus.

\*\*\*\*\* *Multilobata*. — Capitula 5-flora. Invol. squamis 5 æquales subimbriciales. — Folia multifida, lobis linearibus. Caules fruticulosi.

268. *E. ANETHROFOLIUM*, suffruticulosum? ramosissimum adscendens glabrum, ramis teretibus junior. striatis, foliis oppositis petiolatis multipartitis, lobis oppositis linearibus integerrimis ultimis trifidis, corymbi terminalis ramis ramulisque hirtellis, bracteæ setaceo-umbellatis, capitulis pedicellatis 5-floris, invol. squamis 5 glabriusculis oblongo-linearibus subobtusis. ♂ ♀ in Brasiliâ prov. Rio-Grande. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 913 et 949 miss.)

269. *E. MULTIFIDUM*, fruticulosum ramosissimum adscendens, ramis teretibus junioribus setoso-hispidis et pube minimâ glandulosâ subvelutinis, adultis glabris, foliis oppositis subsessilibus multipartitis, lobis oppositis apice trifidis linearibus oblongis integerrimis glabris rarius dorso subsetosis, corymbo composito pedunculato conferto, ramis alternis brevibus, capitulis subsessilibus circ. 5-6-floris, invol. biseriali, ext. foliolis setaceis, inter. squamis 5-6 oblongis obtusis apice ciliatis. ♀ in Brasiliâ prov. Sancti-Pauli. Species distinctissima vix pedalis aut aequipedalis. Folia 6-8 lin. diam. cum invol. subviscoso. Flor. ex axiis albi. Achenium glabrum angulatum. Pappus corollâ dimid. fere brevior. Habitus à genere alienus! (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 508 et 509 miss.)

270. *E. BRASILIENSE* (Spreng. syst. 3. p. 417), caule herbaceo simplicimulo, foliis subsessilibus confertis 3-5-partitis linearifiliformibus glabris, corymbo lateralibus, capitulis sub-3-floris, invol. glabro subimbricato aut simpliciori. — in Brasiliâ. Affine fortè *E. multifido*. Cæter. ign. †.

### † Species non satis notæ.

271. *E. AMYGDALINUM* (Lam. dict. 2. p. 408), caule... tereti simplici glabro, foliis oppositis sessilibus oblongo-lanceolatis subdentatis glabris, paniculâ laxâ subnudâ, capitulis 12-15-floris, invol. ovalis squamis oblongis inæqualibus margine subvillosis. — in Peruvîâ ex h. Juss. Folia inf. 3 poll. longa, summa subalterna.

272. *E. AUSTRALIS* (Thunb. in fl. 1821. p. 332. n. 20), fruticosum, ramis..... foliis..... triplinerviis ellipticis serratis, paniculâ fastigiatâ, capitulis..... ♀ in Brasiliâ. †.

273. *E. BRICKELLIA*, caule herbaceo tenuissimè pubescente apice subtomentoso, foliis oppositis cordatis acuminatis dentatis triplinerviis tenuè pubescentibus, summis subalternis basi obtusis, corymbo paniculato, invol. squamis ext. setaceis, int. linearilanceolatis, achenio subglabro ro-striato. — ad latera collium in Georgiâ occidentali. Brickellia cordifolia Ell. sketch 2. p. 290. Jam suadente cl. Lessing hæc stirpem ab Eupatorio non separare queo, sed ex numero florum potius ad Eupatorii species Mexicanas quàm boreali-Americanas accedere videtur.

274. *E. CHILANÆ* (Molin. chil. ed. gall. p. 335), foliis oppositis amplexicaulis lanceolatis denticulatis, invol. 5-floris. — in Chili. Cæter. ign.

275. *E. COTINIFOLIUM* (Willd. phytogr. 11. t. 7. f. 1. ex sp. pl. 3. p. 1769), fruticosum, foliis alternis petiolatis obovatis mucronatis integerrimis trinerviis, corymbo terminali. ♀ in Martiniâ. Capitula multiflora, involucri nudi rictu ex Spreng. †.

276. *E?* *FARTASTII* (Thunb. in Flor. 1821. p. 332. n. 19), fruticosum scandens, ramis.... foliis.... lanceolatis triplinerviis glabris serratis, paniculis patentibus, capitulis....  $\zeta$  in Brasiliâ.  $\dagger$ .
277. *E. HISPIDUM* (Pers. ench. 2. p. 402. n. 66), caule suffruticoso biapido erecto, foliis alternis ovatis sinuatis denticulatis pilosis, paniculis terminalibus capitulis sub-8-floris, invol. oblongo imbricato.  $\zeta$  in insulâ Afric. Mozambico. *E. sinuatum* Loar. fl. coch. 2. p. 595. non Lam. Flor. purpurascens. Styli longi semibifidi. Recept. concavum. An. Vernoniæ potius species?  $\dagger$ .
278. *E?* *LUDOVICIANUM*. — in Louisianâ. *Cacalia* n. 1. Rob. louis. 432. *Cac. pubescens* Raf. fl. lud. 61.
279. *E. MEGAPOTAMICUM* (Spreng. syst. 3. p. 420), herbaceum, foliis alternis petiolatis ovatis mucronato-dentatis ramisque villosis, paniculâ terminali, capitulis paucifloris, invol. glabro subimbricato aut subsimplici. — in Brasiliâ prov. Rio-Grande. Cæc. ign.  $\dagger$ .
280. *E. MENTHAFOLIUM* (Poepp. ex Spreng. syst. 3. p. 412), caule fruticoso, foliis alternis petiolatis oblongis utrinque attenuatis medio serratis pubescentibus, paniculis axillaribus terminalibusque patentibus, capitulo multifloro, invol. imbricato.  $\zeta$  in Cubâ. Cæc. ign.
281. *E. MONTEVIDEENSE* (Spreng. syst. 3. p. 417), caule herbaceo ramoso, foliis sparsis subsessilibus linearibus margine subserrulato-scrabis, capitulis racemosis paucifloris, pappo rufo. — in Montevideo. Cæc. ign.  $\dagger$ .
282. *E. MYOSOTIFOLIUM* (Jacq. ic. rar. 3. t. 282. coll. 2. p. 341), herbaceum villosulum, caule tereti striato, foliis alternis in petiolum attenuatis obovato-cuneatis obtusis integerrimis, paniculâ diffusâ laxâ, capitulis longè pedicellatis cylindricis multifloris, invol. squamis acutis erectis.  $\textcircled{2}$  in Amer. calid. Cor. purpurea. Cæc. ign.  $\dagger$ .
283. *E. PRARINIANUM* (Spreng. syst. 3. p. 412), fruticosum, foliis alternis petiolatis oblongis utrinque attenuatis obtusiusculis repandis suprâ opacis glabris subtus fusco-tomentosis, paniculâ terminali foliosâ, capitulis multifloris, invol. imbricato oblongo-cylindraceo.  $\zeta$  in Amer. austr. Cæc. ign.  $\dagger$ .
284. *E. POMPEII* (Spreng. syst. 3. p. 415), fruticosum, ramis...., foliis oppositis petiolatis ovato-lanceolatis subintegerrimis subtus subtomentosis, paniculâ corymbosâ terminali, capitulis paucifloris, invol. pauciseriato. — in insulâ Cubâ. Eup. deltoideum Poepp. exs. non Jacq. Cæc. ign. An fortè idem ac *E. cynanchifolium*, sed descr. Spreng. non sufficit.
285. *E. QUATERNUM*, caule suffruticoso subsimplici striato, foliis subquaternis congestis subsessilibus ovato-lanceolatis subserratis rugoso-asperis, corymbo fastigiato, capitulis sub-8-floris, invol. oblongo-cylindrico.  $\zeta$  in agris Cochinchinæ. *E. purpureum* Lonr. fl. coch. 2. p. 595. non Linn. Suffrutex 3-pedalis. Flores purpurei. Recept. subnudum.  $\dagger$ .
286. *E. RETICULATUM* (Desv. in Ham. fl. ind. occ. p. 51. non Hook.) glaberrimum, ramis pentagonis, foliis oppositis obovato-oblongis retosis subundulatis consistentibus subtus reticulatis, paniculâ contractâ apice subumbellatâ, capitulis multifloris. — in Jamaicâ. Cæc. ign.
287. *E. ROSALESIÆ*, caule fruticoso, ramis virgatis apice involucriisque glanduloso-viscosis, foliis oppositis petiolatis ovato-subcordatis crenato-subseratis suprâ asperis subtus subtomentosis, capitulis corymbosis, pedunculis capitula 2-5 gerentibus, invol. imbricato, squamis apice revolutis.  $\zeta$  in Mexici pascuis ad S. José del Corral. *Rosalesia glandulosa* Lallav. et Lex. nov. veg. 1. p. 9. Frutex biorgyialis. Cor. pallide flava, limbo brevissimè 5-dentato. Styli exertè insigniter clavati. Achænia cylindrica striata villosa.
288. *E. SERRATUM* (Spreng. syst. 3. p. 415), fruticosum, ramis...., foliis oppositis subpetiolatis linearilanceolatis argutè serrulatis subtus tomentosis, paniculæ terminalis ramis divaricatis, capitulis paucifloris, invol. paucis seriebus imbricato aut subsimplici.  $\zeta$  in Monte-Video. Cæc. ign.

289. *E. SOPHILEFOLIUM* (Linn. sp. 1175), herbaceum, caule adscendente à basi multiplici, foliis oppositis bipinnatisectis, lobis sinuatis oblongis obtusis, corymbo paniculato, capitulis pedicellatis, invol. cylindrici squamis aretè imbricatis multiserialibus. — in insulis Caribæis. Eup. sophizefolio flore porpureo Plum. amer. ed Barm. t. 128. f. 2. †.
290. *E. SPICATUM* (Lam. dict. 2. p. 409), caule herbaceo? simplici striato subangulato, foliis oppositis lineari-spathulatis supernè rariter dentatis subtilissimè tomentosis, capitulis in spicam terminalem basi interruptam dispositis sessilibus subglomeratis multifloris, invol. ovoidei squamis oblongis margine scariosis. ♀ (Pers.) ♂ (ex Lam.) — in Monte-Video ex Comm. in h. Jan. Folia 2 poll. longa.
291. *E. SPRENGELIANUM*, fruticosum, ramis teretibus pubescentibus, junioribus albo-tomentosis, foliis alternis brevissimè petiolatis oblongis obtusis repandentatis coriaceis rugosis subtùs tomentosis, racemis terminalibus oligocephallis. ♀ in insulis Indiæ occid. E rugosum Spreng. neu entdt. 3. p. 39. non Kunth nec Schrad. Species valde dubia ab auct. ipso in systemate omissa. †.
292. *E. squamosum* (Don prod. fl. nep. 170), caule fruticoso diffuso, ramis teretibus, foliis ovatis acutis petiolatis crenatis 3-nerviis utrinquè tomentosis, corymbo paniculato, capitulis . . . , involucris oblongis pedicellisque scaris, squamis ext. ovatis acutis, intimis oblongis obtusis. ♀ in Nepaliâ. Flores albi.
293. *E. STACHYOPHYLLUM* (Spreng. syst. 3. p. 420), caule herbaceo ramoso hirsuto, foliis oppositis alternisque subsessilibus lanceolatis subcerratis rugosis 3-nerviis pubescentibus, corymbis terminalibus oligocephalis, capitulis pascifloris, invol. squamis obtusis glabris. — in Brasiliâ. Cst. ign.
294. *E. TOMENTELLUM* (Schrad. ind. sem. h. Gott. 1833. p. 3), frutescens, foliis cordatis acuminatis subtùs cano-pubescentibus integris, ramorum juniorum 3-5-lobis villosis, fol. radic. ligulis obovato-ellipticis. ♀ ♂ in Mexico.

*N. B.* Ad Eupatorii genus pertinere videntur stirpes floræ fluminensis vol. 8 sequentes, nempe *Chrysocoma maculata* t. 6, *Chrysocoma alata* t. 48 (affinis *E. quadrangulata* sed caules teretes), *Chrysocoma punctulata* t. 49, *Cacalia emarginata* t. 52, *Cacalia punctulata* t. 66, *Cacalia cor Jean* t. 71<sup>2</sup>, *Cacalia triphylla* t. 74, et *Cacalia lata* t. 76, sed descriptionibus deficientibus non satis nota.

### † Species exclusæ.

- E. ACUMINATUM* Wall. = *Vernonia acuminata*.  
*E. ACUMINATUM* Don. =? *Vernonia multiflora*.  
*E. ALTERNIFOLIUM* Ard. = *Kuhnia eupatorioides*.  
*E. AMARUM* Steud. = *Mikania amara*.  
*E. ARROLATUM* Pers. = *Ethulia conyzoides* var. *ℓ*.  
*E. ARMANI* Balb. = *Orsinia eupatorioides*.  
*E. ASPERUM* Roxb. = *Vernonia aspera*.  
*E. AURICULATUM* Lam. = *Senecio deltoideus*.  
*E. AXILLARE* fl. mex. = *Bulbostylis Cavanillesii*.  
*E. CAPENSE* Spr. = *Vernonia mespilifolia*.  
*E. BICOLOR* Lam. herb. Juss. = *Gynoxis buxifolia*.  
*E. CELEBICUM* Blum. = *Vernonia Celebica*.  
*E. CINERUM* L. f. = *Pteronia incana*.  
*E. COELESTINUM* L. = *Conoclinium coelestinum*.  
*E. COLOBATUM* Willd. = *Decaneurum Senegalense*.  
*E. CONDENSATUM* Reichenb. = *Adenocyclos condensatus*.  
*E. CORDATUM* Lam. non Walt. = *Mikania volubilis*.

- E. CORIACEUM* Vahl non Spr. = *Vernonia fimbrillifera*.  
*E. CORNIFOLIUM* Willd. = *Odontoloma acuminatum*.  
*E. CUBENSIS* Pers. non DC. = *Kuhnia rosmarinifolia*.  
*E. CUSPIDATUM* Willd. = *Dialesta bicolor*.  
*E. CYDONIÆFOLIUM* Willd. = *Vernonia dichocarpa*.  
*E. DALEA* Linn. = *Critonia Dalea*.  
*E. DALEA* H. B. K. = *Critonia pseudodalea*.  
*E. DELTIFOLIUM* Raf. = *Conoclinium celestinum*.  
*E. DENTICULATUM* St. = *Mikania denticulata*.  
*E. DIVARICATUM* Thunb. = *Pteronia divaricata*.  
*E. DISCOLOR* Spr. = *Vernonia discolor*.  
*E. ELEGANS* Vent. = *Liatris elegans*.  
*E. ELEAGNIFOLIUM* Hort. Calc. = *Vernonia eleagnifolia*.  
*E. FERRUGINEUM* Labill. = *Ozothamnus ferrugineus*.  
*E. FLEXUOSUM* Lam. = *Senecio flexuosus*.  
*E. GLABRUM* Heyn. = *Decaneurum glabrum*.  
*E. HOUTONII* Linn. = *Mikania Houstonia*.  
*E. JAVANICUM* Blum. = *Vernonia Javanica*.  
*E. KLEINHOEDII* H. B. K. = *Campuloclinium kleiniioides*.  
*E. LONGICAULE* Wall. n. 2109. = *Vernonia longicaulis*.  
*E. MACROCEPHALUM* Less. = *Campuloclinium macrocephalum*.  
*E. MACROPHYLLUM* Vahl. = *Hebeclinium macrophyllum*.  
*E. MELISSÆFOLIUM* Lam. = *Nothites melissæfolia*.  
*E. MICROPHYLLUM* L. = *Conoclinium microphyllum*.  
*E. MOLLE* Swartz = *Hebeclinium macrophyllum*.  
*E. NEPETÆFOLIUM* H. B. K. = *Bulbostylis nepetæfolia*?  
*E. NERVOSUM* Sieb. non Sw. = *Baccharis nervosa*.  
*E. NUYANSII* H. B. K. = *Bulbostylis secundiflora*?  
*E. OBTUSIFOLIUM* Willd. = *Vernonia Vahlia*.  
*E. OBOVATUM* Willd. = *Vernonia odoratissima*.  
*E. OLIGANTHES* Less. = *Bulbostylis oliganthes*.  
*E. PARVIFLORUM* Aubl. = *Mikania amara*.  
*E. PARVIFLORUM* Swartz. = *Critonia parviflora*.  
*E. PAUCIFLORUM* H. B. et K. = *Bulbostylis*? *pauciflora*.  
*E. PECTINATA* Wall. = *Vernonia pectinata*.  
*E. PENDULUM* Schrad. = *Bulbostylis pendula*.  
*E. POLYANTHES* Spr. = *Vernonia polyanthes*.  
*E. POLYANTHUM* Wall. = *Vernonia Neelgherryana*.  
*E. POLYSTACHYUM* Willd. = *Vernonia mollis*.  
*E. PYRAMIDALE* Don = ? *Vernonia aspera*.  
*E. RETROFRACTUM* Thunb. = *Pegoletia polygalæfolia*.  
*E. RIGIDUM* Swartz = *Ooclinium rigidum*.  
*E. ROSMARINIFOLIUM* Labill. = *Ozothamnus rosmarinifolia*.  
*E. SATURNEÆFOLIUM* Lam. = *Nothites saturniæfolia*.  
*E. SCANDENS* Linn. = *Mikania scandens*.  
*E. SCANDENS* Link = *Senecio deltoides*.  
*E. SCANDENS* Thunb. = *Mikania auriculata*.

- E. scandens* Wall. = *Mikania pubescens*.  
*E. scariosum* Wall. = *Bulbostylis pendula*.  
*E. secundiflorum* Bert. = *Vernonia arborescens* β.  
*E. secundiflorum* Lag. = *Bulbostylis secundiflora*.  
*E. speciosum* Vent. = *Liatris elegans*.  
*E. stipulaceum* Steud. = *Mikania stipulacea*.  
*E. squarrosus* Cav. = *Bulbostylis Cavanillesii*.  
*E. tomentosum* Lam. = *Senecio penicillatus*.  
*E. uniflorum* Moq. = *Monosis salicifolia*.  
*E. urticifolium* L. f. = *Campulodinium urticifolium*.  
*E. vagans* Wall. = *Vernonia vagans*.  
*E. veronicæfolium* H. B. K. = *Bulbostylis veronicæfolia*.  
*E. vestitum* Spreng. = *Salmea hirsuta*.  
*E. vinçæfolium* Lam. = *Mikania amara*.  
*E. volkameriæfolium* Wall. = *Vernonia volkameriæfolia*.  
*E. volubile* Vahl. = *Mikania volubilis*.  
*E. zeylanicum* L. = *Vernonia Zeylanica*.

LXXXVIII. NOTHITES *Cass. dict.* 35 (1825) p. 163. — *Eupatorii* sp. *Lam. Willd. Less.*

Capitulum oblongum 5-florum. Recept. nudum angustum. Invol. 5-phyllyum, squamis 1-seriatis erectis oblongo-lanceolatis. Cor. tubo brevi, fauce elongatâ intus villosâ. Antheræ subinclusæ, appendicibus obtusissimis. Achænia subpentagona hispidiuscula. Pappi setæ 10-20-uniseriales filiformes apice barbellatæ basi membranâ angustissimâ marginatæ. — Herbar. ex Amer. merid. ortæ. Folia opposita rarius alterna, plerumquæ sessilia, dentata. Corymbi terminales. — Genus mihi ignotum et hic ex cl. Cassini omninò expositum, restitutis tamen nominibus arbitrariò mutatis.

\* *Folius omnibus aut plerisque oppositis.*

1. *N. MELISSÆFOLIA*, erecta pubescens, foliis oppositis subsessilibus ellipticis inæqualiter serratis utrinquæ punctato-resinosis triplinerviis reticulatis, corymbo terminali glanduloso-pubescente, pappi paleis 10-12. ♀ in Peruvîâ. *Eupatorium melissæfolium* Lam. dict. 2. p. 411. *Mikania melissæfolia* Willd. sp. 3. p. 1747. *Nothites latifolia* Cass. dict. 35. p. 163. Cor. purpureæ. Folia a poll. longa, sesquipollicem lata.
2. *N. SATURÆÆFOLIA*, erecta suffruticulosa pubescens, foliis suboppositis sessilibus linearibus integerrimis obsolete lanuginosis, corymbis terminalibus. ♀ 5 in Monte-Video. *Eupatorium saturææfolium* Lam. dict. 2. p. 411. *Mikania saturææfolia* Willd. sp. 3. p. 1747. *Nothites angustifolia* Cass. dict. 35. p. 164. Folia pollicem longa, lineam sesquilineamve lata.
3. *N. BREVIFLORA* (Cass. dict. 35. p. 165), erecta pubescens, foliis oppositis subsessilibus oblongo-lanceolatis basi angustatis et integris supernè serratis, corymbo terminali glanduloso-pilosiusculo, pappi paleis circ. 17. ♀ in Brasiliâ. Folia sesquipoll. longa, semipoll. lata.
4. *N. PETIOLATA* (Cass. dict. 35. p. 166), ramosissima glabrituscula, foliis oppositis alternisque petiolatis ovalibus obtusè serratis subpubescentibus glanduloso-punctatis, corymbo terminali pubescente, pappo 12-15-setoso, et in unico cujusque capituli flore coroniformi dentato. — in Peruvîâ circa Limam.

\*\* *Folii alternis.*

**COMARIDEA**, fruticosa virgata glabra, foliis alternis subpetiolatis linearilatis utrinque acuminatis, aliis medio grossè inciso-serratis, aliis rameis axillis fasciculatis integerrimis, paniculâ elongatâ parçè foliaceâ polyâ, pappi setis circ. 20 scaberrimis.  $\S$  in fruticetis apricis colliam circa viso et Quillota Chilenisiam legit cl. Bertero. Decemb. flor. Baccharis? herb. n. 837. Invol. 1-ser. 5-phyllum. Flores 5. Achænia glabra, matura vulnere breviora. Styli longè exserti obtusi subclavati. Cor. ex sicco succo. (v. s.)

**ATZIFOLIA**, fruticulosa minutissimè subvelutina, foliis petiolatis ovatis autis hinc inde sublobatis denticulatis penninerviis, paniculâ polybithryâ, floridis axillaribus, capitulis brevè pedicellatis confertis, pappi setis circ. scabris.  $\S$  in Mexico legit cl. Hænke. Petioli 6 lin. longi. Limb. fol. 27. longus, 20-24 latus. Capitula ferè Mikaniæ sed 5-flora. (v. s. in h. à cl. de Sternberg.)

**L. MIKANIA** Willd. sp. 3. p. 145a. Kunth nov. gen. am. 4. p. 134. ss. syn. 157. non Neck. — Eupatorii sp. Linn. Less. in linnæa 1830.

alium 4-florum. Recept. nudum angustum. Invol. 4-phyllum, aut ad basin aut infrâ basin bracteolâ. Cor. tubus brevis, faux subcampanulata. Antheræ subexsertæ. Achænium angulatum. 1-serialis pilosus scabridus. — Frutices aut herbæ, sæpius oppositæ. Folia constanter opposita, sæpius petiolata, projecturâ reali inter se ligata, rarò ad basin petioli auriculata, limbo cordato aut ovato sed in genere multiformi. Capitula spicata, corymbosa, aut paniculata. Corollæ albide aut pallide flavae. — Caract. gener. levis equidem, sed genus adeò naturale ne subartificiali species hic subungere tantum potui.

§. 1. *Erectæ, caulibus nempè non scandentibus.*

**URARIFOLIA**, herbacea?, erecta parçè ramosa glabra basi radicans, caule ramis sulcato-angulatis, foliis distantibus sessilibus linearibus acutis in- imis, corymbo terminali composito, capitulis pedicellatis, bracteolâ acutâ minutâ, invol. squamis oblongis mucronatis, achæniis subglan- a.  $\mathcal{L}$  in Brasiliæ prov. Sancti-Pauli. Species habitu à cæteris diversissima, rect. non separanda. Fol. 9 lin. longa, vix lineam lata, internodio multò ra. Cor. pallide flavescentes. Pappus rufus. (v. s. in h. Mus. reg. Par. à imp. Bras. sub n. 505 miss.)

**MINUA**, herbacea? erecta, caule tereti, ramis oppositis striatis virgatis s subaphyllis, foliis oppositis sessilibus linearibus integris remotissimis, ubi terminalis subpaniculati ramis ramulisque pubescenti-hirtellis, capi- edicellati bracteolâ setaceâ hirtelli, invol. squamis lanceolatis acumi- glabris.  $\mathcal{L}$  ? in Brasiliæ prov. Rio-Grande. Habitus ferè Cynanchi aphylli. pollicem longa lineam lata. Capit. 4 lin. longa. Pappus rufus. Achænium pubescens. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub p.)

**ZEATILLOBA**, herbacea erecta? glaberrima, caule tereti striato parçè fo- foliis sessilibus pinnatilobatis, lobis lineari-oblongis integerrimis minutè so-ciliatis, corymbo terminali longè pedunculato, pedicellis involucrique ad dorso pubescentibus, achæniis glabris.  $\mathcal{L}$  in Brasiliæ prov. Rio-Grande. is scandens, ad sect. ultimam prope M. apiifoliam transferenda. (v. s. Mus. Par. à Mus. Bras. sub n. 927 miss.)

**AMBIGUA**, collo suffruticoso, caulibus pluribus herbaceis erectis striatis rufis, apice nudo, foliis sessilibus lineari-oblongis acutis aut obtusis aliis

- integerrimis aliis subdentatis glabrescens subtriplinerviis, corymbo paniculato nudo, invol. squamis lineari-oblongis acutis, bracteolâ is æquali et conformi.  $\mathcal{L}$  in Brasiliæ prov. Rio-Grande. Herba 7-8 pollic. Folia 2 poll. longa, 4 lin. lata. Pappus rufescens. (v. s. in h. Mus. reg. Par. & Mus. imp. Bras. sub n. 944 miss.)
5. *M. OBLONGIFOLIA*, subherbacea? erecta glaberrima, caule tereti striatulo, foliis sessilibus oblongis utrinque vix subacutis integerrimis penninerviis subreticulatis utrinque sublucidis rigidulis, paniculâ terminalis ramis erectis angulatis, capitulis secus ramos racemosis pedicellatis, bracteolâ lineari acuminatâ ad pedicelli basin, invol. squamis linearibus acuminatis glabris, achenio juniore rufo-pubero. — in Brasiliæ prov. Sancti-Pauli. Folia sesquipoll. longa, semipoll. lata. Pappus rutilanti-rufus. (v. s. in h. Mus. reg. Par. & Mus. imp. Bras. sub n. 415 miss.)
6. *M. MYRICAFOLIA* (Bojer! in litt. 1835. sub Trixide), fruticosa erecta, ramulis foliorum projecturis exsertis acutè angulatis araneosis, foliis petiolatis oblongis à medio ad apicem dentatis obtusis basi cuneatis utrinque glabris coriaceis, paniculâ terminali corymbosâ, invol. glabro cylindraceo obtuso, bracteolâ minimâ subaraneosâ.  $\mathcal{L}$  in sylvis prov. Saccalavorum insul. Madagascar legit d. Bojer. Frutex 6-8-pedalis Myricæ habita. Species ex projecturis fol. distinctissima. (v. s. flor. nondum expl.)
7. *M. SESSILIFOLIA*, herbacea erecta, caule simplici tereti hirsuto-subhispido, foliis oppositis sessilibus internodio longioribus imbricatis basi cordatis 3-5-nerviis ovato-orbiculatis obtusis subcrenatis suprâ sparè scabris subtus reticulatis tomentoso-hirsutis, ramis apice conferto-corymbosis simplicibus basi foliolosis apice spicatis, capitulis approximatis, bracteolâ acutâ invol. dimidio breviorè, invol. squamis oblongis obtusis puberis, achenio glabro.  $\mathcal{L}$ ? in Brasiliæ prov. Minas General. legit cl. Vauthier (pl. exs. n. 281). Folia annua subalterna, caulina 2 poll. longa, sesquipollicem lata. Capit. 2-2½ lin. longa. (v. s. comm. à cl. inv.)
8. *M. LIXOLENA*, caule herbaceo erecto tereti densè tomentoso-hirsuto, foliis oppositis sessilibus ovatis (infer. basi vix subcordatis) acutis hinc inde subdentatis suprâ pubescentibus subtus tomentoso-hirsutis, paniculis axillaribus paniculatis et in paniculam dispositis puberulis, invol. squamis oblongis obtusis glaberrimis.  $\mathcal{L}$  in Brasiliæ prov. Minas General. ad Serra do Frio legit cl. Vauthier (pl. exs. n. 331!). Affinis *M. sessilifoliz*, sed disp. capitulorum, invol. glabritie, etc. facile differt. (v. s.)
9. *M. NUMMULARIA*, herbacea erecta, ramis teretibus glabrescens aut pulverulento-velutinis, foliis breviter petiolatis orbiculatis basi subcordatis 5-nerviis hinc inde calloso-denticulatis utrinque tomento minimo canescenti-velutinis subtus reticulatis, medio internodio aequalibus, paniculâ laxâ ramis elongatis ramulisque oppositis, capitulis secus ramulos spicato-congestis, bracteolâ minimâ, invol. squamis ovali-oblongis obtusis dorso velutinis, achenio glanduloso.  $\mathcal{L}$  in Brasiliæ prov. Minas Gener. ad Tejuco legit cl. Vauthier (pl. exs. n. 266). Herba rhizomate lignoso, parè ramosa, bipedalis, habita Vernoniz schwenkizfoliz. Folia diam. 7-8 lin. (v. s. comm. à cl. inv.)
10. *M. XEROCaula*, herbacea erecta tota pube minimâ pulverulentâ cinerescens, caule tereti nervis 8 prominulis notato, foliis petiolatis ovalibus basi subcuneatis apice obtusis margine subrepandis aut subdentatis 3-5-plinerviis, paniculâ amplè ramis oppositis folio triplò circ. longioribus, capitulis fasciculatis breviter pedicellatis, bracteolâ ad basin pedicelli vix eo æquali, invol. squamis oblongo-linearibus obtusis, achenio pubero.  $\mathcal{L}$ ? in Brasiliæ prov. Minas Generaliam prope Sabara legit cl. Vauthier (pl. exs. n. 270). Nervi caulis ex utroque folio 3 et 1 utrinque intermedias, sed in ramulis ordo vis conspicuus. Petiol. 4-5 lin. Limbus 15 lin. longus, 10 latus. Folia furè Malmi, sed opposita. (v. s. comm. à cl. inv.)
11. *M. TERNIFOLIA*, erecta basi suffruticosa, caule subsimplici tereti apice strato sparse hispido, foliis ternato-verticillatis petiolatis ovatis acutis grossè incan-

suprà glabriusculis subitùs secus nervos sparè setoso-hispidis inter eos glandulosi, paniculâ terminali oppositè ramosâ, ramis apice corymbosis verticillatis, capitulis confertis brevissimè pedicellatis, bracteolâ pedicellatâ acuminatâ, invol. squamis oblongis acutis glabris.  $\frac{5}{2}$  in prov. Rio-Grande. Petiol. 3-4 lin. Limb. fol. 10-12 lin. longus, 6 lin. a. in b. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 778.)

**ERIZOXOIDES**, erecta herbacea, caule simplici subpuberulo striato, foliis petiolatis hastato-lanceolatis basi cuneatis apice acuminatis inaequaliter lucido-dentatis pubescentibus, corymbo composito fastigiatis elongatis nudis pubescentibus apice corymbulosis capitulis pedicellatis minimè acutâ, invol. squamis lanceolatis acutis glabriusculis.  $\frac{5}{2}$  in prov. Rio-Grande. Herba sesquipedalis. Folia petiolo incluso aequi poll. lata, summa sessilia basi latiora. Pappus rufus. (v. s. a. reg. Par. à Mus. imp. Bras. sub n. 791 miss.)

**AVONAS**, herbacea erecta, caule simplici crasso angulato basi glabro apice pedunculisque villosis, foliis confertis pubescenti-villosis acuminatis et irregulariter dentatis, infer. breviter petiolatis ovatis basi cuneatis subsessilibus basi dilatatis, corymbo composito conferto, ramis panicula breviter pedicellata fasciculata gerentibus, bracteolâ parvâ invol. squamis lanceolatis acutiusculis dorso pubescentibus.  $\frac{2}{2}$  in Bras. v. Rio-Grande. Herba pedalis aut sesquipedalis. Folia inf. 4-5 poll.  $\frac{1}{2}$  poll. lata. Capitula 5 lin. longa. Pappus rufus (v. s. in h. Mus. reg. m. imp. Bras. sub n. 904 miss.)

**SANDRAE**, erecta suffruticosa, caule ramis petiolisque teretibus hirsutis foliis petiolatis ovato-lanceolatis acuminatis integerrimis, suprà tripartitè setoso-scabris, subitùs penninerviis hirsutis, ramis axillaribus basi capitula glomerata subpaniculata gerentibus, bracteolâ ovatâ nuda glabriusculâ, invol. squamis ovali-oblongis obtusis subcariosis  $\frac{5}{2}$  in Brasiliis prov. Sancti-Pauli. Hirsuties flavescenti-rufa. Invol. a. Petiol. 5 lin. longi. Limb. fol. 3 poll. longus, 15 lin. latus (v. s. a. reg. Par. à Mus. imp. Bras. sub n. 424 miss.)

**ELAEOPOLIA**, fruticosa erecta? ramis teretibus densè et brevè velutinis annulisque ferrugineis velutino-hirsutis, foliis petiolatis ovatis basi raris apicem sensim acuminatis integerrimis subquintuplinerviis suprà abeidis subitùs ferrugineo-hirsutis, paniculis terminalibus oppositè annulosisque, capitulis fasciculatis breviter pedicellatis, bracteolâ parvâ, ramis lineari-oblongis obtusis dorso puberulis, achenio villosis.  $\frac{5}{2}$  in prov. Minarum Generalium ad Mariannam legit cl. Vauthier (pl. exs. Petiol. 8 lin. longi. Fol. limb. 3  $\frac{1}{2}$  poll. longus, 2 poll. latus. (v. s. cl. Inv.)

**GRISPOLIA** (Spreng. syst. 3. p. 423), erecta herbacea, foliis oppositis ovato-oblongis serratis subitùs resinoso-pilosis, paniculâ terminali. — in Brasiliâ. Cœt. ign. †.

**KINHALIS** (Mart. in Isis 1824. p. 587), erecta glabra, caule subimbricè decussatis subtriangulari-ovatis sinu grosso cordatis latere dentatis integerrimis cernuis, paniculis corymbosis terminalibus. — in Brasiliâ *pro de Jesu* dicitur. †.

**NEOPOLIA** (Willd. sp. 3. p. 1746). erecta fruticosa hirsuta, foliis petiolato-ovatis acutis trinerviis scabris serratis, paniculis terminalibus, ramis lanceolatis.  $\frac{5}{2}$  in Amer. merid. *Cacalia cordifolia* Mutis in Linn. 351. excl. var. Linnaeus filius qui solus speciem descripsit dicit in diabolica pubescentia et in descript. nunc glabra nunc glaberrima. — *riocicensis* huc à Spreng. relata est diversa.

**OSA** (Spreng. syst. 3. p. 423), erecta fruticosa glabra, ramis nodosis, paniculis subrhombico-ovatis serratis triplinerviis, paniculâ terminali  $\frac{5}{2}$  in ins. Cubâ. Cœt. ign. †.



20. *M. DELTOIDES* (Poepp. ex Spreng. syst. 3. p. 423), erecta fruticosa glabra, ramis teretibus, foliis oppositis deltoideo-ovatis dentatis triplinerviis subtus pallidis, paniculâ terminali corymbosâ.  $\frac{1}{2}$  in ins. Cubâ. Cst. ign.  $\dagger$ .
21. *M. NASRUSA* (Spreng. syst. 3. p. 423), erecta fruticosa, ramis teretibus hirsutis, foliis oppositis oblongis utrinque attenuatis dentato-serratis subtus resinoso-pilosis, paniculâ terminali corymbosâ.  $\frac{1}{2}$  in Bras. Cst. ign.  $\dagger$ .
22. *M. VISOOSA* (Spreng. neu. entdek. 1. p. 277), fruticosa erecta glabra, foliis spathulatis coriaceis apice crenatis confertis venoso-reticulatis, corymbis composito subglutinoso.  $\frac{1}{2}$  in Brasiliâ. Species dubia ab auct. ipso in system. non omissa.  $\dagger$ .

§. 2. *Spiciformes, nempè volubiles, capitulis secus ramos elongatis sessilibus spicatis.*

23. *M. HASTATA* (Willd. sp. 3. p. 1742), fruticosa scandens, canle tereti juniori subpubescente, foliis petiolatis subcordato-hastatis subdentatis acuminatis glabris, capitulis secus pedunculos axillares terminalesque apophyllos spicatis ad axillam bractearum involucri equalis solitariis, invol. squamis 4 oblongis lanceolato ovatis submicronatis, achenio pulverulento-velutino.  $\frac{1}{2}$  inter fruticosa montium Jamaicae. Swartz! obs. 299. Kleinia, etc. Browne jam. 316. t. 34. f. 3. Flores albi suaveolentes. Sapor herbarum amarus. (v. s.)
24. *M. POLYSTACHYA*, fruticosa scandens, ramis teretibus petiolisque hirsutis, foliis petiolatis ovalibus ovali-lanceolatisve acuminatis integerrimis penninerviis supra scabris subtus præcipue secus nervos hirsutis seu villosis, paniculâ terminali compositâ spicis pluribus distantibus oppositis elongatis et terminâ longiore, capitulis secus rachin sessilibus bracteolâ linearî duplò longioribus, invol. squamis 4 lanceolato-linearibus acutis, achenio glanduloso.  $\frac{1}{2}$  in convallibus et paludosis circa Bahiam legerunt cl. Salzmann et Blanchet (n. 1033 et 1885!). Cor. pallide flavescentes. Petiol. 4-9 lin. longi. Limbus fol. 3-4 poll. longus, 12-18 lin. latus. Capit. 4 lin. longa. (v. s.)
25. *M. PSEUDOSTACHYA*, suffruticosa scandens, ramis teretibus petiolisque pube ferrugineâ subvelutinis, foliis petiolatis ellipticis acuminatis basi subacutis integerrimis supra subscabridis nervis villosis subtus puberulis et crebre glandulosis, paniculâ terminali compositâ, spicis pluribus inferne subapproximatis, terminali longiore, capitulis secus rachin distantibus, bracteolâ linearî invol. 3° 4°ve brevior, invol. squamis dorso vix puberis apice mucronatis, mucrone patulo sæpè deciduo, achenio glandulifero.  $\frac{1}{2}$  in Peruvia? ex coll. Poeppig. Affinis *M. polystachyæ* sed pube minore ferrugineâ, spicis, pappo rufo, etc., differt. (v. s. comm. à cl. invent.)
26. *M. SCABRA*, fruticosa scandens, ramis teretibus petiolisque pube ferrugineâ scabro-villosis, foliis petiolatis ovatis basi obtusis apice acuminatis dentato-serratis penninerviis supra junioribus sparse scabris adultis subglabris subtus reticulatis secus nervulos piloso-scabris inter nervos glanduliferis, racemis spicatis elongatis nunc terminalibus subsolitariis nunc in paniculam dispositis oppositis axill. terminalibusque, bracteolis linearibus pedicelli longitudine, invol. squamis apice mucronatis subrecurvis, achenio glabro.  $\frac{1}{2}$  in Ceylone olim legit cl. Patris. Petiol. 4-6 lin. Limb. fol. 3  $\frac{1}{2}$  poll. longus, 1  $\frac{1}{2}$  poll. latus. Pappus rufus. Rami foliaque obscure ferruginei. (v. s.)
27. *M. HOUSTONIA* (Willd. sp. 3. p. 1742), fruticosa glabra scandens, foliis petiolatis ovatis acuminatis integerrimis, paniculâ terminali compositâ, racemulis oppositis elongatis, capitulis pedicellatis, invol. squamis 4 equalibus.  $\frac{1}{2}$  in Vera-Cruce legit olim Houston. Eupatorium Linn. h. cliff. 396. n. 6. Eup. Houstonia Linn. sp. 1172. An species homonyma Swartzii hac aut ad *M. Badii* referenda? An. Eup. Houstonia Mill. dict. n. 10, qui dixit folia cordata et capitula spicato-racemosa, huc pertinet? Spec. mihi dubia.  $\dagger$ .
28. *M. SELLERI* (Spreng. syst. 3. p. 421), fruticosa scandens, ramis angulari ferrugineis, foliis ovato-oblongis acuminatis integerrimis triplinerviis subtus

ferrugineis, spicis paniculato-ramosissimis, capitulis remotiusculis.  $\zeta$   $\cap$  in Brasiliâ. Cct. ign.

§. 3. *Stipulatæ, nempe volubiles, petiolis utrinque ad basin auriculæ foliaceâ quasi stipulatis, capitulis non spicatis sed corymbosis paniculatisve.*

29. *M. STIPULACRA* (Willd. sp. 3. p. 1745), volubilis glabra, caule tereti, ramulis subtriatis, foliis cordato-hastatis, lobis apiceque acuminatis, vix hinc inde denticulatis ad basin petioli utrinque stipulâ foliaceâ donatis, ramis axill. terminalibusque tripartitis corymbiferis, capitulis longè pedicellatis, invol. squamis oblongis aequalibus vix subacutis, bracteolâ minimâ subulatâ ab invol. subdistante, achenio glabro.  $\cap$  in Brasiliâ ad Rio-Janeiro. Eupatorium stipulaceum Vahl symb. 3. p. 94. excl. forsan Lam. syn. p. 108 citat. Capit. 4 lin. longa. Pappus rutilanti-rufus. (v. s.)

30. *M. MANETILE*, volubilis, caule subangulato glabro, ramis basi hexagonis villosiusculis, foliis petiolatis glabris, infer. cordato-hastatis, lobis later. brevibus acuminatis divaricatis, super. cordatis lobis subrotundis, omnibus basi repando-crenatis spicæ acuminatis, auriculis stipulæformibus latis obtusis subdentatis inter petiolos (ut in Rubiaceis) sitis, ramulis floridis axill. terminalibusque corymbum oligocephalum spicæ gerentibus, capitulis pedicellatis, bracteolis linearibus acutis foliaceis pedicellisque pubescentibus, invol. squamis spicæ subscariosis mucrone acuminatis.  $\cap$  in Brasiliâ circa Rio-Janeiro legit cl. Lund. An Eupatorium crenatum..... p. 19. t. 4. f. 1? sed nec folia seniores obtusa nec bracteolas 2 ad basium invol. video. (v. s. comm. à cl. Lund.)

31. *M. FLEXUOSA*, volubilis glabrinacula, ramis striatis, fol. petiolisque alâ foliaceâ repando-undulatâ marginatis basi in auriculam semi-reniformem auctis, limbis latè ovatis acuminatis basi et apice integris cæterum dentatis penninerviis, paniclâ terminali foliis paulò longiore, ramis oppositis, ramulis capitula 3-5 subsessilibus congesta gerentibus, bracteolâ minimâ, invol. squamis lanceolato-linearibus acutis schæniisque glabris.  $\cap$  in Brasiliâ circa Rio-Janeiro legit cl. Lund. Species distinctissima! Petioli 8-9 lin. longi. Limb. fol. 4 poll. longus, 2½ latus. Capit. 2-2½ lin. longa. (v. s. comm. à cl. inv.)

§. 4. *Ecordatæ, nempe volubiles exstipulatæ corymboso-paniculatæ, foliis indivisis basi non-cordatis.*

\* *Foliis oblongis lanceolatisve.*

32. *M. CLEMATIDRA*, scandens?, ramis teretibus striatis tenuissimè puberulis, foliis oppositis alternisve petiolatis ovali-lanceolatis acuminatis basi subcuneatis dentibus utrinque grossis 3-4 acutis notatis basi trinerviis glabrisculis, corymbo trichotomo laxo, capitulis pedicellatis nudis 5-floris, invol. squamis 7-9 ext. obtusis, int. brevioribus subacutis. — in Nepaliâ legit cl. Wallich. Mikania clematidea Wall! cat. n. 3173. comp. n. 283. Pappus albidus. Achenium acutè 5-gonum pilis minimis sparsis subasperum. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)

33. *M. LIGUSTRIFOLIA*, fruticosa scandens glaberrima, ramis teretibus striatis, foliis breviter petiolatis oblongo-lanceolatis acuminatis penninerviis reticulatis supra nitidis, pedunculis axillaribus et terminalibus racemosis in paniculam dispositis, capitulis pedicellatis, bracteolis linearibus ad basin pedicelli, invol. squamis 4 linearibus acutiusculis acheniisque glaberrimis.  $\mathcal{Z}$   $\cap$  in Brasiliæ prov. Sancti-Pauli. Petiol. 2-3 lin. Limb. fol. 3 poll. longus, 6 lin. latus. Pappus rufus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 420 mis.)

34. *M. MYRIOCEPHALA*, fruticosa scandens glabra, ramis teretibus, foliis petiolatis lanceolatis seu ovali-lanceolatis sensim acuminatis basi subobtusis integerrimis penninerviis, paniculæ terminalis ramis pube brevi hirsutulis oppositis, ramulis capitula racemosa conferta innumera gerentibus, bracteolis ovatis

acutis ad basin pedicellulæ, invol. squamis oblongis obtusis acheniisque glabris. 3 in Brasiliâ prope Bahiam ad Ilheos legit cl. Blanchet. Calicis teres nervulis subangulatus sarmentum vitis subreferens. Petiol. 4-5 lin. Limb. fol. 3-3½ poll. longus, 12-15 lin. latus. Cor ex sicco vid. pallidè flavescens. (v. s. comm. à cl. inv.)

35. *M. ANDRACHNE*, caule..., ramis subvolubilibus striatis glabrescentibus, foliis petiolatis ovalibus obtusè acuminatis integerrimis utrinquè lævibus basi quin-  
tuplinerviis, paniculâ laxâ corymbosâ, ramulis subpuberulis, bracteolâ minimis aut nullis, invol. squamis oblongo-linearibus, acheniis angulatis minutè puberulis. 5 in Brasiliâ circa Bahiam ad Ilheos legit cl. Blanchet. Petiol. 6 lin. longi. Limb. fol. 3 poll. longus, 1½ latus, non malè Arbuti *Andrachnis* folia referentia. (v. s.)

36. *M. RUDDLEI-FOLIA*, fruticosa scandens?, ramis teretibus striatis petiolis foliisque subitis pube tomentoso-pulverulentâ canis, foliis petiolatis lanceolatis seu ovato-lanceolatis basi subobtusis apice acuminatis integerrimis penninerviis suprà glabris, paniculâ ramosissimâ polyccephalâ, ramis oppositis, capitulis secus ramulos subsessilibus conferto-racemulosis, bracteolâ oblongâ canescente brevi, invol. squamis oblongis obtusis glabris apice ciliatis, achenio glabro. 5 (?) in Brasiliâ legit cl. Lhotsky. Petiol. 8 lin. longus. Limb. fol. 3 poll. longus, 15-18 lin. latus. Pappus sordidus. (v. s. comm. à cl. invent.)

37. *M. CLAVELLATA*, scandens?, caule tereti sparè puberulo oppositè ramuloso, foliis breviter petiolatis elongato-lanceolatis utrinquè acuminatis distanter serratis trinerviis suprà sparè puberulis subtus pubescenti-hirtellis, ramulis petiolisque villosis-hirtellis, paniculæ ramulorum quemque terminans ramis oppositis subcorymbiferis, capitulis pedicellatis, bracteolâ parvâ acutâ, invol. squamis oblongis obtusis glabrescentibus, achenio pubero. (?) in Brasiliâ prov. Minarum General. ad Mariannam legit cl. Vauthier (pl. ex. n. 287). Stigmata apice insigniter clavellata! Capit. parva. Fol. a poll. longa. 6-7 lin. lata, petiolo 2-3 lin. (v. s. comm. à cl. inv.)

38. *M. KIOCLADA*, volubilis?, ramis teretibus hirsutis confertâ rufâ petiolisque velutino-hirsutis, foliis petiolatis oblongo-seu ovali-lanceolatis acuminatis integerrimis membranaceis pube sparso subglandulosâ suprà secus nervos venasque dispositâ scabridis, paniculâ terminali foliis internixtâ, ramis axillaribus, ramulis capitula 3-5 pedicellata conferta gerentibus, bracteolâ ad basin pedicelli vix eo longioribus, invol. squamis oblongo-linearibus acutis dorso puberulis, acheniis glabris. (?) in Brasiliâ. Cor. ex sicco vid. pallidè flavescens. Petiol. hirsutus 6 lin. longus. Limb. fol. 3½ poll. longus, 15 lin. latus. (v. s.)

39. *M. SMILACINA*, volubilis glaberrima, ramis striatis, foliis breviter petiolatis ovali-lanceolatis callosè-acuminatis rigidis integerrimis basi 5-7 nervis ceterum reticulatis, paniculâ terminali distanter et oppositè ramosâ, ramis apice capitula congesta sessilibus gerentibus, invol. squamis ovalibus obtusis, bracteolâ squamis simili. (?) in Brasiliâ prov. Minarum General. legit cl. Vauthier (pl. ex. n. 290). Petiol. 5 lin. longus. Limb. fol. 3 ¼ poll. longus, 15 lin. latus. Capit. 8-10 arcuè congesta, 3 lin. longa. Pappus albus crassus. (v. s. comm. à cl. inv.)

\*\* foliis obovatis.

40. *M. OBTUSATA*, fruticosa scandens? glabra, ramis teretibus substriatis, foliis breviter petiolatis obovato-oblongis basi cuneatis apice obtusis subintegerrimis subtus pallidis venis reticulatis, paniculæ laxæ ramis oligocephalis oppositis, ramulis ultimis brevibus sæpè trifidis, capitulis breviter pedicellatis, bracteolâ minimâ, invol. squamis oblongis obtusis. 5 (?) in Brasiliâ prov. Sancti-Pauli. Petiol. 2 lin. Folia ferè 15 lin. longa, 5-6 lata. Pappus rufus. Invol. fœca. margine rufa submembranacea. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 472 miss.)

41. *M. OBOVATA*, volubilis glaberrima, ramis teretibus striatis, foliis petiolatis latè obovatis obtusissimis integerrimis subcarnosis, basi cuneatis trinerviis, paniculâ terminali oppositè ramosâ, ramis apice puberis paniculato-corym-

basis, capitulis pedicellatis, bracteolâ lineari invol. dimidiò breviori, invol. squamis oblongo-linearibus subobtusis apice subciliatis, achenio glabro. (in corvallis Brasiliæ circa Bahiam legit ocul. Salzm. Flores albi. Petiol. 4 lin. Limb. fol. bipollic. sesquipollicem latus. (v. s. comm. à cl. inv.)

42. *M. SALZMANNI* FOLIA, fruticosa scandens glaberrima, ramis teretibus, foliis petiolatis obovatis ovalibusve basi subcuneatis apice obtusis integerrimis coriaceis lincidiâ penninerviis subtus reticulatis, paniculæ ramosissimæ laxæ ramulis subangulatis sparsè puberulis, capitulis secus ramos racemosos distantibus breviter pedicellatis, bracteolâ ad basin pedicelli parvulâ, invol. squamis oblongo-linearibus obtusis, achenio glabro.  $\frac{3}{4}$  (in aridis Brasiliæ circa Bahiam legit cl. Salzm. Petiol. 3 lin. longus. Limb. fol. 20 lin. longus, 15 latus. Species affinis *M. obovatæ* sed consist. fol. coriaceâ nec carnosâ et plurib. aliis notis faciliè distincta (v. s. comm. à cl. Salzm.)

43. *M. PSYCHOTRIODES*, fruticosa scandens? glabra, ramis striato-sulcatis, foliis approximatis ovalibus basi breviter cuneatis vix petiolulatis triplinerviis obtusis integerrimis, paniculæ terminalis ramis pubescenti-hirtellis oppositis apice subcorymbosis, capitulis breviter pedicellatis fasciculatis, bracteolâ oblongâ invol. triplò brevior, invol. squamis oblongis obtusis glabrescens, achenio glabro.  $\frac{5}{8}$  (in Brasiliâ circa Bahiam legit cl. Blanchet. Affinis *M. erithalimæ*. Folia sesquipollicem longa, ferè pollicem lata. Pappus pallide rufescens. Cor. subflavescentes. (v. s. comm. à cl. Blanchet.)

44. *M. ERITHALINA*, fruticosa scandens? glaberrima, ramis teretibus, foliis sessilibus obovatis cuneatis subacutis integerrimis, corymbo tripartito terminali, capitulis pedicellatis confertis, bracteolâ lineari acutâ invol. brevior, invol. squamis ovali-lanceolatis mucronato-acuminatis glabris, achenio pubero.  $\frac{5}{8}$  (in Brasiliâ circa Bahiam in monte Serrato legit cl. Blanchet. Folia circ. 3 poll. longa, 15-16 lin. lata. Capit. 4 lin. longa. Invol. subfuscescentia. Habitus rami in herb. ferè *Erithalis fruticosæ*. (v. s. comm. à cl. inv.)

\*\*\* *Foliis latè ovalibus ovatis aut subrotundis.*

45. *M. BANISTERIÆ*, fruticosa scandens? ramis teretibus pube sericeâ mollicanis, foliis oppositis petiolatis ovatis basi obtusis apice sensim acuminate integerrimis penninerviis, supernè junioribus setosis, adultis nervis exceptis glabris, subtus densè sericeis, paniculæ terminalis ramis sericeo-lanatis ramulis capitula pauca subsessilia congesta gerentibus, invol. squamis et bracteolâ ovalibus obtusis glabris.  $\frac{5}{8}$  (in Brasiliâ? Petiol. 4-8 lin. longi. Limb. fol. 5 poll. longus, 2  $\frac{1}{2}$  poll. latus, supernè reticulo prominulo venarum notatus. Pappus rutilans. (v. s. ex coll. cl. Poeppig.)

46. *M. ANGEYRIÆ*, fruticosa scandens? ramis teretibus petiolisque pube densâ hirsutâ confertâ molli velutinis, foliis petiolatis latè ovatis subrotundiave submucatis aut superioribus angustè et acutè cuspidatis basi 5-nerviis aut quintuplinerviis integerrimis suprâ sparsè setulosis secus nervos villosis subtus undiquè sericeo-villosis, paniculâ terminali, ramis oppositis, infer. axillaribus folii longitudine, capitulis ad apices ramulorum subternis subsessilibus, invol. squamis bracteolisque ovatis obtusis glabris.  $\frac{5}{8}$  (in Brasiliæ prov. Sancti-Pauli. Affinis *M. Banisteriæ*. Petiol. 9-10 lin. Limb. fol. 3 poll. longi, 2 poll. lati. (v. s. comm. à cl. Thoremin.)

47. *M. GUACO* (H. et Bonpl. pl. equin. 2 p. 84. t. 105), herbacea volubilis, ramis teretibus sulcatis hirtis, foliis petiolatis ovatis subacuminatis basi breviter angustatis remotè dentatis reticulato-venosis suprâ scabrisculis subtus hirtis, corymbis axillaribus pedunculatis oppositis, capitulis subternis sessilibus, bracteolis linearibus inv. brevioribus, inv. squamis linearibus oblongis obtusis pubescentibus, achenio glabro.  $\frac{3}{4}$  (in calidisimis humidis ad ripam fluvii Magdalens ubi *Guaco* dicitur et antidotum contra serpentum ictus præstans habetur. H. B. et Kunth nov. gen. am. 4. p. 136. *M. Huaco* Rieux in Cav. anal. n. 18. p. 316 ex Spreng.)

48. *M. TAWALLANA* (H. B. et Kunth nov. gen. am. 4. p. 137), herbacea volubi-

lis, ramis exangularibus hirtis, foliis petiolatis elliptico-ovatis acuminatis basi brevissimè angustatis integerrimis reticulato-venosis suprè subteris subtus pubescenti-hirtis, corymbis axillaribus et terminalibus, capitulis ternis sessilibus, bracteolis linearibus inv. brevioribus, invol. squamis lineari-oblongis obtusis pubescentibus, achenio glabro.  $\gamma$  in Peruvia calida humidis ad ripam fluv. Dande prope Guayaquil. Aff. M. Guaco. M. Tafalla Sprung. syst. 3 p. 422.

49. M. TRINITARIA, fruticosa scandens, caule tereti glabro, ramis floridis striato-angulatis apice ferrugineo-subvolutinis, foliis petiolatis ovatis acuminatis integerrimis suprè glabris subtus penninerviis reticulatis inter venas pale brevi ferrugineæ velutinis, paniculâ corymbosâ trichotomâ, ramulis apice tricophyllis, capitulis sessilibus a lateralibus bracteis lanceolatâ brevi donatis, medio ebracteato, invol. squamis 4 lineari-lanceolatis acutis glabrescentibus, achenio glabro.  $\gamma$  in insulâ Trininitatis Sieb. l. fl. Trin. exs. n. 18a. Petiol. 9 lin. longus. Limb. 4 poll. longus, a poll. latus. Capit. 4 lin. longa. Pappus rufus. Nomen alludit ad patriam et ad capitula terna. (v. s. ex coll. Sieb.)

50. M. LÆVIS, fruticosa glabra scandens, caulibus teretibus, ramulis striato-angulatis, foliis petiolatis latè ovatis obtusè acuminatis integerrimis basi subtruncatis trinerviis utrinquè subnitidis, paniculâ ramosissimâ laxâ, ramulis 1-secu 3-cephalis, capitulis pedicellatis minutè bracteolatis aut modico sessi ebracteolato, invol. squamis 4 oblongis vix acutis, achenio junctore subpuberulo.  $\gamma$  in Brasiliâ. Petiol. pollicaris. Limb. 3-3  $\frac{1}{2}$  poll. longus, 2-2  $\frac{1}{2}$  latus acuminè obtusissimo insignis. Capit. a  $\frac{1}{2}$  lin. longa. Pappus pallidè rufescens. (v. s.)

$\beta$ . *angustior*, foliis paulò brevioribus petiolatis angustius ovatis acumin et obtusè acuminatis.  $\gamma$  in montosis Tejucco prope Rio-Janeiro legit d. Lund. Petiol. 8-9 lin. Limb. fol. 3 poll. longus, 15 lin. latus, numpi dimidio angustior quàm var.  $\alpha$ . Inflorescentia similis. (v. s. eodem. à d. Lund.)

51. M. PANICULATA, subscandens glaberrima, ramis teretibus fistulosis, foliis petiolatis ovatis abruptè et obtusiusculè acuminatis integerrimis subtus pallidis, paniculâ terminali ramosissimâ laxâ, ramis oppositis glabris, capitulis acutè ramulos apicatis distantibus, bracteolâ minutâ, invol. squamis lineari-oblongis obtusis achenisque glabris.  $\gamma$  in Brasiliâ circa Rio de Janeiro arborum altissimas paniculis suis involvit. Flores albi odorati ex Lund. l. Camille diam. pollic. Petiol. 4 pollic. Limb. fol. 4 poll. longus, 1  $\frac{1}{2}$  latus. Panicula pedalis. Pappus pallidè rufescens. (v. s.)

52. M. BRACTEOSA, fruticosa scandens glaberrima, ramis teretibus apice striatulis, foliis petiolatis ovatis basi obtusè apice subacuminatis puncto callosè terminatis triplinerviis nunc integerrimis nunc basi repandis denticulato, paniculæ ovalis breviter terminalis ramis oppositis, capitulis apicato-congestis, bracteolâ amplè concavâ ovali invol. sublongiore, invol. squamis oblongo-linearibus vix acutis, achenio glabro.  $\gamma$  in Brasiliâ ita. Senotæ Catharinas leg. d. Gandebaud et Bader. Petiol. pollicares. Limb. fol. a  $\frac{1}{2}$  poll. longus, pollicem latus. Pappus rufus. Cor. ex sicco purpureoscentes. (v. s.)

53. M. RADIX, fruticosa scandens glaberrima, ramis teretibus, foliis petiolatis ovalibus abruptè acuminatis integerrimis subtripli-secu quintriplinerviis reticulatis coriaceis, paniculæ terminalis ramis oppositis apice trifida, ramulis capitula subsessilibus conferta gerentibus, bracteolis linearibus dilatatis, invol. squamis lineari-oblongis apice subobtusè villosè-hirsutis, achenio glabro.  $\gamma$  in Guadalupâ legit olim Badiër; an fortè etiam in Jamaica si Eupat. Houstonii Swartz obs. p. 300 hæc referendum. Species insignis foliis apice abruptè angustatis et venis singulari modo ramoso-subreticulatis. (v. s.)

54. M. LAURIFOLIA (Willd. sp. 3. p. 1746), fruticosa volubilis glaberrima, ramis angulatis, foliis breviter petiolatis ovatis obtusiusculis basi acutis integerrimis reticulato-triplinerviis subcoriaceis nitidis, corymbis paniculatis confertè axill. oppositis et terminalibus, capitulis ternis subsessilibus, bracteolis....

- invol. squamis oblongis obtusis apice ad dorsum puberulis, achenio glabro.  $\zeta$  in Amer. merid. (Mutis), temperatis Novæ Granatæ prope Mariquita (H. et B.) nec in Mexico ut incantâ Persoon indicavit. Hamb. B. et Kunth nov. gen. am. 4. p. 137. t. 350. *Cacalia laurifolia* Mutis in Linn. f. suppl. 351. An stirps Mutisiana que dicitur erecta eadem ac Humboldtiana?
55. *M. CASSAMPHELINA*, scandens glabra, caule tereti, foliis petiolatis ovatis subcordatis obtusis integerrimis, ramis floridis axillaribus terminalibusque apice trifidis corymbiferis, invol. squamis 4 oblongo-linearibus acuminatis, 5<sup>a</sup> ext. linearibus cæteris dimidiò brevioribus, achenio glabro.  $\zeta$  in Sancto Domingo. Capit. 5 lin. longa. Petiol. 9-10 lin. (v. s.)
56. *M. PLATYPHYLLA*, scandens? glabra, ramis teretibus, foliis petiolatis latè ovatis subrotundis apice abruptè et obtusiusculè cuspidatis integerrimis, ramis flor. axillaribus oppositis paniculatis inter capitula tenuiter puberulis, ramulis capitula plurima spicata congesta sessilibus gerentibus, invol. squamis linearibus glabris subobtusis, bracteolâ acutâ brevi, acheniis glabris  $\zeta$  in Peruvia? ex coll. Peppig. Rami floridi in specim. meo incompleto retrofracti, folio vix longiores. Pappus rufus. Petiol. pollicares. Limb. fol. 4 poll. diam., cuspidè 3-4 lin. longo. (v. s. comm. à cl. inv.)
57. *M. ATTENUATA*, caule herbaceo tereti volubili cum petiolis et pedunculis pube minimâ subpulverulentis, foliis petiolatis latè ovalibus utrinquè longè acuminatis integerrimis membranaceis glabris penninerviis, paniculis axillaribus et terminalibus, ramis oppositis, ultimis apice tricephalis, invol. squamis 4 oblongis obtusis bracteolis 2 acutis minimis stipatis, acheniis glabris terebinaculis. — in Peruvia montibus Guanacocensibus legit cl. Hæmke. Folia 4 poll. longa, 2 poll. lata. Pappus rufus. (v. s. in h. Hæmke ab ill. de Sternberg comm.)
58. *M. LINDLEYANA*, scandens glabra, caule tereti, foliis petiolatis ovatis acuminatis integerrimis trinerviis, ramorum floridorum sessilibus multò minoribus, paniculæ ramis oppositis apice corymbiferis, capitulis pedicellatis, bracteolis ovali-oblongis foliaceis invol. aequalibus, invol. squamis ovali-oblongis obtusis mucronatisve acheniisque glabris.  $\zeta$  in Guianâ. Valdè affinis *M. Hookeriana*, sed capitulis pedicellatis et bracteolis faciliè differt. (v. s. in h. Lindl. à cl. Hooker comm.)
59. *M. HOOKERIANA*, suffruticosa scandens? glabra, ramis teretibus vix striatis, foliis petiolatis ovatis ellipticisve basi obtusis apice acuminatis integerrimis trinerviis, paniculæ terminalis ovatis ramis oppositis, capitulis secus ramos sessilibus approximatis, bracteolis parvis ciliolatis, invol. squamis oblongo-linearibus obtusis apice subciliatis, achenio glabro.  $\zeta$  in Guianâ. Petiol. 9-12 lin. longi. Limb. fol. 2 poll. longus, pollicem latus. (v. s. in h. Lindl. à cl. Hooker comm.)
60. *M. AMARA* (Willd. sp. 3. p. 1744), fruticosa scandens, caule tereti glabrescente, foliis petiolatis ovatis subcordatisve acuminatis subintegerrimis suprâ sparè et tenuissimè scabris subtus pubescenti-hirtellis secus nervos subtomentosis, paniculis axill. terminalibusque divaricatis oppositè ramosis, capitulis confertis ad apices ramulorum ternis subsessilibus, bracteolis brevibus acutis, invol. squamis oblongo-linearibus obtusis acheniisque glabris.  $\zeta$  in Guianâ secus rivulos et ex Vahl in insul. Caribæis? Eupat. parviflorum Aubl. guian. n. p. 797. t. 315. sed panic. delineata nimis laxa. Eup. vineæfolium Lam. dict. 2. p. 410, sed folia utrinquè glabra dicuntur. Eupat. amarum Vahl symb. 3. p. 93, sed pubes fol. dicitur stellata. (v. s. in h. Lindl.)
61. *M. LATIFOLIA* (Smith in Rees cycl. 23. n. 8), volubilis glabra, foliis ovatis acuminatis subintegerrimis subtus pallidis, capitulis corymbosis, bracteis lanceolatis seu ellipticis sessilibus subasperis. — in insulâ Sanctæ Lucie. Affinis dicitur *M. amara*, sed descr. non sufficit. †
62. *M. PYRIFOLIA* (Bojer! in litt. 1835 sub *Trixide*), suffruticosa scandens glaberrima, caule tereti, foliis petiolatis ovatis aut subcordatis acuminatis

dentatis coriaceis nitidis penninerviis inter nervos tenuè reticulatis, paniculâ terminali irregulari, capitulis ad apices ramulorum confertis angustis, invol. squamis acutiusculis, bracteolâ minimâ, achæniis glabris eglandulosis.  $\zeta$  in prov. Petition ins. Madagascar legit cl. Bojer. Flores pallidè flavescentes. Capitula 2 lin. longa. (v. s.)

63. *M. sarcocosa*, caulibus suffruticosis volubilibus teretibus, ramis petiolis pedunculis folisque subtus pube confertâ brevi velutinis, foliis longè petiolatis, infer. subcoarctatis, super. ovatis, omnibus acuminatis suprà minutissimè puberulis subtus reticulatis, paniculâ amplâ laxâ trichotomâ, capitulis ternis, bracteolis acutis minimis, invol. squamis lineari-oblongis obtusis dorso subpuberulis, achæniis pentagonis glabris.  $\zeta$  in Peruanis montibus Orinocensibus legit cl. Hænke. Petioli 2 poll. longi. Limb. fol. 6 poll. longus, 3-3  $\frac{1}{2}$  poll. latus. (v. s. in h. Hænke. ab ill. de Sternberg comm.)

§. 5. *Cordiformes, nempe volubiles exstipulatæ corymboso-paniculatæ, foliis basi cordatis integris aut repando-dentatis.*

64. *M. HÆNKEANA*, caule herbaceo volubili glabro, foliis longè petiolatis cordiformibus acutis basi subdentatis membranaceis apicè glabris subtus pubescenti-subtomentosis, ramis floridis axillaribus terminalibus paniculatis, ramulis oppositis subspiciformibus involucriisque minutissimè puberulis, achæniis angulatis glaberrimis. — in montanis Peruanis Orinocensibus legit cl. Hænke. Affinis videtur *M. Orinocensi*. Infior. media inter spiciformes et paniculatas. (v. s. in h. Hænke. ab ill. de Sternberg comm.)
65. *M. ORINOCENSIS* (H. B. et Kunth nov. gen. am. 4. p. 134), fruticosa volubilis glabra, ramis striatis, foliis petiolatis subrotundo-ovatis cordatis acuminatis basim versum angulato-dentatis reticulato-5-nerviis, corymbis paniculatis, capitulis pedicellatis solitariis, bracteolâ invol. dimid. breviorè, invol. squamis oblongo lanceolatis acutis achæniisque glabris.  $\zeta$  in humidis insule Pararuma in Mission. Orinocensibus. Petiol. 8 lin. longus. Limb. fol. sesquipollic. longus, pollicem latus. Valdè affinis *M. micranthæ*.
66. *M. MICRANTHA* (H. B. et Kunth nov. gen. am. 4. p. 134), herbacea volubilis glabra, ramis striatis, foliis longè petiolatis subrotundo-ovatis cordatis acuminatis sinuato-dentatis reticulato-5-nerviis, lobis baseos divaricatis, corymbis paniculatis, capitulis pedicellatis solitariis, bracteolâ invol. duplò breviorè, invol. squamis oblongo-lanceolatis acutis achæniisque glabris.  $\zeta$  in umbræ prope Caripe Cumanensium. Petioli 2  $\frac{1}{2}$  pollices longi. Limb. fol. 2 pollic. longus, sesquipollic. (in textu verisim. error. sesquilineam) latus.
67. *M. STERNBERGIANA*, scandens glabrescens, ramis teretibus striatis, foliis petiolatis subcordatis subacuminatis in parte latiore subrepandis vix tenuissimè pubescentibus, ramis apice corymbiferis, capitulis congestis parvis, invol. squamis submembranaceis dorso subpubescentibus, 3 subacutis, 4<sup>a</sup> longiore magis acutâ, 5<sup>a</sup> exter. omnibus breviorè angustiore, achænio glabrescens subglanduloso.  $\zeta$  in insulâ Trinitatis. Eupatorium Sieb. ! fl. trin. exa. n. 225. Specim. mancum, sed capitulis parvis à plerisque facillè differt. (v. a. ex coll. Sieb.)
68. *M. LOXENSIS* (H. B. et Kunth nov. gen. am. 4. p. 135), herbacea volubilis glabra, ramis sulcatis, foliis petiolatis ovatis angustato-acuminatis breviter cordatis remotè et obsolete dentatis reticulato-trinerviis, corymbis paniculatis oppositis, capitulis pedicellatis solitariis, bracteolâ invol. dimidio breviorè, invol. squamis oblongo-linearibus acutiusculis achæniisque glabris.  $\zeta$  in temperatis inter Alto de Pulla et urbem Loza Novo Granat. Petiol. 7-8 lin. longus. Limb. fol. 2  $\frac{1}{2}$  poll. longus, 15-16 lin. latus.
69. *M. MICROPTERA*, subvolubilis minutè puberula, caule hexagono, angulis in alam angustissimam subfoliaceam productis, foliis petiolatis cordatis sinu lato, auriculis rotundatis, apice acuminatis mucrone calloso, margine subintegerrimis, paniculâ terminali oppositè ramosâ laxâ, capitulis distantibus pedicel-

latis, bracteolâ oblongâ minimâ, invol. squamis oblongo-linearibus obtusis achæniisque glabris. ( in Brasiliâ prope Bahiam (Blanchet! n. 1710) et in Peruvia? (ex coll. Poeppig!). Petioli pollicares. Limb. fol. 2 poll. longus, sesquipollicem latus. Specim. Poeppigianum ferè glabrum sed non videtur diversum. (v. s.)

70. *M. BATATIFOLIA*, scandens glabra, ramis teretibus, foliis petiolatis cordatis acuminatis in parte latiore repando-angulatis, limbo vix petiolo longiore, ramis apice laxè corymbiferis, capitulis pedicellatis, invol. squamis 4 acuminatis, 5<sup>a</sup> ext. cæteris breviorè angustâ adpressâ, achænio glabro. ( in insulâ Cubâ circa Havanam legit indef. Ramon de la Sagra. Formâ foliorum accedit ad *M. congestam*, sed inflorescentiâ faciliè differt. (v. s. comm. à cl. inv.)
71. *M. TAMOIDES*, scandens glaberrima, caule tereti, ramulis striatis, foliis petiolatis cordatis acuminatissimis in parte latiore subrepandis, ramis nudis apice corymbum confertum gerentibus, capitulis breviter pedicellatis, invol. squamis 4 oblongo-linearibus acutis, 5<sup>a</sup> ext. linearì brevi, achænio glabriusculo. ( in ins. Cubâ juxta Havanam legit cl. A. de la Ossa. Petioli 7-8 lin. longi. Species glabritie perfecta et capitulis congestis faciliè à cæteris distincta. (v. s. comm. à cl. inv.)
72. *M. COMEUSTA*, scandens, caule tereti sparè puberulo, ramis striato-sulcatis, foliis petiolatis cordatis acuminatis in parte latiore sinuato-angulatis suprâ glabriusculis subtus secus nervos pubescentibus, sinu exciso semitotundò, pedunculis axill. terminalibusque capitula corymboso-congesta gerentibus, capitulis pedicellatis, invol. squamis 4 mucrone acuminatis, 5<sup>a</sup> ext. etiam acuminatâ cæteris longiorè, achænio glabro. ( in Porto-Ricco legit cl. Bertero. *M. scandens* ex Porto-Ricco Spreng.! syst. 3. p. 421 in h. Balb. ips. teste, sed excl. syn. Capitula vix 3 lin. longa. Limbus intra sinum sæpè cuneatus. Sinus nervis limitatus. (v. s. comm. à cl. Balbis.)
73. *M. STOMEXATA* (Spreng. syst. 3. p. 421), fruticosa scandens, foliis cordatis angulatis glaberrimis coriaceis sub-5-nerviis, lobis baseos divaricatis inæqualibus, capitulis glomerato-paniculatis. 5 ( in Brasiliâ. Cæst. ign. †.
74. *M. ? ALATA*, volubilis glabra, caule striato, foliis petiolatis ovatis profundè cordatis acuminatis integerrimis, cymis compositis, pedicellis ad latus interius angustè alatis, bracteis lanceolatis, invol. squamis 4-5 obtusiusculis triner-viis. 2 ( in nemorosis Guianæ prope Essequibo. Kleinia alata Mey. esseq. 249. Pappus roseus. Achænia sulcata. Ipse cl. auctor de genere dubitat et hîc ex habitu potius referenda videtur.
75. *M. TESTUDINARIA*, scandens glabra, ramis angulatis vix hînc iudè puberulis, foliis breviter petiolatis cordatis acuminatis rigidis integerrimis sinu angustò, nervis 3 subtus prominulis, ramis tripartitis corymbosis, capitulis pedicellatis, bractea ovatâ acutâ paulò ab invol. distante, invol. squamis 4 oblongis acutis glabris, achænio subpilosulo. ( in Brasiliæ prov. Minarum General. prope Mariannam legit cl. Vauthier (pl. exs. n. 330) et circa Rio-Janeiro cl. Lund. Folia ferè Testudinariæ, sed minora et rigidiora. Petiol. 4 lin. Limb. circ. pollicem longus, 7 lin. latus. (v. s. comm. à cl. inv.)
76. *M. CARNOSULA*, fruticosa scandens, ramis teretibus apice villosis, foliis cordatis obtusis seu reniformi-cordatis subcarnosis suprâ sparè puberulis subtus villosò-hirsutis subintegris sinu lato ad basin petioli stipulas foliaceas duas gerentibus, pedunculis apice corymbiferis, capitulis breviter petiolatis, invol. squamis 4 ovali-oblongis dorso pubescentibus, 2 obtusis, 2 subacutis, bractea subulatâ minimâ, achænio glabro. 5 ( in Brasiliæ montibus Macahe prope Rio-Janeiro mense julio flor. legit cl. Lund. Petiol. 6 lin. longi. Limbus 6-8 lin. longus, 10-12 latus. (v. s. comm. à cl. inv.)
77. *M. OPIFARA* (Mart. in Isis 1824. p. 6. p. 583), volubilis glabra, caule angulato, foliis petiolatis cordatis acuminatis repando-dentatis vel subintegerrimis, adultis obtusiusculis, capitulis corymboso-paniculatis pedicellatis, invol. aqua-



- mis oblongis subacutis, bracteolâ lanceolatâ invol. paulò brevior. ( in Brasi-  
liâ ubi ex Mart. dicitur *Erba de Cobra*. (v. s.)
78. *M. DENTICULATA* (Willd. sp. 3. p. 1744), scandens glabrescens, ramis  
acutè angulatis, foliis petiolatis cordatis obtusiusculis subdenticulatis scabris,  
corymbis terminalibus, capitulis umbellatis pedicellatis, bracteolis. . . . invol.  
squamis linearibus. ( in Surinamo (Vahl), in sylvis Xalapensis (Lam.).  
Eupatorium denticulatum Vahl symb. 3. p. 93. Less. in Linnæ 1830. p. 239.  
Spreng. syst. 3. p. 421. excl. Raf. syn.
79. *M. NATALENSIS*, volubilis herbacea? tota breviter hirsuto-velutina, foliis  
petiolatis cordatis acuminatis sinuato-dentatis 5-7-nerviis, auriculis obtusis.  
corymbis axillaribus oppositis pedunculatis trifidis polycephalis folio paulò  
longioribus, invol. squamis glabrescens oblongis mucronato-acutis  $\frac{1}{2}$ ? ( in  
Africa australi-orient. ad Port-Natal legit cl. Dregé! Petioli pollicares. Limb.  
fol. 1  $\frac{1}{2}$  poll. longas, pollicem latus. Pappus pallide rufescens. (v. s.)
80. *M. CARKENSIS*, caule suffruticoso volubili glabro, foliis petiolatis cordatis acu-  
minatis dentatis suprâ glabris subtis hirsutiusculis, corymbis trichotomis,  
invol. squamis oblongo-linearibus subacutis dorso puberulis, acheniis parvis  
glandulosis. — ad. Cap. Bonæ Spei (Dregé!) in sylvis Coloniz distr. orient.  
Burch! cat. geog. n. 3674. in distr. Uitenhage (Eckl!). Krehl pl. esp. exs.  
n. 199. Eupatorium scandens Thunb. fl. esp. 627. non Linn. *Similima*  
*M. scandenti*, sed diversa præsertim hirsutis paginæ fol. inferioris. Prima et  
usque ad nuperrimos, sola generis species hucusque in Africa australi nota.  
(v. s.)
81. *M. OXYOTA*, glaberrima, caule herbaceo volubili, foliis petiolatis acutè ha-  
tato-triangulatis basi subtruncatis quinquenerviis apice acuminatis subto-  
gerrimis, corymbo amplo trichotomo polycephalo, invol. squamis oblongo-  
linearibus lævibus acuminatis, acheniis glabris sparè glandulosis. ( in Africa  
australi ad Van Staades Rivier legit cl. Dregé! Petioli æquipollic. Limb. fol.  
3 poll. longus, 18-20 lin. latus. Cor. albidæ. Pappus pallide rufescens. (v. s.)
82. *M. DIOSCOREA-FOLIA*, scandens herbacea, caule tereti glabrescens, foliis  
petiolatis cordatis acuminatis margine repando-crenatis, sinu subrotundo,  
suprâ glabris seu pube sparâ minimâ vix puberulis, subtis minutissimè gla-  
dulosis, corymbis pedunculatis trichotomis, invol. squamis submembranaceis  
dorso subpuberulis, 3 mucronato-acutis, quarto obtusis, acheniis glandulosis.  
(?) in ins. Madagascar legit cl. Gondot, locis *Ambanivules* dictis. Flores  
nitidè albi. Petiol. pollicem long. Limb. fol. sesquipollicem longus et latus.  
Pappus rufescens. (v. s. comm. ab. amic. B. Delessert.)
- $\beta$ ? *Bojeri*, foliorum sinu subangustiori, floribus flavis ex cl. Bojer. — in  
sylvâ Bé-Fouroun super arbores scandens. An flores potius ochroleuci aut  
albidi? *Trixis*? *dioscorea-folia* Bojer! in litt. 1835. (v. s. comm. à cl. Bojer.)
- $\gamma$ ? *crenata*, caule hinc inde pilosiusculo, foliis papyraceis, floribus ex albo  
sublilacinis — in sylvis cum var.  $\beta$ , sed eâ multò minor. *Trixis*? *crenata*  
Bojer! in litt. 1835. Fortè ut var.  $\beta$  species propria? (v. s. comm. à cl.  
Bojer.)
83. *M. TRUMBURGIANA* (Bojer! in litt. 1835. sub *Trixide*), herbacea scandens  
glabrescens pube nempe minimâ sparâ pubescens præ. in paginâ folii superi.  
caule tenel tereti, foliis petiolatis cordatis acuminatis latè crenato-dentatis,  
pedunculis terminalibus apice corymbosis, invol. squamis 4 acuminatis gla-  
brescens, bracteolâ lineari foliaceâ dimidiam invol. longitudinem æquante,  
achenio angulato glabro. (?) in ins. Madagascar prov. Bé-Fani-Mona in sylvis  
Bé-Fouroun apricis super dumeta scandens. Flores flavi et herba fortè amara ex  
Boj. (v. s.)
84. *M. FLORIBUNDA* (Bojer! in litt. 1835. sub *Trixide*), herbacea scandens.  
caule tereti petiolisque hirsutulis, foliis cordatis subdentatis acuminatis utri-  
que sparè et minutissimè puberulis, paniculâ terminali amplâ ramodivisa  
polycephalâ, invol. squamis 4 acuminatis dorso subpuberulis, bracteolâ linear

foliaceâ invol. dimidium longit. æquante, achænio angulato glabro glandulifero.  $\gamma$  in sylvis umbrosis prov. Bé-Zoussong ins. Madagascar legit cl. Bojer. (v. s.)

86. *M. gonoclada*, fruticosa scandens, ramis acutè hexagonis præsertim ad angulos villosis, foliis petiolatis cordatis acuminatis in sinu subcuneatis hinc inde dentatis suprâ sparse puberis subtus subvillosis, pedunculis axillaribus terminalibusque apice corymbosis, capitulis subfasciculatis, invol. squamis 4 oblongis dorso subvillosis, 3 acutis, 4<sup>a</sup> obtusâ, bracteolâ lineari-acuminatâ, achænio elongato glabro acutangulo.  $\gamma$  in Cubâ juxta Havanam in sylvis humidis legit cl. R. de la Sagra et in Mexico circa Tampico de Tamaulipas cl. Berlandier. Pappus rufescens. A *M. Guaco* differt foliis cordatis et ramis angulatis. (v. s. comm. à cl. inv.)

$\beta$ . *ambigua*, ramis hexagonis, foliis cordatis in sinu non cuneatis, summis vix cordatis, bracteolâ angustiori.  $\gamma$  in Mexico ad Tampico de Tamaulipas (Berl.), in Brasiliâ circa Bahiam (Salm. ! Blanch. !). An species distincta? (v. s.)

86. *M. convolvulacea*, suffruticosa scandens, caule ramisque teretibus junioribus pubescentibus, foliis petiolatis latè cordatis acuminatis subrepando-dentatis suprâ vix puberulis subtus pubescentibus subvillosis, ramis apice corymbosis in paniculam laxè dispositis, capitulis breviter pedicellatis, invol. squamis 4 oblongis acutis dorso pubescentibus, adjectâ 5<sup>a</sup> exteriore lineari foliaceâ cæteris dimidiò breviorè, achænio glabro.  $\gamma$  in Sancto Domingo legit cl. Poiteau. Affinis *M. gonoclada*, sed ramis teretibus facillè differt. (v. s.)

$\beta$ . *Portoricensis*, ramulis striatis, foliis subintegerrimis summis minùs cordatis, ramis pubescenti-hirtellis, capitulis achæniisque panlò minoribus.  $\gamma$  in Porto-Ricco legit cl. Bertero. *M. amara* Bert. ! herb. non Willd. *M. cordifolia* Spreng. ! in h. Balb. syst. 3. p. 423. excl. syn. An fortè species propria? (v. s.)

87. *M. volubilis*, scandens, caule tereti subpubescente, foliis petiolatis cordatis crenatis acutis, lobis basèos æqualibus obtusis, venis subtus petiolisque subpubescentibus, corymbo paniculato trichotomo nudo, capitulis pedicellatis 4-floris, invol. squamis circ. 6-8, inter 4 oblongis acutis, exter. brevioribus subpancioribus, pappo rabente.  $\gamma$  in moderatis dametis et sylvis Ind. or. ad Penang, Singapore, etc. (Wall. !), in Javâ (Blum. !). Eupatorium volubile Vahl symb. 3. p. 93. Eupatorium cordatum Burm. ind. t. 58. f. 2. Mikania volubilis Willd. sp. 3. p. 1743. Blum. ! bijdr. 904. Wall. ! cat. n. 3174. comp. 284. Fl. albi. Achæn. angulata glabra glandulosa. Variat pubescens aut glabra foliis plùs minùs crenatis aut subintegris. An satis à *M. scandenti* diversa? (v. s.)

88. *M. PARKERIANA*, volubilis, caule tereti minutissimè pubescente, ramis striato-sulcatis glabrescentibus, foliis petiolatis cordatis acuminatis membranaceis pubescentibus minutissimè sub lente scabriusculis in parte latiore subangulatis, sinu semirotondo, corymbo terminali composito suprâ ultima folia subsessilibus, capitulis congestis subsessilibus, bracteolâ lineari, invol. squamis ovali-oblongis membranaceis acutè mucronatis achæniisque glabris.  $\gamma$  in Guianâ circa Demerari legit cl. Parker. Affinis *M. scandenti* et pubescenti, sed capitulis in glomerulos plurimos densè congestis et invol. mucronatis distincta. Petioli bipollicares. (v. s. comm. à cl. Parker.)

89. *M. scandens* (Willd. sp. 3. p. 1743. excl. syn. Plum.), caule volubili glaberrimo, foliis petiolatis cordatis acuminatis basi repando-crenatis glabris, ramis axillaribus apice corymbum subfastigiatum gerentibus, invol. squamis 4 acuminatis subciliolatis, adjectâ 5<sup>a</sup> exteriore breviorè, achæniis glabris subglandulosis.  $\gamma$  ad margines rivorum Amer. bor. à Canada australiori? (Hook.) ad Carolinam super. (Michx.) et fortè in Mexico ex specim. Berlandieriano (pl. exs. n. 2272) prope Victoria lectis et Bostonianis similissimis. Pursh fl. 2. p. 517. Nutt. gen. 2. p. 136. Ell. sketch 2. p. 292. Lindl. in trans. hort. soc. Lond. 7. p. 2. p. 250. Eupatorium scandens Linn. sp. 1171. Jacq. Æc. rar. 1.

- t. 169. coll. 1. p. 108. *M. angulosa* Rafin. fl. lud. p. 62? — Pluk. alm. t. 163. f. 3. Flores albid. Pappus albus. (v. s.)
90. *M. RUBESCENS* (Muhl. cat. 71. Nutt. gen. am. 2. p. 136), volubilis, caule tenuiter pubescente, foliis petiolatis cordatis acuminatis basi repando-crenatis angulatisque, ramis axillaribus apice corymbum subfastigiatum gerentibus. invol. squamis 4 acuminatis ciliolatis et pubescentibus, adjectâ 5<sup>a</sup> brevioris ext. angustiore, achenio glabro subglanduloso. (✓) in Carolinâ infer. et Georgia locis humidis. Ell. sketch 2 p. 293. *Eupatorium scandens* Walt. car. 198. ex h. Fraser! Flor. pallidè purpurei odori. Species valdè *M. scandenti* affinis. Stirps Martinicensis huc à Spreng. relata verisim. diversa. (v. s.)
91. *M. LAXA*, volubilis, caule tereti glabro, ramis vix striatulis præsertim in paniculâ pube minimâ rufescente velutinis, foliis petiolatis cordatis acuminatis subintegerrimis trinerviis membranaceis, paniculâ oppositè ramosâ laxâ, capitulis longè pedicellatis secus ramulos subracemosis, bracteolâ oblongo-lineari acutâ invol. dimidiò brevioris, invol. squamis obtusis. (✓) in Peruvâ legit cl. Poeppig. Petioli pollicares. Limb. fol. 3 poll. longus, 2 poll. latus. Pappus rufus. (v. s. ex coll. Poeppig.)
92. *M. MICRODONTA*, suffruticosa scandens, ramis teretibus pube brevi velutino-hirsutulis, foliis petiolo brevi retrorsum hirsutissimo, limbo basi cordato ovato-lanceolato acuminato dentibus parvis raris notato suprâ sparse setoso subtus densè et molliter hirsuto, ramis axillaribus terminalibusque paniculatis oppositè ramosis, ramulis capitula 3-5 congesta subsæcilia gerentibus, invol. squamis bracteolisque ovatis subobtusis glabris. (✓) in Brazilia prov. Minarum General. prope Tejucco legit cl. Vauthier. (pl. exa. n. 291). Petiol. 3 lin. long. Limb. fol. 3-3½ poll. longus, 12-15 lin. latus. (v. s.)
93. *M. MICROCEPHALA*, caule volubili tereti pube brevi subglandulosâ hispidulo, foliis petiolatis cordatis sinu angustato acuminatis dentatis basi 5-nerviis suprâ sparse scabridis subtus præcipuè secus nervos subhirsutis, paniculâ amplâ oppositè ramosâ, capitulis secus ramulos spicato-racemosis subsæcilibus, invol. squamis 4 mucronulatis dorso subpuberis, bracteolâ minimâ, achenio glabro. (✓) in prov. Minarum General. Brazilia legit cl. Vauthier (pl. exa. n. 288). Pars ramorum superior ferè aphylla. Capitula vix 2 lin. longa. Fol. inf. sequipollicem longa, et ferè lata. (v. s. comm. à cl. inv.)
94. *M. HIRsutissima*, ramis scandentibus teretibus pedunculis petiolis folisque subtus villosis-hirsutissimis, foliis petiolatis cordatis acuminatissimis subintegerrimis suprâ sparse puberis, ramis paniculam amplam ramosissimam myriocephalam gerentibus, invol. squamis 4 oblongis subacutis glabris, bracteolâ paulò infrâ involacrum sitâ squamis simili, achenio glabro. (✓) in Brazilia prov. Minarum General. prope Mariannam legit cl. Vauthier (pl. exa. n. 280). Petiol. 2 poll. Limb. fol. 5 poll. longus, 2 poll. latus. (v. s.)
95. *M. TRICHOPIILA*, fruticosa scandens, ramis teretibus pilis rufis confertis petiolis pedunculisque hispidissimis, foliis petiolatis suprâ sparse setosis subtus villosis-hispidis, infer. subcordatis angulatis, super. ovatis, omnibus integerrimis acuminatis, pauculâ ramosissimâ, invol. squamis 4 obtusis vix puberis, bracteolâ subacutâ, achenio glabro. (✓) in Brazilia circa Bahiam legit cl. Blanchet (n. 2097!). Affinis *M. hirsutissima* et forsitan cum eâ necnon cum *M. visniæfoliâ*, eriocladâ et clavellatâ sectionem naturalem format ex ramis hispidis distinctam. Fol. inf. 8 poll. longa, 5-6 lata. (v. s.)
96. *M. MENISPERMEA*, volubilis tota confertè breviterque hirsuta, foliis petiolatis cordatis acuminatis in parte latiore repando-dentatis utrinquè inter hirsutem purè glandulosâ, ramis axill. terminalibusque apice corymbosis, capitulis pedicellatis, invol. squamis 4 acuminatis dorso puberis, bracteolâ linearis-subulata, invol. vix brevioris, achenio glanduloso. (✓) in Mexico circa Bejar mense Augusto flor. legit cl. Berlandier. Capitula 2½ lin. longa. Pappus subrufescens. (v. s. comm. à cl. inv.)
97. *M. POWELLII* (Spreng. syst. 3. p. 422), volubilis, caule subangulato gl.

brinseolo, foliis petiolatis cordatis subhastatis acuminatis repando-crenatis glabruiculis, auriculis subacutis parce divergentibus, pedunculis axillaribus oppositis folio duplò longioribus apice corymbiferis, capitulis confertis brevissimè pedicellatis, bracteolà minimà, invol. squamis ovali-oblongis minutissimè mucronatis, achenio glanduloso.  $\curvearrowright$  in insulâ Cubâ. Mikania denticulata. Poepp. ex Spreng. non Willd. Mikania n. 59. Poepp. pl. exs. Capitula vix 2 lin. longa. Affinis M. menispermæ. (v. s.)

98. M. MOLLIS (H. B. et Kunth nov. gen. am. 4. p. 135), fruticosa volubilis, ramis sexangularibus canescenti-tomentosis, foliis petiolatis subrotundo-ovatis profundè cordatis subacuminatis subintegerrimis reticulato-5-nerviis suprâ pubescenti-hirtis subtus molliter cano-tomentosis, corymbis paniculatis, capitulis semilibus solitariis, bracteolâ acuminatâ invol. breviorè, invol. squamis lineari-oblongis obtusis extus pubescentibus, achenio glabro.  $\curvearrowright$  in temperata Peruvîæ ad ripas flum. Guancabambæ. Petioli æsequipollia. Limb. fol. 2  $\frac{1}{2}$  poll. longus, 2 poll. latus.

99. M. SUAVIOLENS (H. B. et Kunth nov. gen. am. 4. p. 135), fruticosa volubilis, ramis sexangularibus foliisque molliter hispido-pubescentibus petiolatis subrotundo-ovatis profundè cordatis acuminatis grosse dentatis reticulato-5-nerviis, corymbis axillaribus oppositis subpaniculatis, capitulis subpedicellatis subternis, bracteolâ acuminatâ inv. breviorè, invol. squamis lineari-oblongis acutis pedicellis pubescentibus.  $\curvearrowright$  in umbrosis humidis prope Guarumo Novo-Granatensium. Petiol. 1  $\frac{1}{2}$ -2 poll. longus. Limb. fol. 4 poll. longus, 3 poll. latus. Affinis M. scandenti.

100. M. LANUGINOSA, scandens tota ferè hirsutie molli confertâ albâ lanuginosâ, foliis petiolatis cordatis acuminatis except. sinu et acumine dentatis suprâ villosis viridibus subtus lanuginosis canis, paniculis ramos terminantibus oppositè ramosis, capitulis secus ramulos breviter racemosi pedicellatis, bracteolâ ovatâ acutâ glabruiculâ paulò infrâ invol., squamis invol. lanceolatis acutis glabris.  $\curvearrowright$  in Brasiiliæ prov. Sancti-Pauli. Petioli 9 lin. longi. Limb. fol. 2  $\frac{1}{2}$  poll. longus, 2 poll. latus. Invol. rufescentia. Pappus albus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 425 miss.)

101. M. RUBIGINOSA (Smith in Rees cycl. v. 23. n. 12), volubilis?, ramis velutinis, foliis petiolatis cordatis acutis undatis tenuiter velutinis utrinquè globulos resinosos gerentibus, paniculis axillaribus cymosis bracteis spatulatis, invol. squamis obovatis obtusis mucronatisve extus velutinis.  $\curvearrowright$ ? in Amer. merid.? Cacalia cordifolia var. Linn. f. suppl. 352. Numerus florum dubius. †.

§. 6. Angulatæ, nempe volubiles exstipulatæ corymboso-paniculatæ, foliis basi truncatis seu cordatis angulato-dentatis lobatisve.

102. M. CENOPODIFOLIA (Willd. sp. 3. p. 1745 non Wall.), tota glabra, caule herbaceo? scandente tereti striato, ramulis ultimis angulatis, foliis petiolatis triangulari-ovatis acuminatis repando-dentatis 5-nerviis, corymbo panienlato trichotomo laxo, capitulis pedicellatis 4-floris, invol. squamis circ. 8 acuminatis, inter. longioribus.  $\curvearrowright$  in Sierra Leona (Willd.), in arenosis siccis circa Albreda Senegaubiæ (Perrottet!). Affinis volubili. In specim. obviis capitula ferè omnia abortiva, squamis elongatis crispis. In paucissimis vidi achenia teretiuscula, pappum albidum. (v. s. comm. à cl. Perrottet!)

103. M. BRACHYPODA, fruticosa scandens?, ramis teretiusculis apice subangulatis glabris, foliis brevissimè petiolatis acutè triangularibus aut rariùs subrhombeis dentatis 7-nerviis glabris suprâ minutissimè scabridis, corymbo terminali fastigiato laxo, capitulis pedicellatis, bracteolâ lanceolatâ acutâ, invol. squamis lanceolatis mucrone acutis dorso hinc indè subpuberis, achenio glanduloso.  $\curvearrowright$  in Brasiiliæ prov. Sancti-Pauli. Rami tennes rigidi. Petiol. lineam longi. Limb. fol. 8-g lin. longus latusque. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 406 miss.)

104. M. DIVERSIFOLIA, volubilis glabra, ramis feretibus subtiliter striatis, foliis

petiolatis membranaceis, caulinis (seu ramor. præcip.) hastato-triangularibus lobis lateralibus divaricatis terminalique acuminatis integerrimis, ramulis ovatis acuminatis basi truncatis, paniculâ ramis oppositis parvè villosis, capitulis ad apices ramulorum congestis brevè pedicellatis, bracteolâ lineari-subulatâ parvâ, invol. squamis oblongo-linearibus obtusis glabris apice summo puberulis. (in sepibus circa Rio de Janeiro sat rara obs. cl. Lund fl. junio. Invol. straminea. Petiol. fol. caul. 8-9 lin. Fol. 3 poll. longa et lata; petiol. fol. ram. 6 lin. longus, limbus 20-21 lin. longus, 7-8 latus. (v. s. comm. à cl. Lund.)

105. *M. nivoarum*, volubilis, caule tereti vix striatulo glabro, ramis apertè et ad apicem confertè pilosiusculis, foliis sparsè puberis petiolatis trinerviis, caulinis (aut ram. præcip.) subhastato-trilobis lobis et apice acuminatis, lateralibus multò brevioribus, paniculâ terminalis laxè ramis villosiusculis, capitulis pedicellatis, bracteolâ lineari-subulatâ parvâ, invol. squamis linearibus obtusis schenitque glabris. (circa Bahiam Brasiliæ legit cl. Blanchet. Petiol. villosuli fol. caulina. 15 lin. longi, limbus 4 poll. longus, 3 latus; fol. ram. petiol. 5 lin., limbus 2 poll. longus, pollicem latus. Anthera exserta; faux cor. campanulata. (v. s. comm. à cl. inv.)

106. *M. angularis* (H. et Bonpl. pl. equin. 2. p. 87. t. 106), herbacea volubilis glabra, ramis teretibus sulcato-striatis, foliis petiolatis triangulari-hastatis acutis remotè et obsolete denticulatis reticulato-trinerviis, corymbis paniculâ oppositis, capitulis subulterioris pedicellatis, bracteolâ lineari acutâ invol. dimidio breviorè, invol. squamis lanceolatis acutis. (locis humidis in ripa flum. Catamago Novæ-Granatæ H. B. et Kunth nov. gen. am. 4. p. 136. Petiol. bipollis. Folia 7 poll. longa. Species affinis *M. Loxensis* dicitur.

107. *M. scandens*, volubilis glabra, ramis teretibus substriatis, foliis petiolatis trilobis basi truncatis aut subcordatis 3-5-nerviis, lobis lateralibus brevibus terminali acuminato, marginibus integerrimis, paniculâ ramosissimâ amplè ramis ramisque oppositis velutinis, capitulis ad apices ramulorum densè glomeratis subsessilibus, bracteolâ et invol. squamis aequalibus oblongo-linearibus subacutis glabrisculis. (in Brasiliæ sylvis primævis super imas arborum scandens obs. cl. Lhostky. Petiol. 8-9 lin. longi, fol. limb. 2 ½ poll. longus, 2 poll. latus. (v. s. comm. à cl. Lhostky.)

108. *M. redempfolia*, fruticosa volubilis glaberrima, caule striato-sulcato, foliis petiolatis cordatis hastato-trilobis, lobis acuminatis, nervis è nerviâ basi utriusque 2-3 approximatis, paniculâ terminalis ramis ramisque oppositis, capitulis ad apices ramulorum congestis subsessilibus, bracteolâ lineari-subulatâ parvâ, invol. squamis linearibus subacutis schenitque glabris. (in Brasiliâ circa Rio de Janeiro legit cl. Lund, et circa Bahiam cl. Blanchet (n. 3111)). Caulis crassitie digiti. Folia summa ferè triangularia; infer. petioli ferè bipollicares, limb. 3 ½ poll. diam. (v. s. comm. à cl. Lund.)

109. *M. vitifolia*, scandens? herbacea, caulibus fistulosis ramisque teretibus vix striatis sparsè petiolisque pubè articulatis hispidulis, foliis longè petiolatis cordatis 7-nerviis orbiculatis 7-angulatis lobis brevibus acuminatis dentibus callosis exsertis remotis, utriusque secus nervos hirsutis ceterum sparsè scabridis, paniculis amplis oppositè ramosis, ramis ipsis paniculatis, capitulis pedicellatis, bracteolâ lineari acutâ, invol. squamis linearibus obtusis schenitque glabris. — in Brasiliæ prov. Sancti-Pauli. Folia Vitis sed panis majora, ferè potius Sparmanniæ. Paniculæ pedales æquipedales. Petiol. 4 poll. Limb. 4-5 poll. diam. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 426 min.)

**§. 7. Partitæ, nempe volubiles exstipulatæ corymboso-paniculatæ, foliis in segmenta seu partitiones profundè partitis.**

110. *M. apifolia*, suffruticosa scandens glabrescens, caule tereti glabro, ramis subangulatis apice tenuissimè rufo-velutinis, foliis petiolatis trisectis, segmentis petiolulatis ovatis trifidis dentatis acutis glabris, pedunculis axillaribus terminalibusque apice laxè corymbosis, bracteolâ lineari acutâ folliculâ à basi

invol. subdistante, invol. squamis linearibus elongatis acutis schæzizisque glabris.  $\zeta$  in Brasiliæ prov. Sancti-Pauli. Species distinctissima. Pappus rufus. Capit. 5 lin. longa. Cor. ex siccio purpurascens. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 421 mis.)

111. *M. DENTATA* (Spreng. syst. 3. p. 422), herbacea scandens, ramis angulatis, foliis compositis simplicibusque ovatis lanceolatisque angulatis argutè dentatis glabris, paniculis axillaribus diffusis, capitulis solitariis.  $\zeta$  in Brasiliâ. Cact. ign. An. affinis *M. apitifolia*. †.

† *Species dubia.*

*Obs. 1.* Ad Mikania genus referenda videntur stirpes sequentes in vol. 8<sup>vo</sup> floræ fluminensis iconibus nec descriptionibus incompletè illustratæ nempe: *Cacalia cordata* t. 53 (affinis *M. microptera*), *Cac. trilobata* t. 54, *Cacal. angulata* t. 55, *Cac. ternata* t. 56, *Cac. obsoleta* t. 57, *Cac. villosa* t. 58?, *Cac. nitida* t. 59, *Cac. reticulata* t. 60, *Cac. pilosa* t. 61, *Cac. triangularis* t. 62, *Cac. septemata* t. 63?

*Obs. 2.* Specimen incompletum ex Sierra-Leona ortum, *M. tamoidi* affinis vidi in. h. Lindl. Species sola hucusquè in Africâ æquinoctiali nota, verisim. nova.

†† *Species exclusæ.*

*M. ARBORÆA* H. B. K. = *Eupatorium dendroides*.

*M. ARBORÆA* Roxb. = *Lachanodes prenanthiflora*.

*M. ARTEMISIOIDES* Cass. = *Eupatorium leptophyllum* β.

*M. AURICULATA* Willd. = *Senecio deltoides*.

*M. CHENOPODIFOLIA* Wall. = *Senecio chenopodifolius*.

*M. MELISSIFOLIA* Willd. = *Nothites melissæfolia*.

*M. ASPARAGOIDES* Licht. = *Euryopa*.

*M. SATURNEIFOLIA* Willd. = *Nothites satureisefolia*.

*M. SERRATIFOLIA* H. B. K. = *Eupatorium serratifolium*.

*M. TOMENTOSA* Willd. = *Senecio penicillatus*.

XC. *ADENOSTYLES* Cass. dict. 1. suppl. p. 59. Less. syn. 157 non Blum. — *Cacalia* Tourm. inst. p. 451. t. 258. *Fail.* *Adans. Hall. Gærtn. Gaudin. non Cass. nec Less.* — *Cacalia* sp. Linn. Willd.

Capitulum pauciflorum discoideum. Recept. angustum nudum. Invol. cylindraceum 1-seriale oligophyllum. Cor. tubulosæ, limbo campanulato 5-dentato. Styli rami longè exserti semicylindrici undiquè papilloso-glandulosi. Achænium teretiusculum striatum. Pappus pluriserialis setis piliformibus scabris. — Herbe Europæ montanæ perennes. Folia alterna petiolata sæpè basi auriculata reniformia. Capitula in corymbum compositum nudum disposita pedicellata. Flores purpurascens aut pallidi imò albi. — Evidenter ex veterum nomenclaturâ hoc genus *Cacalia* nomen servare debuisset, sed Cassinianam designationem admisi; ne confusionem jam maximam augeam!

1. *λ. GLABRA*, foliis cordatis dentatis utrinquè glabris, corymbis fastigiatis, capitulis 3-6-floris.  $\zeta$  in montanis et subalpinis (nec in alpinis) petrosis irriguis Europæ, non rara in Galliâ, Germaniâ, Italiâ, etc., rara in Siciliâ (Guss.). *Cacalia alpina* var. β Linn. sp. 1170. *Cacalia glabra* Vill. dauph. 3. p. 170. *Cacalia alliarisefolia* Lam. dict. 1. p. 53a. *Cacalia alpina* Jacq. fl. austr. t. 234. Willd. sp. 3. p. 1735. Sturm fl. germ. icon. Tusilago *Cacalia* Scop. carn. n. 1055. *Aden. viridis* Cass. l. c. *Aden. alpina* Bluff et Fing. comp. fl. germ. 2. p. 329. *Senecio glaber* Clairv. man. val. 240. Lob. ic. t. 592. f. 1. Petioli, etiamsi sub-plexicaules, non basi appendiculati. (v. v.)

2. *A. PETASITES* (Bluff et Fing. comp. fl. germ. 2. p. 329), foliis cordatis dentatis subtus incanis superne glabriusculis, corymbis fastigiatis, involucri glabris 3-6-floris.  $\frac{1}{2}$  in pascuis et sylvaticis montium Europæ. *Cacalia alpina* var.  $\alpha$  Linn. sp. 1170 (cum seq. mixta ex syn.). *Cacalia albifrons* Linn. f. suppl. 353. *Cacalia Petasites* Lam. dict. 1. p. 531. DC. fl. fr. n. 3104. *Cacalia alliarie* (Gou. illustr. 65. *Cacalia hirsuta* Vill. dauph. p. 152. *Cac. tomentosa* Jacq. fl. austr. 1. 235. *Adenostyles albida* Cass. l. c. *Adenostyles albifrons* Reich. fl. exc. 2. p. 278. *Senecio hirsutus* Clairv. man. val. 241. — *Clus. hist.* 2. p. 115. f. 1. Petioli sæpius basi utrinque auriculâ appendiculati. (v. v.)  
 $\beta$ . *nuda* (DC. fl. fr. n. 3104  $\beta$ ), petiolis basi exappendiculatis. — *Rarior. Cac. albifrons*  $\beta$  Gaud. fl. helv. 5. p. 216. Invol. 3-6-flora ut in var.  $\alpha$  nec 15-20 ut in sequenti. (v. s.)
3. *A. LEUCOPHYLLA* (Reich. fl. exc. 2. p. 278), foliis cordatis utrinque albomentosis dentatis corymbo subrotando coarctato. involucri dorso lanatis 15-20-floris.  $\frac{1}{2}$  in summorum Alpium petrosis, in Delphinata, Pedemontio, Sabaudia, Valesia. *Cacalia leucophylla* Willd. sp. 3. p. 1736. DC. fl. fr. n. 3105. *Cac. tomentosa* Vill. dauph. 3. p. 171. non Thunb. nec Jacq. *Adenost. candidissima* Cass. l. c. *Senecio tomentosus* Clairv. man. val. 240. non Salisb. *Cacalia alpina* var. *tomentosa* All. ped. n. 644. Petioli basi nudi aut vix subauriculati. (v. s.)
4. *A. HYBRIDA*, foliis cordatis dentatis supra glabriusculis subtus araneosis, petiolis basi sæpius auriculatis, corymbo laxiusculo, involucri squamis dorso puberis 12-18-floris.  $\frac{1}{2}$  in Alpibus Helveticis (Gand!), Delphinensibus (Vill!) et in Sicilia montibus Nebrodensibus (Duby!). *Cacalia tomentosa*  $\beta$  hybrida Vill. dauph. 3. p. 171. DC. fl. fr. n. 3105. Gaud. fl. helv. 5. p. 216. Omnino media inter *Petasitem* et *leucophyllam*, priori foliis, posteriori florum numero affinis! fortè utriusque hybrida progenies? Hanc inter species donec lux fiat admisi quia omnino anxius hæreo an *Petasitis* an *leucophyllæ* sit varietas, et quia reperta fuit in insulâ ubi neutra aliarum adhuc cognita! (v. s.)

#### Subtrib. II. TUSSILAGINÆ Less. syn. 158.

Capitula heterogama aut subdioica.

#### Div. I. PETASITÆ.

Capitula heterogama aut subdioica, corollis femineis etiam tubulosis. — Flores (nunquam flavi) albi aut purpurascens.

XCI. HOMOGYNE Cass. dict. 21. p. 412. Less. syn. 158. — *Tussilago* sect. 2. DC. fl. fr. — *Tussilaginis* sp. Linn.

Capitulum multiflorum subheterogamum, fl. radii 1-serialibus paucis filiformibus femineis, disci tubulosis 5-fidis hermaphroditis. Recept. nudum. Invol. cylindracei squamæ subuniseriales oblongo-lineares acutæ disco paulò breviores. Styli fl. fem. et herm. conformes! Achænium oblongo-cylindraceum sulcatum glabrum. Pappus pilosus multiserialis. — Herbaræ Europæ montanæ perennes. Folia pleræque radicalia cum flore circ. coetanea petiolata cordato-orbiculata dentata aut angulata, 1-2 ad basin scapi. Scapus simplex 1-cephalus imò basi 1-2-foliosus apice parcè squamosus. Flores purpurascens et variantes albi.

1. *H. SYLVESTRIS* (Cass. l. c.), foliis subglabris reniformibus sontè inciso-7-lobis, lobis intermediis 3-dentatis.  $\frac{1}{2}$  in sylvaticis subalpinis Carinthiæ et Carniolæ. *Tussilago sylvestris* Scop. carn. n. 1057. Jacq. austr. 5. app. t. 12. Willd. sp. 3. p. 1967. Scapus circ. pedalis rarissimè 2-3-cephalus (Bluff et Fing.)

cum petiolis et foliorum marginibus parce subhirsutus. Fol. canlis petioli basi vaginantes. (v. s.)

2. *H. ALPINA* (Cass. l. c.), foliis reniformibus denfato-crenatis subtus ad venas pilosis. 2½ in pratis elatis et humidis inter terminum Fagi et lineam nivalem Alpium, Juraasi, montis Generosi (Com.). *Tussilago alpina* Linn. sp. 1312. Jacq. austr. t. 246. Lam. ill. t. 674. f. 7. Schkuhr handb. t. 242. Gaud. fl. helv. 5. p. 272. Baumg. enum. 3. p. 92. *Petasites alpinus* Scop. carn. Herba circ. semipedalis. Folia scapi a vaginantia, limbo subnullo. (v. s.)
3. *H. DISCOLOR* (Cass. l. c.), foliis cordato-orbiculatis undulato-dentatis supra glabris subtus tomentosis. 2½ in alpihus Austriæ (Jacq.), Tyrolis (Hop!) Salzburgi, Carinthiæ, Carniolis super. (Graaf!), etc. *Tussilago discolor* Jacq. austr. t. 247. Sturm deutschl. fl. ic. *Petasites discolor* Baumg. enum. 3. p. 92. *Tussilago alpina* Scop. carn. — Clus. hist. 2. p. 112 fig. inf. et 113 fig. int. Herba vix semipedalis, fortè *H. alpina* var? (v. s.)

**XCII. NARDOSMIA** Cass. dict. 34. p. 186. Less. syn. 139. — *Tussilaginis* sp. auct.

**Capitula** multiflora subdioica — Submascula, radio uniseriali fœmineo ligulato, disco bisexuali et abortu masculo sterili, cor. tubulosis 5-dentatis. — Subfœminea radio pluriseriali fœmineo minutissimè ligulato, disco pauci (1-5) floro, cor. tubulosis 5-dentatis. Recept. nudum plenum. Invol. squamæ 1-seriales floribus æquales aut breviores. Achaemia teretiuscula glabra. Pappus pilosus, fl. masculorum brevior et minus copiosior quàm fœmineorum. — Herbæ Europææ, Asiaticæ aut Boreali-Americanæ, perennes. Folia cum floribus aut paulò post flores nascentia petiolata dentata basi cordata. Scapus squamosus apice thyrsus fastigiatum gerens. Flores purpurascentes aut albi.

1. *N. FRAGRANS* (Reichenb. fl. exc. 2. p. 289), foliis costaneis basi cordatis orbiculatis cartilagineo-dentatis supra glabresculis subtus in nervis puberis, lobis baseos subrotundis, invol. squamis acutis discum æquantibus aut superantibus. 2½ ad Neapolim (Reyn.), in Sicilia (Guss.), Sardinia (Moris!) et Mauritania (Poir.) in vallibus subhumidis et ad rivulos. Nard. denticulata Cass. dict. 34. p. 187. *Tussilago suaveolens* Desf. cat. *Calcia alliarizifolia* Poir. it. num. 2. p. 236. non Linn. ex Poir. dict. 8. p. 151. *Tussilago racemosa* Tausch syll. soc. ratiab. 2. p. 251? ex Ten. syll. 436. Flores hyberni suaveolentes. Herba ex eo culta sub nom. vernaculo *Héliotrope d'hiver*.

\* *Submascula*, ligulis ovali-oblongis nunc invol. longioribus, nunc rariùs invol. ferè brevioribus. — *Tussilago fragrans* Vill. act. soc. h. nat. 1. p. 72. t. 12. (v. v. et s.)

\*\* *Subfœminea*.... nondum cognita.

2. *N. LÆVIGATA*, foliis costaneis basi cordatis orbiculatis cartilagineo-dentatis supra glabresculis subtus in nervis puberulis, lobis baseos subrotundis, invol. squamis obtusis disco brevioribus. 2½ in Sibiria et fortè in Bohemia.

\* *Submascula*, ligulis ovali-oblongis involuero paulò longioribus. *Tussilago* Gmel. fl. sib. 2. t. 69? et ideò *Tussil. lævigata* Willd. sp. 3. p. 1969. Ledeb. fl. alt. 47. p. 54. *Tussil. Bohemica* Hoppe tasch. 1803. p. 60. *Nardosmia straminea* Cass. dict. 34. p. 189. *Petasites lævigatus* Reich. fl. exc. 2. p. 279 hinc essent revocand. etiamsi flores flavescences (forsan ex sicco) dicti! Plantam non vidi.

\*\* *Subfœminea*, lig. fl. fœm. minutissimis. — *Tussilago saxatilis* Turcz. l. in litt. Flores ex eo sordidè rubri. Folia ferè omnino *N. fragrantis*. An discrimen ex invol. petiti ad speciem aut ad sexum pertinet? (v. s. coll. à cl. Turczaninoff in subalpinis montium Kultokensiam et in alpe Chamor.)

3. *N. FRIGIDA* (Hook. fl. bor. am. 1. p. 307. var. a.), foliis cordatis inæqualiter grossè et obtusè dentatis sublobatis supra glabris subtus albo-tomentosis, lobis



bases divergentibus.  $\frac{1}{2}$  in vallibus et montanis alpinis Dahuricis (Turcz.), Sibiricis (Gmel.), Norwegicis (Chr. Smith!), Lapponicis (Linn.), Americis maximè borealis (Pursh). — *Nardosmia angulosa* Cass. dict. 34. p. 188.

\* *Submascula*, ligulis ovali-oblongis stylo longioribus. — *Tussil. frigida* OED. fl. dan. t. 61. — Gmel. sib. 2. p. 150. t. 70. (v. a.)

\*\* *Subfeminea*, ligulis minutissimis, caule et thyrsi duplè longioribus. (v. s. coll. ad salinas Irutianas à cl. Turczaninoff.)

4. *N. CORYMBOSA* (Hook. fl. bor. am. 1. p. 307), foliis cordatis sinuatis et acutè denticulatis oblongis acutis supra glabris subtus tomentosis, lobis hancos divergentibus.  $\frac{1}{2}$  in insulâ Melville (Br.) ad flum. Mackenzie (Hook.). *Tussilago corymbosa* Brown chlor. melv. p. 21. Affinis *N. frigida*.

\* *Submascula*. — adhuc ignota.

\*\* *Subfeminea*, ligulis linearibus minutissimis. — Sola hucusquè descr. *Corymbus pauciflorus*. Invol. squamæ acutæ floribus duplè breviores. (v. s. comm. à cl. Sabine.)

5. *N. SAGITTATA* (Hook. fl. bor. am. 1. p. 307), foliis oblongis acutis sagittatis integerrimis, lobis obtusis.  $\frac{1}{2}$  ad sinum Hudsonis (h. Banks), ad rapem scopulosos, à lacu superiori ad arcidulam Franklini (Hook.). *Tussilago sagittata* Pursh fl. bor. am. 2. p. 531. Thyrsus dicitur ovatus fastigiatus capitulis radiatis. Videtur specim. *submasculum*.

6. *N. PALMATA* (Hook. fl. bor. am. 1. p. 308), foliis subtus tomentosis palmatis-partitis, lobis oblongis grossè pauci-dentatis.  $\frac{1}{2}$  in America bor. ad Labrador et Terram-Novam (Ait.) à lacu Huron ad lacum Bear, in rupibus scopulosis dictis, et ad ostium flum. Columbia (Hook.). *Tussilago palmata* Ait. her. 3. p. 188. t. 11. Specim. *submasculum*. Thyrsus fastigiatus 7-9-cephalus. Ligula oblongæ stylo longiores.

XCIH. PETASITES *Tourn. inst. 451. t. 258. Gært. fr. 2. p. 408. t. 166. Oen. dict. 39. p. 199.* — *Tussilago* sect. *Petasites* DC. fl. fr. Gend. fl. heb. *Tussilaginis* sp. Linn.

Capitula multiflora subdioica. — *Submascula* nempe radio 1-seriali pauci- (1-5-) floro fœmineo, corol. tubulosâ gracili, ovario ovulato, disco bisexuali sed ovario exovulato masculo, cor. tubulosâ fauce subdilatatâ regulariter 5-dentatâ. — *Subfœminea* nempe radio multi-seriali fœmineo fertili, cor. filiformi subtruncatâ, stylo exserto, disco pauci- (1-5-) floro masculo, cor. fauce dilatâtâ regul. 5-dentatâ. Recept. nudum planum. Invol. squamæ subuniseriales floribus breviores bracteolis sæpè basi stipatæ. Achænia cylindracea glabra. Pappus pilosus, marium multò minus copiosior quàm fœmineorum. — Herbes Europæ perennes in humidis crescentes. Folia post flores orta, radicalia petiolata reniformia aut cordata dentata ampla. Scapus sæpius tomentosus squamis membranaceis glabriusculis onustus. Thyrsus polycephalus terminalis. Pedicelli simplices 1-cephali aut in subfœmineis interdum ramosi. Flores purpurascens aut albi.

1. *P. VULGARIS* (Desf. fl. atl. 2. p. 270), foliis profundè cordatis emarginatis inæqualiter dentatis suprâ demum glabris subtus pubescentibus, lobis hancos approximatis. —  $\frac{1}{2}$  in Europæ humidis frequens, etiam Algeriis (Desf.). *Tussilago Petasites* Hopp. tæsch. 1803. p. 35. Willd. sp. 3. p. 1971. DC. fl. fr. n. 316. *Petasites officinalis* Moench.

\* *Submasculus*, thyrsi ovati pedicellis simplicibus. *Tussilago Petasites* Linn. sp. 1215. *Tuss. Petasites hermaphrodita* Poir. Gend. — Lam. ill. t. 64 f. 1. 2. Bull. herb. t. 391. fl. dan. t. 842. (v. v.)

\*\* *Subfœmineus*, thyrsi elongati pedicellis plerique ramosis. *Tussilago* h.

*brida* Linn. sp. 1214. Sow. engl. bot. t. 430. Schkuhr handb. t. 242. Dill. eth. t. 230. (v. v.)

2. *P. ALBUS* (Gertn. fr. 2. p. 406. t. 166), foliis latè cordatis suborbiculatis angulosis argutè dentatis suprà demùm glabratīs subtiſ pubeſcenti-villoſis albidis, nerviſ ſcabriſ, lobis baſeoſ ſubrotundis.  $\frac{1}{2}$  in monteiſ Europæ non raruſ, etiam Algeriæ (Deſf.). *T. alba* Linn. sp. 1214. DC. fl. fr. n. 3166. An hæc aut ad priorem refer. *Petasites glabrescens* Hoppe?

\* *Submaſculus*, thyrſi ſaſtigiatī pedicelliſ ſimplicibuſ. — Fl. dan. t. 524. Hoppe! in Sturm deutſchl. fl. ic. (v. v.)

\*\* *Subſamineuſ*, thyrſi oblongi pedicelliſ pleriſque ramoſiſ. Tuſſa. ramoſa Hoppe! cent. Tuſſa. alba  $\beta$  Vill. act. ſoc. h. nat. par. 1. p. 73. (v. v.)

3. *P. NIVUſ* (Caſſ. dict. 39. p. 202), foliſ latè cordatiſ ſubtriangularibuſ den-tatiſ ſuprà glabriſcuſcuſ ſubtiſ niveo-tomentoiſ, nerviſ concoloribuſ, lobis diſſaricatiſ ſubacutiſ.  $\frac{1}{2}$  in montaniſ Alpiuſ et Juraſi priore rarior; etiam in Roſſia merid. ex ſpecim. maſculo ſub nom. *T. hybridæ*? ab hon. Ac. Sc. Petrop. miſſo. Tuſſilago nivea Vill. act. ſoc. h. nat. Par. 1. p. 73. Willd. ſp. 3. p. 1970. DC. fl. fr. n. 3167. Tuſſilago frigida Vill. dauph. 3. p. 175. Sut. fl. helv. 2. p. 180 non Linn. Pedicelliſ utriuſque ſexuſ ſimpliciſ. Lobi baſeoſ fol. æſpè ſiſſiſ.

\* *Submaſculus*, thyrſi oblongi pedicelliſ brevioribuſ. — Hoppe! in Sturm deutſchl. fl. ic. — Moris. ox. 2. 7. t. 10. f. 4. (v. v.)

\*\* *Subſamineuſ*, thyrſi oblongi pedicelliſ elongatiſ. — Tuſſilago paradoxa Retz obſ. 2. p. 24. t. 3. Hoppe! in Sturm fl. ic. Tuſſil. frigida Sut. fl. helv. 2. p. 180 non Linn. (v. v.)

4. *P. TOMENTOSUſ*, foliſ profundè cordatiſ ovatiſ orbiculatiſve vix dentatiſ ſuprà glabriſcuſcuſ ſubtiſ denſè albo-tomentoiſ, nerviſ concoloribuſ, lobis baſeoſ approximatī ſublobatiſ ſubrotundis.  $\frac{1}{2}$  in Germaniæ et Scaniæ ad ripæ flum. rarior. Tuſſilago ſpuria Willd. ſp. 3. p. 1972? Tuſſa. tomentosa Ehr. beitr. 3. p. 65. Hoppe! taſch. 1803. p. 61. *Petasites ſpuriaſ* Reichenb. fl. exc. 2. p. 279. Pedicelliſ utriuſque ſexuſ ſimpliciſ.

\* *Submaſculus*, thyrſo coarctato, pedicelliſ brevioribuſ.

\*\* *Subſamineuſ*, thyrſo elongato. — Tuſſa. ſpuria. Retz obſ. 1. p. 29. t. 2? (v. s.)

XCIV. *ADENOCAULON* Hook. bot. miſc. 1. p. 19. t. 15. Less. in Linnæ 1831. p. 107. ſyn. 159.

Capitulum 9-10-florum heterogamum, floribuſ tubuloſiſ 4-5-dentatiſ, radiiſ quiniſ 1-ſerialibuſ ſcæmineiſ, diſci 4-5 maſcuſiſ. Invol. uniſerialia demùm reflexum pleiophyllum. Recept. nudum. Achæmia oblongo-obovata glanduliſ ſtipatiſ ſupernè inſtructa. Pappuſ nulluſ. — Herbe in Americâ occidentali habitanteſ perenneſ. Cauleſ albo-tomentoiſ ſupernè glanduliſ pedicellatiſ inſtructi, infernè folioſi, ſupernè nudi laxè paniculati. Folia alterna penninervia ſuprà glabra ſubtiſ tomentosa penninervia petiolata.

1. *A. BECOLOR* (Hook. bot. miſc. 1. p. 19. t. 15. fl. bor. am. 1. p. 308), foliſ ſubtriſlobiſ angulato-dentatiſ.  $\frac{1}{2}$  in ſylviſ denſiſ ad oram occidentalem Ame-ricæ borealiſ prope fretum de Fuca, Fort Vancouver et in Rocky-mountainaſ.
2. *A. CHILENSE* (Less. in Linnæ 1831. p. 107), foliſ ſubrotundo-ellipticiſ cordatiſ integerrimiſ aut ſubintegerrimiſ.  $\frac{1}{2}$  in Chili.

## DIV. II. EUTUSSILAGINÆ.

Capitula heterogama radiata, liguliſ ſcæmineiſ, floribuſ diſci hermaphroditī. — Floreſ ſemper lutei, radii etiam lutei aut rariſſimè roſci.

XCV. TUSSILAGO *Tourn. inst.* 487. t. 276. *Adans. fam.* 2. p. 124. *Gærtn. fr.* 2. t. 170. *Cass. dict.* 34. p. 190 (*excl. spec.*). *Less. syn.* 159. — *Tussilago* sect. *Farfara* DC. *fl. fr.* — *Tussilaginis* sp. *Linn.*

Capitulum multiflorum heterogamum, fl. radii pluriserialibus angustissimè ligulatis fœmineis, disci paucis masculis tubulosis limbo campanulato 5-dentato. Recept. nudum. Invol. subuniseriale, squamis oblongis obtusis. Antheræ subcaudatæ. Stylus disci inclusus abortivus, radii bifidus, ramis teretiusculis. Achænium radii oblongo-cylindraceum glabrum, disci abortivum. Pappus radii pluriserialis, disci uniserialis, setis tenuissimis. — Herba perennis Gerontogea. Rhizoma crassiusculum. Folia post anthesin nascentia petiolata cordata angulata. Scapi 1-cephali tomentosi aphylli sed squamis alternis oblongis adpressis inv. squamas referentibus onusti. Flores flavi.

1. T. FARFARA (*Linn. sp.* 1214). 2. in argillosis petrosis humidis Europæ totius, Byzantii (d'Urv.), Tauriæ (Stev.), Persiæ (Belang.) et in montibus Sibirie (Patr.), Indiæ orient. (Royle!). *OEd. fl. dan.* t. 595. *Sow. engl. bot.* 6. t. 429. *Sturm fl. germ. ic.* T. vulgaris *Lam. fl. fr.* 2. p. 71. T. rupestris *Vall. cat. n.* 2990. comp. n. 100. — Legi ad littora maris et in alpihus ad altit. hexapod. 1300! Variat scapo digitali et pedali, solitario aut multiplici, foliis sævè cum floribus aut post eos natis, membranaceis aut rigidulis, utrinquè subglabris aut subius araneosis densève tomentosis, basi cordatis aut ovatis, angulatis set subindivisis, plus minùs dentatis. Has variationes video et in specim. Europæis et in Asiaticis. (v. v. et s.)

N. B. Quid *Tussilago bifolia* Lavy veg. orig. p. 13?

† *Species exclusæ.*

- T. ALBA L. = *Petasites albus*.
- T. ALBICANS Sw. = *Leria albicans*.
- T. ALPINA L. = *Homogyne alpina*.
- T. ALPINA Scop. = *Homogyne discolor*.
- T. ANANDRIA L. = *Anandria bellidiastrum*.
- T. ANANDRIA Willd. = *Anandria bellidiastrum* β.
- T. BOHEMICA Hopp. = *Nardosmia saxatilis* α.
- T. CACALIA Scop. = *Adenostyles glabra*.
- T. DENTATA L. = *Chaptalia dentata*.
- T. DISCOLOR Jacq. = *Homogyne discolor*.
- T. RESCAPA Pers. = *Loxodon brevipes*.
- T. FRAGRANS Vill. = *Nardosmia fragrans*.
- T. FRIGIDA L. = *Nardosmia frigida*.
- T. HYBRIDA L. = *Petasites vulgaris*. \*\*
- T. JAPONICA L. = *Senecio Kämpferi*.
- T. INTEGRIFOLIA Michx. = *Chaptalia tomentosa*.
- T. LAEVIGATA Willd. = *Nardosmia saxatilis*.
- T. LYRATA Willd. = *Anandria bellidiastrum*.
- T. LYRATA Pers. = *Leria nutans*.
- T. NIVA Hopp. = *Petasites nivens*.
- T. NUTANS L. = *Leria nutans*.
- T. ODORATA Vill. = *Adenostyles glabra*.

- T. *PALMATA* Ait. = *Nardosmia palmata*.  
 T. *PARADOXA* Retz. = *Petasites niveus*. \*\*  
 T. *PARADOXA* Roth. = *Petasites tomentosus*.  
 T. *PETASITES* L. = *Petasites vulgaris*.  
 T. *PILOSELLOIDES* Pers. = *Liberkuhnia piloselloides*.  
 T. *PUMILA* Sw. = *Leria pumila*.  
 T. *RACHIOSA* Tausch. = ? *Nardosmia fragrans*.  
 T. *RAMOSA* Hop. = *Petasites albus*. \*\*  
 T. *RUBRA* Link. = *Senecio cruentus*.  
 T. *SAGITTATA* Pursh. = *Nardosmia sagittata*.  
 T. *SARMENTOSA* Pet. Th. = *Chevreulia stolonifera*.  
 T. *SAXATILIS* Turcz. = *Nardosmia saxatilis*. \*\*  
 T. *SENUATA* Pers. = *Leria pumila*.  
 T. *SPURIA* Willd. = *Petasites tomentosus*.  
 T. *SPURIA* Schrank. = *Petasites niveus*.  
 T. *SUAVEOLENS* Desf. = *Nardosmia fragrans*.  
 T. *SILVESTRI* Jacq. = *Homogyne sylvestris*.  
 T. *TOMENTOSA* Hop. = *Petasites tomentosus*.  
 T. *TRIFURCATA* Forst. = *Senecio trifurcatus*.

XCVI. *CELMISIA* Cass. dict. 37. p. 259 (non 7. p. 357). Gaudich. voy. 470.

Capitulum multiflorum, fl. radii ligulatis fœmineis 1-seriatis, disci tubulosi hermaphr. 5-dentatis. Recept. latum epaleaceum nudum aut alveolare. Invol. squamæ pluriserialis inæquales acutæ disco paulo longiores. Ligulæ stam. rudimentis destitutæ stylum longum ramis obtusis dorso subpapillois gerentes. Antheræ ecaudatæ appendiculatæ. Achænium oblongo-cylindraceum basi attenuatum. Pappus pluriserialis, paleis filiformibus inæqualibus barbellatis. — Herbæ Australasiæ, caule erecto simplici nudo 1-cephalo. Folia radicalia oblonga. Capitulum disco luteo, radio roseo? Contra regulas prioritatis strictas nomen huic nec sequenti generi servavi, quia fuit à cl. Gaudichaud sub hoc nomine sancitum.

1. *C. LONGIFOLIA* (Cass. dict. 37. p. 256), foliis sessilibus linearibus argenteo-lanatis longissimis. ? in Novâ-Hollandiâ orientali ad vallem Jamieson (Gaud.!), in ericetis humidis et excelsis paludibus montium cœruleorum (Cunn.). Gaudich. ! voy. Freye. bot. p. 470. t. 91. Ovarium sessile glabrum. Flosc. ligulati staminibus omnino orhati. Invol. disco longius. Recept. alveolatum. (v. s.)
2. *C. SPATULATA* (A. Cunn. ! in litt.), foliis elliptico-oblongis in petiolum longè attenuatis utrinquè glabris. ? in Novâ-Hollandiæ collibus siccis ad Oyster Harbour prope Regis Georgii sinum legit cl. A. Cunningham. Scapus basi glaber nudus, apice pubescens squamellas lineares paucas gerens, 12-14 lin. longus. Folia petiolo incluso 2-3 poll. longa, 5-6 lin. lata. Invol. basi extus lanatum, squamis acuminatis apice subinduratis. (v. s.)

XCVII. *ALCIOPE*. — *Celmisia* Cass. dict. v. 7. p. 357 (non vol. 37). — *Ligulariæ* sp. Less. — *Arniciæ* sp. Thunb.

Capitulum multiflorum, fl. radii 1-serialibus ligulatis fœmineis sed stam. filamenta subabortiva gerentibus, disci tubulosi 5-dentatis hermaphroditis. Recept. latum epaleaceum. Invol. squamæ pluriserialis subimbricatæ adpressæ. Styli rami breves obtusi parçè papillosi glabri,

in radio et disco sat conformes. Achenium oblongo-cylindraceum subobcompressum. Pappus pluriserialis, setis scabridis imâ basi breviter concretis. — Herbarum suffrutescentes Capenses erectæ ramosæ. Caules teretes tomento denso lanato canescentes. Folia alterna petiolata integra aut denticulata suprâ glabra subtus cum petiolis cano-lanata. Capitula ramos terminantia. Invol. cano-lanatum. Flores flavi. — Genus hinc ad Ligulariam ex ligulis stam. rudimentaria gerentibus, illinc ad Celmisiam ex structurâ stigmatis, accedens. Dixi ex Alciope nymphâ mythologicâ Celmisii matre, alludens ad hoc quod genus primum Celmisia dictum, alterum quasi peperit.

1. *A. TABULARIS*, caule basi fruticoso erecto ramoso, ramis tomentosis elongatis apice 1-cephalis, foliis breviter petiolatis ovalibus obovatisve basi subattenuatis integris aut denticulatis suprâ glabris subtus lanato-tomentosis, invol. squamis oblongo-linearibus subaraneosis, exter. paucis interiores frâ squamulis. § ad Cap. B.-Spei in monte Tabulari (Th. & Dreg. n. 173). *Arnica Tabularis* Thunb. fl. cap. 668. *Celmisia rotundifolia* Cass. dict. 7. p. 357. *Ligularia Tabularis* Less. syn. 390. Folia 1-1  $\frac{1}{2}$  poll. longa, 9-10 lin. lata. (v. s.)

2. *A. LANATA*, caule frutescente erecto subimplici densissime lanato, ramis paucis apice 1-cephalis bracteolis lanatis obtusis, foliis in parte caulis inferioris petiolatis cordato-ovatis obtusis margine subrevolutis coriaceis subtus lanatis, adultis suprâ glabris, invol. 2-3-seriali imbricato dense lanato. § ad Cap. Bonæ-Spei juxta Rode Sand in montibus prope Winterhoek (Th.), in distr. Worcester et Tulbagh (Eckl. n. 1421. 830. 1947). *Arnica lanata* Thunb. fl. cap. 667. *Coryza tomentosa* Burm. prod. 26. *Ligularia lanata* Less. syn. 390. (v. s.)

ß. *grandis*, caule basi frutescente erecto parce ramoso densissime lanato, ramis apice 1-cephalis bracteolis lanatis obtusis, foliis petiolatis ovatis (basi cordatis), margine planis obtusis submembranaceis suprâ glabris subtus dense lanatis, invol. 2-3-seriali imbricato dense lanato. § ad Cap. Bonæ-Spei (Th.), in monte prope Tulbagh (Dreg. n. 1975). *Arnica grandis* Thunb. fl. cap. 668. E. Mey. in Dreg. coll. (v. s.)

XCVIII. BRACHYGLOTTIS Forst. car. gen. n. 46. Cass. dict. 48. p. 464. — Senecionia sp. Less. — Cineraria sp. Willd.

Capitulum pluri- (9-10-) florum heterogamum, fl. radii 1-serialibus paucis fœmineis brevissime ligulatis aut oblique tubulosis disco fere brevioribus, disci tubulosis 5-dentatis hermaphr. Recept. nudum. Invol. oblongo-cylindricum squamis linearibus 1-serialibus extus caliculato-bracteolatum. Styli radii rami exserti revoluti apice clavellati obtusi; disci stylus inclusus subindivisus forte abortivus. Achenia oblonga basi non attenuata. Pappus 1-serialis, setis densis scabris basi in anulum concretis. — Arbores Australasicæ. Folia alterna ovata aut subrotunda petiolata subtus tomentosa. Corymbus paniculatus.

1. *B. REPANDA* (Forst. l. c.), foliis ovatis repando-sinuatis, paniculâ decompositâ divaricatâ. § in Novâ-Zelandiâ (Forst.), frequens in sylvis densis ad littora fluminum ad Kowa-Kowa, etc. (A. Cunn l.). *Cineraria repanda* Willd. sp. 3. p. 2076. non Lour. Arbuscula 12-15-pedalis. Folia petiolo 2-3  $\frac{1}{2}$  poll. instructa subrotunda suprâ glabra subtus adpressè cano-tomentosa penninerviâ rigidula, diam. 5-6 pollic. Panicula amplissima. (v. s. comm. à cl. Cunningh.)

2. *B. ROTUNDIFOLIA* (Forst. l. c.), foliis ovato-subrotundis integerrimis, paniculâ pauciflorâ. § in Novâ-Zelandiâ. Cass. dict. 48. p. 464 descr. Forst. ined. *Cineraria rotundifolia* Willd. sp. 3. p. 2076.

† *Species dubia hic ex patria adnexa.*

3. B. ? FORSTERI, foliis subrotundis angulato-dentatis basi cuneatis subtus lanatis, petiolis alatis basi nudis, corymbo terminali. ♂ in Novâ-Caledoniâ (Spr. pag.), Novis-Hbridis (Spr. syst.). Cineraria Forsteri Spreng. pl. min. eegu. pag. 1. p. 58. syst. 3. p. 545.
4. B. ? CALEDONIÆ, foliis petiolatis ovato-oblongis sinuato-pinnatifidis inæqualiter denticulatis subtus pallidis obsolete lanatis, petiolis amplexicaulibus subulatis, corymbis terminalibus bracteatis. ♂ in Novâ-Caledoniâ. Cineraria Caledoniæ Spreng. syst. 8. p. 547.

Trib. III. *ASTEROIDEÆ* Less. syn. 161. — Asteræ, Inuleæ pleræque et Helianthæ paucæ Cass.

Capitula sæpius heterogama, rariùs homogama aut dioica. Stylus fl. hermaphr. supernè cylindraceus bifidus, ramis longiusculis linearibus extus subplanis sæpius subacuminatis rariùs obtusis, extus æqualiter et minutè puberulis. Stigmatis series angustæ prominulæ usque ad originem pubis externæ extensæ. Corolla pellucida, staminigera tubulosa regulariter dentata. Pollen globosum echinulatum.

## CONSPECTUS TRIBUS.

Subtrib. I. *ASTERINEÆ*. Capitula homo- aut heterogama, sæpius radiata. Anthere ecaudatæ. Folia ferè omnium alterna.

Div. 1. *AMELLEÆ*. Recept. paleaceum. (XCIX-CII.)

Subdiv. 1. *EUAMELLEÆ*. Capitula heterogama radiata omnia recept. paleaceo donata. (XCIX-CI.)

Subdiv. 2. *HETEROETHALAMEÆ*. Capitula dioica, mascula recept. nudo, feminea recept. paleaceo. (CII.)

Div. II. *ASTEREÆ* (1). Recept. epaleaceum. Capitula radiata ligulis heterochromis. (CIII-CLXIII.)

Subdiv. 1. *EUASTEREÆ*. Ligulæ 1-seriales. Pappus pilosus in disco et radio conformis, 1-serialis aut pluriserialis, setis æqualibus. (CIII-CXXIV.)

Subdiv. 2. *DIPLOPAPPEÆ*. Ligulæ 1-seriales. Pappus in disco et radio conformis, biserialis, serie exteriori brevi subpaleacæ, interiori setosâ elongatâ. (CXXV-CXXXI.)

Subdiv. 3. *ERIGEREÆ*. Ligulæ pluriseriales. Pappus pilosus conformis aut inæqualis. (CXXXII-CXLI.)

---

(1) In hac divisione totâ assiduum secutus sum et auream Asterearum enumerationem à cl. Nees (Genera et species Asterearum 8°. Vratisl. 1832) propositam, et observationes sagacissimas manuscriptas à cl. Lindley (conf. Bot. reg. t. 1527) missas. Utrique gratias ex imo pectore perplurimas ago, ob adnotationum speciminumque amicissimas communicationes.

- Subdiv. 4. *HETEROPAPPÆ*. Ligulæ 1-seriales. Pappus diversus in disco et radio. (CXLII-CLI.)
- Subdiv. 5. *BELLIDÆ*. Ligulæ 1-seriales. Pappus paleaceus plurisquamellatus. (CLII-CLIV.)
- Subdiv. 6. *BELLIDÆ*. Ligulæ uni- aut pluriseriales. Pappus nullus aut coroniformis. (CLV-CLXIII.)
- Div. III. *CHYSOCOMEÆ*. Recept. epaleaceum. Capitula lutea, nunc homogama floribus omnibus hermaphroditis, nunc radiata, ligulis cum disco homochronis (1). (CLXIV-CXCIII.)
- Subdiv. 1. *GYMNOSPERMÆ*. Pappus nullus. Ligulæ 1-seriales. (CLXIV-CLXVII.)
- Subdiv. 2. *ACHYRIDÆ*. Pappus paleaceus. Ligulæ 1-seriales. (CLXVIII-CLXXI.)
- Subdiv. 3. *HETEROTHECÆ*. Pappus in disco et radio diversus. Ligulæ 1-seriales. (CLXXII.)
- Subdiv. 4. *PSIADÆ*. Ligulæ pluriseriales. Pappus pilosus in disco et radio conformis. (CLXXIII-CLXXIX.)
- Subdiv. 5. *CHRYSOPSIDÆ*. Ligulæ 1-3-seriales aut nullæ. Pappus biserialis, ext. brevis subpaleaceus, inter. pilosus. (CLXXX-CLXXXII.)
- Subdiv. 6. *SOLIDAGINÆ*. Ligulæ uniseriales aut nullæ. Pappus pilosus in disco et radio conformis, 1-serialis aut pluriserialis, setis æqualibus. (CLXXXIII-CXCIII.)
- Div. IV. *SOLENOGYNÆ*. Capitula discoidea, floribus omnibus tubulosis, exterioribus fœmineis. (CXCIV-CXCVIII.)
- Subtrib. II. *BACCHARIDÆ*. Capitula heterogama aut dioica nunquam radiata. Cor. omnes tubulosæ, fœmineis sæpius in ambitu multiseriales. Antheræ ecaudatæ. Recept. epaleaceum. Folia alterna.
- Div. 1. *CONYZÆ*. Capitula heterogama monoica.
- Subdiv. 1. *SPHÆRANTHÆ*. Capitula in capitulum sphaericum congesta ad bractearum axillas sessilia.
- Subdiv. 2. *GRANGEINÆ*. Capitula in glomerulum non congesta. Pappus nullus aut subnullus.
- Subdiv. 3. *EUCONYZÆ*. Capitula in glomerulum non congesta. Pappus pilosus.
- Div. II. *BACCHARÆ*. Capitula dioica.

---

(1) Si quis autumat me, in divellendis Astereis et Chrysocomeis, colori nimium credidisse, errorum facile corrigat. Etenim in serie homochromâ *Gymnospermæ* representant *Bellidæ*, *Achyridæ* *Bellicas*, *Heterothecæ* *Heteropappæ*, *Psidiæ* *Erigeronæ*, *Chrysopsideæ* *Diplopappæ*, *Solidagineæ* *Euasteronæ*. Divisio nostra à colorum classicâ diversitate petita multo magis cum affinitatibus veris convenit, quam ulla alia mihi cognita.

Subtrib. III. TARCHONANTHEÆ. Capitula aut dioica aut heterogama nunquàm radiata, flor. ambitus fœmineis pluriserialibus tenuissimis, disci ♂ aut ♂ paucioribus majoribus. Anthæræ caudatæ. — Folia alterna.

Div. I. EUTARCHONANTHEÆ. Capitula dioica.

Div. II. PLUCHEINÆ. Capitula heterogama seu monoica.

Subtrib. IV. INULEÆ. Capitula nunquàm dioica, rariùs homogama discoidea, sæpissimè heterogama radiata, ligulis isochromis fœmineis. Recept. (exc. Rhanterio) epaleaceum. Anthæræ caudatæ. — Folia alterna.

Div. I. EUINULEÆ. Capitula pluri- aut multiflora in glomerulum non concreta.

Div. II. CÆSULIÆ. Capitula 1-flora in glomerulum adgregata.

Subtrib. V. BUPHTHALMEÆ. Capitula heterogama (aut rarissimè homogama). Flor. radii fœminei ligulati rariùs tubulosi. Anthæræ longè aut brevissimè caudatæ aut ecaudatæ. Recept. paleaceum. Pappus coroniformis, lacerus aut piloso-dentatus. — Folia alterna.

Subtrib. VI. ECLYPTEÆ. Capitula heterogama, fl. radii fœmineis (excl. Cryphiospermo) ligulatis, disci hermaphroditis. Anthæræ ecaudatæ. Recept. paleaceum. Pappus nullus seu aristatus nunquàm setosus. — Folia opposita.

Subtrib. I. ASTERINÆ *Nees ast. p. 3. excl. gen. pauc. — Asteræ Less. syn. 161 non 162.*

Capitula (exc. Heterothalamo) nunquàm dioica sæpiùs radiata. Recept. sæpiùs epaleaceum. Anthæræ ecaudatæ. — Folia ferè omnium alterna.

#### Div. I. AMELLEÆ.

Recept. paleaceum. — Capitula radiata heterogama aut in Heterothalamo dioica.

Subdiv. I. EUAMELLÆ. — Amelloidæ *Less. syn. 161.*

Capitula heterogama radiata.

XCIX. AMELLUS *Cass. dict. 8. p. 577. 16. p. 210. 37. p. 463 et 489. Nees ast. p. 241. Less. syn. 161. — Amelli sp. Linn. Willd.*

Capitulum multiflorum sæpiùs heterogamum, fl. radii ligulatis sub-uniserialibus fœmineo-fertilibus, disci hermaphr. tubulosis 5-dentatis, rariùs homogamum discoideum. Recept. subconicum paleis confertis inter flores onustum. Invol. ovati squamæ pluriseriales imbricatæ rigidæ acuminatæ, inter. sensim in paleas abeuntes. Achænia cuneiformia compressa, radii subtetragona scabra, disci lævia apice subdentata. Pappus radii paleis brevibus 1-serialibus, disci duplex, exter. paleaceus pappo radii similes, interior 4-setus scaber. — Fruticuli vel herbæ Cœpenses. Folia infer. opposita, super. alterna oblonga, integerrima aut parcè dentata strigilosa aut canescentia. Pedunculi terminales sub-



squamulosi 1-caphali. Capitula radio cœruleo, disco lateo. Cor. disci extus pubescentes.

\* *Fruticosi.*

1. *A. LYCHNITIS* (Linn. sp. 1276. mant. 476), caule fruticoso ramoso subdiffuso, foliis oppositis alternisve spathulato-lanceolatis obtusis integerrimis utrisque ramisque strigis minimis sericeo-canescens. 5 ad Cap. Bonæ-Spei ad Riet-Valley (Ecklon !), etc. Lam. ill. t. 682. f. 1. Jacq. coll. 5. t. 10. f. 1. Verbesina asteroides Linn. spec. ed. 1. p. 902. Buphthalmum Linn. hort. cliff. p. 415. — Bœyn. prod. 3. t. 15. f. 2. ex Willd. (v. v. c. et s. sp.)
2. *A. TENUISOLUS* (Burm. prod. 28. Lam. dict. 1. p. 129), caule fruticoso erecto ramoso, ramis tenuissimè pubescentibus, foliis alternis linearibus subacutis integerrimis involucriisque tenuissimè hirtello-pubescentibus. 5 ad Cap. Bonæ-Spei (Burm. Eckl. ! in h. Dunant), ad Riebeck kastel (Drege !). *A. lychnitis* E. Mey. ! in Drege coll. Frutex pedalis. Folia 1 ½ poll. longa, lineam lata. Capitula omnino *A. Lychnitis*. (v. s.)
3. *A. HISPIDUS*, caule basi suffruticoso ramoso erecto setis patentibus hispidis, foliis alternis lato-linearibus setuloso-asperis aliis integerrimis dente (imo lobulo) utrinque 1-2-notatis, invol. squamis dorso strigillosis apice acuminatis paleas referentibus. 5 in Africa Capensi ad Camisbergen alt. 3-400, et ad Zw. Kous et Gariep alt. 1-200 ped. legit cl. Drege ! Differt ab *A. anisato* Cass. ob paleam adpressam inter setas nullam et caulem fruticosum. (v. s.)  
*β. angustissimus*, pilis patulis confertis aliis brevibus aliis longis setosis, foliis angustissimis omnibus ferè integerrimis. 5 ad Olifant rivier legit cl. Drege. (v. s.)
4. *A. FLOSCULOSUS*, caule fruticoso ramosissimo, ramis folisque pube brevi subvillosis, foliis linearibus integerrimis subconvolutis vix acutis, capitalis terminalibus solitariis subcorymbosis, invol. squamis dorso puberulis acuminatis, ligulis nullis. 5 in Africa austr. ad Klein Namaqualand, Zwarte Kous et Gariep legit cl. Drege ! Similis *A. hispidus* sed discoideus et ex omni parte potius pubescens quàm hispidus. (v. s.)
5. *A. SCABRIDUS*, caule fruticuloso ramoso sparso et subadpresso setuloso, foliis linearibus acutis integerrimis setis subadpressis rigidis sparsis utrinque et ad marginem scabris, invol. squamis dorso hirtello-scabris acuminatis. 5 in distr. Zwellendam legit cl. Burchell (n. 6815), in deserto Karro (Eckl. ! Zeyh.). Caulis rigidulus tenuis. Rami non striati. Folia 7 lin. longa vix lin. lata. Capitula trita anisum spirantia, ferè dimidio minora quàm *A. Lychnitis*. Ligulae pallide. Achenia pleraque compressa margine callosa demum ferè calva. (v. s.)

\*\* *Herbacei annui.*

6. *A. ANISATUS* (Cass. dict. 37. p. 489), caule herbaceo erecto simplici pube minimè adpressa setisque longis articulatis sparsis instructo, foliis oppositis alternisve linearibus integerrimis utrinque pubescenti-scabris. ① ad Cap. Bonæ-Spei. Nees ast. p. 243.
7. *A. STRIGOUS* (Less. in Linnaea 1831. p. 110. syn. p. 162), caule herbaceo adscendente flexuoso debili breviter et patentem piloso, foliis linearibus lanceolatis basi attenuatis integerrimis hirsutis. ① ad Cap. Bonæ-Spei in arenosis campis, Groene Kloof Swartlandia (Thunb.), in regione Trans-Garipiana (Burch. ! cat. geogr. n. 2276), in Karro prope Gauris rivier (Eckl. ! et Zeyh.). Aster strigosus Thunb. fl. cap. 687. (v. s.)
8. *A. COLORATUS*, radice subsimplici, caulibus pluribus herbaceis teretibus pubescentibus, centrali erecto, lateralibus adscendentibus, foliis linearibus integerrimis adpressè pubescentibus, pedunculis sub capitulo incrassatis cavis, invol. squamis adpressis pubescentibus. ① in Africa australi regione Garipiana legit cl. Drege ! Caules 6-8 pollic. Folia 1-1 ½ poll. longa, lineam lata. Ligulae truncatæ cœruleæ. Recept. paleas plerumque flores disci superantes. (v. s.)

9. *A. RANUS*, caule herbaceo à basi ramosissimo caespitoso foliisque confertè et brevè puberulis, foliis elongato-linearì-lanceolatis integerrimis basi attenuatis, capitulis brevissimè pedicellatis foliis 1-2 stipatis, invol. squamis linearibus subtorulosis, ligulis amplis obtusis involucri duplò longioribus, rec. paleis paucis obtusis. (1) in Africæ Capensis regione Carlepinà legit cl. Drege! Radix simplex perpendicularis. Herba 3 poll. Folia infer. 1  $\frac{1}{2}$  poll. longa, 2 lin. lata. Pappus duplex, ext. per anthesin vix perspicuus paleis ovatis, inter actis scabris deciduis. (v. s.)
10. *A. ANNUUS* (Willd. sp. 3. p. 2215), caule herbaceo subadscendente ramoso patentem et mollior pilosinaculo, foliis linearibus pubescentibus utrinquè attenuatis plerisque medio utrinquè unidentatis. (1) ad Cap. Bonæ-Spei. Nees ast. p. 224. excl. syn. Lam. et Spr. Am. alternifolius Roth cat. bot. 2. p. 111. *Kantia amelloides* Zeyh. 1 pl. exs. n. 99. Radix fibrosa caulis à basi ramosus. (v. s. in h. Morie.)
11. *A. TRIDACTYLUS*, caulibus è collo pluribus diffusis, foliis pedunculis involucrique actis patentibus rigidis confertè hispidis, foliis petiolatis tripartitis lobis linearì-oblongis integerrimis. (1) in Africà australi legit cl. Drege! *Leysera tridactyla* E. Mey. 1 in Dreg. coll. Rec. paleas flores disci superantes. (v. s.)
12. *A. MICROGLOSSUS*, caule diffuso ramosissimo tereti parè et patentem puberulo, foliis linearì-oblongis basi attenuatis integerrimis glabrinaculis, invol. squamis linearibus complicatis acuminatis dorso hispidulis, ligulis minimis. (1) in Africà Capensi ad Olifant rivier, Zilverfontain, et Klein Namaqualand legit cl. Drege! Achænia radii et exteriora disci compressa pubescentia pappo brevi paleaceo dentato coronata; disci intima abortiva. Paleas ad latus fl. disci exter. angustissimæ, in centro disci subnullæ. (v. s.)

### † *Species exclusæ.*

- A. CAROLINIANUS* Walt. = *Eolypa brachypoda*.  
*A. DIFFUSUS* Forst. = *Chiliotrichium amelloideum*.  
*A. FLOREBUNDUS* Willd. h. = *Andromachia igniaria*.  
*A. PEDUNCULATUS* Ort. = *Tridax procumbens*.  
*A. ROSEMARINIFOLIUS* Poepp. = *Chiliotrichium amelloideum* γ.  
*A. SPINULOSUS* Pursh. = *Aplopappus spinulosus*.  
*A. UMBELLATUS* Linn. = *Liabum Brownæ*.  
*A. VILLOSUS* Pursh. = *Chrysopsis villosa*.

### C. CORETHROGYNE.

Capitulum multiflorum, fl. radii ligulatis neutris 1-serialibus disci tubo brevi fauce cylindricà apice brevissimè 5-dentatà. Invol. squamæ 2-3-seriales lanceolato-lineares subæquales. Recept. planum paleis linearibus apice scabridis inter flores onustum. Styli rami exserti basi lineares margine glandulosi apice setas rigidas vel penticilliformes gerentes. Achænium turbinatum densè sericeum, villis supremis pappum exteriorem simulantibus. Pappus pluriserialis setosus scabridus. — Herba Californica multicaulis tota molliter incano-tomentosa. Folia alterna acuta, infer. oblongo-lanceolata basi attenuata apice serrata, super. linearia integerrima. Capitula ad apices ramorum solitaria. Invol. parè villosa. Ligulæ angustæ pallidæ ex sicco purpurascentes. Discus flavus. Pappus rubens. — An *Amello*? an *Osmia* affinis? — Nomen ex *αρεθρον*, scopus, et *γυνή*, femina.

1. *C. CALIFORNICA*. γ. in Californiâ anno 1833 legi indef. Douglas. (v. s. comm. à cl. Bentham.)

CL. CHILIOTRICHUM *Cass. bull. philom. 1817. p. 69. dict. 4. suppl. 5-6. R. p. 576. 37. p. 463 et 489. Less. syn. 162. Nees ast. 239. — Tropidolepis Tausch in florâ 12 (1829). p. 68. — Amelli sp. Forst.*

Capitulum multiflorum, fl. radii ligulatis fœmineis 1-seriatis, disci 5-dentatis hermaphr. Invol. squamæ imbricatæ oblongæ acutæ. Recept. convexum paleis linearibus apice barbatis inter flores onustum. Stigmata fl. disci subulato-lineariter elongata pubera. Achænia gracilia cylindracea angulato-striata. Pappus pluriserialis, setis filiformibus scabridis inæqualibus persistentibus. — Frutices in extremâ Amer. austr. spontanei parvi ramosi. Folia alterna sessilia coriacea integerrima margine revoluta supra glabra subtus plus minus tomentosa. Pedunculi solitarii 1-cephali tomentosi. Ligulæ albæ subtus purpurascentes. — *Aster phycioides* H. B. et Kunth fortè ex Nees huic generi adiungendus?

1. *C. AMELLOIDES* (Cass. dict. 8. p. 576), foliis oblongo-obovatis basi angustatis planiusculis. ♂ ad fretum Magellanicum (Forst. Comm. 1), ad portum Egmont (Née!). Amellus diffusus Forst. comm. Gœtt. 9. p. 39. *Tropidolepis diffusa* Tausch in florâ 12. p. 68. *Aster magellanicus* Spreng. syst. 3. p. 526. *Ch. amelloides* γ Nees ast. 240. Folia subtus densè et fulvo-tomentosa. (v. s.)

β. *lanceolatum* (Nees l. c.), foliis lanceolatis acutis basi attenuatis. ♂ in insula Maclovianis ubi arborum gigas interdum 4-5-pedalis, sed in monte Chatelloux vix semipedalis d'Urv. ! fl. mal. n. 80. Gaudich. ! fl. mal. p. 16. *Arbuste à feuilles de romarin* Pern. mal. 2. p. 61. Eadem in Staten-Island legit cl. Webster ex h. Lindl. ! (v. s.)

γ. ? *rosmarinifolium* (Nees l. c.), foliis linearibus intensius margine revolutis basi non angustatis. ♂ in Andibus Chilensium (Poepp. 1), ad Talca-reguè (Cl. Gay !), Amellus rosmarinifolius Poepp. ! exa. *Ch. rosmarinifolium* Less. in linnæâ 1831. p. 109. An species propria? (v. s.)

#### Subdiv. II. HETEROTHALAMÆ.

Capitula dioica; masc. recept. nudum, ligulæ 1-seriales; form. recept. paleis, ligulæ pluriseriales.

CII. HETEROTHALAMUS *Less. in linnæâ 1830. p. 145. 1831. p. 149 et 501. syn. p. 205.*

Capitula multiflora radiata dioica. Involucrum pluriserialiter, squamis siccis lanceolatis æqualibus disco subbrevioribus. Pappus 1-serialis setaceus caducus corollâ brevior. ♂ Recept. planum nudum. Ligulæ uniserialiter fœminæ minimæ, fructu abortivo. Cor. disci tubulosæ 5-dentatæ, antheris ecaudatis, stylo bifido abortivo, pappo clavato. ♀ Recept. conicum paleis invol. similibus onustum. Cor. ferè omnes? ligulatæ breves fœminæ, centrales paucissimæ fortè 5-dentatæ masculæ? Achænia angulata glaberrima basi angustata, pappo apice attenuato. — Frutices Brasilienses glaberrimi erecti Athanasias aut Chrysocomas facie. Folia alterna lineari- aut oblongo-acuta integerrima aut serrata. Capitula subglobosa, ♂ corymbosa parva, ♀ in axillis ramorum sterilium breviter pedunculata duplò majora. Corollæ utriusque sexus flavæ, quo caract. et corollis plurimis ligulatis et recept. ♀ paleaceo à Baccharideis genus alienum videtur.

1. *H. BRUSIIFORMIS* (Less. ! in linnæâ 1831. p. 504), foliis linearibus integerrimis. ♂ in Brasiliâ (Sellow ! ) prov. Rio-Grande (Gaudich. ! ). *Marachalla* (nec *Melananthera*) aliena Spreng. syst. 3. p. 446. (v. s. ♂ et ♀.)

2. *H. PERAZIODES* (Less. in LINNÆA 1831. p. 504), foliis oblongo-obovatis serratis. ♂ in Brasilia (Sellow) prov. Rio-Grande (h. Mus. imp. Bras. n. 958). (v. s. in h. Mus. reg. Par.)

Div. II. ASTERÆÆ.

Recept. epaleaceum nunc nudum nunc alveolatum aut fimbriiferum. Capitula heterogama, fl. radii formineis ligulatis heterochromis (nempe albis purpureis cœruleisve seu cyanicis, disco luteo seu xanthico.)

Subdiv. I. EUASTERÆÆ.

Ligulæ 1-seriales. Pappus conformis pilosus uni- aut pluriserialis. Achænia compresso-plana.

- III. MAIRIA. — *Mairia* et *Zyrphelia* Nees ast. p. 248 et 250. — *Pteropappus* (sect. *Asteria*) et *Zyrphelia* (sect. *Elphegææ*) Less. syn. 166 et 183. — *Asteria Arnicae* et *Leyseræ* sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis fertilibus, disci tubulosis 5-dentatis hermaphroditis nunc fertilibus nunc sterilibus. Recept. nudum subalveolatum. Involucrum pauciserialè subimbricatum. Anthæræ basi omninò muticæ. Achænia plano-compressa glabra aut parcè pilosa erostria. Pappus 1-serialis, setis plumosis. — Herbæ rhizomate suffrutescente aut suffrutices Capenses. Folia alterna oblongo-obovata aut linearia, subrigidula, dentata aut integra. Rami caulève apice subnudi capitula spilaria gerentes. Discus luteus. Radius violaceus aut albus. — Inter nomina 3 æquali jure admittenda Mairiam elegi, ne peregrinator de botanicâ Capensi bene meritus honore debito fraudetur.

Sect. I. PTEROPAPPUS Less. syn. 166. — *Mairia* Nees ast. 248.

Flores disci fertiles. Ligulæ stam. filamenta sterilia nonnulla sæpius gerentes. — Herbacæ rhizomate perenni sublignoso.

1. *M. CRENATA* (Nees ast. 348), foliis subradicalibus obovato-oblongis basi attenuatis carnosiusculis apice grossè crenato-dentatis glabris, pedunculo hirsuto, involucri squamis dorso æqualibus aut longioribus. 2/ ad Cap. Bonæ-Spei in mont. Langkloof (Th.). Tabulari (Mundt et Maire Dreg.), circa urbem et in distr. Zwellendam (Eckl. et Zeyh.) *Arnica crenata* Thunb. fl. cap. 668. Ecklon! it. r. n. 54. b. Lodd. bot. cap. t. 901. *Gerbera crenata* bot. reg. t. 855. Eckl. in h. Duuant. *Aster crenatus* Less. syn. 166. Pedunculus foliis brevior, æqualis aut duplò longior, nempe 1-4 pollicaris. Ligulæ violacæ. Capitulum magn. Asteris alpini. Variat foliis glabris aut pilosiusculis. (v. s.)
2. *M. HIRSUTA*, caule ascendenti erecto subsimplici hirsutissimo, foliis in parte caulis ferè dimidiâ inferiore obovatis in petiolum attenuatis grossè dentatis supra jun. subhirsutis demùm glabris subtus densè hirsutis, superioribus paucis linearibus integerrimis hirsutis, capitulis caulem aut ramos paucos elongatos terminantibus, invol. hirsuto disco æquali. 2/ ad Cap. Bonæ-Spei in distr. Zwellendam in sylvis montosis prope Paspas-Valley leg. cl. Ecklon! et Zeyh. Affinis *M. crenatæ* sed differt caule ferè ad medium folioso, hirsutis omnium partium, et achæniis ipso hirsuto. Ligulæ stam. rudimentis orbatæ. (v. s.)
3. *M. PERAZIODES* (Nees ast. 249), foliis subradicalibus lineari-spathulatis strigilloso-hirsutis subintegerrimis membranaceis, pedunculo-hispido-pubescente, involucri squamis disco sublongioribus. 2/ ad Cap. Bonæ-Spei in Grœnkloof et Swartland (Th.). *Aster perazioides* Less. syn. 167. *Leysera pilosella* Thunb. herb. ex parte juxta Less. Facies *Perezii* squarrozæ.

Sect. II. *ZYRPHELIS* Cass. ann. sc. nat. vol. 17 (1829). p. 420. repet. in diss. segr. p. 36 et dict. sc. nat. 60. p. 597. Nees ast. 259. — *Elphegæ* sp. Less. in *linnæd.*

Flores disci plerique steriles. Ligulæ stam. filamentis destitutæ.

### §. 1. *Acaules.*

4. *M. BURCHELLII*, subacaulis, foliis radicalibus oblongo-linearibus glabris rigidis integerrimis basi attenuatis et hirsuto-ciliatis, caulibus subaphyllis parvis ramosis hirsutulis, invol. squamis acuminatis ciliatis. 2/ ad Cap. Bonæ-Spei ad orientem urbis legit cl. Burchell et mecum comm. sub n. 8200 est. geogr. Folia infernè nervo subtus prominulo subtriquetra apice semicomplicata 4-5 pollic. longa. Caules foliis paulò breviores. Invol. biserial. Achænia subcompressa villosissima basi attenuata. Species ex habita ad sect. primam primè fronte referenda, sed ex disco sterili huc verè pertinens et sic utramque colligens. (v. s.)

### §. 2. *Fruticosa caulescentes.*

5. *M. TAXIFOLIA*, caule fruticuloso erecto ramoso, foliis linearibus acutis basi rigidè ciliatis, invol. squamis acuminatis, achæniis glabris 3 ad Cap. Bonæ-Spei in collibus montis Tabularis (Th. ! Dreg. !). *Aster taxifolius* Linn. sp. 1265. Berg. cap. 286. *Leyssera ciliata* Thunb. fl. cap. 690. *Aster ciliaris* Thunb. ! in h. L'Hér. *Zyrphelis amœna* Cass. dict. 60. p. 598. *Elphegæ ciliata* Less. in *linnæd.* 1831. p. 126. *Zyrphelis taxifolia* Nees ast. p. 250. *Aster filiformis* Eckl. ! coll. exs. n. 103. b. in h. Moric. sed excl. syn. — Burch. ! cat. geogr. n. 592. Achænia matura compressa obovata glabra. Ligulæ subilacino-violaceæ. (v. s.)
6. *M. MICROCEPHALA*, caule fruticuloso erecto ramoso, foliis linearibus acutis basi parvè ciliatis aut pube brevi hirsutulis, invol. squamis interioribus obtusis, achæniis pilosiusculis. 3 ad Cap. Bonæ-Spei prope Kamnario (Mundt et Maire), ad orientem urbis (Burch. ! cat. geogr. n. 8159), ad Zeederberge (Dreg. !) in Zwelldam (Eckl. ! et Zeyh.). *Elphegæ microcephala* Less. syn. 184. *Zyrphelis microcephala* Nees syn. 252. *Aster nanus* Ern. Mey. ! in Dreg. coll. Folia .4 lin. longa. Capitula dimidiò minora quàm prioris cui simillima. (v. s.)
7. *M. LASIOCARPA*, caule fruticuloso erecto ramoso, foliis linearibus acutis basi rigidè ciliatis, invol. squamis acuminatis, achæniis sericeo-villosis. 3 ad Cap. Bonæ-Spei in distr. Caledon leg. cl. Ecklon ! et Zeyh. Simillima *M. taxifolia*, sed achæniis densè et adpressè sericeo-villosis facillè differt. (v. s.)

CIV. *FELICIA*. — *Felicia* et *Polyarrhena* Cass. bull. philom. 1818. p. 165. dict. 37. p. 464. et 490. 56. p. 172. Nees ast. 207 et 205. — *Polylepis* (1) (sect. *Asteris*) et *Polyarrhena* (sect. *Elphegæ*) Less. syn. 173 et 184. — *Asteris* sp. Linn. et auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis foemineis 1-serialibus, disci tubulosis 5-dentatis hermaphr. sæpius omnibus fertilibus. Recept. nudum areolatum. Invol. squamæ bi- aut pluriseriales adpressæ angustæ. Achænia compressa glabra aut pubescentia. Pappus 1-serialis brevis caducus, setis flexuosis filiformibus serrulatis. — Herbariis aut frutices Capenses. Caules ramosi. Rami 1-cephali sæpè fastigiati. Folia alterna angusta crassiuscula. Discus luteus, radius cœruleus aut albus.

---

(1) *Polylepis* fl. Per. est *Rosacea-Sanguisorbea*, et *Polylepis* Cass. est sect. *Spharanthi*.

Sect. I. *HEBECARPEÆ*. — *Felicia* Cass. seu *Polylepidis* Less. species plurimæ.

*Achænia pilosiuscula* aut *subhirtella*. — *Folia linearia*.

*Herbaceæ, annuæ, radice fibrosâ.*

1. *F. TENELLA* (Nees ast. 208), herbacea, foliis sparsis linearibus patulis setuloso-ciliatis, ramis floridis apice nudis 1-cephalis, invol. squamis linearibus acutis dorso subhirsutis, achæniis parvis et minutè pilosiusculis. (1) ad Cap. Bonæ-Spei, in collibus circa urbem ad Paarlberg (Dreg.), in reg. Gariepina (Dreg.), in distr. Caledon et Stellenbosch (Eckl. et Zeyh.). *Aster tenellus* Linn. mant. 471. Jacq. obs. 4. p. 88. Curt. bot. mag. t. 33. Less. syn. 173. Sieb. pl. exs. 253. *Felicia fragilis* Cass. dict. 16. p. 315. *Aster deustus* et *Ast. tenellus* Thunb. fl. cap. 686 *Kaulfussia ciliata* Spreng. f. suppl. syst. p. 25. Zeyh. pl. exs. n. 36a. *Gineraria tenella* Link. enum. 2. p. 334. — Pluk alm. t. 271. f. 4. Variat caule erecto aut diffuso, foliis remotè aut crebrè serrato-ciliatis, radio pallidiùs aut intensiùs cæruleo. Fortè hæc pertinet *Cotula linifolia* Burm. prod. fl. cap. 27. certè ex herb. radio donata. (v. sp. et v. c.)

♂? *longifolia*, caule basi suffruticuloso, ramis glabris teretibus, florifera apice nudis 1-cephalis, foliis angustè linearibus sparsis vix acutis setulas brevès dentiformes rigidas hinc inde ad margines gerentes, invol. squamis linearibus acutis dorso subhirsutis, achæniis maturis minutè et parvis pilosiusculis. ♂ ♀ in Africa Capensæ ad Bergvriër et Giffthurg legit cl. Dregel ad Stellenbosch cl. Ecklon et Zeyh. Variat foliis 1-1½ longis, semilineam latis, plus minus setuloso-ciliatis; pedicellis 1-4 poll. longis. An var. *F. tenellæ frutescens*? An spec. propria? (v. s.)

2. *F. COTULOIDES*, caule herbaceo tenello glabro à basi ramosissimo, ramis apice longè demodatis 1-cephalis, foliis angustè-linearibus acutis glabresculis integerrimis, invol. squamis linearibus glabris acutis, achæniis vix sub lente puberulis, pappo multiset. (1) in Africa Capensæ ad Bergvriër (Dreg.) in distr. Worcester (Eckl. et Zeyh.). Radix exilis. Herba 5-6 pollic. Folia 5-6 lin. longa 1½ lin. lata. Affinis *F. tenellæ*, sed præter not. ind. capitula duplè ferè minor. (v. s.)
3. *F. MICROGONIMA*, radice simplici, caulibus è collo pluribus herbaceis erectis teretibus versus apices glanduloso-puberulis viscosis (?) basi indivisis apice corymbosis, foliis linearibus integerrimis obtusiusculis, invol. squamis linearibus acutis adnatisque glabresculis. (1) ? in Africa Capensæ regione Gariepina legit cl. Dregel Radix et collum sublignescens. Caules 8-12 pollic. Folia 9-10 lin. longa, semilineam lata. Capitula *F. tenellæ*, sed laxè corymbosa. Achænia pube brevi sparsa adpressa donata. (v. s.)
4. *F. ADPUNIS* (Nees ast. 209), herbacea, foliis obovato-oblongis hirsutis, inferioribus oppositis. (1) ad Cap. Bon.-Spei. *Aster adfinis* Less. syn. 174. Proxima *F. tenellæ*, sed caulis totus hirsutus nec basi glaber. Folia latiora basi angustata utrinquè hirsuta; invol. squamæ 3-4-seriales nec biseriales. Pappus disco corollæ æqualis nec eà brevior. Fel. dubia Cass. dict. 16. p. 315. (v. s. in h. Cass. nunc Mus. reg. Par.)

\*\* *Caule herbaceo simplicissimo rhizomate perenni orto.*

5. *F. MACRORHIZA*, caule herbaceo simplicissimo è rhizomate perenni orto, foliis alternis strigilloso-hirsutis obovato-linearibus confertissimis integerrimis, achæniis pilosis ciliatis. ♀ ad Cap. Bonæ-Spei. *Aster macrorhizus* Thunb. fl. cap. 687. Less. syn. 175. Nees ast. 3a. Habitus *Erigerontis* graminæ seu *Mairia perzeioidis*.
6. *F. SARCOGONIMUS*, caulibus herbaceis erectis teretibus tenuissimè pubescentibus, foliis elliptico-oblongis utrinquè glabris minutissimè serrato-ciliatis reticulato-subtrinerviis, axillis foliiferis, ramis apice nudis 1-cephalis in paniculam ferè

dispositis, invol. squamis acutis glabriusculis, acheniis puberis.  $\frac{7}{8}$  in Africa austr. orient. ad Port-Natal legit cl. Drege! Habitu accedit ad *Erigeron* fetidum. Folia poll. longa, 3 lin. lata. (v. s.)

\*\*\* *Caule fruticoso.*

7. *F. FASCICULARIS*, caule fruticoso erecto ramoso glabriusculo, foliis linearifiliformibus patulis integerrimis rigidulis setis sparsis plus minus scabris axillis folia novella fasciculata gerentibus, capitulis pedicellatis solitariis, invol. squamis linearibus acutis dorso pedicellisque minutè glandulosis.  $\frac{5}{8}$  in Africa Capensis distr. Carro (Dreg.! Eckl.! et Zeyh.) in Uitenhage et territorio cesso (Eckl. et Zeyh.). *Aster fascicularis* E. Mey.! in Dreg. coll. *Felicia unariata* Nees ast. 210? *Aster filifolius* Zeyh.! exa. in h. Moric. *Soffrutex* circ. *scirpoidalis* spithameusve. Folia var. ferè glabra, caulina 3 lin., axill. 2 lin. longa achenium villosum. Pappus scaber. (v. s.)

$\beta$ . *pubescens*, ramis pubescentibus elongatis, foliis ramulis linearibus, ad axillas nudis.  $\frac{5}{8}$  in distr. Uitenhage (Eckl.! et Zeyh.) et prope Becho (Dreg!). (v. s.)

8. *F. ANGUSTIFOLIA*, caule fruticoso erecto, ramis pubescentibus aut laevib., foliis linearibus planis ciliatis acutiusculis adultis glabris, capitulis longè pedunculatis  $\frac{5}{8}$  ad Cap. Bon.-Spei.

$\alpha$ . *hirsutifolia*, ramis canescentibus, foliis junioribus pube brevi molli subhirsutis.  $\frac{5}{8}$  Burm.! Dreg.! Eckl.! et Zeyh. *Aster hysopifolius* Berg. cap. 287. Less. syn. 175. *Felicia hysopifolia* Nees ast. 211. *Aster villosus* et *hirtus* Thunb. fl. cap. 687 et 688. ex Less. *Aster frutescens* Burm.! prod. 27. *Conyza chrysocomoides* Desf. arb. 1. p. 232. excl. syn. *Aster chrysocomoides* Desf.! cat. h. par. 1815. p. 121. non Willd. *Aster conyzoides* Desf.! cat. h. par. 1829. p. 175 non Willd. (v. s.)

$\beta$ . *glabra*, ramis vix pubescentibus, foliis etiam junioribus glabris.  $\frac{5}{8}$  ad Zoederbergen et Stormberg Dreg.! *Aster angustifolius* Jacq. h. Schumbr. t. 370. Less. syn. 177. *Felicia angustifolia* Nees ast. 213. (v. s. et  $\alpha$ .)

9. *F. HIRUTA*, caule fruticoso, ramis virgatis hirsuto-canescens, foliis linearibus planis obtusiusculis utrinquè strigilloso-hirsutis, capitulis solitariis breviter pedunculatis, acheniis hirsutiusculis.  $\frac{5}{8}$  ad Cap. Bonæ-Spei in regione Trans-Gariepinâ (Borch.! cat. geogr. n. 2331), in territorio cesso et Graf Ruyner (Eckl.! et Zeyh.) ad Klaarprivier (Dreg.). Folia 5-6 lin. longa erecta per totos ramos disposita, apice ramuli breviter denudato. Invol. campanulatum, squamis 2-3-ser. exter. dorso hirtis omnibus margine membranaceis. An forti *Aster lanuginosus* Wendt beytr. 2. p. 9. (v. s.)

10. *F. RIGIDULA*, caule fruticoso, ramis subadgregatis rigidulis puberis, foliis linearibus planis obtusis utrinquè adpressè strigilloso-hirsutis, capitulis solitariis pedunculatis, invol. squamis dorso pulverulento-macronatis diocam ferè superantibus, acheniis puberulis  $\frac{5}{8}$  ad Cap. Bonæ-Spei (h. Bolus!) in distr. Uitenhage (Eckl.! et Zeyh.). Valdè ex caracteribus accedit ad *F. hirsutam*. sed habitus satis diversus et inv. squamæ extus non hirsutæ divisionem asserunt. (v. s.)

$\beta$ . *subcanescens*, foliis ramulisque pube confertiore subcanescentibus.  $\frac{5}{8}$  in Africa Capensi (Dreg!), in distr. Uitenhage (Eckl.! et Zeyh.). (v. s.)

11. *F. CANA*, caule fruticoso adscendente ramosissimo, ramis folisque pube confertâ brevi velutino-canis, foliis linearifiliformibus obtusis integerrimis, axillis sæpè foliiferis, pedicellis elongatis nudis 1-cephalis pubescentibus, invol. squamis linearibus acutis puberis, acheniis sparè pilosiusculis.  $\frac{5}{8}$  in Africa Capensis distr. Zwelendani et deserto Karro leg. cl. Ecklon! et Zeyh. *Soffrutex* 5-6 pollic. totus canus. Folia 5 lin. longa. Ligula albæ lineares. (v. s.)

12. *F. ZYTHREI* (Nees ast. 215), caule fruticoso, ramis adultis glabris, foliis oblongo-cuneiformibus planis glabris densè strigilloso-ciliatis, capitulis longè pedunculatis, invol. squamis glabris disco subbreioribus acuminatis, acheniis

2. *H. psalmioides* (Less. in *linnaea* 1831. p. 504), foliis oblongo-obovatis serratis.  $\frac{3}{4}$  in Brasiliæ (Sellow) prov. Rio-Grande (h. Mus. imp. Bras. n. 958). (v. s. in h. Mus. reg. Par.)

### Div. II. ASTERÆÆ.

Recept. epaleecrum nunc nudum nunc alveolatum aut simbrilliferum. Capitula heterogama, fl. radii fœmineis ligulatis heterochromis (nempe albis purpureis cœruleisve seu cyanicis, disco luteo seu xanthico.)

### Subdiv. I. EUASTERÆÆ.

Ligulæ 1-seriales. Pappus conformis pilosus uni- aut pluriserialis. Achænia compresso-plana.

- CIII. MAIRIA. — Mairia et Zyrphelis *Nees ast. p. 248 et 250.* — Pteropappus (sect. Asteria) et Zyrphelis (sect. Elphægæ) *Less. syn. 166 et 183.* — Asteris Arnicae et *Leysseræ sp. auct.*

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis fertilibus, disci tubulosis 5-dentatis hermaphroditiis nunc fertilibus nunc sterilibus. Recept. nudum subalveolatum. Involucrum pauciserialia subimbricatum. Anthæræ basi omnino muticæ. Achænia plano-compressa glabra aut parcè pilosa erostris. Pappus 1-serialis, setis plumosis. — Herbæ rhizomate suffrutescente aut suffrutices Capenses. Folia alterna oblongo-obovata aut linearia, subrigidula, dentata aut integra. Rami caulesve apice subnudi capitula æolitaria gerentes. Discus luteus. Radius violaceus aut albus. — Inter nomina 3 æquali jure admittenda Mairiam elegi, ne peregrinator de botanicâ Capensi bene meritis honore debito fraudetur.

### Sect. I. PTEROPAPPUS *Less. syn. 166.* — Mairia *Nees ast. 248.*

Flores disci fertiles. Ligulæ stam. filamenta sterilia nonnulla sæpius gerentes. — Herbacæ rhizomate pereuni sublignoso.

1. *M. CRENATA* (Nees ast. 348), foliis subradicalibus obovato-oblongis basi attenuatis carnosiusculis apice grossè crenato-dentatis glabris, pedunculo hirsuto, involucri squamis dorso æqualibus aut longioribus.  $\frac{1}{2}$  ad Cap. Bonæ-Spei in mont. Langkloof (Th.). Tabulari (Mundt et Maire Dreg.), circa urbem et in distr. Zwellendam (Eckl. et Zeyh.) Arnica crenata Thunb. fl. cap. 668. Ecklon! it. r. n. 54. b. Lodd. bot. cap. t. 901. Gerbera crenata bot. reg. t. 855. Eckl. in h. Duuant. Aster crenatus Less. syn. 166. Pedunculus foliis brevior, æqualis aut duplò longior, nempe 1-4 pollicaris. Ligulæ violacæ. Capitulum magn. Asteris alpini. Variat foliis glabris aut pilosiusculis. (v. s.)
2. *M. HIRSUTA*, caule adscendenti erecto subsimplici hirsutissimo, foliis in parte caulis ferè dimidiâ inferiore obovatis in petiolum attenuatis grossè dentatis supra junc. subhirsutis demùm glabris subitis densè hirsutis, superioribus paucis linearibus integerrimis hirsutis, capitulis caulem aut ramos paucos elongatos terminantibus, invol. hirsuto disco æquali.  $\frac{1}{2}$  ad Cap. Bonæ-Spei in distr. Zwellendam in sylvis montosis prope Puspas-Valley leg. cl. Ecklon! et Zeyh. Affinis *M. crenatæ* sed differt caule ferè ad medium folioso, hirsutis omnium partium, et achæniis ipso hirsuto. Ligulæ stam. rudimentis orbatæ. (v. s.)
3. *M. PEREZIODES* (Nees ast. 249), foliis subradicalibus linearispathnatis strigilloso-hirsutis subintegerrimis membranaceis, pedunculo-hispido-pubescente, involucri squamis disco sublongioribus.  $\frac{1}{2}$  ad Cap. Bonæ-Spei in Grænekloof et Swartland (Th.). Aster perzioides Less. syn. 167. *Leyssera pilosella* Thunb. herb. ex parte juxta Less. Facies Perezizæ squarrosæ.



dispositis, invol. squamis acutis glabriusculis, acheniis puberis.  $\frac{1}{2}$  in Africa austr. orient. ad Port-Natal legit cl. Drege! Habitu accedit ad *Erigeron* frigidum. Folia poll. longa, 3 lin. lata. (v. s.)

\*\*\* *Caule fruticoso.*

7. *F. fascicularis*, caule fruticoso erecto ramoso glabriusculo, foliis linearifiliformibus patulis integerrimis rigidulis setis sparsis plus minus acutis axillis folia novella fasciculata gerentibus, capitulis pedicellatis solitariis, invol. squamis linearibus acutis dorso pedicellisque minutè glandulosis.  $\frac{1}{2}$  in Africa Capensis distr. Carro (Dreg.! Eckl.! et Zeyh.) in Uitenhage et territorio cesso (Eckl. et Zeyh.). *Aster fascicularis* E. Mey.! in Dreg. coll. *Felicia maritima* Nees ast. 210? *Aster filifolius* Zeyh.! exa. in h. Moric. *Suffrutex* circ. semipedalis spithameusve. Folia var. serè glabra, caulina 3 lin., axill. 2 lin. long. achenium villosum. Pappus scaber. (v. s.)

$\beta$ . *pubescens*, ramis pubescentibus elongatis, foliis rametis linearibus, ad axillas nudis.  $\frac{1}{2}$  in distr. Uitenhage (Eckl.! et Zeyh.) et prope Bode (Dreg!). (v. s.)

8. *F. angustifolia*, caule fruticoso erecto, ramis pubescentibus aut laevibus, foliis linearibus planis ciliatis acutiusculis adultis glabris, capitulis longi pedunculatis  $\frac{1}{2}$  ad Cap. Bon.-Spei.

$\alpha$ . *hypsipifolia*, ramis canescentibus, foliis junioribus pube brevi molli subhirsutis.  $\frac{1}{2}$  Barm.! Dreg.! Eckl.! et Zeyh. *Aster hypsipifolius* Berg. cap. 287. Less. syn. 175. *Felicia hypsipifolia* Nees ast. 211. *Aster villosus* et hirtus Thunb. fl. cap. 687 et 688. ex Less. *Aster fruticosus* Barm.! prod. 27. *Conyza chrysocomoides* Desf. arb. 1. p. 232. excl. syn. *Aster chrysocomoides* Desf.! cat. h. par. 1815. p. 121. non Willd. *Aster conyzoides* Desf.! cat. h. par. 1829. p. 175 non Willd. (v. s.)

$\beta$ . *glabra*, ramis vix pubescentibus, foliis etiam junioribus glabris.  $\frac{1}{2}$  ad Zeederbergen et Stormberg Dreg.! *Aster angustifolius* Jacq. h. Schomb. t. 370. Less. syn. 177. *Felicia angustifolia* Nees ast. 213. (v. s. et c.)

9. *F. hirsuta*, caule fruticoso, ramis virgatis hirsuto-canescens, foliis linearibus planis obtusiusculis utrinquè strigilloso-hirsutis, capitulis solitariis breviter pedunculatis, acheniis hirsutiusculis.  $\frac{1}{2}$  ad Cap. Bon.-Spei in regione Trans-Gariepinâ (Burch.! cat. geogr. n. 2331, in territorio cesso et Graf Reynet (Eckl.! et Zeyh.) ad Klaarpriver (Dreg.!). Folia 5-6 lin. longa erecta per totos ramos disposita, apice ramuli breviter denudato. Invol. campanulatus, squamis 2-3-ser. exter. dorso hirtis omnibus margine membranacea. An forte *Aster lanuginosus* Wendl. beytr. 2. p. 9. (v. s.)

10. *F. rigidula*, caule fruticoso, ramis subadgregatis rigidulis puberis, foliis linearibus planis obtusis utrinquè adpressè strigilloso-hirsutis, capitulis solitariis pedunculatis, invol. squamis dorso pulverulento-mucronatis diacna fere superantibus, acheniis puberulis  $\frac{1}{2}$  ad Cap. Bon.-Spei (h. Boiss.!) in distr. Uitenhage (Eckl.! et Zeyh.). Valdè ex characteribus accedit ad *F. hirsutam*, sed habitus satis diversus et invol. squamæ extus non hirsutæ divisionem asserunt. (v. s.)

$\beta$ . *subcanescens*, foliis ramulisque pube confertiore subcanescentibus.  $\frac{1}{2}$  in Africâ Capensi (Dreg.!), in distr. Uitenhage (Eckl.! et Zeyh.). (v. s.)

11. *F. cana*, caule fruticoso adscendente ramosissimo, ramis foliisque pube confertâ brevi velutino-canis, foliis linearifiliformibus obtusis integerrimis, axillis sæpè soliferis, pedicellis elongatis nodis 1-cephalis pubescentibus, invol. squamis linearibus acutis puberis, acheniis sparè pilosiusculis.  $\frac{1}{2}$  in Africa Capensis distr. Zwelendani et deserto Karro leg. cl. Ecklon! et Zeyh. *Suffrutex* 5-6 pollic. totus caesus. Folia 5 lin. longa. Ligulæ albæ lineares. (v. s.)

12. *F. Zeyheri* (Nees ast. 215), caule fruticoso, ramis adultis glabris, foliis oblongo-ovatoformibus planis glabris densè strigilloso-ciliatis, capitulis longi pedunculatis, invol. squamis glabris disco subbrevioribus acuminatis, acheniis

dentato-sinuat hispidulis, involucri 2-3-seriali, floribus disci fertilibus, ligulis apice grossè dentatis. 3 ad Cap. Bonæ-Spei, Cineraria hirsuta Vent. malum. n. et t. 95.

2. *M. CYMBALARIA* (Nees ast. 218), herbacea, foliis petiolatis ovatis dentatis, floribus capituli omnibus fertilibus, invol. glabro subbiseriali. 2 ad Caput Bonæ-Spei inter saxa montis Tabularis (Mundt et Mair. Dreg.) etc. Aster cymbalarie Thunb. prod. fl. cap. 689. Felicia brachyglossa Cass. dict. 25. p. 97. Mun. brachyglossa Cass. dict. 37. p. 483. Aster oppositifolius Linn. amœn. 5. p. 369? (v. s.)

β. *microcephala*, capitulis paulò minoribus, floribus centralibus subabortivis, caule laxè diffuso. — Burch. ! cat. geogr. n. 647 et 8466. (v. s.)

3. *M. CYNALAROIDES*, foliis oblongis subdentatis utrinquè attenuatis, floribus disci abortu sterilibus, involucri uniseriali. ① ad Cap. Bonæ-Spei in arenosis lapideis jugi Leonis (Eckl. ! in h. Dunant. it. 2. n. 1050). Folia plerumque radialis basi angustata in petiolum latum trinervium, ciliis longis distantibus ornatum. Pedunculus scapiformis. Achænia radii plano-compressa subelliptica nigrescentia margine cillato-villosa ceterum vix pilosuscula, omnino ut in priore, ad fl. disci abortivi. An fortè Felicia dubia Cass. dict. 16. p. 315? (v. s.)

**CVL AGATHEA** Cass. *bull. philom.* 1815. p. 175. *dict. i. suppl.* p. 77. 13. p. 116. et 37. p. 463 et 489. — Detris *Adans. fam.* 2. p. 131. — Detrioides (sect. *Elphegea*) et Detris (sect. *Asteris*) Less. — Detridium et Agathæa Nees. — *Asteris* sp. Mill. — Cinerarie sp. Linn.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis, disci tubulosi 5-dentati hermaphr. fertilibus aut rarius sterilibus. Recept. nudum planum vix alveolatum. Invol. campanulati squamæ 1-2 seriales adpressæ complicato-carinatæ subherbaceo-rigidulæ. Achænia matura plano-compressa nervo tumidulo cincta, pilis rigidulis scabra (in *A. microphylla* lævia), sterilia glabra. Pappus 1-serialis, setis caducis aut persistentibus serrato-scabris. — Herbæ aut suffrutices Capenses. Folia omnia aut saltem infima opposita, integra aut parcè dentata, hispidula aut rarius glabra. Pedunculi nudi 1-cephali. Discus luteus. Radius cœruleus.

**Sect. I. DETRIDIIUM** — Detridium Nees syn. 255. — Elphegeæ sect. Detrioides Less. syn. 182.

Flores disci steriles. Pappus disci et radii pauci- (5-6) setus.

1. *A. BERGERIANA*, caule herbaceo simplici aut ramoso erecto molliter hispido, foliis plerisque alternis oblongo-obovatis acutis integris, capitulis longè pedunculatis. (1 ad Cap. Bonæ-Spei in arenosis lapideis jugi Leontini. Cineraria Bergeriana Spreng. syst. 3. p. 551. Elphegea Bergiana Less. syn. 182. Detridium Bergianum Nees syn. 256. Kaulfussia amelloides Eckl. ! in h. Dunant. Variat caule simplici 1-cephalo et parcè ramoso 2-7-cephalo. Habitus omnino sequentium. (v. s. in h. Bois. et Dunant.)

**Sect. II. DETRIS.** — Detris *Adans. l. c.* — Asteris sect. Detris Less. syn. 168. — Agathæa Cass. *ll. cc.* Nees syn. 257.

Flores disci pleriquè fertiles. Pappus multi- (10-15) setosus.

\* *Herbaceæ annuæ.*

2. *A. ROTUNDIFOLIA* (Nees ast. 258), caule herbaceo adscendente hirsutiusculo, foliis ovato-ellipticis obtusis integerrimis strigillosis uninerviis. (1 ad Cap. Bonæ-Spei. Aster rotundifolius Thunb. fl. cap. 689. Less. syn. 172. Achænia flavo-fusca.
3. *A. SERRATA* (Nees ast. 259), caule herbaceo adscendente infernè glabro superne hispidulo, ramis usquè ad capitula densè foliosis, foliis verticillatis lan-

ceolatis acutis callosis-serrulatis glabris trinerviis. ① ad Cap. Bonæ-Spei. Aster serratus Thunb. fl. cap. 689. Less. syn. 173. Similis priori et magis adhuc ex et habitu Heteropappo hispido.

4. *A. diffusa*, caule herbaceo tenui ramoso diffuso dense et breviter hirtello, foliis oppositis sessilibus elliptico-oblongis integerrimis hirtellis, pedicellis elongatis 1-cephalis nudis parvis et brevissimè puberulis, invol. squamis linearibus acutis dorso hirtellis, achæniis maturis in disco utroque fere lævibus. ①? in Africa Capensi legit cl. Drege! Pappus radii parvus caducissimus. Folia 2-3 lin. longa, lineam lata. (v. s.)
5. *A. stricta*, caule herbaceo erecto basi simplici sparse hispido, foliis oppositis lineari-lanceolatis acutis basi attenuatis remotè serratis utrinquè hispidis, ramis subpaniculatis confertè et brevè hirtellis, involucri squamis acutis subhirtellis, ligulis parvis. ① in Africa Capensi ad Riebbekastel legit cl. Drege! Ex habitu stricto et radice exili fibrosa ab *A. hispida* et *A. diffusa* facillè differt. Ligula minores quàm in aliis speciebus. Achænia juniora pallida puberula. (v. s.)
6. *A. tenera*, caule herbaceo erecto oppositè ramoso hispidulo, foliis oppositis remotis sessilibus trinerviis ovatis seu ovato-oblongis distanter et callosis-serratis membranaceis subhispidulis, pedicellis elongatis nudis hispidulis, invol. squamis lineari-lanceolatis dorso hispidis apice membranaceis glabris. ① in Africa Capensis regione Gariopina legit cl. Drege n. 2802! Herba 8-10 pedic. Capitula magnit. Agathæ amelloidis. Achænium oblongum pubero-villosum. (v. s.)

•• *Frustrantes.*

7. *A. Zeyheri* (Nees ast. 260), fruticosa, foliis oppositis alternis linearilanceolatis pellucidis-punctatis setoso-serratis, involucri squamis trinerviis, achæniis scabris. ③ ad Cap. Bonæ-Spei. Cineraria linifolia Zeyh. coll. ex. n. 53 in h. Moric. Kaulfussia amelloides var. Spreng. ex Nees. Folia 7-9 lin. longa, lineam sequilineamve lata glabrescens. Ligula lineares elongata. (v. s. in h. Moric.)
8. *A. barbata*, caule fruticoso erecto ramoso glabro, ramulis setis longis hispidis, foliis oppositis rariùs alternis lineari-oblongis basi attenuatis integerrimis margine et hinc inde subtus setis longis patulis ciliatis, ramis floridis basi foliolosis apice nudis glanduloso-hirtellis et subsetosis, invol. squamis dorso hirtellis margine subscariosis, achæniis utrinquè hirtis. ③ in Africa Capensi ad Graaf-Reynet alt. 2-300 ped. legit cl. Drege! Fruticulus vix pedalis. Folia 6-8 lin. longa, 1-2 lata. Capitula vix *A. amelloidi* minora. (v. s.)  
*♀? subaibescens*, setis paucioribus, foliis pube adpressâ præter setas confertè canescentibus minoribus. ③ ad Sneeuwbergen alt. 4-500 ped. legit cl. Drege! (v. s.)
9. *A. hispida*, caule imò basi fruticuloso ramosissimo cum foliis pedunculisque undiquè setis longis patulis rigidulis hispido, foliis alternis (imis paucis oppositis) lineari-subcuneatis obtusiusculis basi attenuatis integerrimis, invol. squamis setis et pube pulverulentâ extus scabris. ③ in Africa Capensi ad Zeederbergen alt. 2-300 ped. legit cl. Drege! Cineraria hispida E. Mey. in Dreg. coll. Achænium atrum in disco utroque sparse puberum. Invol. disco aequale. Ligula lineares invol. duplò longiores. (v. s.)  
*♀. barbigerà*, pedunculis densius hispidis, invol. squamis apice floccoso-barbatis. ③ in regione Gariopina (Dreg.!). Flocci squamas terminantes etiam in var. *♀* hinc inde obvii, sed in var. *♀* majores et constantes videntur. (v. s.)
10. *A. nuda*, caule fruticoso adscendente ramoso, ramis elongatis teretibus internè foliosis pube minutissimâ et setis patulis hispidis, superne longe aphyllis 1-cephalis pube glandulosâ et setulis rigidis brevibus hirtellis, foliis linearilanceolatis integerrimis margine subrevolutis setis rigidis ciliatis et hirtis, infer. oppositis, super. alternis, invol. squamis setalis ut in pedicello coloratis hirtis

lis, acheniis subglandulosis.  $\frac{5}{8}$  in Africâ Capensi ad Breederivier legit cl. Drege! Pedicelli 5 poll. longi. Folia 5-7 lin. longa. 2 lin. lata. (v. s.)

11. *A. ECKLONIS* (Nees ast. 262), fruticosa, foliis alternis elliptico-oblongis subamplexicaulibus integerrimis subciliatis substrigosis suprâ asperis, acheniis hirtulis, involucri squamis trinerviis.  $\frac{5}{8}$  ad Cap. Bonæ-Spei in calcariis inter Krede rivier et Duyreuhoks rivier. Aster Ecklonis Less. syn. 170. Setulae paucæ secus folior. margines ut in *A. Zeyeri* cujus fortè varietas latifolia (v. s. in h. Dussant comm. à cl. Ecklon.)
12. *A. SCABRIDA*, caule fruticoso erecto ramoso glabro, ramulis pedicellisque minutè pulverulento-scabridis, foliis alternis cuneatis basi attenuatis integris ciliatis apice subtruncatis 3-5-dentatis utrinquè echinulato-scabridis, pedicellis foliolis linearis integra paucæ gerentibus, invol. glabriusculis  $\frac{5}{8}$  in Africâ Capensi legit cl. Drege! Aster scabridus E. Mey. in Drege. coll. (v. s.)
13. *A. BRUVIFOLIA*, caule fruticoso erecto ramoso glabro, ramis nascentibus subvelutinis, senescentibus subspinosis, floridis apice nudis 1-cephalis, foliis alternis pube confertâ brevi velutinis basi cuneatis apice obtusè 3-5-dentatis ferè lobatis, invol. squamis 3-seriatis linearibus glabriusculis, acheniis maturis parè in disco pilosiusculis.  $\frac{5}{8}$  in Africâ Capensi ad Zilverfontain legit cl. Drege! Frutex 1-1  $\frac{1}{2}$ -pedalis rigidus. Folia 2 lin. longa. Cor. disci apice ru-bentes. Stigmata valdè hirta. Pappus rigidulus longus scabridus. (v. s.)
14. *A. LEOCARPA*, caulibus à collo pluribus diffusis basi suffruticosis hispidis, foliis alternis oblongo-cuneatis apice acutè 3-5-dentatis basi integris sessilibus utrinquè puberulis basi ciliatis pedicellis glanduloso-puberulis folio triplò longiori-bus, invol. squamis 1-2-seriatis lineari-lanceolatis acutis vix puberis, acheniis maturis glaberrimis.  $\frac{5}{8}$   $\frac{1}{2}$  in Africâ Capensi ad Drakestenberge alt. 3-400 ped. legit cl. Drege! Ligulae cœruleo-purpureæ. Achenia disci sterilia puberula, radii matura pappo deciduo calvescentia. Folia 7-9 lin. longa. (v. s.)
15. *A. MICROPHYLLA* (Cass. bull. philom. 1817. p. 183. dict. suppl. p. 63), fruticosa, foliis oppositis alternisque obovato-oblongis sessilibus suprâ et subtiùs ad nervum strigosis, acheniis glaberrimis, invol. squamis trinerviis.  $\frac{5}{8}$  ad Cap. Bonæ-Spei, inter saxa montis Tabularis (Eckl.!) ad Zuccereberge (Dreg.!) in distr. Caledon et circa urbem (Eckl. et Zeyh.) et in distr. Albany (Burch. cat. geog. n. 3966) Nees ast. 261. Aster æthiopicus Burm. prod. 27. Less. syn. 169. Cineraria trachyphylla Spreng. syst. 3. p. 547. Cineraria microphylla Vahl herb. ex Less. sed habui ex eod. herb. sub nom. Cineraria amelloïdia. Habitus sat varius. Rami hispidolii aut glabri. Folia interdum subglabra, sed acheniis glaberrimis imò levibus à genere distinctissima. (v. s.)
16. *A. STRIGOSA* (Nees ast. 263), fruticosa, foliis oppositis oblongo-cuneatis integerrimis erecto-adpressis apice recurvis utrinquè ramisque densè strigosis, acheniis scabris, pappi setia subplumosis, invol. squamis 1-nerviis.  $\frac{5}{8}$  ad Cap. Bonæ-Spei in Roggeweld, Karro, Uitenhagen, etc. Leyseria ovata Thunb. fl. cap. 692. Kaulfussia strigosa Spreng. f. suppl. syst. p. 25. Zeyh. coll. ex n. 270. Aster strigosus Licht. herb. non Thunb. Aster Charieis Less. syn. 171. (v. s. in h. Moric.)
17. *A. AMELLOIDES*, fruticosa, foliis oppositis ovato-vel orbiculato-ellipticis in petiolum brevem attenuatis integerrimis ramisque strigosis, acheniis punctatis et scabris, invol. squamis 1-nerviis.  $\frac{5}{8}$  ad Cap. Bonæ-Spei in lateribus montis Tabularis. Frequens in hortis Europæis. Cineraria amelloïdes Linn. sp. 1245. Berg. cap. 290. Agathæa cœlestis Cass. bull. philom. 1815. p. 175. dict. 1. suppl. p. 77. opusc. phyt. 6. Nees ast. 264. Aster Capensis Less. syn. 168. — Mill. icon. t. 76. f. 2. (v. v. c.)
18. *A. ELONGATA* (Nees ast. 266), fruticosa, ramis alternis glanduloso-hispidis, foliis oppositis sessilibus oblongis acutis integerrimis utrinquè patentè-hirsutis acheniis hirsutiusculis, invol. squamis angustis uninerviis ligulis unicoloribus.  $\frac{5}{8}$  ad Cap. Bonæ-Spei in arenosis Swartlandiæ et Saldanhabay (Th.) ad flum. Gariep (Dreg.!). Aster elongatus Thunb. fl. cap. 688. Less. syn. 170. Ciner-

ria hirta Vahl herb. ex Less. Invol. squamæ ext. hispidulæ, inter. glabræ submembranaceæ. (v. s.)

19. *A. tarrocon* (Nees ast. 267), fruticosa?, foliis oppositis oblongo-lanceolatis integerrimis utrinque patentibus-hirsutis, invol. squamis angustis 1-nerviis, acheniis setuloso-scabris, ligulis bicoloribus. ♂? ad Cap. Bonæ-Spei.

**CVII. BELLIDIASTRUM** Mich. nov. gen. t. 29. Cass. dict. sc. nat. 4. suppl. p. 70. 37. p. 494 non Vaill. nec Less. — *Margarita* Gaud. fl. helv. 5. p. 335. — *Dorenicia* sp. Linn. — *Arnica* sp. Hall. — *Asteris* sp. Less.

Capitulum radiatum, ligulis 1-2-seriatis heterochromis fœmineis, disci tubulosi hermaphroditi. Involucrum subcampanulatum, squamis 1-2-seriatis linearibus acutis subsoliaceis. Recept. conicum nudum punctatum. Stam. in fœmineis omnino abortiva. Achenia omnia oblonga subcompressa hispidula. Pappus 1-2-serialis, pilis numerosis seruosus scabris. — Herba Europæa perennis, trunco brevissimo radiciformi, foliis radicalibus petiolatis obovato-oblongis dentatis, scapo foliis longiore villosa 1-cephalo. Capitulum ferè Bellidis. — Cum *Astere* fortè non immeritò jungunt cl. Lessing et Lindley.

1. *B. MICHELII* (Cass. l. c.). 2. in sylvis montosis Europæ. *Dorenicum* Bellidiastrum Linn. sp. 1247. Jacq. fl. austr. t. 400. *Arnica* Bellidiastrum Vill. daph. 3. p. 212. Sturm deutschl. fl. f. 38. t. 15. Lam. l et DC. fl. fr. n. 3201. *Aster* Bellidiastrum Scop. carn. n. 1044. Less. syn. 166. *Margarita* Bellidiastrum Gaud. l. c. Bellidiastrum montanum Hoppe! pl. exa. 1832. Variet ligulis albis aut extus pilis minutis, interdum omnino rubris. (v. s.)

**CVIII. ASTER** Nees ast. p. 16. Lindl. adn. mss. in ast. Nees. — *Asteris* sp. Linn. Cass. Less. — *Amellus* Adans. non Linn.

Capitulum radiatum, fl. radii ligulatis fertilibus 1-serialibus, disci hermaphr. 5-dentatis. Recept. planum alveolatum, alveolorum marginibus plüs minüs dentatis. Involucrum squamæ pluriserialis laxè vel imbricatè apicè plüs minüs herbacæ inò interdum foliaceæ. Achenium compressum. Pappus pilosus persistens pluriserialis, setis scabridis subinæqualibus cæterum inter se similibus. — Herba perennes pleræque ex Americâ boreali, rariùs ex orbe veteri aut ex Amer. austr. ortæ, interdum suffruticosæ aut scaposæ. Folia alterna simplicia integra aut dentata. Capitula solitaria aut plurima corymbosa seu paniculata. Discus flavus demùm interdum purpurascens. Radius albus cœruleus purpureusve.

**Ser. I. ALPIGENI** Nees ast. p. 24. — *Oritrophium* sect. *Asteris* Kunth nov. gen. am. 4. p. 91.

Invol. squamæ herbacæ angustæ longitudine subæquales. Recept. nudum scrobiculatum. Pappus pauciserialis. Achenia sæpiùs hirta. Stigmata brevia triangularia. — Caules mono- aut oligocephali, sæpiùs hirsuti. Capitula majuscula.

1. *A.?* *NOLOZARICUS* (Forst. prod. p. 56. n. 295), foliis subtis albo-tomentosis, radicalibus oblongo-lanceolatis angustè serratis, caulinis linearibus integerrimis, caule 1-cephalo compresso, invol. squamis lanceolato-linearibus laxis. 2. in Novâ-Zelandiâ ad rupes muscosas in Duskybay. Nees ast. p. 25. Rich. et Less. fl. nov. Zeel. p. 248. Capitulum a poll. diam. Invol. squamæ non nudi

glutinosæ et lanuginosæ discum sequentes. Ligulæ pollicares lineæ angustiores purpureæ. Pappus ferrugineus.

2. *A. ? CORIACEUS* (Forst. prod. p. 56. n. 297), foliis radicalibus lato-lanceolatis sessilibus coriaceis integerrimis subplicatis subtus candido-pellitis, caule lanato 1-cephalo supernè squamato, invol. squamis laxis lanceolatis longissimè acuminatis apice pilosis.  $\mathcal{L}$  in Novâ-Zeelandiâ. Spreng. syst. 3. p. 528. Nees ast. p. 26. n. 3. Valdè affinis *A. holosericeo* et cum eo subsectionem ferè formans ex Nees. An uterque ad *Euribiam* referendus?
3. *A. ALPINUS* (Linn. sp. 1226), caule 1-cephalo, foliis radicalibus apathulato-oblongis caulinisque integerrimis lanceolatis pilosis, invol. squamis lanceolatis laxis ciliatis dorso glabriusculis obtusis aut acutiusculis disco æqualibus, acheniis hirsutis.  $\mathcal{L}$  frequens in pratis apricis montium Pyrenæorum etiam Hispanorum, Jurassi, Alpium omnium, Caucasi, Sibiriz in pratis montis Karagiol prov. Karabagh (Szow. ex Fisch. et Mey.). Nees ast. p. 26. n. 4. Jacq. fl. austr. t. 88. Sturm deutschl. fl. ic. Ligulæ cœruleæ lineam latè obtusiusculæ 6-8 lin. longæ. Var. glabriusculus, pilosus aut hirsutus (*A. hirsutus* Host fl. austr. 2. p. 485). (v. v. et s.)
  - $\beta$ . *brachyglussus*, humilis, foliis linearibus, radicalibus vix apice latioribus, ligulis brevissimis discum non superantibus.  $\mathcal{L}$  in Vallisæ Alpe excelâ dictâ Fluve ad glacies æternas. Habitus ab *A. alpino* diversissimus, sed post iterata examina ad hanc speciem reduxi. (v. s. comm. à cl. Phil. Duman.)
  - $\gamma$ . *nivalis*, foliis angustioribus magis canescentibus, capitulo minore, ligulis angustioribus, invol. squamis canescentibus brevibus obtusis apice purpurascens.  $\mathcal{L}$  in Sibiria orient. *A. nivalis* Adams in Web. et Mohr beytr. 1. p. 67. *A. pulchellus* quorumd. Conf. Nees ast. p. 27.  $\beta$ . excl. secundo Willd. syn.
4. *A. ARGUNNENSIS*, caule 1-cephalo subnudo sparse villosa, foliis radicalibus oblongis basi angustatis obtusiusculis integerrimis villosis, caulinis paucis oblongis acuminatis, invol. squamis biserialibus dorso lanuginosis acuminatis, exter. lanceolatis, int. linearibus, acheniis glabris.  $\mathcal{L}$  in Dahuriâ ad flum. Argunnum legit cl. Razgildew. Habitus ferè Hieracii. Invol. purpurascens. Ligulæ lineares angustæ 9 lin. longæ, lineam latæ. Involucrum *Erig.* uniflori Linn. Ligulæ *Asteris alpini*. (v. s. comm. à cl. Turczaninow.)
5. *A. ELLENIFOLIUS* (Turcz. in litt. 1834), glabriusculus, caule erecto simplici 1-cephalo apice subnudo, foliis infer. oblongo-linearibus basi attenuatis, super. linearibus attenuatis, omnibus integerrimis 1-nerviis, invol. squamia linearibus acutis æqualibus puberulis, ligulis linearibus disco duplò longioribus.  $\mathcal{L}$  in Sibiria orient. inter Ochotiam et Saculiam legit cl. Turczaninow. Herba semipedalis ad *A. pulchellum* habitu paulò accedens. Ligulæ ex sicco vid. cœruleæ. (v. s. comm. à cl. inv.)
6. *A. GAYANUS*, pube setulosâ adpressâ scabriusculus, caule erecto apice subcorymboso, ramis nudiusculis 1-cephalis, foliis oblongo-linearibus acutis sessilibus integerrimis, invol. squamis linearibus acutis disco æqualibus pauciserialibus dorso piloso-scabris, ligulis linearibus disco duplò longioribus.  $\mathcal{L}$  in excelsis Andibus Chilensium legit indef. Cl. Gay. Habeo etiam ex itin. Nesi sine loci designatione. Ligulæ ex sicco videntur albæ aut pallidè purpurascens. Achenia juniora vix pilosiuscula. Variat ramis laxè racemosis ramulis 1-cephalis et ramis simplicibus necnon caulibus foliisque glabriusculis aut scabris. (v. s.)
7. *A. GRAMINIFOLIUS* (Pursh fl. am. sept. 2. p. 545), glaber, caule surperne ramo, foliis angusto-linearibus enerviis integerrimis erectis ramulis nudiusculis 1-cephalis, invol. squamis laxis discum æquantibus.  $\mathcal{L}$  ad sinum Hudsonis (Pursh) in Canadâ (Lindl.) Nees ast. 277, sed ad *Asteres alpinos* nec ad *Tripolia* referendus ex Lindley adn. mes. Ligulæ pallidè purpureæ.
8. *A. SARANS* (H. B. et Kunth nov. gen. 4. p. 90. t. 33a bis), rhizomate repente, foliis radicalibus lanceolatis obtusis integerrimis reticulato-venosis coriaceis

- glabris, caulibus minutis, caule scapiformi erecto 1-cephalo, invol. squamis lanceolato-acuminatis laxè imbricatis.  $\frac{1}{2}$  in frondosis circa urbem Loxa Novo-Granatensium. Nees ast. 30. Ligulæ albæ. Rhizoma lanuginosum. Habitus Bellidis.
9. *A. caocivoltus* (H. B. et Kunth nov. gen. am. 4. p. 89. t. 332. n. 1), foliis radicalibus angustè linearibus uninerviis integerrimis margine revolutis subtus cano-tomentosis coriaceis, caulibus linearibus multò minoribus, caule scapiformi 1-cephalo, invol. squamis laxè imbricatis lanceolato-linearibus lanatociliatis.  $\frac{1}{2}$  in frondosis circa Loxam Novo-Granat. Nees ast. p. 31. Ligulæ albæ. Habitus Bellidis.
10. *A. pellitus* (H. B. et Kunth nov. gen. am. 4. p. 91), caule erecto simplici 1-cephalo basi lanato, foliis lineari-lanceolatis acutis integerrimis margine revolutis supra glabris subtus albido-lanatis radicalibus petiolatis, invol. lanatis squamis linearibus obtusis imbricatis.  $\frac{1}{2}$  in montosis Andium Novo-Granatensium locis uliginosis. Nees ast. p. 31. Herba 6 pollic. Capitulum magn. Inula britannicæ. Ligulæ albæ. Achænia pubentia. Pappus rufescens.
11. *A. pygæus* (Lindl. adn. mas.), caule villosa 1-cephalo, foliis ovato-oblongis obtusis integerrimis glabris, invol. hemisphærici squamis laxis villosissimis linearibus obtusis.  $\frac{1}{2}$  in Americâ arcticâ. Caules biunciales. Capitula maxima. (Lindl.)
12. *A. californicus* (Less. in linnæâ 1831. p. 33), caule ascendente oligocephalo, foliis inferioribus spatulatis crenato-serratis margine pubescentibus, caulibus lingulatis integerrimis, invol. subæqualis squamis lineari subulatis hirsutis.  $\frac{1}{2}$  in Californiâ. Nees ast. p. 33. Herba facie Bellidis. Ligulæ purpureæ.
13. *A. japonicus* (Less. in Nees ast. p. 34), caule erecto 1-cephalo superne nudo magis piloso, foliis elliptico-acuminatis integerrimis pilosis multinerviis, invol. pauciserialis squamis lineari-acuminatis hirsutis, exterioribus subquarrosis.  $\frac{1}{2}$  in Japoniâ. Inula dubia Thunb. fl. jap. 318. Rhizoma repens? Caulis teres striatus. Capitulum A. Sibirici. Antheræ ecaudatæ. Ovaria compressa hirsuta. Ligulæ cœruleæ?
14. *A. Unalascensis* (Less. in linnæâ 1831. p. 122), caule erecto 1-3-cephalo supernè sparè villosa folioso, foliis infimis obovatis obtusis, cæteris lanceolatis ovato lanceolatis acuminatis subserratis pubescenti-scabris ciliatis, invol. subæqualis squamis lineari-lanceolatis herbaceis villosis.  $\frac{1}{2}$  in insulâ Unalascâ (Less.) et in Amer. bor. montibus scopulosis (Lindl. mas.). Nees ast. p. 34. (v. s. ex comm. ab Acad. imp. Petrop.)
15. *A. poliacus* (Lindl. adn. rus.), caule simplici? racemoso, foliis oblongo-lanceolatis amplexicaulibus subserratis glabris, capitulis ramulos axillares terminantibus, foliis caulibus imò ramealibus superatis, invol. squamis patentibus foliaceis glabris.  $\frac{1}{2}$  in Unalascâ (Fisch.). Species notabilis involucri squamis folia ramealia referentibus, pedunculo tomentoso. — in specim. meo etiam ex Unalascâ à Fischero misso, invol. squamæ erectæ nec patentēs, ciliatæ nec omninò glabræ, folia basi obtusa, caules pilosiusculi, pilis hinc indè subserratis. Fortè *A. peregrinus* Less. in linnæâ 1831. p. 123 huc excl. syn. referendus? (v. s.)
16. *A. parviflorus* (Pursh fl. bor. am. 2. p. 556. non Less.), caule erecto subsimplici glabro supernè hirsuto summitate 1-3-floro, foliis sessilibus remotis oblongo-lanceolatis sensum attenuatis glabris, invol. squamis lanceolato-linearibus villosis acutissimis laxis longitudine disci, lignis numerosis.  $\frac{1}{2}$  ad Oros occid. Americæ borealis et in insulâ Unalascâ. Nees ast. 33. Ast. Tillæi Wikstr. in act. holm. 1822. p. 13. Pili caulibus ex Less. in linnæâ 1831. p. 123? seriatis sæpius trifariam dispositi. Pappus flocculi disci æqualis ex Nees, sed valde dubium mihi videtur an stirps in Unalascâ crescens eadem ac Americânâ?
17. *A. Eschschæensis* (Nees ast. p. 36. n. 19), caule simplici vel subcorymbose hirsuto basi repente ascendente, foliis oblongo-lanceolatis acutis argutè mucronato-serratis subtus villosis, invol. squamis subæqualibus lanceolatis acumi-

natis patulis villosis, achæniis elongatis.  $\frac{1}{2}$  ad unum Elsholtzii in prom. Es-  
penberg et Kotzebue Sund legit cl. Chamisso. Aster salsuginosus? Cham. ! in  
litt. Less. in Linnæa 1831. p. 124. Capit. A. Alpini sed squamæ inv. magis  
foliaceæ et patulæ. (v. s.)

$\beta$ . ? *dubius*, invol. squamis lanceolatis æqualibus erectis villosis.  $\frac{1}{2}$  in Una-  
laschkâ. A. peregrinus herb. Acad. imp. Petrop. ! sed non videtur A. pere-  
grinus Pursh ob caulem hirsutum nec seriatim pilosum. (v. s.)

18. A. *Pyrenæus* (DC. fl. fr. n. 3139), caule erecto hispido striato rarò simplici  
1-cephalo sæpius apice ramoso corymboso, ramis 1-cephalis foliosis, foliis  
lanceolatis semiaplexicaulibus acuminatis apice remotè serratis utrinquè  
hispido-scabris, invol. squamis laxi lanceolato-linearibus acuminatis dorso  
pubero-scabris.  $\frac{1}{2}$  in Pyrenæis montibus (Tourn.), ad montem Esquierr  
(Lapeyr.), nec certissimè in Sibiria (Lindl.). Desf. ! cat. h. par. 102. Lapeyr.  
abr. 514. Ast. Sibiricus Lam. dict. 1. p. 305 et pler. hort. bot. non Linn. Ast.  
Sibiricus var.  $\alpha$  et  $\gamma$  Nees ast. p. 37. ex syn. Ast. Pyrenæus, etc. Tourn. iust.  
482. n. 3. Jam à tempore Tourn. in hort. bot. Paris. cultus. (v. v. o.)
19. A. *Quitensis* (Willd. in Spreng. syst. 3. p. 533), caule sulcato villosa dupli-  
cato-racemoso oligocephalo, foliis cordato-oblongis amplexicaulibus basi api-  
ceque denticulatis pubescenti-hirtis, invol. laxi æqualis squamis 3-serialibus  
lineari-subulatis discum superantibus, achæniis pubescentibus.  $\frac{1}{2}$  in regno  
Quitensi Humb. et Bonpl. Nees ast. p. 38.

## Ser. II. AMELLI Nees ast. 39.

Invol. squamæ ciliatæ squarrosæ, exteriores herbaceæ, interiores totæ  
vel apice saltem membranaceæ coloratæ. Recept. alveolatum marginibus  
licero-dentatis. Achænia hirta rariùs glabra. Stigmata apice lanceolata. —  
Caulis corymbosus aut racemosus rigens. Folia scabrida subintegra. Capitu-  
la ampla.

20. A. *ISCUTIANUS*, caule erecto tereti sparsè hirsuto 1-5-cephalo sub capitulis  
tomentoso-hirsuto vix incrassato, foliis lanceolatis utrinquè acuminatis à me-  
dio ad apicem serratis suprâ glabris subtùs puberalis, invol. squamis lineari-  
lanceolatis subhirsutulis acutis nunc erectis nunc apice patente squarrosis  
radio imò disco brevioribus, ligulis circ. 20, achæniis pilosis.  $\frac{1}{2}$  in sabulosis  
Dahuriæ ad ripam fl. Iræuti legit cl. Turczaninow. Aster conspicuus Turcz. ! in  
litt. 1830 non Fisch. Ligulæ purpureæ 4-5 lin. longæ, lineam latæ. Affinis  
A. Richardsonii sed ex hujus descr. facilè distinctus videtur. (v. comm. à cl.  
Turczaninow.)
21. A. *RICHARDSONII* (Spreng. syst. 3. p. 528. Nees ast. 30), caule erecto ramoso  
1-2-cephalo sulcato villosa-canescente sub capitulis incrassato tomentoso,  
foliis latè oblongis remotè dentatis subtùs breviter pilosis, invol. squamis lineari-  
lanceolatis squarroso-patulis villosis, interioribus radium subæquantibus, li-  
gulis circ. 30 angustis, achæniis pilosis. — in Amer. borealis desertis à Point  
Lake ad mare Polare (Richards.) et in Sibiria (Br. ex h. Pall.). Aster montanus  
Brown in Frankl. it. p. 749. non Nutt. A. Ammanni Lindl. adn. mss. ipso doc.  
in litt. 1835. Folia sesquipollicem longa. Radix repens. Ligulæ ex Nees albæ ?
22. A. *SALSUGINOSUS* (Richards. ex Brown fl. reg. pol. in Frankl. it. p. 749.  
app. ed. 2. p. 32), caulibus simplicibus 1- (seu oligo-) cephalis, foliis lineari-  
obovatis acutis subintegerrimis venosis, invol. squamis laxè imbricatis lineari-  
bus acutis disco ferè duplò brevioribus, achæniis hirtis.  $\frac{1}{2}$  in regione sylvestri  
inter 54 et 64° lat. bor. Americæ sept. Hook. in bot. mag. t. 2942. Nees ast. 29.  
Ligulæ purpureæ lineares 6-8 lin. longæ sæpè emarginatæ. Pappus scaber sub-  
fuscescens. Species invol. brevissimo præsertim distincta.
23. A. *HYLORUS* (Michx. fl. bor. am. 2. p. 114), caule glabro sæpius dicephalo,  
foliis lanceolatis serratis scabris, invol. squamis arcuè imbricatis ovatis.  $\frac{1}{2}$  in  
Labradoriâ ad sinum Hudsonianum et in Pensylvaniæ montibus altissimis. Ast.



- biflorus Nees syn. p. 15. ast. p. 39. *Aster strictus* Pursh fl. 2. p. 556. Nutt. gen. am. 2. p. 157. Mey. labr. p. 62. Ligulæ pallide lilacine.
24. *A. suaviculosus* (Michx. fl. bor. am. 2. p. 113), caule simplici subpubescente laxo apice oligocephalo, foliis lineari-lanceolatis acuminatis setaceo-serrulatis margine scabris, infer. in petiolum attenuatis, super. vaginanti-amplexicaulibus, invol. squamis cuneiformi-spathulatis mucronatis apice patulis ramulique pubescentibus.  $\frac{1}{2}$  in sylvis Carolinæ septentr. (Michx.), in marginibus pratorum paludosorum in Tennessee et Virginia (Nutt.). Nees ast. 40. Radix stolonifera. Capit. A. smelli. Ligulæ violaceæ. Pappus rufescens.
- $\beta$ . Foliis obovato-lanceolatis. — in pinetis humidis Carolinæ circa Parysburg. Ast. surculosus Ell. sketch 2. p. 352. Ast. liatroides Muhl. An i var. a distinctus?
25. *A. conspicuus* (Lindl. adn. mss.), caule stricto apice corymboso, ramis erectis pubescentibus 1-cephalis aphyllis, foliis oblongis acutis basi angustatis grosse serratis pubescentibus ambitu scabris, invol. hemispharici squamis squarrosis acuminatis, infer. minoribus.  $\frac{1}{2}$  in America arctica. *A. conspicuus* Lindl. adn. mss. 1833 non Fisch. nec Turcz. Ad *A. spectabilem* accedit, sed foliis latius ferè omnibus grosse serratis faciliè distinguendus. Radix corulea. (Lindl. mss.)
26. *A. spectabilis* (Ait. kew. 3. p. 209), caule stricto apice corymboso, ramis rigidis incurvis corymbosis, foliis oblongo-lanceolatis acutis scabris cæcilibus inferioribus serratis, invol. squamis obtusis copiosissimis squarrosis ramulique glanduloso-pubescentibus.  $\frac{1}{2}$  in Amer. bor. (Ait.), in Novâ-Cæsareâ vulgaris (Nutt.). Nees ast. p. 41. Lindl. bot. reg. t. 1527. *A. elegans* Willd. sp. 3. p. 2048 ex parte et *A. spectabilis* Willd. sp. 3. p. 2048 excl. syn. ex Nees. An *A. speciosus* Horn. h. hafn. 2. p. 816? Ligulæ coruleo-purpureæ. (v. v. c.)
- $\beta$ . *bellidifolius* (Nutt. gen. am. 2. p. 157), foliis oblongo-obovatis serratis corymbo subsimplici ramulis 1-cephalis.  $\frac{1}{2}$  in Novâ-Cæsareâ.
27. *A. RADULA* (Ait. kew. ed. 1. v. 3. p. 210. non Less.), caule angulato glabriusculo corymboso, ramis oligocephalis nudiusculis, foliis lanceolatis utrinque attenuatis mediè argutè serratis rugosis scaberrimis, invol. squamis pluri-fariam imbricatis oblongis acutiusculis apice patulis.  $\frac{1}{2}$  in Novâ-Scotiâ (Ait.), montibus altioribus Novi-Eboraci et Pennsylvania (Pursh). Nees ast. p. 43 Lindl. pl. exs. Ligulæ 5-nerviæ ex Pursh albæ, ex Nees pallide lilacine. — Ast. nudiflorus Nutt. gen. 2. p. 157 fortè ex Nees non differt? (v. s.)
28. *A. nudiflorus* (Nutt. gen. am. 2. p. 157), caule simplici glabro, foliis sessilibus ovato-lanceolatis subacuminatis et argutè serratis, corymbo simplici oligocephalo, pedunculis pubescentibus nudis sub-1-cephalis, invol. hemispharici disco brevioris squamis densè imbricatis lineari-oblongis ciliatis.  $\frac{1}{2}$  in paludibus Novæ-Cæsareæ rarior. Capitula magna, ligula pallide violaceis. Affinis *A. radulæ* et ab eo vix distinctus ex Nees, sed patria diversissima.
29. *A. LIMA* (Lindl. adn. mss.), caule scabro virgato sub-1-cephalo, foliis linearibus scaberrimis acutis ciliatis sensim in involucrium decreascentibus, invol. squamis linearibus rigidis acutis acutè imbricatis basi albidis.  $\frac{1}{2}$  in Mexico ex herb. Bentham. Radius purpureus. Habitus Diplopappi linearifolii sed pappus involucrium et achenium *Asteris* Amelli. An frustula descripta sint caules aut rami? (Lindl. mss.)
30. *A. TATARICUS* (Linn. f. suppl. 373), caule erecto striato hispido apice ramoso-corymboso, ramis subnudis, foliis lanceolatis seu ovali-lanceolatis basi attenuatis integerrimis apice apicibus grosse serratis acuminatis superne glabriusculis subtus hispidis summis linearibus integerrimis parvis, invol. squamis linearibus glabriusculis subæqualibus margine et apice purpurascens.  $\frac{1}{2}$  in Sibiria à Jeniseo fluviò ad Kamtschatkam (Gmel.), in montosis apricis Dahuriæ (Turcz.). *A. rhomboides* Lindl. in adn. mss. 1833 et ad *A. Tataricum* rel. anno 1835. Gmel. sib. 2. p. 186. t. 80. f. 1? *A. Sibiricus*  $\beta$  Nees ast. p. 37. An *Grindelia Sibirica* Link. enum. 2. p. 336? Ligulæ coruleæ aut albæ? (v. s. comm. à el. Turcz.)

31. *A. SPATHULATUS* (Lindl. adn. ms.), caule simpliciter ramoso, foliis spathulatis obtusis integerrimis ramorum basi dilatatis amplexicaulis, pedunculis subaphyllis foliis brevioribus, invol. squamis linearibus acutis erectis tomentosis.  $\frac{1}{2}$  in America arctica. Facies *Ast. pulchelli* (Lindl.)
32. *A. ASCENDENS* (Lindl. adn. ms.), caulibus ascendentibus simpliciter racemosis vel corymbosis, foliis oblongo-linearibus subspathulatis elongatis integerrimis glabris margine cillato-scabris, involucri glabri squamis numerosis acutis patulis.  $\frac{1}{2}$  in America arctica. (Lindl. ms.)
33. *A. SERRICUS* (Linn. sp. 1226), caule pubescente apice corymboso, ramis simplicibus 1-cephalis vel compositis apice incanis, foliis oblongis grosse serratis pubescentibus, superioribus sessilibus, involucri squamis linearibus acuminatis, omnibus imbricatis squarrosis.  $\frac{1}{2}$  in Sibiria (Prescott.) *Aster inuloides* Fisch. nec Don. *A. lacerus* Lindl. adn. ms. 1833 et ad *A. Sibiricum* redactus 1835. — Gmel. sib. 2. p. 186. t. 80. f. 1? Species pulchra, invol. squamis interioribus purpureis squarrosis, radio verosimiliter albo. (Lindl. ms.)
34. *A. PRESCOTTII* (Lindl. adn. ms. 1835), caule simplici scabro adscendente 1-2-cephalo apice incano, foliis oblongo-lanceolatis argutè serratis basi angustatis scabris, superioribus sessilibus integerrimis, invol. squamis linearibus acuminatis, omnibus imbricatis squarrosis.  $\frac{1}{2}$  in Dahuria prope Irkutsk (Prescott.). *Aster pauciflorus* Lindl. adn. ms. 1833 non Nutt. *Ast. biflorus* Presc. nec Michx. *Asteri lacero* proximus, foliis inflorescentiâ radioque purpureo diversus. (v. s. comm. à el. Lindl.)
35. *A. AMELLUS* (Linn. sp. 1226), caule erecto 1-cephalo aut sæpius simpliciter corymboso, foliis oblongo-lanceolatis acutis subserratis subtriplinerviis pubescenti-scabris, invol. squamis 4-5-seriatis oblongis patulis obtusis aut vix acutis apice membranaceis.  $\frac{1}{2}$  in Europæ mediæ et australis Asiæque occid. siccis. Nees ast. p. 44. Adest ex Lindl. adn. ms. in herb. Linn. sub nominibus *A. Amelli* et *A. Sibirici* var. Variat 1 aut polycephalus, caule breviter scabro aut hispido, foliis infer. obtusis aut mucronatis, etc. Præcipuæ varietates aut fortè species distinguendæ sunt:
- a. *communis*, foliis plerisque integerrimis obtusis cum mucrone, invol. squamis obtusis.  $\frac{1}{2}$  *A. Amellus* Jacq. fl. austr. t. 435. DC. fl. fr. n. 3136. *A. amelloides* Roem. arch. 2. p. 298. Wender. ind. sem. h. marb. app. n. 3? Koch! et Ziz obs. *A. elegans* Nees syn. 20. (v. v. sp.)
  - ß. *latifolius*, foliis inf. spathulatis obovatis obtusis, invol. squamis obtusis ciliatis.  $\frac{1}{2}$  *A. Amellus* bot. reg. t. 340. *A. amelloides* Reichenb. fl. exc. 246. *A. pseudo-amellus* DC. fl. fr. suppl. n. 3136 in adn. (v. v. c.)
  - γ. *hispidus*, foliis infer. oblongis basi attenuatis cauleque patentim hispidis. — in Thuringiâ ex Wallroth. (v. s.)
  - δ. *Bessarabicus*, foliis oblongis basi attenuatis utrinque cauleque breviter scaberrimis, invol. squamis subobtusis. — *Ast. amelloides* Bess.! enum. p. 33 et ic. 8. *Ast. Bessarabicus* Bernh. ex Reichenb. fl. exc. 246. (v. s.)
  - ε. *Ibericus*, foliis integerrimis acutis, invol. squamis interioribus vix coloratis. *Ast. Ibericus* Stev. in Bieb. fl. n. 1732.
36. *A. MONSTROSUS* (Lindl. adn. ms.), caule simplici folioso, corymbo paucifloro foliis supremis brevioribus, foliis lanceolatis acuminatis grosse serratis glabris, invol. squamis linearibus acuminatis.  $\frac{1}{2}$  in America arctica. *A. brumalem* nomen refert sed affinitate ad *Ast. Novæ-Angliæ* proximè accedit. (Lindl. ms.)
37. *A. NOVÆ-ANGLIÆ* (Ait. kew. 3. p. 201. Nees ast. 46), caule stricto corymboso-composito, ramis arcuatis, foliis lanceolatis attenuatis auriculato-amplexantibus integerrimis hispido-scabris, invol. squamis laxis angustis.  $\frac{1}{2}$  in America. ad rivulorum ripas et margines à Canada ad Carolinam. *Ast. altissimus* Mæench. Ligulæ purpureo-violaceæ. Achænia hirta.
- a. *agrestis* (Nees l. c.), caule humiliori polycephalo, ramulis elongatis, invol. squamis lanceolatis venoso-trinerviis.  $\frac{1}{2}$  *Ast. spurius* Willd. ap. n. p. 2032. *Ast. amplexicaulis* Lam. dict. 1. p. 304. non Willd. nec Michx.

- A. Nova-Angliæ Ell. sketch 2. p. 35r. Lindl.! pl. exs. A. grandiflorus hortul. (v. s.)
- β. *hortensis* (Nees 1. c.), caule altiori oligocephalo, ramulis brevibus ad apicem confertis, invol. squamis lineari-lanceolatis. ♀ Aster Nova-Angliæ Linn. sp. 1229. Lindl.! bot. reg. t. 183. — Asteripholis Pont. dia. 224. — Volk. fl. norib. p. 50. ic. Knorr del. 1. p. 5. 2. Herm. parad. Lond. b. p. 98. ic. Variat in hortis caule purpureo subviscido, etc. (v. v. c.)
- γ. *roseus*, ligulis roseis. Ast. roseus Desf.! cat. h. par. ed. 3. p. 401. non Stev. — bot. reg. t. 183. f. d. (v. v. c.)
38. A. CAROLINIANUS (Walt. car. 208), caule suffruticoso flexuoso ramosisimo pubescente, foliis oblongis acuminatis subintegerrimis in petiolam brevem auriculato-amplexicaulem contractis pubescenti-scabris, ramis apice simpliciter racemosis, invol. squamis apice reflexis, acheniis levibus 10-striatis. ♀ in dumetis umbrosis et paludibus Carolinæ et Georgiæ. Nees ast. p. 48. Michx. fl. bor. am. 2. p. 111. Ell. sketch 2. p. 353. Ast. scandens Michx.! herb. L'Her. Jacq. f. in Spreng. syst. 3. p. 524. Ligula late purpurea roseæ aut carnea. Achenia pubescentia aut adulta glabra. Folia dicuntur integerrima, sed in spec. meo hinc indè acutè serrulata. (v. s. sine flor.)
39. A. OBLONGIFOLIUS (Nutt.! gen. am. 2. p. 156), caule herbaceo scabro divaricato-ramoso, ramis subcorymbosis 1-cephalis, foliis lineari-oblongis subamplexicaulibus acutis integerrimis scabris, invol. squamis lineari-oblongis foliaceis squarrosis, acheniis brevè villosis. ♀ in Amer. bor. ad flum. Mississippi Nees ast. p. 48. A. graveolens Nutt.! in Journ. ac. nat. sc. philad. 1821. oct. p. 116. in Linnæa 1829. p. 39.! Ramuli fastigiati. Ligula pallide purpurea. Herba viscosa graveolens. (v. s. comm. à cl. Nutt.)
40. A. PATENS (Ait. kew. 3. p. 201), caule ramoso hirtio, ramis elongatis oligocephalis, foliis oblongo-ovatis integerrimis cordatis amplexicaulibus scabris, pedunculis 1-cephalis subnullis, invol. squamis imbricatis lanceolatis patulis. ♀ in Virginia et Pensylvaniâ Willd. sp. 3. p. 2034, in sylvis circa Bostoniam (Big.). A. patens & Nees ast. 49. excl. Michx. syn. Ast. amplexicaulis Michx. fl. 2. p. 114. Big.! fl. hort. ed. 2. p. 312. Ast. amplexicaulis Michx. est omnino A. patens ex Lindl. A. undulatus Linn. sp. ed. 1. p. 1228 ex Sm. Ell.! sketch 2. p. 361. Folia subobtusè minute mucronulata basi non angustata, auriculis obtusis. Pappus rufus. Achenia sericeo-villosa. (v. s. sp.)
41. A. PHLOGIFOLIUS (Muhl. in Willd. sp. 3. p. 2034), caule simplicissimo pubescente apice paniculato, foliis lanceolatis integerrimis utrinquè attenuatis infernè dilatatis cordato-amplexicaulibus subtus pubescentibus, invol. squamis laxis imbricatis lanceolatis, acheniis pilosiusculis. ♀ in Pensylvaniâ (Muhl.) à Novâ-Cæsareâ ad Virginiam (Pursh?). Ast. patens β. Nees ast. p. 49. excl. syn. Michx. sed ab A. patente satis diversus mihi videtur ex foliis ad A. Carolinianum accedentibus, molliter pubescentibus nec scabris, achenio multò minùs villosò, capitalis majoribus. (v. s. et v. c.)
42. A. PATENTISSIMUS (Lindl. adn. mss.), caule racemoso-composito, ramis longissimis foliosis, foliis ovato-oblongis cordatis amplexicaulibus suprâ glabris subtus pilosiusculis, pedicellis brevibus, involucri turbinati lanati squamis obtusis. ♀ in Louisiana. Radli cyanei. A. patente proximus. (v. s. comm. à cl. Lindl.)
43. A. AURITUS (Lindl.! adn. mss.), caule racemoso-composito, ramis rigidis apice densè racemosis, foliis oblongis acuminatis basi cordatis amplexicaulibus suprâ levibus subtus pubescentibus, involucri parùm imbricati squamis linearibus acuminatis ramulique glandulosi. ♀ in America sept.? Cultus in hort. anglicis sed rarius. Radius discolor. Species A. patentissimo proxima. (v. s. comm. à cl. Lindl.)
44. A. GRANDIFLORUS (Linn. sp. 1231), caule erecto hispido racemoso, ramis elongatis 1-cephalis foliosis, foliis lanceolato-ligulatis mucronatis integerrimis hispidis reflexis, invol. squamis lineari-lanceolatis squarrosis, acheniis striguloso-hirtis. ♀ in sylvis siccis sabulosis à Virginia ad Carolinam (Pursh) bot.

reg. t. 273. Hoffm. phyt. t. 65. t. A. f. 1. Nees ast. p. 50. — Mill. ic. t. 282. Dill. eith. p. 41. t. 36. f. 41. Capitula ampla ligulis violaceis. (v. v. c.)

45. *A. sericeus* (Vent. h. Cels. n. et t. 33), basi suffruticulosus totus pube sericeâ adpressâ argenteus, ramis laxis apice 1-cephalis, foliis lanceolatis integerrimis, invol. squamis foliaceis imbricatis squarrosis argenteis, acheniis glabris.  $\frac{1}{2}$   $\frac{3}{4}$  ad ripas fluminum præruptas inter rupes in prov. Illinoensi et ad flum. Mississipi et Missouri. Nees ast. p. 51. Lindl. ! pl. exs. *Aster argenteus* Michx. fl. bor. am. 2. p. 111. Ligulæ cœruleæ. (v. v. c.)

$\beta$ . *microphyllus*, foliis capitalisque triplè minoribus.  $\frac{3}{4}$  in Mexici distr. orientalibus prov. Texas legit cl. Berlandier (pl. exs. n. 1876). An species propria? (v. v. c.)

46. *A. montanus* (Nutt. gen. am. 2. p. 156), caule erecto gracili, ramis laxis 1-cephalis, foliis lineari-lanceolatis integerrimis subsericeo-villosis obsoletè 3-nerviis, invol. hemisphaerici squamis imbricatis lineari-lanceolatis erectis. — in montibus distr. Tennassée et Carolinæ septentr. prope Ashville et Morganton rarior (Nutt.). Nees ast. p. 51. Valdè accedit ad *A. sericeum*. Ligulæ 2-5 violaceo-purpureæ.

### Ser. III. GENUINI *Nees ast. p. 52.*

Invol. squamæ imbricatæ vel laxæ, margine magis minùsve membranaceæ, membranacæ parte hasin versus incremente et totam ferè basin in iis quæ fortius sibi incumbunt occupante. Recept. alveolato-dentatum. Achenia glabra vel puberula. Pappus pluriserialis. — Caulis altior compositè corymboeus racemosus paniculatusve.

#### A. Heterophylli *Nees ast. p. 52.*

Folia radicalia seu inferiora petiolata cordata vel rariùs rotundata.

47. *A. urophyllus* (Lindl. adn. mss.), caule racemoso-paniculato, ramis thyrsoides, foliis ovato-lanceolatis acuminatissimis argutè crenato-serratis suprâ scaberrimis subtùs pilosis, invol. imbricati squamis subulatis.  $\frac{1}{2}$  in Louisianâ. Radii albi involucreo longiores. Discus purpureus. Species *A. hirtello* proxima. (Lindl. mss.)
48. *A. hirtellus* (Lindl. adn. mss.), caule racemoso-paniculato, ramis apice conferto-racemosis paucifoliis, foliis cordato-ovatis mediò crenato-serratis suprâ scaberrimis subtùs pilosis, involucreo laxè imbricato.  $\frac{1}{2}$  in Louisianâ. Radius verosimiliter lilacinus, discus purpureus. (Lindl. mss.)
49. *A. cordifolius* (Linn. sp. 1229. non Lam. nec Nutt.), caule erecto racemoso-paniculato subhirsuto, foliis petiolatis cordatis acuminatis argutè serratis suprâ scabris subtùs pilosis, ramis laxis paucifoliis, invol. squamis arcuè imbricatis.  $\frac{1}{2}$  in sylvis umbrosis montium à Canadâ ad Floridam. Nees! ast. p. 52. Lindl. ! bot. reg. 1597. *Ast. cordifolius* var. foliis subtùs pubescentibus Desf. ! in hort. Par. 1815. *Solidago cordifolia* Moench. Caulis rubens nunc ferè glaber nunc subhirsutus. Folia etiam superiora cordata, petiolis subhirsutis. Ligulæ pallidè purpureascentes aut albidæ. Achenia glabra seu ad costas setulosa. (v. v. c. et s. p.)
50. *A. paniculatus* (Ait. kew. ed. 1. v. 3. p. 207), caule glabro stricto ramisque thyrsoides foliosis, foliis petiolatis infer. cordatis summis, ovato-lanceolatis, omnibus acuminatis serratis suprâ scabris subtùs glabris, invol. squamis imbricatis,  $\frac{1}{2}$  in pratis, sæpius et sylvarum marginibus à Canadâ ad Carolinam. Nees ast. p. 53. Ell. sketch 2. p. 363. Lindl. ! pl. exs. *Ast. cordifolius* Lam. diet. 1. p. 307. *A. cordifolius* var. foliis subtùs glabris Desf. ! hort. Par. *Ast. heterophyllus* Nees syn. p. 15. non Willd. — Moris. ox. 3. a. 7. t. 22. f. 34. Ligulæ violacæ. (v. v. c.)
51. *A. heterophyllus* (Willd. sp. 3. p. 55), caule racemoso-decomposito patulo, foliis radicalibus cauliniisque infer. cordatis elatè petiolatis serratis apice

integerrimis suprà in ambitu totiusve scabris subtus glabris, supremis lanceolato-linearibus minutis, invol. acutè imbricato.  $\frac{1}{2}$  in fruticosis Pennsylvaniae Nees! ast. p. 55. Lindl. ! pl. exs. Ast. cordifolius Nutt. gen. am. a. p. 156. non Linn. A. diversifolius Spreng. syst. 3. p. 533. non Michx. (v. v. c.)

*ß. glabriusculus*, caule piloso-lineato, foliis glabris vel subpubescentibus. — A. pubescens Horn. h. hafn. suppl. 98. Nees! syn. p. 16. (v. v. c.)

$\gamma$ . *scaber*, caule hirsuto, foliis suprà scaberrimis, capitulis majoribus. An ex Nees species propria?

52. A. *sagittifolius* (Wedem. in Willd. sp. 3. p. 2055), caule glabro racemosocomposito, foliis radicalibus petiolatis cordato-subcuneatis (vel et profunde cordatis) oblongis glabris inæqualiter serratis margine scabris, caulinis ovato-oblongis in petiolum alatum contractis serratis glabris, involucri arcuè imbricato.  $\frac{1}{2}$  in Americæ bor. (Wed.), distr. superioribus utriusque Carolinae (Ell. sketch 2. p. 362). Ligulae cœruleæ (Willd.), lilacinæ (Nees ast. p. 56).

53. A. *Daumenoides* (Lindl. adn. mas.), totus incanus, caule ramisque racemoso-thyrsoideis strictis, foliis cordato-ovatis crenato-serratis, supremis sessilibus, ramulorum linearibus oblongis acuminatis, invol. squamis subulatis.  $\frac{1}{2}$  in Louisiana. Inter A. paniculatum et undulatum quasi medius, canitie facillimè distinguendus. Radii verosimiliter cyanei. (Lindl. mas.)

54. A. *diversifolius* (Michx. fl. bor. am. 2. p. 113), caule puberulo racemoso-thyrsoideo, ramulis virgatis apice secundifloris, foliis radicalibus longè petiolatis cordatis crenato-serratis acutis pubescentibus margine scabris, petiolo dilatato basi amplectente, superioribus sessilibus ovato-lanceolatis ramulorum subulatis, invol. arcuè imbricato.  $\frac{1}{2}$  in Amer. bor. à Canadâ ad Carolinam, in Novâ-Angliâ (Pursh), in siccis Carolinae frequens (Ell. sketch 2. p. 361), in Canadâ (Michx.), circa Bostoniam (Big. fl. boat. ed. 2. p. 312. Lindl. adn. mas. A. undulatus Linn. sp. ed. 2. p. 1228. Nees ast. p. 57. A. scaber Ell. sketch 2. p. 363. non Thunb. Ligulae cœruleæ. (v. s. sp.)

### B. Homophylli Nees ast. p. 58.

Folia radicalia basi acuta magis minùve attenuata.

#### \* Petiolati Nees ast. p. 58.

Folia latiuscula serrata, radicalia et caulina inferiora in petiolum attenuata, summa basi angustiore sessilia.

55. A. *parvus* (Lam. dict. 1. p. 308), caule glabro racemoso-paniculato, ramis patulis, foliis oblongis profunde serratis in petiolum attenuatis suprà pubescentibus subtus glabris margine scabris, invol. laxè imbricato.  $\frac{1}{2}$  in pratis sepius sylvarumque marginibus à Canadâ ad Virginiam usquè. Desf. ! cat. h. par. 102. A. paniculatus Willd. sp. 3. p. 2035, ex parte Nees ast. p. 16. excl. syn. A. Cornuti Wendl. in Nees ast. p. 58. nec Mull. A. Tradescanti Hofm. phyt. bl. p. 68. t. d. f. 2. ex Nees. Ligula cœrulea. Nomen multò vetustius, nec recentius et verisim. decipiens, admisi. (v. v. c.)

*ß. Radio pallido, foliis glabriusculis. — A. pallens Willd. enumer. suppl. p. 58. Lindl. bot. reg. 1509. A. tardiflorus Linn. 1. p. 1231, hæc aut ad var. ß pertinet ex cl. Lindley.*

56. A. *abbreviatus* (Nees! syn. p. 16), caule erecto glabro seu serotimè piloso racemoso, ramis brevibus thyrsoideis simplicibusve, foliis infer. ovali-lanceolatis serratis secus petiolum adnato-decurrentibus suprà scabris subtus levibus, sup. oblongo-linearibus subintegerrimis, invol. squamis laxè imbricatis lanceolato-linearibus.  $\frac{1}{2}$  in Amer. bor. sed ex confus. syn. locos proprios non citare audeo. A. Cornuti Mull. ex Nees. A. acuminatus Nees Ast. p. 60. excl. syn. Variat ramis abbreviatis aut in plantâ cultâ elongatis. Ligulae pallide cœruleæ. (v. v. c. à cl. Nees ad hort. Genev. misa.)

57. A. *fruticulosus* (Muhl. in Willd. sp. 3. p. 2046), caule paniculato, ramis

- reg. t. 273. Hoffm. phyt. t. 65. t. A. f. 1. Nees ast. p. 50. — Mill. ic. t. 282. Dill. elth. p. 41. t. 36. f. 41. Capitula ampla ligulis violaceis. (v. v. c.)
45. *A. sericeus* (Vent. h. Cels. n. et t. 33), basi suffrutescens totus pube sericea adpressâ argentea, ramis laxis apice 1-cephalis, foliis lanceolatis integerrimis, invol. squamis foliaceis imbricatis squarrosis argenteis, acheniis glabris.  $\mathcal{L}$  3 ad ripas fluminum præruptas inter rupes in prov. Illinoensi et ad flum. Mississipi et Missouri. Nees ast. p. 51. Lindl. ! pl. exs. *Aster argenteus* Michx. fl. bor. am. 2. p. 111. Ligulæ coruleæ. (v. v. c.)
- $\beta$ . *microphyllus*, foliis capitalisque triplè minoribus.  $\mathcal{L}$  in Mexici distr. orientaliibus prov. Texas legit cl. Berlandier (pl. exs. n. 1876). An species propria? (v. a.)
46. *A. morranus* (Nutt. gen. am. 2. p. 156), caule erecto gracili, ramis laxis 1-cephalis, foliis lineari-lanceolatis integerrimis subsericeo-villosis obsolete 3-nerviis, invol. hemisphaerici squamis imbricatis lineari-lanceolatis erectis. — in montibus distr. Tennesæ et Carolinæ septentr. prope Ashville et Morganton rarior (Nutt.). Nees ast. p. 51. Valde accedit ad *A. sericeum*. Ligulæ 2-5 violaceo-purpureæ.

### Ser. III. GENUINI *Nees ast. p. 52.*

Invol. squamæ imbricatæ vel laxæ, margine magis minùse membranaceæ, membranaceâ parte basin versus incremente et totam ferè basin in iis quæ fortius sibi incumbunt occupante. Recept. alveolato-dentatum. Achenia glabra vel puberula. Pappus pluriserialis. — Caulis altior compositè corymbosus racemosus paniculatusve.

#### A. Heterophylli *Nees ast. p. 52.*

Folia radicalia seu inferiora petiolata cordata vel rariùs rotundata.

47. *A. urophyllus* (Lindl. adn. mss.), caule racemoso-paniculato, ramis thyrsoides, foliis ovato-lanceolatis acuminatissimis argutè crenato-serratis suprâ scaberrimis subtùs pilosis, invol. imbricati squamis subulatis.  $\mathcal{L}$  in Louisiana. Radii albi involucri longiores. Discus purpureus. Species *A. hirtello* proxima. (Lindl. mss.)
48. *A. hirtellus* (Lindl. adn. mss.), caule racemoso-paniculato, ramis apice conferto-racemosis paucifoliis, foliis cordato-ovatis mediò crenato-serratis suprâ scaberrimis subtùs pilosis, involucri laxè imbricato.  $\mathcal{L}$  in Louisiana. Radii verosimiliter lilacini, discus purpureus. (Lindl. mss.)
49. *A. cordifolius* (Linn. sp. 1229. non Lam. nec Nutt.), caule erecto racemoso-paniculato subhirsuto, foliis petiolatis cordatis acuminatis argutè serratis suprâ scabris subtùs pilosis, ramis laxis paucifoliis, invol. squamis arcè imbricatis.  $\mathcal{L}$  in sylvis umbrosis montium à Canadâ ad Floridam. Nees! ast. p. 52. Lindl. ! bot. reg. 1597. *Ast. cordifolius* var. foliis subtùs pubescentibus Desf. ! in hort. Par. 1815. *Solidago cordifolia* Mœnch. Caulis rubens nunc ferè glaber nunc subhirsutus. Folia etiam superiora cordata, petiolis subhirsutis. Ligulæ pallidè purpurascens ant albidæ. Achenia glabra seu ad costas setulosa. (v. v. c. et a. sp.)
50. *A. paniculatus* (Ait. kew. ed. 1. v. 3. p. 207), caule glabro stricto ramisque thyrsoides foliosis, foliis petiolatis infer. cordatis summis, ovato-lanceolatis, omnibus acuminatis serratis suprâ scabris subtùs glabris, invol. squamis imbricatis,  $\mathcal{L}$  in pratibus, sepibus et sylvarum marginibus à Canadâ ad Carolinam. Nees ast. p. 53. Ell. sketch 2. p. 363. Lindl. ! pl. exs. *Ast. cordifolius* Lam. dict. 1. p. 307. *A. cordifolius* var. foliis subtùs glabris Desf. ! hort. Par. *Ast. heterophyllus* Nees syn. p. 15. non Willd. — Moris. ox. 3. a. 7. t. 22. f. 34. Ligulæ violaceæ. (v. v. c.)
51. *A. heterophyllus* (Willd. sp. 3. p. 55), caule racemoso-decomposito patulo, foliis radicalibus caulinisque infer. cordatis alatè petiolatis serratis apice

integerrimis suprà in ambitu totiusve scabris subtus glabris, supremis lanceolato-linearibus minutis, invol. arcuè imbricato.  $\mathcal{L}$  in fraticosis Pennsylvaniae Nees! ast. p. 55. Lindl. ! pl. exs. Ast. cordifolius Nutt. gen. am. a. p. 156. non Linn. A. diversifolius Spreng. syst. 3. p. 533. non Michx. (v. v. c.)

$\beta$ . *glabriusculus*, caule piloso-lineato, foliis glabris vel subpubescentibus. — A. pubescens Horn. h. hafn. suppl. 98. Nees! syn. p. 16. (v. v. c.)

$\gamma$ . *scaber*, caule hirsuto, foliis suprà scaberrimis, capitulis majoribus. An ex Nees species propria?

52. A. *SAGITTIFOLIUS* (Wedem. in Willd. sp. 3. p. 2055), caule glabro racemoso-composito, foliis radicalibus petiolatis cordato-sagittatis (vel et profunde cordatis) oblongis glabris inæqualiter serratis margine scabris, caulinis ovato-oblongis in petiolum alatum contractis serratis glabris, involucre arcuè imbricato.  $\mathcal{L}$  in Americæ bor. (Wed.), distr. superioribus utriusque Carolinæ (Ell. sketch 2. p. 362). Ligulæ cœruleæ (Willd.), lilacinæ (Nees ast. p. 56).

53. A. *DRUMMONDII* (Lindl. edn. mas.), totus incanus, caule ramisque racemoso-thyrsoideis strictis, foliis cordato-ovatis crenato-serratis, supremis sessilibus, ramulorum lineari-oblongis acuminatis, invol. squamis subulatis.  $\mathcal{L}$  in Louisiana. Inter A. paniculatum et undulatum quasi medius, canitie facillimè distinguendus. Radii verosimiliter cyanei. (Lindl. mas.)

54. A. *DIVERSIFOLIUS* (Michx. fl. bor. am. 2. p. 113), caule puberulo racemoso-thyrsoideo, ramulis virgatis apice secundifloris, foliis radicalibus longè petiolatis cordatis crenato-serratis acutis pubescentibus margine scabris, petiolo dilatato basi amplectente, superioribus sessilibus ovato-lanceolatis ramulorum subulatis, invol. arcuè imbricato.  $\mathcal{L}$  in Amer. bor. à Canadâ ad Carolinam, in Novâ-Angliâ (Pursh), in siccis Carolinæ frequens (Ell. sketch 2. p. 361), in Canadâ (Michx.), circa Bostoniam (Big. fl. bost. ed. 2. p. 312. Lindl. adn. mas. A. undulatus Linn. sp. ed. 2. p. 1228. Nees ast. p. 57. A. scaber Ell. sketch 1 p. 363. non Thunb. Ligulæ cœruleæ. (v. s. sp.)

### B. Homophylli Nees ast. p. 58.

Folia radicalia basi acuta magis minùve attenuata.

#### \* Petiolati Nees ast. p. 58.

Folia latiuscula serrata, radicalia et caulina inferiora in petiolum attenuata, summa basi angustiore sessilia.

55. A. *PATULUS* (Lam. dict. 1. p. 308), caule glabro racemoso-paniculato, ramis patulis, foliis oblongis profundè serratis in petiolum attenuatis suprà parte minutissimâ aspersis glabrisve subtus glabris margine scabris, invol. laxè imbricato.  $\mathcal{L}$  in pratis sepius sylvarumque marginibus à Canadâ ad Virginiam usquè. Desf. ! cat. h. par. 102. A. paniculatus Willd. sp. 3. p. 2035, ex parte Nees ast. p. 16. excl. syn. A. Cornuti Wendl. in Nees ast. p. 58. nec Mall. A. Tradescanti Hofm. phyt. bl. p. 68. t. d. f. 2. ex Nees. Ligulæ cœruleæ. Nomen multò vetustius, nec recentius et verisim. decipiens, admisi. (v. v. c.)

$\beta$ . Radio pallido, foliis glabriusculis. — A. pallens Willd. enum. suppl. p. 58. Lindl. bot. reg. 1509. A. tardiflorus Linn. 1. p. 1231, hic ast. id var.  $\beta$  pertinet ex cl. Lindley.

56. A. *ABBRUVIATUS* (Nees! syn. p. 16), caule erecto glabro seu seriatim piloso racemoso, ramis brevibus thyrsoideis simplicibusve, foliis infer. ovali-lanceolatis serratis secus petiolum adnato-decurrentibus suprà scabris subtus levibus, sup. oblongo-lanceolatis subintegerrimis, invol. squamis laxè imbricatis lanceolato-linearibus.  $\mathcal{L}$  in Amer. bor. sed ex confus. syn. locos proprios non citare audeo. A. Cornuti Mull. ex Nees. A. acuminatus Nees Ast. p. 60. exd. syn. Variat ramis abbreviatis aut in plantâ cultâ elongatis. Ligulæ pallide cœruleæ. (v. v. c. à cl. Nees ad hort. Genev. miss.)

57. A. *FERNATOIDES* (Muhl. in Willd. sp. 3. p. 2046), caule paniculato. ramis

♂. caule toto glabro foliis saturatius viridi-lucidis infer. oblongo-lanceolatis, radio disco paulò longiore. — In solo pingui.

1. *PAETRIANNA*, caule piloso-lineato racemoso virgato, ramis sub-1-cephalis, foliis lanceolatis utrinque modicè angustatis acutis subintegerrimis marginibus scabris, ramorum ramulorumque conformibus, invol. laxi squamis linearibus subæqualibus. ♀ in Hungariâ circa Pesthinum. A. riparius Nees ast. 73. nec p. 280. Ligulæ purpurascentes? *Achænia hirta*. Species dubia ex m.

1. *MULTICARPUS* (Lindl. adn. mss.), undiquè glanduloso-pubescent, caule ramoso, ramis erectis 1-cephalis, foliis oblongo-linearibus acutis pubescentibus serratis, invol. squamis linearibus subulatis basi albidis squarrosis. In Louisiana. Species distinctissima nulli cognita acutè affinis. Amelli secunda diceret si squamæ invol. interiores membranaceæ et coloratæ essent. Lindl. mss.)

\*\* Versiformes Nees ast. p. 73.

1. *sessilis*, lanceolata vel oblongo-lanceolata inf. basi attenuata sup. basi a. invol. squamæ erecto-patulae laxæ erectæ vel recurvæ. Capitula medio-radio pallidè cæruleo vel albo. — Sectio vix à præcedente distincta.

1. *MURIVUS* (Ait. kew. ed. 1. v. 3. p. 203), caule paniculato-ramoso vel ramoso, ramulis sparsis laxis 1-cephalis, foliis lanceolatis ciliatis, radicalibus pinnato-serratis, caulinis integerrimis, invol. angustis obconicis squamis integerrimis subulatis. ♀ in Amer. bor. uliginosis siccis sylvis caducis à Noveboraco ad Pennsylvaniam. Pursh fl. bor. am. 2. p. 549. Nees ast. p. 74. Lindl. ! mss. Ligulæ albidæ. *Achænia puberula* basi attenuata. In specim. vivo à cl. meo ad hort. Genev. misso caulis equidem glaber ut vult cl. auct. sed rami casto-hirsuti ut innuit Willd. sp. 3. p. 2030. (v. v. c.)

1. *SUMERICATUS* (Nees ast. p. 74). caule piloso-lineato, ramis virgatis ramulis hirsutis subspicatis, foliis oblongo-lanceolatis acutis adpresso-serratis glaucis margine scabris amplexicaulibus, invol. subæqualis laxi squamis ext. stipulato-lanceolatis latioribus. ♀ in promontorio Mullgrave ad oram bot. cid. Americæ sept. An eadem ac *A. blandus* Pursh? Ligulæ purpurascentes. Humum densè incano-strigulosum.

1. *BLANDUS* (Pursh fl. bor. am. 2. p. 555), caule pyramidato-ramoso, ramis axillaribus racemiferis folio vix longioribus, foliis subamplexicaulibus longo-lanceolatis acuminatis serratis glabris, pedunculis tomentosis nudis, velutris laxis subæqualibus disco brevioribus. ♀ in Canadâ ex herb. Banks. mss. ast. p. 76. Ligulæ pallidè purpureæ.

1. *OBLIQUUS* (Nees syn. 1818. 27. ast. p. 76), caule infernè glabro supernè cernoso-composito stricto, ramis apice subcorymbosis, foliis linearilanceolatis sessilibus mucronatis subintegerrimis suprâ scabris obliquis, ramulorum tentibus, invol. laxi squamis ext. majoribus spatulato-lanceolatis patulis. In Amer. bor. ad Noveboracum. A. obliquus Lindl. pl. exs. A. junceus 2 lanceolatus Nutt. gen. 2. p. 158. A. rigidulus Desf. cat. h. par. 1815. p. 122. a. A. lanceolatus auct. americ. ? Ligulæ lilacinæ et var. albæ. (v. v. c.)

1. *caule humiliore oligocephalo corymboso, pappo rufescente* Nees l. c.

1. *CAUSAPHA* (Lindl. adn. mss.), caule infernè glabro supernè pubescente cernoso-composito, capitulis ramorum secundis, foliis angusto-lanceolatis subulatis acuminatis uniformibus suprâ per totam superficiem scabris, superioribus gradatim minoribus, involucri hemisphærici squamis linearibus acutis nec conspicuè brevioribus, exterioribus laxis. ♀ in Louisiana. Ab A. obliquo distinctissimus foliis latioribus uniformibus semper ut videtur integerrimis, gmelicis majoribus. Ast. tenuifolium etiam mentitur. (Lindl. mss.)

1. *MINIMALIS* (Nees ast. p. 77), caule glabro laxo racemoso-composito, foliis sessililanceolatis amplexicantibus acutis integerrimis margine scabris, ramulis linearibus, invol. laxi squamis lanceolatis subæqualibus pluriseriatis discum



aequantibus ciliatis.  $\frac{1}{2}$  patr. dub. verisim Amer. bor. In hortis altissimis *See* A. salicifolius Nees syn. 26. nec alior. A. laevis Vahl herb. ex h. Puer. — 1 elth. t. p. 39. t. 35. f. 39. Ligulae albae subviolaceae. Achænia pubescentibris. (v. s.)

79. A. TARDIFLORUS (Nees! ast. 78 non Linn. ex Lindl.), caule glabro corboso-divaricato, foliis oblongo-lanceolatis obliquè amplexantibus apicem serratis circa margines scabris ramulorum obtusis, invol. squamis imbricatis squarroso-patulis crassis, infer. sæpè elongatis subradiantibus.  $\frac{1}{2}$  in Amer. locis depressis inundatis à Noveboraco ad Virginiam (Pursh.). Ligula lilaei Achænia glabra. — Ab hoc, monente cl. Lindley in litt. et confirmante *See* authent. à cl. Neesio comm. non differt Symphyotrichum unctuosum Nees! ast. p. 135. Species insignis ob pappi setas in anulum solubilem suberectas (v. v. et s. c.)

$\beta$ . *cæspitosus* (Lindl. adu. mss.), caule humiliore, ramis densè corymbosis ramorum caulinis parum minoribus.  $\frac{1}{2}$  cult. in hort. Angl. A. adulterinus Lindl. bot. reg. t. 1571 non Nees. Symphyotrichum unctuosum Nees! ast. p. 135. ex. Lindl. Achænia paucissimas setulas subparentia. Pappus in anulum solubilem basi coherens ex Nees.

80. A. NOVI-BELGII (Nees! ast. p. 79), caule corymboso-composito vel simpliciter composito, ramis pilosulis rigidis, foliis lanceolatis amplexantibus ac circa margines scabridis, inferioribus medio serratis, invol. laxi squamis imbricatis acuminatis.  $\frac{1}{2}$  in Amer. bor. sponte et hinc inde in Europâ ex hortis emigra

a. *ampliflorus* (Nees l. c.), ramis simpliciter racemosis subcorymbosis foliis oblongo-lanceolatis lanceolatisve, capitalis amplifloribus, lig. densis latioribus, invol. squamis laxi erectis sæpè aequalibus lanceolatis omnibus acutis. — A. Novi-Belgii Linn. sp. 1231. A. acrotinus Nees! 21. A. floribundus Nutt. gen. 2. p. 158. — A. Nova-Belgii, etc. Nees! h. Lagd. b. p. 69. ic. (v. v. c.)

$\beta$ . *squarrosus* (Nees l. c.), invol. squamis exterior. squarroso-patulis s. foliaceo-elongatis. — Cætera ut in var. a.

$\gamma$ . *serus* (Nees l. c.), caule exaltato, ligulis densis carneis discum venis bidis.

$\delta$ . *floribundus*, caule humiliore, foliis lanceolatis attenuatis, ramis cæpè dichotomo-corymbosis polycephalis, radio minus denso breviori, imbricatis imbricatis squamis inæqualibus.  $\frac{1}{2}$  A. floribundus Willd. sp. 3. p. 20 Nees! syn. 21. A. Novi-Belgii minor Nees! ast. p. 81. (v. v. c.)

81. A. LUXURIANS (Nees! ast. 83), caule paniculato, ramis apice corymbosis, ramulis 1-cephalis, foliis ovali-oblongis vel lanceolatis acuminatis attenuatis amplexantibus, invol. squamis laxè imbricatis lineari-lanceolatis acutis.  $\frac{1}{2}$  in Amer. bor. sepibus et agris neglectis à Novâ Angliâ ad Virginiam (Nees!) A. Novi-Belgii Nutt. gen. 2. p. 158. A. prenanthoides Nees! syn. p. (v. v. c.)

$\beta$ . caule magis composito magisque piloso-lineato. A. luxurians *See* syst. 3. p. 538. excl. syn. ex Nees.

82. A. ASPER (Nees! ast. p. 84), caule superne paniculato-composito glabro tuberculis aspero, ramis racemoso-corymbosis, foliis oblongo-lanceolatis serratis amplexantibus supra scabris, omnibus in medio serratis, invol. imbricatis squamis apice squarroso-patulis.  $\frac{1}{2}$  patr. ign. verisim. Amer. h. Ligulae ferè duplici serie pallide cæruleæ seu lilacine. Achænia obsoletè berula.

83. A. ADULTERINUS (Willd. enum. 2. p. 884), caule à basi paniculato-composito, ramis corymboso-racemosis, foliis oblongo-lanceolatis mucronatis amplexantibus levibus margine scabris, invol. obovatis subaequalibus squamatis spatulato-lanceolatis squarrosis post florescentiam magis auctis.  $\frac{1}{2}$  in Americâ septentr. Nees! syn. 22. ast. 85. Ligulae albae, demum pallide violaceae Achænia glabra. An var. A. Novi-Belgii ex Nees? (v. v. c.)

84. *A. squarrosulus* (Nees ast. p. 86), caule corymboso-decomposito, ramis apice corymbosis erectis, foliis lanceolato-acuminatis subamplexantibus medio argutè serratis in ambitu superne scabris, invol. ovati squamis angustis imbricatis apice recurvis.  $\mathcal{U}$  in Americâ boreali. *A. spectabilis* Willd. sp. 3. p. 2048. Nees syn. 21. *A. mutabilis* Linn. sp. 1280? ex Nees. *A. recurvatus* Spreng. in Schrad. Journ. bot. 1800. p. 195.

$\beta$ . *albiflorus*, elatior, radio albo dein subviolaceo, disco mox saturatè purpureo. Nees l. c.

85. *A. longitrolus* (Lam. l. dict. 1. 1782. p. 306), caule glabro paniculato patulo, ramis apice simpliciter corymbosis, foliis lanceolato-acuminatis subamplexantibus, infer. in medio argutè serratis, supra in ambitu latè scabris, invol. turbinati subæqualis squamis linearilanceolatis patulis.  $\mathcal{U}$  in Amer. bor. paludosis à Novæboracæ ad Carolinam. *A. junceus* Ait. kew. 3. (1789). p. 204. Nees! spec. min. *A. eminens* Nees ast. p. 87. (v. v. c.)

$\beta$ . *major* (Willd.), caule 3-4-pedali glabro, paniculâ densiore, foliis ramorum ovato-lanceolatis, disco latè cœruleo. Nees.

$\gamma$ . *eminens* (Nees ast. 87), caule 3-4-pedali laxè paniculato ad basin lineis à foliis decurrentibus exasperato, fol. ram. lanceolatis, radio parciore pallidiore. Ast. *eminens* Willd. enum. (1809) p. 886. Nees syn. 22.

$\delta$ . *densiflorus*, paniculâ densiore, caule 2-3-pedali subpurpureascente. Nees l. c.

$\epsilon$ . *lævigatus*, caule elato 5-6-pedali, foliis oblongo-lanceolatis lævibus solo margine scabris rigidis saturatè viridi-lucidis. Nees ast. 88 (excl. verisim. syn. Ait. alibi ut species propria admissio). Ast. *lævigatus* Pursh fl. bor. am. a. p. 553. nec. Willd.

$\zeta$ . *breviramus*, caule racemoso, ramis abbreviatis Nees l. c.

$\alpha$ . *albiflorus*, ligulis sæpius albis, rarius ad violaceo-rubrum tendentibus, foliis latiusculis supra magis scabris. Ast. *virginens* Nees! syn. p. 22. Ast. *bicolor* Dietr. ined. Ast. *eminens virginens* Lindl. bot. reg. t. 1656. ex foliis etiam superioribus grossè serratis et invol. squamis erectis nec patulis videtur altera species. (v. v. c.)

86. *A. Douglasii* (Lindl. adn. mss.), caule glabro racemoso-composito, ramis parè foliosis oligocephalis laxè corymbosis, foliis linearilanceolatis versua basin angustatis glabris, omnibus supremis exceptis serratis, invol. laxi subæqualis squamis linearibus acutis.  $\mathcal{U}$  in Amer. septentr. occid. campis apricis prope flum. Columbia confluentem legit cl. Douglas.

$\alpha$ . ramis abbreviatis pleiocephalis, capitulis, minoribus.

$\beta$ . ramis gracilibus elongatis oligocephalis subaphyllis, capitulis majoribus. (v. s. comm. à Soc. hort. Lond.)

87. *A. salicifolius* (Willd. sp. 3. p. 2040), caule glabro paniculato-racemoso, ramis virgatis apice ramulique corymbosis, foliis lanceolatis margine scabris infer. basi attenuatis in medio serratis, superioribus subamplexantibus, ramulorum paucis linearibus erectis, invol. laxi squamis subæqualibus.  $\mathcal{U}$  in Europæ mediæ secus fluminum ripas, in Alastiâ, Germaniâ, Hungariâ. Nees! ast. p. 90. DC. fl. fr. suppl. 470. *A. salicifolius* Scholl. fl. barb. suppl. 328. cum ic. ex W. et N. Ast. *Hungaricus* Poir. suppl. 1. p. 496. *Ligulæ alba*. (v. s.)

\*\*\* *Leucanthi*. — *Leucanthi* et *ericoidei* Nees ast. p. 91 et 106.

Invol. squamæ laxè imbricatæ lineares vel linearilanceolatæ acutæ parùm inæquales, intimæ longiores angustiores. — Caulis laxè corymbosus paniculatusve. Folia caulina oblongo-lanceolata, linearia aut subulata, ramulorum minima. Capit. parva seu mediocria, ligulis albis seu pallidis.

88. *A. simplex* (Willd. enum. 2. p. 887), caule glaberrimo racemoso-decomposito, ramis apice subcorymbosis ramulisque contractis oligocephalis, foliis lanceolatis acuminatis lævisimis margine scabris, infimis subserratis, invol. laxi squamis linearilanceolatis imbricatis.  $\mathcal{U}$  in Amer. septentr. à Novâ-Cæsareâ

ad Virginiam Nees! syn. 27. ast. p. 91. Spreng. syst. 3. p. 535. excl. syn. Similis *A. saligno*. Ligula alba. *Achenia puberula*. (v. v. c.)

89. *A. lanceolatus* (Willd. sp. 3. p. 2050), caule piloso-lineato à basi racemoso-decomposito, ramis erectis, ramulis corymbosis, foliis lanceolato-acuminatis sessilibus margine serrulato-scabris, infimis argutè serrulatis, ramulorum patulis, invol. laxi squamis imbricatis lineari-subulatis. ♀ in America boreali? Nees! syn. 27. ast. 92. Ligula albæ. (v. v. c.)

90. *A. arcuatus* (Nees ast. 93), caule piloso-lineato à basi racemoso-decomposito recurvato, ramulis apice subcorymbosis, foliis angustè lanceolatis acuminatis sessilibus margine scabris, caulinis rameisque argutè serrulatis, ramulorum integerrimis patentibus, involucri laxè imbricati squamis linearibus acutis. ♀ in paludosa Pennsylvania. An *Ast. recurvatus* Willd. et auct. ab hoc diversus? An ab *Ast. lanceolato* satis distinctus? †.

91. *A. leucanthemus* (Desf.! cat. h. par. ed. 3. p. 401), caule piloso-lineato racemoso-decomposito, ramulis circ. dicephalis laxis, foliis lineari-lanceolatis acuminatis sessilibus in ambitu scabris, inferioribus medio argutè subserrulatis, invol. squamis adpressè imbricatis apice subrecurvis. ♀ in America bor. depressa à Novâ-Cæsareâ ad Carolinam, nunc ex hortis migr. in Belgio et Germania hinc inde. Nees ast. 93. *Ast. dracunculoides* Willd. sp. 3. p. 2050. Nees syn. 28. non Lam. *A. bellidiflorus* Nees syn. 21, non Willd. *Ast. lanceolatus* Lejeune! fl. Spa 175. non Willd. *A. artemisiiflorus* Poir. suppl. 1. p. 500. Capitula parva, ligulis albis. (v. v. c. et s. ex Belgio.)

92. *A. latus* (Willd. enum. 2. p. 857), caule glabro racemoso-composito vti racemoso-decomposito, ramis laxis apice subcorymbosis, ramulis elongatis, foliis angustè lanceolatis acuminatis ambitu scabris, inferioribus subserratis, ramulorum linearibus oblique patentibus, capituli terminalis squamis invol. laxis subaequalibus linearibus, reliquorum imbricatis apice reflexis. ♀ in Amer. bor. sabulosis demissis à Novâ-Cæsareâ ad Virginiam. Nees! syn. 27. ast. 95. Ligula pallidè carnea subalbæ. *Achen. puberula*. (v. v. c.)

β. caule humiliori supernè racemoso-composito, foliis caulinis summis rameisque basi dilatato-amplexantibus. Nees l. c.

93. *A. bellidiflorus* (Willd. enum. 886), caule stricto racemoso-decomposito seu suprâ decomposito angusto infernè glaberrimo albo-variegato purpureo-cente, ramis apice racemosis racemoso-compositis, foliis lanceolatis adpresso-serrulatis margine scabris, ramulorum lanceolato-acuminatis subulatisve apice recurvis, capitulis confertis, invol. squamis imbricatis linearibus acutis inter. sæpè apice coloratis. ♀ in Amer. bor. Ligula albidæ, dein lilicæ in γ amonè purpureæ.

α. *rubricaulis*, caule purpureo scabro, ramis brevibus apice simpliciter racemosis, capitulis approximatis, radio specioso, foliis pedicellorum subimbricatis linearibus. — *A. bellidiflorus* Willd. enum. 2. p. 886. Nees! pl. viv. miss. Lindl. pl. exs. (v. v. c. et s.)

β. *angustus* (Nees l. c.), caule viridi alboque striato, ramis brevibus apice simpliciter racemosis, foliis ramulorum et pedicellorum acuminato-subulatis remotioribus. — *A. angustus* Nees syn. 26. (v. s. ex h. Gœtt.)

γ. *rigidulus* (Nees l. c.), ramis longioribus racemoso-decompositis et suprâ decompositis. Cœt. ut in var. δ. *Ast. rigidulus* Nees! syn. 29. (v. v. c.)

δ. *salicinus*, racemo terminali subsimplici, foliis caulinis ramorumque inferioribus latioribus. Nees l. c.

94. *A. parviflorus* (Nees syn. 29. ast. 99), caule pubescente racemoso-decomposito ramisque apice corymbosis strictis, foliis lanceolatis acuminatis medio argutè serratis enprâ in ambitu scabris, ramulorum linearibus acutis patentibus involucri squamis imbricatis linearibus acutis, ligulis linearibus involucri longioribus. ♀ in Amer. bor. à Canadâ ad Novam-Angliam (Pursh). Michx. (Raf.). *Aster miser* Ait. Kew. ed. 1. v. 3. p. 205. non *Lina pallida* lilicæ dein purpureæ. *Achenia pubescenti-scabra*. (v. s. ex fl. Raf. et α. à cl. Lindl.)

95. *A. LAMARCHIANUS* (Nees ast. 100), caule piloso-lineato racemoso-decomposito coarctato, ramis paniculato-racemosis, foliis lanceolatis acuminatis sessilibus adpresso-serratis supra scaberrimis, ramulorum lanceolato-acuminatis patentibus, invol. laxè subæqualis squamis linearibus.  $\frac{1}{2}$  in Canadâ (Lam.). *Ast. paniculatus* Lam. dict. 1. p. 306. non Ait. *A. Tradescanti* Nees! syn. 28. Affinis *A. divergenti*. Ligulæ pallidæ, ex Nees invol. breviores, in plantâ vivâ ab eo minis evidenter eo longiores. (v. v. c.)
96. *A. DRYGAEUS* (Ait. kew. ed. 1. v. 3. p. 205), caule piloso-lineato racemoso-composito, ramis æqualibus patentibus, foliis lanceolatis acuminatis medio mucronato-serratis in ambitu scabris, ramulorum oblongo-lanceolatis basi attenuatis integerrimis mucronulatis patentibus, invol. squamis imbricatis lanceolatis.  $\frac{1}{2}$  in agris neglectis et sylvis siccis Amer. bor. à Novo-Eboraco ad Carolinam (Pursh), et in dumetis prope Jalapam (Schied. et Dep.). Nees syn. 28. ast. p. 101. *A. divaricatus* Raf. in litt.  $\frac{1}{2}$  (v. s.)
- $\beta$ . *humilior*, foliis brevioribus magis confertis, paniculæ ramis infer. elongatis. Nees l. c.
97. *A. FRAGILIS* (Willd. sp. 3. p. 2051), caule piloso-lineato racemoso-decomposito, ramis subpaniculato-racemosis, foliis infer. oblongis adpresso-serratis, super. linearibus, omnibus in ambitu scabris, ramulorum oblongo-lanceolatis integerrimis mucronulatis patentibus, invol. arcuè imbricati brevis squamis lanceolato-linearibus.  $\frac{1}{2}$  in Americâ bor. (Muhl.), in paludosis pinetis Novæ Cæsareæ et Alabamæ (Torr.) Lindl. bot. reg. t. 1257. Bart. fl. philad. 2. p. 117. *Ast. scoparius* Nees syn. 28.
- $\beta$ . *subinteger*, foliis omnibus linearibus elongatis undulatis subintegerrimis. — *A. fragilis* Nees syn. p. 29. Desf. cat. h. Par. ed. 3. p. 176. (v. s.)
98. *A. FALCATUS* (Lindl. adn. mss.), caule piloso angusto racemoso-subcomposito, ramulis brevibus 1-cephalis paucifoliis, foliis linearibus obtusis falcatis pilosisculis, invol. laxi parum imbricati squamis linearibus sessinæculis.  $\frac{1}{2}$  in Americâ arcticâ. Facies omnino *A. hiemalis* sed affinitas vera cum *A. fragili*. Folia infer. Bupleuro falcato similis. Radius albus. (Lindl. mss.)
99. *A. TRADESCANTI* (Linn. sp. 1230), caule glabresculo racemoso-subcomposito patulo apice simpliciter racemoso, ramis virgatis, ramulis subsecundis, foliis linearibus acuminatis margine ciliato-scabris, infer. à medio argutè remotè serratis, ramorum ramulorumque patentibus mucronatis, involuero imbricato.  $\frac{1}{2}$  in Amer. bor. sæpibus et paludibus exsiccatis à Canadâ ad Virginiam. Nees ast. p. 103. *Ast. vimineus* Lam. dict. 1. p. 306. Desf. cat. h. Par. 176. *Ast. pinifolius* Nees! syn. 29. Ligulæ albæ demum subviolacæ. Capitula parva. (v. v. c.)
100. *A. CORDIFOLIUS* (Michx. fl. bor. am. 2. p. 112), caule glabro racemoso-composito decompositoque, ramis patulis intricatissimis, foliis glabris, inferioribus lineari-lanceolatis subserrulatis, rameis depauperatis squamiformibus patuloreflexis, ramis apice 1-cephalis, invol. imbricati squamis linearibus obtusiusculis, ligulis acutis, ovario glabro.  $\frac{1}{2}$  in aridis sylvarum Carolinæ infer. (Michx. Boes! Lindl.!), circa Noveboracum (Don) Lindl. bot. reg. t. 1487. Desf. cat. h. Par. ed. 2. p. 122. *A. foliolosus* Ell. sketch 2. p. 345. excl.? syn. Species Willd. sp. 3. p. 2028, ex Nees syn. p. 30, est fortè diversa. Ligulæ pallidè carnes. Pappus flosc. disci æqualis. (v. v. c.)
101. *A. DUMOSUS* (Linn. sp. 1227), caule glabresculo racemoso-composito decompositoque patulo, ramis apice corymbosis, foliis linearibus integerrimis margine serrulato-scabris, super. internodiis longioribus, rameis ramulorumque parvis acutis, invol. squamis imbricatis linearibus acutis.  $\frac{1}{2}$  in Amer. bor. campis siccis sterilibus à Novâ Angliâ ad Carolinam (Pursh). Nees ast. p. 99. *Ast. foliolosus* Nees syn. 30. *A. foliosus* Pers. ench. n. 46? *A. Novæ Angliæ*, etc. Herm. parad. p. 95. icon opt. Capitula parva. Ligulæ albæ. (v. v. c.)
102. *A. HALICOIDES* (Linn. sp. 1227) caule glabro racemoso-composito, ramis virgatis, ramulis subsecundis ab apice remotis, foliis linearibus angustis inte-

- gerrimis glaberrimis, invol. brevis squamis imbricatis subulato-patulis infer. cuspidatis.  $\frac{1}{2}$  in Amer. bor. sterilibus et ad margines viarum à Canadâ ad Carolinam (Pursh), in Kentucky (Raf.!) — Nees syn. 31. ast. 107. Ast. tenuifolius Willd. sp. 2. p. 2026. excl. syn. ex Nees. Rami apice non corymboid ad producti et 1-cephali. Capitula parva, ligulis albis. (v. s. comm. à cl. Rafin.)
103. A. GLABELLUS (Nees syn. 31. ast. 107), caule racemoso-subcomposito, ramis filiformibus, foliis caulinis infer. lanceolatis serratis, super. linearibus subulatis integerrimis recurvatis, invol. subhemisphaerici squamis imbricatis à basi oblongo-cuspidatis subciliatis.  $\frac{1}{2}$  in campis petrosis Amer. bor., in Pennsylvania, Novo Eboraco.
- a. caule foliisque infer. suprâ pilosis. — A. sparsiflorus Willd. enum. 880 ex parte sec. Nees.
- β. glaber. — A. damosus Nees syn. 31 non Linn. A. sparsiflorus Michx. fl. 2 p. 112. non Willd. A. ramosissimus et ni fallor A. leptophyllus et A. subulatus hort. Par. (v. s.)
104. A. PILOsus (Willd. sp. 3. p. 2025), caule villosa racemoso-composito, foliis linearibus lanceolatis integerrimis albo-villosis, capitulis secundis, invol. oblongi squamis lanceolato-subulatis imbricato-patulis.  $\frac{1}{2}$  in terrâ Illinoisensi Amer. bor. Nees syn. 30. ast. 109 Ast. villosus Michx. fl. bor. am. 2. p. 113. Cl. Nees dubitat num reverâ ab A. glabello differat.
105. A. HARECLADUS, caule fruticuloso gracili pube patulâ confertâ brevi hirsuto, ramis laxis parce ramosis ramulisque foliosis 1-cephalis, foliis sessilibus linearibus integerrimis mucronato-acutis utrinquè villosa-scabris, super. ciliatis, invol. laxè imbricatis squamis linearibus puberulis acutis.  $\frac{1}{2}$  in Mexicanæ prov. Texas districtibus orientalibus legit cl. Berlandier (pl. exs. n. 2085). Affinis A. scopario sed rami hirsuti laxiores oligocephali, capitulo triplò ferè minore (v. s. comm. à cl. inv.)
106. A. SCOPARIUS (DC. non Nees), caule fruticuloso erecto ramosissimo glabro apice summo vix ac ne vix puberulo, ramis erectis racemiferis, foliis sessilibus linearibus scabridis, super. ciliatis, ramulis foliolosis, invol. squamis laxè imbricatis linearibus acutis ciliatis.  $\frac{1}{2}$  in Mexicanæ prov. Texas districtibus orientalibus legit cl. Berlandier (pl. exs. n. 1868, 1871 et 1894). Habitat ere Ericæ scopariæ. Folia 4-5 lin. longa semilineam lata. Pappus rufus. Achromia puberulum. (v. s. comm. à cl. inv.)
- \*\*\* Densiflori Nees ast. 110.
- Invol. squamæ arcuè imbricatæ. Rami densè racemiflori, capitulis confertis subsecundis. Folia integra aut serrata.
107. A. HIRsutICAULIS (Lindl. adn. mss.), caule racemoso hirsutissimo, ramis foliis brevioribus vel parùm longioribus, foliis canlinis linearibus lanceolatis integerrimis acuminatis, costâ subtus hirsutissimâ, capitulis subsecundis, invol. squamis arcuè imbricatis subulatis.  $\frac{1}{2}$  in Amer. septentr. juxta Albany (Torrey.). Radix alba. Discus purpureus. (Lindl.)
108. A. DIFFUSUS (Ait. kew. ed. 1. v. 3. p. 205), caule pubescente racemoso-composito, ramis recurvo-patentibus, foliis caulinis lanceolatis, ramulis oblongo-lanceolatis medio serratis suprâ scabriusculis margine scaberrimis, capitulis subsecundis confertis, invol. squamis imbricatis acutis.  $\frac{1}{2}$  in Amer. bor. agris et sepius à Canadâ ad Carolinam (Pursh). Nees syn. 28. ast. 110. Lindl. pl. exs. Spreng. syst. 3. p. 586 excl. syn. A. divergens β Nutt. gen. 2. p. 150. A. secundiflorus Desf. h. par. 1815. Capitula parva, ligulis albis seu pallide lilacinis, disco luteo mox purpur. Achromia ad costas pubescenti-scabra (Nees). (v. v. c.)
109. A. STENOPHYLLUS (Lindl. adn. mss.), caule glabriusculo racemoso, ramis patentibus apice densissimè racemosis, foliis canlinis linearibus lanceolatis acuminatis scabriusculis, capitulis secundis. invol. squamis linearibus acuminatis

tis, interioribus membranaceis coloratis.  $\mathcal{N}$  in Amer. septentr. ? in hortis anglicis occurrit. Radii dilute carnei. Discus mutabilis (Lindl. mss.)

110. *A. myrica* (Linn. sp. 1232. excl. syn.), caule pubescenti-scabro racemoso-composito, ramis erectis, foliis caulinis oblongo-cuneiformibus acuminatissimis medio serratis rameisque ovalibus subintegerrimis suprâ scabris, capitulis subsecundis confertis, invol. laxè imbricati squamis acuminatis.  $\mathcal{N}$  in pratis odis ad Philadelphiam. Nutt. gen. sm. 2. p. 158. Nees ast. p. 111. non alior. Capit. a lin. lata. Ligulæ lineares albæ. Achæn. pubescentia.

$\beta$ . *myrsifolius*, caule altiori, capitulis majoribus.  $\mathcal{N}$  in hortis cultas. *A. myrsifolius* Willd. enum. suppl. 59.

$\gamma$ . *abbreviatus*, ramis abbreviatis capitulisque ideò magis confertis Nees l. c.

111. *A. pendulus* (Ait. kew. 3. p. 205), caule pubescenti-hirto laxo diffuse racemoso-composito, ramis elongatis flaccidis, foliis elliptico-lanceolatis pubescenti-scabris, caulinis medio adpresso-serratis, ramulorum ellipticis integerrimis, invol. imbricatis.  $\mathcal{N}$  in Amer. bor. agris neglectis (Pursh) ad Novæboracum (Bern.) Nees syn. 28. Ast. 114. Hoffm. phyt. bl. p. 80. t. c. f. 2. Ast. horizontalis Desf. ! cat. h. par. ed. 3. p. 402 ? Capitula parva, radio pallidè purpurascens seu albo. Achænia pubescenti-scabra. (v. v. c. in h. Par.)

112. *A. nyctons* (Lindl. adn. mss.), caule scabro laxo racemoso, ramis apice racemosis, foliis ovali-lanceolatis suprâ scabris ferè omnibus serratis, capitulis secundis, invol. squamis subulatis.  $\mathcal{N}$  in Amer. septentr. *A. pendulo* affinis, sed ramulis oligocephalis et capitulis amplioribus distinctus. (Nees in litt. ad Lindl.)

113. *A. Menziesii* (Lindl. adn. mss.) caule incano apice racemoso, foliis oblongo-lanceolatis sessilibus integerrimis scabrinisculis incanis, inv. squamis obtusis squarrosis.  $\mathcal{N}$  in America septentr. occid. legit cl. Menzies. Ab *A. concolor* vix differt nisi pilis rigidis nec mollibus, necnon invol. squamis squarrosis nec adpressis. (Lindl. mss.)

114. *A. concolor* (Linn. sp. 1228), caule pubescente apice racemoso-composito, foliis oblongo-lanceolatis sessilibus integerrimis cano-pubescentibus, capitulis ad axillas fol. ram. solitariis subsessilibus racemos foliosos constituentibus, invol. squamis imbricatis lanceolatis sericeis.  $\mathcal{N}$  in Amer. bor. pinetis quercetisque littoreis arenosis à Novâ Cæsareâ ad Floridam. Nees syn. 30. ast. 114. Lindl. ! pl. exs. Ell. sketch 2. p. 350. *Virgaria concolor* Raf. ! in litt. 1831. Radix in arenosis tuberosa. Ligulæ 12-15 latè cœruleæ. Discus etiam cœruleus (Ell.). Radius discusque purpureus (Nees). Achænium densè sericeo-villosum. Pappus rufus. (v. s. sp. ex Carolinâ et Ohio.)

115. *A. squarrosus* (Walt. car. 209), caule scabro composito-racemoso, ramis oligocephalis foliis minutis è basi cordatâ amplexicanli acuminatis reflexis margine scaberrimis, invol. imbricati squamis lineari-lingulatis mucronatis laxè adpressis apice subfoliaceis.  $\mathcal{N}$  in pinetis sterilibus Carolinæ infer. (Pursh) et Georgiæ (herb. Duby ! ) Nees ast. 115. Ell. sketch 2. p. 350. Rami virgati tenues. Folia parva rigidula. Ligulæ 15-20 latè cœruleæ. Achænia pilosa. (v. s. sp. coll. à cl. Bosc.)

116. *A. ramulosus* (Lindl. adn. mss. non Labill.), caule piloso erecto angusto racemoso, ramulis corymbosis 1-cephaliæve, foliis linearibus serrulato-scabris basi dilatatis sessilibus, ramulorum rectis minoribus, invol. squamis acutissimis mucronatis squarrosis.  $\mathcal{N}$  ? in America arcticâ. Capitula duplò majora quàm *A. multiflori*.

$\beta$ . *incano-pilosus*. — in Amer. arcticâ. (Lindl. mss.)

117. *A. multiflorus* (Ait. kew. ed. 1. v. 3. p. 116), caule pubescente racemoso-composito diffuso, foliis linearibus integerrimis margine serrulato-scabris subtrinerviis, ramulorum recurvis, invol. squamis laxè imbricatis apice recurvis subspathulatis mucronatis.  $\mathcal{N}$  in Amer. bor. campis aridis ad viarum margines à Canadâ ad Carolinam (Pursh). Nees ast. p. 116. Lindl. ! pl. exs. *A. ericoides* Lam. dict. 1. p. 304. Michx. fl. bor. am. 2. p. 113. ex Lindl. *A. ericoides multiflorus* Pers. ench. 2. p. 443. *A. multiflorus* et *A. ciliatus* Nees syn.

3o et 29. — Dill. elth. 4o. t. 36. f. 40. Capitula parva ligulis albis. Achæn. pubescenti-scabra. (v. s.)

β. *ciliatus* (Nees l. c.), foliis latioribus ramealibus patulis nec recurvis, omnibus ciliatis, caule stricto. — *A. ciliatus* Willd. sp. 3. p. 2027. (v. s. comm. à cl. Lindl.)

118. *A. RACEMOSUS* (Ell. sketch 2. p. 348), caule leviter pubescente diffuso, ramis gracilibus elongatis, foliis lineari-lanceolatis subtus subpubescentibus margine scabris, capitulis subsessilibus confertis juxta summitatem ramorum, invol. squamis arcuè imbricatis lineari-lanceolatis. ½ in solo pingui insula Paris Amer. bor. Capitula parva. Ligulæ pallide purpureæ. Affinis vid. *A. multifloro*.

119. *A. CANESCENS* (Pursh flor. bor. am. 2. p. 547), caule subsimplici racemoso-villoso, ramis 1-cephalis, foliis linearibus integerrimis margine serrulato-scabris subtrinerviis, ramulorum recurvis, invol. squamis acuminatis squarrosis. ② in Massachusetts (Lindl.), ad ripas fl. Missouri (Pursh), in argillois denudatis ab Arikarees ad Fort-Mandan (Nutt.). Aster biennis Nutt. gen. am. 2. p. 155. non Ledeb. Calimeris biennis Nees ast. 231. Capit. magnit. *Helidii*. Ligulæ circ. 20 purpurascenti-cœruleæ. Hic nec ad Calimeridem relatus ex aut. cl. Lindley in adn. mss.

120. *A. MICROPHYLLUS* (Torr. in Lindl. adn. mss. non Vent.), caule glaberrimo composito racemoso, ramis 1-cephalis strictissimis, foliis minutis ovatis glaberrimis cauli adpressis, invol. turbinatis squamis linearibus acutis basi albidis arcuè imbricatis. ①? in Amer. bor. ditioe Alabamæ legit cl. Torrey. Radii cyanei (Lindl. mss.)

\*\*\*\*\* Concinni *Nees ast. p. 118.*

Invol. squamis imbricatis basi latiores pallidæ apice virides ciliatis. Recept. alveolæ dentiformes. Achænia sparsim puberula. Folia latiuscula. Radices speciosus raro albus.

121. *A. PURPURATUS* (Nees ast. p. 118), caule subsimpliciter racemoso-virgato, ramis elongatis 1-cephalis, foliis angusto-lanceolatis amplexicaulibus medio argutè serratis margine scabris, super. ovato-lanceolatis, ramorum pluribus lanceolatis patulis integerrimis æquè distantibus æqualibusque, invol. turbinatis squamis imbricatis lanceolatis apice coloratis. ½ in Amer. septentr. ? Aster miser hort. Par. Lam. dict. 1. p. 308. excl. syn. Caulis spec. mei glaber nec pilis seriatim scaber. Discus mox purpureus. Ligulæ saturatè purpureæ. Achænia glabra. (v. s. coll. in h. Par. anno 1798. sub nom. *A. miseri*.)

122. *A. AZUREUS* (Lindl. adn. mss.), caule racemoso-composito virgato, ramulis elongatis 1-cephalis, foliis lanceolatis utrinquè acuminatis scaberrimis suberratatis, superioribus integris, ramorum subulatis, invol. hemisphærici squamis imbricatis apice tantum patulis. ½ in Louisianâ. Radius azureus. Habitus *A. purpurati*. (Lindl. mss.)

123. *A. RETROFLEXUS* (Lindl. adn. mss.), caule composito-racemoso, ramis brevibus 1-cephalis, foliis lineari-lanceolatis integerrimis glaberrimis, ramulorum lineari-subulatis recurvis distantibus, invol. hemisphærici squamis lineari-lanceolatis squarrosis. ½ in Amer. septentr. Discus albidus parum mutabilis. Radii cœrulei. *A. squarrosus* hort. Angl. est species diversissima. (v. s. comm. à cl. Lindl.)

124. *A. TURBINELLUS* (Lindl. adn. mss.), caule subsimpliciter racemoso-virgato, ramis elongatis filiformibus sub-1-cephalis, foliis oblongo-lanceolatis subsimplicicaulibus integerrimis, ramorum oblongis obtusis sensim in subulata de-crescentibus, invol. turbinatis squamis linearibus obtusiusculis apice herbacea. ½ in Louisianâ. Species ab *A. purpurato* diversa foliorum formâ, involacro magis turbinato, ramisque filiformibus nec rigidis strictis. (Lindl. mss.)

125. *A. TENUIFOLIUS* (Linn. sp. 1227), caule glabro racemoso-decomposito 11-

minque virgatis, foliis angusto-lanceolatis margine scabris, infer. medio subserratis, ramorum pluribus lanceolatis integerrimis patentibus aequè distantibus aequalibusque, ramulis 1-cephalis, invol. ovato-hemisphaerico squamis imbricatis è basi lineari-oblonga acutis apice patulis.  $\mathcal{L}$  in Americâ septentr. Nees ! ast. p. 119. Lindl. ! pl. exs. Ast. polyphyllus Willd. enum. a. p. 338. Ast. junceus Nees syn. 26. Ast. junceus et polyphyllus Pursh. Nutt. Ligula albo-violacea. Achenia pubescentia. Caules erecti. (v. v. c. in h. Par. et a. ex h. Gœtt.)

$\beta$ . caule recurvato, ramis adscendentibus, invol. laxiore radio densiore. Ast. recurvatus Nees syn. 26. A. tennifolius Mōss handb. 2. p. 1485 ex Nees.

$\gamma$ . caule ramisque erectis, capitulis paulò minoribus, invol. magis imbricato. Nees l. c.

126. *A. concinnus* (Willd. enum. 384), caule glabriusculo laxo subcorymboso, ramis virgatis dichotomo-paniculatis, foliis lanceolatis amplectentibus remotè argutè serratis, ramulorum oblongis integerrimis, invol. arcuè imbricati squamis margine albidis.  $\mathcal{L}$  in Amer. bor. campis et sylvis à Novo-Eboraco ad Pensylvaniam (Pursh). Nees syn. 21. ast. 121. Lindl. bot. reg. t. 1619. Colla hort. rip. app. 3. p. 22. t. 12. act. ac. Tur. 33. p. 134. t. 12. Capit. mediocria, radio pulchrè cœruleo.

$\beta$ . ramis strictioribus magis racemosis, foliis parvis crebrioribus. Nees l. c. (v. a. c. in h. Par. 1815.)

127. *A. onustus* (Nees ast. 122), caule glabriusculo supernè seriatiim hirsutulo racemoso-decomposito, ramis apice ramosis laxis, foliis lanceolatis utrinquè attenuatis amplectentibus medio adpresso-serratis margine scabris, capitulis subsecundis breviter pedicellatis, invol. brevis squamis imbricatis lanceolatis disco subæquantibus.  $\mathcal{L}$  in Amer. bor. ? *A. confertus* Desf. ! cat. h. Par. ed. 3. p. 401 non Nees. Ligulae albae violaceo subinctæ. Achenia glabra. Stricto jure nomen Desf. antiquius servandum esset, sed nomen Neesianum admisi ne confusio generis augeatur. (v. a. c.)

$\beta$ . *squarrosus* (Lindl. adn. mss.), foliis ramulorum linearibus squarrosis. Caltus in hortis Angliæ sub nomine *A. Tradescanti cœrulei*. (Lindl. mss.)

128. *A. chilensis* (Nees ast. 123), caule basi glabro apice seriatiim piloso racemoso-decomposito, ramis apice densè corymbosis racemosive, foliis lanceolatis amplexicaulibus acutis crenato-serratis utrinquè scabris, pedicellorum parvis oblongis squarrosis, invol. squamis arcuè imbricatis oblongis obtusis  $\mathcal{L}$ ? in regione montanà agri Chilensis (Hænk.), in Californiâ (Cham.) *A. Radula* Less. in linnæa 1831. p. 125. excl. syn. ex Nees.

129. *A. strictus* (Poir. suppl. 1. p. 498), caule glabro stricto ramoso-decomposito, ramis strictis coarctatis, foliis lanceolatis attenuatis subamplectentibus medio subserratis ambitu spinoso-scabris, invol. arcuè imbricato.  $\mathcal{L}$  in Amer. bor. Ligula elongatæ lilacinæ. Similis dicitur *A. præalto*. Nees ast. p. 124. *A. strictus* Pursh. *A. strictus* Wender. et *A. strictus* hortul. sunt diversi.

$\beta$ . *angustifolius* (Lindl. adn. mss.), foliis lineari-lanceolatis vix serratis in Amer. arcticâ.

130. *A. mutabilis* (Ait. kew. ed. 1. v. 3. p. 205. an Linn. ?), caule lævi à basi paniculato decomposito patulo, ramis densè corymbosis, foliis oblongo-lanceolatis acuminatis lævibus margine scabris subutroque glaucis, infer. medio serratis, super. amplexicaulibus integerrimis, invol. disco brevioris squamis imbricatis lanceolatis.  $\mathcal{L}$  in Amer. bor. agris neglectis et sylvis à Pensylvaniâ ad Virginiam (Pursh). Nees syn. 22. ast. 125. Willd. sp. 3. p. 2045. excl. syn. Pluk. *A. versicolor* hortul. — Herm. lugd. 6. p. 67. f. c. Variat foliis omnibus aut serratis aut integerrimis. Discus luteus mox purpureus. Ligulae cœruleæ seu purpureæ. (v. v. c.)

131. *A. confertus* (Nees ! syn. 21. ast. 126. non Desf.), caule lævi apice subcorymboso-decomposito, ramis confertissimis corymbosis, foliis oblongo-lanceolatis acuminatis basi cordato-amplectentibus medio adpresso serratis mar-



- gine scabris, invol. imbricato.  $\frac{1}{2}$  in Amer. bor. Ligulae albæ. Achenia glabra. Variat foliis magis distinctè serratis. (v. v. c. ab hort. Bonn. miss.)
132. *A. vasicolora* (Willd. sp. 3. p. 2045), caule lævi paniculato-composito, ramis apice simpliciter corymbosis densis, foliis oblongo-lanceolatis acuminatis levissimis concoloribus, infer. medio serratis, super. amplexantibus integerrimis, invol. squamis imbricatis lanceolatis.  $\frac{1}{2}$  in Amer. bor. campis neglectis et ad margines sylvarum à Novâ-Cesareâ ad Carolinam. Nees! syn. 22. ast. 127. Spreng. syst. 3. p. 335, excl. syn. Capitula majuscula radio primum albo demum saturatè violaceo; in iisdem indiv. adsunt flores violacei et albi. Ades proles monstruosa radio abbreviato lacero à Nees obs. (v. v. c. ab h. Bonn. miss.)
133. *A. LÆVIS* (Linn. sp. 1230), caule lævi racemoso-composito, ramulis brevibus in apice ramorum approximatis, foliis lanceolatis amplexantibus glabris ambitu scabris viridibus, inferioribus subserratis, invol. imbricato disco brevioribus.  $\frac{1}{2}$  in Amer. bor. sylvæ et ad paludum latera à Novo-Eboraco ad Virginiam (Pursh). Nees syn. 24. ast. p. 128. Ait. kew. 3. p. 206. Lindl. bot. reg. t. 1500. Capitula ampla, ligulis cœruleis. Achenia pubescentia. Nomen subdecipiens, nam folia non sunt omnino lævia. (v. v. c.)
134. *A. LÆVIGATUS* (Willd. sp. 3. p. 2046), caule lævi racemoso-composito virgato, foliis oblongo-lanceolatis amplexantibus lævibus margine scabris saturatè viridibus, infer. in petiolum alatum attenuatis parçè serratis, super. senim minoribus integerrimis, invol. discum æquantis squamis imbricatis lanceolatis.  $\frac{1}{2}$  in Amer. bor. paludosis sylvisque uliginosis à Pensylvaniâ ad Virginiam (Pursh). Nees syn. 24. ast. 129. Hook. bot. mag. t. 2995. Caulis purpureus. Ligulae elongatæ cœruleæ. Sæpè in hortis nunc cum lævi nunc cum instabili confusus. (v. v. c.)
135. *A. AMPLIXICAULIS* (Muhl. in Willd. sp. 3. p. 2046), caule lævi paniculato-corymboso, ramis dichotomo-corymbosis coarctatis, foliis oblongis acutis basi cordatè amplexantibus lucidis margine scabris infer. subserratis, invol. artè imbricati squamis latè linearibus.  $\frac{1}{2}$  in Pensylvaniâ (Muhl.), in sabulosis ad margines sylvarum à Novâ-Cesareâ ad Virginiam (Pursh). Nees syn. 22. ast. 130. Capitula mediocria, ligulis saturatè cœruleis.
136. *A. RUBRICAULIS* (Lam. dict. 1. p. 305), caule latè stricto racemoso-composito-subpaniculato, foliis oblongo-lanceolatis amplexantibus margine scabris glaucis, radicalibus ovatis in petiolum contractis obtusis; caulibus infer. basi attenuatis subserratis, super. integerrimis, invol. disco brevioris squamis imbricatis lanceolatis.  $\frac{1}{2}$  in Amer. septentr. (Lam.). Nees ast. p. 132. excl. forte var. æ. Lindl. ! pl. exs. Capitula saturatè violacea. Recept. simbrilliferam, simbrillis in pilum desinentibus. Achenia glabra. (v. v. c.)
137. *A. CYANEUS* (Hoffm. phyt. bl. 1. p. 71. t. B. f. 1), purpurascens-glaucus. caule lævi racemoso-composito subpaniculato, foliis oblongo-lanceolatis amplexantibus lævibus margine scabris glaucis, super. integerrimis, radicalibus spatulatis obtusis crenato-serratis, invol. disco brevioris squamis artè imbricatis ex ovato-triangularibus.  $\frac{1}{2}$  in America boreali. Nees ast. 182. Lindl. bot. reg. t. 1495. Recept. alveolato-dentatum epilosum. Ligulae cœruleæ.
- α. invol. squamis angustioribus. — *A. glaucus* Nees syn. 23. *A. deserti* Desf. ined. ex Nees.
- β. invol. squamis latioribus, ext. ovato-dilatatis. — *A. politus* Nees syn. 23 et *A. cyaneus* ibid. *A. rubricaulis cyaneus* Poir. suppl. 1. p. 493.

*Species non satis notæ.*

138. *A. ANSIF* (Mart. in act. sc. Monac. phil. 6. p. 195), fruticosus, foliis sessilibus angusto-linearibus acutis subtus hirsutis, capitulis solitariis axillaribus subsessilibus.  $\frac{1}{2}$  in monte Kurmâ Arabiæ. Aster ericæfolius Forst. arab. p. 150 *Ansif*, arab.
139. *A. CARICIFOLIUS* (H. B. et Kunth nov. gen. am. 4. p. 92. t. 833), cauli

- CERTASTERNIFOLIUS** hortul. = *Diploappus filifolius*.  
**CHRYSOPHYLLUS** Cunn. = *Eurybia chrysophylla*.  
**CHRYSOCONOIDES** Desf. = *Felicia angustifolia*.  
**CILIARIS** Thunb. *ms.* = *Mairia taxifolia*.  
**CONCOLOR** Linn. = *Sericocarpus concolor*.  
**CONYZOIDES** Desf. = *Felicia angustifolia*.  
**CONYZOIDES** Willd. = *Sericocarpus conyzoides*.  
**CORYMBOLIUS** Michx. = *Biotia corymbosa*.  
**CORYMBOLIUS** hortul. = *Biotia macrophylla*.  
**CORIACEUS** hortul. = *Diplostephium extenuatum*.  
**CORNIFOLIUS** Muhl. = *Diplostephium cornifolium*.  
**CORYMBOSUS** Ait. = *Biotia corymbosa*.  
**CORYMBOSUS** Moench. = *Inula germanica*.  
**CRENATUS** Less. = *Mairia crenata*.  
**CRINITUS** Linn. = *Athrixia heterophylla*.  
**CRINITUS** Thunb. = *Athrixia Capensis*.  
**CRISPUS** Forst. = *Franseria crispa*.  
**CYDONIAEFOLIUS** Cunn. = *Eurybia cydoniaefolia*.  
**CYMBALARIS** Thunb. = *Munychia Cymbalaris*.  
**DEBILIS** Boj. = *Athrixia debilis*.  
**DECURRENS** Moench. = *Inula bifrons*.  
**DECURRENS** Cunn. = *Eurybia decurrens*.  
**DENTATUS** Thunb. = *Felicia tenella*.  
**DENTATUS** Andr. = *Olearia dentata*.  
**DICHOTOMUS** Ell. = *Diplostephium dichotomum*.  
**DIVARICATUS** Linn. = *Biotia macrophylla*.  
**DIVARICATUS** Spreng. = *Diplostephium amygdalinum*.  
**DIFFUSUS ACUMINATUS** Pers. = *Diplostephium acuminatum*.  
**DRACUNCULOIDES** Lam. = *Galatella dracunculoides*.  
**DRACUNCULOIDES MINOR** Ledeb. = *Linosyris glabrata*.  
**DRACUNCULOIDES** Bess. = *Galatella punctata*.  
**DYSENTERICUS** All. = *Pulicaria dysenterica*.  
**ECHINATUS** Less. = *Felicia echinata*.  
**Ecklonia** Less. = *Agathæa Ecklonia*.  
**ELATUS** Bert. = *Biotia corymbosa*.  
**ELLIPTICUS** Cunn. = *Olearia elliptica*.  
**ELÆAGNIFOLIUS** Cunn. = *Eurybia farfureacea*.  
**ELÆOPHYLLUS** Cunn. = *Eurybia elæophylla*.  
**ELONGATUS** Thunb. = *Agathæa elongata*.  
**ENSIFOLIUS** All. = *Inula ensifolia*.  
**ERICEFOLIUS** Forst. = *Athrixia ericifolia*.  
**ERUBESCENS** Steb. = *Eurybia erubescens*.  
**EXASPERATUS** Linn. = *Eurybia aculeata*.  
**FASTIGIATUS** Fisch. = *Turczaninovia fastigiata*.  
**FASTIGIATUS** Lehm. = *Galatella fastigiata*.  
**FASTIGIATUS** Ledeb. = *Galatella Hauptii*.  
**FERRUGINEUS** Wendl. = *Olearia dentata*.  
**FILIFOLIUS** Vent. = *Diploappus filifolius*.

oblongis utrinquē acutis sessilibus margine revolutis, subhis reticulato-venosis triplinerviis, pedunculis subaphyllis, invol. imbricati laxiusculi squamis acutissimis. 2℥ in paludibus exsiccatis Georgiæ et Carolinæ (Parsh). Nees est. 277. Disci radii albi. Capit. mediocria. An Galatellæ sp.?

150. A. RIPARIUS (H. B. et Kunth nov. gen. am. 4. p. 92), herbaceus caespitosus procumbens ramosus microphyllus, ramis glabris floriferis elongatis, foliis sessilibus linearibus acutis integerrimis 1-nerviis glabris, capitulis solitariis terminalibus, invol. hemisphærici squamis lanceolato-linearibus imbricatis, ext. subquarrosis. 2℥ in humidis Mexici juxta lacum Cuiseo. A. riparius Nees est. p. 280 non p. 73. Ligulæ 20-25 albæ. Pappus rufescens pilosus. Achæn. ign.

### †† Species exclusæ.

- A. ACRIIS Linn. herb. = Tripolium vulgare.
- A. ACRIIS Linn. = Galatella punctata æ.
- A. ACRIIS Willd. = Galatella rigida.
- A. ACRIIS Nees. = Galatella dracunculoides.
- A. ACULEATUS Labill. = Eurybia aculeata.
- A. ACUMINATUS Michx. = Diplostephium acuminatum.
- A. ADFINIS Less. = Felicia adfinis.
- A. ÆTHIOPICUS Burm. = Agathæa microphylla.
- A. ALBESCENS Wall. = Amphirepis albescens.
- A. ALBUS Willd. = Heleastrum album.
- A. ALPINUS γ Poir. = Galatella Lusitanica.
- A. ALTAICUS Willd. = Calimeris Altaica.
- A. ALYSOIDES Turcz. = Calimeris alysoides.
- A. AMBIGUUS Bernh. = Biotia Schreberi.
- A. AMYGDALINUS Lam. = Diplostephium amygdalinum.
- A. AMYGDALINUS Ell. = Diplostephium umbellatum.
- A. ANGUSTIFOLIUS Jacq. = Felicia angustifolia β.
- A. ANNUUS Linn. = Stenactis annua.
- A. ARGOPHYLLUS Labill. = Eurybia argophylla.
- A. ARAGONENSIS Asso. = Galatella Aragonensis.
- A. ASPER Ham. = Calimeris asperissima.
- A. ASPERULUS Wall. = Stenactis asperula.
- A. ASPERRIMUS Wall. = Calimeris asperissima.
- A. AURANTIUS Linn. = Clomenoclema aurantia.
- A. BELLIDIATRUM Scop. = Bellidiastrum Michellii.
- A. BELLIDIODES Don. = Stenactis bellidioides.
- A. BERNBERIVOLIUS Cunn. = Eurybia erubescens.
- A. BIENNIS Ledeb. = Callistephus? biennis.
- A. BIFIDUS Nees. = Galatella dracunculoides.
- A. BIFRONS All. = Inula bifrons.
- A. BONARIENSIS Spreng. = Erigeron glaucum.
- A. CANUS W. et Kit. = Galatella çana.
- A. CAFRORUM Less. = Felicia Cafrorum.
- A. CAPENSIS Less. = Agathæa amelloides.
- A. CAUCASICUS Muss. = Diplopappus Caucasicus.
- A. CHARIS Less. = Agathæa strigosa.
- A. CHAMÆDRIFOLIUS Lam. = Printzia Bergii.
- A. CHINENSIS Linn. = Callistephus Chinensis.

- A. MACRORHIZUS Cunn. = *Eurybiopsis macrorhiza*.
- A. MACROPHYLLUS Raf. = *Biotia latifolia*.
- A. MACRORHIZUS Thunb. = *Felicia macrorhiza*.
- A. MAGELLANICUS Lam. = *Clarionea magellanica*.
- A. MAGELLANICUS Spreng. = *Chilotrichium amelloideum*.
- A. MARYLANDICUS Michx. = *Sericocarpus conyzoides*.
- A. MAXIMUS Less. = *Erigeron maximum*.
- A. MESPILIFOLIUS Less. = *Munychia*? *mespilifolia*.
- A. MICROPHYLLUS Vent. = *Eurybia microphylla*.
- A. MICROPHYLLUS Labill. = *Eurybia lepidophylla*.
- A. MONTANUS All. = *Inula montana*.
- A. MORANENSIS H. B. K. = *Diplostephium Moranense*.
- A. MONTICOLA Lindl. = *Erigeron monticola*.
- A. MURICATUS Thunb. = *Felicia muricata*.
- A. MYRSINOIDES Labill. = *Eurybia myrsinoides*.
- A. NEMORALIS Ait. = *Galatella nemoralis*.
- A. NEMORALIS Willd. = *Galatella rigida*.
- A. NUDICAULIS Lam. = *Lagenophora Commersonii*.
- A. OBOVATUS Ell. = *Diplostephium obovatum*.
- A. OBTUSATUS Thunb. = *Diplopappus obtusatus*.
- A. OBTUSATUS Willd. = *Diplostephium obtusatum*.
- A. OCHRACEUS H. B. K. = *Tetramolopium ochraceum*.
- A. ODORUS All. = *Pulicaria odora*.
- A. ODONTOPHYLLUS Wall. = *Doronicum linifolium*.
- A. OFFICINALIS All. = *Inula Helenium*.
- A. OPPOSITIFOLIUS Linn. = *Munychia Cymbalariae*?
- A. PALUDOSUS Ait. = *Heleastrum paludosum*.
- A. PALUSTRIS Lam. = *Tripolium vulgare*.
- A. PANNONICUS Jacq. = *Tripolium vulgare* β.
- A. PAPPOUS Willd. = *Printzia Bergii*.
- A. PAUCIFLORUS Nutt. = *Tripolium pauciflorum*.
- A. PEDUNCULARIS Wall. = *Amphirepis peduncularis*.
- A. PEREZIODES Less. = *Mairia perexioides*.
- A. PERSONIODES Cunn. = *Eurybia personioides*.
- A. PHLOGOPAPPUS Labill. = *Olearia phlogopappa*.
- A. PHLOGOTRICHUS Spreng. = *Olearia phlogopappa*.
- A. PHYLOIDES H. B. K. = *Tetramolopium phyticoides*.
- A. PINULEOIDES Cunn. = *Eurybia pinuleoides*.
- A. PINNATUS Cav. = *Clomenocoma*? *pinnata*.
- A. PLANTAGINEUS Poepp. = *Senecio plantagineus*.
- A. POLIFOLIUS Linn. = *Printzia Bergii*.
- A. PROPINQUUS Cunn. = *Eurybia propinqua*.
- A. PUBESCENS Moench. = *Inula squarrosa*.
- A. PULCHERRIMUS Lodd. = *Diplopappus linariifolius*.
- A. PULICARIUS Scop. = *Pulicaria vulgaris*.
- A. PULVERULENTUS Hoffmann. = *Eurybia lirata*.
- A. PUMILUS Fisch. = *Calimeria Altaica*.
- A. PUNCTATUS W. et Kit. = *Galatella punctata* γ.

- A. PYRRAICUS Pourr. = *Jasonia sicula* β.  
 A. QUERCIFOLIUS Sieb. = *Olearia quercifolia*.  
 A. RAMULOSUS Labill. = *Euryhia ramulosa*.  
 A. REFLEXUS Michx. = *Felicia reflexa*.  
 A. RIGIDUS Michx. = *Diplopappus linarifolius*.  
 A. RIGIDUS Linn. = *Diplopappus rigidus*.  
 A. RIGIDUS Mœnch. = *Inula salicina*.  
 A. ROSEUS Stev. = *Calimeris rosea*.  
 A. ROSMARINIFOLIUS Spreng. = *Diplopappus obtusatus*.  
 A. ROTUNDIFOLIUS Thunb. = *Agathæa rotundifolia*.  
 A. RUPESTRIS H. B. K. = *Tetramolopium rupestre*.  
 A. RUTABODORUS Cunn. = *Euryhia glandulosa*.  
 A. SALICINUS All. = *Inula salicina*.  
 A. SALINUS Schrad. = *Tripolium vulgare* δ.  
 A. SATURNIÆFOLIUS Banks. = *Lorentea astureioides*.  
 A. SCABER Thunb. = *Doellingeria scabra*.  
 A. SCHREBERI Nees. = *Biotia Schreberi*.  
 A. SCORPIOIDES Scop. = *Aronicum scorpioides*.  
 A. SEDIFOLIUS Linn. = *Galatella punctata* α.  
 A. SERRATUS Thunb. = *Agathæa serrata*.  
 A. SERRATUS Lag. = *Grindelia inuloides*.  
 A. SINUATUS Cunn. = *Olearia quercifolia*.  
 A. SOLIDAGINEUS Michx. = *Sericocarpus solidagineus*.  
 A. SPATULARIS Cav. = *Grindelia*.  
 A. SPATULATUS Cels. = *Inula salicina* β.  
 A. SQUARROSUS All. = *Inula spiræifolia*.  
 A. SQUARROSUS Banks. = *Diplopappus linarifolius*.  
 A. STELLULATUS Labill. = *Olearia stellulata*.  
 A. STRICTUS Wendl. = *Galatella dracunculoides*.  
 A. STRIGOSUS Thunb. = *Amellus strigosus*.  
 A. STRIGOSUS Licht. = *Agathæa strigosa*.  
 A. SUBULATUS Michx. = *Tripolium subulatum*.  
 A. SUBCYMOSUS Bernh. = *Biotia Schreberi*.  
 A. TANACETIFOLIUS H. B. K. = *Machæranthera tanacetifolia*.  
 A. TATARICUS Linn. f. = *Calimeris incisa* ?  
 A. TAXIFOLIA Linn. = *Mairia taxifolia*.  
 A. TENELLUS Linn. = *Felicia tenella*.  
 A. TENERRIMUS Less. = *Tetramolopium tenerrimum*.  
 A. TOMENTOSUS Schr. et Wendl. = *Olearia tomentosa*.  
 A. TORTIFOLIUS Michx. = *Sericocarpus tortifolius*.  
 A. TRINERVIS Desf. = *Galatella rigida*.  
 A. TRINERVIS α Nees. = *Galatella punctata*.  
 A. TRINERVIUS Roxb. = *Galatella asperrima*.  
 A. TRIPOLIUM Linn. = *Tripolium vulgare*.  
 A. TRIPOLIUM Walt. = *Galatella flexuosa*.  
 A. UMBELLATUS Ait. = *Diplostephium umbellatum*.  
 A. UNDULATUS Mœnch. = *Pulicaria dysenterica*.  
 A. UNIFLORUS Michx. = *Galatella nemoralis* β.

**VERUS** Linn. = *Stenactis verna*.

**VILLOsus** Thunb. = *Felicia angustifolia* a.

**VISCOsus** Labill. = *Eurybia viscosa*.

**ZEYHERI** Less. = *Felicia Zeyheri*.

1. **TRIPOLIUM** Nees ast. 152. Lindl. obs. mss. — *Aster Adams. fam.* 2. p. 124. non Nees. — *Asteris* sp. Linn.

capitulum radiatum, ligulis foemineis uni-rariissime biseriatis, disco maphr. Invol. squamæ jam secus pedunculum ortæ bracteiformes sim aut imbricatæ aut subbiseriales, inter. longiores læves. Recept. solatum, alveolis dentatis. Styli fl. herm. rami apice crassiores subul. Achænia oblonga compressa glabra aut sparsim pilosa basi setulis ta. Pappus simplex pluriserialis, setis tenuibus remotè denticulatis. Herbæ subcarnosæ perennes rarius annuæ, in uliginosis salsuginosis icentes, glabræ aut tantum ad margines foliorum scabræ. Folia rna angusta integra aut vix serrata. Capitula corymbosa. Ligulæ ulæ aut albæ. — Genus recognoscendum, habitu plus quàm ca- teribus distinctum, hîc juxta Lindleyi admonitiones dispositum.

1. *Involucro subbiseriali, squamis omnibus obtusis, exterioribus brevioribus.* — *Tripolia vera*.

f. **VULGARIS** (Nees ast. 153), caulibus simplicibus ramosive diffusis aut adscen- dentibus glabris, foliis subcarnosis lineari-lanceolatis margine subcæbris basi filatis, radicalibus latioribus triplinerviis obtusis in petiolum attenuatis, ramis corymbosis, invol. squamis imbricatis interioribus maximis obtusis lævibus. f. Linn. Schrad. ② Willd. In paludosis salsis Europæ totius. *Aster Tripolium* Lam. sp. 1226. fl. dan. t. 615. engl. bot. t. 87. DC. fl. fr. n. 3137. *Aster acris* Lam. herb. mon. cl. Lindley. *Aster palustris* Lam. fl. fr. 2. p. 143. Ligulæ lilae, rarius albæ. (v. v.)

f. **Pannonicum** (Nees ast. 153), foliis infer. lanceolatis basi attenuatis, cau- linis angustis ciliato-serratis, caule adscendente aut erecto ramosissimo, invol. coloratis. — in paludosis subsalsis Hungariæ et Moraviæ. *Aster Pannonicus* Jacq. h. vind. 1. p. 3. t. 8. (v. v.)

γ. **diffusum**, foliis infer. lanceolatis basi attenuatis, caulibus angustis ciliato- serratis, caulibus à basi ramosissimis diffusis, invol. coloratis. — in palu- dosis salsis agri Chersonensis. *Aster Tripolium* Bieb. fl. taur. n. 1733. *Tripolium vulgare* f. Nees ast. 154? (v. s. comm. à cl. Beaupré).

δ. **salinum**, caule erecto subsimplici, foliis etiam infer. lanceolatis aut lineari- lanceolatis ciliato-serratis acutis, invol. colorato obtuso. ② in salsuginosis Sibiriz et Dahuriæ. *Aster salinus* Schrad. hort. Goett. 1. t. 3. neu journ. 1810. st. 2. p. 138. *Aster Tripolium* Ledeb. fl. alt. 4. p. 99. — Gmel. fl. sib. t. 80. f. 2. Ligulæ cyanæ paulò angustiores quàm priorum. An fortè spec. propr.? (v. s.)

ε. **longicaule**, caule erecto subsimplici, foliis etiam infer. lineari-lanceolatis acutis margine ciliato-serratis, invol. squamis subcoloratis apice obtusius- culis. — in Hispaniæ salsis maritimis loco dicto La Rapila legit cl. L. Du- four! et rep. etiam in Græciâ (ex h. ac. Monac.). *Aster longicaulis* Dufour! in litt. 1824. Fol. sup. valde acuminata. Pappus rufus. Vix à var. δ differt. (v. s.)

2. *Invol. pluriseriali, squamis imbricatis, interioribus saltem aut omnibus acuminatis.* — *Oxytripolia*.

**T. angustum** (Lindl. adn. mss.), caule angusto strictissimo, ramis erectis ra- cemosis, foliis linearibus serrulatis scabris glabris, invol. hemisphærici imbricati squamis linearibus acutis, ligulis inconspicuis. — in Amer. sept. Δ T. subulato

- A. *PRÆNANS* Poir. = *Jasonia stœlia* β.  
 A. *QUERCIFOLIUS* Sieb. = *Olearia quercifolia*.  
 A. *RAMULOSUS* Labill. = *Eurybia ramulosa*.  
 A. *REFLEXUS* Michx. = *Felicla reflexa*.  
 A. *RIGIDUS* Michx. = *Diplopappus linearifolius*.  
 A. *RIGIDUS* Linn. = *Diplopappus rigidus*.  
 A. *RIGIDUS* Moench. = *Inula salicina*.  
 A. *ROSEUS* Stev. = *Callimeris rosea*.  
 A. *ROSMARINIFOLIUS* Spreng. = *Diplopappus obtusatus*.  
 A. *ROTUNDIFOLIUS* Thunb. = *Agathaea rotundifolia*.  
 A. *RUFESTRIS* H. B. K. = *Tetramolopium rupestre*.  
 A. *RUTABODORUS* Cunn. = *Eurybia glandulosa*.  
 A. *SALICINUS* All. = *Inula salicina*.  
 A. *SALINUS* Schrad. = *Tripolium vulgare* β.  
 A. *SATURNIÆFOLIUS* Banks. = *Lorentea satnreoides*.  
 A. *SCABER* Thunb. = *Duellingeria scabra*.  
 A. *SCHREBERI* Nees. = *Biotia Schreberi*.  
 A. *SCORPIOIDES* Scop. = *Aronium scorpioides*.  
 A. *SEDFOLIUS* Linn. = *Galatella punctata* α.  
 A. *SERRATUS* Thunb. = *Agathaea serrata*.  
 A. *SERRATUS* Lag. = *Grindelia lauloides*.  
 A. *SINUATUS* Cunn. = *Olearia quercifolia*.  
 A. *SOLIDAGINEUS* Michx. = *Sericocarpus solidaginens*.  
 A. *SPATHULARIS* Cav. = *Grindelia*.  
 A. *SPATHULATUS* Cels. = *Inula salicina* β.  
 A. *SQUARROSUS* All. = *Inula spiracifolia*.  
 A. *SQUARROSUS* Banks. = *Diplopappus linearifolius*.  
 A. *STELLULATUS* Labill. = *Olearia stellulata*.  
 A. *STRICTUS* Wendl. = *Galatella dracunculoides*.  
 A. *STRIGOSUS* Thunb. = *Amellus strigosus*.  
 A. *STRIGOSUS* Licht. = *Agathaea strigosa*.  
 A. *SUBULATUS* Michx. = *Tripolium subulatum*.  
 A. *SUBCYMOSUS* Bernh. = *Biotia Schreberi*.  
 A. *TACHETIFOLIUS* H. B. K. = *Machæranthera tanacetifolia*.  
 A. *TATARICUS* Linn. f. = *Callimeris incisa*?  
 A. *TAXIFOLIA* Linn. = *Mairia taxifolia*.  
 A. *TENELLUS* Linn. = *Felicla tenella*.  
 A. *TENERIMUS* Less. = *Tetramolopium tenerimum*.  
 A. *TOMENTOSUS* Schr. et Wendl. = *Olearia tomentosa*.  
 A. *TORTIFOLIUS* Michx. = *Sericocarpus tortifolius*.  
 A. *TRINERVIS* Desf. = *Galatella rigida*.  
 A. *TRINERVIS* α Nees. = *Galatella punctata*.  
 A. *TRINERVIUS* Roxb. = *Galatella asperima*.  
 A. *TRIPOLIUM* Linn. = *Tripolium vulgare*.  
 A. *TRIPOLIUM* Walt. = *Galatella flexuosa*.  
 A. *UMBELLATUS* Ait. = *Diplostephium umbellatum*.  
 A. *UNDULATUS* Moench. = *Fulcaria dysenterica*.  
 A. *UNIFLORUS* Michx. = *Galatella nemoralis* β.

apice corymboso-composito, ramis ramulisque gracilibus patentibus foliolosis, invol. squamis ovatis obtusiusculis ciliatis.  $\frac{1}{2}$  in *Fastigiatus* Lehm. mss. ex Nees.

**AST.** (Nees ast. 173), caule puberulo scabro apice corymboso, folio-linearibus integerrimis nerviis impunctatis scabris, invol. scabris acutis pauciseriatis, ligulis elongatis. — in *sphagnosis Amer.* ad Novam Caesaream. *Aster nemoralis* Ait. kew. v. 3. p. 198. *fls* Parsh fl. 2. p. 544.

**aphyllus.** — *Aster uniflorus* Michx. fl. 2. p. 110. ex Nees et Lindl. *alp.* suth. vid. (v. s.)

**ma**, glabra, caule apice angulato, foliis lineari-lanceolatis acuminatis margine et sæpe costâ mediâ serrato-scabris impunctatis, subulatis in squamas parvas subulatas secus pedunculum desinenti-imbricatâ squamis infer. acuminatissimis, intim. longioribus margine cili lanceolatis minüs acuminatis. — in Sibiriz jugo Altaico et ocul legit cl. Patrin. Variat caule simplici et ramoso, ramis paucis ligulæ ample cœruleæ. Styli rami acuminati longè exserti. (v. s. unic. Delessert.)

**roLIA** (Lindl. ! adn. mss.), caule erecto tenui glabro simplici x-cephale subcorymboso, foliis linearibus glabris vix margine subulatis aut infimis trinerviis, ramulis folia subulata parva subimbricata, invol. squamis lanceolatis acutis subciliatis siccis, ligulis elongatis.  $\frac{1}{2}$  (h. Lindl.!). in regione Kirghisorum (Moric.). *Aster tenuifolius* ex Lindl. ! *Aster leptophyllus* Fisch. ex Lindl., sed non est *Aster* à Neesio citatus. Valde accedit ad priorem sed folia vix  $1-1\frac{1}{2}$  lin. lata. Styli rami minüs exserti obtusiusculi. (v. s.)

**A** (Lindl. ! adn. mss.), caule stricto flosco subfastigiato, ramulis x-cephalis, foliis linearibus integerrimis nerviis impunctatis, invol. hemisphaericis squamis linearibus acutis disco vix brevioribus.  $\frac{1}{2}$  septentrionali legit cl. Royle sub n. 239. (v. s.)

**zona** (Lindl. adn. mss.), caule corymboso, ramulis elongatis aphyllis lineari-lanceolatis acuminatis margine scabris, involuero brevissimo,  $\frac{1}{2}$  triplo longioribus.  $\frac{1}{2}$  in Sibiria. Amman. Ramulis longè aphyllis non cognoscitur. Cert. mihi ign.

*non satis notæ (fortè, mon. cl. Lindl., ad Asteres alpinos rejicienda?)*

**PANICA** (Nees ast. 165), caule erecto pubescente 1-3-cephalo, foliis  $\frac{1}{2}$  radicalibus spathulatis apice dentatis, caulinis linearibus integerrimis supra bisulcis, invol. squamis laxè imbricatis lanceolatis discum superantibus.  $\frac{1}{2}$  in dumetis Lusitaniz Brot. fl. lus. 1. p. 385. *Aster Lusitanicus* Brot. phyt. 63. *Alpinus*  $\gamma$  *fugax* Poir. suppl. 1. p. 489. Ligulæ cœruleæ aut purpureæ steriles. Achenia subvillosa. †

**ROMANENSIS** (Nees ast. 167), caule erecto apice corymboso oligocephalo  $\frac{1}{2}$  radicalibus spathulatis integerrimis, caulinis linearibus acutis  $\frac{1}{2}$  bisulcis, invol. squamis imbricatis discum subsequantibus, ligulis elongatis. Arragoniâ. *Aster Arragonensis* Asso fl. arr. 121. t. 8. f. 2. Lam. bot. ill. t. 681. f. 5. Ligulæ cœruleæ. †

†† *Species exclusa.*

**ma** Nees. = *Eurybia glandulosa*.

## CXI. TURCZANINOVIA.

**ma** pluriflorum radiatum, ligulis 7-8 uniserialibus femineis



differt caule nullo modo flexuoso, foliis serrulatis, squamis invol. capitalesque duplò majoribus. (Lindl.)

α. ramis oligocephalis. — in montibus saxosis.

β. ramis polyccephalis. — ad lacum Slave dictum.

3. *T. FLEXUOSUM* (Nees ast. p. 155), glabrum, caule erecto tereti flexuoso, foliis linearis-subulatis crassis laevibus impunctatis, ramis 1-cephalis squamisque invol. squamis imbricatis lanceolatis acutis, infer. subpatulis, ligulis invol. circiter aequalibus, achenio glabresculo. ½ in paludibus salis Amer. bor. à Novâ-Caesarâ ad Novaboracum (Nutt.), in Carolinâ (Nutt. Ell.). *Aster Tripolium* Walt. car. 210 non L. *A. sparsiflorus* Pursh fl. 2. p. 547 non Michx. *Ast. flexuosus* Nutt. gen. am. 2, p. 145 non Fisch. Foliis carnosius accedit ad *Tr. vulgare*, sed invol. regulariter imbricato ligulisque brevissimis facile differt. Folia infer. trinervia. (v. s.)

4. *T. PAUCIFLORUM* (Nees ast. 167. Lindl. obs. mas. 1835), caule humili simplici 3-6-cephalo, ramis 1-cephalis viscido-pubescentibus, foliis linearibus subcor-nosis glabris, caulinis subulatis amplexicaulibus, invol. hemisphaerici squamis subaequalibus acutis. ½ ad margines fontium aquae salis prope castellum Missis-sippi ad Missouri fluvium. *Aster pauciflorus* Nutt. gen. am. 2. p. 154. Habitat Tripolii sed caractere diversus, potius fortè ad *Asteres alpinos* referendus ex Lindl. adn. mas. 1834.

5. *T. SUBULATUM* (Nees ast. 157. Lindl. adn. mas.), caule erecto racemoso-decomposito, foliis linearibus acuminatis integerrimis, summis ferè subulatis, paniculâ laxâ, invol. squamis imbricatis linearibus acuminatis exter. lacin-culis, ligulis biserialibus vix involucre longioribus. ① in maritimis paludosis Americæ. Achenia compressa oblonga puberula. Pappus 1-serialis. Recept. nudum. Invol. demum reflexum. Species in orbe novo dispersa et polymorpha, nempe:

\* *boreale*; à Pensylvaniâ ad Carolinam. *Aster subulatus* Michx. fl. bor. am. 2. p. 111. *Aster longifolius* Desf. cat. h. par. ed. 3. p. 400. Poir. dict. 7. p. 480. (v. s. ex Carol. comm.)

\*\* *Mexicanum*. — circa urbem Matamoros legit Berlandier pl. exa. n. 2339. (v. s.)

\*\*\* *Cubense*. — in insulâ Cubâ (Poepp. La Sagra!), *Erigeron expansus* Spr. syst. 3. p. 518. (v. s.)

\*\*\*\* *Domingense*. — in Sancto-Domingo ad flum. Yague. *Erig. limifolium* Bert. herb. (v. s.)

\*\*\*\*\* *Brasilianum*. — in Brazilæ prov. Rio-Grande. (h. Mus. Par. n. 859) et in Bandâ orientale (Bacle!). (v. s.)

\*\*\*\*\* — in ins. O-Wahu (Cham.). *Aster subulatus* Less. in Linn. 1831 p. 120. Capitula paulò minora. (v. s.)

6. *T. CONSPICUUM* (Lindl. obs. mas. 1835), caule erecto racemoso, foliis linearibus acuminatis canaliculatis, ramis distantibus subaequilongis 3-6-cephalis rarius decompositis, bracteis ramorum et pedicellorum acutis oncullatis squamiformibus, invol. turbinato, squamis interioribus basi pallidis apice herbaceo vel coloratis acutis. — in Chili legit cl. Bridges pl. exa. n. 183. Caulis bipedalis glaber, internodiis inflorescentiæ ramis longioribus vel parum brevioribus. — Specim. Bridg. non vidi. Huc verisim. referenda *Baccharis asteroides* Bert. n. 83 et 828, in pascuis et hortis circa Rancagua et Valparaiso lecta. (v. s.)

### † Species exclusa.

*T. PALUDOSUM* Nees. = *Doellingeria paludosa*.

CX. *GALATELLA* Cass. dict. 37 (1817) p. 463 et 488. Nees ast. 158. Lindl. obs. mas. — *Galatea* Cass. dict. 18. p. 56. bull. philom. 1818. p. 165. Less. syn. 187. — *Asteris* sp. auct.

Capitulum radiatum, ligulis 1-serialibus neutris aut organa foemine abortiva gerentibus, fl. disci hermaphroditis tubulosis fertilibus. Invol.

ius squamis imbricatis inappendiculatis, exter. sæpè tri-  
cept. alveolatum, alv. marginibus dentatis. Styli fl. herm.  
spatulati obtusi. Achænia densè hirsuta vel strigoso-sericea.  
triserialis, setis filiformibus rigidis scabridis. — Herbar. pe-  
hemisphaerium boreale sparsæ. Caulis simplex apice corym-  
bia alterna integra oblonga aut linearia omnia basi saltem  
sessilia, sæpè punctata, punctis in superficie superâ præser-  
impressis subglandulosis. Capitula ferè Asteris sed ligulæ  
antes et invol. breve. Discus flavus. Ligulæ cœrulescens pur-  
pureæ aut albæ.

\* *Folius punctatis.*

**LYA** (Nees ast. p. 170), caule stricto tereti apice corymboso; foliis  
integerrimis utrinquè punctatis subcæbris 1-nerviis, ramulis 1-ce-  
pt. lineari-subulata parva gerentibus, invol. squamis lanceolatis acutis  
abmembranaceis disco brevioribus, ligulis paucis discum non super-  
antibus in Amer. bor. umbrosis siccis à Virginia ad Pensylvaniam. Aster  
Linn. sp. 1228. Galatella albiflora Cass. dict. 18. p. 58. Chrysopsis  
att. gen. sm. 2. p. 152. Ligulæ steriles albæ. Cl. Nees qui optimè  
et syn. ast. p. 18 designavit, incautè in ast. folia impunctata dixit.

**TRITIA** (Nees ast. p. 160), caule stricto apice coarctato-corymboso,  
ob-corymbosis, foliis lanceolato-linearibus acutis integerrimis trinerviis  
scabris, ramulorum lineari-subulatis, invol. squamis omnibus  
sessilibus elongatis. 2. in campis sabulosis et sylvis à Nova-Cæsarea  
ad Aster hyssopifolius Linn. mant. 114 (non Cav.) et ejus herb. ex  
L. sketch 2. p. 342. Ligulæ albæ purpureo tinctæ, 5 ex Ell., 10 in  
L. obviis. (v. s. comm. à cl. Lindl.)

**ATA**, caule erecto tereti scabrido simplici aut apice corymboso, foliis  
aut lineari-lanceolatis trinerviis aut summis 1-nerviis punctatis sub-  
nullis folia parva lineari-subulata gerentibus, invol. obconici squa-  
matis acutis margine membranaceis disco brevioribus, ligulis invol.  
longioribus. 2. in Europâ ab Hispaniâ ad Caucasum. Hæc mihi  
rursus Aster acris Linn. (conf. fl. fr. n. 3138 et suppl. p. 469) ex syn.  
et acretudine, sed cum altera species sit in ejus herb. servata et plu-  
rimo nomen ad alter. spec. transtulerint, hoc omnino rejicendum  
stat, ne confusio augeatur. Suad. cl. Lindley huc trahi var. sequentes:  
**sedifolia**, foliis omnibus linearibus angustis. — in apricis Hispaniæ,  
et Mediterraneæ, Italiæ. A. acris Linn. sp. ed. 2. p. 122 (excl. syn. et  
Hungariæ) non Willd. A. hyssopifolius Cav. ic. 3. p. 17. t. 232 excl.  
A. sedifolius Linn. sp. ed. 1. p. 874. Chrysocoma Linosyris Asso! fl.  
p. 116 excl. syn. — Lob. ic. t. 349. f. 2. Barr. ic. t. 606. Simillima  
Gallia. Ligulæ lilacinæ. (v. v. et s.)

**media**, foliis lineari-lanceolatis basi trinerviis, ramis linearibus 1-  
nerviis. — in Italiæ et Galliæ austr. collibus. Aster acris hort. Par. Reich.  
n. 1. p. 246. Galatea intermedia Cass. dict. 18. p. 58. Galat. punctata  
ast. p. 161. — Garid. Aix t. 11. Pluk. alm. t. 271. f. 3. Videtur om-  
nem in solo pinguiori orta. (v. s. et v.)

**iposa**, foliis latius linearibus lanceolatis ferè omnibus evidentius trinerviis  
— in salis subuliginosis Hungariæ (W. et K. Graaf!), Podoliæ austr.  
et Tauriæ (Bieb.), Caucasi (Mey.), Persiæ prov. Karabagh  
et. A. punctatus Waldst. et Kit. hung. 2. t. 109. Willd. sp. 3.  
22. excl. syn. Cav. Aster dracunculoides Bess. l. enum. 33. Galatella  
lata Cass. dict. 18. p. 57. Galatella insculpta Nees ast. p. 162. Aster  
rvis var. α Nees syn. 19 (v. s. sp. et v. c.)

(Nees ast. 163), tota lanngine molli canescens, caule erecto à medio

- ramoso-corymboso, foliis lanceolato-oblongis mucronatis integerrimis trinerviis punctatis, ramis subcorymbosis virgatis, invol. squamis 4-seriatis ovato-lanceolatis acutis vix margine membranaceis, ligulis disco ferè duplò longioribus.  $\frac{1}{2}$  in fruticosis Banatùs Hungariæ. Aster canus W. et K. 1. p. 28. t. 36. Gal. canescens Cass. dict. 18. p. 57. Ligulæ illacinae 8-10. Achænia albo-lanata (v. s. sp. et v. c.)
5. *G. DAMURICA*, caule glabriusculo subangulato erecto apice corymboso, ramis foliosis, foliis lineari-lanceolatis acuminatis trinerviis margine scabris punctulatis, invol. hemisphaerici squamis pluriseriatis imbricatis acuminatis subulatis disco paulò brevioribus, ligulis 20-30 elongatis.  $\frac{1}{2}$  in sabulosis Dahuriæ ad ripas Ircuti flum. detexit cl. Turczaninow. Aster Hauptii Turcz. l in litt. nos alior. Folia 3 lin. lata. Achænia villosa. Ligulæ purpurascences, ab omnibus prioribus monente cl. Lindl. diff. invol. hemisphaerico nec obconico. (v. s.)
6. *G. DRACUNCULOIDES*, caule stricto tereti puberulo apice breviter densèque corymboso, foliis lineari-lanceolatis mucrone calloso apiculatis integerrimis trinerviis utrinquè punctatis subtus scabridis, ramealibus angustis subulatis, ramulis di- aut ramoso-polycephalis, invol. squamis margine scariosis, ext. acutis int. obtusis, ligulis paucis disco duplò longioribus aut nullis.  $\frac{1}{2}$  in campestribus apricis Sibiriae, Tauriae, Caucasi. Aster dracunculoides Ledeb. fl. alt. 4. p. 97.
- a. *radiata*, ligulis paucis. — Aster dracunculoides Lam. dict. 1. p. 363. Aster acris Nees syn. 19. Galatella pauciflora Nees ast. p. 164. Ast. strictus Wend. act. soc. Marb. 2. f. 6. p. 54 ex Nees. Ligulæ corollae lineares (v. s.)
- β. *discoidea*, ligulis nullis. — Chrysocoma biflora Linn. sp. 1178. Chrysocoma dracunculoides Lam. dict. 2. p. 192. Crinitaria biflora Cass. dict. 37. p. 476. Galatella biflora Nees ast. p. 159. Aster bifidus Nees syn. 18. — Gmel. sib. 2. p. 189. t. 82. f. 1. Cl. Ledebour monet ligulas in iisdem individuis nunc paucas nunc nullas.
- \*\* *Folii impunctatis.*
7. *G. RIGIDA* (Cass. dict. 18. p. 58. sub Galatella), caule erecto substriato puberulo apice corymboso, foliis lanceolato-linearibus mucronato-acutis integerrimis trinerviis impunctatis subtus margineque scabris, ramis flor. subtrichis pedicellisque rigidis, invol. squamis lanceolatis subacutis margine membranaceis, ligulis disco longioribus.  $\frac{1}{2}$  in vineis circa Meldas Occitanorum legit cl. Prost. Aster acris Willd. sp. 3. p. 2023. non Linn. Gall. acris Nees ast. 171. Aster trinervis hort. Par. l Nees syn. 18. Aster acris cl. Lam. dict. 1. p. 363. excl. ubiquè synonym. hic non citatis. Ligulæ corollae. Herba junior subarch-noides. (v. v. c.)
8. *G. LEPTOPHYLLA* (Nees ast. 167), caule angulato apice simpliciter corymboso juniore lanuginoso, foliis lanceolato-linearibus integerrimis acutis trinerviis carinatis impunctatis, juvenilibus ad costas marginesque sublanuginosis densim scabris, ramis gracilibus minutè foliosis, invol. squamis ovatis obtusis ciliatis, ligulis disco longioribus, achæniis hirtis. — in America septentrionali? Ligulæ saturatè corollae. Aster leptophyllus a cl. Desfontaines ipso mecum comm. est diversissimus, nec in Sprengelii opere etiamsi à Neesio citatus reperitur. Forte eadem ac *G. rigida*. Species valdè obscura. †
9. *G. HAUPTII* (Lindl. adn. mss.), foliis oblongo-linearibus acuminatis trinerviis margine scabris, caule basi subsimplici supernè in ramos corymboso-fastigiatos monocephalos densè foliosos diviso, invol. turbinati squamis pluriseriatis, exter. linearibus acuminatis, intermed. latioribus acutis, intus membranaceis rubescentibus, omnibus costâ fuscâ notatis.  $\frac{1}{2}$  in Sibiriae altis collibus et locis apricis frequens. Aster fastigiatus Ledeb. ic. fl. ross. alt. t. 161 non Fisch. nec. Lehm. Aster Hauptii Ledeb. fl. alt. 4. p. 100.
10. *G. FASTIGIATA* (Nees ast. 169), foliis angustè lanceolatis integerrimis subulato-acuminatis trinerviis planis impunctatis ad costam subtus marginesque

- scabris, caule apice corymboso-composito, ramis ramulisque gracilibus patentibus apice minutè foliolosis, invol. squamis ovatis obtusiusculis ciliatis.  $\frac{1}{2}$  in Sibiria. Aster fastigiatus Lehm. mss. ex Nees.
11. G. *MEMORIALIS* (Nees ast. 173), caule puberulo scabro apice corymboso, foliis lanceolato-linearibus integerrimis enerviis impunctatis scabris, invol. squamis linearibus acutis pauciseriatis, ligulis elongatis. — in sphagnosis Amer. bor. à Canadâ ad Novam Caesaream. Aster nemoralis Ait. kew. v. 3. p. 198. Aster ledifolius Pursh fl. 2. p. 544.
- $\beta$ . *monocephalus*. — Aster uniflorus Michx. fl. 2. p. 110. ex Nees et Lindl. qui specim. auth. vid. (v. s.)
12. G. *SQUAMOSA*, glabra, caule apice angulato, foliis lineari-lanceolatis acuminatis trinerviis margine et sæpè costâ mediâ serrato-scabris impunctatis, summis lineari-subulatis in squamas parvas subnatas secus pedunculum desinentibus, invol. imbricatè squamis infer. acuminatissimis, intim. longioribus margine membranaceis lanceolatis minùs acuminatis. — in Sibiria jugo Altaico et circa Barnaul legit cl. Patrin. Variat caule simplici et ramoso, ramis paucis 1-cephalis. Ligulæ amplæ cœruleæ. Styli rami acuminatè longè exserti. (v. s. comm. ab amic. Delessert.)
13. G. *TENUIFOLIA* (Lindl. ! adn. mss.), caule erecto tenui glabro simplici 1-cephalo aut apice subcorymboso, foliis linearibus glabris vix margine subscabris uninerviis aut infimis trinerviis, ramulis folia subulata parva subimbricata gerentibus, invol. squamis lanceolatis acutis subciliatis siccis, ligulis elongatis.  $\frac{1}{2}$  in Sibiria (h. Lindl. !), in regione Kirghisorum (Morici. !). Aster tenuifolius Linn. herb. ex Lindl. ! Aster leptophyllus Fisch. ex Lindl., sed non est Aster homonymus à Neesio citatus. Valde accedit ad priorem sed folia vix 1-1  $\frac{1}{2}$  lin. lata sive glabra. Styli rami minùs exserti obtusiusculi. (v. s.)
14. G. *JUNCICA* (Lindl. ! adn. mss.), caule stricto junceo subfastigiato, ramulis subaphyllis 1-cephalis, foliis linearibus integerrimis enerviis impunctatis laeviusculis, invol. hemisphaerici squamis linearibus acutis disco vix brevioribus.  $\frac{1}{2}$  ? in India septentrionali legit cl. Royle sub n. 139. (v. s.)
15. G. *SUDIFLORA* (Lindl. adn. mss.), caule corymboso, ramulis elongatis aphyllis, foliis lineari-lanceolatis acuminatis margine scabris, involuero brevissimo, ligulis invol. triplò longioribus.  $\frac{1}{2}$  in Sibiria. Amman. Ramulis longè aphyllis primo aspectu cognoscitur. Cert. mihi igno.

† *Species non satis notæ (fortè, mon. cl. Lindl., ad Asteres alpinos rejicienda?)*

16. G. ? *LUSITANICA* (Nees ast. 165), caule erecto pubescente 1-3-cephalo, foliis glabriusculis, radicalibus spathulatis apice dentatis, caulinis linearibus integerrimis punctatis suprâ bisulcis, invol. squamis laxè imbricatis lanceolatis discum subæquantibus, ligulis lanceolatis discum superantibus.  $\frac{1}{2}$  in dumetis Lusitaniae. Aster fugax Brot. fl. lus. 1. p. 385. Aster Lusitanicus Brot. phyt. 63. t. 29. Aster Alpinus  $\gamma$  fugax Poir. suppl. 1. p. 489. Ligulæ cœruleæ aut purpureæ sæpè steriles. Achænia subvillosa. †
17. G. ? *ARRAGONENSIS* (Nees ast. 167), caule erecto apice corymboso oligocephalo lanato, foliis radicalibus spathulatis integerrimis, caulinis linearibus acutis subitis bisulcis, invol. squamis imbricatis discum subæquantibus, ligulis elongatis.  $\frac{1}{2}$  in Arragonia. Aster Arragonensis Asso fl. arr. 121. t. 8. f. 2. Lam. diet. 1. p. 302. ill. t. 681. f. 5. Ligulæ cœruleæ. †

†† *Species exclusæ.*

G. *GLANDULOSA* Nees. = Eurybia glandulosa.

## CXI. TURCZANINOVIA.

Capitulum pluriflorum radiatum, ligulis 7-8 uniserialibus fœmineis

fertilibus integris apice obliquis, disci 7-8 tubulosi 5-dentatis hermaphroditis fertilibus paucisque sterilibus mixti. Recept. alveolatum. Invol. squamæ circ. biseriales lineares obtusiusculæ disco multò breviores. Stigm. radii elongata, disci brevia inclusa. Achænia compressa glabra colorata. Pappus 1-serialis setosus, setis sericeis subcabris. — Herba Sibirica aut Dahurica simplex erecta apice corymboso-fastigiata, basi glabra, apice pubescenti-scabrida. Folia alterna lineari-lanceolata integerrima acuta uninervia. Corymbus compositus fastigiatus polycephalus. Ligulæ albæ. Discus flavus. — Habitus Galatellæ sed ligulæ fertiles et achænia glaberrima. Caract. ferè Asteris sed involucrium disco multò brevius pauci-seriale, squamis non foliaceis, etc. — Dicavi cl. N. Turczaninof, qui Dahuriæ regionumque vicinarum botanicam egregie illustrat et mecum hanc ipsam stirpem cum pluribus aliis rarissimis communicavit.

1. *T. FASTIGIATA*. (2) seu  $\frac{1}{2}$  in Sibiria (Fisch.), in Dahuriæ paludosis ad flum Argunum (Turcz.). Aster fastigiatus Fisch. mem. soc. mosc. 3. p. 74 non Ledeb. (v. s. comm. à cl. Turczaninof.)

CXII. *CALIMERIS* Nees ast. p. 225. — *Kalimeris* Cass. dict. 24. p. 324. 37. p. 464 et 491. — *Asteris* spec. auct.

Capitulum radiatum, radio fœmineo fertili, ligulis 1-serialibus latiusculis, disco hermaphr. fertili. Stigm. apice conico lanceolata angustiora. Invol. squamæ 2-4-seriales subæquales herbaceæ margine membranaceæ laxè unbricatæ. Recept. convexi vel conici alveolæ subquadratae ad angulos obtusè dentatæ. Achænia compressa marginata hirta. Pappus sub-1-serialis setaceus inæqualis scaber. — Herba perennes Asiaticæ. Caulis angulatus apice corymbosus. Folia integerrima vel inciso-dentata. Discus flavus. Ligulæ albæ seu cœruleæ.

2. *C. FLEXUOSA* (Royle! in Lindl. adn. mss.), caule flexuoso glabro (aut hirsuto-scabro), foliis ovato-lanceolatis basi attenuatis apice acuminatis grossè serratis supra et margine setoso-scabris subtus pallidis secus nervos vix scabris, invol. squamis lanceolatis biserialis margine hyalino-scariosis, achæniis spontè hirtis.  $\frac{1}{2}$  in Indiæ orient. prov. Mussonæ legit cl. Royle pl. exa. n. 113. Ligulæ cyanææ lineares obtusæ. Pappus subrufescens corollulæ ferè æqualis. (v. s. comm. à cl. Royle.)

3. *C. INCISA*, caule erecto striato secus angulos scabro aut (in pl. cultâ) hirsuto apice corymboso, foliis lanceolatis utrinquè attenuatis remotè inciso-serratis margine subtusque scabris, invol. squamis apice rufo-marginatis, pappo corollæ duplò brevioris inæquali.  $\frac{1}{2}$  in Sibiria (Nees) et Dahuriæ (Turcz.) hemididis. Aster incisus Fisch. mem. soc. mosc. 3. p. 76 (1812). *Grindelia incisa* Spreng. syst. 3. p. 575. *Kalimeris platycephala* Cass. dict. 24 (1822). p. 325. Nees ast. 226. Caulis 1-2-pedalis. Ligulæ cœruleæ 6 lin. longæ, lineam latè. Variet ex Lindl. adn. mss., invol. squamis immarginatis. (v. s. et v. e.)

4. *C. ALTAICA* (Nees ast. 228. Lindl. adn. mss.), caule pubescenti-scabro semihæpidulo irregulariter corymboso, foliis super. lineari-lanceolatis integerrimis ciliatis, capitulis paucis subcorymbosis, invol. pubescentis squamis 3-4-seriatis lanceolatis acuminatis albo-marginatis, pappo piloso corollæ brevioris.  $\frac{1}{2}$  in montibus Sibiria Altaicæ (Ledeb.), Tanaicæ? (Fisch.), ad flum. Scherelle et in campis Barabensibus (Pall. ex Lindl. mss.). Aster pumilus Fisch.! est. h. Cov p. 67. A. Altaicus Willd. enum. p. 881. Ledeb. fl. alt. 4. p. 99. qui fortè à seq. non separat. Ligulæ cœruleæ angustissimæ. Pappus subviolaceus sericeus. Folia var. longiora et obtusiora. (v. s. comm. à cl. Fischer qui etiam ad Willd. misit.)

4. *C. CANESCENS* (Nees ast. 229), pubescenti-cana, caule erecto irregulariter corymboso inferne fasciculato-folioso, foliis lanceolato-linearibus obtusis integerrimis rariis subserratis, ramulis foliosis, capitulis solitariis, invol. squamis glabriusculis lanceolatis acuminatis albo-marginatis, pappo piloso corollâ paulò brevior. 2½ in Sibiria Altaica (Gebl. ex Lindl. mas.) ad Sagai (Fisch.), in arenosis ad Selengam (Pall. ex Lindl.). Aster Gebleri Bess. in litt. A. lithospermifolius hort. Par. 1821. Aster Altaicus plur. auct. et fortè à Cal. Altaica vix sætis differt. Capitula paulò majora. Ligulæ latiores. (v. s.)  
 γ? *latifolia* (Lindl. adn. mas.), caule erectiusculo scabrido-hispidulo, foliis caulinis oblongis retusis, ligulis latioribus. 2½ in India orient. ad Kumbawur legit cl. Royle (pl. exs. n. 103). Folia 8-10 lin. longa, 2-3 lin. lata, distantia. (v. s. comm. à cl. Royle.)
5. *C. TATARICA* (Lindl. adn. mas.), caule corymboso scabriusculo, foliis linearibus acutis pubescentibus margine ciliato-scabris, ramis patentibus sæpius simplicibus 1-cephalis, invol. squamis acuminatissimis hirsutis herbaceis (nec albo-marginatis). 2½ in Dahuria (Fisch. Presc. ex Lindl.), in arenosis Mongolice Chinensis (Turcz.!). Ligulæ azureæ 7-8 lin. longæ. (v. s. comm. à cl. Turcz.)
6. *C. HISPIDA* (Lindl. adn. mas. non Nees), caule striato apice composito-ramoso folisque linearibus acutis integerrimis hirsutis, invol. hispido-villosissimis squamis acuminatis, exterioribus squarrosis. 2½? in Sibiria (Gmel. ex Lindl.). Calimeris hispida Nees est *Heteropappi* sp. Capitula magnitudine et facie *C. Altaicae*. Invol. squamæ interiores vix membranaceæ.
7. *C. EXILIS* (Lindl. adn. mas.), caulibus caespitosis simplicibus 1-cephalis corymbosive pubescentibus, foliis adpressis linearibus acutis pubescenti-scabris incanis sensim in invol. squamas laxas angustas subhipidas abeuntibus. 2½ in Sibiria. *C. Altaica* proxima; foliis acutis, invol. squamis acuminatis capitulisque minimis, radio breviori diversa. (Lindl.)
8. *C. INTEGRIFOLIA* (Turcz. in litt. 1834), caule erecto simplici apice corymboso, ramis 5-7 foliosis 1-cephalis, foliis oblongo-linearibus sessilibus acutis integerrimis cauleque pube brevi pulverulento-velutinis, invol. squamis laxis glabriusculis acuminatis, ligulis ovali-oblongis. ① ②? in Sibiria ad flumen Amur legit cl. Turczaninof et ad flum. Argunum cl. Razgildeef, etiam è China in herb. Stanton video. Ligulæ cœruleæ latiores et breviores quàm *C. Altaicae*. (v. s.)
9. *C. ROSA*, glabriuscula basi suffruticosa, caulibus erectis ramosissimis, foliis linearibus integerrimis acutis, ramis 1-cephalis, invol. squamis circ. biserialis lanceolatis acuminatis margine membranaceis ciliatis, achænio sericeo-villoso, pappo piloso. 3 2½ in Caucasii orientalis monte Schadagh et in saxis circa Schindur legit cl. Steven. Ast. roseus Stev. act. mosc. 3. p. 245. et 4. p. 64. Bieb. suppl. n. 1727. Nees ast. p. 29 non hort. Par. Caulis 8-9 poll. longus. Folia ferè 2 poll. longa, sesquilineam lata. Capit. Ast. alpino vix minùs. Ligulæ 5 lin. longæ, lineam latæ. (v. s. comm. à cl. inv.)
10. *C. ALYSSOIDES*, fruticulosa, ramis foliosis adpressè cano-tomentosis, foliis oblongis integerrimis araneoso-tomentosis subtus canis, infer. obtusis, cæteris mucrone acutis, capitulis solitariis subcorymbosis, invol. squamis oblongis acuminatis subaraneosis erectis margine subacariosis uninerviis, achæniis paucis, pappo piloso. 3 in Mongolia Chinensis locis subarenosis detexit cl. Turczaninof. Aster alyssoides Turcz. chin. n. 21. Ligulæ albæ 5-7, 3-4 lin. longæ, lineam latæ. Folia 3-4 lin. longa, vix ultrâ lineam lata. (v. s. comm. à cl. inv.)

† *Species exclusæ.*

*C. BEAKNS* Nees. = *Aster canescens*.

*C. HISPIDA* Nees. = *Heteropappus hispidus*.

## CXIII. EURYBIOPSIS.

Capitulum pluriflorum radiatum, ligulis fœmineis 1-serialibus, fl. disci tubulosis 5-dent. hermaphr. Invol. squamæ pauciseriales imbricatæ lineares acuminatæ. Recept. angustum alveolatum. Achænia elongata compressa basi subangustata hispidula. Pappus 1-serialis multisetus, setis scabridis acuminatis. — Herba Australasica glabra multicaulis, rhizomate crasso lignoso, caulibus tenuibus virgatis parçè foliosis demùm frutescentibus. Folia alterna integerrima, radicalia lineari-oblonga obtusa, caulina linearia subacuta. Capitula ad apices caulium solitaria. — Fructus compressus Asteris, cætera Eurybiæ.

1. *E. MACRORHIZA*.  $\frac{1}{2}$   $\frac{3}{4}$  in Australiæ orâ boreali occid. ad radices collium prope Prince Regent's rivier, octob. flor. legit cl. A. Cunningham. Aster? *macrorhiza* A. Cann. in litt. 1834 non Thunb. (v. s. comm. à cl. invent.)

CXIV. *PODOCOMA* Cass. bull. philom. 1817. p. 137. dict. 37. p. 462 et 484. 42. p. 60. Less. syn. 187. — *Erigerontis* sp. Poir.

Capitulum radiatum, ligulis fœmineis 1-serialibus (Cass.) seu 1-2-ser. (Less.), disco hermaphrodito regulari. Recept. nudum planum. Invol. irregul. imbricatum disco brevius. Achænium oblongum compressum rostratum. Pappus pilosus pluriserialis conformis. — Herba perennes multicaules Bonarienses. Folia infima obovata dentata pubescentia, caulina sparsa oblongo-lanceolata. Capitula ramulos terminantia subcorymbosa. Ligulæ lineares albæ (Less.), forsan flavæ (Cass.). Pappus rufus. — Plures ex Less. adsunt in herbariis species in-descriptæ.

1. *P. HIERACIFOLIA* (Cass. dict. 42. p. 60), capitulis subcorymbosis.  $\frac{1}{2}$  in Bonariâ (Comm.) *Erigeron hieracifolium* Poir. dict. 8. p. 491.  
2. *P. PRIMULÆFOLIA* (Cass. l. c. p. 61), capitulis majoribus solitariis. In Mont-Video. †.

## CXV. ASTEROPSIS Less. syn. p. 188.

Capitulum radiatum, ligulis fœmineis 1-serialibus, disco hermaphrodito regulari. Recept. epaleaceum. Invol. pauciserialia. Achænium breviter rostratum. Pappus pilosus 1-serialis conformis. — Herba Brasiliensis. Folia alterna integra. Capitula solitaria.

1. *A. MACROCEPHALA* (Less. l. c.). — in Brasiliâ (Sellow). Cat. ign.

CXVI. *ARCTOGERON*. — *Erigerontis* sp. Linn. — *Asteris* sp. Gmel. Ann.

Capitulum radiatum, ligulis fœmineis 1-serialibus, disco hermaphr. fertili. Recept. angustum planum subalveolatum. Invol. imbricatum squamis 3-serialibus adpressis lanceolatis acuminatis carinâ virgibus et puberulis margine albo-scariosis. Ligulæ ovali-oblongæ apice dentatæ invol. duplò longiores. Stigm. disci et radii crassa brevia. Achænium oblongum subcompressum sericeo-villosissimum. Pappus simplex pluriserialis persistens, setis subinæqualibus scabris. — Herba Sibirica habitu *Armeriæ*, rhizomate caudiciformi frutescente multiplici brev. Folia radicalia conferta lineari-subulata margine scabro-ciliolata cæterum glabra rigida erecta pollicaria semilineam lata. Scapi vix foliis longiores hirsuti monocephali. Discus flavus. Radius albus. — Nomen et s, septentrion, et γειν, senex, quasi *Erigeron boreale*.

1. *A. GRAMINEUM*.  $\mathcal{L}$  in campis sabulosis Sibiria Transbaikalenſis (Gmel.) et Dahuriæ (Fisch.), ad Tchita (Patr.), frequens in desertis Selenge, Uda, Ononis et Arguni fl. adjacentibus (Amm.). *Erigeron gramineum* Linn. sp. 1212. Lam. ill. t. 681. f. 4. Poir. dict. 7. p. 490. — Gmel. fl. sib. 2. p. 174. t. 76. f. 2. — Amm. ruth. 215. *Erig. gramineus* est omnino diversus. (v. s. ex Selenga et Tchita à cl. Turca. et Patr. coll.)

CXVII. *SERICOCARPUS* Nees ast. p. 148. — *Asteris* sect. 1. Ell. sketch 2. p. 340. — *Asteris* sp. auct.

Capitulum radiatum, ligulis 1-serialibus paucis latiusculis fœmineis fertilibus, disco hermaphrodito paucifloro fertili. Stigmata apice subulata. Invol. imbricatum ovatum vel oblongum, squamis multiserialibus latis basi cartilagineis enerviis apice herbaceis. Recept. alveolæ sæptis lacero-ciliatæ. Achænia obpyramidata brevia dense sericeo-strigosa. Pappus simplex panicularis rigidus scaber. — Herba perennes ex America boreali ortæ. Caulis apice corymbosus. Folia impunctata integerrima basi attenuata. Capitula ad apices ramorum adgregata conyzoides. Ligulæ albæ.

1. *S. SOLIDAGINEUS* (Nees ast. 149. Lindl. ! adn. mas.), caule erecto tereti nervis prominulis subangulato glabro, foliis lineari-lanceolatis basi attenuatis apice obtusis margine scabris obscure trinerviis glabris, corymbo fastigiato, invol. oblongi squamis adpressis apice obtusis viridi-callois, ligulis 5 linearibus invol. duplò longioribus.  $\mathcal{L}$  in pinguibus paludosis et marginibus sylvarum Amer. boreal. à Pennsylvania ad Carolinam. *Conyza linifolia* Linn. sp. 1205. *Aster solidaginoides* Willd. sp. 3. p. 2024. Ell. sketch 2. p. 340 non Raf. *Aster solidagineus* Michx. fl. bor. am. 2. p. 108. — Pluk. alm. t. 79. f. 2. bene. (v. s. olim ex sem. à Michx. missis sub nom. *A. corymbosi*.)
2. *S. CONYZOIDES* (Nees ast. 150), caule erecto subangulato puberulo, foliis cuneato-oblongis subtrinerviis apice serratis acutis subtus sparſe et suprâ secus nervos puberulo-scabris, corymbi compositi ramis apice 3-cephalis, capitulis subsessilibus, invol. obovato-oblongi squamis ext. squarrosis int. erectis acutis, ligulis involucri aequalibus.  $\mathcal{L}$  in campis et dumetis à Pennsylvania ad Floridam (Pursh). *Conyza asteroides* Linn. sp. 1206. *Aster Marylandicus* Michx. fl. bor. am. 2. p. 108. *Aster conyzoides* Willd. sp. 3. p. 2043 non Desf.

$\beta$ ? *plantaginifolius* (Nees ast. 150), foliis radicalibus spatulatis, capitulis subpedicellatis, ligulis invol. aequalibus aut longioribus.  $\mathcal{L}$  in sylvis Novæ-Cæsaræ (Nutt.) et ex Noveboraco (Mitch.), Albany, Niagara et Montreal (Janin!), Kentucky (Raf.), recepi. *Aster conyzoides*  $\beta$  Nutt. l. c. Hanc semper ex America recepi sub nomine *A. conyzoidis*. An species propria? An ligulæ longitudine variantes? (v. s.)

3. *S. RIGIDUS* (Lindl. adn. mas.), foliis obovatis subtrinerviis integris cauleque scaberrimis, corymbo terminali simplici vel composito contracto, involucri turbinati squamis adpressis basi nudis, radio pappo brevior aut nullo.  $\mathcal{L}$  in America septentr.-occident. *S. conyzoides* proximus. (Lindl.)
4. *S. TORTIOSOLUS* (Nees ast. 151), totus pube brevi densâ rigidulâ canescens, caule erecto tereti apice corymboso, foliis obovato-cuneiformibus mucronatis integerrimis subaveniis, capitulis pedicellatis basi folia duo parva gerentibus, invol. obovati squamis oblongis imbricatis acutiusculis basi pallidis, ligulis involucri paulò longioribus.  $\mathcal{L}$  in Carolina infer. et Georgia. *Conyza bifoliata* Walt. car. 204. *Aster tortifolius* Michx. fl. 2. p. 109. Ell. sketch 2. p. 341. (v. s.)

CXVIII. *MACHERANTHERA* Nees ast. 224.

Capitulum radiatum, ligulis fœmineis (neutrisve?) linearibus 3-nerviis. Invol. pluriserialle imbricatum, squamis herbaceis margine pallidis



apice patulis. Recept. nudum? Stigmata disci apice lineari elongato sterili. Antheræ corollæ longiores basi muticæ appendice terminali cultriformi. Achænia compressa (conica?) sericeo-hirsuta, pilis pap-pum exter. simulantibus. Pappus pilosus inæqualis scaber. — Herba perennis pilis capitatis pubescens facie Anthemidis. Folia alterna pinna-tifida lobis dentatis. Capitula terminalia solitaria ramo apice nudo pe-dunculata. Ligulæ albæ ex H. et B., in sicco purpurascentes. Discus luteus.

1. *M. TANACETIFOLIA* (Nees l. c.). 2/ in hort. Mexic. culta. *Aster tanacetifolius* H. B. et Kunth nov. gen. am. 4. p. 95. *Aster chrysanthemoides* Willd. herb. Spreng. syst. 3. p. 538. Affinis dicitur *A. pinnato* Cav.

#### CXIX. TETRAMOLOPIUM *Nees ast. 202.*

Capitulum radiatum, ligulis angustis fœmineis fertilibus 1-serialibus, fl. disci hermaphr. abortivis? masculisve. Recept. nudum punctatum. Invol. imbricatum, squamis pauciserialibus angustis. Achænia 4-nervia, radii compressa cuneiformia, disci lineari-tetragona. Pappus 1-serialis setosus scabridus. — Species habitu variæ, fortè non omninò con-generes.

##### §. 1. *Involucro pauciseriali.*

1. *T. TENERRIMUM* (Nees l. c. p. 203), herbaceum caespitosum, foliis inferioribus confertis lineari-spathulatis acutis ciliato-serratis, ramis apice nodis. 2/ in insulâ O-Wahu. *Aster tenerrimus* Less. in Linnæâ 1829. p. 120. Ligulæ albæ? Capitula 3 lin. alta et lata.

##### §. 2. *Involucro multiseriali. — An species verè congeneres?*

2. *T. ? RUPESTRIS* (Nees l. c.), fruticosum erectum infernè nudum cicatrisatum, foliis lineari-lanceolatis margine revolutis subtus tomentosis. 3/ in frigidissimis montium Rucu-Pichinchensis et Antisana. *Aster rupestris* H. B. et Kunth nov. gen. am. 4. p. 94. t. 334. *Aster Pichinchensis* Willd. herb. ex Nees. Invol. squamæ multiseriales. Ligulæ circ. 30 albæ. Achænia glanduloso-punctata.
3. *T. ? PHYLLICOIDES*, fruticosum ramosissimum, ramis cano-lanatis, foliis appropiatis patenti-reflexis sessilibus oblongo-linearibus margine revolutis supra glabris subtus lanatis, capitulis corymboso-congestis, invol. campanulati squamis imbricatis lanceolato-linearibus lanato-ciliatis. 3/ in Mexico? *Aster phyllioides* H. B. et Kunth nov. gen. am. 4. p. 98. Capit. magnit. *Solidaginis Virgæ-aureæ*. Ligulæ circ. 20 albæ. Discus 8-10-florus, fl. masculis et hermaphr. mixtis. Ovarium lineare compressum hispidulum. Pappus pilosus rufescens scaber. — Affine ex Kunth *A. Chiliotricho* sed rec. epaleaceum, aut potius *Tetramolopio rupestri*.
4. *T. ? OCHRACEUM*, fruticosum, ramulis angulatis tomentosis, foliis petiolatis lanceolato-oblongis remotè et obsolete denticulatis supra glabris subtus ochra-ceo-tomentosis, corymbis ramosis, invol. subhemisphærici squamis laxè im-bricatis margine diaphanis. 3/ in montibus Quitensibus? *Diplostephium?* *ochraceum* Nees ast. 201. *Aster ochraceus* H. B. et Kunth nov. gen. am. 4. p. 94. Ligulæ circ. 5 albæ. Discus 20-florus, fl. masc. et hermaphr. intermixtis. Achænia obovata compressa, sterilia linearia. Pappus pilosus scaber. *Habitus omninò Eurybie lyratæ*.

CXX. *HENRICIA* *Cass. bull. phil. 1817. p. 11. et 1818. p. 123. dict. 20. p. 567 et 37. p. 491. Nees ast. 219.*

Capitulum radiatum, ligulis fœmineis fertilibus 1-serialibus, disco hermaphr. Invol. hemisphæricum disco æquale, squamis biserialis

s. adpressis, ext. foliaceis ovali-acutis, inter, scariosis obtusis convexum nudum. Achenium pilosiusculum cylindraceum. Aerialis setosus scabridus. — Suffrutex Madagascariensis ramis alterna breviter petiolata ovato-lanceolata serrata rigida setuloso-scarba. Corymbus aphyllus oligocephalus. Pedicelli v. l. subhispidum. Discus flavus. Ligula alba.

FRAXINUM (Cass. l. c.). ♂ in Madagascr. (v. s. in h. Juss.)

HELLINGERIA. — Dellingeria et Tripoli sp. Nees. — Asteris sp. auct.

lum radiatum, flor. radii 1-ser. ligulatis foemineis, disci her-  
5-dentatis. Recept. alveolatum. Involucrum imbricatum.  
disci brevis apice obtusa extus subhispidi. Achenia angulata  
6. Pappus biserialis pilosus, setis scabridis, aliis filiformibus  
7. incrassato-clavellatis. — Herbes rhizocarpica, Japonica aut  
s. Folia alterna serrata aut integra. Capitula fore Asterum ex  
- Genus certe apud cl. Nees species heterogeneas amplectens,  
scabrâ stabilitum.

RA (Nees ast. 183), caule herbaceo tereti glabro supernè dichotomo-  
so, foliis ovalibus utrinquè attenuatis medio grossè serratis rigidulis  
versè subhis secus nervos scabris, invol. squamis imbricatis obtusis,  
ser. 3, acheniis glabris. ♀ in Japoniâ. Aster scaber Thunb. fl. jap.  
Pappus scaber Less. in Nees. Pappus pallidè rufus plurisetosus, setis  
8. aliis apice incrassatis, aliis acutis. Stigmata brevia apice obtusa contin-  
ua. Achenium angulatum. (v. s. à Thunb. olim miss. in h. Delacour.)

SCABRA, caule herbaceo tereti, ramis pedicellisque vestino-scarbis,  
9. tubo-oblongis utrinquè attenuatis latè dentatis suprâ mucronato-scar-  
bis secus nervos nervulosque villosis subhirs, capitulis secus ramulos  
10. latè racemosis longè pedicellatis, invol. squamis oblongo-linearibus  
latè subobtusis, ligulis circ. 12, acheniis pilosis. ♀ in Chinâ legit  
11. Pappus pallidè rufescens, setis aliis filiformibus, aliis parvis subincrass-  
atis. Habitu ad D. scabram accedit, sed ex pappo media inter Dellin-  
12. gium et Asterem. (v. s. in h. Lindley.)

### † Species exclusæ.

ALINA Nees. = Diplostephium amygdalinum.

ALTA Nees. = Diplostephium cornifolium.

ALTA Nees. = Diplostephium obovatum.

ALTA Nees. = Helcastrum album.

ALTA Nees. = Diplostephium umbellatum.

### CXXII. HELEASTRUM. — Asteris sp. auct.

lum radiatum, ligulis foemineis 1-serialibus, fl. disci her-  
nvol. imbricatum, squamis laxis acuminatis apice foliaceis  
rufis. Recept. alveolatum, alveolis exsertis acutè dentatis.  
13. disci elongati acuminatissimi dorso hispidi. Achenia oblonga  
utrinquè attenuata glabra. Pappus biserialis setis longitudine  
s. aliis filiformibus, aliis in ensem subclavellatam dilatatis.  
14. boreali-Americane perennes. Caules simplices. Folia alterna  
acuminata integerrima. Ligula coerulesc aut alba. — Differt à  
15. rufâ stigmatè acuminatissimo nec obtuso, à cæteris Eustereis

pappo biformi setis aliis subclavellatis, à Diplopappeis pappo exteriori interiore sequante. — Nomen ex *ἰλος*, *ἰλιος*, palus, et *αστρον*, astrum.

1. *H. PALUDOSUM*, caule apice subcorymboso et tenuissimè subpuberulo, foliis linearibus amplexicanalibus subconcavis apice subulatis, invol. squamis subaquarrosis.  $\frac{1}{2}$  in paludibus et uliginosis Carolinæ infer. (Pursh), Georgiæ prope Savannah (Herb. in h. Duby), à Novâ-Cæsareâ ad Floridam (Nutt.). Aster paludosus Ait. kew. ed. 1. v. 3. p. 201. Nutt. gen. 2. p. 155. Ell. sketch 2. p. 343. Aster grandiflorus Walt. ex Ell. Tripolium paludosum Nees ast. 155. Diplopappus paludosus Lindl. adn. mss. Capitula 3-5. Ligulæ latè coruleæ nimmerosæ. Caulis pubescens. Folia margine scabra. (v. s.)
2. *H. ALBUM*, caule apice in ramos 1-cephalos corymboso-fastigiatos divise, foliis oblongo-lanceolatis scabris infernè attenuatis, involucrio levî.  $\frac{1}{2}$  in Amer. borealis planitie Missouriensis (Nutt.), frequens videtur in Canadâ et Amer. arctica (Lindl.). Chrysopsis alba Nutt. gen. am. 2. p. 152. Aster albus Willd. herb. ex Spreng. syst. 3. p. 528. Doellingeria ptarmicoides Nees ast. 183. Diplopappus ptarmicoides Lindl. adn. mss. Ligulæ lineari-oblongæ albæ (Nutt.). Pappus biformis clavatus et setaceus (Lindl.).

$\beta$ . caule 1-cephalo. — ad lacum superiorem.

$\gamma$ . corymbo composito, foliis subdentatis. — cum var.  $\beta$  legit Pücher ex Lindl. mss.

CXXIII. *BIOTIA DC. non Cass.* — *Eurybiæ* sp. *Cass. dict.* 37. — *Eurybia genuina* Nees ast. 137. — *Asteris* sp. *aut.*

Capitulum radiatum, ligulis fœmineis fertilibus 1-serialibus latiusculis, disco hermaphr. fertili. Recept. alveolæ obtusè dentatæ. Invol. ovati squamæ pluriseriales arcuè imbricatæ adpressæ non appendiculatæ, int. gradatim longiores et angustiores. Stigmata disci hispida acuminata retorta. Achænia glabra vel pubescenti-scabrida elongata tricostrata. Pappus, setis rigidulis filiformibus scabridis subinaequalibus. — Herbæ rhizocarpicæ Boreali-Americanae Asteris facie. Caules erecti apice corymbosi. Folia serrata, radicalia ampla petiolata, caulina alato-petiolata ovata vel oblonga. Ligulæ pallidæ. Discus flavus. — *Biotia Cassinii* est monstrositas *Madia*. *Eurybia* Cass. genuina (dict. v. 16) ex speciebus Novo-Hollandicis tantum constat. Genus ex Asteribus sejunctum dicavi cl. amiciss. Biot qui ex Astrorum studio ad Botanicam physiologicam feliciter migravit.

1. *B. SCHREBERI*, caule glabro apice decomposito-corymboso, foliis grossè serratis suprâ subscabris subtus glabris, radicalibus longè petiolatis ovatis aut suborbiculatis, caulinis inf. secus petiolum subdecurrentibus, super. sessilibus ovatis acutis, capitulis pedicellatis, invol. squamis arcuè imbricatis oblongè obtusis, ligulis invol. duplò longioribus.  $\frac{1}{2}$  in Amer. bor. (Beruh.). Aster Schreberi Nees! syn. (1818). p. 16. Aster subcymosus et A. ambiguus Beruh. ex Nees et Cav. cat. h. taur. 1821. p. 12. A. macrophyllus var. Van Roy. hort. lugd. ex h. Delaroche. *Eurybia Schreberi* Nees! ast. 137. Sæpè cum *B. macrophylla* confusa sed caulis glabritie faciliè distincta, ab etiam et à *B. latifolia* foliis radicalibus non cordatis differt. (v. v. c. à cl. Nees mss.)
2. *B. LATIFOLIA*, caule glaberrimo apice corymboso, foliis glabris, radicalibus caulinisque infer. petiolatis cordatis, super. alato-petiolatis sessilibusve, omnibus serratis acuminatis, capitulis pedicellatis, invol. squamis arcuè imbricatis oblongo-lanceolatis obtusis, ligulis elongatis.  $\frac{1}{2}$  in Amer. bor. (Desf.!) ad montes Alleghany (Raf.!) et circa Bostoniam (Lindl.!) herb.). Aster latifolius Desf. cat. h. par. p. 176. Aster macrophyllus Raf. in litt. *Eurybia macrophylla*  $\gamma$  Nees ast. 140? fortè suad. Lindley mera *B. Schreberi* var. foliis cordatis donata? (v. s.)

3. *B. GLOMERATA*, caule glabro apice thyrsoides, foliis infer. alius lato cordatis alato-petiolatis, super. ovato-ellipticis subsessilibus, omnibus serratis suprâ in ambitu subtus ad costas scabris, capitulis subsessilibus, invol. squamis arcuè imbricatis ovatis obtusis, ligulis vix involucri aequantibus.  $\frac{1}{2}$  in Pennsylvania (Bernh.). *Aster glomeratus* Bernh. ex Nees. *Eurybia glomerata* Nees ast. 139. — *Aster* in hort. Par. anno 1818 sub nom. erroneo *A. corymbosi* cultus huc pertinere videtur, sed ligulæ parum invol. longiores nec breviores. (v. s.)

4. *B. MACROPHYLLA*, caule plus minus hirsuto apice patulo-corymboso, foliis radicalibus petiolatis cordatis, caulinis alato-petiolatis sessilibusve ovatis, omnibus subscabris serratis acuminatis, capitulis pedicellatis, invol. squamis arcuè imbricatis oblongo-lanceolatis obtusis, ligulis elongatis.  $\frac{1}{2}$  in Amer. bor. umbrosis rupestribus à Canadâ ad Virginiam. *Ast. macrophyllus* Linn. sp. 1232. Michx. fl. bor. am. 114. *Eurybia macrophylla* Cass. dict. 37. p. 487. Nees ast. 140. Desf. hort. Par. ed. 3. p. 176. *A. cordifolius* hort. plur. Folia floralia aenta. (v. v. et s.)

$\beta$ . *divaricata*, foliis floralibus rotundatis.  $\frac{1}{2}$  in Amer. bor. (Linn.). *Ast. divaricatus* Linn. sp. 1226. *Ast. macrophyllus carneus* hort. angl. ex herb. Linn. obs. Lindl. in adn. mss.

5. *B. COMMIXTA*, caule infernè glabro supernè pubescenti-scabro patulo-corymboso, foliis ovatis serratis, radic. petiolatis, caulinis super. oblongo-lanceolatis sessilibus viscido-scabris, capitulis pedicellatis, invol. squamis imbricatis subsquarrosis acutis, ligulis elongatis.  $\frac{1}{2}$  patr. ign. verisim. Amer. bor. *Eurybia commixta* Nees ast. 142. An. E. Jussiei Cass. dict. 37. p. 487? Ligulæ lilacinæ. (v. s. c. olim in hort. angl. sub nom. *A. ornati*.)

6. *B. CORYMBOSA*, caule glabro gracili apice laxè corymboso, foliis petiolatis serratis acuminatis subtus secus nervos subsetosis, infer. petiolatis cordatis, mediis ovatis, sup. alato-petiolatis aut subsessilibus, capitulis pedicellatis, invol. squamis imbricatis obtusis, exter. ovatis, ligulis 6-8 invol. duplè longioribus.  $\frac{1}{2}$  in sylvis montanis à Canadâ ad Virginiam Lindl. bot. reg. t. 1532. *Aster corymbosus* Ait. kew. ed. 1. v. 3. p. 207. Nees syn. 17. *Eurybia corymbosa* Cass. dict. 37. p. 487. Nees ast. 143. *Aster cordifolius* Michx. fl. bor. am. 2. p. 114. ex ejus herb. obs. cl. Lindley. (v. s.)

$\beta$ . *alata* (Bartl. fl. phil. 2. p. 114), petiolis latius et frequentius alatis. Varietas ex cl. auct. permanens.

CXXIV. *EURYBIA* Cass. bull. philom. 1818. p. 166. dict. 16. p. 486. non fl. mex. — *Eurybie* sp. Cass. dict. 37. p. 485. — *Eurybie* sp. et *Spongotrichum* Nees. — *Asteris* sp. Labill.

Capitulum pluriflorum radiatum, ligulis 1-serialibus foemineis fertilibus, fl. disci tubulosi 5-dent. hermaphr. Recept. parvum planum aut subconvexum alveolatum. Invol. oblongum imbricatum, squamis chartaceo-membranaceis. Styli radii ramis elongatis erectis obtusis vix hispidulis. Achænium nunquam compressum obovatum aut oblongo-cylindraceum striato-angulatum aut angulato-alatum glabrum vel apice puberulum. Pappus uniserialis, setis scabris præsertim ad apicem, subæqualibus, interdum apice incrassatis. — Frutices omnes è Novâ-Hollandiâ orti. Folia alterna (in 2-3 speciebus opposita) coriacea petiolata aut sessilia, integerrima dentata aut sinuata, sæpè subtus tomentosa. Inflorescentia varia. Discus flavus. Ligulæ albæ violacæ aut carneæ. — *Eurybia* in veterum mythologiâ erat stellarum mater.

#### Sect. 1. BRACHYGLOSSA.

Ligulæ stylos suos non excedentes.

1. *E. BRACHYGLOSSA*, fruticosa, foliis alternis subsessilibus obovatis margine re-

volatis subintegerrimis suprà glabris subtùs tomentosis, capitulis ad axillas foliorum solitariis subsessilibus, ligulis circ. 3 stylo suo fere brevioribus. § in Novâ Hollandiâ orâ orientali. Simillima icon. t. 202 Labill. sed diversa capitulis solitariis nec ternis, sessilibus nec pedunculatis, ligulis brevissimis, invol. squamis nec acuminatis. Achenium glabrum. Pappus 1-serialis. (v. s. comm. à Mus. reg. Par.)

2. *E. AXILLARIS*, fruticosa, ramis tomentosis, foliis alternis oblongo-canescitis obtusis integerrimis, junioribus suprà araneosis demùm glabris subtùs tomentosis, capitulis in axillis fol. solitariis subsessilibus paucifloris, ligulis 3-4 stylum non superantibus. § in Novâ Hollandiâ littore orientali. Folia 9-10 lin. longa, 2 lin. lata. Capitula parva. Folia minima obtusa cava extus involvari bracteiformis. Invol. squamæ obtusæ. Pappus pallidè rufescens subuniserialis. (v. s. comm. à Mus. reg. Par. ex itin. Baudin.)
3. *E. LINEARIFOLIA*, fruticosa, ramulis tomentoso-canis, foliis alternis linearibus utrinquè acutiusculis integerrimis suprà demùm glabris subtùs tomentosis, capitulis in axillis foliorum subsessilibus solitariis et secus ramos foliatis racemum longum constituentibus, ligulis stylo fere brevioribus. § in Novâ Hollandiâ australi. Rami conferti. Folia 10 lin. longa, lineam lata, margine revoluta. Invol. squamæ oblongo-lineares apice summo ad dorsum tomentosæ. Pappus pallidè rufescens. Capit. triplò majus quàm in axillari, 9-florum. Ligulæ 4 parvæ. Achenium villosissimum. (v. s. comm. à Mus. reg. Par. ex itin. Baudin.)
4. *E. CAPITELLATA*, fruticosa, ramulis tomentoso-canis, foliis linearibus basi attenuatis vix acutiusculis integerrimis suprà junioribus subpubeo-mentosis adultis glabris subtùs tomentosis, ramulis apice capitula parva pauciflora distinctè paniculata pauca axillaria et terminalia gerentibus, ligulis paucis stylum non superantibus. § in Novâ Hollandiâ orientali. Valdè accedit ad *E. axillarem* sed capitulis distinctissimè pedicellatis egregiè differt. Invol. squamæ extus subcanescentes lineares obtusæ demùm patentes. Recept. planum vix acrobolatum. (v. s. comm. à Mus. reg. Par. ex itin. Baudin.)
5. *E. DAMPIERI* (A. Cunningham! in litt. 1834. sub Conyza), fruticosa, ramulis tomentoso-canis, foliis alternis linearibus obtusis integerrimis margine subrevolutis suprà glabris subtùs tomentosis, capitulis axillaribus breviter pedicellatis solitariis geminisve, invol. squamis obtusis, ligulis stylo subequalibus. § in insulis Archipelagi Dampieri frequens (King voy. 1818), in ins. Enderby uni ex insulis Rosemary anglis dictis (A. Cunn!), Conyza Novæ-Hollandiæ angustis rosmarini foliis Woodward in Damp. voy. ed. 1729. vol. 3. p. 111. t. 4. f. 3. ex Cunn. Species foliis pollicem longis vix lineam latis ab omnibus affinis distincta! (v. s. comm. à cl. Cunn.)
6. *E. OLIGANTHA*, fruticosa, ramis canescentibus, foliis oblongo-canescitis obtusis integerrimis suprà glabris subtùs canescentibus, capitulis 3-4-floris ad axillas foliorum solitariis subsessilibus. § in Novâ Hollandiâ ad sinum Canum meritorum legit cl. Gaudichaud. Specimina valdè manca ex ejus herb. habet et his tantùm ex habitus analogiâ refero. (v. s. sine fl. et fr.)

Sect. II. ARGOPHYLLÆ. — Eurybiæ heteroclytæ Nees ast. 144.

Ligulæ stylos suos longè superantes. Achenium subcylindricum.

7. *E. CHRYSOPHYLLA*, fruticosa, foliis oppositis petiolatis oblongis basi attenuatis apice acutiusculis integerrimis suprà glabris nitidis sublaevibus anthesis flavo-sericeo-tomentosis, pedunculis axillaribus folio duplò longioribus 1-capitis in corymbum laxum dispositis. § in montanis Novæ-Cambriz australis ad septentr. urbis Bathurst, mense septemb. 1822. omninò defloratam legit cl. All. Cunningham. Aster chrysophyllus Cunn! in litt. 1834. Frutex robustus 6-pedalis. affinis *E. viscosa* sed foliis suprà nec reticulatis nec viscosis et pedicellis 1-cephalis differt. (v. s. comm. à cl. invent.)
8. *E. VISCOsa* (Cass. dict. 37. p. 487. Nees ast. 146), fruticosa, foliis oppositis

petiis oblongiave utrinque attenuatis apice obtusiusculis integerrimis reticulatis subtus albo-tomentosis, corymbis terminalibus subtus villosissimis.  $\S$  in Novâ Hollandiâ ad oram merid. et ins. (Cunn.!) ad Hobart-Town (Cunn.!). Aster viscosus (Labill.!) p. 2. p. 53. t. 203. Balisia Caledoniæ Spreng. syst. 3. p. 569. viscosa Dietr. Folia recentia supra viscosa (Labill.) 3 poll. longa, t. petiol. 2-3 lin., apice obtusa aut vix acuta. Capitula parva paucisquamis lineares acutæ. Ligulæ violaceæ apice 3-dentata. Achenia pendulis villosissima teretiuscula. (v. s.)

causa, fruticosa, foliis alternis et suboppositis obovatis basi cuneatis petiolatis integerrimis coriaceis supra venosis glaberrimis lucubris sericeâ adpressâ confertâ canis, capitulis pedicellatis paucisq. invol. squamis ciliatis, achenio hirtello.  $\S$  in petrosis insulis (Cunn.!) ad montem Wellington 4200 ped. alt. supra mare (Aster persoonioides Cunn.!) in litt. 1834. Affinis E. myrsinoidi. Suffrag. ut videtur humilis. Rami angulati canescentes. Folia sesquipol. lin. lata, obtusa aut submucronata. Pappus albidus copiosus. Flosc. in invol. pauci. (v. s.)

causa, fruticosa, foliis alternis petiolatis ovatis ovalibusve coriaceis supra glabris viridibus subtus tomento adpressissimo argenteis, oppositis.  $\S$  in Novâ Zeelandiâ ad ripas fluminum et spec. ad flumen ferunt cl. D'Urville et Lesson et A. Cunningham qui mecum human. pecuniam. Aster furfuraceus Rich. et Less. fl. Nov. Zeel. p. 216. stipulatus Cunn. in litt. 1834. Species elegans 6-10-petalis E. argentea sed foliis integris faciliè distincta, affinis etiam E. chrysophyllæ sed folia alterna. Invol. ferè turbidatum, squamis 4-seriatis p. ceta apice longius barbata. (v. s.)

causa (Cass. dict. 37. p. 487), fruticosa, foliis alternis petiolatis angulatis acuminatis inæqualiter et argutè serratis coriaceis supra albis ramisque argenteo-sericeis, capitulis paniculato-corymbosis. Umbrosis montanis Nov. Holland. oræ merid. et orientalis et ad Menapi. Aster argophyllus Labill. Nov. Holl. spec. 2. p. 52. t. 201. sag. t. 1563. Arbor 20 ped. alta, truncus 8 poll. diam. (Cunn.!). Odorem moschi redolent! Capitula parva. Invol. squamis imbricatis extus velutinis. Ligulæ 3-5 albe, stylo evid. exserto bifido. Cor. l. apud 5-fida, lobis linearibus. Ovaria glabra. Pappus rufescens l. bi-, disci uni-serialis. Recept. angustum alveolatum. (v. v. c. et

causa, fruticosa, foliis alternis petiolatis ellipticis obtusis basi integerrimis supra glaberrimis subtus petiolis ramis pedunculis invol. lulliter velutino-tomentosis, pedunculis squamulosis parè ramosis, basi squamis imbricatis multiserialibus, inter. gradatim longioribus  $\S$  ad oram orientalem insulâ borealis Novæ Zeelandiæ octob. 1826. Cunningham. Aster cydoniifolius Cunn. in litt. 1834. Flores in flores mihi non satis noti. (v. s. comm. à cl. invent.)

causa, fruticosa, ramis junioribus tomentosis adultis lirato-striatis, de petiolatis lanceolatis basi obtusis apice longè acuminatis repando supra glabris subtus tomentosis, corymbo paniculato.  $\S$  in Novâ Aster liratus (nec lyratus) Sims bot. mag. t. 1509. Diplostegium p. 201. 189. Aster pulverulentus Hoffmannsegg vers. pfl. 1834. p. 165. E. longa, 5-6 lin. lata, petiolo 3 lin. Ligulæ albe circ. 15 lineares, ovata pentagona glabra strigulis minimis secus angulos apice intr. l. meo sensu 1-serialis. (v. s. c.)

causa, fruticosa, foliis alternis vix petiolatis oblongis seu obovatis sub serratis coriaceis supra glabris reticulatis subtus ramulosis, capitulis 15-20-floris paniculatis.  $\S$  in Novâ Hollandiâ a collibus petrosis montium Coraleorum (Cunn.!) et in ins. Van-

Diemen prope Hobart-Town (D'Urville). Aster cruescens Sieb. ! n. 389. Aster herberifolius Cunn. ! in litt. Folia 12-15 lin. longa, petiolo lineam longo. Tomentum rufescens. Pedunculi axillares et paniculati foliis parvis oblongis subintegris ad ramific. instructi. Pappus E. argophyllæ. Ligulæ paucæ lineares fœminæ ex sicco albi 5-dentati nec 5-fidi. Achænium teretiusculum 8-10-striatum, adultum (v. s.)

*β. ilicifolia*, foliis oblongo-lanceolatis utrinquè acutè attenuatis. ! ad septentr. urbem Bathurst in Novâ Cambriâ australi legit cl. ham nov. 1822. Aster ilicifolius Cunn. in litt. 1834. Folia 2-3 p. 6-8 lin. lata. Specim. Sieberianum omnino medium inter *cl. Cunninghami*. (v. s. comm. à cl. inv.)

15. E. MYRSINOIDES (Nees ast. 146. excl. Sieb. syn.), fruticosa, foliis subsessilibus obovatis margine revolutis et subcrenatis suprà glabris, tomentosis, pedunculis 3-cephalis folio paulò longioribus 5-6-floris stylos suos duplò ferè excedentibus. *β* in Novâ Hollandiâ ad Cap. Van Aster myrsinoides Labill. Nov. Holl. sp. 2. p. 53. t. 202. Folia 4-5 l. 2. lin. lata. Invol. imbricatam. Flores 6-8. Ligulæ angustæ acutæ. glabra.

*β. serrata*, foliis argutè serratis subtùs sericeo-tomentosis, ligulis : cor. disci duas multò superantibus. *β* in insulâ Van-Diemen Cunn. An spec. propria ? (v. s. comm. à cl. Lindley.)

16. E. PIMELEOIDES, fruticosa, foliis alternis obovatis obtusis basi cune integris subrevolutis suprà glabris subtùs ramulisque tomentosis terminalibus solitariis, invol. subhemisphærici multiflori squamis ! ext. brevibus subtomentosis, inter. longioribus scariosis glabris, 10 disco duplò longioribus, achæniis sericeo-villosissimis. *β* in atromosis propè Peel's range in Nov. Holl. inter. ad occid. urbem Port-Jack cl. Cunningham jun. 1817. Aster pimeleoides Cunn. ! in litt. 1834. Pimeleæ. Folia referunt icon. Labill. t. 202. sed capitula solitaria non achænia villosissima, etc. Villi achæniis superiores pappum exteñt lant. (v. s. comm. à cl. Cunningham.)

17. E. SUBREPANDA, fruticosa, ramis tomentoso-canis, foliis alternis linearibus obtusis basi attenuatis margine obtusissimè subrepandis ang ribus pulverulento-scabris demùm glabris subtùs tomentosis, cap cellatis ad apices ramorum racemoso-corymbosis. *β* in Novâ H. King prope fretum Bass (itin. Baudin !) et in insulâ Van-Diemen pro Town (D'Urville ! Cunn !). Folia 8-9 lin. longa, lineam lata. Pediculi toti 3 lin. longi. Invol. squamæ oblongæ acutæ dorso tomentosæ flaviores. Pappus rufescens. Achænium pubescens. Ligulæ 8-10. (v. s.)

18. E. GUNNIANA, fruticosa, ramis tomentoso-canescens, foliis al longo-linearibus obtusè sinuato-dentatis subcoriaceis suprà glabris v culatis subtùs densè tomentoso-canis, capitulis longè pedicellatis ramulorum racemoso-corymbosis *β* in insulâ Van-Diemen legit Valde accedit ad E. subrepandam sed folia longiora, etiam juniora ang regulariter dentata, 15 lin. longa, 3-4 lin. lata, capitula dimidiò m gius pedicellata, ligulæ 12-13 (albæ) paulò breviores, fl. disci paucos invol. squamæ ferè glabræ. (v. s. comm. à cl. Lindley.)

19. E. ROSEMARINFOLIA, fruticosa, ramulis canescenti-tomentosis, foliis linearibus mucronatis margine revolutis integerrimis suprà sub subtùs tomentosis, pedunculis corymbosis ultimis trichotomis, palle gatis, invol. squamis lanceolato-linearibus subcarinatis apice calli Novâ Hollandiâ prope Bathurst in collibus calcareis mense novemb Cunningham. Aster rosmarinifolius Cunn. ! mas. 1822. An fortè Soli marinfolia Cass. dict. 56. p. 167 ? Folia 1 ; 2 poll. longa. Capitula gule albæ. Achænium glabrum. (v. s. comm. à cl. invent.)

20. *E. LEDIFOLIA*, fruticosa, ramis glabris subangulatis junioribus pube adpressâ subsericeis, foliis alternis margine revolutis integerrimis obtusis subcylindricis suprâ glabris subtus in sulco sericeo-subhirsutis, pedicellis paucis 1-cephalis, invol. squamis oblongis patulis dorso subhirsutis,  $\frac{5}{2}$  in insulâ Van-Diemen (Gunn!) in rupestribus montis Wellington januario flor. (Cunn.!). *Aster ledifolius* Cunn. in litt. 1834. Capitula majora paucissima et folia dimidiò breviora quàm *E. rosmarinifolia*. Pappus sordide albus pleriserialis, setis exterioribus nonnullis brevissimis. Achænia hispidula. (v. s.)
21. *E. DECURRENS*, fruticosa glabra glutinosa, ramis angulatis, foliis remotis basi angustissimè decurrentibus linearibus coriaceis nunc integerrimis nunc versus apicem dentibus 1-2 repando-dentatis, capitulis ramulos terminantibus subracemosis, invol. squamis linearibus glabris, flosculis paucis, achæniis turbatis sericeo-villosis.  $\frac{5}{2}$  in sterilibus partium Novæ Hollandiæ interiorum ad austr.-occid. flum. Lachlan legit cl. A. Cunningham. *Aster? decurrens* Cunn. in litt. 1834. Fruticulus ferè Ephedræ facie. Stigmata longè exserta apice lanceolata. (v. s. comm. à cl. invent.)

Sect. III. SPONGOTRICHUM *Nees ast.* 176.

*Ligulæ* stylos suos superantes. Achæniuni tetragono-conoideum scabrum. Capitulum campanulatum. Invol. squamæ glanduloso-ciliatæ. Suffrutices foliis linearibus. — An genus proprium?

22. *E. GLANDULOSA*, fruticulosa glabra, foliis alternis sessilibus linearibus margine glandulas protuberantes gerentibus, capitulis pedicellatis corymbosis, invol. squamis margine angustè membranaceis tenuissimè ciliato-serratis.  $\frac{5}{2}$  in paludosis Novæ-Hollandiæ prope lacum Georgii ad Cap Van-Diemen et fretum d'Entrecasteaux. *Aster glandulosus* Labill. Nov. Holl. sp. 2. p. 50. t. 197. *Gallatella glandulosa* Nees ast. 174. *Aster rutæodorus* Cunn. in litt. Folia pollicem et ultrâ longa vix semi-lineam lata patentia. Ligulæ 10-20 albæ oblongæ integerrimæ. Achænia cuneiformi-tetragona pubescenti-scabra. Pappus 1-serialis pallide rufescens, setis apice paulò longius barbellatis. (v. s.)
23. *E. HYGROPHILA*, suffruticosa, caule infernè glabro secus ramos apice pubescente, foliis alternis linearibus utrinquè attenuatis medio paucidentatis aut subrepandis suprâ glabris nervo medio impresso, subtus pube stellatâ brevi velutinis margine subrevolutis, capitulis longè pedicellatis longè racemoso-paniculatis, invol. squamis acutis, pappo biseriali vix inæquali.  $\frac{5}{2}$  abundè in fossâ paludosâ Novæ Hollandiæ, ad littus boreale insulæ Stradbroke (lat. 27  $\frac{1}{2}$ ) mense septembr. floridam legit cl. Cunningham. *Aster hygrophilus* Cunn. in mm. 1828. Ligulæ albæ. Achæniuni glabrum. Caulis rubescens. (v. s. comm. à cl. invent.)
24. *E. TENUIFOLIA*, fruticulosa araneosa aut glabrata, foliis alternis sessilibus eglandulosis integerrimis, ramulis foliosis apice 1-cephalis subcorymbosis dispositis, invol. squamis linearibus acuminatis puberulis.  $\frac{5}{2}$  in Novâ Hollandiâ orientali (h. Mss. Par.) et interiori ad Cæcegangrivier (Cunn.). Accedit ob *E. glandulosam* sed folia glandulis orbata. Ligulæ 9-10-lineares elongatæ. Achænia scabra. Pappus 2-serialis subrufescens, serie ext. brevi multisetâ. (v. s.)
- ß. Bachurstiana*, invol. squamis interioribus subobtusis acutiusculisve nec acuminatis.  $\frac{5}{2}$  in sterilibus circa Bathurst legit cl. A. Cunningham. Frutex floribundus fortè ex cl. invent. distinguendus. (v. s. comm. à cl. Cunningham.)
25. *E. ELAOPHYLLA*, suffruticosa glabra, caule erecto, ramis striato-angulatis, foliis alternis linearibus integerrimis margine subrevolutis, capitulis ad apices ramorum ramulorumque solitariis, invol. squamis lanceolatis sentis dorso subpuberulis, achæniis pubescente subangulato.  $\frac{5}{2}$  in paludosis ad ripas lacus dicti Lagoon in sinu Regis Georgii ad oram merid.-occid. Novæ Hollandiæ, januario flor. legit cl. Cunningham. *Aster elæophilus* Cunn. in litt. 1834. Affinis *E. tenuifoliæ*. Capitula dimidiò minora. (v. s. comm. à cl. inv.)



26. *E. RAMOSISSIMA*, fruticosa ramosissima, ramulis villosis-subtomentosis, foliis lineari-lanceolatis integerrimis acutis minimis supra glabris subtus tomentosis, capitulis terminalibus solitariis sessilibus, invol. squamis imbricatis apice ovato-lanceolatis dorso sublanatis margine nigro ciliato-serratis, achenio villosissimo, pappo biseriali.  $\zeta$  in Novâ Hollandiâ prope Port-Jackson legit cl. Gaudichaud, et in Nov. Camb. interiore ad septentr. flum. Dumaresq in climat ferè tropico collegit cl. Cunningham. Species distinctissima! Folia 2-3 lin. longa. Capitula 5-6 lin. longa et lata. Ligulæ 10-12. Recept. alveolatum. Pappus copiosus albus dentato-scaber. Achenium subangulato-turbatum vili albis erectis copiosis onustum. (v. s.)
27. *E. LEPIDOPHYLLA*, fruticosa, foliis alternis conferto-imbricatis exiguis oblongis crassis margine revolutis integerrimis coriaceis subtus tomentosis, ramulorum abortivorum fasciculatis ramos obtegentibus, capitulis ad apices ramulorum brevissimorum solitariis.  $\zeta$  in Novâ Hollandiâ ad littus australe et Cap. Van Diemen. Aster microphyllus Labill.! Nov. Holl. sp. 2. p. 51. t. 199 non Vent. Aster lepidophyllus Pers. ench. 2. p. 442. Invol. squamæ vix acutæ. Achen. villosum. Pappus albidus, setis exter. dimidiò brevioribus paucioribus scabridis ut interiores. (v. s.)
28. *E. MICROPHYLLA*, fruticosa, ramis setuloso-hirtis, foliis obovatis integerrimis obtusis basi attenuatis supra muricatis subtus tomentosis vix subinermis, capitulis ad apices ramulorum axillarum brevissimorum solitariis secus ramos foliosos racemosis, pappo exteriori brevissimo.  $\zeta$  in Novâ Hollandiâ orientali. Aster microphyllus Vent. malm. 83 (non Labill.). Sieb.! pl. exa. Nov. Holl. n. 338. Diplostegium microphyllum Nees ast. 191. Folia ex ramulis axill. abortivis fasciculata lineam sesquilineamve longa. Capit. 2 lin. longa. Invol. squamæ subobtusæ apice dorso puberulæ. Ligulæ albae angustæ. Achenium puberulum. Pappus duplex, ext. brevis. (v. s. ex Sieb. Gaudich. et in Baudin.)
29. *E. PROPINQUA*, fruticosa, ramis glabris teretibus, ramulis pubescenti-subnervescens, foliis alternis minimis linearibus integerrimis margine subrevolutis supra glabris subtus subpubescentibus, capitulis ramulos terminantibus glabris, squamis linearibus vix acutis, achenio villosis.  $\zeta$  in dumosis planis. Liverpool Novæ Hollandiæ legit cl. Cunningham julio flor. Eandem fort. habeo ex orâ merid. Novæ Holl. in Baudin. expedit. lectam. Aster propinqua Cunn. in litt. 1834. Media inter *E. microphyllum* et *E. ramulosam*. (v. s.)
30. *E. RAMULOSA*, fruticosa, ramis pubescenti-scabris, foliis alternis linearibus lanceolatis acutis integerrimis supra muricato-scabridis subtus tomentosis albidis, capitulis ad apices ramulorum axill. brevissimorum solitariis secus ramos foliatis racemosis, pappo subuniseriali.  $\zeta$  in Novæ Hollandiæ littore merid. ad Port-Jackson, circa Bathurst et Cap. Van-Diemen. Aster ramulosus Labill. Nov. Holl. sp. 2. p. 50. t. 196. Diplostegium ramulosum Nees ast. 193. Aster australis Hortulanorum. Folia 4 lin. longa, semilin. lata. Capit. 2 lin. longa. Invol. squamæ lanceolatæ acutæ membranaceæ dorso vix pubescentes. Ligulæ 6-7 albae ferè lineares. Cor. disci 7-8 albae! Pappi setæ ext. 1-3. (v. s. sp. c. v. c.)
31. *E. ACULEATA*, fruticosa, ramis setuloso-hirtis, foliis alternis linearibus lanceolatis integerrimis supra muricatis subtus tomentosis, capitulis ad apices ramulorum axill. brevissimorum solitariis secus ramos foliatis densè racemosis, pappo subuniseriali.  $\zeta$  in Novæ Holl. Cap. Van-Diemen. Aster aculeatus Labill. Nov. Holl. sp. 2. p. 52. t. 200. Aster exasperatus Link. enam. 1 p. 328. Aster foliosus Hortul. Maxtonia foliosa hort. Bonn. Diplostegium aculeatum Nees ast. 192. Invol. squamæ oblongo-lineares obtusiusculæ apice ciliatæ. Ligulæ paucæ linearibus lanceolatæ. Pappus albus. (v. s.)
32. *E. EPILYRIA*, fruticosa, ramis tomentoso-canis, foliis alternis linearibus lanceolatis basi attenuatis integerrimis supra glabris levibus nervo medio impresso subtus tomentosis canis, capitulis ad apices ramulorum axill. brevissimorum solitariis secus ramos foliatis densè racemosis, pappo subuniseriali.  $\zeta$  in Novæ

Holl. terrâ Van-Diemen. Aster aculeatus herb. Mus. Par. Accedit reverà ad *E. aculeatum* sed ramis tomentosius nec hirtellis, foliis suprà lævibus nec muricatis paulò majoribus etc. differt. Achænium glaberrimum. Pappus ferè 1-ser., setis nempè ext. paucissimis subnullis. (v. s. comm. à Mus. reg. Par.)

† *Species exclusæ.*

- E. COMMIXTA* Nees. = *Biotia commixta*.  
*E. CORYMBOSA* Cass. = *Biotia corymbosa*.  
*E. FULVIDA* Cass. = *Olearia stellulata*.  
*E. JUSSIEUI* Cass. = *Biotia commixta*.  
*E. GLOMERATA* Nees. = *Biotia glomerata*.  
*E. QUERCIFOLIA* Cass. = *Olearia phlogopappa*.  
*E. SCHREBERI* Nees. = *Biotia Schreberi*.

Subdiv. 2. DIPLOPAPPÆ.

Radius 1-serialis. Pappus disci et radii conformis duplex, exterior squamosus, interior pilosus.

CXXV. *OLEARIA* *Manch suppl.* 254. *Nees ast.* 184. — *Diplostephia eurybioidea* *Nees ast.* 193. — *Diplopappi* sp. *Less.* — *Haxtonia hortul.* — *Asteria*, sp. *anc.*

Capitulum multiflorum heterogamum, fl. radii 1-seriatis ligulatis fœmineis, disci tubulosis hermaphroditis 5-dentatis. Recept. alveolatum planiusculum pedunculo apice cavo obconico insidens. Invol. squamæ imbricatæ adpressæ post fructif. patentes. Stam. radii abortiva, disci filamentis brevissimis donata. Stigmata disci cochleariformia. Achænium cylindraceo-tetragonum villosum. Pappus duplex, exterior brevis paleaceo-setosus, interior setosus longus scaber. — Frutices Novo-Hollandici facie Eurybiæ. Folia alterna petiolata ovata repanda seu obtusè dentata aut integra suprà sparsè villosa subtùs ramique densè tomentosa. Pedunculi 1-cephali rigiduli ad apices ramorum 1-3. Discus flavus. Radius albus seu carneus.

\* *Foliis integerrimis subtùs lepidotis.*

1. *O. ELLIPTICA*, fruticosa, foliis alternis petiolatis ellipticis vix acutis basi attenuatis integerrimis suprà junioribus araneosis demùm glabris subtùs adpressè lepidotis subalbidis, corymbo terminali, capitulis longè pedicellatis. 5 in sylvis siccis umbrosis jugi montium Illawara Novæ Hollandiæ detexit cl. A Cunningham. Aster ellipticus A. Cunn. ! in litt. 1834. Habitus ferè Eurybiæ viscosæ sed pappus duplex Oleariæ (v. s. comm. à cl. invent.)

\*\* *Foliis dentatis sinuatisve subtùs tomentosis.*

2. *O. DENTATA* (*Manch l. c.* *Nees l. c.*) fruticosa, foliis alternis petiolatis ovatis repando-dentatis crassis, pedunculis paucis 1-cephalis folio brevioribus. 5 in Novâ Hollandiâ, Aster dentatus Andr. bot. rep. 1. t. 61. Aster tomentosus Schrad. et Wendl. sert. hannov. 8. t. 24. Sieb. ! pl. exs. Nov. Holl. n. 341. Aster ferrugineus Wendl. in florâ 1819. p. 676. Hoffmans. verz. 1826. p. 61. (v. v. c. et s. sp.)
3. *O. ROTUNDFOLIA*, fruticosa, foliis alternis breviter petiolatis ovato-rotundis, inferioribus basi cordatis suprà sparsè hirsutis subtùs ramulisque densè strigiloso-hirsutis hinc indè subrepandis, pedunculis paucis ad apices ramorum 1-cephalis folio triplò longioribus. 5 in Novâ Hollandiâ ad Port-Jackson legit

- cl. Gandichaud. *Diplopappus rotundifolius* Less. in *linnæa* 1831. p. 116. Proximè ex habitu ad priorem accedit. (v. s. comm. à cl. Gandichaud.)
4. *O. QUERCIFOLIA*, fruticosa, foliis alternis subsessilibus obovato-oblongis obtusis suprâ glabris rugosis subtus pube stellatâ tomentosis sinuato-dentatis, dentibus latis obtusis, pedunculis axillaribus folio longioribus 1-3-cephalis. 3 in Novâ Hollandiâ (Sieb.!), in paludosis rori cataractæ Campbell in montibus Cœruleis expositis (Cunn.!). *Aster quercifolius* Sieb.! pl. exs. Nov. Holl. n. 340. *Aster sinuatus* A. Cunn.! in litt. 1834. Folia 15-18 lin. longa, 6-8 lin. ad apicem lata. Pedunculi bipollic. Achænia acutè pentagona adulta glabra. Pappi setæ subbiser. pallide rufescentes scabræ, setæ exter. brevès paucissimæ. (v. s.)
5. *O. PHLOGOPAPPA*, fruticosa, foliis alternis breviter petiolatis oblongo-lanceolatis suprâ glabris subtus tomentosis sinuato-dentatis, dentibus obtusis, corymbis paniculatis. 3 in Novâ Hollandiâ ad Cap. Van-Diemen. *Aster phlogopappus* Labill.! Nov. Holl. sp. 2. p. 49. t. 195. *Aster phlogotrichus* Spreng. syst. 3. p. 525. *Eurybia quercifolia* Cass. dict. 16. p. 47. *Diplostephium phlogotrichum* Nees ast. 186. Ligulæ circ. 10 violaceæ 3 dentatæ. Achænia 5-angulata pubescenti-scabra. Pappus rufescens biserialis scaber. (v. s.)
6. *O. STELLULATA*, fruticosa, foliis alternis in petiolum brevem attenuatis oblongo-lanceolatis acuminatis suprâ subscabridis subtus pube stellatâ densè tomentosis grossè dentatis, dentibus mucronatis, corymbis paniculatis. 3 in Novâ Hollandiâ ad Cap. Van-Diemen. *Aster stellulatus* Labill.! Nov. Holl. sp. 2. p. 50. t. 196. *Eurybia fulvida* Cass. dict. 16. p. 48. *Diplostephium stellulatum* Nees ast. 187. Proximè ad *O. phlogopappam* accedit. Folia 3-4 poll. longa, 4-6 lin. lata. Invol. squamæ imbricatæ lineares acutæ. Ligulæ ex Labill. circ. 15, et tubuli circ. 20. Achænia matura pauciora obovato-teretiuscula obtusè tetragona villosa. Pappus, setis biserialibus scabris subrufescentibus, exter. brevibus subpaleaceis. Recept. parçè convexum alveolatum. (v. s.)

CXXVI. *DIPLOSTEPHIUM* Cass. dict. 37. p. 486. — Doellingeria sp. Nees. — *Diplopappi* sp. Less. — *Asteris* sp. auct.

Capitulum multiflorum radiatum, ligulis fœmineis fertilibus 1-seriilibus, disco hermaphrodito. Invol. subcylindraceum disco brevius, squamis pauciseriatis imbricatis adpressis coriaceis margine membranaceis. Recept. planum alveolatum, alveolis dentatis. Stigmata subulata hispida. Achænia elongata angulata hispidula. Pappus duplex, ext. setulosus brevis, inter. elongatus filiformis aut apice clavellatus scaber. — Herbæ perennes aut frutices asteriformes. Species omnes Americane. Folia alterna. Discus flavus. Ligulæ albæ aut cœruleæ. Sectiones duæ fortè in posterum genericè distinguendæ.

#### Sect. 1. EUDIPLOSTEPHIUM.

Pappi interioris setæ plures apice clavellatæ. — Herbæ rhizocarpion Boreali-Americane facie Asteris.

1. *D. UMBELLATUM*, caule angulato scabro apice corymboso composito, foliis oblongo-lanceolatis utrinquè attenuatis margine scabridis, invol. imbricatis, achæniis pubero-hirtis. 2 in paludosis et ad fluminum ripas à Canadâ ad Carolinam. *Aster umbellatus* Ait. kew. ed. 1. v. 3. p. 199. Hoffm. phytogr. II. p. 74. t. b. f. 2. ex Pers. *Chrysopsis amygdalina* Nutt. gen. 2. p. 263. anal. syn. *Aster amygdalinus* Ell. sketch 2. p. 367. *Doellingeria umbellata* Nees ast. 178. Pappus rufus. Ligulæ albæ. (v. s.)
2. *D. AMYGDALINUM* (Cass. dict. 37. p. 486), caule subangulato glabro apice corymboso-paniculato, foliis lanceolatis oblongiæ margine scabris, invol. laxo, ligulis 8-10, achæniis hirtellis. 2 in umbrosis petrosis à Canadâ ad Ca-

rolinam. *Aster amygdalinus* Lam. dict. 1. p. 305. Lindl. bot. reg. t. 1517. *Chrysopsis humilis* Nutt. gen. 2. p. 153. excl. syn. *Aster humilis* Willd. sp. 3. p. 2038. *Aster divaricatus* Spreng. syst. 3. p. 529. excl. syn. *Doellingeria amygdalina* Nees ast. 179. Ligulæ albæ. Pappus rufus, setis plerisque apice incrassatis. (v. v. et s.)

2. *humilis*, caule angulato glabro simplici dense folioso, foliis sessilibus utrinque attenuatis, corymbo simplici oligocephalo. 2/ in Terrâ Novâ legit el. de Lapilayc. (v. s.)

3. *D. cornifolium*, caule tereti glabro parce paniculato, foliis elliptico-oblongis utrinque acuminatis breviter petiolatis ciliatis integerrimis glabris, corymbi ramis nudis 2-3-cephalis, invol. laxè imbricato, acheniis glabris. 2/ in Pennsylvania (Muhl.), à Canadâ ad Carolinam (Michx.). *Aster cornifolius* Muhl. in Willd. sp. 3. p. 2039. *Aster infirmus* Michx. fl. Bor. Am. 2. p. 109 (certè ex Lindl.). *Aster humilis* Ell. sketch 2. p. 366. — Pluk. alm. t. 79. f. 1. Ligulæ 7-8 albide. Pappus rufus, ext. brevis. E fol. basi decurrentiæ tenuissime interdum puberulæ. Achænia obovata subcompressa glabra. (v. s. comm. à cl. Torrey.)

4. *D. acuminatum*, caule simplici angulato-flexuoso scabro, foliis lato-lanceolatis basi angustatis et integerrimis apice grossè serratis acuminatis margine scabridis ceterum glabrisculis subsessilibus, paniculâ corymbosâ laxâ, pedunculis nudis folio longioribus, invol. squamis linearibus laxis disco brevioribus, achenio sparse glanduloso-pubero. 2/ in sylvis Amer. bor., in Canadâ (Michx.), agro Bostoniensi (Big.), Noveboracensi (Tor.), ad saltus Niagarae (Janin!) in montibus Alleghany (Raf.). *Aster acuminatus* Michx. fl. Bor. Am. 2. p. 109. Hook. bot. mag. t. 2707. *A. diffusus acuminatus* Pers. Ligulæ albæ. Pappus pallide rufus, setis exter. brevibus paucis. Affinis *D. cornifolio*. (v. s.)

5. *D. dichotomum*, caule pubescente, foliis arcè sessilibus ovalibus obtusis pubescentibus, corymbi dichotomi ramulis nudis elongatis, invol. squamis linearilanceolatis pubescentibus achenio valde hirsuto vix longioribus. — in humidis pinguibus insulæ Paris Amer. bor. *Aster dichotomus* Ell. sketch 2. p. 366. Ligulæ 10-16 albæ purpureo tinctæ. Pappus duplex.

6. *D. obovatum*, totus brevè et molliter pubescens, caule tereti subsimpliciter corymboso, foliis obovatis obtusis integerrimis subsessilibus margine scabris, invol. imbricato pubescente, ligulis circ. 10, achenio angulato hispido. 2/ in Georgiae fruticetis ad margines paludum propè Savannah. *Chrysopsis obovata* Nutt. gen. 2. p. 152. *Diplostephium boreale* Spreng. syst. 3. p. 544. excl. syn. *Doellingeria obovata* Nees ast. 182. *Aster obovatus* Ell. sketch 2. p. 368. Ligulæ et disci cor. ochroleucæ. Pappus ferrugineus duplex.

## Sect. 11. AMPHISTEPHIUM. — *Diplostephium* Kunth nov. gen. am. 4.

p. 97.

Pappi interioris setæ filiformes nec clavellatæ. — Frutices Mexicani foliis linearibus. — Sectio media inter *Diplostephium* et *Diplopappum*, fortè genericè separanda, sed mihi non satis nota.

7. *D. MORANENSE* (Nees ast. 198), fruticosum, ramis foliosis glabrisculis, foliis singulato-linearibus callosomucronatis planis subcarnosis, infer. apicem versus mucronato-dentatis, ramis florigeris subcorymbosis minute foliosis, invol. hemisphaerici imbricati disco brevioris squamis inf. subulatis, cæter. lanceolato-cuspidatis. 5 in montibus Mexicanorum propè fodinam Moran rarissimum. *Aster Moranensis* H. B. et Kunth nov. gen. am. 4. p. 93. Ligulæ paucæ albæ siccitate revolutæ. Achænia compresso-quinquangularia hirsuto-sericea. Pappus duplex exterior setulosus.

8. *D. LAVANDULÆFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 97. t. 335), fruticosum, foliis linearilanceolatis margine revolutis integerrimis subtus caule involucriaque albo-tomentosis, capitulis racemoso-confertis. 3 in arenosis

exstis ad radices montium ignivomorum Iliniæ et Cotopaxi prope Mula. Nees ast. 189. Aster lavendulaceus Willd. herb. Frutex orgyalia. Capit. 9-10 lin. diam. Ligula 30-35 albæ. Discus pallidè luteus. Stigm. lineari-subulatis. Pappus duplex, ext. paleaceus brevis, int. setosus ferrugineus. Achæn. variisimil. non compressa. An hujus generis?

† *Species exclusæ.*

- D. ACULEATUM Nees = Eurybia aculeata.
- D. EXTENUATUM Nees = Diplopappus extenuatus.
- D. FILIFOLIUM Nees = Diplopappus filifolius.
- D. FRUTICULOSUM Nees = Diplopappus fruticosus.
- D. GRANDIFLORUM Spr. = Andromachia grandiflora.
- D. HISPIDUM Nees = Diplopappus hispidus.
- D. IGNIARIUM Spreng. = Andromachia igniaria.
- D. LIRATUM Nees = Eurybia lirata.
- D. MICROPHYLLUM Nees = Eurybia microphylla.
- D. MELASTOMOIDES Spreng. = Andromachia melastomoides.
- D. NUBIGENUM Spreng. = Andromachia nubigena.
- D. OBTUSATUM Nees = Diplopappus obtusatus.
- D. OCHRACEUM Nees = Tetramolopium ochraceum.
- D. PHLOGOTRICUM Nees. = Olearia phlogopappa.
- D. RAMULOSUM Nees = Eurybia ramulosa.
- D. SÆSILIFLORUM Spr. = Andromachia sæsiliflora.
- D. SOLIDAGINEUM Spr. = Andromachia solidaginea.
- D. STELLULATUM Nees = Olearia stellulata.
- D. TRANTIFOLIUM Nees = Diplopappus filifolius β.
- D. VOLUBILE Spr. = Andromachia volubilis.

CXXVII. CALLISTEPHUS Cass. dict. 37. p. 464 et 491. Nees ast. 221. — Callistemma Cass. bull. philom. 1817. p. 32. dict. 6. suppl. 45. ast. et op. phyt. 4. 7. — Asteris sp. Linn. — Diplopappi sp. Less. — Chrysopsidis sp. Nutt.?

Capitulum multiflorum heterogamum, fl. radii ligulatis femininis in statu natur. 1-serialibus, disci hermaphroditis tubulosi 5-dentatis. Recept. latum subconvexum breviter alveolatum. Invol. 3-4-seriale patens, squamis ciliatis obtusis, bracteis foliaceis involucri brevioribus cinctum. Stigmata disci apice ovata. Achænia obovato-cuneiformia compressa hirtella. Pappus duplex uterque 1-serialis, exter. brevissimus paleaceo-setaceus setis in coronam subconcretis, inter. setis longis filiformibus scabris deciduis. — Herba Chinensis annua erecta ramosa, ramis apice 1-cephalis. Folia alterna sessilia. Capitula ampla aut floribus omnibus ligulatis aut omnibus tubuloso-amplatis plena, disco flavo, ligulis tubulosis amplatis omnes colores flavo excepto exhibentibus.

§. 1. *Invol. squamis exterioribus patulis.*

1. C. CHINENSIS (Nees l. c.), foliis infer. spathulatis petiolatis, medillis rhombico-lanceolatis, summis oblongis, omnibus grosse dentatis plus minus ciliatis, ramis 1-cephalis elongatis, invol. squamis foliaceis subquarternis longe ciliatis. (1) in China et Japonia, nunc in hortis totius orbis cultus, gallicè *Reine Marguerite*. Aster Chinensis Linn. sp. 1232. Callistemma hortense Cass. bull. phi-

- lom. 1817. p. 39, dict. et opusc. atl. t. 7. *Callistephus hortensis* Cass. dict. 37. p. 491. *Diplopappus Chinensis* Less. syn. 165. — Dill. elth. t. 34. (v. v. c.)
2. *C. ? nuxensis* (Lindl. in litt. 1835), foliis integerrimis oblongo-linearibus ciliatis, ramis elongatis monocephalis, invol. squamis lanceolato-linearibus acuminatis ciliatis aequalibus disco longioribus squarroso-patentibus. (2) in Sibiria. *Aster* Gmel. fl. sib. 2. p. 189. t. 81 excl. descr. *Species mihi planè ignota*. †.

§. 2. *Invol. squamis exterioribus erectis*. — *Poloa DC. in Guill. ann. 1833, v. 2. p. 514 et in Wight contr. p. 10.*

3. *C. WIGHTIANUS*, foliis sessilibus oblongo-linearibus integris aut subserratis breviter mucronatis, ramis foliosis apice compressis minutè puberulis, invol. squamis exter. foliaceis lineari-oblongis eciliatis vix interiores superantibus. — in Ind. orient. ad Dindigul legit cl. Wight. *Erigeron Wightianum* Wall.! cat. n. 3284. comp. n. 394. *Erigeron dubium* Spreng. in litt. ad Wall. *Buphtbalmum Wight!* herb. *Poloa Wightiana* DC. l. c. Variet glabriusculus aut subpubescens. Ligula fertiles aut rariùs substeriles, pappo ext. nunc cupuliformi nunc subnullo. (v. a. comm. ab hon. Coet. merc. angl. Ind. or.)
4. *C. HEYNEANUS*, foliis sessilibus oblongis acutè serratis pube minutissima utrinquè ramisque hirtellis, invol. squamis exter. foliaceis lineari-oblongis eciliatis inter. paulò superantibus. — in India orient. legit cl. Heyne. *Aster Indicus* Heyne in Wall.! cat. n. 2971. comp. n. 81 non Linn. *Species priori proxima sed hirtutis brevi et confertà, foliis serratis, ramis non compressis et habitu diversa*. (v. a. comm. ab hon. Coet. merc. angl. Ind. or.)

† *Species exclusa.*

*C. INDICUM* Don. = *Asteromea Indica*.

CCXVIII. DIPLOPAPPUS. — *Diplopappi* sp. Cass. dict. 13. p. 308. Less. syn. 163. — *Rhinactina* Less. syn. 162, non Willd. — *Asteris* sp. auct. — *Diplostephii* sp. Nees.

Capitulum multiflorum radiatum, ligulis 1-serialibus fœmineis, fl. disci hermaphr. 5-dentatis. Recept. planum subfoveolatum. Invol. squamæ imbricatæ. Achænia oblonga compressa. Pappus disci et radii duplex scabridus, ext. brevis setulosus, inter. longus setosus. — Herbæ aut suffrutices patriâ et habitu varii. Genus (forsan ideò in posterum dividendum) differt à *Diplostephio* achæniis compresso-planis nec teretiusculo-angulatis; à *Callistepho* achænio evidentius compresso et pappo exter. setoso nec coroniformi; à *Chrysopside* pappo exter. potius setoso quàm paleaceo et ligulis à disco discoloribus albis cœruleis seu purpureis nec concoloribus flavis. — An fortè *Asterosperma* Less. syn. 389 mihi ignotum huc etiam referendum?

- §. 1. *Pedunculares*. — *Diplostephia peduncularia* Nees ast. 195; *caulibus fruticosis, invol. imbricati squamis adpressis lined dorsali elevatâ et glandulis 1-2 secus lineam donatis margine subscariosis foliis 1-nerviis glabris integerrimis*. (1)
2. *D. PAUCICULOSUS* (Less. in linnæâ 1830. p. 171. syn. 164), foliis lineari-cuneiformibus integerrimis planis glanduloso-punctatis glabris, ramulis lateralibus apice nudis 1-cephalis, pedunculis folia triplo superantibus, involuero discum æquante, ligulis lanceolatis involuero duplo longioribus. 3 ad Cap. Bonæ-Spei

(1) Si aliquis vellet aut quinque hujus divisionis species in unicam conjungere aut ex iis genus proprium conficere, non repugnarem.

(Eckl. et Zeyh.). *Diplostephium fruticosum* Nees ast. 194. *Diplostephium longipes* Cass. dict. 56. p. 171. *Aster fruticosus* Linn. sp. 1225. Thunb. fl. cap. 687. Burm. ! prod. 27. *Aster fruticosus* Willd. sp. 3. p. 2018. Jacq. fragm. p. 9. t. 5. f. 4. Sims bot. mag. t. 2286. — Comm. h. Amst. 2. t. 27. Pedunculi pollicares ramos praeritum laterales terminantes fastigiatim dispositi. Ligulae lilacinæ seu albidæ. (v. s.)

℘. ligulis lateraliter ad apicem dente acuto utrinquē incisus elongatis linearibus. — Hook. bot. mag. t. 2718.

2. *D. EXTENUATUS* (Nees ast. 195 sub *Diplostephio*), foliis lineari-cuneiformibus integerrimis planis glabris grossè glanduloso-punctatis, ramulis lateralibus apice nudis 1-cephalis, pedunculis folia triplò superantibus, invol. disco breviori, ligulis lanceolatis involucri duplò longioribus. ℥ ad Cap. Bonæ-Spei ad Glorikloof (herb. Deless.) et in monte Tabulari (Eckl. in herb. Danant). *Aster fruticosus* var. auct. *Aster coriaceus* hortul. fortè *Diplostephium fruticosum* vera varietas loco pinguiori nata. Huc potius quàm *D. obtusatum* pertinet icon Plak. mant. t. 3 fo. f. 19. (v. s.)

3. *D. OBTUSATUS* (Less. in linnæa 1831. p. 118), foliis lineari-cuneatis integerrimis planis glabris pellucido-punctatis, ramulis lateralibus apice 1-cephalis, pedunculis subnudis vix folia superantibus, involucri disco breviori, ligulis linearibus oblongis involucri vix longioribus. ℥ ad Cap. Bonæ-Spei ad Blear-Bery ex herb. Deless. *Diplostephium obtusatum* Nees ast. 196. *Aster obtusatus* Willd. sp. 3. p. 2018. excl. syn. Thunb. *Aster rosmarinifolius* Spreng. syst. 3. p. 521. — *Aster fruticosus* ℘ Willd. sp. 3. p. 2018 et reverà fortè *D. externati* varietas brevi-pedunculata. Variat foliis obtusis aut acutiusculis. (v. s.)

4. *D. FILIFOLIUS*, foliis lineari-filiformibus glanduloso-punctatis glabris, ramulis lateralibus apice 1-cephalis, pedunculis subnudis folia duplò superantibus, involucri disco subæquante, ligulis involucri duplò longioribus. ℥ ad Cap. Bonæ-Spei (Eckl. et Zeyh.) Dreg. ! *Diplostephium filifolium* Nees ast. 197. *Aster filifolius* Vent. malm. (1804). t. 82. — *Aster* Eckl. ! coll. exa. Ligulae albæ. Folia 6-8 lin. longa. Pedunculi pollic. et sesquipollic. Glandulae foliorum et scrobuli tantum apparent. (v. s.)

℘. *teretifolius*, foliis pedunculis ligulisve brevioribus. ℥ ad Cap. Bonæ-Spei. *Diplost. teretifolium* Nees ast. 198. *Diplopappus teretifolius* Less. in linnæa 1831. p. 118. *Aster fruticosus* Krebs ! coll. exa. n. 192. *Aster fruticosus* var. Thunb. fl. cap. 687. (v. s.)

5. *D. ELONGATUS*, foliis lineari-filiformibus eglandulosis glabris, axillis andè ramis apice longè denudatis monocephalis folia triplò superantibus, invol. squamis dorso ad apicem glandulosis, ligulis involucri duplò longioribus. ℥ ad Cap. Bonæ-Spei ad Zneenroberge alt. 200-300 ped. legit cl. Dreg. ! Affinis *D. filifolio* sed folia non fasciculata pollicem nec semipollicem longa. Pedunculi tripollicares. Invol. disco dimidio brevius, etc. (v. s.)

§. 2. *Calimeridei, caulibus herbaceis, foliis latis sæpius trinerviis et dentatis, invol. squamis pauciserialibus vix imbricatis eglandulosis.*

6. *D. ASPER* (Less. syn. 163), caule herbaceo adscendenti-erecto simplici acutè apice longè aphylo 1-cephalo, foliis oblongo- aut ovali-lanceolatis triplinerviis integerrimis aut dente grosso 1-2 hinc inde notatis utrinquē asperis ciliatis. invol. squamis disco subæqualibus planis acuminatis margine scariosis. ℥ ad Cap. Bonæ-Spei in graminosis prope Galgebosch (Th.), prope Hangklipp, Mueb. et Mair.), in Cafraria (Krebs ! coll. n. 173), in col. orient. non longè à fana Gauritz (Burch. n. 4740), in distr. Albany et in territorio cesso (Eckl. et Zeyh.), in Zneenroberge et Willberge (Dreg. !). *Calendula hispida* Thunb. cap. 704 et *Leyssera pilosella* ejusd. herb. sp. dextr. et lav. ex Less. *Diplostephium hispidum* Nees ast. 163. Capitulum ferè *Asteris alpini*. (v. s.)

7. *D. ROYLEI*, caule herbaceo erecto subsimplici pedunculisque hispidis. fl.

oblongis acuminatis sessilibus aliis integerrimis aliis apice 3-5-dentatis supra glabris subtus et margine scabris obscure trinerviis, invol. squamis glabris disco brevioribus margine scariosis. — in Indiæ prov. bor. occid. legit cl. Royle. Valdè accedit ad *D. asperum*. Ligulæ extus revolutæ ex sicco albide. Achænia villosa. Pappus crassè barbellatus. Cl. Lindley in adn. mss. habet *Diplopappum*, quemdam Roylei fortè à nostro diversum sed mihi ignotum. (v. s. comm. à cl. invent.)

8. *D. molluscus* (Lindl. adn. mss.), caulibus simplicibus adscendentibus moncephalis pubescenti-scabris, foliis oblongis integerrimis scabriusculis subtus strigoso-pilosis, capitulis longè pedunculatis, invol. hemisphærici squamis acuminatissimis, exter. basi pubescentibus, inter. membranaceis.  $\frac{1}{2}$  in Indiâ orient. ad Sirmore. Aster molluscus Wall.! cat. et herb. n. 2972. comp. n. 82. (v. s. comm. ab hon. Cœt. merc. angl. Ind. or.)
  9. *D. asperulus*, caule herbaceo erecto dichotomè ramoso tereti glabro, foliis ovatis acuminatis grossè serratis penninerviis ciliatis supra sparè scabris, radicalibus petiolatis, caul. basi cuneatis, corymbo laxo, pedicellis nudis, invol. squamis oblongo-linearibus obtusis glabris, ext. foliaceis laxis, int. disco brevioribus subscariosis.  $\frac{1}{2}$  in montibus Indiæ orientalis ad Kamaon. Aster asperulus Wall.! cat. et herb. n. 2968. comp. n. 78. Nees ast. 281. Achænia sparè villosa. Ligulæ elongatæ. (v. s. comm. ab hon. Cœt. merc. angl. Ind. or.)
  10. *D. asperimus*, caule herbaceo stricto ramoso scabro, foliis elliptico-ovatis lanceolatisve acuminatis sessilibus trinerviis utrinquè scaberrimis, corymbo patulo foliato, invol. squamis oblongis obtusis adpressis ciliatis, achæniis sericeo-villosis, pappi setis exter. paucis brevibus.  $\frac{1}{2}$  ad Narrain-Hotty et alibi in Nepaliâ legit cl. Wallich. Aster asperimus Wall.! cat. et herb. n. 2970. comp. n. 80. Aster trinervius Roxb. cat. h. calc. 61. Don prod. fl. nep. 177. Aster asper Ham. herb. Galatella asperima Nees ast. n. 173. Cineraria canescens Wall.! herb. Ligulæ fertiles à Galatellâ excludunt. Recept. in centro alveolato-timbrilliferum. Achænia plano-compressa nec ut in *Diplostephio* angulata. Pappus rufus primâ fronte 1-serialis sed, mon. cl. Lindl. verè sed obscure biserialis, setulis inter achæniæ villos super. subconfusis. Habitus ferè *Dœllingeriæ* sed setæ nec clavellatæ nec 1-seriales. (v. s.)
  11. *D. caucasicus*, caule herbaceo erecto striato pubescenti-scabro, foliis ovato-sen oblongo-lanceolatis serratis acuminatis basi trinerviis utrinquè (præc. margine et secus nervos) scabridis, capitulis 1-7 caulem ramoque terminantibus, invol. subæqualis squamis lineari-acuminatis glabriusculis ciliatis, achæniis subtrigonis hirtis, pappi setis exter. paucis brevibus.  $\frac{1}{2}$  in alpinis Caucasi. (M. Puschk.) ad Gurjel (Szow.! ex Fisch. et M.). Aster caucasicus Willd. sp. 3. p. 2020. Bieb. fl. et suppl. n. 1729. Nees ast. p. 35. Pappus duplex ritè observ. cl. Meyer. Ligulæ purpurascens. Herba pedalis. (v. s. comm. à cl. Fisch. et Mey.)
- §. 3. *Amelloidei*, Nees ast. p. 199, sub *Diplostephio*; caule herbaceo-suffrutescente, foliis linearibus rigidulis ciliato-scabris, involucris eglandulosis imbricatis, capitulis subcorymbosis.
12. *D. linearifolius* (Lindl.! adn. mss.), caule stricto apice racemoso, foliis linearibus rigidis glabris margine scaberrimis serrulato-ciliatis, invol. turbinatis squamis linearibus acutis patulo-imbricatis margine coloratis.  $\frac{1}{2}$  in Amer. bor. spec. in agro Noveboracensi. Aster linearifolius Linn. sp. 1227. Aster rigidus Michx. fl. bor. am. 2. p. 110. *Diplostephium* linearifolium var. Nees ast. 199. *Chrysopsis* linearifolia Nutt. gen. 2. p. 152? Ligulæ cœruleæ. Achænia compressa villosa. Caulis imâ basi suffruticulosus. (v. s.)
  13. *D. rigidus* (Lindl. adn. mss.), caule stricto racemoso, ramis elongatis 1-cephalis, foliis linearibus rigidis margine scaberrimis serrulato-ciliatis, invol. hemisphærici squamis linearibus acutis, exterioribus laxis à foliis pedicelli parum diversis.  $\frac{1}{2}$  in Amer. bor., spec. ex Carolinâ et Georgiâ recepti. Aster



*rigidus* Linn. sp. 1228. *Aster linarifolius* Michx. fl. 2. p. 110. *Aster pulcherrimus* Lodd. bot. cab. 1. t. 6. *Diplostephium linarifolium* var. Nees est. 199. — Leon Pluk. t. 14. f. 7. hinc fortè pertinet. — Cl. Lindleyum qui specim. *Linnaeanum* vidit secutus sum in dist. hujus et *linarifolii*. (v. s.)

14. *D. INCANUS* (Lindl. bot. reg. t. 1693), totus incanus, caule suffruticoso, ramis longis corymbose 1-cephalis apice nudis, foliis linearibus mollibus distantibus obtusis mucronatis, invol. ovati glandulosi squamis linearibus multiseriatis, exter. squarrosis.  $\frac{1}{2}$  in Californiâ et ad oram occid. Americæ bor. Hook. bot. mag. t. 338a. Caulis a  $\frac{1}{2}$ -pedalis à basi ramosus pube brevi mollis, foliis valde denudatis. Ligulæ amonè violaceæ. Proximus *D. rigido* et *linariæ-folio*. Ovarium teres villosum, undè potius ad *Diplostephium* referendus? (v. s.)

15. *D. LUTESCENS* (Lindl. adn. mss.), caule simplici corymbose, ramulis sub-1-cephalis subaphyllis, foliis oblongo-linearibus basi longè angustatis subintegris scabris, invol. pubescentis squamis oblongo-lanceolatis obtusis.  $\frac{1}{2}$  in Americâ arcticâ. Radius lutescens. Proximus *Heleastro albo* ex Lindl. An fortè *Chrysopsis* sp.?

16. *D. LEUCOPHYLLUS* (Lindl. adn. mss.), fruticosus? undiquè lanatus, ramis brevibus 1-cephalis, foliis crassis ovalibus acutis crenatis in petiolum angustatis, invol. squamosi squamis linearibus membranaceis, superioribus ciliatis? in Californiâ. Pappus exter. brevis, inter. valde inæqualis subulna *Achenia* tomentosa fusiformia. (Lindl.)

### † Species exclusæ.

- D. CHAMISSONIS* Less. = *Aplopappus Chamissonis*.  
*D. CHINENSIS* Less. = *Callistephus Chinensis*.  
*D. CHRYSANTHEMIFOLIUS* Less. = *Aplopappus chrysanthemifolius*.  
*D. CORONOPIFOLIUS* Less. = *Aplopappus coronopifolius*.  
*D. DELPHINIFOLIUS* Cass. = *Polyactidium delphinifolium*.  
*D. DUBIUS* Cass. = *Stenactis annua*.  
*D. ERICOIDES* Less. = *Aplopappus ericoides*.  
*D. GRAMINIFOLIUS* Less. = *Neja gracilis*.  
*D. PRUNELLOIDES* Less. = *Aplopappus prunelloides*.  
*D. ROTUNDIFOLIUS* Less. = *Olearia rotundifolia*.  
*D. SERICEUS* Less. = *Aplopappus sericeus*.

### CXXXIX. RHINACTINA Less. syn. 162 non Willd.

Omnia *Diplopappi* sed corollæ disci inæqualiter 5-dentatæ et idè ferè subbilabiata, pappi serie externâ paucisetâ. — Herbariæ Sibiricæ perennes, rhizomate lignoso. Folia glaucescentia brevissimè subhirtella integerrima, radicalia obovata in petiolum attenuata obtusa aut submucronata, caulina pauca. Caulis simplex 1-cephalus aut parcè ramosus oligocephalus. Invol. squamæ siccæ glabræ. *Achenia* villosa. Ligulæ ceruleæ. — An genus à *Diplopappo* satis diversus? *Habitus* ferè *Asterum* alpigenorum sed pappus duplex.

1. *R. LIMONIFOLIA* (Less. in linnæa 1831. p. 119. syn. p. 162), foliis radicalibus obovatis obtusis trinerviis longè petiolatis, caulinis ovali-oblongis sessilibus. caulibus capitula 2-4 longè pedicellata gerentibus, invol. squamis obtusisacutis.  $\frac{1}{2}$  in rupestribus graminosis Sibiricæ Altaicæ ad montem Kartscham. *Aster* obovatus Mey. in Ledeb. fl. alt. 4. p. 95. Ledeb. ic. alt. 5. p. 492. (v. s. com. à cl. Meyer.)

1. *UNIFLORA* (Bunge ex herb. Acad. imp. Petrop.), foliis radicalibus oblongo-bovatis acutis uninerviis breviter petiolatis, caulibus paucis oblongis basi squamatis, caulibus simplicibus 1-cephalis, invol. squamis acuminatis.  $\frac{1}{2}$  in montibus Altaicis. Caules floridi vix bipollicares folia radicalia paulatim succedentes.

## CXXX. NOTICASTRAUM.

Capitulum radiatum, ligulis 1-serialibus fœmineis fertilibus linearibus, fl. disci tubulosis 5-dentatis bisexualibus sterilibus. Invol. 3-seriale, squamis linearibus acuminatis, extimis brevibus, intermed. coloratis nitidis subscariosis disco ferè longioribus. Recept. planum alveolatum. Antheræ ecaudatæ. Styli rami in radio et in disco elongati subdorso hispidi. Achænia oblongo-linearibus pube sericeâ villosa sessilibus subrostrata, radii fertilia compressa nervoso-striata, disci angustissima. Pappus pluriserialis persistens, setis rigidulis scabris extimis brevioribus. — Herba perennis Chilensis puberulo-glandulosa. Rhizoma frutescens. Caules subsimplices teretes adscendentes 6-7 poll. in radicalia in petiolum attenuata, caulina sessilia subsemiamplexilina lanceolata mucrone calloso apiculata integerrima. Capitula paucapices ramulorum solitaria subcorymbosa. Ligulæ rubentes disco vix longiores integræ, juniores convolutæ. Discus flavus. — Genus disco sterili et stigmatibus subulatis distinctissimum. Affinitas subina. Nomen ex *NOTICASTRA*, australis, et *ASTER*, quasi *Aster australis*.

1. *ADSCENDENS*.  $\frac{1}{2}$  in Cordilleris Chilensibus legit indef. Claud. Gay et mecum commun. (v. s.)

## CXXXI. DISTASIS.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis sineis, disci hermaphrod. tubulosis 5-dentatis. Recept. angustum vix alveolatum. Invol. squamæ 2-3-seriales subæquales lineares. Antheræ ecaudatæ? Achænia subcompressa hirtella. Pappus plex uterquè 4-5-paleaceus, ext. paleis ovali-oblongis brevibus, int. setis piliformibus scabris. — Herba Mexicana ramosissima pulcherrima verisim. annua. Folia caulina alterna sessilia linearia obtusissima basi attenuata. Capitula ramos terminantia solitaria laxè corymbosa. Ligulæ albidæ. Discus flavus. — Genus si verè herbarum ecaudatarum affine Neje sed diversum ligulis uni- nec pluriserialibus, pappo pauciseti et habitu: si anth. forsan caudatarum affine Pentstemon et Francoeuria. Ex hac dubitatione Distasis (græcè dubitatio) est.

1. *MODESTA*. (1) in Mexico ad Laredo legit cl. Berlandier (pl. exs. n. 1416). Capitula vix 3 lin. diam. Folia 3-4 lin. longa. Ligulæ revolutæ. (v. s. comm. à l. hav.)

## Subdiv. 3. ERIGERÆÆ.

Ligulæ pluriseriales. Pappus pilosus, nunc 1-serialis, nunc biserialis, setis externis aequali aut breviori.

## CXXXII. MELANODENDRON. — Solidaginis sp. Rozb.

Capitulum multiflorum heterogamum, fl. radii pluri- (3-5-) serialibus fœmineis angustè ligulatis, disci tubulosis hermaphrod. 5-dentatis. Recept. planum ebracteolatum tenuiter alveolare. Invol. pluriserialis brevius, squamis lanceolatis acuminatis disco subbrevioribus. Stigm.

exserta Asterearum. Anthæræ basi muticæ. Achænium compresso-tetragonum glabrum erostre. Pappus 1-serialis, setis scabridis. — Arbor insulæ Sanctæ-Helenæ maxima glabra. Folia alterna oblonga basi attenuata integerrima obtusa coriacea obscure viridia et ideò arbor anglicè dicta *Black-Cabbage-tree* undè græcè *Melanodendron* dixi. Capitula numerosa pedicellata in corymbum amplum disposita. Ligulæ albæ. — Genus *Erigeronti* valdè affine sed habitu diversissimum et achæniis classicè distinctum.

1. *M. INTRARIFOLIUM*. 5 in insulæ Sanctæ-Helenæ arenosis. *Solidago integrifolia* Roxb. in *Beats* app. 323 non Desf. Arbor insulæ maxima, trunco 5-6 ped. ambita, comâ ramosâ ampliâ, ramis lœvibus, foliis nigricantibus, floribus albis, ligulis 20-30. Achænia subtetragona glabra. (v. s. comm. à cl. Burchell.)

CXXXIII. *LEPTOCOMA* Less. in *linnæa* 1831. p. 130. syn. p. 188. *Nees* et. p. 220. — *Laveniæ* sp. Wall.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus femineis minimis ligulatis integris, disci paucis tubulosis 5?-dentatis. Recept. planum brevissimè fimbriiferum. Invol. campanulatum subbiserialè, squamis ellipticis acutis. Stylus ♀ bifidus ligulæ aequalis, ♂ brevis inclusus. Achænia plano compressa glaberrima costâ marginatâ disco parvo calloso, marginalia rostrata, centralia erostrata. Pappus 2-4-setis tenuissimis flexuosis deciduis constans. — Herba erecta rigidab vix puberula Nepalensis. Folia alterna brevissimè petiolatâ elliptico-lanceolata utrinquè acuminatâ integerrima aut hinc indè calloso-serrulata triplinervia suprà sparsè puberula subtus pallidiora glanduloso-punctatâ in nervis pubentia. Pedunculi axillares distantes 1-cephali juniores breves reflexi, adulti longiores patentes racemum foliaceum rigidum terminalem constituentes, sed capitula infer. seriùs florida! Flores albi parvi. — Genus affine *Lagenophoræ* et *Brachycomi* sed pappo diversum.

1. *L. RACEMOSA* (Less. et Nees ll. ec.). 2½ in Nepaliâ detexit cl. Wallich. *Lavenia* rigida Wall! cat. et herb. n. 325. Rhizoma lignosum subterraneum. (v. s. comm. ab hon. Cœtu merc. angl. Ind. or. et à cl. Royle.)

CXXXIV. *VITTADINIA* Rich. et Lesson fl. nov. Zeel. 250. — *Brachycome* Gaudich. ? non Cass.

Capitulum multilorum heterogamum, fl. radii 1? 2-3-serialibus ligulatis fœmineis, disci tubulosis cylindricis 5-dentatis hermaphroditis aut styli abortu masculis. Recept. planiusculum nudum. Invol. subbiserialè, squamis subequalibus linearibus margine scariosis. Anthæræ basi ecaudatæ. Achænium elongatum basi attenuatò substipitatum tertiusculum striatum. Pappus 1-serialis, setis confertissimis scabridis. — Herbæ Australasiæ diffusæ aut erectæ. Folia alterna dentato-lobatâ rariùs integrâ. Rami apice 1-cephali. Ligulæ purpurascences aut albæ. Pappus rufus.

§. 1. *Novo-Zeelandicæ, ligulis fortè 1-serialibus.*

1. *V. AUSTRALIS* (Rich. et Less. fl. nov. Zeel. p. 250), caule à basi ramoso diffuso. foliis obovato-spathulatis grossè dentatis basi attenuatis ciliatis, summis subintegerrimis, ramis floridis foliosis. 2½ in Novâ-Zeelandiâ loco dicto *Passe-de-Français* (Lesson!) et in valle Wangarua (A. Cunn.). *Brachycome* spathulatu

indich. voy. Freyc. bot. p. 467? Ligulæ 1-2-seriales. Herba variat særè glabra & puberula. (v. s.)

§. 2. Novo-Hollandicæ, *ligulis 2-3-serialibus*.

1. *TRILoba*, caule erecto ramoso puberulo-hirtallo, foliis glabrinaculis basi sessatis apice trifidis, lobis oblongis integerrimis submucronulatis, ramis floribus apice nudis. ½ in Novæ-Hollandiæ orâ orientali prope Port-Jackson. *achycome triloba* Gaudich. voy. Freyc. bot. p. 467. (v. s.)

2. *CUNEATA*, caule erecto ramosissimo vix pubescente, foliis angustè cuneatis sessatis integerrimis pubescentibus, pedunculis apice nudis. ½ in herbosis ad mœnem Lachlan in parte interiore Novæ-Cambiæ australis legit cl. A. Cunningham aprili flor. Ligulæ parvæ 2-3-seriales. Pappus pallescens. Folia 3-6 lin. longa ad axillas sæpè fasciculata. (v. s. comm. à cl. A. Cunningham.)

3. *SCABRA*, caule adscendente minutè puberulo, foliis linearibus aut infer. angustato-lanceolatis integerrimis aut infer. denticulatis confertis utrinquè pubescentibus, ramis floribus subnudis. ½ in collibus scopulosis ad Cleveland-bay Novæ-Hollandiæ tropicâ junio flor. legit cl. A. Cunningham. Ligulæ 2-3-seriales fortè albæ. Folia pollicem et ultra longa, 1-2 lin. lata. (v. s. comm. à cl. Cunningham.)

4. *DENTATA*, caule . . . , foliis linearibus integerrimis vel remotè denticulatis sess. tricuspidatis sublanatis. — cum priorè. *Brachycome dentata* Gaudich. l. c.

CXXXV. FULLARTONIA.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus fœminis angustè ligulatis, disci tubulosi 5-dentati hermaphrod. Recept. nudum. Invol. polyphyllum squamis linearibus acuminatis 1-2-seriatis sub bracteis foliaceis proximè cinctum. Antheræ ecaudatæ. Stylus in apice bifidus glaber, disci indivisus apice capitatus truncatus barbatus. Achænia erostris teretiuscula, radii glabra calva, disci ovula pappo 1-seriali scabro rufo coronata. — Herba ex Indiâ boiss. occid. orta erecta supernè pilis sparsis apice glandulosi hispida. Folia alterna semiamplexicaulia ovalia, super. oblonga grossè serrata. Capitula ad apicem caulis et ramorum axillarum solitaria. Flores ex sicco pallidè flavi seu albi. — An ex ligulis pluriserialibus achænia radii calva ad Astereas referenda? An ex stigmate apice serrato et subbarbato potiùs ad Senecioneas? — Nomen ex cl. Fullerton peregrinatore Anglo qui Indiam totam et speciatim prov. Kamaonensem peragravit et sæpè à Geographis citatur.

1. *KAMAONENSIS*. — in Kamaon prov. Indicâ ad radices Himalayæ. (v. s. sine nom. et nom. comm. ab ill. Anlâ merc. Ind. or.)

XXVI. POLYACTIDIUM. — *Polyactis* Less. syn. (1832) p. 188 non Link 1809. — *Stenactis* Cass. dict. 5o. p. 483 excl. sp. plur.

Capitulum multiflorum radiatum, ligulis fœmineis angustissimis numerosissimis circ. 2-serialibus, disco hermaphr. Recept. nudum planum minutissimè punctulatum. Invol. squamæ lineares 2-3-seriales æquales. Achænium compressum, disco epigyno minuto. Pappus simplex, ext. brevissimus coroniformis dentatus setulis subadpressis, prior paucisetosus elongatus scaber. — Herba Mexicana strictè erecta pedicula. Folia alterna, pleraque pinnatipartita, lobis linearibus densis, superiorum integris, summa linearia. Capitula corymboso-paniculata. Radius albus seu pallidè cœruleus. Discus flavus. — Differt ab

Erigeronte ut *Diplopappus* ab *Astere*. Valdè affine *Stenactidi* sed distinctum ob florum omnium pappos conformes.

1. *P. DELPHINIFOLIUM*. (2) in Mexico (Alam.! Mairer!) circa urbem (Berl. pl. exs. n. 659), in Amer. merid.? (Humb. et Bonpl.). *Erigeron delphinifolius* Willd. hort. berol. n. et t. go enum. p. 873. *Diplopappus delphinifolius* Cass. dict. 13. p. 96. *Stenactis delphinifolia* Cass. dict. 50 p. 483. *Polyactis delphinifolia* Less. l. c. (v. s. sp. et c.)

#### CXXXXVII. HETEROCHÆTA.

Capitulum multiflorum radiatum, ligulis pluriserialibus linearibus fœmineis, disco hermaphr. 5-dentato. Recept. nudum puncticulatum. Invol. squamæ 2-3-seriales acuminatæ subæquales. Achænia pubera compressa? Pappus disci et radii duplex, ext. brevis paleaceus, int. setiformis scaber longus. — Herbæ Indicæ (et fortè Americane) pilosæ villosæ aut canescentes. Capitula solitaria terminalia. Ligulæ albæ aut purpurascentes. — Genus medium inter *Stenactidem* et *Erigeropsidem*, sed differt pappo disci et radii conformi. Dist. ab *Erigeronte* pappo duplici, à *Diplopappo* et *Diplostephio* ligulis pluriserialibus. Nomen ab *στερος*, diversus, et *χαιτα*, seta.

#### §. 1. Species gerontogæ.

1. *H. DIPLOSTEPHIOIDES*, caule erecto folioso hispido tereti striato, foliis radice.... caulibus linear-lanceolatis sessilibus acuminatis integerrimis utrinquè sparsè hispidis, invol. squamis subbiserialibus dorso hispidis lanceolatis acuminatis disco longioribus, achæniis plano-compressis obovatis hispidis, pappi serie externâ paleis lanceolatis constante. ♀ in Indiæ bor. occid. montibus legit cl. Royle. Capitulum ligulis exceptis diam. pollicare. Ligulæ angustæ pollicem longæ purpureæ numerosissimæ. (v. s. comm. à cl. Royle sub n. 147.)
2. *H. HISPIDA*, caule erecto ramoso folioso setis patentibus rigidis patulis hispido, foliis lato-linearibus acutis integerrimis secus margines et nervum médium setis sparè hispidis, invol. confertè et brevius hispido, achæniis plano-compressis glabrisculis, pappi paleis ext. squamiformibus lanceolatis. — Specim. habeo ex itin. Nemesio, an ex Amer. calid.? an ex insulis Philippinis? Species distinctissima. Capitula ad apices ramorum denudatorum solitaria. (v. s.)
3. *H. ERIGEROIDES*, caespitosum totum hirtello-canescens, caulibus erectis scapiformibus 1-cephalis basi tantùm foliosis, foliis radicalibus petiolatis obovato-oblongis obtusis aut calloso-submucronatis integerrimis, caulinis paucis sessilibus oblongis, invol. hirsutissimi squamis biserialibus linearibus acuminatis disco longioribus, achænio villosa, pappi serie externâ multipaleacea brevi. ♀ in montibus Indiæ boreali-occidentalis legit cl. Royle. Habitus fortè *Erigeri* Alpini aut uniflori. Ligulæ numerosæ angustissimæ. Collum suffruticosum multiceps. Caulis 3-4 pollic. (v. s. comm. à cl. Royle sub n. 114. 1.)
4. *H. ASTEROIDES*, caule erecto 1-cephalo folioso hirsutissimo, fol. radice.... caulinis ovalibus sessilibus apice callosis integerrimis utrinquè hirsutis, infer. basi attenuatis, invol. squamis 2-3-seriatis lanceolatis linearibusque puberis disco longioribus subcoloratis, ligulis subuniseriatis linearibus obtusiusculis, achænio villosa, pappo ext. brevissimo subcoroniformi. ♀ in montibus Indiæ boreali-occidentalis. Habitus omninò *Asteris* Alpini. (v. s. comm. à cl. Royle sub n. 114. 2.)

#### §. 2. Americane, congeneres dubiæ.

5. *H. ? GNAPHALIOIDES*, caule erecto infernè simplici apice paniculato polycephalo albo-lanato, foliis sessilibus lanceolato-linearibus apice dentato-serratis subtus tomentoso-lanatis albis, radicalibus suprâ glabris, caulinis lanato-pubescentibus. ♀ in Mexici convalle Guanajuatensi et ad montem Cubilete. Eriger-

*nephalioides* H. B. et Kunth nov. gen. am. 4. p. 88. t. 331. *Stenactis* lodes Cass. dict. 50. p. 484. Ligula pappo breviores albida circ. 100, set. hæc et sequens differunt à *Stenactide*.

*rescens*, caule erecto ramoso polycephalo folisque pubescentibus, setis petiolatis oblongis lobato-dentatis, caulinis sessilibus integerrimis, longe pedunculatis. ? in temperatis juxta Moran Mexicanorum. *Eribesceus* H. B. et Kunth nov. gen. am. 4. p. 88. *Stenactis pubescens* ex. 50. p. 484. Affine *Polyactidio* ex Kunth, *Stenactidi* ex Cass. Ligula alba subtus pallide violacea.

## CXXXVIII. THEROGERON.

*Erigerontis* sed pappi dispares, ligularum pluriserialium pilosus, disci duplex, ext. paleis brevibus lanceolatis acutis inter. setis 5-7 elongatis. — Suffrutices erecti ramosi glabri in Hollandiâ interiore spontanei. Capitula ramos terminantia solitaria alba tenuia. — Nomen ex *θερος*, ætas, et *γερων*, senex, *geron* æstivus.

*truncatum*, foliis linearibus apice sublatioribus dentibus 3-5 notatis. Nova-Hollandiâ interiore in Molle's plantis prope Lechlan flumen 179. flor. legit cl. A. Cunningham. Caulis tenuis teres. Ramuli apice sola indivisa acuta paucissima gerentes. Invol. squamæ lineares acute. lati innumeri, tubulosi pauci latiores aspius steriles. Achænia radii non glabra. (v. s. comm. à cl. A. Cunningham.)

*monanthum*, foliis lanceolato-linearibus utrinque acuminatis integerrimis plantis apricis Novæ Hollandiæ interioris rarissima ad Lechlan julio 1817 flor. legit cl. A. Cunningham. Caulis teres. Folia ferè lineariter. Rami apice nudi vix foliosi. Invol. squamæ lineares acuminatæ numerosissimæ. Flores tubulosos verisim. paucissimos in capitulo uni, sed ex analogiâ cum priore summâ hæc refero. (v. s. comm. à cl. Cunningham.)

## XIX. ERIGERON Linn. gen. n. 951. (excl. spec.) Less. syn. 190.

*nudum* multiflorum radiatum, ligulis fœmineis pluriserialibus in disco æqualibus aut longioribus, fl. disci tubulosis regularibus omnibus hermaphr., nunc exterioribus fœmineis centralibus steroditis, nunc fortè abortu submasculis. Recept. nudum foveolatum (nec nisi in *E. maximo*) alveolato-fimbriiferum. Invol. compressum erostre. Pappus pilosus scaber 1-serialis (aut 2?, serie interiore brevissimâ?). — Herbariis rarius suffrutices. terna integra dentata aut lobata. Capitula subhemisphærica, invol. ligulis albis cœruleis purpurascensibusve, rarissimè ochroleis. Genus apud auctores heterogeneas species non satis notas colligens.

**HERIGERON.** — *Erigeron* Cass. dict. 15, p. 181. 37. p. 462 et 482.

disco longiores aut eo æquales. Pappi setæ in anulum basi non con-

§. 1. *Ligulis disco longioribus.*

\* *Fruticosa seu perennia.*

*fruticosum*, glabrum, caule fruticoso ramoso, foliis oblongo-lanceolatis acuminatis in petiolum attenuatis medio dentibus magnis acutis utrin-

quæ 2-3 incisæ, capitulis pedicellatis corymbosis, invol. squamis linearibus acutis imbricatis pedicellisque pubescentibus, ligulis disco duplò ferè longioribus.  $\S$  in insulâ Juan Fernandez legit cl. Larrain qui ad cl. Bertero misit. Affine *E. paucifloro* sed faciliè foliis inciso-dentatis differt. Ligulæ albæ. (v. s. comm. à cl. Bertero.)

2. *E. PAUCIFLORUM* (Hook. et Arn. bot. Beech. f. 2. p. 87), caule fruticoso ramoso glabro, ramulis pubescentibus, foliis confertis glabris lineari-lanceolatis basi attenuatis atque ciliatis integerrimis aut hinc inde serratis, junioribus subpubescentibus, paniculæ terminalis corymbosæ ramis 2-4-cephalis, ligulis lineari-oblongis disco paulò longioribus, squamis invol. oblongis acutis disco subæquantibus.  $\S$  in insulis Sandwich speciatim in ins. Onahu. Habitus *Radiarda* linearis. Folia pollicem longa sesquilineam lata. Ligulæ lineares albæ obtusiusculæ pappo subrufescente longiores. Stigmata oblonga linearia papillosa. (v. s. coll. à cl. Macrae ab hon. Soc. hortie. Londin. comm.)

3. *E. LARIDOTUM* (Less. in Linnæa 1831. p. 502), fruticosum, ramis striatis junioribus ad apicem foliatis, foliis linearibus acutis integris serratisve glabris adpressè ciliatis lepidotis, corymbis compositis pedicellis filiformibus hirsutopubescentibus, invol. campanulati squamis linearibus acuminatis.  $\S$  in Onahu ins. Sandwichensium. Achænia hirsuta punctata. Ligulæ reflexæ albæ?

4. *E. SULCATUM*, caule erecto striato-sulcato pubero-scabrido apice ramoso, foliis sessilibus auriculis 2 obtusis semiamplexicaulibus ovalibus acuminatis grossè et distanter mucronato-dentatis utrinquè setulis sparsis aspero-scabris summis lineari-lanceolatis integerrimis, capitulis ad apices ramorum solitariis subcorymbosis, invol. squamis glabris lanceolatis inter. scariosis acuminatis, ligulis disco 3-4-ve longioribus.  $\mathcal{L}$  ? in Brasiliæ paucis uliginosis legit cl. Lund. Ligulæ biseriales lineares albæ. Cor. disci flavæ. Anth. ecandata. Pappus 1-serialis. Achænia compressa glabra. Folia 6-7 pollic. longa,  $1\frac{1}{2}$  poll. lata. Habitus Inula aut *Doronici*. Car. Asteris aut *Erigeri*. Species distinctissima, socialis, putridè fetens, 8-12 pedes alta. Radix ex Lund subglobosa fibrosa (v. s. comm. à cl. inventore et ex prov. Sancti-Pauli in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 401 miss.)

$\beta$ . caule glabriusculo, foliis involucriisque vix pilosis.  $\mathcal{L}$  in Brasiliæ prov. Sancti-Pauli. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. sub n. 413.)

5. *E. GLAUCUM* (Ker. bot. reg. t. 10), caulibus adscendenti-erectis pilosis, foliis glaucis viscosis ciliato-villosis, infer. obovatis in petiolum alatum attenuatis paucidentatis, super. oblongis sessilibus, capitulo terminali subsolitario, ligulis circ. 100 linearibus disco triplò ferè longioribus.  $\mathcal{L}$  in Americâ ad Buenos-Ayres? (Ker.). Aster Bonariensis Spreng. syst. 3. p. 528. *Stenactis glauca* Nees ast. 275. Habitus *Asterum* alpigenorum. Rhizoma caudiciforme breve persistens. Herba ingrâtè odora. Ligulæ lilacinæ 2-3-seriales. Capitulum 15-18 lin. diam. Pappus simplex mihi videtur. (v. v. c.)

6. *E. AFRICOSUM*, caule erecto ramoso pilis eglandulosis sparsis distanter hispido, foliis integerrimis ciliatis glabris, radicalibus spathulatis in petiolum attenuatis, caulinis sessilibus acuminatis, capitulis ramos apice nudos terminantibus, invol. hispidociliato, ligulis numerosissimis (circ. 120) disco triplò ferè longioribus.  $\mathcal{L}$  in Californiâ. *Stenactis speciosa* Lindl. bot. reg. t. 1577. Valde affinis *E. glauco* et *E. glabello*. Capit. diam. bipollic. Ligulæ atro-violacæ.

7. *E. GLABELLUM* (Nutt. gen. am. 2. p. 148), caule erecto vix sparsè piloso, foliis integerrimis glabris eciliatis, radicalibus obovato-oblongis spathulatis, super. sessilibus lanceolatis, capitulis 3-5 pedicellatis corymbosis, invol. pubescente, ligulis numerosissimis (circ. 100) linearibus disco duplò longioribus.  $\mathcal{L}$  in planitiebus ad fl. Missouri prope Fort Mandan (Nutt.), in sylvis Amer. arcticæ (Richards. app. p. 30). Hook. bot. mag. t. 2923, sed caulibus pilis capitulis hispidulus pingitur! quod in meis non occurrit. Ligulæ pallidæ caruleo-purpureæ. Affine *E. glauco* et specioso. Capit. diam. pollicari. (v. v. c.)

8. *E. MAXIMUM* (Otto hort. berol.), caule herbaceo erecto striato setulis callosis scabro, foliis amplexicaulibus dentato-serratis suprâ exasperatis, radicalibus

elongato-lanceolatis, sup. cordato-lanceolatis acuminatis, capitulis ad apices pedunculorum solitariis, invol. squamis biserialibus acuminatis disco subæqualibus, ligulis biserialis, recept. alveolato-fimbriifero.  $\frac{1}{2}$  in Mexico. *Leptostelma maximum* Don in Sweet br. fl. gard. ser. 2. t. 38. *Aster maximus* Less. syn. 182. Rec. non est paleaceum sed fimbriiferum et ideo genus ab *Erigerone* non satis differt. Ligulæ ex albo purpurascentes circ. semipoll. longæ.

γ. *E. LONGIFRUS*, caule simplici imâ basi folioso longissimè aphylo 1-cephalo basi hispido apice pubescente, foliis subradicalibus sessilibus obovato-cuneatis grossè dentatis utrinquè sparsè hispidis, dentibus mucronatis, invol. squamis linearibus dorso villosis, ligulis numerosissimis disco duplò longioribus.  $\frac{1}{2}$  in Mexico legit cl. Karvinski. Caulis 12-13 poll. longus per 10-11 nudus. Ligulæ albae lineares. (v. s. in h. acad. reg. Monac.)

δ. *E. PHILADELPHICUM* (Linn. sp. 1211 non Willd.), pubescens, caule adscendenti-erecto laxè corymboso, foliis infer. cuneato-obovatis sinuato-dentatis, caulibus oblongo-lanceolatis semiamplexicaulibus, capitulis longè pedicellatis, ligulis numerosissimis ultra 100 tenuissimis disco duplò longioribus.  $\frac{1}{2}$  in Amer. bor. agris et pascuis à Canadâ ad Carolinam. Pursh fl. bor. am. 2. p. 533. Ell. sketch 2. p. 396, descr. optima. Er. amplexicaule Poir. suppl. 5. p. 464. Willdenowius pessimè phrasim Linnæanam correxit et pluribus dubiis ansam dedit. Herba circ. pedalis. Invol. post anthesim reflexum. Ligulæ pallidè purpureæ. (v. v. c. et s. sp.)

β. *quercinum*, foliis infer. ultra medium pinnatifidis sublyratis, lobis hinc indè dentatis.  $\frac{1}{2}$  in Carolinâ (Ell.) et Floridâ (Pursh). *Erig. quercifolium* Pursh fl. bor. am. 2. p. 533 (excl. syn. Lam.). Ell. sketch 2. p. 397. Var. *E. Philadelphici* monente opt. Elliot. (v. s.)

ι. *E. KARVINSKIANUM*, caule virgato tereti suffruticulosò puberulo, foliis canlinis inf. obovatis longè cuneatis medio triplinerviis utrinquè glabris basi ciliatis apice tri-rarius quinquedentatis, dentibus grossis obtusis mucronatis, super. oblongis subintegerrimis, capitulis pedunculatis laxè corymbosis, invol. squamis glabrisculis linearibus subulatis, ligulis disco duplò longioribus biserialibus.  $\frac{3}{4}$  in Mexico legit cl. Karvinski. (v. s. in h. acad. reg. Monac.)

α. *E. QUERCIFOLIUM* (Lam. ill. t. 681. f. 4), glabrinseculum, caulibus erectis à basi ramosis, foliis inferioribus lyrato-dentatis basi in petiolum attenuatis, superioribus sessilibus oblongo-linearibus integerrimis, capitulis pedicellatis laxè corymbosis, invol. squamis linearibus acutis glabrisculis, ligulis tennissimis circ. 100 disco duplò longioribus.  $\frac{1}{2}$  in Floridâ (Lam.), in territorio Arkansano (Nutt.) Poir. dict. 7. p. 491. Proximè accedit ad *E. Philadelphicum* sed glabritie ferè totâ jam primâ fronte differt. (v. s. comm. à cl. Nuttall.)

ζ. *E. MUCRONATUM*, caule erecto tereti glabrinseculo aut subpubero ramoso, foliis lanceolatis basi attenuatis glabrisculis ciliatis integerrimis aut utrinquè lobulo ad medium actis apice lobulise mucrone-callosò longo superatis, capitulis paucis pedunculatis, invol. squamis linearibus subulatis puberulis, ligulis biserialibus disco duplò longioribus.  $\frac{1}{2}$  in Mexico legit. cl. Karvinski. Ligulæ albæ. (v. s. in h. acad. reg. Monac.)

4. *E. BELLIDIFOLIUM* (Muhl. in Willd. sp. 3. p. 1958), hirsutum, caule florido erecto paucifolio è basi stolonifero, foliis radicalibus obovatis subhærratis, caulibus distantibus sessilibus oblongo-lanceolatis, capitulis 3-5 pedicellatis corymbosis, ligulis linearibus (circ. 50) disco duplò longioribus.  $\frac{1}{2}$  in siccis umbrosis Amer. bor. à Canadâ ad Carolinam. Pursh fl. bor. am. 2. p. 532. Ell. sketch 2. p. 393. Sims bot. mag. t. 2402. Er. palchellum Michx. fl. bor. am. 2. p. 124. Variat incanum aut minus hirsutum. Ligulæ numerosissimæ pallidè cœruleæ. Pedicelli sub capitulis obconici. (v. s. sp.)

β. *scariosum*, hirsuto-incanum, invol. squamis ferè omninò albo-scariosis vix basi ad carinam pilosiusculis.  $\frac{1}{2}$  in Kentucky legit cl. Rafinesque. Er. bellidifolium Raf. in litt. (v. s. sp.)

5. *E. PURPUREUM* (Ait. kew. ed. 1. v. 3. p. 186), caule erecto sparsè piloso,



- foliis caulinis oblongis dentatis amplexicaulibus, summis integerrimis, pedunculis corymbosis sub capitulo paulò incrassatis, invol. squamis coriâ pilosis margine membranaceis, ligulis tenuissimis numerosissimis (ultra 200) disco duplò longioribus.  $\frac{1}{2}$  in pascuis et fertilibus collibus ad ripas fluminum Amer. bor., in Canadâ ad sinum Hudsonis (Ait.), à Canadâ ad Virginiam (Pursh 2. p. 583), in Kentucky (Raf.!), Noveboraco (Lod. Bonap.!) ad ripas fl. Mississippi (Berl.!), in California (Douglas!). Erig. amplexicaule Torr. f. in litt. Ligula purpurea (Pursh) in sicco pallida imò albæ. (v. a. sp.)
16. *E. ASPERUM* (Nutt. gen. am. 2. p. 147), hirsuto-scabrum, caule gracili circ. 2-cephalo, foliis lanceolatis acutis integris, radicalibus subspathulatis, pedunculis 2-3-subterminalibus, capitulis hemisphericis. — in Amer. bor. planitiibus ad fl. Missouri. Ligula albæ. Pappus duplex, int. simpliciter pilosus circ. 20-setus achenio levi multò longior leviter rufescentes. An ex pappo dupli Stenactis aut Heterochaeta?
17. *E. INTEGRIFOLIUM* (Rigel. fl. bust. ed. 2. p. 302), caule simpliciter foliaceo glabro, foliis lanceolatis integris 3-nerviis pubescentibus, capitulis corymbosis, inv. hemisphericis squamis acutis.  $\frac{1}{2}$  circa Bostoniam in sylvis. Ligula albæ. Inter *E. bellidifolium* et *E. strigosum* collocat cl. auctor.
18. *E. PUMILUM* (Nutt. gen. am. 2. p. 147), hirsutum, caulibus à collo plurimis adgregatis 1-cephalis, foliis oblongo-linearibus integris sessilibus, capitulis hemisphericis ante anthesin nutantibus, involacro hirsutissimo. — in Amer. bor. planitiibus ad flum. Missouri. Erigeron hirsutum Pursh fl. suppl. 242, non Lour. Ligula albæ invol. duplò longiores. Capit. magna. Bellidis perennis. Pappus duplex, inter. brevis 12-setus.
19. *E. HUMILE* (Grah. in Jam. new. phil. Journ. Ed. 1828. decemb. p. 175). caule brevissimo 1-cephalo, foliis integerrimis ciliatis sursum pubescentibus patulis, radicalibus aspathulatis basi longè attenuatis, caulinis linearibus lanceolatis acuminatis, invol. campanulati calyculati squamis 6 laxis subimbricatis pilosis, ligulis invol. aequalibus disco longioribus, tubulis disci 4-dentatis.  $\frac{1}{2}$  in Americæ arcticæ littoribus (Richards). Ligula albæ.
20. *E. BERBERIANUM*, caule gracili erecto foliisque setis minutissimis adpressis puberulo apice laxè paniculato, ramis subfoliosis apice 1-3-cephalis, foliis distantibus linearibus integerrimis, invol. squamis glabriscaulis linearibus acuminatis, inter. margine membranaceis disco aequalibus, ligulis disco duplò ferè longioribus. — in fruticosis sabulosis collum prope Valparaiso Chiliensium jan. fl. legit cl. Bertero. Habitu accedit hinc ad *Asterem Gayanum* illius ad *Er. spinulosum*, à priori ligulis 2-3-seriatis, à poster. ligulis disco duplò longioribus facilè distinctum. (v. a. comm. à cl. Bertero.)
21. *E. WIGHTII*, caule erecto hispidò breviter ramoso, foliis oblongis, infer. basi attenuatis subserratis obtusiusculis, sup. integerrimis acutis, omnibus utrinquè puberulo-hirtis, capitulis pedicellatis subracemosis, invol. squamis dorso scabris lineari-subulatis disco aequalibus, ligulis tenuissimis disco longioribus, achenio glabrisculo.  $\frac{1}{2}$ ? in peninsula Indis orient. legit cl. Wight (herb. f.). Ligula pallide purpurascens multiseriales. Rami valdè hispidi. Herba cinerascens (v. a. in h. Wight.)
22. *E. AUSTERIACUM* (Lindl. 1. edn. mss.), caule basi suffruticoso ascendente ramosissimo, ramis apice aphyllis 1-cephalis, foliis linearibus integerrimis glabris, invol. squamis linearibus acuminatis disco brevioribus, inter. membranaceo-marginatis, ligulis angustè linearibus biserialibus, achenio villosulo.  $\frac{3}{4}$  in Indis or. montibus Pœrpanjal legit cl. Royle. Palmaris. Ligula cyanea. (v. a. comm. à cl. Royle sub n. 115.)
23. *E. MORTICOLA* (Wall. 1. cat. et herb. n. 2977. comp. n. 87), caule limà basi frutescente erecto simpliciter cum foliis sparè villosis, foliis radicalibus linearibus lanceolatis basi attenuatis, caulinis sessilibus linearibus, omnibus subintegerrimis acutiusculis, capitulis 2-5 pedicellatis corymbosis, invol. squamis linearibus acutis disco brevioribus, ligulis linearibus multiserialibus, fl. tubulos

paucis in centro.  $\frac{1}{2}$  in montibus Nepalie ad Kamaon et Deyra D'hoon. Er. humilis Spreng. in lit. ad Wall. Ast. monticola Lindl. adn. mas. Herba 3-6 poll. longa. Ligulæ albae. Achenia oblonga compressa puberula. Pappus rufus. (v. s. comm. ab ill. Cur. merc. angl. Ind. or. et à cl. Royle.)

24. *E. Pourretii*, totum hispidò-scabrum, caule erecto subangulato racemoso, foliis lineari-lanceolatis elongatis acutis integerrimis, infer. basi in petiolum attenuatis, super. sessilibus, ramis erectis paucifoliis apice nudis 1-cephalis, invol. squamis linearibus acutis dorso hispidò-scabris disco brevioribus, ligulis disco duplò ferè longioribus, acheniis elongatis glaberrimis.  $\frac{1}{2}$  in Chili? legit cl. Poeppig (pl. exs. n. 21). Herba ferè pedalis. Capitula circ. 5 distantia. Ligulæ ex sicco albidæ. Valdè accedit præsertim ex acheniis formâ ad *E. Andicola*. (v. s. ex coll. Poeppig.)

25. *E. Andicola*, cæspitosum, caulibus plurimis erectis molliter hispidis 1-cephalis apice subaphyllis, foliis linearibus acutis longè ciliatis et hispidis integerrimis, radicalibus erectis basi attenuatis, caulinis sessilibus paucis, invol. squamis lanceolatis-linearibus acuminatis hispidis disco aequalibus, ligulis linearibus disco duplò longioribus, acheniis elongatis glaberrimis.  $\frac{1}{2}$  in excel-sis Andibus Chilensium legit cl. Gay. Invol. purpurascens setis albis hispidum. Ligulæ ex sicco albae. Folia radic. a poll. longa, a lin. lata. Caulis 3-4 pollic. (v. s. comm. à cl. Gay.)

*ß. pubescens*, setis omnibus brevissimis, capitulo minore minùs hispidò. — in Andibus Chil. etiam legit cl. Gay. An spec. propria? An *E. Poeppigii* var. 1-cephala? (v. s.)

26. *E. glaberrimum*, glaberrimum subcaule, rhizomate suffruticuloso, foliis radicalibus lineari-oblongis basi subattenuatis, caulinis paucis semiamplexicaulis, omnibus integerrimis acutis, caule adscendente 1-cephalo, invol. squamis disco aequalibus oblongis obtusis adpressis, ligulis disco duplò longioribus.  $\frac{1}{2}$  in summis Andibus Chilensium legit cl. Gay. Etiam adest sub n. 638. in plant. Poeppigianis. Habitus Leontodontium alpinorum. Accedit ad *E. Andicola* sed distinctissimum. Caulis 2-3 pollic. Folia radic. pollicem longa, 3 lin. lata. Ligulæ albidæ ex sicco. Pappus albus. Achenia puberò-hirtella. (v. s.)

27. *E. scarorum*, caule basi decumbente surculoso-ramoso folioso, apice longè demumdato 1-cephalo puberulo, foliis pubescenti-hirtellis, infer. subcongestis oblongo-lanceolatis dentatis, summis distantibus paucis semiamplexicaulis lanceolatis integerrimis, invol. squamis linearibus acuminatis dorso hirtellis disco aequalibus, ligulis disco duplò longioribus.  $\frac{1}{2}$  in Mexico circa Toluocam spreßi flor. legit cl. Andrieux (pl. exs. n. 277). Folia infer. sesquipollic. longa, 3-4 lin. lata. Caulis floridus spithameus. Ligulæ albidæ. Discus in sicco fuscus. (v. s.)

*ß. latifolium*, foliis lato-oblongis minùs hirtellis, capitulo majore, invol. squamis glabresculis sen minùs hirtellis.  $\frac{1}{2}$  circa Mexico legit cl. Berlandier pl. exs. n. 375. (v. s.)

28. *E. pulchellum* (DC. non Michx.), caulibus basi subcæspitosis simplicibus 1-cephalis sparè pilosis apice confertè et longè hirsutis, foliis glabresculis ciliatis, infer. obovatis spatulatisve, caulinis sessilibus oblongis, invol. hirsutissimo subcolorato, ligulis tenuissimis disco duplò longioribus.  $\frac{1}{2}$  in Alpibus Iberico-Caucasicis (Stev.), Sibirie et Dahuriæ speciatim ad montem Schebet (Turczaninoff!). Aster pulchellus Willd. sp. 3. p. 2099. Aster Alwartensis Ledeb. cat. 1807. ex Sims bot. mag. t. 2321. Erig. alpinus  $\beta$  ericalyx Ledeb. fl. alt. 2. p. 91. Ligulæ purpureæ. Fl. fem. tubulosi nulli, undè ab *E. alpino* facillè differt. (v. s.)

*ß. Unalaschense*, minor, foliis radicalibus petiolatis obovatis obtusis, caulinis linearibus acuminatissimis hispidis, caule ad apicem involucrique setis longis rufidis patulo-tortuosis hirsutis, ligulis albidis.  $\frac{1}{2}$  in ins. Unalaschka et ad sinum Sancti-Laurentii legit cl. de Chamisso. Erig. Alpinus Less. in linnæa 1831. p. 126. Fl. fem. tubulosi ut in var.  $\alpha$  nulli. (v. s.)

29. *E. COMPOSITUM* (Pursh fl. bor. am. 2. p. 535), subcaule, foliis radicalibus longè petiolatis triplicato-tripartitis, lobis linearibus divaricatis petiolesque puberis, caule folia 1-2 linearia distantia gerente vix puberulo 1-cephalo. ① in Amer. bor. ad ripas fl. Kooakosky (Lewis ex Pursh), in reg. arcticis ad Point-Lake (Rich. ex Hook. in trans. Linn. soc. 14. p. 374. t. 13). Cineraria Lewisii Richards. in Frankl. journ. app. p. 148 ex Hook. Capitulum Bellidis. Ligula ex albo demùm rubellæ invol. duplò longiores.
30. *E. PAPPICHRONA* (Labill! nov. holl. 2. p. 47. t. 193), glabrum acule. foliis radicalibus obovatis subserratis basi attenuatis ciliatis, caulinis integris linearibus distantibus, caulibus scapiformibus simplicissimis 1-cephalis. ① in insuli Van-Diemen. Erigeron phlogotrichus Spreng. syst. 3, p. 520. Herba 3 pollic. Lignæ dilutè violacæ disco numerosiores circ. 2-seriales. Achenia glabra compressa. Pappus rufus. An certè hujus generis? (v. s. comm. à cl. Labill.)

\*\*\* *Herbaceæ annuæ.*

31. *E. REVULARE* (Swartz prod. 113. fl. ind. occ. 1346), caule erecto parè ramoso pilosiusculo folioso oligocephalo, foliis radicalibus caulinisve oblongo-cuneatis membranaceis ciliatis puberulis, aliis integerrimis, aliis dentatis, aliis pinnatifidis, dentibus lobisve mucronulatis, capitulis pedicellatis subcorymbosis, invol. squamis linearibus disci longitudine, ligulis linearibus disco paulò longioribus. ① in petrosis secus rivulos Jamaicæ (Sw.). Erig. runcinatum Bert! herb. Folia ferè Capsellæ Bursæ-Pastoris. Species E. Jamaicensi affinis. Ligulæ albæ parvæ. (v. s. ex h. Bert. sine loci design.)
32. *E. JAMAICENSIS* (Linn. amœn. 5. p. 406), caule erectiusculo pubescenti-subvillosa paucifolio et oligocephalo, foliis radic. rosulatis puberulis membranaceis cuneiformi-lanceolatis apice paucidentatis, caulinis distantibus squameformibus, invol. squamis linearibus glabrisculis disci longitudine, ligulis linearibus disco paulò longioribus. ① in agris Jamaicæ (Sloan. jam. 1. p. 260. t. 152. f. 3. Swartz obs. 305. t. 8. f. 2) et Sancti-Domingi (Bert!). Herba semipedalis. Folia radical. 2 poll. longa 6-7 lin. lata. Capit. parva, ligulis albis. (v. s. ex Sancto-Domingo.)
33. *E. CUNEIFOLIUM*, rhizomate obliquo, caulibus erectis oligocephalis rigidis nodis glabris vix basi puberulis, foliis radicalibus obovato-cuneatis obtusè subintegerrimis glabrisculis, caulinis distantibus squameformibus, invol. squamis linearibus disci longitudine, exter. patulis, ligulis linearibus disco paulò longioribus. ① in ins. Porto-Ricco legit cl. Wydler (pl. exa. n. 303). Affine E. Jamaicensi. Folia pollicem longa, 5 lin. lata. Caulis 7-8 pollic. Capitula longè pedicellata parva. Ligulæ in sicco purpurascences. (v. s.)
34. *E. BELLIOIDES*, caulibus debilibus adscendentibus gracillimis 1-2-cephalis glabris, foliis radic. rosulatis orbiculatis ovatisve subdentatis glabrisculis membranaceis in petiolum alatum longè cuneato-attenuatis, capitulis longe pedicellatis minimis, invol. squamis linearibus disci longitudine. ① in Porto-Ricco decemb. flor. legit cl. Bertero. Erig. revulare Spreng. in h. Bert. Herba pusilla Bellii seu Bellidis annuæ facie. E. Jamaicensi affinis sed distinctissima. Folia 9 lin. longa 3 lin. apice lata. Caules 2-3 pollic. (v. s. ex h. Bert.)
35. *E. TENELLUM*, pubescens, caule à basi ramoso gracili, ramis apice sub 1-cephalis, foliis oblongis integerrimis, infer. obtusis basi attenuatis, super. sessilibus acutis, invol. squamis linearibus puberulis disco subaequalibus, ligulis disco duplò longioribus. ① in Mexico circa Tamaulipas in campis Matamoros legit cl. Berlandier martio flor. Herba 2-4 pollic. Bellidis annuæ fere facie sed pubescens hinc indè villosa. Ligulæ albedo-cærulesc. (v. s. comm. à cl. inv.)
36. *E. LEPTORHIZON*, caule herbaceo patentim bispido apice ramoso, foliis membranaceis infernè ciliatis utrinquè puberulis, infer. basi in petiolum attenuatis obovatis seu subspathulatis apice parè dentatis, super. sessilibus oblongo-lanceolatis basi cuneatis apice grossè serratis, capitulis pedicellatis paucis subcorymbosis, invol. squamis linearibus puberulis disco aequalibus.

lis disco duplò longioribus. ① in Amer. anstr. (Demb.!) circa Limam adia!) Herba semipedalis. Folia 2 et ultrà pollices longa, 5-6 lin. lata. Inflo. albae. Pappus paucisetus. An fortè *Stenactidia* species, sed in capit. juniorum externam non ritè vidi. (v. s.)

**AFFINE**, caule erecto basi patentim hirsuto supernè pubero simplici apice nudos 1-2 nudos elongatos 1-cephalos diviso, foliis infer. elliptico-oblongis longè cuneatis integris apice dentibus grossis mucronatis utrinquè ciliis utrinquè pubero-hirtis, summis semiamplexicaulibus serratis acuminatis, invol. pubescenti-hirto, ligulis circ. 100 linearibus disco duplò longioribus. ① circa urbem Mexico junio flor. legit cl. Berlandier (pl. exs. n. 522). Philadelphico affine sed caules 1-2-cephali, capitula duplò majora circ. 2 lin. perennis. (v. s. comm. à cl. inv.)

### §. 2. *Ligulis discum non superantibus.*

**STRICTUM**, caule strictè erecto simplici sparsè setifero tereti, foliis confertis linearibus glabris secus nervum subtus hinc inde setosis margine longè plicè ciliatis aut demùm subserulatis, paniculæ spiciformis confertæ terminalis ramulis brevibus foliolosis, invol. squamis glabris linearibus, ligulis brevibus. — in Mexico circa Bejar augusto flor. legit cl. Berlandier (pl. exs. n. 50). Habitus ferè *Linarie*. Caract. *Erig. Canadensis* sed panicula contracta expansa, folia stricta magis ciliata, etc. (v. s. comm. à cl. Berlandier.)

**SPICULOSUM** (Hook. et Arn. bot. Beech. 1. p. 32), caule herbaceo erecto laxè corymboso-paniculato hispido, setis rigidis infernè sparsis patulis adicellis confertis adpressis, foliis lineari-oblongis acutis integris margine limbo setis rigidulis præsertim scabro, invol. squamis circ. discum æquantibus, ligulis minimis disco æqualibus. — in Chili (Bridg.), in pascuis sterilibus circa Valparaiso Chil. (Bert.!). *E. fasciculatus* Colla pl. chil. f. 2. n. 67. Affine videtur *E. Canadensis* et Bonariensis sed setis rigidis ferè ciliosis facillè differt. (v. s. comm. à cl. Bert.)

**CANADENSE** (Linn. sp. 1211), caule erecto hispido paniculatim ramosissimis, foliis lineari-lanceolatis ciliatis, invol. cylindricis, ligulis confertis invol. vix longioribus, achenio oblongo breviter pilosulo. ① frequens in pratis et agris Americæ borealis; etiam crescit in Mexico (Alam.!) ad Xalap. (Less.) in Brasiliâ (Ch.), Cubâ (La Sagra!), Guadalupâ (Bert.!), ins. Antis (Sieb.), Maderâ (Low.!), Indiâ orient.? (Lamb.!), Persiâ boreali v. l., Africâ austr.-orient. (Dreg.!) et Europâ totâ! Fl. dan. t. 292. *Senebaliatus* Walt. fl. car. 208? *Erig. paniculatum* Lam. fl. fr. Herba peregrina morphæ. Variet erecta aut diffusa, caule plus minùs hispido interdum glaberrimè racemiformi aut laxè, etc. An species plures hic confusæ? (v. v.)

**PAUILLUM** (Nutt. gen. am. 2. p. 148), gracile glabrum, foliis lineari-solatis, omnibus integris margine scabris, paniculæ subsimplicis ramulis ciliatis, ligulis paucis, achenio oblongo vix puberulo. ① frequens in Novâ Angliâ et Pensylvaniâ (Nutt.!) in Carolinâ? (Ell. sketch 2. p. 398). *E. Canadense* ß paucillum Bert. fl. phil. 2. p. 108. An *Erig. Canadensis* var. macilentâ? in 4-5 poll. longa. Flores disci 4-dentati ex Nutt. (v. s.)

**DECAUCATUM** (Michx. fl. bor. am. 2. p. 123), decumbens ramosissimum, ramulis fastigiatis, foliis subulatis, ligulis brevissimis, corollis disci 8. ① in pratis Illinoisensium (Michx.). In Kentucky et Louisiana (Nutt. 2. p. 140). Affine *E. Canadensis* et *E. pusillo* dicitur. An *E. ramosum* Raf. id. p. 66. satis differt?

**BONARIENSE** (Linn. sp. 1211), caule hirsuto erecto, foliis pubescenti-hirsutis elongato-lanceolatis utrinquè attenuatis, infer. grossè inciso-dentatis, r. integerrimis, paniculâ elongatâ racemiformi, ligulis tenuissimis discum superantibus. ① in Amer. austr. ad Bonariam (Dill. Bac!.), in Brasiliâ Rio-Janeiro in cultis et ad vias (Lund!). *Erig. undulatum* Moench meth.

598. — Dill. elth. 2. f. 334. Herba subpurpurascens seu obscure virescens. Folia aëpe ad basin margine reflexa aut plana. Pappus simplex. Aff. *E. Canadensi*. (v. v. c. et s. sp.)
44. *E. MULTIFLORUM* (Hook. et Arn. bot. Beech. f. 2. p. 87), glabrum, caule herbaceo, foliis lineari-lanceolatis basi in petiolum longum attenuatis integerrimis, paniculâ ramosissimâ compactâ, ramis nudis, ramulis squematis, squamis involucri oblongo-lanceolatis radium subæquantibus. — in insulis Sandwich. An fortè *E. Canadense* in ins. Owhau à cl. Less. citatum in *Linna* 1831. p. 126 ?
45. *E. HIRTELLUM*, caule suffruticoso ramoso tereti setis sparsis patulis strigillois hirtello, foliis lineari-oblongis integerrimis ad margines et subtus in nervo adpressè hirtello-ciliatis, capitulis paucis pedicellatis, invol. squamis linearibus glabriusculis, acheniis pubescentibus ligulis vix disco subæqualibus.  $\S$  in Chili circa Valparaiso legit cl. Gaudichaud.
46. *E. CHINENSIS* (Jacq. h. Schoenb. 3. p. 30. t. 303), caule erecto hispido apice paniculato, foliis villosis lyratis in petiolum attenuatis, lobis inf. parvis paucis terminali maximo ovali dentato acuto, summis lanceolatis, invol. hemispharico demum reflexo. — in Chinâ. Ligulæ albæ breves disco vix longiores.
47. *E. ? TRIDACTYLUM*, caule fruticoso brevi, ramis ferè à collo ortis erectis teretibus apice nudis 1-cephalis foliis involucriisque pubescentibus, foliis in parte ramorum infer. confertis in petiolum attenuatis apice trifidis, lobis et parte indivisâ linearibus, summis remotis linearibus acutis, invol. squamis linearibus 1-2-seriatis disco brevioribus, ligulis minimis, acheniis setigero-villosis, pappo scabrido corollis longiore.  $\S$  in Brasiliæ provinciâ Rio-Grande (h. imp. Bras. n. 1054 !). Fortè genus proprium ? Pappus rufus multisetus. Cor. ex sicco purpurascens; ligulæ paucæ. (v. s. in h. Mus. reg. Par.)

Sect. II. TRIMORPHÆA. — *Trimorphæa* Cass. dict. 37. p. 462 et 482. 55. p. 348.

Flor. fœm. innumeri, exteriores 1-3-seriales angustissimè ligulati, interiores pluriseries corollâ tubulosâ tenuissimâ truncatâ donati. Flor. centrales tubulosæ hermaphrod. 5-fidi. — *Erig.* Alpini var. aliæ plurimis aliæ paucissimis floribus fœmineis tubulosis donatæ demonstrant strictam hujus sectionis cum priorè affinitatem.

48. *E. KAMTSCHATICUM*, caule erecto glabriusculo paniculato, ramis erectis apice racemosis, foliis infimis dentatis, inferioribus oblongis obtusis basi longè in petiolum alatum et ciliatum attenuatis, superioribus sessilibus lanceolatis. invol. squamis acuminatis puberulis pappum æquantibus, ligulis disco æqualibus.  $\mathcal{L}$  in Kamtschatkâ. *E. Villarsii* Willd. sp. 3. p. 1958. excl. syn. et patr. Herba ferè bipedalis. Caulis purpurascens. Folia vix triplivenia. Capitula triplo minora quàm *E. Villarsii* veri. (v. s. comm. ab ill. Acad. sc. Petrop.)
49. *E. ACRE* (Linn. sp. 1211), piloso-hirsutiusculum, caule erecto folioso superne racemoso, pedunculis 1-cephalis, foliis oblongis integerrimis inferioribus basi angustatis, super. sessilibus, pappo achenio duplò longiore. (3) seu  $\mathcal{L}$  in scis apricis montosis Europæ totius, Caucasii et imp. Rutheniæ usque ad Dabariam. DC. fl. fr. n. 3131. *Er. vulgare* Linn. fl. lepp. 308. *Trimorphæa vulgaris* Cass. dict. 55. p. 324. Utraque varietas variat plus minus hirsuta, ligulis purpurascens corollis albidisve, pappo rufo aut albedo.

a. *brachyglossum*, radio discum non superante. *Er. acre* Sow. engl. bot. t. 1158. Schkuhr handb. t. 241. Bess. ! enum. p. 33. Ledeb. fl. alt. 3. p. 89. *E. murale* Lapeyr. suppl. 133. (ex specim. mecum ut Lapeyr. à cl. Dufour comm.). *Er. Philadelphicum* Willd. sp. 3. p. 1958 ? (v. v. et s. olim in hortis sub nom. *E. Philadelphici* cult.)

$\beta$ . *asteroides*, radio discum superante. *Er. asteroides* Andr. ! ex Bess. Lin. enum. h. berol. 2. p. 324 ? *Er. Podolicus* Bess. ! enum. p. 33. Ledeb. fl.

alt. 4. p. 190. var. *a*. Er. Dräbachiense Retz prod. n. 1010 ex fl. dan. t. 874. — Col. ocephr. 2. p. 25. t. 26. f. 2. (v. v.)

50. *E. ELONGATUM* (Ledeb. ic. alt. t. 31. fl. alt. 4. p. 91), glabrum, caule simpliciter aut paniculato, foliis integerrimis, infer. spatulatis obtusis, super. lanceolatis oblongisve acutis, pedunculis elongatis axillaribus 1-cephalis, invol. glabro, ligulis disco paulò longioribus. (2) seu 2. in siccis lapideis Sibiriae Altaicæ (Ledeb.) et Baicalensis (Turcz.). Affine *E. acris* et ab eo vix diversum mon. cl. Fischer in litt. sed glabritie distinctum. *E. elongatum* Mönch suppl. 247. est verisimè diversum sed non satis notum. (v. s. sp. et c.)

51. *E. ARMERIFOLIUM* (Turcz. in litt. 1830), caespitosum basi multicaulum humile, caulibus ramiis pubescentibus 1-cephalis, foliis omnibus linearibus integerrimis glabris basi setoso-ciliatis, invol. glabriusculi squamis linearibus apice coloratis acuminatis, ligulis disco paulò longioribus. 2. in uliginosis subsalsis prope Ircutiam in Dahuriâ legit cl. Turczaninof et in deserto Kuraico cl. de Bunge. *Erigeron podolicus*  $\beta$  pusillus Ledeb. fl. alt. 4. p. 90. (ex sp. à cl. Meyer misso). *Erig. gramineus* Stev. et Falk. non Linn. ex Ledeb. Variat caulibus foliosis brevibus et apice elongatis subnudis. Ligulae pallidae tenuissimae. Cor. fl. foem. tenuissimè truncata vix à stylo distinguendae. (v. s. comm. à cl. inv.)

52. *E. ALPINUM* (Lam. fl. fr. 2. p. 140. et DC. ed. 3. n. 313a), caulibus basi subcaespitosis pilosis mono- seu oligocephalis, foliis lanceolatis acutiusculis villosis ciliatisve integerrimis, invol. squamis apice subulatis plus minus hirtis, ligulis disco duplò longioribus, pappo paulò achænio longiore. 2. in pascuis montanis imò alpinis Europæ et Sibiriae, imò fortè Americæ arcticæ. *Stenactis alpina* Cass. dict. 50. p. 485. Species polymorpha nec tamen meo sensu divellenda.

*a.* caule 1-cephalo, invol. glabriusculo, foliis villosis, cor. tubulosis femineis numerosis. — frequens in pascuis elatis apricis. *Erig. Alpinum* Sow. engl. bot. t. 464. Hæc et sequens var. sub *Er. alpino* Linn. sp. 1211. Sturm fl. germ., etc. militat. (v. v. et s.)

*β.* caule sæpius oligocephalo, invol. glabriusculo, foliis glabris ciliatis, cor. tubulosis femineis paucis. — in pascuis pinguioribus. *Erig. glabratum* Hoppe! pl. exs. et *E. rupestre* ejusd. vix differre videtur. (v. v. et s.)

*γ.* caule 1-cephalo, invol. hirsutissimo, cor. tubulosis femineis paucis. — in pascuis editis siccoribus. *Erig. uniflorum* Linn. sp. 1212. fl. 1app. t. 9. f. 3. (v. v. et s.)

*δ.* caule oligocephalo, invol. hirsutissimo, cor. tubulosis femineis paucis. — in pratis et agris montanis. *Erig. hirsutum* et *Erig. grandiflorum* Hoppe! pl. exs. (v. v. et s.)

53. *E. VILLARSII* (Bell. app. ad fl. ped. p. 38. t. 7.), caule erecto pubescente subviscoso apice racemoso, foliis lanceolatis triplinerviis integerrimis scabridis sessilibus, ramulis 1-cephalis folio longioribus, invol. scabrido, ligulis disco duplò longioribus, pappo achænio vix longiore. 2. in collibus montanis Delphinatûs (Will.), Galloprov. super. (Honorat!), Pedemontii (Bell.), Sabaudis (J. DC.), Vallis (Schl.). DC. fl. fr. n. 3137. *Erig. atticum* Vill. dauph. 3. p. 237. Herba subviscosa pedalis polycephala rigida. Species Willd. hortensis videtur diversa! Villars junior nomen suum Villars scribebat undè Bellardi dixit *E. Villarii*. (v. v. et s.)

54. *E. SPATULATUM* (Vest St. Cr. p. 303), caule erecto simplici hispido folioso apice corymboso, foliis infer. in petiolum subulatis attenuatis ovalibus (subspatulatisve) grossè dentatis, sup. gradatim subcassilibus serratis acutis, omnibus ciliatis villosis subhirsutis, invol. squamis linearibus acuminatis diaci longitud., ligulis minutissimis. — in ins. Sancti Tomasi locis siccis aprili flor. legit cl. Bertero et fortè etiam in Porto-Ricco? St. Crucis (Vest). *Erig. incisum* var. caribæa Spreng. syst. 3. p. 516 (excl. syn. ad spec. 3. refer.). Accedit ad *E. Chinense* Jacq. sed folia infernè non lyrata. Ligulae albæ vix disco longiores. Cor. foem. tubulosas tenues, hermaphrod. latiores. Invol. demum reflexum. Ambigit inter *Nidorellam* et *Trimorphæam*. Nomen specif. minùs aptum tamen retinui. (v. s. ex h. Bert.)

55. *E. SEMIBARBATUM*, caule erecto subimplici hispido, foliis oblongis callosomucronatis, infer. basi attenuatis glabriusculis, cæteris sessilibus semimplexicaulibus ciliatis et pubescenti-hirtis dentatis integerrimis, capitalis 1-5 ad apicem caulis ramorumve solitariis subcorymbosis, invol. pedicellisque hispidissimis, corollis fœmineis ligulatis tubulosive longè barbatis, hermaphr. glabriusculis. — in Indiâ orient. legit cl. Royle. Herba vix pedalis. Folia 2 poll. longa, 4-5 lin. lata. Capit. 9-10 lin. diam. Ligulæ cœruleæ disco duplò longiores. Cor. tubulosæ fœmineæ brevissimæ, herm. 5-dentatæ glabræ. Achenia præ. radii villosissima. Pappus basi setis squamulisve cinctus; an ex achenio productus, an duplex? (v. s. comm. à cl. Royle sub n. 138.)
56. *E. ROYLEI*, caule erecto hispidulo, foliis ovali-oblongis subserratis callosomucronatis ciliatis scabridis triplinerviis, infer. basi attenuatis, super. sessilibus, capitulis multifloris pedicellatis racemosis, invol. pedicelloque hispidissimo, ligulis tubulæque fœmineis parçè barbatis, corollis hermaphr. hirsutis, acheniis puberulis. — in Indiâ orientali legit cl. Royle. Folia 2 poll. longa, 6-7 lin. lata. Ramuli erecti vix folio suo longiores. Ligulæ ex sicco albidæ. Pappus rufus cor. tubulosus æqualis. (v. s. comm. à cl. Royle sub n. 140.)
57. *E. LESCHENAUTII*, totum piloso-hispidum, caulibus herbaceis erectis teretibus striatis, foliis sessilibus oblongo-linearibus lingulatisve obtusiusculis mucronatis integerrimis, pedunculis axillaribus folio paulò longioribus 1-cephalis. summis brevioribus oligocephalis subcorymbosis, invol. squamis linearibus disco brevioribus, acheniis villosis. — in Indiâ australis montibus Neelgherry dictis legit cl. Leschenault. Villi fructus superiores calyculum circa pappum interdum simulant. (v. s. comm. à cl. Leschenault.)
58. *E. MULTICAULE* (Wall. cat. et herb. n. 3286. excl. litt. B.), glaberrimum, caulibus herbaceis à collo multiplicibus erectis teretibus, foliis lanceolatis basi attenuatis sessilibus acutis integerrimis, paniculâ terminali laxâ, ramis floridis elongatis capitula solitaria paucave gerentibus, invol. squamis linearibus acuminate, ligulis linearibus 1-serial. invol. paulò longioribus, tubulis form. pluriserialibus, masc. circ. 20. ①? in Indiâ orient. prov. Kamaon legit cl. Wallich. Herba 2-pedalis. Caulis subpurpurascens. Folia 12-15 lin. longa, 3 lin. lata. Ligulæ albæ. Pappus rufescens. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
- β. *Madurensis* (DC. in Wight contrib. bot. ind. p. 9), caule folisque pube brevi sparâ scabridis, foliis super. integerrimis, capitalis omnibus solitariis. — in provinciâ Madurensi legit cl. Wight. *Erigeron multicaule* C. Wall. cat. n. 3286. (v. s. in h. Wight.)
- γ? *Sirmorensis*, caule folisque pube brevi sparâ scabridis, foliis super. serratis. — in agro Sirmorensi legit cl. Gerard. *Er. multicaule* B. Wall. l. l. c. Fortè species propriæ? (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
59. *E. HISPIDUM* (DC. in Wight contrib. p. 9), caule basi suffrutescente erecto ramoso hispido, foliis oblongis hinc inde grossè dentatis subpuberulis, inferioribus obtusis basi attenuatis, superioribus acutis basi dilatatis cordato-amplexicaulibus, capitulis ad apices ramorum 1-2, invol. squamis linearibus seminatis, ligulis innumeris angustissimè linearibus elongatis 1-2-serialis, 2. sum. pluriserialis vix ligulatis, centralibus hermaphrod. 5-dentatis. 3. ② in Benghalo et fortè agro Maderaspetano in arenosis (Wight!). *Aster Benghalensis* Haya! herb. *Erig. hirsutum* Wall. cat. et herb. n. 2976. comp. n. 86. non Lour. nec Pursh et *Conyza graveolens* Wall. cat. n. 3106. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
60. *E. SUBLYRATUM* (Roxb. ined. ex Wight! herb. DC. in Wight contr. bot. ind. p. 9), caule ascendenti-erecto molliter hispido subimplici, foliis utriusque villosis obovatis grossè inciso-dentatis dentibus obtusis mucronatis, infer. basi attenuatis incisè sublyratis, summis cordato-amplexicaulibus, corymbis terminali 7-9-cephalo, capitulis pedicellatis, invol. squamis linearibus acuminate glabriusculis disco subæqualibus. ③ in arenosis et apricis ad Negapatam prov. Tanjore legit cl. Wight, ad Rampur in Rohilkund et in Nepaliâ Wall. cat. et herb. n. 3007. comp. 117. *Conyza hirsuta* Wall. non Linn. *Conyza laeta*

mm. herb. Ligulæ albæ Achenia glabra parva elliptica compressa. (v. s. mm. ab hon. Car. merc. angl. Ind. or.)

#### Sect. ? IV. PTERIGERON.

ligulæ pluriseriales disco longiores femineæ. Fl. centrales pauci tubulosi utati hermaphrod. aut stylo subindiviso abortientes et ferè masculi? Cor. losses, fauce non dilatata. Anther. basi ecaudatæ coadunatæ. Stylus fl. herm. mm. apice vix bifidus. Pappus scaber rigidulus. Recept. alveolatum. Habitus Conyzearum decurrentium. Sectio dubia.

L. ? DECURRENS, sparsè hirsutum, caule suffruticulosus?, foliis breviter adto-decurrentibus ovatis acutis integris aut ad basin dentatis, capitulis ad loss ramulorum solitariis subpaniculatis dispositis, invol. squamis imbricis plariserialibus seminatis, ext. ovatis dorso scabris, int. scariosis glabris, mentis sericeo-villosis. — in Novâ Hollandiâ orient. aut in Timor. Ligulæ sares pappo scabro longiores apice tridentatæ pluriseriales. Fl. herm. seu to indiviso masculi centrales pauci tubulosi fauce vix dilatata. Antheræ acutæ. Recept. alveolatum. Pappus densè barbellatus. An prioris congener? genus proprium? (v. s. comm. à Mus. reg. Par. ex itin. Bandin.)

#### Sect. v ? ELEUTHEROGERON.

ligulæ pluriseriales disco longiores femineæ. Fl. centr. pauci hermaphr., tubo, fauce dilatata campanulata 5-dentata. Anth. liberæ. Styli rami subforcipati hispiduli. Recept. punctatum. Achenia villosa, pappo barbellulato. — breviter decurrentia.

L. ANOMALUM, caule erecto glabriusculo, ramis puberulo-scabris, foliis affinis semiamplexicaulibus breviter decurrentibus obovato-oblongis basi attenuatis grossè serratis utrinquè subscabridis, summis oblongis non decurrentibus, capitulis breviter pedicellatis ad apices ramorum subcorymbosis, invol. s-serialis squamis oblongo-linearibus subequalibus, ligulis disco paulò majoribus. — in Indiâ orientali legit cl. Royle. Flor. ligulati feminei 20-25, mâ lineariter integrè vix stylo longiore. Fl. hermaphr. 5, fortè stig. abortiva hermaphr. Antheræ liberæ certè ecaudatæ. Faux cor. abruptè dilatata. Recept. mm. sensim alveolatum. Achenia fertilia compresso-plana obovato-oblonga disco pubera. (v. s. comm. à cl. Royle sub n. 144.)

#### † Species non satis notæ.

L. ? GANESCHENS (Willd. sp. 3. p. 1953), caule erecto ramoso cano-pubescente, fls lanceolatis integerrimis pubescentibus, pedunculis axillaribus 1-cephalis. Patr. ign. Cor. radiata flava et ideo à genere verisim. removenda?

L. CAUCASICUM (Stev. act. soc. h. nat. mosc. 4. p. 63), hispidum, caule procephalo, foliis integerrimis obtusè mucronatis, radicalibus oblongis petiolatis, caulinis cordato-ovatis sessilibus, capitulis longè pedunculatis, achenio pappo longitudine pappi. — in Caucasi alpe Kaischaur. Invol. valdè hirtum lute coloratum. Capit. E. acri majus.

L. ? COCHINCHINENSIS (Lour. fl. coch. 2. p. 611. in obs. Pers. ench. 2 p. 431), mâ erecto ramoso polyccephalo, foliis lanceolatis sinuatis subserratis, caulibus semiamplexicaulibus, paniculâ terminali, capitulis oblongis. — in Cochinchinâ. Erig. Philadelphicum Lour. l. c. non Linn. Flores radii tenuissimi (an alati?) feminei erecti cum disco lutei.

L. DIVERHAMPOLIUM (Cav. anal. scienc. nat. 1801. v. 4. p. 88), caule herbaceo atro, foliis inferioribus lato-lanceolatis serratis petiolatis, rameis linearibus sessilibus. — in insulâ Teneriffâ.

L. RUSTIUM (Spreng. syst. 3. p. 519), caule erecto simplici villosa, foliis sessilibus linearibus acutis pilosiusculis, infimis subserratis, pedunculis paucis



- erectis 1-cephalis, radio discum superante. — ad flum. Magnum (Rio-Grande) Brasilie. Cæt. ign.
68. *E. FASTIGIATUM* (Horn. h. hafn. add. p. 971), foliis spathulatis basi subpinnatifidis apice remote serratis, ramis paniculatis, pedunculis 1-cephalis. ① patr. et cæt. car. ign. †.
69. *E. FILIFORME* (Spreng. syst. 3. p. 520), caule herbaceo subramoso, foliis filiformibus subflexuosis longè pilosis, pedunculo solitario 1-cephalo. — ad flum. Magnum (Rio-Grande) Brasilie (Sell.). Cæt. ign. An fortè *Stenactis pilosa*?
70. *E. ? FRAGRANS* (Spreng. syst. 3. p. 518), caule ramoso folisque lanceolatis serratis incano-villosis, ramis 1-cephalis subviscosis. — patr. et cæt. car. ign. *Senecio fragrans* Lehm. ex Spreng. †.
71. *E. HIRSUTUM* (Lour. fl. coch. ed. Willd. 2. p. 611. non Pursh), caule erecto hispido, foliis lineari-lanceolatis subserratis pilosissimis, capitulis paniculatis, invol. subsquarroso. — circa Cantonem Sinarum. Herba 2-pedalis. Discus latus. Ligulæ cœruleæ lineares multæ.
72. *E. LÆVIGATUM* (Rich. in act. soc. h. nat. Par. 1790. p. 105), caule angulato debili piloso, ramis paniculatis, foliis subspathulato-linearibus integris lævigatis subpilis ciliatis. ② (ex Horn. h. hafn. 2. p. 807). Hab. in Cayennâ.
73. *E. LEHMANNI* (Spreng. syst. 3. p. 517), caule humili arachnoideo-lanato, foliis petiolatis spathulatis mucronatis integerrimis subtus subtomentosis, capitulis corymbosis adgregatis densè lanatis. ① in Ægypto ex hort. angl. Er. psillum Lehm. sem. h. hamb. 1823. p. 8. non Nutt. Er. Lehmanni Lehm. in act. nov. nat. cur. 14. p. 2. p. 809. Recept. convexum. Flor. et fr. indescr.
74. *E. LINEARIFOLIUM* (Cav. anal. scienc. nat. 1801. v. 4. p. 89), herbaceum villosum, foliis linearibus, pedunculis foliosis mono- aut polycephalis. — Circa Mogador. †.
75. *E. MONTEVIDEENSIS* (Spreng. syst. 3. p. 519), fruticosum, ramis 1-cephalis, foliis confertis lineari-filiformibus flexuosis strigillosis, pedunculis villosis, radio discum superante. — in Montevideo (Sell.). Cæt. ign.
76. *E. MYRSOTIS* (Pers. ench. 2. p. 431. n. 33), caule simplici sub-1-cephalo foliisque linearibus hirsutis. — ad fretum Magellanicum. Er. myrsotoides Jamb. herb. Caulis ferè pedalis. Flores purpurascetes. Cæt. ign. †.
77. *E. ? PILOSUM* (Pers. ench. 2. p. 431. n. 10. non Walt.), caule simplicinervulo piloso, foliis brevi-ovatis petiolatis grossè serratis pubescentibus, capitulis subpaniculatis. — patr. et cæt. car. ign. †.
78. *E. ? PINIFOLIUM* (Poir. dict. 8. p. 490), glabrum, caule adscendente perçi ramoso, foliis confertis filiformibus acutis submucronulatis integerrimis, pedunculis nudis 1-cephalis, invol. hemisphærico vix pilosiusculo, ligulis invol. longioribus. — in Bonariâ (Comm.). Flores lutei. Certè à genere videtur alienum.
79. *E. ? RESINOSUM* (Spreng. syst. 3. p. 519), villosum granis resinosis asperum, foliis sessilibus linearibus acutis paucidentatis, pedunculis paucis erectis, radio discum superante. — ad flum. Magnum (Rio-Grande) Brasilie. Ad *Asteris* genus relatum cum dubio à cl. Less. syn. 182.
80. *E. RETROFLEXUM* (Poir. suppl. 5. p. 464), caule glabro paniculato-ramoso, foliis angustissime linearibus glabris vetustate retroflexis, pedunculis erectis subpubescentibus alternis sub-1-cephalis, invol. brevi imbricato glaberrimo. ½ in Amer. bor. Flor. et fruct. indescr.
81. *E. ? ROTTLERI* (Spreng. syst. 3. p. 518), suffruticosum, foliis oblongo-lanceolatis subdentatis hispidiusculis, caulinis amplexicaulibus, ramis elongatis filiformibus 1-cephalis, invol. glabro. — in Indiâ orient. An *Conyza* sive *Blumeæ* species?
82. *E. ? SCANDENS* (Thunb. fl. jap. 313), caule scandente, foliis ovatis serratis

villosis, capitulis axillaribus. — in Japoniâ. Certissimè alius generis est, sed frustula suppetentia in herb. Thunb. nimis manca sunt. (Less. syn. 190.)

83. E.? SUMATRANÆ (Retz. obs. 5. p. 28), tomentosum, foliis sessilibus lanceolatis dentatis, paniculâ racemosâ. — in Sumatrâ. Flosc. radii dicuntur capitulares undè verisim. species ad Conyzam seu ad Blumeam referenda videtur, sed vix nota.

84. E. WARLII (Gaudich. ann. sc. nat. 1825 mai p. 104. voy. Freyc. 468), caulibus herbaceis subsimplicibus erectis folisque glabris obovato-lanceolatis acutiusculis remotè serratis infernè angustatis basi vaginatis. — in insul. Macloviania. Capit. ignota.

†† *Species exclusæ.*

- E. ÆGYPTIACUM Linn. = Conyza ægyptiaca.
- E. ALATUM Don = Blumea alata.
- E. AMBIGUUM Nutt. = Stenactis ambigua.
- E. ANNUUM Linn. = Stenactis annua.
- E. ASTEROIDES Roxb. = Blumea pubiflora.
- E. CAMPHORATUM Linn. = Blumea camphorata.
- E. CANADENSE Ten. = Conyza ambigua.
- E. CAPENSÆ Houtt. = Webbia pinifolia.
- E. CAROLINIANUM Linn. = Phalacroloma obtusifolium.
- E. CILIATUM Ledeb. = Conyza Altaica.
- E. COMTORTUM Desf. = Conyza ambigua.
- E. CRASPUM Pourr. = Conyza ambigua.
- E. DELPHINIFOLIUM H. et B. = Polyactidium delphinifolium.
- E. DIFFUSUM Pers. = Aplopappus diffusus.
- E. DUBIUM Spr. = Callistephus Wightianus.
- E. FOLIOSUM Don = Conyza flava.
- E. GLANDULOSUM Poir. = Chrysopsis graminifolia.
- E. GLUTINOSUM Linn. = Jasonia glutinosa.
- E. GLUTINOSUM Desf. = Psiadia glutinosa.
- E. GNAPHALIOIDES H. B. K. = Heterochaeta gnaphalioides.
- E. GOUANI Linn. = Conyza Gouani.
- E. GOUANI Jacq. = Conyza ægyptiaca.
- E. GRAMINEUM Linn. = Arctogeron gramineum.
- E. GRAVEOLENS Linn. = Inula graveolens.
- E. HETEROPHYLLUM Willd. = Stenactis annua.
- E. HIERACIFOLIUM Don = Blumea hieracifolia.
- E. HIRTUM Thunb. = Conyza incisa β.
- E. HYSSOPIFOLIUM Michx. = Phalacroloma obtusifolium.
- E. INCANUM Wahl = Conyza incana.
- E. INCISUM Thunb. = Conyza incisa.
- E. INULOIDES Poir. = Pulicaria inuloides.
- E. JAPONICUM Thunb. = Conyza Japonica.
- E. JUMLEI Pers. = Conyza aspera.
- E. LEUCANTHEMUM Don = Blumea leucanthema.
- E. LINIFOLIUM Willd. = Conyza ambigua.
- E. LINOSYRIS Clairv. = Linosyris vulgaris.
- E. LONGIFOLIUM Desf. = Jasonia longifolia.

- E. MOLLE* Don = *Blumea trichophora*.  
*E. HERVOSUM* Willd. = *Chrysopsis graminifolia*.  
*E. RUDICAULE* Michx. = *Stenactis verna*.  
*E. ORLIQUUM* Linn. = *Blumea amplexens*.  
*E. PINNATIFIDUM* Don = *Conyza abainthifolia*.  
*E. PINNATIFIDUM* Thunb. = *Conyza pinnatifida*.  
*E. PINNATUM* Linn. f. = *Conyza pinnatilobata*.  
*E. PUBESCENS* H. B. K. = *Heterochaeta pubescens*.  
*E. RUTILUM* Poir. = *Conyza callocephala sericea*.  
*E. SCABRUM* Thunb. = *Pulicaria erigeroides*.  
*E. SCABRUM* Pers. = *Conyza aspera*.  
*E. SCARIOSUM* Moench. = *Conyza Gonani*.  
*E. SERRATUM* Forst. = *Conyza ægyptiaca*.  
*E. SICULUM* Linn. = *Jasonia sicula*.  
*E. SQUARROSUM* Clairv. = *Inula conyza*.  
*E. STIPULATUM* Schum. = *Blumea Guineensis*.  
*E. TRICUNEATUM* Linn. = *Baccharis cuneata*.  
*E. TRISULCUM* Don = *Blumea trisulca*.  
*E. TUBEROSUM* Linn. = *Jasonia tuberosa*.  
*E. VISCOSUM* Linn. = *Inula viscosa*.  
*E. VISCOSUM* Desf. = *Psadia glutinosa*.  
*E. WIGANTIUM* Wall. = *Callistephus Wightianus*.

CXL. RHYNCHOSPERMUM *Reinw. in Blume Bijdr. p. 300.*

Capitulum multiflorum radiatum, ligulis 2-3 serialibus fuscis in disco longioribus, fl. disci tubulosi 5-dent. herm. Invol. hemisphaericum, squamis numerosis imbricatis adpressis. Recept. subconvexum acrobiculatum. Antheræ ecaudatæ. Styli rami oblongi breves. Achænium compressa glabra basi attenuata apice in rostrum brevem producta. Pappus 1-serialis setosus, setis apice tortilibus seu inflexis. — Herba perennis Javanica glabriuscula. Caulis teres, ramis ultimis verticillatis. Folia alterna lanceolata utrinque acuminata subpetiolulata remotè serrulata minutissimè scabra. Capitula ad axillas foliorum solitaria brevè pedunculata, summa nuda subterminalia. Color flor. in siccis pallidus. — Fructu rostrato accedit ad Myriactidem, sed pappo facili differt.

1. *R. VERTICILLATUM* (Reinw. et Blum. l. c.). 2. In montanis provinciis Javanicæ Tjanjor flor. aëg. Planta rarissima et distinctissima. Folia 3 poll. longa, 5 lin. lata. (v. s. comm. ab amic. Blume.)

CXLI. MICROGYNE *Less. syn. 190. non Cass. (1)*

Capitulum multiflorum..... flor. in ambitu pluriserialibus fuscis, in disco hermaphroditis regularibus. Achænium erostre. Pappus conformis pluriserialis. — Herba Brasiliensis annua ramosa canescens. Folia alterna trifurcata, laciniis linearibus. Capitula solitaria ramosa

---

(1) *MICROGYNE* Cass. = *Eriocephalis* sect. *Cryptogyne*.

supernè aphyllis terminantia. Pappus rufescens. — An fl. radii ligulati ut in *Erigereis* aut filiformes ut in *Solenogyneis*?

1. *M. TRIFURCATA* (Less. l. c.). ① in Brasiliâ (Sellow.). Cst. ign.

#### Subdiv. 4. HETEROPAPPÆ.

Radix 1-serialis. Pappus diversus in disco et radio.

#### CXLII. SIMBLOCLINE.

Capitulum radiatum, ligulis 1-serialibus fœmineis linearibus elongatis, fl. disci tubulosi 5-dent. hermaphr. aut verisim. abortu styli sterilibus. Invol. multiseriale, squamis exter. brevibus lanceolatis acutis dorso tomentosis imbricatis, inter. multò longioribus glabris discum superantibus squarrosis glabris. Recept. alveolatum, alveolis elevatis dentatis ferè fimbriiferis. Styli radii elongati apice bifidi glabri, disci inclusi brevè bifidi. Anthæræ ecaudatæ. Achænia radii oblonga subcompressa parcè pilosa et subglandulosa, disci tenuia. Pappus radii 1-serialis setosus scabridus, disci biserialis, serie utrâque setiformi exter. brevi, inter. setis radii simili. — Frutex Peruvianus. Folia oblongo-lanceolata integerrima supernè glabra atro-viridia subtus ramulis rufescenti-tomentosa. Capitula pedicellata laxè corymbosa. Flor. ex sicco vid. purpurascens? Ligulæ numerosæ circ. 30-40. — Genus differt à *Diplostephio* sect. *Amphistephio* pappo radii simplici nec cum centrali conformi, à *Tetramolopio* pappo disci duplici nec simplici et recept. profundè alveolato. — Nomen ex *σμβλος*, alveus apum, et *κλίμα*, receptaculum.

1. *S. HANKEI*. ♂ in Peruvîâ legit cl. Hanke ex herb. reg. Acad. Monac. Folia a poll. (petiolo 3 lin. incl.) ferè longa, 6 lin. lata. (v. s. in h. suprâ cit.)

CXLIII. HETEROPAPPUS Less. syn. 189. — *Asteris* sp. Thunb. — *Calimeria* sp. Nees.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus fœmineis ligulatis, disci tubulosi 5-dentati hermaphroditis. Recept. .... Invol. biserialis squamis patenti-divergentibus. Achæniæ erostre. Pappus radii coroniformis, disci pilosus 1-serialis. — Herba Japonica erecta ramosa hispida. Folia alterna, infer. oblongo-obovata crenata, super. lanceolata. Capitula terminalia solitaria parva, radio albo.

1. *H. HISPIDUS* (Less. l. c.). — in Japoniâ. *Aster hispidus* Thunb. fl. jap. 315. Banks ic. Kämpf. t. 29. non Lam. — *Keitsja* Kämpf. am. ex. 5. p. 876. *Calimeria hispida* Nees et. p. 227.

CXLIV. PHALACROLOMA Cass. dict. 39. p. 404 (excl. sp. 2). Less. syn. 188. — *Erigerontis* sp. Linn.

Capitulum pluriflorum radiatum, ligulis fœmineis 1-serialibus disco hermaphrodito fertili regulari multò longioribus. Recept. nudum planiusculum. Invol. cylindræo-campanulatum 2-3-seriale imbricatum. Achænia compressa. Pappus radii coroniformis brevis 1-serialis, disci biserialis, exterior radiali conformis, interior setis multis longis scabris. — Herba boreali-americana erecta supernè ramosa. Folia alterna remota sessilia oblonga aut linearia integerrima hispidula. Capitula laxè paniculata. Discus flavus. Ligulæ ex Dill. et Pursh flavæ, ex Cass. et Less. forsan albæ.

1. *P. OBTUSIFOLIUM* (Cass. l. c.).  $\mathcal{N}$  in demissis Amer. bor. ad sinum Hudsonis et juxta lacum Mistassin (Michx.), à Canadâ ad Carolinam (Parah), sed ab Elliot in Carolinâ non indicatur. Virga-Aurea Caroliniana Dill. cith. 2. p. 412. f. 394. *Erigeron Carolinianum* Linn. sp. 1210. *Erigeron hyssopifolium* Michx. fl. bor. am. 2. p. 123. An stirps Cassiniana certè eadem ac Dilleniana sive Linnaeana? Dillenius flores ligulatos pappo eodem ac tubulosos donatos exhibet!

† *Species exclusa.*

*P. ACUTIFOLIUM* Cass. = *Stenactis annua*.

CXLV. MINURIA.

Capitulum multiflorum, fl. radii 1-serialibus foemineis ligulatis, disci tubulosi 5-dentatis styli abortu masculis. Invol. pauciseriale imbricatum, squamis linearibus acuminatis margine scariosis. Recept. planum nudum. Achænia disci subtruncata sericeo-villosissima, pappo multisetiformi scabro, villis achæniæ summis pappum duplicem simulantibus; disci gracilia glabra inania pappo 1-seriali, paleis aliis brevibus inter se concretis aliis paucis setiformibus longis scabris. — Suffrutices Novæ-Hollandiæ interioris indigenæ erecti glabri ramosissimi. Folia alterna acerosa seu linearia acuta integerrima. Rami apice summo plus minus nudi 1-cephali. Ligulæ albæ. — Genus pappo heterogeneo distinctissimum, habitu *Erigeropsidis*, caractere ferè *Stenactidis*. Nomen ex *μινυρος*, exilis.

1. *M. LEPTOPHYLLA*, ramis usque ad capitulum folia parca gerentibus, ligulis disco triplo longioribus.  $\mathcal{N}$  ad apices collium sterilibus Novæ-Hollandiæ interioris ad flumen Lachlan aprili flor. legit cl. Cunningham. Ligulæ circ. 20 lineares. (v. s. comm. à cl. A. Cunningham.)
2. *M. TENUISSIMA*, ramis apice longiusculè nudis, ligulis disco duplò longioribus.  $\mathcal{N}$  in collibus scopulosis sterilibus Novæ-Hollandiæ interioris ad flum. Lachlan aprili flor. legit cl. Cunningham et mecum comm. Ligulæ pauciores et paulò longiores ex sicco videntur. Capitula dimidiò ferè minora. (v. s.)

CXLVI. *STENACTIS* Nees ast. p. 273. Less. syn. 189. — *Diplopappi* et postea *Stenactidis* et *Phalacroloomatis* sp. Cass. dict. 37. p. 462 et 485 et 39. p. 405. — *Asteris* sp. Linn. — *Erigerontis* sp. Ait. — *Policaris* sp. Gærtn.

Capitulum multiflorum heterogamum, fl. radii uniserialibus foemineis angustè ligulatis, disci hermaphr. 5-dentatis. Recept. nudum punctulatum planum seu convexum. Invol. squamæ 2-3-seriales imbricatæ angustæ. Achæniium oblongum compressum. Pappus radii 1-serialis setaceus deciduus, disci duplex, exter. brevissimus, interior pappo radii similis. — Herbx erectæ ramosæ. Folia alterna dentata. Capitula ad apices ramorum solitaria disco luteo, radio albo aut pallidè violaceo. — Nomen ex *στενός*, angustus, et *ακτίς*, radius.

\* *Americana.*

1. *S. ANNUA* (Nees ast. p. 273), caule erecto apice polyccephalo corymboso, foliis infer. ovatis, super. lanceolatis dentato-serratis pilosiusculis, invol. setose. ① in America bor. secus vias et in ruderalis frequens, undè migravit in Europam. Habeo lectam circa Bassano à fratre meo, inter Petaviam et Vicentiam à cl. Duby, indic. in Belgio à Dekin et Passy fl. brux. p. 55. *Diplopappus dubius* Cass. bull. philom. 1817. p. 137. 1818. p. 77. *Stenactis dubia* Cass. dict. 37. p. 485. *Phalacroloma acutifolium* Cass. dict. 39. p. 405. *Policaria annua* Gærtn. fr. 2. p. 462. *Cineraria corymbosa* Moench. *Erigeron annuum* Pers. ench. 2

. 431. *Erigeron heterophyllum* Willd. sp. 3. p. 456. Nutt. l. gen. 147. *Aster nanus* Linn. sp. 1229. fl. dan. t. 485. — Corn. can. 193. t. 194. Herba valde modè habitum et formas diversas foliorum varians. Ligulæ albæ. (v. v. et s.)

3. *AMBIGUA*, vix pubescenti-scabrida, caule erecto apice ramoso subcorymoso, foliis elongatis oblongo-linearibus, infer. basi in petiolum longè attenuatis subserratis, super. linearibus integerrimis, invol. squamis puberulo-abris. — in Georgia (Nutt. l. gen.) et in territorio Arkansano (Nutt. l. in litt.). *Erigeron ambiguus* Nutt. gen. am. 2. p. 147. Ell. sketch 2. p. 395? Ligulæ numerosæ 1-seriales. Pappus radii brevis, disci duplex, ext. paleaceus brevis. v. a. comm. à cl. Nutt.)

. *PILOSA*, caule erecto apice corymboso-ramoso tereti pilis patulis subhispido, liliis linearibus acutis integerrimis pilis longis et pube breviori hirtellis, invol. squamis acuminatis margine membranaceis dorso puberis, acheniis villosis. ) in Brasiliæ prov. Rio-Grande. Ligulæ albide tenues. Pappus exter. disci areæ conspicuus. Herba 6-10 pollic. (v. s. in h. Mus. reg. Par. à Mus. imp. res. sub n. 1046 miss.)

4. *VERNA* (Nees ast. 275), caule subaphyllo glabriusculo composito-subcorymoso, foliis radicalibus rosulatis oblongis acutis glabris subdentatis, caulinis instantibus linearibus integerrimis, invol. vix pubescente. 2/ in depressis humidis Carolinæ et Georgiæ. *Aster vernus* Linn. sp. 1230. *Erigeron nudicaule* Michx. fl. bor. am. 2. p. 134. Ell. sketch 2. p. 332. *Aster vernus* et *Erigeron nudicaule* Pers. ench. *Doronicum lævifolium* Walt. fl. car. p. 205? Ligulæ lineares 1-seriales albæ ex Ell., purpureæ in spec. meo Georgico, non invol. æquales dicunt cl. Nees, Michx., etc., sed duplò longiores ut dixit cl. Elliot. An ideò agitudine varia? An duæ species confusæ? Pappus duplex, in radio depauperatus. (v. s.)

5. *STRIGOSA*, caule erecto sparso et patentem piloso, foliis lanceolatis utrinque attenuatis dentatis strigoso-hispidis, capitulis laxè corymboso-paniculis, invol. glaberrimo, ligulis numerosis filiformibus. (2) ? in Pennsylvania (Muhl.), Noveboraco (Torrey!). *Erigeron strigosus* Muhl. in Willd. sp. 3. 1256. *Erig. australe* Horn. ex Spreng. An *Erig. strigosus* Ell. sketch 2. 304 et *Doronicum ramosum* Walt. fl. car. 205 in Carolinâ cresc. ad hanc sectionem referenda? (v. s. comm. à cl. Torr.)

#### \*\* Asiaticæ.

6. *ELLIPSOIDES* (Nees ast. p. 274), caule adscendente angulato-striato oligophallo pilosiusculo, foliis lanceolatis acutis apice subserrulatis pubescenti-abridis, infer. hinc indè grossè serratis, summis sublinearibus integerrimis, vol. setoso. 2/ in Nepalæ superioris montanis apricis et in Kamaon. *Aster ellipoides* Don prod. fl. nep. p. 177. Wall. l. pl. asiat. rar. 2. p. 1. t. 101. cat. 2978. comp. 88. Ligulæ lilacinæ. (v. s. comm. ab hon. Cæst. merc. angl. d. or.)

. *MULTIRADIATA* (Lindl. in litt. 1835), caulibus à basi ramosis erectis infer. abris sup. villosis sulcatis, foliis sessilibus ovalibus basi subcuneatis et interrimis apice sparso et grossè serratis mucrone calloso terminatis utrinque iberulis, ramis 1-cephalis sub capitulis brevi spatio nudis, invol. squamis serratis acuminatis hirtellis disco æqualibus, ligulis linearibus numerosissimis, acheniis puberulis. 2/ in India montibus boreali-occid. leger. cl. Wallich Royle. *Aster multiradiatus* Wall. l. cat. et herb. n. 2969. comp. n. 79. *Erigeron illoides* Spreng. in litt. ad Wall. *Aster involoides* Don prod. fl. nep. 178? Nees t. 36? sed ex Lindl. adn. descr. Neesii fortè è pluribus speciebus petita? (v. a.)

#### † Species exclusæ.

. *PIXA* Cass. = *Erigeron alpinum*.

. *ALPINITOXA* Cass. = *Polyactidium delphinifolium*.

. *AUCA* Nees = *Erigeron glaucum*.

5. *GNAPHALIOIDES* Cass. = *Heterochæta gnaphalioides*.

5. *PUBESCENS* Cass. = *Heterochæta pubescens*.

5. *SPICIOSA* Lindl. = *Erigeron speciosum*.

CXLVII. *GYMNOSTEPHIUM* Less. *syn.* 185. *Nees ast.* 253.

Capitulum multiflorum heterogamum, fl. radii ligulatis femineis uniserialibus, disci tubulosis 5-dentatis masculis sterilibus. Recept. nudum areolatum. Invol. squamæ imbricatæ subtriseriales. Achænia radii plano-compressa erostria subpilosa, disci linearia incana. Pappus disci obsoletus paucisetosus, setis plumoso-serratis, radii nullus. — Plantæ herbacæ Capenses ramosæ. Folia alterna linearia integerrima. Capitula pedunculata solitaria parva, radio cœruleo, disco luteo. Involucra sæpius glandulis oblongis ut in *Tagete* notata. — Affine dicitur *Chariæ*, sed differt disco masculino. Affinis fortè *Felicis* et *Mairis*, sed differt achæniis radii epapposis, disci paucisetis. — Genus fortè nimis artificiale?

§. 1. *Glabra, invol. glandulis oblongis notatis.*

1. *G. GRACILE* (Less. l. c.), herbaceum? glabrum, ramis floridis superne aphyllis, invol. squamis ellipticis glandulis oblongis notatis. 2? ad Cap. Bonæ-Spei in Thonow. Facies *Mairis* microcephalæ.

2. *G. FRUTICOSUM*, glabrum fruticosum, ramis floridis apice villosis fortè ad apicem folia seu squamas elongatas gerentibus, invol. squamis linearibus-oblongis ext. acutis, inter. obtusis et glandulas oblongas gerentibus. 5 in Africa Capensi ad Zwillingendam alt. 200-300 ped. legit cl. Dreg. Fruticula circ. pedalis. Folia 9-12 lin. longa, 1-1 ½ lin. lata margine subrevoluta, uniuersa. (v. s.)

§. 2. *Hirsuta, involucris eglandulosis.*

3. *G. HIRSUM* (Less. l. c.), herbaceum? hirtum, invol. squamis linearibus cæminatis. 2? ad Cap. Bonæ-Spei in Steenboks rivier ad Swarteberga. Capitula magna. Bellidis perennis.

CXLVIII. *CHARIÆ* Cass. *bull. philom.* 1817. p. 68 et 1821. p. 22. *Dis. l.* p. 191. 24. p. 369. 37. p. 463 et 489. *Less. syn.* 389. — *Kaulfussia* *Nees herb. ber.* p. 53. t. 11. *ast. p.* 268 non *Blume*.

Capitulum multiflorum, heterogamum, fl. radii ligulatis femineis, disci hermaphr. tubulosis 5-dentatis. Recept. alveolatum, alveolorum marginibus subdentatis. Invol. 2-seriale campanulatum, squamis ext. planis, interioribus complicato-carinatis. Stylus disci ramis apice quasi trapezoides depressis et ibi dorso subpuberulis. Achænia obovato-cuneiformia compressa nervo calloso cincta cæterum subnerviis hirtella, radii sæpè inania semper epapposa aut 1-setosa. Pappus disci setis capillaribus plumosis 1-serialibus. — Herba monocarpica Capensis. Folia infima opposita, cætera alterna, oblonga lanceolatave basi trinerviis cum ramis subhirtellis. Pedunculi terminales aphylli. Capitula nascentia subflava demum cœrulea, disco et radio sæpè discoloribus aut concoloribus. Potius ex habitu et stigmate ad *Astereas* quàm ad *Seneciones* meo sensu pertinere videtur.

1. *C. HETEROPHYLLA* (Cass. ll. cc.). ① ② ad Cap. Bonæ-Spei (Nees), ad Zilverfontain (Dreg.). Char. cœrulea *Nees ast. p.* 269. cum var. 3 nempte:

- α. caule erecto oligocephalo, disco lateo ligulis angustioribus. — *C. heterophylla* Cass.  
 β. caule erecto oligocephalo disco radioque cœruleis, ligulis latioribus. —  
*C. cœrulea* Cass. dict. 24. p. 370.  
 γ. caule ramosiore sæpè diffuso, disco radioque cœruleis, ligulis latioribus.  
*C. Neesii* Cass. dict. 24. p. 372. *Kaulfussia amelloides* Nees hor. ber. 53.  
 t. 11. (v. v. c. et s.)

CXLIX. CHÆTOPAPPA. — *Chætophora* Nutt. in litt. 1825 non Schrank nec Brid., nec Hornsch. (1).

Capitulum multiflorum radiatum, fl. radii 1-serialibus fœmineis ligulatis, disci hermaphrod. tubulosi 5-dentatis. Recept. angustum nudum. Invol. squamæ laxè imbricatæ apice scariosæ acuminatæ. Achænia teretiuscula. Pappus nunc duplex nunc subduplex, ext. latè et hyalinè paleaceo, in fl. disci brevi in fl. radii longiore, inter. setis circ. 5 rigidis scabris constante. — Herba pusilla annua Boreali-Americana. Tota pubescenti scabra, sæpius multicaulis ramosa, rarius in siccis simplex. Folia caulina alterna linearia aut lineari-spathulata integerrima, radicalia obovato-spathulata in petiolum attenuata. Rami apice nudi 1-cephali. Invol. glabrum nitens. Ligulæ albæ demùm subpurpurascentes. Discus flavus. — *Boltoniæ* paulò affinis videtur.

1. *C. ASTEROIDES* (Nutt. l. in litt. sub *Chætophora*). ① in territorio Arkansano. Herba 2-5 poll. longa. (v. s. comm. à cl. invent.)

CL. BOLTONIA *L'Hér. sert. angl. p. 27. Cass. dict. 37. p. 464 et 491. Nees est. 234.* — *Madea Soland. ined. ex L'Hér. mss.* — *Matricaria* sp. *Linn.* — *Chrysanthemi* sp. *Walt.*

Capitulum radiatum, ligulis 1-serialibus linearibus fœmineis fertilibus, disco hermaphr. fertili. Receptaculum hemisphæricum alveolatum. Involucri squamæ discum sequantes biseriales adpressæ margine membranaceæ. Achænia plano-compressa calloso-marginata glabriuscula seu hispidula. Pappus brevissimus setulosus nunc subæqualis, nunc in floribus disci setis duabus elongatis subulatis crassioribus constans. — Herba rhizocarpica Boreali-Americana erecta ramosa Asteris facie. Folia alterna subverticalia lanceolata glaucescentia lævia margine scabra integerrima aut inferiora dentata. Capitula laxè corymbosa. Discus luteus. Radius albus aut pallidè purpurascens.

1. *B. GLASTIFOLIA* (*L'Hér. l. c.*), foliis lanceolatis glaucescentibus, inferioribus serratis, capitulis brevibus pedicellatis, achæniis obovatis, pappi disci setis plurimis inæqualibus, 2 oppositis crassis elongatis rigidulis. ¼ in Amer. bor. à Pennsylvania ad Carolinam ad ripas rivulorum et paludum. Willd. sp. 3. p. 2162. Nees est. 235. bot. mag. t. 2381. Ell. sketch 2. p. 399 (excl. signo?). *Madea glauca* Soland. in hort. Lee ex *L'Hér.* l. *Matricaria glastifolia* Hill. h. kew. 19. t. 3? (v. v. c.)  
 2. *B. DIFFUSA* (Ell. sketch 2. p. 399), glaberrima, foliis lineari-lanceolatis integerrimis margine scabris, paniculæ diffusæ pedunculis elongatis, achæniis obovatis, pappi disci setis plurimis inæqualibus 2 oppositis crassis elongatis rigidulis. ¼ in Amer. bor. pinguibus humidis inter Chatahouchie et Alabama. *Anonymos asteroides* Michx. l. herb. *Boltonia asteroides* bot. mag. t. 2554 excl. syn. Capitula minora quàm *B. glastifoliæ*, cui accedit ex pappo disci, sed sæpè

(1) *Chætophora* Schrank est genus à plerisque inter Algas admissum. *Chætophora* Brid. est *Leskeæ* sp. *Chætophora* Hornsch est *Hookeriæ* sp.



ob folia integerrima et capitula longè pedicellata, in hortis cum *B. asteroides* confusa. (v. s.)

3. *B. asteroides* (L'Hér. l. c.), foliis integerrimis lineari-lanceolatis, capitulis longè pedicellatis laxè subcorymbosis, achæniis ovalibus glabris, pappo brevissimo deciduo in radio et disco subæquali.  $\frac{1}{2}$  in paludosis et ad lacuum ripas Carolinæ et terræ Illinoensium? Willd. sp. 3. p. 2162. Nees ast. 236. *Matricaria asteroides* Linn. mant. 116. *Chrysanthemum Carolinianum* Walt. fl. car. 204. (v. v. c.)

CLL. SOMMERFELTIA Less. gen. 180 non Schum. (1)

Capitulum radiatum, ligulis fœmineis 1-serialibus, disco sterili regulari. Recept. epaleaceum. Invol. imbricatum. Achænium erostre. Pappus pilosus, radii pluri-, disci 1-serialis. — Fruticulus Brasiliensis. Folia conferta pinnatisecta rigida glabra, segmentis filiformi-subulatis spinescentibus. Capitula solitaria ramos supernè aphyllous terminantia.

1. *S. spinulosa* (Less. l. c.).  $\frac{3}{4}$  in Brasiliâ. *Conyza spinulosa* Spreng. syst. 3. p. 510. Folia cinerea. Pedunculi adgregati (Spr.).

Subdiv. 5. BELLIÆ.

Radius 1-serialis. Pappus squamellatus.

CLII. CALOTIS Brown in bot. reg. n. 504.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus fœmineis, disci tubulosi 5-dentatis masculis. Recept. epaleaceum, convexum scrobiculatum. Invol. floribus disci subæquale, squamis 1-2-seriatis distinctis. Anthæræ basi muticæ. Stigmata 2 gracilia divergentia. Achænium verticaliter compressum cuneatum coronatum paleis 2 auriculæformibus aristisque 1-3-glochidiatis. — Herbæ Novo-Hollandicæ ramos pilosiusculæ. Folia alterna sessilia inciso-dentata. Capitula ramos apice aphyllous terminantia solitaria. Discus flavus. Radius cœruleus.

1. *C. CUNEIFOLIA* (Brown in bot. reg. t. 504), glabriuscula, foliis cuneatis apice inciso-dentatis, invol. biseriali circ. 15-16-phylo, aristis per totam ferè longitudinem retrorsum serratis. — in Novâ-Hollandiâ interiore ad ripas fluminis Lachlan Lodd. bot. cab. t. 825.
2. *C. DENTEX* (Brown in bot. reg. t. 504 in obs.), glabriuscula, foliis oblongo-lanceolatis utrinquè incisis 2-3 profundis acutis notatis, invol. subbiseriali 8-9-phylo, aristis apice summo glochidiatis. — in Novâ-Hollandiâ circa Port-Jackson. (v. s. ex hort. kew.)
3. *C. DILATATA* (Cunn. l. in litt. 1834), tota pubescens, foliis latè subrhomboidis grossè dentatis basi cuneatis in petiolum basi obtusè auriculatum attenuatis, involucri biseriali, aristis lævibus.  $\frac{1}{2}$  in collibus lapideis Novæ-Hollandiæ interioris ad Peel's rage, Exley, etc. junio 1817 florent. legit cl. A. Cunningham. Folia pollicem lata. (v. s. comm. à cl. invent.)

CLIII. ASTEROMEA Blum. bijdr. p. 901. Nees ast. 232. Less. syn. 191. — *Asteris* sp. Linn.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci hermaphr. tubulosi 5-dentatis. Recept. convexum alveolatum. Invol. biserialis, squamis subæqualibus margine membra-

(1) SOMMERFELTIA Schum. = *Drepanocarpus*. Conf. Prod. 2. p. 420.

is apice herbaceis. Stigm. disci apice lanceolata. Achænium 4-cos-  
m compressiusculum basi attenuatum subpuberulo-glandulosum.  
pus paleaceus 1-serialis, paleis brevibus basi subcoactis apice  
300-multifidis. — Herba Asiatica erecta glabriuscula. Folia alterna  
i-oblonga utrinquè acuminata sessilia integra aut hinc inde serrata  
ca obovata. Capitula ad apices ramorum solitaria. Ligulæ cœruleæ  
albæ, discus luteus. — Habitus Asteris.

1. *INDICA* (Blum! l. c.). — in Indiâ or.? (Linn.), in imp. Chinensi? (Nees),  
ita in hortis Indicis. Aster indicus Linn. sp. 1230. Burm. fl. ind. 181. Willd.  
. 3. p. 2041 excl. descr. et car. ex Nees ast. 233. Aster Javanensis Garc.  
rb. ex Burm. Callistemma indicum G. Don in Loud. hort. brit. t. 348.  
. a.)

2. *BELLIUM* Linn. mant. p. 157. DC. fl. fr. 4. p. 923. Cass. dict. 4. suppl.  
47. et 37. p. 454. Less. syn. 191. — Bellidis sp. Tourn. — Pectidis sp.  
kreb.

spitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis  
lineis, disci hermaphr. tubulosis 4- (rarius-5) dentatis. Rec. ovoï-  
conicum nudum. Invol. 1-2-seriale, squamis oblongis. Anth. inclusæ.  
m. subforficata. Achænium compressum obovale subpuberulum.  
pus squamis 4-5 membranaceis truncatis et setis totidem cum  
amis alternantibus constans. — Herbæ in Mediterraneis regionibus  
igenæ annuæ aut perennes. Folia subradicalia obovata integra aut  
us dentata. Pedunculi scaposi 1-cephali. Capit. discus luteus, radius  
15. — Habitus Bellidis.

1. *BELLIDIODES* (Linn. mant. 285. excl. syn. Triumf.), caule stolonifero,  
lis radicalibus spathulatis integris basi longè attenuatis, scapis folia ferè  
lplò superantibus. ① in rupestribus humidis maritimis Corsicæ!, Sardinie  
Moris), ins. Balearicarum (Camb. bal. 104), Therasiæ (D'Urv.), Chalci et Rhodi  
libth.) Viv. ! fragm. f. ital. 1. p. 8. t. 10. fl. 1. Lam. ill. t. 684. DC. fl. fr. 4.  
. 923. non Desf. Bellis droseræfolia Gou. ill. 69. — Bocc. mus. t. 107. (v. v.  
. a.)

2. *STIVALE* (Requien! in ann. sc. nat. 5. p. 383), caule non stolonifero, foliis  
adicalibus spathulatis integris breviter attenuatis, scapo apice villosa, foliis  
sulò longioribus aut iis subæqualibus. ①? in editioribus montibus Corsicæ  
gerunt cl. Requien et Soleirol, et in Sardinia cl. Badaro. Herbula pollicaris  
smilis. An ex cl. Duby (bot. gall. 1. p. 266) à priori satis differt? (v. s.)

3. *CRASSIFOLIUM* (Moris! pl. sard. elench. 1. p. 26), caule non stolonifero  
nai multicipiti subadscendente, foliis subradicalibus crassis obovatis integerrimis  
basi longè attenuatis subpubescentibus, scapis folia triplò quadruplove  
perantibus pubescentibus. 2/ in rupestribus maritimis Sardinie ad S. Eliam  
r Sancti-Petri ad Cala-Vinagra legit cl. Moris. Species distinctissima habitu  
ellidis perennis. (v. s.)

4. *ROTUNDFOLIUM*, caule non stolonifero, foliis radicalibus obovato-subro-  
radiis crenato-dentatis in petiolum longum desinentibus pubescentibus, scapo  
lis triplò quadruplove longiore pubescente. 2/ in Atlante. Doronicum rotun-  
dum Desf. fl. atl. 2. p. 279. t. 235. f. 1. Arnica rotundifolia Willd. sp. 3.  
. 2110. Bellium giganteum Cass. dict. 37. p. 456. Habitus ferè Bellidiastri sed  
appus paleis 5 setisque totidem alternantibus ut in Bellio.

5. *MINUTUM* (Linn. mant. 286), caule folioso capillari adscendente lævi, fo-  
lis obovatis integerrimis petiolatis. ① ad Orientis fontes. Pectis minuta Linn.  
p. 2. p. 1250. Schreb. nov. act. ups. 1. p. 81. t. 5. f. 2. Bellis cretica fontana  
minium minima Tourn. cor. 37.

† *Species exclusæ.*B. ANNUUM Less. = *Bellis annua*?B. DENTATUM Viv. = *Bellis dentata*.

## Subdiv. 6. BELLIDÆ.

Radius 1- aut rarissimè 2-serialis. Pappus nullus aut coroniformis.

CLV. BELLIS. Linn. gen. 962. Gaertn. fr. 2. t. 168. Cass. dict. 37. p. 493. Less. syn. 192. — *Bellis* et *Kyberia* Neck. elem. n. 80 et 81. — *Bellidis* sp. Tournef. Willd.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci tubulosis hermaphrod. 4-5-dentatis. Recept. conicum. Invol. campanulatum 1-2-seriale, squamis obtusis foliaceis æqualibus. Achænia compressa obovata hispidula omnino epapposa. Herbe Europæe caules caulescentesve. Folia radicalia aut alterna oblonga aut obovata sæpius dentata in petiolum attenuata. Pedunculi axillares aut scapiformes 1-cephali. Discus luteus. Ligulæ albæ extus subroseæ aut subviolaceæ ante explicationem capituli in globum conniventes.

\* *Species caulescentes verisim. annuæ.* — *Kyberia* Neck. elem. n. 81.

1. B. DENTATA, caule sæpius basi ramoso diffuso hispidulo, foliis obovatis grossi dentatis basi cuneatis, invol. squamis vix acutis sæpè pilosis, corollis omnibus ad basin tubi pilorum articulorum fasciculo barbatis. (1)? in humidis Italis ad Pisas (Savi!), Veronam (Reich.), Neapolim (Guss!), Siciliam (Guss!), Melitæ (Beaupré!), Milli (Beaupré!), Leucadam (Parol!), Constantinopolem (Castagne!), Barbariam (Desf.). *Bellium dentatum* Viv. fragm. p. 8. t. 10. f. 2. (excl. syn. plerisq. et descr. fract.). *Bellium bellidioides* Desf. fl. atl. 1. p. 279. non Linn. — *Bellis minima annua* Triumf. obs. p. 80. ic. Similissima sequenti sed corollâ imâ basi barbigerâ undè incautè ad *Bellium* hinc inde relata. (v. s.)

2. B. ANNUA (Linn. sp. 1249), caule sæpè à basi ramoso diffuso hispidulo, foliis obovatis integris subdentatis basi in petiolum attenuatis, invol. squamis vix acutis sæpè pilosis, corollis basi nudis. (1)? in pratis humidis Occidentis, Galloprov., ins. Balearicarum (Camb.!), Melitæ (D'Urv.!), Corsicæ (Thom.), Teneriffæ (Brouss.), Barbariæ (Desf.), Orientis, Italiæ, Hispaniæ (Willd.). DC. fl. fr. n. 3220. — Bocc. mus. 2. t. 35. B. ramosa et B. repens Lam. & fr. 2. p. 122. *Bellium annuum* Less. syn. 192? Habitus varius: acaulis, erectis aut sæpius diffusa. (v. v.)

β *minuta*, caule basi multicipiti, scapis erectis folio quadruplò longioribus. — circa Byzantium locis humidis legit cl. L. Castagne et in Græciâ (h. anal. Monac!). Habitus omnino *Bellii bellidioidis* sed achænia calva. (v. s.)

3. B. INTEGRIFOLIA (Michx. fl. bor. am. 2. p. 131. non Lam.), caulescens divaricato-ramosa, foliis integerrimis longè ciliatis inferioribus obovatis, super. lanceolatis, invol. squamis glabriusculis levibus acuminatissimis et quasi in aristam desinentibus. (1) ad ripas rivulorum et in collibus umbrosis Tennessee (Michx.), in Kentucky occidentali (Raf.!) in agro Arkansasano (Nutt.!). Radix exilis. *Eclipta integrifolia* Spreng. syst. 3. p. 602. ex syn. (v. s.)

\*\* *Species subacaules perennes.* — *Bellis* Neck. elem. n. 80.

4. B. PERENNIS (Linn. sp. 1248), foliis subradicalibus spatulato-obovatis crenato-dentatis pubescentibus venosis invol. squamis pilosis vix acutis. ? in Europæ pascuis vulgatissima usquè ad Tauriam (Bieb.), etiam in Madagascariâ dicitur introducta (Low.!) Lam. ill. t. 677. fl. Dan. t. 502. engl. bot. t. 424.

Schkuhr handb. t. 251. Variat glabriuscula aut hispidula, aculis aut subcaulescens, etc. Reperitur, in hortis præcipuè, scapis fasciatis, capitulis proliferis (J. Bauh. hist. 3. p. t. p. 113. f. 2), flosculis omnibus ligulatis vulgò (bot. mag. t. 228), et rarius floec. omnibus tubulosis ampliatis (Blackw. t. 530), ligulis albis aut purpureascentibus, etc. (v. v.)

*B. integrifolia* (DC. fl. fr. ed. 3. v. 4. p. 185. var. s), caulescens elongata, foliis integerrimis, capitulis minoribus. — *Bellis integrifolia* Lam. ! dict. 5. p. 6. non Michx. diù quasi spontanea in colle horti Parisiensis. (v. s.)

5. *B. sylvestris* (Cyr. pl. rar. 2. p. 22. t. 4), foliis subradicalibus obovatis crenatis trinerviis, invol. squamis pilosis vix acutis.  $\frac{1}{2}$  in campis apricis Siciliae (Guss.!), agri Neapolitani (Cyr. Ten.!), Pisani (Savi!), Nicænsis (C. Maun.!), Galloprov. (Rob.!), Avenionensis (Req.!), in Istriâ (Reich.), in Apenninis Appuanis (Berol.), Sardinia (Moris!), agro Byzantino (Cast.!), Smyrnæ (Parol.!), et in Mauritania Mogadorienæ (Brouss.!). Pedunculi longiores et capitulum majus quàm *B. perennis*. Invol. squamæ magis foliaceæ et disco paulò longiores. (v. s.)

*B. hybrida* (Ten. ! syll. p. 436. fl. med. 2. p. 64), caule basi subramoso, foliis elliptico-oblongis magnitud. *B. perennis*, invol. squamis disco subbrevioribus.  $\frac{1}{2}$  cum. var. *a.* Omnino media inter *B. perennem* et *B. sylvestrem* et fortè vera hybrida? (v. s.)

† *Species exclusæ.*

*B. ciliaris* Labill. = *Brachycome ciliaris*.

*B. droserifolia* Gon. = *Bellium bellidioides*.

*B. graminea* Labill. = *Paquerina graminea*.

*B. javanica* Blum. = *Lagenophora javanica*.

*B. magellanica* Lam. = *Lagenophora Commersonii*.

*B. pilosa* Blum. = *Lagenophora pilosa*.

*B. spinosa* G. alp. = *Plagus ageratifolius*.

*B. stipitata* Labill. = *Lagenophora Billardieri*.

CLVI. *BRACHYCOME* Cass. dict. 37. p. 464. et 491. — *Brachycomes* sp. Less. syn. 192. — *Brachyscome* Cass. bull. philom. 1816. p. 199. dict. v. 5. suppl. p. 63. — *Bellidis* sp. Labill.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis 1-seriatis, disci tubulosi 5-dentati hermaphroditis. Recept. conicum subalveolatum epaleaceum. Invol. campanulatum pauciseriale, squamis margine submembranaceis. Achænium lateraliter plano-compressum erostre pappo subsetiformi brevissimo coronatum. — *Herbæ Novo-Hollandicæ*. Folia alterna glabra parcè dentata pinnatilobata aut trifida. Discus luteus. Radius albus.

1. *B. stricta*, caule strictè erecto folisque pube brevi hirtellis, foliis oblongis basi attenuato-cuneatis grossè dentatis, invol. squamis apice membranaceis obtusissimis, fructu alà integrà marginato in disco subtuberculato.  $\frac{1}{2}$  in Novæ-Hollandiæ terrâ Van-Leuwin. *Bellis aculeata* Labill. ! Nov. Holl. specim. 3. p. 55. t. 206. *Brach. aculeata* Less. syn. 192. excl. syn. Sieb. *Herba rigidula erecta tota pubero-hirta*. Rami apice aphylli. Achænia vix tuberculata nec verè aculeata. (v. s. comm. à cl. Labill.)

2. *B. setzerodonta*, glabriuscula, caule adscendente aut erecto, foliis radicalibus petiolatis pinnatifidis, caulinis sessilibus oblongis apice grossè et acutè trilobatis basi subdenticulatis medio integerrimis, invol. squamis apice membranaceis acutis, fructu alà dentatà marginato in disco tuberculato.  $\frac{1}{2}$  in planis humidis Novæ-Hollandiæ interioris circa Bathurst et ad Lachlan river legit cl. A.

- Cunningham. Br. Billardieri Cass. dict. 5 suppl. p. 63. excl. syn. Folia forma et incisuris valde varians. Pappus setis longioribus et confertioribus quam ceter. specierum. (v. s. comm. à cl. invent.)
3. *B. SCAPIFORMIS*, caule erecto 1-cephalo apice nudo glabro sub capitulo confertè et brevè velutino, foliis radicalibus spathulatis obtusis crenatis in petiolum attenuatis, caulinis sessilibus oblongis linearibus, infer. apice serratis, ceteris integerrimis, invol. squamis margine membranaceis acutis, achænio obovato integro et tuberculato.  $\mathcal{L}$ .
- $\alpha$ . *puberula*, foliis pube brevi scabridis. — in Smith's Plains ad Lecklan river in Novâ-Hollandiâ inter. junio flor. legit cl. Cunningham. (v. s. comm. à cl. inv.)
- $\beta$ . *glabra*, foliis glabris. — in insulâ Van-Diemen legit cl. Gunn. (v. s. comm. à cl. Lindl.)
4. *B. SIEBERT*, glabra adscendens, foliis oblongis basi attenuatis acutè paucidentatis, invol. squamis subobtusis vix membranaceis, achænio obovato margine calloso integro. — in Novâ-Hollandiâ, Sieb. coll. exs. n. 485. *B. aculeata* Less. syn. 192. excl. syn. Labill. — Priori affinis sed differt glabritie totâ, caule adscendente debili gracili, capitulis dimidiò minoribus, invol. squamis angustis vix obtusis. Achænia obovato-compressa puberula margine callosa integra. (v. s.)
- $\beta$ . *Gunnii*, herba tota brevissimè subpuberula.  $\mathcal{L}$  Cl. Gunn legit in insulâ Van-Diemen. (v. s. comm. à cl. Lindl.)
5. *B. CILIARIS* (Less. syn. 192), glabra, foliis pinnatipartitis, lobis linearibus distantibus integerrimis, squamis invol. obtusis apice membranaceis, achæniis margine ciliatis. — in Novâ-Hollandiâ terrâ Van-Leuwin (Labill.) ad ora orientalem (Mus. Par.). Bellis ciliaris Labill. Nov. Holl. specim. 2. p. 56. t. 27. In specim. juniore achænia ciliata non vidi et ideo species mihi dubia. (v. s. comm. à Mus. reg. Par.)
6. *B. MULTIFIDA*, erecta glaberrima, foliis pinnatipartitis, lobis lineari-subulatis integerrimis acutis, invol. squamis obtusis, achænio oblongo immarginato utrinquè tuberculato.  $\mathcal{L}$  in umbrosis rupestribus Novæ-Hollandiæ int. ad Peel's Range junio flor. legit cl. A. Cunningham. Valdè accedit ad iconem 207 Labill. sed achænia certissimè non sunt ciliata, lobi foliorum tenuiores acutiores. (v. s. comm. à cl. invent.)
7. *B. ANGUSTIFOLIA* (Cunn. ! in litt. 1834), glabra adscendens aut erecta, foliis oblongo-linearibus integerrimis 1-nerviis, nervo valido, invol. squamis acutis, achænio integerrimo tuberculato, pappo brevissimo.  $\mathcal{L}$  in siccis glareosis apricis Novæ-Hollandiæ ad Goulburn Plains, maio flor. legit cl. Cunningham, et in ius. Van-Diemen cl. Gunn. Folia rigidula nervo prominulo 2 poll. longa, 2 line lata. Pappus minutissimus. (v. s. comm. à cl. Cunn. et Lindl.)
8. *B. LINEARIFOLIA*, glabra, foliis linearibus integerrimis indivisis. — in Novâ-Hollandiâ. Omnino refert figuram sinistram tab. 204. sp. fl. Nov. Holl. Labill. sed achænia certè pappo prioribus simili coronata nec nuda. (v. s. comm. à cl. Lambert. et Thibaud.)

†. *Species exclusæ.*

- B. DENTATA* Gaudich. = Vittadinia dentata.
- B. SPATULATA* Gaudich. = Vittadinia spathulata.
- B. TRILOBA* Gaudich. = Vittadinia triloba.
- B. XANTHOCOMOIDES* Less. = Keeria linearifolia.

CLVII. PAQUERINA Cass. dict. 37. p. 464 et 492. Less. syn. 192 — Bellidis  $\mathcal{L}$  Labill.

Capitulum multiflorum heterogamum, fl. radii ligulatis semine 1-seriatis, disci hermaphroditi tubulosi 5-dentatis. Recept. conicum

viter paleaceo-fimbrilliferum. Invol. campanulati squamæ subuni-ales. Achænium erostre epapposum plano-compressum margine sinuatum è scrobiculis recept. oriundum. — Herba Novo-Hollan-za glabra gracilis, tota foliata, simplex aut ramosa, ramis 1-cephalis. lina alterna lineari-oblonga integerrima basi attenuata.

P. GRAMINEA (Cass. l. c.). — in Novâ-Hollandiâ ad prom. Van-Diemen (Labill.) t ad fretum d'Entrecasteaux. Bellis graminea Labill. Nov. Holl. sp. 2. p. 54. 204. excl. fortè fig. sinistra. (v. s. comm. à Mus. reg. Par.)

VIII. LAGENOPHORA Cass. bull. philom. mart. 1818. Less. syn. 193. — agenifera Cass. bull. philom. 1816. decemb. — Microcalia A. Rich. — Bellidis p. DC. — Asteris sp. Lam. — Calendulæ sp. Forst.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis eriat, disci tubulosis 5-dentatis hermaphroditis masculisve steris; antheræ subliberæ. Recept. planum nudum. Invol. disco sub-ale, squamis circ. biseriatis acutis basi coriaceis adpressis, apice membranaceis. Achænia radii plano-compressa oblonga epapposa trata, disci abortiva. — Herbæ graciles, ad collum folia pauca propè in gerentes, scapo squamulato aphylo 1-cephalo. Folia basi atte-ata aut petiolata. Flores disci lutei, radii purpurei.

L. FORSTARI, foliis glabris obovato-orbiculatis acutè dentatis petiolatis, pe-iolis ciliatis. 2/ in Novâ-Zeelandiâ (Forst.) ad flumen Hawa-Hawa (D'Urv. l), oco gallicè dicto Havre de l'Astrolabe (Lesson!). Calendula pusilla Forst. prod. 05. Willd. sp. 3. p. 2344. Microcalia australis A. Rich. l. fl. Nov. Zeel. p. 231. oy. astr. 30. Petiolus limbo paulò longior, pilis articulatis rigidulis basi præ-ertum instructus. Limbi dentes infimi profundiores. Capitulum majus quàm in eque. (v. s. comm. à cl. D'Urville, Lesson et Rich.)

L. COMMENSALIS (Cass. dict. 25. p. 110), foliis glabris obovatis obtusis inte-gerimis aut apice pauci-dentatis in petiolum brevem attenuatis. (?) ? ad fretum agellanicum (Comm. l), in Staten-Island (Webster. l in h. Lindl.), in sum-is Andibus Chilensibus (C. Gay!), et in insulâ Tristan d'Acunhâ (Pet. Th.). later nudiculis Comm. herb., Lam. l. dict. 1. p. 308. ill. t. 681. f. 4. Calendula pusilla var. ß. Forst. prod. 305. Calendula Magellanica Willd. sp. 3. p. 2344. Calendula pusilla Pet. Th. flor. Trist. d'Ac. p. 40. t. 9. Bellis Magellanica DC. n Lam. dict. enc. 5. p. 7. (v. s.)

L. BILLARDIERI (Cass. dict. 25 p. 111), foliis utrinquè hirtellis oblongis basi a petiolum brevem attenuatis sinuato-dentatis. 2/ ? in Novâ-Hollandiâ ad prom. Van-Diemen et alibi. Fortè species plures hic commixtæ nempe :

a. pusilla, foliis utrinquè hirsutis vix basi attenuatis, scapo sesquipollicari, invol. squamis exterioribus lineari-subulatis subpatulis. — Specim. habeo à cl. Labillardière sub nom. Bellidis stipitata. (v. s.)

ß. media, foliis utrinquè sparsè hirtellis basi sensim attenuatis, scapo 4-5-pollic., invol. squamis linearibus subacutis subadpressis. Bellis stipitata Labill. Nov. Holl. spec. 55. t. 205. — Habui ex coll. Sieb. l. n. 505. ex herb. Lamb. et cult. in hort. Kew. (v. s.)

γ. glabrata, foliis utrinquè glabresculis basi sensim attenuatis et verè petiolatis, scapo 4-6 pollic., invol. squamis erecto-adpressis. (v. s. comm. à cl. Lambert.)

L. HIRSUTA (Pœpp. ex Less. in. linnæâ 1831. p. 131), foliis utrinquè hirsutis petiolatis obovatis grossè dentatis, dentibus mucronatis, scapo vix foliis du-plò longiore. 2/ in Chili Pœpp. l. coll. exs. n. 715. Folia cum petiolo 9-10 lin-terè bipollicaria, 9-10 lin. lata membranacea. Invol. squamæ erectæ acuminatæ disco paulò longiores. Achæni rostrum glanduloso-viscosum. (v. s.)

CLIX. IXAUCHENUS *Cass. dict.* 56. p. 176. — *Ixiauchenus Less. syn.* 193.

Capitulum multiflorum heterogamum, fl. radii biseriatis fœmineis ligulatis longissimis, disci tubulosi 5-dentati hermaphroditis aut abortu masculis. Recept. nudum latum planum. Invol. subhemisphaericum disco æquale, squamis pluriseriatis laxè adpressis foliaceis. Achænia ovali-oblonga compressa glabra in collum brevem glutinosum attenuata epapposa. — Herba Novæ-Hollandiæ?, glabra basi tantum foliata. Canis scapiformis aphyllus 1-cephalus. Folia alterna subradicalia valdè inæqualia semiamplexicaulia ferè lyrata. Discus flavus. Radius albus. An genus à *Lagenophorâ* aut à *Myriactide* satis diversum?

1. I. SUBLYRATUS (Cass. l. c.). — in Novâ Hollandiâ propè Port-Jackson? *Ixiauchenus lyratus Less. l. c.*

CLX. MYRIACTIS *Less. in Linnæi* 1831. p. 127. *syn.* 193. — *Lavania sp. Wall. litt.* — *Balsamita sp. Wall. herb.* 1821.

Capitulum multiflorum heterogamum, fl. radii bi- aut multiserialibus fœmineis angustè ligulatis integerrimis, fl. disci hermaphr. tubulosi 5-lobis. Recept. nudum. Invol. 1-2-seriale, squamis linearibus acutis. Achænia compresso-plana calva glabra erostria apice sæpè glandulifera. — Herbæ Indiæ aut Persiæ erectæ subdichotomo-ramosæ. Folia alterna ovata aut lanceolata dentato-serrata, suprà pube minimâ scabrida. Pedunculi elongati 1-cephali subpaniculatim dispositi. Capitula globosa albida aut flavida.

§. 1. *Ligulis circiter biseriatis, achænio apice vix subattenuato. — Species Indiæ australioris aut insularis, mediæ inter Lagenophoram et Myriactidem veram.*

1. M. WIGHTII (DC. in Wight contrib. bot. ind. p. 10), parçè pilosa, foliis inferioribus ovatis basi longè cuneato-attenuatis grossè inciso-dentatis, superioribus oblongis integris sessilibus dentium et folii ipsius apicibus calloso-imbricatis ½ in Indiâ orientali ad montes Neelgherry legit cl. Wight. *Carpesium nov. sp. Wight! herb.* Folia radicalia in petiolum attenuata ovata, caulin. infer. basi cuneata parçè dentata, summa subsessilia integra utrinquè acuminata. Capitula terminalia solitaria 4 lin. diam. Invol. hirsutiusculum post anthesin reflexum Ligulæ albidæ subbiseriales per exsic. revolutæ. (v. s. in h. Wight.)

2. M. JAVANICA, caule erecto aut adscendente glabriusculo apice ramoso, foliis ovalibus in petiolum attenuatis distanter et calloso-serratis sparè pilosis ciliatisve. ½ in insulæ Javæ sylvis altioribus montis Gede legit cl. Blume. *Bellis Javanica Blum.! bijdr. p. 917. Lagenophora Javanica Blum.! in litt. 1831.* Herba sesquipedalis. Rami subaphylli. Capitula ferè priora. Ligulæ plerimæ Recept. convexum. (v. s. comm. à cl. Blume.)

3. M. PILOSA (Blum.! in litt. 1831), tota pilosa, caule suffruticuloso adscendente, foliis ovato-lanceolatis acuminatis profundè dentatis, racemo terminali paucifloro. — in summo montis Gede ins. Javæ. *Bellis pilosa Blum.! bijdr. 917. Lagenophora pilosa Blum.! in litt. 1831.* Capitula priora sed minora. Recept. minus elevatum. (v. s. sine flor. comm. à cl. Blume.)

§. 2. *Ligulis pluriserialibus, achænio apice omninò erostrato. — Species Nepalenses.*

4. M. NEPALENSIS (Less. ll. cc. in lin. t. 2. f. F. 1), caule glabro, involucri biseriali capitulo multò breviori, receptaculo convexo ad centrum subconico.

— in Nepaliâ, *Lavenia sphaerantha* Wall.! cat. et herb. n. 327. Capitula globosa Pisi magnitudine. (v. s.)

**M. WALLICHII** (Less. II. cc.), caule hirsutiusculo, involucri subniseriali capitulo subequali, receptaculo plano. — in montibus excelsis Nepaliâ borealis. *Balsamita dentata* Wall.! cat. n. 326. Capitula dimidiò quàm prioris minora. (v. s.)

β. *hispidior*, caule et foliis utrinquè pilis sparsis hispidissimo. — in montibus Indiâ bor. occid. legit cl. Royle. (v. s.)

**M. Gmelini**, erecta pilis articulatis setulisque parè hirsuta, foliis petiolatis membranaceis grossè dentatis, invol. squamis 2-3-seriatis lineari-oblongis ciliato-fimbriatis, achæniis basi glandulosis in collum globulo glanduloso terminatum productis. 2½ in Persiâ prov. Gilan legit Gmelin jun. et in sylvis montium Talsch rev. Hohenbaker. *Botriyadenia* Gmelini Fisch. et Mey. in litt. 1835. Folia ferè *Callistephi* Chinensis. Habitus *Dicrocephalæ* latifoliæ. Descrip. ex F. et M. litt. (v. s. capitulum à cl. auct. miss.)

**XL. GARULUM** Cass. dict. 18. p. 162. 30. p. 324, Less. syn. 194. excl. syn. Tourn. — *Osteospermum* sp. Ait. *L'Hér. Thunb.*

Capitula radiata, fl. radii longè ligulatis fœmineis, disci tubulosi dentatis masculis, ovulis abortivis. Recept. convexum nudum. Invol. uamæ biserialis. Stylus disci ramis apice liberis divergentibus, extus liberis, intus ad margines glandulosus. Ovaria disci oblonga. Achænia dii obovato-oblonga 3-4-costata erostrata epapposa sicco-coriacea, ricarpio tenui extus rugoso et subaspero, disci plano-compressa læva. — Herbæ suffrutescentes Capenses erectæ glabræ aut subviscidolabrescentes. Folia alterna pinnatisecta, lobis plùs minùs dentatis. Capitula terminalia pedunculata, radio cœruleo, disco luteo, corollis uti puberulis.

**G. PINNATIDUM**, foliis pinnatifidis, lobis ovatis dentatis. 3 ad Cap. Bonæ-Spei (Th.) ad Sneeuobergen alt. 4-500 ped. (Dreg.). *Osteospermum pinnatifidum* Thunb. prod. 167. fl. cap. 717. *L'Hér.*! st. nov. 11. t. 6. Ost. cœruleum Ait. kew. 3. p. 275. Jacq. ic. rar. 1. t. 179. coll. 1. p. 78. *Garaleum viscosum* Cass. dict. 18. p. 325. *Chrysanthemoides cœrulea* Moench. meth. 584. — *Zeyh.*! pl. exs. cap. n. 102. in h. Moric. Herbæ odor gravis. (v. v. c.)

**G. BIPINNATUM** (Less. l. c.), foliis bipinnatisectis, lobis lineari-setaceis acutè lobulatis aut integris. 3 ad Cap. Bonæ-Spei, in Hantum (Th.), in Carro (Dreg.! Eckl.!), ad Zondag rivier et Graaf rivier (Dreg.!). Achænia radii trigona irregulariter transversè sulcata, disci inania plano-compressa lævia. *Osteospermum bipinnatum* Thunb. fl. cap. 717. (v. s.)

CLXII. KEERLIA. — *Brachycomes* sp. Less.

Capitulum multiflorum radiatum, ligulis circ. 15-25 fœmineis 1-2-rialibus linearibus, fl. disci hermaphr. aut abortu masculis 5-dentatis revibus. Invol. latè campanulatum imbricatum pluriseriale, squamis pœcolatis acuminatis margine submembranaceis. Recept. planum nudum. Styli radii breves obtusi, disci erecti dorso subpuberuli. Achænia abra obpyramidata seu cylindracea sæpius longitudinaliter sulcata. Pappus paleis brevissimis pluribus acutis basi subconcretis et ideò ferè ironam minutissimam conficientibus. — Herbæ Mexicanæ glabræ aut abe brevi velutino-villosulæ erectæ parè ramosæ. Folia alterna oblonga aut cuneata sessilia seu semiamplexicaulia integræ aut apice dentata. Rami apice nudi 1-cephali. Flores albi aut flavi. — Genus à *Brachycomi* (confirmante habitationum diversitate) recept. plano et achæniis teretiustulis nec lateraliter plano-compressis distinctum, dicavi cl.



F. W. Keerl peregrinatori germano qui unam generis speciem primus collegit.

1. *K. LINNÆIFOLIA*, glabra, caule imâ basi suffruticuloso subradicante, foliis linearibus acutis integerrimis, invol. squamis coriaceis, ligulis circ. 15-18 ovali-oblongis, corollarum omnium tubo basi attenuato.  $\frac{1}{2}$   $\frac{3}{4}$  in Mexico legit d. Alaman. Ligulæ ex sicco videntur flavæ undè verisimil. huc referenda *Brachycome xanthocomoides* Less. syn. 192, et in linnæa 1834. p. 265. *Achæmia* (juniora tantum obvia) teretia non aut nondum sulcata. (v. a.)
2. *K. RAMOSA*, subpubescens, caule imâ basi suffruticuloso erecto ramoso, foliis semiamplexicaulibus sublinearibus integerrimis, invol. squamis margine huius membranaceis, ligulis 20-25 linearibus, corollarum omnium tubo basi attenuato.  $\frac{1}{2}$   $\frac{3}{4}$  in Mexico circa Tlapojahua legit d. F. W. Keerl. Ligulæ albe parco subtus admissio rubore. *Achæmia* sulcata ovato-subteretia. (v. s. in herb. Mart.)
3. *K. SKIRRHOBASIS*, pube brevi confertâ velutino-pubescenti, caule herbaceo erecto parce ramoso aut simplici, foliis sessilibus obtusis oblongis basi attenuatis integerrimis aut apice dentatis, invol. squamis margine membranaceis, ligulis 20-25 linearibus, corollarum omnium tubo basi demum incrassato et indurato. (1) in Mexico inter Bejar et flum. Triñtatis junio florentem legit d. Berlandier (pl. exs. n. 1755). Priori similissima sed radix exilis evidenter annua et car. supra indic. facile ab eâ separant. (v. s.)

### CLXIII. APHANOSTEPHUS.

Capitulum multiflorum heterogamum, fl. radii ligulatis 1-serialibus fœmineis, disci hermaphr. 5-dentatis. Recept. valdè convexum nudum. Invol. squamæ 2-seriales lanceolatæ acuminatæ margine membranaceæ. Achæniû teres striatulum pappo parvo coronæformi integro superatum. — Herbæ Americanæ erectæ ramosæ pubescentes. Folia sessilia alterna inciso-dentata aut parce lobata. Rami apice nudi 1-cephali. Discus flavus. Ligulæ albæ invol. duplò longiores. — An *Brachycome xanthocomoides* Less. syn. 192. huc referenda? sed non satis nota et recept. dicitur planiusculum. An *Chrysanthemeis*, an *Bellideis* adjungendum?

1. *A. RAMOSISSIMUS*. (1) in Mexico circa Bejar et Laredo legit d. Berlandier (pl. exs. n. 2061). Habitus ferè *Erigerontis* aut *Distaseos*. Radix simplex exilis. (v. s.)

### Div. III. CHRYSOCOMEÆ.

Recept. epaleaceum nunc nudum, nunc alveolatum aut fimbrilliferum. Capitula lutea nunc homogama corollis omnibus tubulosis hermaphroditis, nunc radiata ligulis fœmineis aut neutris cum disco homochromis.

### Subdiv. 1. GYMNOSPERMEÆ.

Pappus nullus. — Subdiv. analogæ *Bellideis*.

### CLXIV. XANTHOCOMA H. B. et Kunth nov. gen. am. 4. p. 311. t. 112. Less. syn. 192.

Capitulum multiflorum radiatum, fl. radii 1-seriatis fœmineis ligulatis, disci hermaphrod. tubulosis 5-dentatis. Recept. vix convexum subfoveolatum. Invol. campanulati squamæ basi imbricatæ apice subfoliaceæ laxiusculæ. Ligulæ disco duplò longiores. Achænia compressa calva. — Herba Mexicana cæspitosa glabra. Folia alterna linearia integerrima. Capitula solitaria terminalia. Cor. omnes flavæ.

1. *X. HUMILIS* (H. B. et Kunth l. c.).  $\frac{1}{2}$  in humidis Mexici circa urbem (Berl. pl. exa. n. 374 et 538) inter pagum Carpio et lacum Sanoti-Christophori (H. et B.). *Chrysanthemum humile* Spreng. syst. 3. p. 584. *Chrysanthemum arvense* herb. Mexic. anon. Caples diffusis interdum proliferi. Fol. radice lanceolato-linearia in petiolam attenuata. (v. s.)

## CLXV. XEROTHAMNUS.

Capitulum multiflorum monoicum, fl. radii 1-serialibus ligulatis femineis, disci tubulosi 5-dentatis sterilibus. Invol. imbricatum, squamis lanceolato-subulatis rigidulis apice subdivergentibus. Recept. nudum areolatum. Ligulæ revolutæ 4-nerviæ. Cor. disci tubo brevi angustissimo, fauce cylindrica, dentibus erectis acutis. Anth. ecaudatæ. Stylus radii bifidus lobis acuminatis glabris, radii indivisus apice capitellatus. Achænia radii subtrigona dorso convexo curvato, areolâ terminali ad lateris interni mediam longitudinem, disci abortiva recta obovata, mucrone brevi superata. Pappus nullus. — Suffrutex Capensis rigidulus ramosus erectus Stypheliæ seu Epacridis facies. Folia alterna conferta subtriquetra acuta dorso versus apicem et margine muricato-acabra sicca patulo-squarrosa, Pedunculi versus apices ramorum 1-3 erecti, squamis paucis distantibus instructi, 1-cephali. Flores lutei. — Ordinatio obscura! An *Asterea*?

1. *X. ECKLONIANUS*.  $\frac{1}{2}$  ad Cap. Bonæ-Spei in distr. Uitenhage legit. cl. Ecklon! et Zeyher. fl. martio. Fruticulus semipedalis. Folia 4 lin. longa, vix lineam lata. (v. s.)

## CLXVI. ANAGLYPHEA.

Capitulum multiflorum heterogamum, fl. radii 1-seriatim ligulatis femineis, disci tubulosi 5-dentatis hermaphrod. Invol. 2-seriale, squamis aequalibus disco paulo longioribus acuminatissimis. Recept. plenum alveolatum. Cor. tubus hirsutus. Anth..... Styl..... Achænia obovatum puberum calvum. — Suffrutex Capensis. Caules et præsertim ramorum apices pube confertâ brevi glanduloso-subvelutini. Folia alterna linearia, apice recurva acuminata, utrinque striato-sulcata, ciliis rigidis patulis secus margines et subhis versus apicem in nervis scaberrima. Capitula parva terminalia solitaria. Flores disci et radii flavi. — Genus ex unico specimine stabilitum et recognoscendum!

1. *A. ASPERA*.  $\frac{1}{2}$  in Africa Capensis distr. Albany legit cl. Dregé! Habitus *Mairia* taxifolia seu *Felicia* ciliaris, sed car. gen. diversi, ligulæ luteæ 4-nerviæ, folia striato-sulcata multo magis setis aspera, etc. Folia 7 lin. longa vix lineam lata rigidula. (v. s.)

## CLXVII. GYMNOSPERMA Less. syn. 194. — Sellow sp. Spreng. non Kunth.

Capitulum pauciflorum subradiatum, fl. radii 3-5-femineis 1-serialibus angustissimis brevissimè ligulatis interdum nullis, fl. disci 3-10 regularibus hermaphroditis aut sterilibus. Recept. nudum angustum. Invol. oblongum, squamis imbricatis adpressis scarioso-coriaceis. Achænia crostria subcompressa omnino calva. — Suffrutescentes Americani glabri erecti corymboso-ramosi. Folia sessilia integra oblonga aut linearia acuta punctata sæpè glutinosa. Capitula oblonga ad apices ramulorum terna aut aggregata sæpius corymboso-fastigiata. Cor. luteæ. — Genus accedens hinc ad *Brachyridem* sed pappus nullus, illinc ad *Madiam* sed ligulæ minutissimæ, invol. palæ achænia non involventes, etc.

§. 1. *Foliis alternis.*

1. *G. GLUTINOSUM* (Less. l. c.), suffruticosum, ramis teretibus, foliis lineari-lanceolatis integerrimis trinerviis glutinosis, paniculâ fasciculatâ, ligulis minimis, disco sterili. ♀ in Brasiliâ. *Selloa glutinosa* Spreng. pug. p. 36. n. 35. nov. prov. 36. syst. 3. p. 496. Ker. bot. reg. t. 462. Molina viscosa hort. berol. ex Spreng.
2. *G. CORYMBOSUM*, fruticosum, ramulis subangulatis dichotomo-corymbosis, foliis oblongis utrinquè attenuatis subviscosis trinerviis, venis lateralibus tenuissimis, capitulis ad apicem ramuli cujusque subterais 8-floris, ligulis 5, disco circ. 3-floro. ♀ in Mexico circa Matamoras legit cl. Berlandier (pl. ex. n. 2296). Folia 12-15 lin. longa, 2 lin. lata. (v. s.)
3. *G. MULTIFLORUM*, fruticosum, glutinosum, ramulis angulatis, foliis linearibus acutis trinerviis venis lateralibus tenuissimis, capitulis ad apices ramulorum 3-5 confertissimè corymbosis 13-14-floris, ligulis 5, fl. disci 7-8. ♀ in Mexico (Alam.! Mairet!) circa urbem (Berl. pl. ex. n. 50). Flor. septemb. Molina glutinosa Alam. in litt. Priori affine, Folia 2 poll. longa, 1½-2 lin. lata. (v. s.)
4. *G. SCOPARIUM*, fruticosum subviscosum ramosisimum, ramis erectis confertis apice paniculato-corymbosis, foliis oblongo-linearibus acutis trinerviis, venis lateralibus tenuibus, capitulis ad apices ramulorum plurimis confertis circ. 10-floris, ligulis et tubulis 5. ♀ in Mexicanæ confederationis statu Sancti-Ludovici de Potosi legit cl. Berlandier (pl. ex. n. 1338). Fol. 9-10 lin. longa, 1½-2 lata. (v. s.)

§. 2. *Foliis oppositis.*

5. *G. NUDATUM*, caule herbaceo lævi angulari, foliis oppositis crassius integerrimis linearibus, capitulis in corymbum compositum trichotomum dispositis, bracteis filiformibus, tubulis 5-8, ligulis subnullis. ♀ in Floridâ orientali. *Selloa nudata* Nutt. in Sillim amer. journ. 5. 1822. p. 300. Affinis dicitur *Chrysocoma* et potius ut videtur *Bigelowiæ*.
6. *G. ? OPPOSITIFOLIUM*, caule herbaceo lævi tereti, foliis oppositis linearibus submembranaceis integerrimis aut remotè denticulatis, capitulis in corymbum compositum trichotomum dispositis, bracteis lineari-filiformibus, floribus 10 omnibus tubulosis. — in Mexico inter San-Ferrado et Santander legit cl. Berlandier (pl. ex. n. 2263!). Herba erecta 1-1½-pedalis. Folia 3-4 poll. longa, 3 lin. lata. Corymbus polycephalus flavus. Habitus omninò sect. 1, sed foliis oppositis et capitula discoidea. (v. s.)

Subdiv. 2. *ACHRYDEÆ.*

*Pappus paleaceus multiradiatus* (ut in *Bellieis*).

CLXVIII. *BRACHYRIS* Nutt. gen. am. 2. p. 163. Less. syn. 191. — *Brachyachrys* Spreng. syst. 3. p. 574.

Capitulum pluriflorum radiatum, fl. radii 5-10 ligulatis 1-serialibus fœmineis, disci 3-10 tubulosis 5-dentatis hermaphroditis. Recept. nudum. Invol. ovati aut cylindracei squamæ arcuè imbricatæ. Achænia obconica erostria. Pappus paleis 5-8 oblongis persistentibus. — Herbe Occident.-Americanae rhizocarpicae aut suffruticosae glabrae erectae. Folia alterna linearia seu lanceolata integra, sæpè punctata. Capitula solitaria aut terna ad apices ramulorum corymbosa. Invol. squamæ pallidè apice maculâ viridè notatæ. Cor. luteæ. — Habitus hinc *Euthamiae*, illinc *Bigelowiæ* sed pappo paleaceo nec piloso distincta, ex *Ααχρὶς*, brevis, et *αχρὺς*, palea.

Sect. 1. *EUBRACHYRIS.*

Achænia radii pappo fl. disci simili donata. Fl. disci numero ligularum aequali aut pauciores.

\* *Suffruticosæ.*

**EPHIALA**, caule suffruticuloso corymboso ramoso tereti vix striatulo, stissimè linearibus patulis margine subrevolutis, corymbo fastigiato, invol. paulò longioribus, fl. disci 1-3.  $\zeta$  in Mexico prope locum *lito* legit cl. Berlandier januar. florent. Invol. parva oblonga glutinosa in sicco revolutæ. Folia pollicem longa vix semilin. lata. (v. s.)

**LATA**, caule suffruticoso paniculato ramoso teretiusculo, foliis linearibus punctatis planis crassiusculis, capitulis oblongis in paniculam fl. radii 5 invol. paulò longioribus, disci 5-6.  $\zeta$  in Chili circa Copiã legit cl. Gaudichaud et circa Acapulco cl. Henke (ex h. Sternb.). Etiam in loci designatione olim à cl. Née lectam. Folia 7-9 lin. longa, ferè s. (v. s.)

\*\* *Herbaceæ.*

**MLA** (Nutt. gen. am. 2. p. 163), caulibus numerosis marginato-scabris, foliis adproximatis angustè linearibus integris punctatis corymbo dichotomo fastigiato, fl. radii 5 invol. circ. longitudine, fl. in collibus aridis ad fl. Missouri, ab Arikarees ad montes. Solidago Pursh fl. bor. am. 2. p. 540. Brachyachyris Euthamiæ Spreng. l. c. semica subfetida strenuè diuretica.

**ANICA**, caulibus teretibus apice subpaniculatis, foliis linearibus acutis serratis et subciliatis planis crassiusculis, paniculâ racemosâ oligocephalâ ovatis, ligulis circ. 8-10 invol. paulò longioribus, fl. disci 8-10.  $\sigma$ nia legit cl. Douglas. Folia pollicem longa, vix lineam lata. (v. s. bon. soc. hortie. Londin.)

**IA**, rhizomate lignoso, caulibus pluribus subangulatis apice monophthalis, foliis linearibus acutis crassiusculis viscosis, infer. oblongo, capitulis obovatis ligulis 8-10 disco duplò ferè longioribus, fl. disci abeo ex herb. Neei sine loci natalis design. verisim. ex Mexico aut ex cl. squamæ minùs quàm in prioribus adpressæ. Herba vix digiti crans. Folia pollicem longa, 1-2 lin. lata. (v. s.)

**OLIA**, herbacea, caule erecto tereti, foliis ovatis acutis petiolatis serratis, capitulis subsessilibus glomeratis in racemum elongatum subsimplicem unilateralem dispositis.  $\mathcal{L}$  in agro Kentuckensi ad ripas legit cl. Rafinesque. Solidago sphacelata Raf. l. a. n. 1820 ex specim. Species distinctissima! Capit. oblongum squamæ inv. subsphacelataridi non notatæ. Ligulæ 5. Fl. disci circ. 4-5. Pappus paleis 5-6.

## Sect. II. AMPHIACHYRIS DC. not. VII.

adii ferè calva aut coronâ brevissimâ donatæ. Flor. disci ligulis un-

**SCULOIDES** (DC. not. VII. jard. gen. p. 1. t. 1), caulibus erectis rathibus, foliis lanceolato-linearibus punctatis uninerviis aut basi subnervine scabridis, corymbo laxo, ramis oligocephalis, invol. obovato la bracteato, ligulis 10-11 ovali-oblongis invol. longioribus, fl. disci in territorio Arkansano semina legit cl. de Pourtalès. Herba priorihabitu Galatellæ aut Euthamiæ graminifoliæ. Folia 3 lin. longa. revolutum. (v. v. in hort. Genev. comm. à cl. de Charpentier.)

**SSISIMA**, foliis linearibus angustissimis uninerviis.  $\mathcal{L}$  in terr. Arkansiam semina legit cl. de Pourtalès. Folia lineam lata punctata. Capit. id prioris. (v. v. in h. Gen.)

## CLXIX. HEMIACHYRIS.

in pluriflorum, fl. radii circ. 5 ligulatis fœminicis, disci 7-8 dentatis hermaphr. Invol. ovatum, squamis oblongis adnatis. Recept. nudum. Styli fl. disci appendiculati hirtelli.

Achænia obovato-teretiuscula villosiuscula, radii calva, disci squamellis ovatis subacutis brevibus coronata. — Herba suffruticosa Texana erecta ramosissima glabra. Rami juniores sulcato-angulati, adulti teretes. Folia alterna linearia integerrima uninervia, summa ferè subulata. Capitula parva ramulos terminantia solitaria. Flores lutei. — Affinis Brachyridi sed pappi paleæ brevissimæ squamiformes nec lineares. An Euheleniis potiùs referenda? — Nomen alludit ad achænia alia calva, alia squamellas gerentia.

1. H. TEXANA. ♂ in Mexici prov. Texas legit cl. Berlandier prope Bojer. Flor. aug. (Berl.) pl. exs. n. 1763 et 1765). Caulis  $1\frac{1}{2}$ -pedalis parèe foliosus. Folia infer. pollicem longa,  $1\frac{1}{2}$  lin. lata, summa multò minora. (v. s.)

CLXX. LEPIDOPHYLLUM Cass. bull. philom. 1816. p. 199. dict. 26. p. 36. Less. syn. 191. — Conyzæ sp. Lam.

Capitulum pluriflorum radiatum, ligulis 2-3 fœmineis brevibus 1-serialibus, disco 4-6-floro hermaphr. regulari. Invol. oblongo-cylindraceum, squamis paucis obtusis imbricatis. Recept. parvum nudum. Achænia oblonga striata. Pappus 1-serialis, setis paleaceis confertis inæqualibus margine sublimbriatis. — Frutex glaber ramosus habitu ferè Pteroniæ, caract. Brachyridis, ex Amer. austr. extremâ ortus. Folia opposita quadrifariam imbricata ovato-subtrigona coriacea parva integerrima. Capitula ramulos terminantia sessilia solitaria. Cor. flavæ.

1. L. CUPRESSIFORME (Cass. dict. 26. p. 37). ♂ ad fretum Magellanicum inter oram Patagonicam et sinum Boucaut leg. Commerson, et ad portum Desideratum Née! Conyzæ cupressiformis Lam. dict. 2. p. 91. ill. t. 697. f. 3 Baccharis cupressiformis Pers. ench. 2. p. 425. n. 50. Athanasia? cupressiformis Comm. ined. (v. s.)

CLXXI. GRINDELIA Willd. mag. naturforsch. Berl. 1807. p. 261. Dunal in mem. mus. h. n. Paris. 5. p. 48. — Demetria Lag. gen. nov. in el. h. madr. 1814. p. 30. — Donia Brown in h. kew. ed. 2 (1813) v. 5. p. 82 non Don. — Grindelia et Aurelia Cass. dict. 37. p. 468. — Grindelia et Donia Desf. Link, etc. — Grindelia et Doniæ sp. Less. — Asteris sp. Cav.

Capitulum multiflorum heterogamum, fl. radii 1-seriatis ligulatis (in unicâ spec. nullis), disci tubulosis 5-dent. hermaphr. Recept. foveolatum. Invol. basi hemisphæricum, squamis pluriserialibus imbricatis. Antheræ basi nudæ nec caudatæ. Achænia obovata aut subangulata glabra. Pappus pauciaristatus deciduus, aristis crassis rigidis. — Suffrutices aut Herbæ Americanæ ad apices et præsertim ad involucri glutiniferæ. Folia alterna serrata aut integra, radicalia sæpitis spatulata, caulina sessilia aut semiamplexicaulia. Capitula ad apices ramorum solitaria. Flores flavi. — Genus naturalissimum etiãsi ab auctoribus etiã gravissimis variè vexatum.

1. G. GLUTINOSA (Dunal in mem. mus. h. n. Par. 5. p. 49), fruticosa glabra apice glutinosa, foliis sessilibus breviter decurrentibus obovatis obtusis apice serratis punctato-pellucidis, capitulis terminalibus solitariis præsertim ante explicationem valde glutinosis, pappo 5-8-seto. ♂ in Mexico. Aster glutinosus Cav. ic. 2. p. 53. t. 168. Doronicum glutinosum Willd. sp. 3. p. 2115. Inula glutinosa Pers. ench. 2. p. 452. Donia glutinosa Brown h. kew. ed. 2. v. 5. p. 82. Demetria glutinosa Lag. gen. in elench. h. madr. 1814. p. 30. Aurelia decurrens Cass. dict. 37. p. 468. Pappi setæ rigide triquetrae secus angulo acabræ. (v. v. c.)

2. *G. squarrosa* (Dunal l. c. p. 50), herbacea glabra, caule parçè ramoso. ramis apice 1-cephalis, foliis semiamplexicaulibus oblongis serrulatis, invol. squamis glutinosi in acumen filiforme recurvato-squarrosos productis, pappo 2-4-setoso. ♀ in pratis apricis secus rivos fluminis Missouri (Pursh). *Donia squarrosa* Pursh fl. bor. am. 2. p. 559. bot. mag. t. 1706. *Aurelia amplexicaulis* Cass. dict. 37. p. 468. (v. v. c.)
3. *G. subdecurrens*, suffruticulosa glabra parçè ramosa, ramis 1-cephalis, foliis ovato-oblongis per totam longitudinem calloso-serratis, infimis breviter decurrentibus, super. cordato-sessilibus, capitulis ebracteatis glutinosis, invol. squamis lineari-subulatis squarrosis, pappo 2-4-seto. ♂ in Mexico circa Villalpando legit cl. Mendez. Suffrutex circ. pedalis. Rami albi glabri. Pappus caducissimus. Habitus hinc *G. squarrosa*, illinc *G. pulchella*. (v. s.)
4. *G. ruizoides* (Willd. mag. nat. Berl. 1807. p. 261. enum. 894), caule basi suffruticoso apice pubescente aut sparè hirsuto, foliis radicalibus spatulatis, caulinis subovatis serratis basi latioribus subamplexicaulibus, ramis apice 1-cephalis subhirtellis, invol. squamis glabris in acumen lineari-subulatum patulum productis, pappo 1-3-seto. ♂ in Mexico (Lag.), planitie altà (H. et B.), inter Vittoria et Tula (Berl. n. 2186), in ditone Texas (Berl. n. 1542 et 2186) et Oaxaca (Andr. n. 271). Dunal l. c. p. 50. t. 1. Cass. dict. 37. p. 468. bot. reg. t. 248. Ast. spatularis Brouss. est serratus Lag. ex Steud. Grind. spatulata Link enum. h. ber. 2. p. 336. *Aster spatulatus* hort. madr. 1805. *Inula serrata* Pers. ench. 2. p. 451. *Demetria spatulata* Lag. el. h. madr. 1814. p. 30. (v. v. c. et a. sp.)
5. *G. microcephala*, caule basi suffruticuloso glabro, ramis apice parçè pubescentibus, foliis sessilibus subsemiamplexicaulibus ovato-oblongis serratis subpubescentibus, capitulis ramos terminantibus solitariis, invol. squamis glabris in acumen lineari-subulatum productis, pappo disci 4-5-seto, acheniis radii ovatis vix angulatis levibus. ♂ in sylvia prov. Mexicanæ Texas legit cl. Berlandier (pl. exs. n. 2057) Affinis *G. inuloidi* sed folia et capitula dimidiò minor. Flores disci steriles in specim. meo videntur. Achenia radii matura grossa calva. (v. s.)
6. *G. anguta* (Schrader in litt.), herbacea subcaespitosa pubescenti-hirtella, foliis infer. spatulatis in petiolum attenuatis, super. oblongis basi angustatis semiamplexicaulibus, omnibus distanter et acutè dentatis, capitulis terminalibus solitariis pedicellatis, invol. squamis linearibus acutis, pappo trisetato. ♀ in Mexico circa Valladolid de Mechoacan et Pazuaro (H. et B.). *Grindelia angustifolia* H. B. et Kunth nov. gen. am. 4. (1820) p. 309. bot. reg. t. 781. non Dunal. An *Grind. spatulata* Bern. in Link enum. h. ber. 2. p. 336. sed nostra est nec annua nec glabra. (v. s. c.)
7. *G. angustifolia* (DC. in Dunal l. c. 1819. p. 51. t. 3. non Kunth), caule subherbaceo glabro, foliis angustis integerrimis, radicalibus spatulatis, caulinis distantibus ovatis acutis sessilibus, capitulis pedunculatis, invol. squamis lanceolatis adpressis. ♀ ♂ in Mexico. Asteris sp. Moq. fl. mex. ined. *Grindelia Duvalii* Spreng. syst. 3. p. 575. (ubi Duval loco Dunalii ter citatur.)
8. *G. pulchella* (Dunal l. c. p. 51. t. 2), suffruticosa glabra erecta apice glutinosa, ramis teretibus, foliis caulinis sessilibus oblongis acutis hinc indè acutè dentatis, capitulis ad apices ramorum solitariis foliis 2-3 proximè bracteatis, invol. glutinosi squamis lanceolatis acutis adpressis, pappo 3-aristato. ♀ in Amer. legit cl. Née! circa Mendoza Chilensium ex herb. hort. monsp. sine loci design. in h. Thib. nunc meo. An verè Chilensis? (v. s. ex itin. Née.)
9. *G. integrifolia*, caule herbaceo erecto ramoso sparè hirsuto, foliis caulinis semiamplexicaulibus ovato aut oblongo-lanceolatis acutis integerrimis membranaceis puberulis margine scabridis, capitulis ad apices ramorum solitariis inter folia suprema bracteantia sessilibus, invol. squamis extus puberis intus glutinosis in acumen filiforme subulatum productis, pappo 1-3-aristato. ♀ in America borealis parte occid. (v. s. ex hort. Soc. hortic. Lond. à filio lect.)

10. *G. RUBRICAULIS*, caule herbaceo basi glabro apice sparse piloso, foliis sessilibus semiamplexicaulibus oblongis rigidis grosse inciso-serratis glabris, invol. squamis intimis confertè glutinosis, exter. laxiusculis, pappo oligosetato caducissimo, ligulis circ. 20 basi angustatis.  $\mathcal{U}$  in Californiâ legit et. Douglas. Caulis ruber. Folia circ. 3 poll. longa 8-9 lin. lata. (v. s. comm. ab. hon. Soc. hort. Lond.)
11. *G. SUPRATHALMOIDES*, fruticosa glabra, foliis oblongo-cuneatis obtusis serratis secus ramos floridos confertis, capitulis terminalibus solitariis sessilibus, invol. squamis acuminatis exterioribus amplis foliaceis, recept. demum convexo vix areolato, pappi setis circ. 10 corollæ disci longitudine.  $\mathcal{U}$  in Brasiliæ prov. Rio-Grande. Species patriâ et formâ distinctissima. Ligulæ amplæ flavæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1027 miss.)
12. *G.?* *CORONOPHOLIA* (Lehm. ind. sem. h. Hamb. 1828. p. 16. et in Linnæ 1830. p. 376 descr.), herbacea, foliis sessilibus crassiusculis linearibus pinnatifido-dentatis rugosis glabris pellucido-venosis, capitulis solitariis, invol. glutinoso-squamis ext. linearibus apice patentibus, pappo coronæformi membranaceo setaceo aristato.  $\mathcal{U}$  in terris Mexicanis. Capitula magn. Matricariæ Chamomillæ. Achenium obovatum compresso-quadrangulum. An ex pappi structuri omnino congener?
13. *G.?* *ANOMALA*, fruticosa glabra, foliis lineari-lanceolatis utriusque acuminatis acutè et remotè serratis subviscosis, capitulis ramos terminantibus solitariis discoideis, invol. squamis glutinosis in appendicem filiformem disco æqualem setaceo aristato.  $\mathcal{U}$  in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1031!). Folia 1  $\frac{1}{2}$  poll. longa, 2-3 lin. lata. (v. s. in h. Mus. reg. Par.)

† *Species dubiæ.*

*G. FRUTICOSA* (Desf. cat. h. par. ed. 2. p. 269), sub *Donia* indescripta et in editi 3<sup>a</sup> omissa.

*G. LAMBERTII* (G. Don in Lond. hort. brit. p. 350).  $\mathcal{U}$  in Mexico. Descripsi nomen non novi.

†† *Species exclusæ.*

*G. INCISA* Spreng. = *Calimeris incisa*.

*G. SIBIRICA* Link =? *Aster conspicuus* Fisch.

*DONIA CILIATA* Nutt. = *Aplopappus ciliatus*.

*DONIA CANARIENSIS* Less. = *Jasonia lævigata*.

Subdiv. 3. *HETEROTHECÆ.*

Pappus diversus in disco et in radio. Ligulæ uniseriales.

CLXXII. *HETEROTHECA* Cass. bull. philom. 1817. septemb. dict. 21. (1823: p. 130. 51. (1827) p. 460. — Calycium Ell. sketch 2. (1824) p. 339. non Pres. — *Diplocoma* Don in Sweet br. fl. gard. (1828) ser. 1. t. 246. Less. syn. 162 — *Doronici* sp. Cerv. Link.

Capitulum radiatum, fl. radii ligulatis foemineis 1-serialibus, disci tubulosi 5-dentati hermaphroditi. Rec. planum alveolatum. Invol. squamæ pauciseriales imbricatæ lineares adpressæ apice subpatulæ. Stigmata radii teretiuscula glabra, disci apice lanceolata hispida. Achenia radii oblonga lævia apice calva, disci cuneiformia villosa pappo aut pauciseta, pappo duplici, exter. brevissimo setoso, inter. longo setoso subscabro. — Herbæ Bor.-Americanae perennes (biennese?) erectæ paniculato-ramosæ villosæ aut pilosæ. Folia caulina sessilia integra rarius dentata. Flores flavi. Pappus fl. disci rufus, ext. brevis albidus.





1-serialis setiformis subscaber. — Herba Oronocensis erecta Polymniam ferè referens. Caulis glaber tetragonus. Folia opposita, petiolis basi latè auriculatis, auriculis in vaginam connatis, limbo latè ovato acuminato grossè inciso-dentato basi 5-7-nervio, suprà juniora adpressè villosa adulta glabra, subtus secus nervos adpressè pubera. Capitula in corymbum terminalem compositum densum disposita. Ligulæ ex sicco flavæ, cor. tubi colore intentiores. — Dixi ex nomine musæ Eratois alludens ad habitum Polymniæ affinem.

1. E. POLYMNIOIDES. *U* in Peruviam montanis Oronocensibus legit cl. Hænk. Folia petiolo 2-3 pollic. longo, limbo 5-6 poll. longo, 2-4 lato. (v. a. in h. Hænk. à cl. de Sternberg miss.)

## CLXXIV. WOODVILLEA.

Capitulum multi- (80-100) florum radiatum, ligulis circ. 3-serialibus linearibus fœmineis sterilibus disco et invol. duplò longioribus, fl. disci tubulosis 5-dent. hermaphr. Invol. campanulatum, squamis 2-3-serialibus lineari-oblongis æqualibus. Recept. epaleaceum. Ligulæ basi longè tubulosæ hispidæ. Antheræ ecaudatæ. Stylus radii longè bifidus glabriusculus, disci ramis brevibus crassiusculis, dorso puberis. Achænia villosa, radii linearia sterilia, disci oblonga compressa. Pappus setis pluriserialibus tenuissimis subscabris. — Herba Californica Calendulæ facie glanduloso-villosa. Caulis adscendens teres. Folia alterna oblonga basi attenuata sessilia obtusa integerrima subtiliter trivenia. Capitula solitaria terminalia magn. Calendulæ. Ligulæ verisim. flavæ. — Locus in tribu subdubius, artificiali normâ admissus. — Dixi in memoriam cl. Gullielmi Woodville auct. operis meritissimi cui titulus *Medical Botany*. (3 vol. 4°. 1790-1793.)

1. W. CALENDULACHA. — in Californiâ legit cl Douglas. Herba epithamææ. Folia 2 poll. longa, 6-7 lin. lata. Ligulæ 6 lin. longæ, lineam latæ. (v. a. comm. ab hon. Soc. hort. Londin.)

CLXXV PSIADIA Jacq. hort. Schœnbr. 2. p. 13 t. 152. Less. syn. 190. — Psiadia et Elphegea Cass. dict. 43. p. 503 et 14. p. 361. — Conyzæ et Baccharidis sp. Lam. — Thonaria Vent. herb. — Alix et Glutinaria Comm. herb.

Capitulum multiflorum, heterogamum, primâ fronte discoideum reverà radiatum, fl. radii pluriserialibus fœmineis ligulâ brevissimâ integrâ, disci tubulosis 5-dentatis hermaphroditis et ovarii abortu masculis paucis interdum subnullis. Recept. epaleatum subalveolatum. Invol. imbricatum pauciseriale squamis margine subscariosis. Achænia radii oblonga glabra erostria. Pappus conformis 1-serialis setosus scabriusculus. — Frutices Mascarenenses junioribus partibus glutinosi seu viscosi aut Madagascarienses non viscosi. Folia alterna petiolata oblonga grossè dentata aut integra. Corymbi compositi polycephali subaphylli.

## Sect. 1. GLUTINARIA Comm. mss.

Frutices Mauritanii glutinosi, foliis eglandulosis sæpius triplinerviis, ligulâ numerosis pluriserialibus brevissimis.

\* Foliis serratis aut dentatis.

1. P. GLUTINOSA (Jacq. h. schœnbr. 2. t. 152), ramis teretiusculis, foliis prælatâ lanceolatis utriusque acuminatis serratis penninerviis, squamis integerrimis.

is, capitulis 25-30-floris. ♂ in insulâ Mauritiî (Lam. Alt.), sed specim. laurit. nunquam vidi. *Solidago viscosa* Schrad. sert. hannov. 12. t. 6. *Conyza glutinosa* Lam. dict. 2. p. 89. *Erigeron viscosum* et nunc *Erigeron glutinosum* ort. Par. ex Desf.! cat. p. 174. Folia eglandulosa juniora valdè glutinosa, -4 poll. longa, pollicem lata petiolata. (v. v. c.)

♀. *BALSAMICA* (DC. in Wight contrib. bot. ind. p. 10), ramulis angulatis, his vix petiolatis basi cuneatis trinerviis integerrimis apice penninerviis serraturis grossis et paucis instructis, capitulis 25-30-floris. ♂ in insulis Mauritanis. Variat foliis lanceolatis glabris (*Baccharis arguta* Pers. ench. 2. p. 423. *Conyza Balsamica* Wall.! cat. n. 3068. comp. n. 178. *Eupatorium flexuosum* Aub. fl. ind. 3. p. 415), lanceolatis villosulo-glutinosi, ovali-lanceolatis subcortis (*Elphegea crenata* Cass. dict. 14. p. 362?) et obovatis subretusis (*Elphegea latifolia* Cass.! dict. 14. p. 362). Folia et ramuli viscidii. Similis Psiad. glutinosæ nec certissimè ab eâ separanda. Vulgò in hortis *Baume africain*. (v. s. sp.)

♀. *RETUSA* (Lam. dict. 2. p. 90. sub *Conyzâ*), suffruticosa pilis sparsis subirsutula, foliis longè cuneatis retusis infernè integerrimis triplinerviis apice ematis, pedunculis axillaribus terminalibusque folio paulò brevioribus corymbum 5-7-cephalum gerentibus, invol. globosi squamis subglabris margine utè membranaceis obtusis. ♂ in insulâ Borboniæ maritimis (Comm.! Bontou!). Folia sapore salino donata et aceto ad usum culinarem præparata. Alix salicilla Comm. mss. et ic. Vulg. *la saliette*, *la bien salée*. Folia 3 poll. longa, 1 lin. lata. (v. s.)

♀. *DENTATA*, ramis teretiusculis, foliis breviter petiolatis coriaceis linearimucronatis basi cuneatis integerrimis et subtrinerviis apice serraturis utrinquè -5 notatis, capitulis multifloris. ♂ in montanis insulæ Borboniæ (Bontou!). *Elphegea dentata* Cass.! dict. 14. p. 363. Folia 1-3 pollic. longa, 3-5 lin. lata viscosa. (v. s.)

\*\* *Foliis integerrimis.*

♀. *LINEARIFOLIA*, foliis petiolatis lineari-lanceolatis utrinquè acuminatis triplinerviis integerrimis, capitulis 10-12-floris. ♂ in insulis Mauritanis, nempe in ins. Borboniæ (Comm.), Mauritiî (Bontou in h. Lindl.). *Conyza linearifolia* Pers. ench. 2. p. 92. *Baccharis linearifolia* Pers. ench. 2. p. 425. *Elphegea mucrolata* Cass.! dict. 14. p. 562. Species distinctissima evidenter duabus rioribus affinis et caract. Psiadiæ exhibens. Ligulæ paulò majores quàm in rioribus. Flores flavi. Folia et ramuli glutinosi. Folia cum petiolo 3 pollicaria, 4-5 lin. lata. Sequentis var. videtur ex Bojer obs. in litt. (v. s. in h. Indley.)

♀. *INTEGERRIMA* (DC. in Wight contrib. p. 10), foliis distinctè petiolatis obovatis mucronatis integerrimis subtripplinerviis, capitulis 8-12-floris. in montanis insulæ Mauritiî et culta? in Indiâ orientali (Wight) et Moluccis (D'Urv.) Ligulæ albidæ paulò majores quàm prioria. Fl. tubulosi paucissimi imò nulli, undè stirps subdioica evadit. Petioli circ. semipollicares.

a. *minor*, foliis triplinerviis, nervo medio vix ac ne vix nervulos pinnatos gerente. *Baccharis viscosa* Lam. dict. 1. p. 345. journ. phys. 1. p. 62. t. 3. *Elphegea minor* Cass. dict. 14. p. 362. (v. s.)

β. *trinervia*, foliis triplinerviis nervo medio penninervio. Psiadia trinervia Willd. mag. naturf. ges. Berl. 1807. p. 139 huc pertinere videtur etiam si folia dicantur opposita cum in nostrâ evidenter alterna.

γ? *quinquenervia*, foliis elliptico-lanceolatis integerrimis 5-nerviis, corymbis densis, capitalis multifloris. ♂ in ins. Mauritanis. *Elphegea quinquenervia* Cass. dict. 14. p. 363 huc relata ex auct. cl. Bojer in litt. sed mihi ignota.

♀. *PENNINERVIA*, foliis distinctè petiolatis ellipticis integerrimis penninerviis, capitulis 10-12-floris. ♂ in montibus ins. Mauritiî legit cl. Bojer. Psiadia glutinosa Bojer! in litt. Differt à Psiadiâ glutinosâ Jacq. et à divis. prim. foliis in-

tegerrimis nec serratis, à P. integerrimâ et linearifoliâ foliis penninerviis nec tri aut triplinerviis (v. s. comm. à cl. Bojer.)

### Sect. II. GLANDULARIA DC. non Gmel.

Species Madagascariensis, herbacea? non viscosa, foliis sessilibus penniveniis subtus minutè glanduliferis, pedunculis axillaribus, floribus purpurascens, ligulis paucis oblongis 1-serialibus.

8. *P. MADAGASCARIENSIS*, foliis sessilibus oblongo-aut ovali-lanceolatis acuminatis subintegerrimis penniveniis glaberrimis subtus sub lente glandulosis, pedunculis axillaribus apice 8-10-cephalis folio circ. longitudine aequalibus in paniculam interruptam digestis.  $\mathcal{U}$  in insulâ Madagascar. *Conyza Madagascariensis* Lam. dict. 2. p. 85. *Baccharis Madagascariensis* Pers. ench. 2. p. 423. Folia imâ basi angustè semiamplexicaulia ex sicco non glutinosa. Invol. squamæ glaberrimæ margine membranaceæ. Ligulæ oblongæ. Pappus rufescens. (v. s.)

### CLXXVI. MICROGLOSSA. — Conyza sp. auct.

Capitulum multiflorum minutissimè radiatum, flor. radii pluriseriis ligulatis foemineis, ligulis linearibus pappo brevioribus, fl. disci paucis tubulosis 3-5-fidis abortu masculis. Recept. nudum. Invol. imbricatum, squamis lanceolato-linearibus subscariosis erectis. Anthera basi ecaudatæ. Achænia oblonga compressa? glabra. Pappus pilosus 1-serialis. — Suffrutices Indici aut Africani, erecti aut subvolubiles glabriusculi apice ramosi quasi proliferi. Folia petiolata ovata acuminata integra aut denticulis exsertis acutis exsertè subserrata subtus pubescentia et glandulosa adulta subglabrata. Corymbi aphylli fastigiati divaricati terminales. Pappus rufescens. — Differt à Conyzâ fl. radii non tubulosis sed brevissimè ligulatis, ab Erigeronte floribus disci masculis paucis et habitu toto.

\* Volubiles.

1. *M. VOLUBILIS*, caule volubili, foliis petiolatis ovatis acuminatis, ligulis stylo suo ferè brevioribus, foliis ramulisque glabriusculis.  $\mathcal{J}$  in Indiæ orient. montanis humidis, in insulis Javâ (Blum!) et Penang (Wall!). Contra prolifera Blum! bijdr. 896. *Conyza volubilis* Wall! cat. et herb. n. 303. comp. n. 167. *Sonchus volubilis* Rumph. amb. 5. t. 104. f. 1 (v. s.)

$\beta$ . *ambigua*, ligulis stylo suo longioribus, ramulis foliisque subtus brevissimè pubescenti-hirtellis. — in Sierra-Leonâ? Simillima speciminibus prioris à Wallichio in ins. Penang lectis, sed folia subtus densius velutis et confertius glandulosa. An certè diversa? (v. s. ex h. L'Hér.)

$\gamma$ . *Madagascariensis*, ligulis stylo suo brevioribus, ramulis foliisque subtus brevissimè pubescenti-hirtellis. — in insul. Madagascar ad margins riv. varum legit cl. Bojer. (v. s.)

*N. B.* An fortè *Conyza pyrifolia* Lam. dict. 2. p. 89 etiam ex ins. Javâ ortu esset hujus generis altera species? *Conyza prolifera* Lam. videtur diversissima et Decaisn. novv. ann. mus. p. 408.

2. *M. ASSLIFOLIA*, volubilis glabra, ramis teretibus subherbaceis, foliis sessilibus basi obtusis ellipticis acuminatis integerrimis, ligulis stylo suo circ. aequalibus.  $\mathcal{J}$  in insulæ Madagascar (Forb! in Lindl! herb.) sylvis super arbore scandens (Bojer.). Folia 2  $\frac{1}{2}$  poll. longa, pollicem lata. Rami super. axillares et terminales subpaniculati. Invol. squamæ acutæ. Flores lutei. Pappus rufescens. (v. s. comm. à cl. Bojer.)

\*\* Non volubiles.

3. *M. ALTISSIMA*, arborea glabra viscosa, ramis confertis flexilibus subangulatis, foliis sessilibus lanceolatis basi attenuatis, inf. serratis, super. integerrimis.

- corymbis compositis irregulariter subfastigiatis, invol. squamis obtusis, ligulis stylo suo æqualibus. ♀ in insulæ Madagascar prov. Emirnenis legit cl. Bojer. Baccharis altissima Bojer! in litt. 1835. Arbor 25-30-pedalis speciosa. Folia super. 2 poll. longa, 4 lin. lata, cum ramulis glutine viscoso illinita. Flores lateri. (v. s. comm. à cl. Bojer.)
4. *M. HISPIDA*, fruticosa, ramis erectis teretibus subherbaceis hispidis, foliis ovato-oblongis grossè dentatis hispidis in petiolum alatum basi obtusè auritum attenuatis, corymbo composito laxo, invol. squamis acuminatis, recept. convexo alveolato, ligulis stylo suo æqualibus. ♀ in insulæ Madagascar prov. Tananarivou circa urbem ad fossarum margines legit cl. Bojer. Frutex 4-5-pedalis. (v. s.)
5. *M. PETIOLARIS*, fruticosa, ramis teretibus anbtiter striatulis et puberulis, foliis petiolatis ovatis acuminatis subintegerrimis supra secus nervos subpuberulis subius pallidis pubescentibus, corymbo composito fastigiato, invol. squamis glabris linearibus subacutis, ligulis stylo subæqualibus. ♀ in Sierrâ-Leonâ legit cl. Don. Petioli 6 lin. longi. Folia 2 poll. longa, 10 lin. lata. (v. s. in h. Lindl.)

CLXXXVII. NIDORELLA Cass. dict. 37. p. 459 et 469. v. 56. p. 166. — Erigerontis, Conyzæ, etc., sp. auct.

Capitulum multiflorum heterogamum, fl. homochromis, radii fœmineis minutissimè ligulatis uni- bi- aut pluriseriatis, disci circ. 20 tubulosi 5-fidis bisexualibus, centralibus interdum sterilibus. Invol. campanulatum imbricatum. Recept. alveolatum. Anth. ecaudatæ. Achænia oblonga teretiuscula aut subcompressa, sæpius minutè puberula. Pappus 1-serialis, setis subscabridis basi in anulum subconcretis. — Herbæ aut suffrutices Australi-Africana, et una ex Arabiâ. Caulis erecti. Folia alterna. Capitula corymbosa aut rariùs solitaria. Corollæ omnes flavæ, ligulis interdum disco pallidioribus.

### §. 1. Genuinæ, corymbis polycephalis compositis.

\* *Foliis incisis dentatisæ.*

1. *N. MESPILOFOLIA*, suffruticosa ramosa glabriuscula, foliis petiolatis ovatis sæu cordatis profundè et acutè inciso-dentatis, capitulis corymbosis. ♀ in Africâ australi inter dumeta et sylvas, in distr. Uitenhage, Stellenbosch et Albany (Eckl. et Zeyh.), ad Zneenreberge et Omsanculo (Drege!). Aster mespilifolius Less. syn. 180. Eupatorium Eckl. l Rami teretes. Folia aut glabra aut pabe minutissimè scabrida. Corymbi compositi 15-20-cephali. Invol. squamæ oblongo-lineares imbricatæ. Recept. foveolatum. Cor. omnes pallidè flavæ, radii ligulatæ stylo longiores, disci fertiles. Achænia pube minutissimè scabrida. (v. s.)
2. *N. PINNATILOBATA*, basi suffruticosa subsimplex tota puberulo-hirtella, foliis in petiolum attenuatis pinnatilobatis, lobis utrinquè 2-4 lanceolatis acutis, corymbo composito polycephalo. ♀ in Africâ austr.-orient. ad Zw. Key et Basche legit cl. Drege! Folia pollicem longa, inf. pinnatifida, sup. pinnatipartita lobis angustioribus. Invol. squamæ vix acutæ dorso subscabræ margine angustè scariosæ. Ligulæ minimæ obovatæ stylo longiores. Achænia teretiuscula pubescentia. (v. s.)
3. *N. TRILOBA*, rhizomate suffruticoso, caulibus ramisque erectis foliisque pubescentibus, foliis 3-5-lobatis basi in petiolum angustatis, lobis oblongis terminali (præs. in fol. infer.) majore obtusiore, corymbo subpaniculato laxo, invol. squamis puberis oblongis acutis ext. dorso puberis, int. subscariosis ciliatis, omnibus demum reflexis ♀ 2 l. in desertis circa montem Sinai legit cl. Bove (n. 101!), mens. junio flor. Arabis dicitur *Nefflé*. An Erigeron incisus Sieb. pal. à Spreng. citat.? Conyza triloba Decaisn. ann. scienc. nat. 2. p. 267. Herba

*semipedalis pallida, invol. substraminea. Cor. flava. Ligulæ brevissima vix perspicue transeunt gradatim in tabos, nudæ species media inter Trimorphum et Nidorellam. Pappi setæ albae basi in annulum ab echenio separabilem concretæ. (v. s. comm. à cl. inv.)*

4. *N. CHRYSOCOMA*, herbacea, ramis teretibus virgatis villosis, foliis ramulis basi obtusè aurito-subordatis lineari-oblongis grossè serratis aut subincisè pubescentibus, corymbo composito polycephalo, invol. squamis dorso puberulis apice acuminatis. ① in ins. Madagascar ad agros cultos legit cl. Bojer. Chrysocoma aurea Bojer! in litt. 1835. Herba strictè erecta speciosa 2-3 pedes. Flores aurei, ligulæ minutissimæ. Affinis *N. pinnatilobata* sed herbacea, foliis vix incisa potius dentata, invol. squamæ multò magis acuminatæ. (v. s. comm. à cl. inv.)
5. *N. RESERIFOLIA*, herbacea erecta, caule foliorumque petiolis parè strigillosis, foliis subpuberulis in petiolum longè attenuatis pinnatipartitis, lobis 1-2-jugis oblongis, terminali obovato-oblongo paulò majore, omnibus mucronatis, corymbo composito, invol. squamis elliptico-oblongis obtusiusculis subciliatis. — in Africæ australis regione Gariepina, Klein Namaqualand legit cl. Drege! Ligulæ minutæ pallidius quàm tubuli flavae ovals subcutæ. Achenis gibbriuscula. (v. s.)
6. *N. HIRTA*, herbacea subimplex erectiuscula, caule strigilloso-hirto, foliis hirtello-scabris, infer. oblongis in petiolum longè attenuatis integris aut his inde lobulatis, cæteris basi dilatato-amplexicanibus pinnatipartitis, lobis 1-2-jugis oblongo-linearibus, corymbo composito conferto, invol. squamis linearibus acutis dorso subhirtellis. — in Africæ australi ad Stormberg alt. 5-600 ped legit cl. Drege! Affinis *N. resedifolia*. Cor. etiam radii saturatius flava. Ligulæ minimæ. Folia in axillis sæpè fasciculata. (v. s.)
7. *N. AURICULATA*, herbacea erecta subimplex, caule tereti confertè hirtello, foliis basi rotundè auriculatis pube minimè pubescenti-scabris, infer. petiolis latè ovatis grossè dentatis, super. basi suprè auriculas attenuatis oblongis integris, corymbo composito fastigiato, capitulis congestis, invol. squamis obtusiusculis subciliatis dorso subhirtellis. — in Africæ austr.-orient. ad Zw. Omtz et Omsamwabo legit cl. Drege! Folia inf. petiolo incl. 4 poll. longa, ferè poll. lata. Corymbus ferè *N. fetida*. Ligulæ minimæ flavae. (v. s.)
8. *N. OBOVATA*, herbacea erecta tota brevè et confertè hirtella, foliis obovatis basi cuneato-attenuatis subsessilibus grossè et obtusè dentatis reticulato-serratis confertis, corymbo composito polycephalo, capitulis confertis, invol. squamis oblongis ciliolatis obtusis. — in Africæ austr. distr. Kafferland legit cl. Drege! Folia sesquipollicem longa, 9-10 lin. lata, summa oblonga integra Ligulæ minimæ obtusæ flavae. (v. s.)
9. *N. SENECEIOMEA*, herbacea erecta tota minutè puberulo-hirtella, foliis oblongis infernè attenuatis imò basi vix auriculatis, infer. grossè paucidentatis, super. integris, corymbo composito fastigiato, invol. squamis oblongis subcutè dorso puberis et subciliatis. ② in Africæ Capensis distr. Uitenhage et Caledon legit cl. Ecklon! et Zeyher. Ligulæ minutæ flavae. Folia sesquipollic. long. 4 lin. lata. (v. s.)
10. *N. ALBANENSIS*, foliis obovatis magis dentatis in axillis confertis. — in distr. Albany. Eckl! (v. s.)
10. *N. LINIFOLIA*, herbacea erecta simplex glabra, foliis lineari-lanceolatis acuminatis, infer. basi longè attenuatis grossè serratis, super. sessilibus integerrimis, corymbo composito, invol. squamis acutis dorso glabris margine subciliatis. ①? in Africæ austr.-orient. prope Port-Natal legit cl. Drege! Folia infer. 3 pollic. longa, 3 lin. lata. Ligulæ minimæ obtusæ. (v. s.)

\*\* *Foliis integerrimis.*

11. *N. SOLIDAGINEA*, collo sublignoso, caulibus herbaceis erectis subimplex infernè sparè setosis supernè minutè et confertè puberulis, foliis linearibus

integris acutis minutè puberulis basi subciliatis, infer. basi attenuatis, super. basi subdilatis, corymbo composito, pedicellis involucrique minutè puberulis, ligulis angustis acutis invol. paulò longioribus.  $\mathcal{U}$  in Africâ australi ad Witthergen et Stormberg alt. 4-500 ped. legit cl. Drege. Capitula in genere majora nempe 3-4 lin. diam. Folia 2 poll. ferè longa, 2 lin. lata. (v. s.)

12. *N. HORRENTORICA*, caule basi sublignoso erecto villosulo-sublanuginoso, foliis integerrimis subtus laxè lanatis suprâ villosis ant subglabris, infimis obovatis in petiolum attenuatis mediis oblongo-cuneatis obtusis basi angustis, summis oblongis basi latiusculis apice mucronatis, corymbo composito.  $\mathcal{Z}$  ad Cap. Bonæ-Spei, longè ab urbe in regione Transgariepina inter Klaarwater et Litakun prope Kosi-Fountain legit cl. Burchell et sub n. 2528. cat. geogr. mecum comm. Capitula omninò E. foetidi sed habitu et lanugine valde differt. (v. s.)
13. *N. POLYCEPHALA*, herbacea erecta paniculato-ramosissima tota pube brevi cinerascens-pubescent, foliis caulinis auriculato-semiamplexicaulis elongatis lineari-lanceolatis triplinerviis, ramis linearibus uninerviis, omnibus integerrimis acuminatis, corymbo cujusque rami composito congestifloro, invol. squamis acutis subhirtellis. — in Africâ australi legit cl. Drege! Herba bipedalis. Folia caul. 3 poll. longa 3 lin. lata, ramea pollicaria lineam lata. Ligulæ parvæ discoloriores parvæ obtusæ subtridentatæ. (v. s.)
14. *N. MUCRONATA*, caule herbaceo tereti striato villosulo, foliis integerrimis lineari-oblongis mucronatis basi obtusè subauritis sessilibus glabriusculis 1-nerviis, corymbo composito congestifloro, invol. squamis obtusiusculis disco dimidiò brevioribus. ( $\mathcal{U}$ ) in insulâ Madagascar savannis ad radices montis Autoungun legit cl. Bojer. Chrysocoma lappulina Bojer! in litt. 1835. Affinis *N. rapunculoidi*. (v. s. comm. à cl. Bojer.)
15. *N. RAPUNCULOIDES*, caule herbaceo erecto simplici infernè setis hispido superne glabriusculo aut minutissimè velutino, foliis elongatis lineari-cuneatis obtusis basi longè attenuatis integerrimis minutè mucronatis basi ciliatis cæst. sparè puberulis, corymbo composito, invol. squamis ovatis obtusiusculis, ligulis obtusis integris minimis.  $\mathcal{U}$ ? in Africâ australi legit cl. Drege. Baccharis rapunculoides E. Mey.! in Dreg. coll. Radix obliqua subsimplex. Caulis bipedalis. Capitula parva. Ligulæ disco pallidiores. Folia 2  $\frac{1}{2}$  poll. longa 4 lin. lata. (v. s.)
16. *N. HYSSOPIFOLIA*, caule suffruticuloso ramoso tereti plus minùs hirtello, foliis lineari-cuneatis integerrimis obtusis basi attenuatis minutissimis pubescentibus, corymbis compositis, invol. squamis oblongis acutis dorso subhirtellis, ligulis minimis obtusis subcucullatis.  $\mathcal{Z}$  in Africâ Capensi (Drege!) ad Albany, Uitenhage, territ. cessum et circa urbem (Eckl.! et Zeyh.). Baccharis hyssopifolia E. Mey.! in Dreg. coll. Valde accedit ad *N. foetidam* sed folia remotiora, capitula minora, etc. (v. s.)  
*f. glabrata*, caule folisque glabris. — in Uitenhage. Eckl.! (v. s.)
17. *N. FOETIDA*, caule herbaceo erecto simplici pilis hispido, foliis confertis linearibus basi cuneato-attenuatis obtusis integerrimis submucronatis utrinquè subscabridis, axillis ferè omnibus foliiferis, corymbis pluribus compositis pedunculatis confertis, invol. squamis linearibus acutis dorso subhirtellis, ligulis minimis obtusis.  $\mathcal{U}$  in Africâ (Pluk. Herm.), ad Cap. Bonæ-Spei (Th. L'Hér.! herb. Burm.! herb. Eckl.!). Erigeron foetidum Linn. sp. 1213. Less. syn. 190. Inula foetida Linn. sp. 1241 (ex Bocc. syn.), Willd. sp. 3. p. 2105. Thunb. fl. cap. 667. Burm.! prod. 26. Solidago retusa Burm.! prod. 27. Nidorella foliosa Cass. dict. 37. p. 469. — Mill. dict. t. 233. Pluk. t. 223. f. 3. Herm. lugd. b. t. 662. Stirps Melitensis Bocconi huc falsò à pluribus relata est Jasonia glutinosa. (v. s.)
18. *N. PUNCTULATA*, glaberrima, caule fruticoso, ramis viscosis, foliis in petiolum attenuatis elongato-lanceolatis acuminatis integerrimis utrinquè punctis glandulos minimissimis notatis, corymbo composito, pedunculis compressis, involucrio pappo et etiam disco breviorè, squamis obtusiusculis.  $\mathcal{Z}$  ad Cap.

semipedalis pallida, invol. substraminea. Cor. flavidæ. Ligulæ brevissima vix perspicua; transeunt gradatim in tubos, nudæ species media inter Trimorphazam et Nidorellam. Pappi setæ albæ basi in annulum ab achænio separabilem coarctatæ. (v. s. comm. à cl. inv.)

4. *N. CHRYSOCOMA*, herbacea, ramis teretibus virgatis villosis, foliis rameis basi obtusè aurito-subcordatis lineari-oblongis grossè serratis aut subincisus pubescentibus, corymbo composito polycephalo, invol. squamis dorso puberulis apice acuminatis. ① in ins. Madagascar ad agros cultos legit cl. Bojer. Chrysocoma aurea Bojer! in litt. 1835. Herba strictè erecta speciosa 2-3 pedalis. Flores aurei, ligulæ minutissimæ. Affinis *N. pinnatilobata* sed herbacea, folia vix incisa potius dentata, invol. squamæ multò magis acuminatæ. (v. s. comm. à cl. inv.)
  5. *N. RESIDUOLIA*, herbacea erecta, caule foliorumque petiolis parçè strigillosis, foliis subpuberulis in petiolum longè attenuatis pinnatipartitis, lobis 1-2-jugis oblongis, terminali obovato-oblongo paulò majore, omnibus mucronatis, corymbo composito, invol. squamis elliptico-oblongis obtusiusculis subciliatis. — in Africæ australis regione Gariëpinâ, Klein Namaqualand legit cl. Drege! Ligulæ minutæ pallidius quàm tubuli flavidæ ovales subacutæ. Achænia glabriscula. (v. s.)
  6. *N. HIRTA*, herbacea subsimplex erectiuscula, caule strigilloso-hirto, foliis hirtello-scabris, infer. oblongis in petiolum longè attenuatis integris aut hinc indè lobulatis, cæteris basi dilatato-amplexicaulibus pinnatipartitis, lobis 1-2-jugis oblongo-linearibus, corymbo composito conferto, invol. squamis linearibus acutis dorso subhirtellis. — in Africâ australi ad Stormberg alt. 5-600 ped. legit cl. Drege! Affinis *N. residuoliz*. Cor. etiam radii saturatius flavæ. Ligulæ minimæ. Folia in axillis sæpè fasciculata. (v. s.)
  7. *N. AURICULATA*, herbacea erecta subsimplex, caule tereti confertè hirtello, foliis basi rotundè auriculatis pube minimâ pubescenti-scabris, infer. petiolatis latè ovatis grossè dentatis, super. basi suprâ auriculari attenuatis oblongis integris, corymbo composito fastigiato, capitulis congestis, invol. squamis obtusiusculis subciliatis dorso subhirtellis. — in Africâ austr.-orient. ad Zw. Omtata et Omsamwabo legit cl. Drege! Folia inf. petiolo incl. 4 poll. longa, ferè 7 poll. lata. Corymbus ferè *N. foetida*. Ligulæ minimæ flavæ. (v. s.)
  8. *N. OBOVATA*, herbacea erecta tota brevè et confertè hirtella, foliis obovatis basi cuneato-attenuatis subsessilibus grossè et obtusè dentatis reticulato-nervosis confertis, corymbo composito polycephalo, capitulis confertis, invol. squamis oblongis ciliatis obtusis. — in Africâ austr. distr. Kafferland legit cl. Drege! Folia sesquipollicem longa, 9-10 lin. lata, summa oblonga integra Ligulæ minimæ obtusæ flavæ. (v. s.)
  9. *N. AURICIONA*, herbacea erecta tota minutè puberulo-hirtella, foliis oblongis infernè attenuatis imâ basi vix auriculatis, infer. grossè paucidentatis, super. integris, corymbo composito fastigiato, invol. squamis oblongis subacutis dorso puberis et subciliatis. 2L in Africâ Capensis distr. Uitenhage et Caledon legit cl. Ecklon! et Zeyher. Ligulæ minutæ flavæ. Folia sesquipollic. longa, 4 lin. lata. (v. s.)
- ß. *Albanensis*, foliis obovatis magis dentatis in axillis confertis. — in distr. Albany. Eckl. (v. s.)
10. *N. LINIFOLIA*, herbacea erecta simplex glabra, foliis lineari-lanceolatis acuminatis, infer. basi longè attenuatis grossè serratis, super. semilibus integerrimis, corymbo composito, invol. squamis acutis dorso glabris margine subciliatis. ①? in Africâ austr.-orient. prope Port-Natal legit cl. Drege! Folia infer. 3 pollic. longa, 3 lin. lata. Ligulæ minimæ obtusæ. (v. s.)

\*\* *Foliis integerrimis.*

11. *N. SOLIDAGINEA*, collo sublignoso, caulibus herbaceis erectis subsimplicibus infernè sparè setosis supernè minutè et confertè puberulis, foliis linearibus

- integris acutis minutè puberulis basi subciliatis, infer. basi attenuatis, super. basi subdilatatis, corymbo composito, pedicellis involucrique minutè puberulis, ligulis angustis acutis invol. paulò longioribus.  $\mathcal{L}$  in Africâ australi ad Wittbergen et Stormberg alt. 4-500 ped. legit cl. Drege. Capitula in genere majora nempe 3-4 lin. diam. Folia 2 poll. ferè longa, 2 lin. lata. (v. s.)
12. *N. HORTENTOTICA*, caule basi sublignoso erecto villosulo-sublanuginoso, foliis integerrimis subtus laxè lanatis suprâ villosis ant subglabris, infimis obovatis in petiolum attenuatis mediis oblongo-cuneatis obtusis basi angustis, summis oblongis basi latiusculis apice mucronatis, corymbo composito.  $\mathcal{J}$  ad Cap. Bonæ-Spei, longè ab urbe in regione Transgariepina inter Klaarwater et Litakun prope Kosi-Fountain legit cl. Burchell et sub n. 2528. cat. geogr. mecum comm. Capitula omnino E. foetidi sed habitu et lanugine valdè differt. (v. s.)
13. *N. POLYCEPHALA*, herbacea erecta paniculato-ramosissima tota pube brevi cinerascens-pubescentis, foliis caulinis auriculato-semiamplexicaulis elongatis lineari-lanceolatis triplinerviis, rameis linearibus uninerviis, omnibus integerrimis acuminatis, corymbo cujusque rami composito congestifloro, invol. squamis acutis subhirtellis. — in Africâ australi legit cl. Drege! Herba bipedalis. Folia caul. 3 poll. longa 3 lin. lata, ramea pollicaria lineam lata. Ligulæ parvæ pallidiores parvæ obtusæ subtridentatæ. (v. s.)
14. *N. MUCRONATA*, caule herbaceo tereti striato villosulo, foliis integerrimis lineari-oblongis mucronatis basi obtusè subauritis sessilibus glabriusculis 1-nerviis, corymbo composito congestifloro, invol. squamis obtusiusculis disco dimidiò brevioribus. ( $\mathcal{L}$ ) in insulæ Madagascar savennis ad radices montis Antonngun legit cl. Bojer. Chrysocoma lappulina Bojer! in litt. 1835. Affinis *N. rapunculoidi*. (v. s. comm. à cl. Bojer.)
15. *N. RAPUNCULOIDES*, caule herbaceo erecto simplici infernè setis hispido superè glabriusculo ant minutissimè velutino, foliis elongatis lineari-cuneatis obtusis basi longè attenuatis integerrimis minutè mucronulatis basi ciliatis cat. sparsè puberulis, corymbo composito, invol. squamis ovatis obtusiusculis, ligulis obtusis integris minimis.  $\mathcal{L}$ ? in Africâ australi legit cl. Drege. Baccharis rapunculoides E. Mey.! in Dreg. coll. Radix obliqua subsimplex. Caulis bipedalis. Capitula parva. Ligulæ disco pallidiores. Folia 2  $\frac{1}{2}$  poll. longa 4 lin. lata. (v. s.)
16. *N. HYSSOPIFOLIA*, caule suffruticuloso ramoso tereti plus minus hirtello, foliis lineari-cuneatis integerrimis obtusis basi attenuatis minutissimis pubescentibus, corymbis compositis, invol. squamis oblongis acutis dorso subhirtellis, ligulis minimis obtusis subcucullatis.  $\mathcal{J}$  in Africâ Capens! (Drege!) ad Albany, Uitenhage, terit. cessum et circa urbem (Eckl.! et Zeyh.). Baccharis hyssopifolia E. Mey.! in Dreg. coll. Valdè accedit ad *N. foetidam* sed folia remotiora, capitula minora, etc. (v. s.)
- $\beta$ . glabrata*, caule foliisque glabris. — in Uitenhage. Eckl.! (v. s.)
17. *N. FOETIDA*, caule herbaceo erecto simplici pilis hispido, foliis confertis linearibus basi cuneato-attenuatis obtusis integerrimis submucronatis utrinque subcabridis, axillis ferè omnibus foliiferis, corymbis pluribus compositis pedunculatis confertis, invol. squamis linearibus acutis dorso subhirtellis, ligulis minimis obtusis.  $\mathcal{L}$  in Africâ (Pluk. Herm.), ad Cap. Bonæ-Spei (Th. L'Hér.! herb. Burm.! herb. Eckl.!). Erigeron foetidum Linn. sp. 1213. Less. syn. 190. Inula foetida Linn. sp. 1241 (ex Boccon. syn.). Willd. sp. 3. p. 2105. Thunb. fl. cap. 667. Burm.! prod. 26. Solidago retusa Burm.! prod. 27. Nidorella foliosa Cass. dict. 37. p. 469. — Mill. dict. t. 233. Pluk. t. 223. f. 3. Herm. lugd. b. t. 662. Stirps Melitensis Bocconi huc falsò à pluribus relata est Jasonia glutinosa. (v. s.)
18. *N. PUNCTULATA*, glaberrima, caule fruticoso, ramis viscosis, foliis in petiolum attenuatis elongato-lanceolatis acuminatis integerrimis utrinquè punctis glandulosis minutissimis notatis, corymbo composito, pedunculis compressis, involucrio pappo et etiam disco breviorè, squamis obtusiusculis.  $\mathcal{J}$  ad Cap.



Bonæ-Spei in regione Transgariëpinâ inter montes Asbestinos et Witte-water legit cl. Burchell (cat. geog. n. 2061). Species distinctissima. Folia 3-4 poll. longa, 3-5 lin. lata, crassiuscula. Flores radii feminei brevissimè ligulati. (v. s.)

19. *N. AMPLEXICAULIS*, caule herbaceo glabro simplici apice corymboso, foliis oblongo-ligulatis basi obtusè auriculato-cordatis breviter decurrentibus apice subacutis hinc inde subdenticulatis glabris, corymbi compositi ramis pubescentibus. 2½ ad Cap. Bonæ-Spei ad orientem urbis legit cl. Burchell et necum sub n. 7135. cat. geogr. comm. Herba bipedalis. Folia radicalia basi subangustata, caulina infer. longius decurrentia, 4 poll. longa, 6-8 lin. lata, summa vix decurrentia acuminata. Capitula parva. (v. s.)
20. *N. LONGIFOLIA*, herbacea erecta glaberrima, caule ad collum subadscendente apice angulato, foliis integerrimis radicalibus longissimis oblongis obtusis basi longè attenuatis, caulinis sessilibus semiamplexicanibus acutis, corymbo composito polycephalo, invol. squamis acutis, ligulis minimis. 2 in Africâ australi ad Zneenrebergen legit cl. Drege! et fortè in Kafferland d. Ecklon et Zeyher. Herba ferè bipedalis, similis *N. amplexicauli*, sed folia radicalia pedem et ultra longa, 10-13 lin. lata. (v. s.)

§. 2. *Spuriæ, capitulis solitariis terminalibus, foliis sessilibus linearibus integerrimis hirtis.*

21. *N. EXILIS*, caule herbaceo basi suffruticoso erecto exili apice subnudo, ramis paucis elongatis 1-cephalis, foliis linearibus subacutis integerrimis, omnibus ramisque pube minutâ subglanduliformi puberulis, inferioribus majoribus setas longas patulas sparsas insuper gerentibus, capitulis longè pediculis solitariis, invol. squamis subimbricatis planis linearibus acutis disco aequalibus, lignis 8-10 in sicco revolutis, acheniis villosis. 3 et fortè 1 in Africâ Capensi ad Clam-Williams leg. cl. Ecklon et Zeyh. Achenia disci in specim. meo pleraque abortiva. Stylorum disci rami conici undiquè hirtelli. Pappus 1 serialis plumosus. Habitus Senecionis puberuli sed genericè diversissima. Flores disci et radii flavi isochromi. (v. s.)
22. *N. ? GARIËPINA*, caule fruticoso erecto divaricato-ramoso tereti hirtello, foliis linearibus integerrimis utrinquè hirtis, ramis floridis apice breviter densatis, invol. squamis linearibus acuminatis dorso hirtis apice ad marginem scariosis ciliatis, acheniis vix puberis, ligulis minimis. 3 in Africâ Capensi regione Gariëpinâ locis asperis legit cl. Drege. Recept. omnino nudum, lvs biserialæ demum reflexum. Folia 6 lin. longa, semilineam lata. (v. s.)

CLXXXVIII. HOMOCHROMA.

Capitulum multiflorum, fl. radii ligulatis 1-seriatis homochromis. disci tubulosi 5-fidis bisexualibus sed sterilibus. Invol. 2-5-seriale. squamis linearibus acuminatis adpressis. Recept. subconvexum foveolatum. Anth. basi ecaudatæ. Stylus Asteris. Achenia radii plano-compressa obovata villosa, disci filiformia glabra. Pappus 1-serialis setosus. setis radii circ. 15, disci circ. 10, omnibus præsertim ad apicem plumosis. — Suffrutex Capensis ramosus hirtellus, ramis foliosis apice nudis 1-cephalis glanduloso-subhirtellis. Folia alterna sessilia oblongo-inæria integerrima hirtella et ciliata. Capitula flava ligulis majoribus quàm in Nidorellâ.

1. *H. ECKLONIS*. 3 in Africâ Capensi ad Clam-Williams leg. cl. Ecklon et Zeyher. Folia 6 lin. longa, lineam aut sesquilineam lata. (v. s.)

*NEJA D. Don in Sweet brit. fl. gard. ser. 2. n. et t. 78. — Diplopappi sp. Less.*

luna multiflorum heterogamum, fl. radii 1-3-serialibus angustè æmineis, disci 5-dentatis hermaphrod. Recept. planum favu-  
vol. latè hemisphæricum imbricatum 3-seriale, squamis lineari-  
Cor. extus papilloso-pilosæ, stigm. radii obtusa lavia, disci  
cuta. Achænium lineari-oblongum compressum sericeo-villo-  
pus duplex, ext. brevis angustè paleaceus, interior longior  
aber. — Suffrutices Americani, pilis longis sparsis patulis  
lami foliosi aut apice nudi 1-cephali. Folia sparsa linearia  
in callosa-mucronata integerrima. Invol. hirtellum. Cor. om-  
Pappus Diplopappi. Ligulæ Erigeri. Color florum Chrysopsi-  
sus proprius.

#### Sect. I. PODONEJA.

vix favulosum. Cor. papilloso-pilosæ. Capitula longè pedunculata. —  
nctata.

*NEJA* (Don in Sweet! l. c.), ramosa, foliis subulato-linearibus parè et  
pilosus impunctatis, ramis apice longè aphyllis, involucri glabresculo  
è breviorè, ligulis 2-3-seriatis.  $\S$  in Mexico. Lodd. bot. cab. t. 1814.  
pus graminifolius Less. syn. p. 165. Folia angustissima ferè subulata.  
in 8-9 lin. diam. (v. s. ex hort. Barclay et Sweet.)

#### Sect. II. PHYLLONEJA.

alveolat. Corollæ glabræ. Capitula sessilia. Folia punctata.

*ROCEPHALA*, caule subsimplici per totam longitudinem densè folioso,  
o-linearibus punctatis molliter villosis, involucri villonissimo disco  
ligulis 2-3-seriatis elongatis.  $\S$  in Brasiliæ provinciâ Rio-Grande.  
re. pedalis foliis obtectus. Folia 12-15 lin. longa,  $1\frac{1}{2}$  lata. Invol.  
multiserialis elongato-acuminatæ, apice purpurascente rigidulo. Capit.  
è sesquipollicari. Recept. breviter alveolatum. Pappus ext. brevissimus.  
ferè Inularum sed antheræ omninò ecaudatæ. (v. s. in h. Mus. reg.  
lus. imp. Bras. sub n. 1001, 1006 et 1047 miss.)

#### Sect. III. MONOGYRIA.

-seriales. Capitula ad apices ramorum apice nudorum solitaria. Cor,  
Sectio Chrysopsidi affinis sed achænia oblonga nec obovata.

*HELLOIDES*, caule basi fruticoso multicauli cæspitoso, ramis adacen-  
molliter subhispidis ferè ad apicem usque foliosis 1-cephalis, foliis  
is aut lineari-sublanceolatis ciliato-subhispidis et glanduloso-puberis,  
quam linearibus subpuberulis pappo ext. parco brevi setiformi.  
Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1048!). Pars rami  
liformis vix foliis longior. Folia pollicem longa aut paulò longiora,  
lata. Pappi series ext. vix à pilis achæniæ distinguenda. (v. s. in h. Mus.  
)

*HELLOSA*, caule fruticoso ramoso, ramis villis subadpressis subhirsuto-  
pice longè nudatis vix hirtellis 1-cephalis, foliis subvillosis linearibus  
invol. squamis linearibus glabresculis, pappo ext. brevi subpaleaceo.  
siliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1015!). Sequenti affinis  
pecim. manca satis certa. Villositas caulis et foliorum et ciliorum vero-  
rentia satis notabiles characteres. (v. s. in h. Mus. reg. Par.)

*RAMIFOLIA*, caule fruticoso ramoso glabro, ramis hispidulis apice sub-  
aphyllis 1-cephalis, foliis linearibus elongatis hinc inde ciliatis, invol.

squamis linearibus subhirtellis, pappo ext. minuto vix paleaceo. ♂ in *Brasiliæ* prov. Rio-Grande (h. Mus. imp. Bras. n. 1034 et 1045). Primâ fronte videtur var. *N. tenuifoliæ* sed folia planiuscula linearia ferè pollicem longa, et pappus exterior vix oculo nudo facîle perspicuus triplò prioris ferè brevior. (v. s. in h. Mus. reg. Par.)

6. *N. tenuifolia*, caulibus fruticosis ramosis glabris, ramis longè foliosis apice nudis vix subpilosiusculis 1-cephalis, foliis lineari-subulatis tortilibus angustissimis longè ciliatis, invol. glabriusculo, ligulis linearibus, pappo utroque circ. 10-seto. ♂ in *Brasiliæ* prov. Rio-Grande (h. Mus. imp. Bras. n. 1009). Valdè *N. ciliari* affinis, sed folia 8-9 lin. longa subtortilia nec stricta. Pedunculi glabriusculi. Caulis magis fruticosus. Pappus pauciorisetus. (v. s. in h. Mus. reg. Par.)

7. *N. ciliaris*, basi suffruticosa multicaulis, caulibus adscendenti-erectis inferè glabratis apice subaphyllis hirsutis 1-cephalis, foliis lineari-subulatis scemilibus acutis erectis præsertim ad basin longè ciliatis inv. squamis linearibus dorso hirtellis disco paulò brevioribus, ligulis angustè linearibus. ♂ ♀ in *Brasiliæ* prov. Rio-Grande (h. Mus. imp. Bras. n. 1004!). Caulis subcæspitosi semipedales. Folia 5 lin. longa. Pappus duplex uterque circ. 15-setosus, ext. paleaceus, inter. setiformis. Capit. ferè *Nejæ* gracilis sed ligulæ 1-ser. (v. s. in h. Mus. reg. Par.)

#### Subdiv. 5. CHRYSOPSIDÆ.

Pappus duplex, exter. brevis subpaleaceus, inter. setosus. Ligulæ 1-seriales aut rarissimè 2-3-seriales.

CLXXX. *CHRYSOPSIS* Nutt. *gen. am.* 2. p. 150 (sect. *Inulæ*). *Ell. sketch* 2. p. 333. *excl. sp.* — *Diplogon Raf. non Brown.* — *Diplopappi sp. Less.* — *Inula sp. Pursh.*

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci herm. 5-dent. Recept. planum subalveolatum. Invol. imbricatum, squamis linearibus acuminatis. Anth. ecaudatæ. Achænia obovata compressa. Pappus disci et radii duplex, ext. brevis subpaleaceus, int. setiformis scaber elongatus. — *Herbæ* *Americanae rhizocarpicæ* *Inuloideæ* sæpiùs ramosæ et villosæ. Folia integra aut dentata alterna. Capitula ramos terminantia. Cor. omnes luteæ. — Differt ab *Inulâ* antheris ecaudatis, ab *Heterothecâ* achæniis omnibus eodem pappo donatis, ab *Oleariâ* achæniis compressis, à *Diplopappo* ligulis flavis.

#### §. 1. *Herbaceæ.*

1. *C. PINIFOLIA* (Ell. sketch 2. p. 335), glaberrima, caule rigido, foliis linearibus confertis rigidis integerrimis acutis, pedunculis corymbosis 1-cephalis, invol. squamis margine submembranaceis apice pubescenti-lanatis, achænio oblongo villosio. ♀ in collibus arenosis *Carolinæ* inter flumina *Flint* et *Chataboocæ* legit cl. *Elliot*. Pappus int. rufus, ext. brevis albidus. (v. s. comm. à cl. *inv.*)
2. *C. GRAMINIFOLIA* (Ell. sketch 2. p. 334), tota adpressè sericeo-villosa nitens, caule supernè etiam folioso, foliis lanceolato-linearibus acuminatis erectis integerrimis nervosis, corymbo composito, invol. squamis acuminatis dorso subpubescentibus et distinctè glandulosis, achænio oblongo utrinquè attenuato villosio. ♀ in arenosis alceis *Carolinæ* et *Georgiæ*, *Floridæ* (Nutt.) *Inula graminifolia* Michx. fl. bor. am. 2. p. 122. Nutt. gen. 2. p. 151. *Erigeron nervosum* Willd. sp. 3. p. 1953. *Erigeron glandulosum* Poir. dict. 8. p. 487 non Walt. Pappus rufus, ext. brevis setiformis. Caulis apice polycephalus, laxè corymbosus. Specimen habeo è *Georgiâ* simplex 1-cephalum. An dist. var.? (v. s.)
3. *C. ARGENTEA* (Ell. sketch 2. p. 334), tota adpressè sericeo-villosa nitens, caule supernè subnudo, foliis lanceolato-linearibus erectis acuminatis integerrimis vix subtiliter nervulosis, corymbo subpaniculato, invol. squamis acumi-

natis dorso pubescentibus non aut vix glandulosis, achænio oblongo villosio.  $\frac{1}{2}$  in sicca Georgia, Carolina (Ell.), agri Kentuckensis orient. (Raf.!), à Virginia ad Floridam (Nutt.) Inula argentea Pers. ench. 2. p. 452. Nutt. gen. 2. p. 151. Diplogon stenophyllum Raf. in litt. Valdè accedit ad C. graminifolium. Nomina facile decipiunt. Hæc potiùs graminea, et altera sericea dicendæ. Nostra interdum forè glabra lucens et tunc verè graminum folia refert. (v. s.)

C. FALCATA (Ell. sketch 2. p. 336 in adn. n. 4), lanato-villosa, foliis sessilibus linearibus acutissimis subfalcato-patentibus nervosis utrinquè pilosis, pedunculis paucis axillaribus corymbosis involucriisque villosis.  $\frac{1}{2}$  in arenosis pinetis Novæ-Cæsareæ. Inula falcata Pursh fl. bor. am. 2. p. 532. Aster gracilentus Banks herb. ex Pursh. Inula Mariana var.  $\beta$ . Nutt. est verisim. diversa. Herba 7-9 poll. longa, meo sensu C. graminifoliæ potiùs quàm Mariæ affinis. (v. s. comm. à cl. Torrey.)

C. MARIANA (Ell. sketch 2. p. 335), caule villosa, foliis sessilibus oblongo-ellipticis inf. subovatis obtusis distanter dentatis mucronulatis laxè subsericeo-villosis, corymbo simplici, pedunculis fastigiatis, invol. squamis linearibus acutis viscido-pubescentibus, achænio obovato villosa.  $\frac{1}{2}$  in arenosis siccis Novæ-Cæsareæ (Raf.! Torr.), Carolina (Ell.), Georgia, etc. Inula Mariana Linn. sp. 3. p. 1240 excl. syn. Pluk. Pursh fl. 2. p. 531. Inula Mariana var.  $\alpha$  Nutt. gen. am. 2. p. 151. Diplogon Marianus Raf. in litt. Inula glandulosa Lam. dict. 3. p. 259 ex syn. — Mill. dict. ic. t. 57. Nomen fortè falsum nam provenit ex syn. Pluk. nunc excluso et incertum manet an stirps in Marylandiâ cresc. (v. s.)

C. TRICOPHYLLA (Ell. sketch 2. p. 336), caule basi laxè lanuginoso superne glaberrimo, foliis oblongis obtusis laxè lanuginosis margine anscabris, superioribus minoribus glabriusculis, pedunculis corymbosis fastigiatis 1-cephalis glaberrimis, invol. hemisphærici squamis oblongo-linearibus glabris, apice subfoliaceo parè recurvo, ligulis 14-16, achænio oblongo villosa.  $\frac{1}{2}$  in Carolinâ infer. (Fraser!) et Georgiâ (Nutt.). Inula trichophylla Nutt. gen. 2. p. 150. Valdè accedit ad C. gossypinam sed facilè glabritie partium super. differt. (v. s.)

C. GOSSYPINA (Ell. sketch 2. p. 237), tota laxè lanuginoso-candicans, foliis sessilibus oblongo-spathulatis obtusis integerrimis, pedunculis corymbosis fastigiatis 1-cephalis puberalis, invol. squamis linearibus acutis puberis, achænio oblongo villosa-hirtello.  $\frac{1}{2}$  à Virginia ad Floridam et Georgiam, frequens in pinetis. Inula gossypina Michx. fl. 2. p. 122. Nutt. gen. 2. p. 150. Inula glandulosa Lam. dict. 3. p. 259 ex herb. sed excl. syn. Habitus Inularum. Ligulæ 20-25. Pappus int. rufus, ext. brevis albus. (v. s.)

C. DENTATA (Ell. sketch 2. p. 337), tota lanuginosa, foliis cuneato-obovatis obtusis sinuato-dentatis, superioribus oblongo-ovalibus integris, corymbo simplici, ligulis 30-35 nervosis, achænio hispidoblongo.  $\frac{1}{2}$  in Georgiâ circa Louisville. Affinis C. gossypinæ.

C. VILLOSA (Nutt. gen. 2. p. 151), subdecumbens hirsuto-villosa, caule ramoso corymboso, foliis sessilibus integris lineari-oblongis subspathulatis acutis, infernè ciliatis margine scabris, capitulis fastigiatis, achænio obovato-villosa.  $\frac{1}{2}$  in planitie Missouriianâ à fonte ad confluentem (Nutt.). Anellus villosus Pursh fl. 2. p. 564. Sideranthus integrifolius Fras. ex Steud. Species ex Nutt. distinctissima ob canlis pubescentiam duplicem villosam nempe et pilosam. Ligulæ flavæ circ. 25 lineari-oblongæ.

## §. 2. Fruticosæ.

1. C. HIRTELLA, caule basi fruticuloso ascendente ramoso, foliis ad basin ramosum confertis linearibus integerrimis hirtellis, ramis basi hispidulis foliosis apice glabriusculis aphyllis 1-cephalis, invol. demum omnino reflexi squamis linearibus dorso hispidulis.  $\frac{1}{2}$  in Amer. bor. occident. ad flumen Columbia legit cl. Douglas. Herba 5-6 pollic. Folia 1  $\frac{1}{2}$  poll. longa,  $\frac{1}{2}$ -1 lin. lata. Ligulæ circ. 25 lineares flavæ. Achænia oblonga compressa parè pubera. Pappus duplex

uterque paucisetosus. Recept. sine alveol. rudim. nudum, setulam ex areolis præ. marginalibus proferens. (v. s. comm. ab hon. soc. hort. Lond.)

11. *C. CANESCENS*, caule basi fruticoso erecto ramosissimo, ramis junioribus tomentoso-canis adultis foliisque adpressè puberis cinereo-subcanescentibus, foliis lineari-subulatis integerrimis infernè confertis supernè nudis, pedunculis 1-cephalis subaphyllis laxè corymbosis, invol. squamis linearibus acutis disco brevioribus dorso puberis.  $\frac{5}{8}$  in Amer. boreal. occid. ad flumen Columbia legit cl. Douglas. Folia æquipoll. longa. Ligulæ circ. 25 lineares ex sicco pallidè flavæ disco duplò triplòve longiores. Achænia tenuia sparè pilosa. Recept. scrobiculatum. (v. s. comm. ab hon. soc. hort. Lond.)

† *Species exclusæ.*

- C. ALBA* Nutt. = *Dœllingeria ptarmicoides*.  
*C. AMYGDALINA* Nutt. = *Diplostephium umbellatum*.  
*C. DIVARICATA* Nutt. = *Heterotheca Lamarckii*.  
*C. HUMILIS* Nutt. = *Diplostephium amygdalinum*.  
*C. LINARIFOLIA* Nutt. = *Diplopappus linarifolius*.  
*C. LINIFOLIA* Nutt. = *Galatella linifolia*.  
*C. OBOVATA* Nutt. = *Diplostephium obovatum*.  
*C. SCABRA* Ell. = *Heterotheca scabra*.

CLXXXI. FRESENIA.

Capitulum multi- (15-16) florum homogamum. Invol. squamæ circ. 3-seriales imbricatæ lineares margine submembranaceæ dorso glandulas 1-3-ariatis dispositas gerentes. Recept. areolatum angustum. Cor. tubulosæ glabræ 5-dentatæ fauce non ampliata. Anth. ecaudatæ. Stylus Chrysocomæ. Achænia compressa erostria sericeo-villosa. Pappus duplex (more Vernoniæ), exter. brevis paleiformis, inter. setaceus scabridus longus. — Suffrutices Australi-Africani glabri, ramis oppositis apice nudis 1-cephalis. Folia opposita linearia acuta ferè subulata integerrima sæpè in axillis fasciculata. Cor. pallidè ex sicco flavæ. — Genus omninò *Diplopappis* peduncularibus simile, diversum ligulis nullis et foliis oppositis, cæterùm à *Pteroniâ* et *Chrysocomâ* pappo biseriali inæquali facilè distinctum, à *Vernoniâ* stigmatè Asteroideo, à *Pegolettiâ* antheris ecaudatis diversum, dicavi cl. Fresenio botanico Francofurtensi qui *Ruppellii* plantas Arabicas et Africanas egregiè descripsit.

1. *F. LEPTOPHYLLA*, foliis oppositis lineari-subulatis glabris.  $\frac{5}{8}$  in Africa Capens. ad Zeederbergen legit cl. Drege. Folia 3-4 lin. longa,  $\frac{1}{4}$  lin. lata. (v. s.)  
 2. *F. ? SCAROSA*, foliis alternis ad basin ramorum confertis linearibus adpressè villosis.  $\frac{5}{8}$  in Africa australi legit cl. Drege! Canlis fruticosos demissus brevis. Rami veteres reliquiis foliorum vestiti. Folia 6-8 lin. longa aut omninò linearia aut versùs apicem sublata. Rami floridi 3 pollic. 1-cephali molliter hirsutuli. Invol. biserialè squamis linearibus dorso villosis. Cor. flavæ glabræ. Achænium oblongum sericeo-villosum. Pappus duplex scaber, exter. brevis. An *Diplopappi* sp. discoidea? An *Freseniæ* sp. alternifolia? An genus proprium? (v. s.)

Subdiv. 6. SOLIDAGINÆ. — Chrysocomearum et Asterearum gen. Less

Ligulæ nunc nullæ nunc 1-seriales. Pappus disci et radii conformis pilosus 1-serialis. Achænia (ferè omnium) angulata seu teretiuscula nec compresso-plana. — Generum discoideorum et radiatorum junctionem jubent 1°. Specier. ligulis

panceisve imò solitariis cum discoideis analogia (*Euthamia*, *Bigelovia*); ationes in toto ordine frequentes inter capitula discoidea et radiata sæpè a generibus imò iisdem speciebus obviæ, præsertim in capitalis homomorphiæ.

II. *BIGELOWIA* DC. non Spreng. nec Smith nec Raf. (1). — *Chrysocoma* Americanæ auct.

tulum 3-5-florum homogamum aut ligulâ unicâ fœmineâ heterostam. Recept. angustum squamam setaceam basi latiusculam achænia inter flores centralem gerens. Invol. oblongum, squamis imbricatis erectis. Achænia oblonga subangulosa pubescenti-pappus pilosus 1-serialis rigidus scaber. — *Herbæ Borealiæ perennes*. Folia alterna integra, caulina oblonga aut linearia in corymbum disposita. Cor. lutæ. — Genus differt à *Linosyppo* 1-seriali, à *Chrysocomâ* invol. squamis non foliaceis, ab *capitulis* paucis nec multifloris, achæniis teretiusculis nec com-  
pactis, pappo rigido, recept. subsquamigero, etc. A *Chrysocomâ* um dicavi cl. J. Bigelow qui floræ Americanæ auream coronam *ostoniensi* et medicâ addidit.

§. 1. *Genuinæ, glaberrimæ, capitulis 3-4-floris.*

*NUDATA*, herbacea glaberrima, foliis radicalibus spathulato-lanceolatis obtenuiter 3-nerviis, caulinis distantibus linearibus, corymbo composito fasciculato, capitulis 3-4-floris. 2. ad paludum margines à Virginia ad Floridam. *Chrysocoma nudata* Michx. fl. bor. am. 11. p. 101. Nutt. gen. 2. p. 137. Ell. 1. 2. p. 309. (v. s.)

*LINEATA*, herbacea glaberrima, foliis angustis linearibus enerviis, caule to ramoso, ramis corymbiferis fastigiatis, capitulis oblongis 3-4-floris, ocri squamis glutinosis adpressis. 2. ad paludum margines in Novâ-  
reâ (Nutt.) et in Floridâ prope Savannah (h. Dnb.). *Chrysocoma virgata* gen. am. 2. p. 137. (v. s.)

*IPURIÆ, pubescenti-tomentosæ, capitulis 5-floris.* — *An certè congeneres?*

*MISSOURIENSIS*, herbacea tomentoso-pubescenti, foliis angustis linearibus, sub laxiusculo, invol. 5-floris, squamis glabris, exter. patulis. 2. ad ripas nis Missouri. *Chrysocoma nanseosa* Pursh fl. bor. am. 2. p. 517. Nutt. am. 2. p. 137.

*SACUNCULOIDES*, suffruticosa, ramis albidis pulverulento-tomentosis, foliis ribus 3-nerviis impunctatis, corymbo fastigiato, capitulis 5-floris, invol. lato levi. 3. in locis denudatis ad ripas flum. Missouri. *Chrysocoma draculoides* Pursh fl. 2. p. 517 non Lam. *Chrysocoma graveolens* Nutt. gen. 1. p. 136. Odor rutaceus.

*l. Solidagineæ, glabra, capitulis 5-floris, flor. 4 tubulosis, 1 ligulato.* — *An Solidagini an Bigelowiæ associanda?*

*UNILIGULATA*, glabra, foliis lanceolatis utrinquè acuminatis serratis, par-  
tibus compositis multifloris, fasciculis erectis, invol. angusto-oblongis 5-floris, 1 unicâ. 2. in Novâ-Cæsareâ (Greene), in arenosis à Virginia ad Carolinam

---

*BIGELOWIA* Smith cycl. = *Borya* Willd. — *BIGELOWIA* Spr. non. entd. est ex  
st. *Samyda* spec. e. Spreng. syst. est eadem ac *Borreria*. — *BIGELOWIA* Raf.  
ibus ignota.

(Pursh). *Solidago pauciflosculosa* Michx. fl. bor. 2. p. 116. Pursh fl. 2. p. 53: huc pertinere videtur etiamsi nostra non habeat folia obtusa enervia nec caulem fruticosum. Vereor ne cl. Michx. confuderit Big. nudatam et B. pauciflosculosam. Pursh qui ex Michx. dicit caulem fruticosum addit signum  $\gamma$ , unde caract. non videtur satis notus. (v. s. comm. à cl. Greene.)

CLXXXIII. *SOLIDAGO* Linn. gen. n. 955. Gärtn. fr. 2. p. 447. t. 170. *Nun. gen. am.* 2. p. 159. — *Virga-aurea* Tourn. inst. p. 483. t. 275. *Vaill. act. ac. Par.* 1720. — *Doria Adans. fam.* 2. p. 124. — *Solidago* et *Euthamia* Cass. Less.

Capitulum pluri- aut multiflorum radiatum, ligulis 5-15 oblongis fœmineis 1-serialibus sæpè inter se distantibus, fl. disci hermaphr. 5-dentatis. Recept. epaleaceum nunc omninò nudum nunc alveolatum seu fimbrilliferum. Invol. adpressè imbricatum polyphyllum sæpè oblongum. Achænium teretiusculum multicostatum. Pappus 1-serialis pilosus scabridus. — Herbæ pleræque Americanæ, rhizocarpicæ, rariogerontogæ, interdum basi suffrutescentes elatæ rusticæ. Folia alterna integra aut serrata. Racemi nunc terminales nunc axillares, erecti patientes imò reflexi, pedicellis secundis et cymas subscorpioides confidentibus aut verè racemosis. Ligulæ semper luteæ exceptâ S. bicolor. Discus semper luteus.

Sect. I. *VIRGAUREA*. — *Virga-aurea* Tourn. *Vaill.* — *Solidago* Cass. dict. 37. p. 459 et 472. Less. syn. 163.

Involucrum imbricatum, squamis liberis. Flor. ligulati pauciores quàm fl. disci. Recept. nudum aut alveolatum, non fimbrilliferum. — Racemi paniculati aut simplices, non corymbosi, capitulis ad apices pedicellorum solitariis.

### Ser. I. UNILATERALES Loud. hort. brit.

Racemi secundi subscorpioides.

\* *Triplinerviæ, foliis triplinerviis.*

1. *S. CANADENSIS* (Linn. sp. 1233), caule villosa erecto, foliis lanceolatis serratis triplinerviis scabris, racemis paniculatis secundis recurvis, ligulis abbreviatis  $\gamma$  in sepibus et agris Amer. bor. à Canadâ ad Pensylvaniam (Pursh) et in montibus Carolinæ (Ell.). Variat foliis nudiusculis et scabris, serratis aut sub integerrimis, racemis elongatis et patentibus. — Pluk. alm. t. 263. f. 1. Hec S. humilis Mill. dict. n. 16. Planta polymorpha, ligulis brevissimis inter omnes ferè congeneres distincta. Confr. diss. Tauschii in florâ 1828. p. 497. in qua et plantis hortensibus plures varietates aut species huic et seq. affines distinguuntur. (v. s. ex Kentucky à cl. Rafin. commun. et v. c.)
2. *S. PROCERA* (Ait. kew. ed. 1. v. 3. p. 211), caule hirsuto erecto, foliis lanceolatis serratis triplinerviis scabris subtus villosis, racemis spiciformibus erectis. innuptis nutantibus, ligulis abbreviatis.  $\gamma$  in sepibus et paludosis Amer. bor. à Canadâ ad Virginiam (Pursh) et fortè in Georgiâ occid. (Ell. sketch 1 p. 369?). Herba 4-6-pedalis. Olim in hort. Par. culta sub nom. *S. hirsuta* (L'Hér. herb.). *S. procera* Desf. cat. Affinis *S. Canadensi* sed capitula multo minora. (v. v. c.)
3. *S. LONGIFOLIA* (Schrad. hort. Gœtt. ex litt. 1832), caule erecto tereti villosa paniculato-ramoso, foliis lineari-lanceolatis acuminatis triplinerviis super scabris subtus pubescentibus, inf. serratis, sup. integerrimis, racemis elongatis patulis secundis in paniculam dispositis, invol. glabri squamis subscutis, ligulis linearibus vix disco longioribus.  $\gamma$  in Amer. bor. ? (v. s.)
4. *S. NUTANS* (Desf. cat. h. Par. ed. 3. p. 402), caule pubescenti subhirsuto erecto, foliis lanceolatis acutis argutè serrulatis triplinerviis super scabris sub-

abscentibus, paniculâ ante anthesin nutante, ligulis abbreviatis.  $\frac{1}{2}$  in bor. ex testim. hortul. Panicula ampla. Capitula magnitudine media inter occeram et canadensem. (v. v. c. in h. Par. jam ann. 1815. sub hoc nom.)

**REFLEXA** (Ait. kew. ed. 1. v. 3. p. 211), caule erecto subvillosa, foliis oboatis acuminatis versus apicem argutè serratis scabris margine reflexis, subsecundis brevibus in paniculam angustam confertis, ligulis linearibus disco ferè duplò longioribus.  $\frac{1}{2}$  in pinetis et agris vetustis Amer. bor. à Casarea ad Carolinam (Pursh). (v. v. c.)

**SAGRANS** (Willd. enum. suppl. 59), caule erecto glabro, foliis glabris utrinquè attenuatis subtriplinerviis, infer. subserratis, superior. integris, racemis subsecundis, pedicellis pubescentibus, ligulis oblongis disco longioribus.  $\frac{1}{2}$  in Amer. bor. (Link. enum. 2. p. 531). Bracteæ lineariter arcuato-recurvæ. Invol. squamæ subvirescentes. (v. s. comm. à cl. Ard.)

**STEREOPHYLLA** (Linn. sp. 1234), caule erecto pilosiusculo, foliis lanceolatis triplinerviis glabris margine scabris inferioribus serratis, racemis paniculatis brevibus subsecundis, ligulis linearibus oblongis disco vix duplò longioribus.  $\frac{1}{2}$  in campis et sylvis siccis Amer. bor. à Canadâ ad Carolinam. h. fl. 530. Ell. sketch 370. Herba dimidiò minor quàm S. canadensis. Can. latiora. (v. v. c.)

**ANGULATA** (Muhl. ex Schrad! in litt. 1813. Mart! enum. h. Erlang. 1814), erecto angulato apice summo cum pedicellis scabrido, foliis inf. ovatis integris semiamplexicaulibus penninerviis, super. sessilibus lanceolatis triplinerviis, omnibus serratis suprâ scabris subtus lævibus, racemis brevibus utrobis secundis, ligulis paucis linearibus disco ferè brevioribus.  $\frac{1}{2}$  in r. septentr. Invol. squamæ glabræ submembranaceæ lineares. Achænia fuscula. Ligulæ 3-4, fl. disci 9-10. (v. s. comm. à cl. Schrader et Mart.)

**GLABRA** (Desf! cat. h. Par. ed. 3. p. 402), glaberrima erecta, foliis infer. oboatis, sup. linearibus lanceolatis, omnibus acuminatis argutè serrulatis concet brevè ciliatis triplinerviis, racemis secundis revolutis, ligulis 10-12 linearibus disco paulò superantibus.  $\frac{1}{2}$  in Amer. bor. In sylvis et sepibus à h. Angliâ ad Pensylvaniam (Pursh). Rami paniculæ subpubescentes. Folia basi attenuata. S. serotina Ait. kew. ed. 1. v. 3. p. 211? S. altissima R. L'Hér! herb. (v. v. c. in h. Par. anno 1819, sed altera species sub hoc anno 1829 colebatur.)

**GIGANTEA** (Ait. kew. ed. 1. v. 3. p. 211), caule erecto glabro, foliis lanceolatis utrinquè attenuatis serratis margine scabris triplinerviis, racemis patentibus secundis, pedunculis hirtis, ligulis linearibus oblongis disco duplò longioribus.  $\frac{1}{2}$  in Amer. bor. In apricis ad latera collium fertilem à h. Angliâ ad Virginiam (Pursh). Valdè accedit ad S. serotinam. Nervi foliorum laterales validiores. (v. s. ex Noveboraco comm. à cl. Torrey.)

**CILIARIS** (Muhl. in Willd. sp. 3. p. 2056), caule erecto glabro, foliis glabris, infer. ovalibus acuminatis basi longè attenuatis ciliatis serratis penninerviis, super. lanceolatis subintegris triplinerviis, pedunculis glabris, bracteis linearibus, ligulis linearibus oblongis disco paulò longioribus.  $\frac{1}{2}$  in Pensylvaniâ (Muhl.), Noveboraco (Torr!). S. glabra hort. Par. 1829. non Desf. (v. s. et )

**GLIELIA**, caule erecto glabro tereti, foliis oblongo-lanceolatis sessilibus utrinquè attenuatis argutè serratis medio triplinerviis utrinquè et ad marginibus glabris et lævibus, racemis axillaribus folio sæpè brevioribus patentibus, capitulis longè pedicellatis, pedicellis glabrisculis, ligulis disco duplò longioribus.  $\frac{1}{2}$  patr. ign. S. dubia Scop. ex hort. Ital. non Schult. Affidetur S. patulæ et S. argutæ sed folia triplinervia. Folia radicalia oblonga obovata in petiolum attenuata. (v. v. c. anno 1808 in hort. cl. Glielie Griedi prope Genavam.)

**SCABRIDA**, caule erecto basi glabro tereti apice puberulo angulato, foliis



oblongo-lanceolatis mucrone apiculatis integerrimis aut hinc inde adpressis triplinerviis utrinque subscabridis secus nervum et ad margines rigidis scabro-ciliatis, racemis secundis, bracteolis subulatis, invol. squamis glabris acutis, ligulis lineari-oblongis disco paulò longioribus. 2 $\frac{1}{2}$  in Mexico legit cl. Alaman. Affinis *S. Canadensi* et *S. fragranti*. (v. s.)

14. *S. VELUTINA*, tota pube brevi conferta pulverulento-velutina, caule erecto tereti apice paniculato, foliis oblongo-lanceolatis mucrone subapiculatis integerrimis triplinerviis, racemis erectis vix subsecundis parè bracteolatis, invol. squamis linearibus acutis, ligulis disco paulò longioribus. 2 $\frac{1}{2}$  in Mexico circa urbem (Berl. Alam.) et var. paniculà contractà ad Real del monte (Hank.). Sol. Mexicana Berl. pl. exs. n. 930. à bot. Mexic. plurimis cum *S. limonifolia* et *S. paniculatà* sæpè confusa sed facillè distincta. (v. s.)

15. *S. DECEMFLORA*, caule erecto tereti foliisque pube brevissimà subscabris, foliis oblongis mucronulatis integerrimis subtriplinerviis, radicalibus basi attenuatis, racemis subsecundis in paniculam corymbosam dispositis, invol. glabri squamis linearibus, capitulis 10-floris, ligulis 5 minutissimis. 2 $\frac{1}{2}$  in Mexicana prov. Texas districtibus orientalibus legit cl. Berlandier (pl. exs. n. 1924). Achænia sericeo-villosa. Herba pedalis. (v. s.)

16. *S. PUNCTICULATA*, caule erecto tereti glabro, ramis apice subscabridis, foliis oblongo-linearibus margine subscabris integerrimis subtrinerviis glabris utrinque sub lente opacè puncticulatis, racemis subsecundis in paniculam oblongam dispositis, invol. glabri squamis linearibus, capitulis 10-floris, ligulis 4-5 minutis. 2 $\frac{1}{2}$  in Mexico ad Rio de la Guadalupe legit cl. Berlandier (pl. exs. n. 1900). Affinis *S. decemfloræ* sed præter notas indic. achænia glabrinscula (v. s.)

17. *S. POLYGLOSSA*, caule erecto tereti hirsuto, foliis elongato-lanceolatis mucrone acuminatis, super. apice parè et acutè serratis triplinerviis utrinque pubescenti-hirtellis, racemis secundis patulis subarcuatis in paniculam dispositis, invol. glabri squamis linearibus acutis, floribus circ. 20, ligulis 12-14 disco paulò longioribus. 2 $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Habitus et folia ferè *S. giganteæ* aut *S. notanensis*. Capitula ferè *S. microglossæ*. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 860 missa.)

18. *S. MICROGLOSSA*, caule basi suffruticuloso erecto tereti subhirsuto, foliis sessilibus elongato-lanceolatis acuminatis puberulis trinerviis, infer. adpressis et distanter serrulatis, super. integerrimis, racemis secundis in paniculam oblongam dispositis, pedicellis bracteolatis, invol. squamis linearibus glabris, floribus circ. 20-25, ligulis 10-15 disco æqualibus. 3 $\frac{1}{2}$  in Brasiliæ parte dictà Banda orientale legit cl. Bacle. Achænia gracilia glabrinscula. (v. s. comm. à cl. Bacle.)

β. *Megapotamica*, paniculæ ramis longioribus magis patulis. 2 $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Priori similis sed major et panicula magis diffusa. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 846, 854 et 857.)

19. *S. MARGINELLA*, caule suffruticuloso? erecto tereti vix puberulo, foliis linearibus integerrimis acuminatis ciliolato-scabris, infer. basi attenuatis subtriplinerviis, racemis secundis patulis, pedunculis puberis, invol. squamis linearibus utrinque lineà scariosà albà marginatis, floribus circ. 25, ligulis 15 disco æqualibus. 3 $\frac{1}{2}$  in Brasiliæ parte dictà Banda orientale legit cl. Bacle. Pappus rufus. Achænia glabra. Stirps vix pedalis. Affinis *S. microglossæ*. Specimen notabilis pilis rigidis brevibus nervulis secus marginem folii circ. biserialibus (v. s. comm. à cl. Bacle.)

γ. *sublanceolata*, caule elatiori glaberrimo, foliis lineari-lanceolatis utrinque glabris, paniculæ ramis puberis, ligulis minimis. 2 $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Herba sesqui- aut bipedalis. Folia 2-3 lin. lata. Pappus albus (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 870 missa.)

\*\* Venosæ, foliis penninerviis aut venosis.

20. *S. ASPERA* Ait. kew. ed. 1. v. 3. p. 212.), caule erecto tereti piloso, foliis

ato-ellipticis scaberrimis rugosis serratis, infer. penninerviis, sup. enerviis, ramis paniculatis secundis, ligulis lineari-oblongis disco paulò longioribus. in Amer. bor. campis siccis et sylvis à Noveboraco ad Carolinam (Pursh 530). Verisim. secus montes nec in planitie (Ell. sketch 2. p. 371). — Dill. h. 2. t. 305. f. 392. opt. (v. v. c.)

1. **ALTISSIMA** (Linn. sp. 1233. excl. Dill. syn. omnibus), caule erecto hirtio, foliis lanceolatis penninerviis profundè serratis scaberrimis rugosis, paniculis nudis, ligulis 7-10 lineari-oblongis disco vix longioribus. 2/ in Amer. bor. siccis et agris frequens. — Mart. cent. t. 14. S. rigidula Bosc! herb. et hort. ris. ann. 1828. Huc ex cl. Ait. pertinent S. altissima, pilosa, recurvata et giniana Mill. dict. Species valdè varians. S. Canadensis affinis. (v. v. c.)

2. **RUGOSA** (Mill. dict. n. 25. Willd. sp. 3. p. 2058), caule erecto hirtio, foliis infer. ovatis dentato-serratis, super. lanceolatis adpressè serratis omnibus bris rugosis acutis, racemis paniculatis secundis patentissimis, ligulis lineari-oblongis disco vix duplò longioribus. 2/ in Amer. bor. siccis à Canadà ad Virginiam (Pursh. fl. 537), in pinguibus Carolinæ (Ell. sketch 2. p. 372). An altissima var. ? ex Ell. — S. altissima E. Ait. kew. 3. p. 213. — Dill. elth. 108. f. 396. (v. s.)

3. **VILLOSA** (Pursh fl. 2. p. 538), caule villis patulis hispido tereti, foliis ser. sessilibus ovato-lanceolatis mucronatis integerrimis sæpè subtus secus nervum hispidis, infer. serrulatis, omnibus 1-nerviis margine scabridis, racem. secundis folio paulò longioribus in paniculam dispositis, invol. glabri ramis acutis, ligulis vix disco longioribus. 2/ in Amer. bor. sylvis et agris quous (Pursh), in Carolinà (Ell. sketch. 2. p. 372). Habeo ex Carolinà (McC!) et ex Georgià (h. Duby!), sed specim. Georgic. habet capitula minorà. altissima Bosc. herb. S. Bosciana Moret! rel. Bosc. ined. Achania vix pulula. (v. s.)

4. **SCABRA** (Muhl. in Willd. sp. 3. p. 2059), caule erecto hirtio sulcato, foliis oblongis utriusquè attenuatis acuminatis suprà glabris subtus rugosis scabris disco adpressè serratis, racemis secundis, ligulis lineari-oblongis disco paulò longioribus. 2/ in agris et limitibus Amer. bor. à Pennsylvania ad Virginiam (Pursh fl. 537), in insulà longà (Rafin!). Folia subtus secus nervos nervulosos penninervios scabra. (v. s.)

5. **MEMORIALIS** (Ait. kew. ed. 1. v. 3. p. 213), caule erecto pube brevissimà nitido, foliis radicalibus obovato-cuneatis serratis basi longè attenuatis, lineis lanceolatis integerrimis scabris, racemis paniculatis secundis, ligulis. in arenosis et agris siccis Amer. bor. à Canadà ad Carolinam (Pursh fl. 537. sketch 2. p. 373.), in Kentucky (Raf.). Herba tota subcinerascens. (v. s.)

6. **PUMPERULA** (Nutt. gen. am. 2. p. 162), caule simplici tereti subpubescente, foliis lanceolatis integris pubescentibus utrinquè attenuatis, radicalibus subnatis, racemis spicatis axillaribus erectis condensatis, invol. squamis lineari-oblongis acutis, ligulis circ. 10 elongatis. 2/ in Novà-Cesareà (Green!) circa iboy, etc. (Nutt.). Habitus S. memorialis quoad folia sed in flor. diversa. a. comm à cl. Greene.)

7. **PATULA** (Muhl. in Willd. sp. 3. p. 2059), caule erecto glabro striato, foliis ellipticis serratis glabris, radicalibus oblongo-spathulatis, racemis paniculatis secundis patentibus, pedunculis pubescentibus. 2/ in Pennsylvania. Forradic. exactè Lencanthei vulgaris.

8. **ULMIFOLIA** (Muhl. in Willd. sp. 3. p. 2060), caule erecto glabro, foliis ellipticis profundè serratis acuminatis subtus villosis, radicalibus obovatis, ramis paniculatis secundis, ligulis abbreviatis. 2/ in paludosis et umbrosis sylvis à Noveboraco ad Virginiam (Pursh fl. 2. p. 538), in Georgià occid. (Ell. sketch 2. p. 375), ex quo et ex Muhl. caulis villosus striatus, folia suprà subbra, pedunculi villosi.

9. **ANGUSTA** (Ait. kew. ed. 1. v. 3. p. 313), caule erecto glabro, foliis glabris

ad marginem subscabrido-ciliolatis argutè et inæqualiter serratis, radicalibus ovato-oblongis basi attenuatis, caulinis ellipticis acutis, racemis paniculatis secundis, ligulis lineari-oblongis elongatis.  $\frac{1}{2}$  in sylvis et pratis Amer. bor. à Canadâ ad Virginiam (Pursh), in Carolinâ (Ell.), ad Noveboracum (Torr.). S. argentea Horn. suppl. p. 99 (saltem stirps sub hoc nom. in hortis vagans), non differt, ni fortè capitulis paulò majoribus. (v. s. sp. et v. c.)

30. S. CINERASCENS (Schwein in Ell. sketch 2. p. 375), caule erecto gracili pubescente, foliis elongatis lineari-lanceolatis basi attenuatis serratis utrinque subscabris pubescentibus, racemis secundis recurvis, pedunculis ligulisque elongatis.  $\frac{1}{2}$  in Carolinâ boreali prope Salem et in Georgiâ occid. (Ell.)
31. S. JUNCEA (Ait. kew. ed. 1. v. 3. p. 213), caule erecto glabro, foliis lanceolatis glabris margine scabris, infer. serratis, racemis paniculatis secundis.  $\frac{1}{2}$  in agris et sylvis arenosis à Novâ-Cæsareâ ad Carolinam (Pursh fl. bor. am. 2. p. 538), in Carolinâ superiori (Ell. sketch 2. p. 375 \*), in Georgiâ circa Savannah (h. Duby!). Invol. squamæ plerumque apice virides et obtusæ. Ligulae paucæ (1-4) parvæ disco paulò longiores. Achænia sublentè pubescentia subcompressa. Habitus ferè Erigerontis subulati sed ligulæ paucæ flavæ. (v. s.)
32. S. ELLIPTICA (Ait. kew. ed. 1. v. 3. p. 2060), caule erecto glabro, foliis ellipticis laevibus serratis penninerviis, nervis utrinque 1-2 validioribus, racemis paniculatis secundis, ligulis oblongis disco duplò longioribus.  $\frac{1}{2}$  in umbrosis rupestribus Amer. bor. à Noveboraco ad Virginiam (Pursh fl. 538). S. latissimifolia Mill. ex Ait. S. plantaginea Desf. cat. h. Par. ed. 3. p. 402. An S. elliptica Ell. sketch 2. p. 376, in pinguibus insulæ Paris lecta omninò eadem? Folia sup. subtriplinervia. Racemi suberecti. (v. v. c.)
33. S. VERRUCOSA (Schrud. hort. Gœtt. p. 12. t. 6. et in neu. jour. 1810. p. 146), caule verrucoso glabro erecto apice reflexo, foliis ovato-lanceolatis acutis serratis glabris, super. angustioribus, racemis axillaribus folio aequalibus compositis apice reflexis, ligulis ellipticis.  $\frac{1}{2}$  in Amer. bor. Affinis S. argutæ et S. ellipticæ dicitur.
34. S. RECURVATA (Willd. enum. p. 889), caule erecto pubescente, foliis lanceolatis serratis acuminatis margine scabris suprâ glabris subtus æcus nervo subscabridis, racemis elongatis secundis recurvatis paniculatis, ligulis oblongo-linearibus disco paulò longioribus.  $\frac{1}{2}$  in sylvis umbrosis Amer. bor., in Pensylvaniâ et Virginiâ (Pursh). (v. s. c. in h. Gœtt. à cl. Schrad. comm.)
35. S. TORTIFOLIA (Ell. ! sketch 2. p. 377), caule erecto pubescente, foliis lineari-lanceolatis subserratis patulis tortuosis suprâ nervoque scabris subtus subglabris, paniculâ pyramidatâ, racemis recurvis, capitulis 8-10-floris, ligulis lineari-oblongis disco ferè duplò longioribus.  $\frac{1}{2}$  in pascuis siccis Carolinæ et Georgiæ. S. odora Michx. fl. 2. p. 118. non Ait. Affinis S. odore sed folia minora tortilia non pellucido-punctata sed venis ramosis pellucidis notata. (v. s.)
36. S. PYRAMIDATA (Pursh fl. bor. am. 2. p. 537), caule erecto tereti hirs. foliis oblongis acutis subsemialexicauli-sessilibus glabris margine scabris rariter obsolete dentatis, paniculâ nudâ secundâ pyramidatâ, ramis reflexis pedunculis glabris, ligulis minutis.  $\frac{1}{2}$  in pinetis Georgiæ (Pursh). Vix à S. retrorsâ differt Nutt. gen. 2. p. 159. Ell. sketch 2. p. 378. Achænia lævis.
37. S. ODORA (Ait. kew. 3. p. 214), caule erecto pubescente, foliis lineari-lanceolatis integerrimis glabris pellucido-punctatis margine scabris, racemis paniculatis secundis, ligulis oblongo-linearibus disco paulò longioribus.  $\frac{1}{2}$  in siccis collibus Amer. bor. à Canadâ ad Carolinam (Pursh fl. 539), in montibus Alleghany (Raf.). Big. med. bot. 1. p. 188. t. 20. fl. boat. 304. non Willd. et Michx.? Sol. lanceolata Bosc! herb. S. retrorsa Michx. fl. 2. p. 117. Ell. ! sketch 2. p. 377. Vulgò dicitur *Goldenrod* et ab Anglo-Americanis ut *Thee sacerdarum* admittitur. Folia fragrantia, distillata oleum volatile aromaticum præbent — Pluk. alm. t. 116. f. 6. (v. s.)
38. S. GONOLADA, caule erecto tereti glabro, ramis angulatis glabriusculis p-

niculatis, foliis elongato-linearibus sessilibus integerrimis acuminatis subpenninerviis glabris margine scabris pellucido-venosis, racemis secundis, invol. squamis glabris acutis, ligulis vix disco longioribus.  $\mathcal{L}$  in urbis Mexici valle legit cl. Berlandier (pl. exs. n. 728). Folia infer. 3-4 poll. longa, 5-6 lin. lata sup. sensim angustiora et breviora. Invol. stramineum. (v. s.)

39. *S. AMPLEXICAULIS* (Linn. sp. 1232), caule erecto glabro, foliis glabris lævibus subcarnosis margine scabris integerrimis, radicalibus ovalibus basi in petiolum attenuatis, canlinis lineari-lanceolatis, racemis paniculatis secundis, pedunculis puberulis, ligulis lineari-oblongis disco duplò longioribus.  $\mathcal{L}$  in humidis et ad ripas rivorum à Canadà ad Pensylvaniam (Pursh), in Noveboraco (Linn. Torr.), in Carolinà (Ell.). *S. Noveboracensis* et *carnosa* Mill. *S. tardiflora* Moench. — Pluk. alm. t. 235. f. 5. Moris. ox. s. 7. t. 23. f. 15. Caulis ruber orgyalis. (v. v. et s.)
40. *S. CORYMBOSA* (Ell. sketch 2. p. 378), caule erecto glabro, ramulis hispidis, foliis inferioribus oblongo-lanceolatis, super. ovatis, omnibus carnosius rigidis glabris margine asperissimis ciliatisque, racemis secundis corymbosis, inferioribus recurvis, ligulis elongatis.  $\mathcal{L}$  in Georgià medià, circa Louisville (Ell.). Affinis *S. lævigatæ* et *S. limonifoliæ*. An *S. corymbosa* Poir. suppl. 5. p. 461 eadem aut diversa?

## Ser. II. ERECTÆ Loud. loc. cit.

Racemi erecti nec secundi.

\* *Serratæ, foliis serratis.*

41. *S. BICOLOR* (Linn. mant. 114), caule erecto pilosinsculo, foliis utrinquè pilosis, infer. ellipticis acutis dentatis in petiolum attenuatis, super. elliptico-oblongis mucronatis integerrimis, racemis erectis brevibus, invol. squamis obtusis, ligulis ovalibus disco paulò longioribus, achæniis glabris.  $\mathcal{L}$  in sylvis et collibus siccis Amer. bor. à Canadà ad Carolinam (Pursh). — Pluk. alm. t. 114. f. 3. Achænia in iisdem capitulis compressa et teretiuscula. Racemi nunc foliis paulò longiores nunc breves in axillis conferti. Ligulæ albæ, quæ notà ab omnibus *Solid.* differt et ad *Asterem* vergit. (v. v. c.)
42. *S. AXILLARIS* (Pursh fl. bor. am. 2. p. 542), caule glabro tereti stricto, foliis lanceolatis acuminatis remotè serratis glabris margine scabris, racemis in foliorum axillis sessilibus subglobosis erectis, ligulis ovali-oblongis disco duplò longioribus.  $\mathcal{L}$  in sylvis umbrosis à Canadà ad Virginiam (Pursh), in Kentucky (Raf.). *S. lateriflora* Raf. in litt. Dispositione capitulorum accedit ad *S. bicolor.*, sed ligulæ flavæ et folia diversissima. Ad hanc alludere videtur Ell. sketch 2. p. 387. in notà sub *S. flexicauli.* (v. s.)
43. *S. PETIOLARIS* (Ait. kew. ed. 1. v. 3. p. 216), caule erecto villosa, foliis ellipticis scabriusculis petiolatis, racemis erectis, ligulis elongatis.  $\mathcal{L}$  in siccis arenosis à Novà-Cæsareà ad Carolinam (Pursh fl. 2. p. 539), in montanis Carolinæ. (Ell. sketch 2. p. 383.)
44. *S. FLEXICAULIS* (Linn. sp. 1234), caule erecto flexuoso glabro angulato, foliis ovatis acuminatis serratis glabris, racemis axillaribus erectis folio brevioribus, summis in paniculam spiciformem confertis, ligulis oblongis disco paulò longioribus.  $\mathcal{L}$  in sylvis et agris Amer. bor. à Canadà (Pursh) ad Georgiam (Ell.). *S. flexicaulis* var.  $\alpha$  Willd. sp. 3. p. 2064. Pursh fl. 542. — Pluk. alm. t. 235. f. 3. Variat foliis angustè ovatis et latis imò interdum latioribus quàm *S. latifoliæ*. Caule regulariter flexuoso facillè distincta. Folia secus canlem in nervos 2 decurrentia. *S. scrophulariæfolia* Mill. hujus var. ex Steud. (v. v. c.)
- $\beta$  ? *ciliata*, foliis latè ovatis profundè argutè et inæqualiter serratis villosociliatis infernè longius acuminatis, racemis folio multò longioribus.  $\mathcal{L}$  in Amer. bor. Noveboraco (Torr.) et Canadà (Bosc.) *S. flexicaulis*  $\beta$  ? latifolia Torr. in litt. An fortè spec. propria? (v. s.)
45. *S. LATIFOLIA* (Linn. sp. 1234), caule erecto glabro angulato non flexuoso, foliis ellipticis utrinquè attenuatis serratis glabris, racemis axillaribus folio

brevioribus, ultimis in paniculam angustam concretis, ligulis oblongis disco paulò longioribus.  $\frac{1}{2}$  in Amer. boreali ad Canada (Pluk. alm. t. 335. f. 4). Videtur à *S. flexuosa* (jam mon. Linn. Desf., etc.) sat distincta, caule stricto, foliorum dentibus minoribus magis adpressis. Icones Pluk. optimè hujus et flexicaulis habitum expriment. (v. v. c.)

46. *S. HUMILIS* (Desf.! cat. h. Par. ed. 3. p. 402, non Pursh), caule erecto subangulato villis patulis hispido, foliis sessilibus confertis ovato-lanceolatis acuminatis nervoso-rugosis serratis suprà glabris subtus secus nervos villosis, racemis erectis in paniculam dispositis, invol. squamis glabris subacutis, ligulis disco vix longioribus.  $\frac{1}{2}$  in Amer. bor. ad Terram-Novam legit cl. Laphaie. In specim. spontanei rami breviores conferti, in hortensibus longiores nunc erecti nunc expansi imò reflexi. (v. v. c. et s. sp.)
47. *S. FLABELLIFORMIS* (Wendl. ex hort. Gœtt.), caule erecto glabro striato, foliis caulinis oblongo-lanceolatis utrinquè acuminatis ferè petiolatis subtriplicinerviis glabris margine scabridis hinc indè serratis aut integerrimis, racemis axillaribus erectis paniculatis folio brevioribus, pedicellis glabrisculis bracteolatis, invol. squamis linearibus margine subhyalinis, ligulis ovali-oblongis disco longioribus.  $\frac{1}{2}$  patr. ign. Involucra glabra substraminea. (v. s. comm. à cl. Wallroth.)
48. *S. AMBIGUA* (Ait. kew. ed. 1. v. 3. p. 217), caule erecto subflexuoso glabro angulato ramoso, foliis oblongo-lanceolatis densè serratis subtus pilosiusculis, racemis erectis in paniculam subaphyllam dispositis, ligulis oblongis disco duplò longioribus.  $\frac{1}{2}$  patr. ign. verisim. Amer. bor. Willd. sp. 3. p. 206, enum. h. Ber. 891. (v. s. ex hort. Berol. comm. à cl. Wallroth.)
49. *S. MULTIFLORA* (Desf.! cat. h. Par. ed. 1804. p. 103. ed. 3. p. 402), caule erecto glabro tereti ramosissimo, ramis sparsè pubescentibus in paniculam dispositis, foliis sessilibus lanceolatis acuminatis serratis glabris aut summis secus nervos subpuberis, racemis erectis, invol. squamis glabris acutis, ligulis disco paulò longioribus.  $\frac{1}{2}$  in Amer. bor. Pers. ench. 2. p. 449. n. 34. Poir. suppl. 5 p. 461. uterq. ex hort. Paris. (v. v. c.)
50. *S. GRACILIS* (Poir. dict. 8. p. 476? DC. pl. rar. j. de Genev. not. 6. p. 6. n. 2), caule erecto glabro subsimplici, foliis lanceolatis acuminatis serratis glabris ciliolatis penninerviis, infer. ovali-lanceolatis basi attenuatis, summis subintegerrimis, pedunculis puberulis, racemis erectis oligocephalis in paniculam angustam dispositis, invol. squamis vix acutis, floribus 9, ligulis 4 disco duplò ferè longioribus.  $\frac{1}{2}$  Patr. ign. verisim. Amer. bor. Herba sesquipedalis gracilis, caule rubente. Cl. Poir. dixit folia integerrima, sed in hortis sub nomine Poir. ubiquè occurrit. (v. v. c.)
51. *S. SCHRADERI*, caule erecto tereti sparsè pubero paniculato-ramoso, foliis lineari-lanceolatis acuminatis apice serratis glabris margine subscabris, summis integerrimis, racemis erectis brevissimis bracteolatis, invol. squamis acutis, ligulis linearibus disco æqualibus.  $\frac{1}{2}$  Patr. ign. verisim. Amer. bor. *S. gracilis* Schrad.! hort. Gœtt. ex litt. 1832. Affinis *S. gracili* sed folia longiora, angustiora paulò rigidiora. Rami longi in paniculam dispositi et racemulos brevè axillares longè gerentes. Ligulæ multò breviores, etc. (v. v. c.)
52. *S. SALICINA* (Elliot sketch 2. p. 389), caule elato gracili supernè pubescenti scabrinusculo, ramis virgatis elongatis erectis, foliis lanceolatis suprà scaberrimis subtus glabris, inferioribus serratis, racemis subsecundis, ramulis breviter recurvis.  $\frac{1}{2}$  in Georgiâ occid. Ligulæ 5 tenues. Achænia ferè glabra.
53. *S. CÆSIA* (Linn. sp. 1234), caule erecto lævi, foliis lanceolatis acuminatis serratis glabris, racemis axillaribus erectis, ligulis disco paulò longioribus.  $\frac{1}{2}$  in sylvis et agris demissis Amer. bor. à Novâ-Angliâ ad Carolinam et Georgiam. Pursh fl. 540. Ell. sketch 2. p. 385. Ait. kew. ed. 1. v. 3. p. 217. — DC. elth. 2. t. 307. f. 395. Caulis purpurascens-cæsius. Achænia ferè glabra. Invol. squama obtusiuscula. (v. s. comm. à cl. Torrey.)
54. *S. LIVIDA* (Willd. enum. 891), caule glabro paniculato, foliis lanceo-

serratis glabris margine scabris, ramis apice racemosis, ligulis elongatis.  $\mathcal{L}$  in renosis agris et sylvis à Noveboraco ad Virginiam. (Pursh fl. 2. p. 541.)

**S. HISPIDA** (Muhl. in Willd. sp. 3. p. 2063), caule erecto hispido scabro, foliis lanceolatis scabris integerrimis, radicalibus serratis, racemis erectis, ligulis medioeribus.  $\mathcal{L}$  in campis graminosis et sylvis Amer. bor. à Novâ-Cæsareâ ad Virginiam (Pursh fl. 541). Similis *S. cæsiæ* sed folia caulina integerrima.

**S. HIRTA** (Willd. enum. p. 891), caule paniculato hirta, foliis lanceolatis trinque scabris, caulinis serratis, rameis integerrimis, racemis erectis, ligulis longatis.  $\mathcal{L}$  in Amer. bor. Cæst. ign.

**S. GRANDIFLORA** (Desf. cat. h. Par. ed. 3. p. 403), caule erecto villosa-scabro pice paniculato, foliis utrinque villosa-scabris omnibus serratis acutis, infer. ellipticis basi attenuatis, super. oblongo-lanceolatis, racemis erectis compactis, invol. squamis obtusis, ligulis 8-10 elongatis.  $\mathcal{L}$  patr. ign. verisim. Amer. bor. Valde accedit ad *S. rigidam* sed præter. car. ind. differt foliis radic. vix petiolatis, imbo basi attenuato nec obtuso, acheniis puberis, fol. superioribus rariter serratis sed non integerrimis, etc. (v. v. c.)

**S. RIGIDA** (Linn. sp. 1235), caule erecto piloso-scabro apice paniculato, foliis rigidis scabris, radic. petiolatis ovalibus crenato-dentatis, caulinis sessilibus ovato-oblongis integerrimis, racemis erectis compactis, invol. squamis obtusis, ligulis 8-10 elongatis.  $\mathcal{L}$  in montanis Amer. bor. à Novâ-Cæsareâ ad Carolinam. Pursh fl. 2. p. 543. — Herm. parad. t. 243. Capitula ampla multilora. Achenia glabra. (v. s. sp. et v. c.)

♂? *microcephala*, capitulis multo minoribus, ligulis 11-13, disco abortivo.

— Hanc varietatem verisim. monstrosam in horto Parisino lectam à cl. Adr. de Jussieu sub nom. *S. asperulæ* recepi, sed non respondi. phrasi Desf. (cat. p. 302). *S. asperula* mihi manet ignota. (v. s.)

**S. GLOMERATA** (Michx. fl. bor. am. 2. p. 117), caule humili simplicissimo, foliis glabris oblongo-lanceolatis serratis, racemo simplici glomerulis axillaribus superioribus capitato-congestis constante, invol. turgidis multifloris.  $\mathcal{L}$  in montibus Carolinæ (Michx.), prope Salem (Schwein.). Folia infer. ovalia cuminata (Nutt.). Affinis *Asteri* (Nutt.). Distincta serraturis profundis approximatis et racemis capitatis. (Ell. sketch 2. p. 387.)

**S. MACROPHYLLA** (Pursh fl. bor. am. 2. p. 542), caule. . . . , foliis infer. ovatis cuminatis attenuatis inequaliter argute serratis, caulinis lanceolatis utrinque attenuatis subsessilibus serratis, racemis axillaribus pedunculatis foliosis longitudine foliorum, invol. oblongis turgidis multifloris, ligulis subelongatis.  $\mathcal{L}$  in Canadâ. Media inter *Solid.* et *Asterem* (Pursh). Vix à *S. glomeratâ* distincta (Nutt.). †.

**S. ELATA** (Ell. sketch 2. p. 389), caule piloso tereti supernè tomentoso, foliis ovali-lanceolatis acutis subintegerrimis venosis subtus tomentoso-pubescentibus, racemis erectis paniculatis, ligulis 7-10 elongatis.  $\mathcal{L}$  in pinetis Georgianæ et Carolinæ borealis. An *S. elata* Pursh fl. 2. p. 543 omnino eadem?

**S. SQUARROSA** (Nutt. gen. am. 2. p. 161), caule crasso robusto supernè pubescente, foliis levibus, inferioribus latissimis spatulato-ovalibus serratis acutis margine scabris, super. sessilibus lanceolato-ellipticis integris racemis glomeratis rigidis pubescentibus, invol. multifloro squarroso!, ligulis elongatis 10-12.  $\mathcal{L}$  in rupestribus Novæ-Cæsareæ (Nutt.), Pennsylvaniae et Georgiæ (Muhl.). sed fortè ex Ell. sk. 2. p. 389, stirps Georgica differt à specie Nuttalianâ verâ Affinis dicitur *S. strictæ*. Achenia glabra.

**S. CARINATA** (Schrader in litt. 1832), caule erecto apice angulato subpubescente, foliis radic. longè petiolatis oblongis basi attenuatis apice obtusis et mucronatis mucronato-serratis, super. sessilibus linearibus subserratis, omnibus glabris margine scabris nervo medio subtus prominulo, racemis erectis in paniculam dispositis, ligulis 7-9 disco longioribus.  $\mathcal{L}$  patr. ign. verisim. Amer. bor. Folia radicalia petiolo incluso 7 poll. longa, 8-10 lin. lata. Pedicelli pauculi bracteolati. (v. s. c. ex hort. Goett.)

75. *S. FUSCATA* (Desf.! cat. h. Par. ed. 3. p. 402), glaberrima, caule erecto fusco lævi, foliis lanceolatis utrinque integerrimis, racemis erectis, pedicellis brevibus bracteolatis, invol. squamis oblongis glabris vix acutis, ligulis linearibus disco paulo longioribus.  $\frac{1}{2}$  in Amer. bor. ? Ligulæ 5-6, fl. disci 6-7. Habitus ferè *S. ambigua*, sed caulis rubescens, folia integerrima, etc. (v. v. c. in hort. Par.)
76. *S. VIMINEA* (Ait. kew. ed. 1. v. 3. p. 215), caule erecto subpubescente, foliis lineari-lanceolatis membranaceis basi attenuatis glabris margine scabris, infimis suberratis, racemis erectis, ligulis elongatis.  $\frac{1}{2}$  ad ripas fluminum Amer. bor. à Canadâ ad Virginiam (Pursh fl. 541). *S. integerrima* Mill. dict. ex Ait. Affinis *S. limonifolia* sed caulis subpubescens, folia non carnosâ.
77. *S. LIMONIFOLIA* (Pers. ench. 2. n. 36), glaberrima, caule erecto-obliquo, foliis subcarnosis integerrimis lævibus, infer. ellipticis basi longè attenuatis, cæteris lanceolatis sessilibus semiamplexicaulibus, racemis paniculatis erectis, pedunculis squamosis glabris, invol. squamis linearibus acutis, ligulis elongatis.  $\frac{1}{2}$  in salsis et ad ripas rivulorum Amer. bor. (Willd.), in Carolinâ (Ell.), nec in Mexico. (Nutt. gen. am. 2. p. 150). *S. Mexicana* Linn. sp. 1234. *S. obliqua* Moench. — Dod. icon. 4. t. 219. Pluk. alm. t. 235. f. 2. malè. Nomen *Limonifolia* veteris admissum jam præeuntibus Nutt. 2. p. 159 et Ell. 2. p. 380 recepit, cum planta in Mexico non crescat. Affinis *S. sempervirens* sed racemi non secundi, *S. lævigata* sed pediculi glabri. (v. v. c.)
78. *S. LÆVIGATA* (Ait. kew. ed. 1. v. 3. p. 215), caule erecto lævi, foliis sessilibus lanceolatis carnosis integerrimis nudique lævibus, racemis paniculatis erectis, pedunculis squamosis villosis, invol. squamis linearibus acuminatis, ligulis angustis elongatis.  $\frac{1}{2}$  in Amer. bor. paludosis salis à Canadâ ad Virginiam Pursh fl. 2. p. 541. Affinis *S. limonifolia* sed si specim. meum cum *Aitoniana* specie omnino congruit, facile differt foliis triplinerviis et notis indicatis. (v. v. c. ex hort. Hafn.)
79. *S. INTEGRIFOLIA* (Desf.! cat. h. Par. ed. 1804. p. 103. ed. 3. p. 402. n. Roxb.), caule erecto simplici puberulo, foliis lanceolatis sessilibus acuminatis tenuè et obsolete triplinerviis glabris, infer. parè denticulatis, cæteris integerrimis, racemis axillaribus erectis subpaniculatis, pedicellis puberis, invol. squamis acutis, ligulis elongatis.  $\frac{1}{2}$  in Amer. bor. (Desf.). Pers. ench. 2. p. 449 n. 35. ex hort. Par. An *S. speciosa* Nutt. gen. am. 2. p. 160? et ex eo *S. sempervirens* Michx. ? Affinis *S. lævigata* et *limonifolia*. (v. v. c.)
80. *S. PUBESCENS* (Ell. sketch 2. p. 381), caule erecto ramoso pubescente, foliis longè lanceolatis basi attenuatis pubescentibus, inferioribus serratis, racemis erectis paniculatis, ligulis 7-10 mediocribus.  $\frac{1}{2}$  in humidis Georgiæ circa Louisville. Achænia minutè pubescentia. Habitus *S. speciosa* Nutt. (*integrifolia* Desf. ?)
81. *S. PANICULATA*, glaberrima. caule erecto apice multiramoso paniculato, foliis lineari-lanceolatis acuminatis sessilibus integerrimis margine scabris triplinerviis, venis lateralibus nervo medio longè parallelis, racemis erectis, pedicellis puberulis, invol. squamis linearibus subacutis, ligulis disco paulo longioribus.  $\frac{1}{2}$  in Mexico circa urbem. *S. Mexicana* H. B. et Kunth nov. gen. am. 4. p. 104? Allam. ! in litt. An *Virga-aurea Mexicana* Benth. pin. 51? Certè à *S. limonifolia* diversa. (v. s.)
82. *S. STRICTA* (Ait. kew. ed. 1. v. 3. p. 216), caule erecto glabro, foliis elongato-lanceolatis acutis glabris margine scabris triplinerviis, radic. basi longe attenuatis serratis, super. integerrimis, racemis paniculatis erectis, pedunculis glabris, invol. squamis acutis, ligulis elongatis.  $\frac{1}{2}$  in Americæ borealis sive arenosis à Novâ-Cæsareâ ad Carolinam (Pursh fl. 2. p. 540). Herba 2-3-pedalis stricta parè foliosa. (v. v. c.)
83. *S. STRICTA* (Pursh fl. 2. p. 542), caule erecto tereti basi glabro apice ter pedunculis pubescente, foliis lanceolatis subæqualibus glabris venosis integre margine fimbriato-scabris, racemis erectis brevibus confertis, invol. squamis

linearibus obtusiusculis, ligulis 5-7 oblongis disco vix longioribus.  $\frac{1}{2}$  in humida Carolinæ. Nutt. gen. 2. p. 161. Ell. sketch 2. p. 385. *S. viminea* Bosc! herb. Species botan. amer. ipsa dubia. Specim. meum desor. Ell. sat convenit, à descr. Nutt. differt foliis minoribus sed folia omnia Sphæria quadam parasitica deformata. (v. s. ex h. Bosc.)

84. *S. ANGUSTIFOLIA* (Ell. sketch 2. p. 388. non Mill.), caule erecto glabro, foliis subulato-linearibus integerrimis glabris, racemis erectis paniculatis, invol. squamis acutis lineari-lanceolatis, ligulis oblongis 7-10 disco paulò longioribus.  $\frac{1}{2}$  in insulâ Paris circa Beaufort (Ell.) et in Georgiâ (h. Duby). Affinis *S. vimineæ*. Rami graciles elongati. Folia sup. erecta. (v. s.)

85. *S. LINEARIFOLIA*, caule (ramisve) tereti puberulo, foliis linearibus acutis sessilibus integerrimis tenuè trinerviis glabris margine ciliolato-scabridia, racemis brevibus erectis in paniculam racemiformem confertis, invol. squamis linearibus glabris subscariosis, floribus circ. 20, ligulis 7-8 vix disco longioribus. — in campis Chilensibus dictis *Pampas* olim legit cl. Née. Ex specim. non satis apparet an sit fruticulosa sen herbacea. Invol. pallidè rufescens. Achænia glabra. (v. s.)

✧? *Pæppigii*, caule glabro, foliis margine magis scabrido-ciliatis, paniculæ ramis magis distantibus, capitulis minoribus. — in Amer. austr. legit cl. Pæppig et ex ejus coll. recepti sub. n. CCXXVII sine loci designatione. An spec. propr.? (v. s.)

86. *S. DOMINGENSIS* (Spreng. ! syst. 3. p. 639), caule erecto subangulato glabro apice paniculato, foliis lato-linearibus rigidiusculis integerrimis obsolete trinerviis lævibus, racemis strictè erectis in paniculam dispositis foliosis, ligulis paucis minutissimis.  $\frac{1}{2}$  in Sancto-Domingo legit cl. Bertero. Folia apice callosa sesquipollicem longa, 2 lin. lata, juniora fortè viscosa. Invol. glaberrima pauciflora. (v. s.)

Sect. II. EUTHAMIA. — *Euthamia* Nutt. gen. am. 2. p. 162. Cass. dict. 37. p. 471.

Capitula oblonga. Invol. imbricata squamis subagglutinatis. Flor. ligulati 10-20 nempe floribus disci numerosiores. Recept. fimbriiferum. Achænia oblongo-ovoidea villosa. — Herbar., capitulis ad apices ramulorum sessilibus adgregatis in corymbum dispositis. Folia linearia integerrima. — An genus proprium ut innuit Nuttall et asserit Cassini? Habitus medius hinc inter Solidaginem et Nidorellam, illinc Brachyridem et Bigelowiam.

87. *S. GRAMINIFOLIA* (Ell. sketch 2. p. 391), caule erecto glabro angulato ramossissimo, foliis lanceolato-linearibus 3-5-nerviis secus nervos et margines scabriusculis, axillis nudis, corymbis terminalibus fastigiatis, fl. disci 8-10, ligulis 15-20 longitudine disci.  $\frac{1}{2}$  in agris et pascuis ad latera sylvarum et flum. Americæ bor. à Canadâ ad Pensylvaniam (Pursh), in Carolinâ (Ell.), Novæboraco (Torr.!). *Chrysocoma graminifolia* Linn. sp. 1178. *Solidago lanceolata* Willd. sp. 3. p. 2060. *S. lanceolata* α major Michx. fl. 2. p. 116. *Euthamia graminifolia* Nutt. gen. am. 2. p. 162. *Chrysocoma Virginiana* olim in h. Par. (v. v. c. et s.)

88. *S. TENUIFOLIA* (Pursh fl. bor. am. 2. p. 540), caule erecto subscabro angulato corymboso-ramoso, foliis angustissimè linearibus patulis 1- sen rariùs 3-nerviis scabris, axillis foliosis, corymbis terminalibus fastigiatis, fl. disci 5-6, ligulis 10 disco paulò longioribus.  $\frac{1}{2}$  in pinetis Amer. bor. à Novâ-Cæsareâ ad Floridam (Pursh), in pascuis siccis Carolinæ (Ell. sketch 2. p. 392) et Georgiæ (h. Duby!). *Euthamia tenuifolia* Nutt. gen. am. 2. p. 162. *Sol. lanceolata* β minor Michx. fl. bor. am. 2. p. 540. Caulis humilis. Folia atomis resinosis tecta. (v. s.)

† *Species non satis notæ.*

89. *S. CANTONIENSIS* (Lour. coch. ed. W. 2. p. 612), caule herbaceo simplicissimo, foliis lanceolatis sessilibus integerrimis glabris, pedunculis polycephalis



Nepalensis Don prod. nep. p. 177. Herba stricta variat nunc 4 poll. oligocephala, nunc bipedalis polyecephala.

\*\* *Capitulis solitariis longè pedunculatis.*

6. A. PEDUNCULARIS, caulibus herbaceis pluribus erectis subsimplicibus basi angulatis glabris supernè teretibus hispidis, foliis longè petiolatis ovatis subcordatis acuminatis integerrimis suprà petioloque setoso-hispidis margine ciliatis subtùs pallidis glabrescentibus, pedunculo elongato hispidulo nudo 1-cephalo, invol. squamis paucis foliaceis oblongis dorso hispidis, ligulis circ. 20 linearibus, achæniis villosis.  $\frac{1}{2}$  in Indiâ orientali ad Kamaon. Aster peduncularis Wall. ! cat. n. 2967. comp. n. 77. Cineraria ovata Spreng. in litt. ad Wall. Flores ex sicco vid. flavi. Species distinctissima à prioribus habitu sat diversa Rhizoma vid. repens. (v. s. comm. ab hon. Cæta merc. angl. Ind. or.)

CLXXXV. COMMIDENDRUM DC. in Guill. arch. bot. 1833. v. 2. p. 334. — Conyzæ et Solidaginis spec. Roxb.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci tubulosis hermaphr. 5-dentatis. Recept. nudum angustum. Invol. imbricatum, squamis linearibus pauciseriatis acuminatis. Anthere ecaudate appendiculatæ. Stylus Asterearum. Achænium compresso-planum erostre glabrum. Pappus 1-serialis pilosus scaber. — Arbusculæ Helenianæ gummiferæ (et ideò *Gum-wood* angl. dictæ, unde cl. Burchell nomen Commidendri finxit). Rami cicatricibus post foliorum lapsum crassis callosis notati. Pedunculi corymbosi polycephali ex axillis supremis aut apice ramorum orti. Flores albi.

§. 1. *Foliis glabris, pedunculis apice ramosis corymbiferis. — Solidaginis sp. Roxb.*

1. C. ROTUNDFOLIUM, foliis latè obovatis basi cuneatis subdentatis longè petiolatis utrinquè glaberrimis, junioribus vernicosis, corymbis compositis polyecephalis, pedunculis apice ramosis, pedicellis circ. capituli longitudine.  $\frac{5}{8}$  in insulâ Sanctæ-Helenæ. Solidago rotundifolia Roxb. in Beate. app. p. 324. *Bastard cabbage, Gum-Wood et Bastard-Tree* vern. angl. Petioli limbo sublongiores. Ligulæ 3-10. Capitula parva. (v. s. comm. à cl. Burchell et Wallich.)
2. C. SPURIMUM, foliis cuneato-lanceolatis basi attenuatis integris subsimilibus apice serratis utrinquè glabris, corymbis polycephalis compositis, pedunculis apice ramosis, pedicellis ideò capitulo vix longioribus.  $\frac{5}{8}$  in insulâ Sanctæ-Helenæ. Conyza arborescens Forst. comm. Gært. g. p. 68? Willd. sp. 3. p. 2055 excl. Ait. syn. Folia suprema 2 poll. longa, 8 lin. lata. Ramuli vix subpauciruli. Ligulæ 5-6. Fl. tubulosi 6-7. (v. s. comm. à cl. Burchell.)

§. 2. *Pedunculis axillaribus à basi ramosis in pedicellos valde elongatos 1-cephalos aut simplicibus. — Conyzæ sp. Roxb.*

3. C. GUMMIFERUM, foliis cuneato-lanceolatis basi integris attenuatis subpetiolatis apice serratis subrugosis junioribus suprà sparè puberulis subtus tomentosis, pedunculis axillaribus ferè à basi in pedicellos valde elongatos monoecephalos divisis.  $\frac{5}{8}$  in insulâ Sanctæ-Helenæ montanis. Conyza gummifera Roxb. in Beate. app. p. 304. *Gum-Wood-Tree* vern. angl. Arbor ramis 2-3-chorum. Folia ad apices ramorum conferta, juniora gummitera Roxb. Pedicelli foliis breviores. (v. s. comm. à cl. Burchell.)
4. C. ROBustum, foliis cuneato-lanceolatis basi attenuatis integris subpetiolatis apice crenato-dentatis rugosis suprà araneosis subtus tomentosis, pedunculis ex imâ basi in pedicellos elongatos 1-cephalos divisis, involucro arachnoideo  $\frac{5}{8}$  in insulâ Sanctæ-Helenæ prope Longwood. Conyza robusta Roxb. in Beate

app. p. 304. Arbor 20-pedalis. Capitula paulò majora et pauciora prioris ad quam valde accedit. (v. s. comm. à cl. Burchell et Wallich.)

5. *C. mucosum*, foliis cuneatis ferè petiolatis acutis apice grossè serratis suprà pubescentibus subtùs hirsuto-villosis reticulato-rugosis, pedunculis axillaribus 1-cephalis hirsutis simplicibus folii circ. longitudine, invol. squamis glabris linearibus.  $\mathfrak{J}$  in insulâ Sanctæ-Helenæ. *Conyza rugosa* Ait. kew. 3. p. 184 non Vahl ed. 2. v. 5. p. 30 excl. syn. Willd. *Aster glutinosus* Roxb. in Beats. app. p. 303 ex Burch. in litt. (v. s. in h. Deless. olim à fratre suo in hort. Kew. lect. et nom. Aitoniano insig.)

## CLXXXVI. STEIRACTIS.

Omnia *Solidaginis* sed ligulæ steriles. — Frutex arboreus Novo-Zeelandicus. Folia petiolata ovali-orbiculata glabra repanda subtùs albedo-purpurea. Capitula paniculata. Invol. subcylindraceum, squamis linearibus imbricatis margine membranaceis. Ovarium teres pilosiusculum. Pappus flavicans fimbriato-pilosus. (Car. ex Rich.)

1. *S. ARBORESCENS*.  $\mathfrak{J}$  in Novâ-Zeelandiâ legit Forster. *Solidago arborescens* Forst. prod. 298. Rich. fl. Nov.-Zeel. 252.

## CLXXXVII. ROCHONIA.

Capitulum multiflorum radiatum, ligulis 1-serialibus homochromis fœmineis sterilibus, fl. disci tubulosis hermaphr. 5-dentatis. Invol. imbricatum, squamis subscariosis linearibus 1-nerviis. Recept. alveolato-fimbrilliferum. Styli rami elongati semiteretes glabriusculi. Achænia teretiuscula subvillosa. Pappus setosus 1-serialis subscaber. — Frutices Madagascarienses ramulis foliisque subtùs canis. Folia alterna sessilia coriacea integerrima 1-nervia. Capitula pauca aut solitaria ramulos terminantia. Flores flavi. — Genus affine *Aplopappo* dicavi cl. Alex. Rochon physico Gallo qui insul. Madagascar adiit et descripsit. (Conf. Roch. voy. Par., in-8. 1791. et vol. 3. 1802.)

1. *R. CINERARIOIDES*, foliis ellipticis suprà glabris læviusculis subtùs cano-tomentosis subreticulatis, capitulis 7-8 pedicellatis in racemum dispositis.  $\mathfrak{J}$  in savannis interioribus insulæ Madagascar legit cl. Bojer. *Conyza cinerarioides* Bojer in litt. 1835. Frutex circ. orgyalis. Invol. squamæ extimæ tomentosæ, int. scariosæ flavidæ glabriusculæ. Folia 7-9 lin. longa, 4 lin. lata. Pedicelli tomentosi. (v. s. comm. à cl. Bojer.)

*ℓ. subreticulata*, foliis ovatis supernè reticulo vix elevato notatis, capitulis racemoso-corymbosis.  $\mathfrak{J}$  cum priore legit cl. Bojer. An fortè species distincta? (v. s.)

2. *R. CUNEATA*, foliis cuneatis suprà glabris nervulis elevatis reticulatis subtùs adpressissimè canis, capitulis terminalibus solitariis.  $\mathfrak{J}$  in savannis interioribus insulæ Madagascar legit cl. Bojer. *Conyza cuneata* Bojer! in litt. 1835. Frutex octopedalis. Folia 10 lin. longa, 3 lin. lata. (v. s. comm. à cl. Bojer.)

CLXXXVIII. *APLOPAPPUS* Cass. dict. 56. p. 169. — *Diplopappus* Less. in *linnæâ* 1831. p. 110. — *Diplopappi* et *Doniæ* sp. auct.

Capitulum multiflorum radiatum, ligulis 1-serialibus fœmineis (in unic. sp. nullis), fl. disci hermaphroditis 5-dentatis. Recept. planiusculum soveolatum seu alveolato-subfimbrilliferum. Invol. squamæ imbricatæ sublineares acutæ. Achænia oblonga teretiuscula aut turbinata densè sericea rarius glabra. Pappus setosus uni-aut pluriserialis subinæqualis sed setis inter se conformibus. — Herbæ aut suffrutices

14. *A. serotoculatus*, fruticulosus, ramis procumbentibus striatis, foliis spatulatis indivis dentatis, pedunculis elongatis filia parva pappi angusta serratis, invol. squamis lanceolatis serotoculatis canaliculis glutinosis - v. n. achaniis strigosis-sericeis.  $\frac{1}{2}$  in Cordilleris Chilensibus. Adm. glutinosus Poir. et Less. *Diplopappus serotoculatus* Nees. det. p. 284.
15. *A. Berterii*, caule suffruticuloso adscendente striato glabro apice apicul. foliis subdistansibus obovatis basi cuneatis apice obtusis margine serratis ciliatis et grasse serratis tenue reticulatis, capitulo terminali erecto. invol. squamis linearibus acuminatis margine membranaceis ciliatis, achaniis sericeo-villosissimo.  $\frac{1}{2}$  in petronis et rupestribus maritimis prope Valparaiso Chilensium legit cl. Bertero. *Grindelia glutinosa* Bert. Merc. cat. 1799. n. p. 646. non Donal. Capitulum fere *A. macrocephali*. Ligula glabra n. n. *A. Chamissonis* medio pilosa. Caulis apex subviscosus glaber. v. s. comm. beat. et amic. Bertero.
- $\frac{1}{2}$  lanceolatis, caule sub capitulo puberulo, foliis lanceolatis basi cuneatis apice acutiusculis intensius reticulatis, invol. squamis subpuberulis et magnioribus.  $\frac{1}{2}$  in rupestribus maritimis prope Valparaiso legit Bertero et cum priore sub n. 1038 cum sign. comm. Folia fere *A. pulchelli* et capitulum *A. Berterii* aut *macrocephali*. v. s.
16. *A. macrocephalus*, herbaceus, caule simplici erecto glutinoso tereti densae foliis apice apicul. 1-cephalo. foliis subcoriacis spatulatis oblongis serratis margine hispidiusculis reticulato-venosis, invol. late campanulatis squamis linearibus mucronato-acuminatis viscosis hispidiusculis, achaniis sericeo-villosissimo.  $\frac{1}{2}$  in Chili Porpp., prope Rancagua in monte La Leca (Bertero). *Diplopappus macrocephalus* Porpp. ex Less. in linnæa 1831. p. 114. Caulis 10-pollic. Pappus flavescens. Corollæ glabræ luteæ. v. s. comm. et Bertero sub n. 315.
17. *A. grindelioides*, herbaceus, caulibus basi ramosis teretibus apice subapophyllis hirsutis, foliis sessilibus subsemiamplexicaulibus obovatis acutis serratis glabriusculis, capitulis solitariis, invol. campanulatis squamis linearibus acuminatis subpuberulis, ligulis elongatis, achaniis hirsuto-canaliculatis.  $\frac{1}{2}$  in Chili legit cl. Porppig. *Diplopappus grindelioides* Less. in linnæa 1831. p. 115. Capitulum fere *Grindelia*. Pappus sordidus. (v. s. comm. et cl. Porppig sub n. 916.)
18. *A. chrysanthemifolius*, herbaceus?, ramis striatis florigeris apice subundulatis, foliis sessilibus membranaceis glabris subciliatis, infer. obovatis, super. lanceolato-linearibus mucronato-serratis et summis integerrimis, capitulis solitariis, invol. campanulatis squamis linearibus acuminatis glaberrimis, achaniis bracteato-villosis.  $\frac{1}{2}$  in rupibus maritimis prope Concon Chilensium. *Andromeda alternifolia* Kunze in Porpp. coll. ex. 1. n. 243. *Diplopappus chrysanthemifolius* Less. in linnæa 1831. p. 116. Pappus rufescens. Cor. glabræ, tubularum hispidulo.
19. *A. hispidulus*, fruticulosus, ramis teretibus, junioribus foliisque pube breviter confertis hispidulis, adultis glabris, foliis linearibus confertis linearibus obtusis integerrimis, ramis apice apophyllis 1-cephalis, invol. squamis linearibus acuminatis hispidulis, achaniis parvis sericeo-villosis.  $\frac{1}{2}$  in Chili prope Coquimbó legit cl. Gandichaud. Folia pollicem fere longa, semilineam lata, minoribus axillis glomerata. Pedunculus pollicaris. Ligula angustæ flavæ. Accedit ex foliis ad sect. iv, ex ligulis ad sect. v. (v. s. comm. et cl. Gandichaud sub n. 87.)

## Sect. IV. LEUCOPSIS.

Achania villosa aut hirsuta elongata. Pappus corollis disci longior. — Folia integra villosa-canescuntia. — An genus proprium?

20. *A. acuminatus*, herbaceus, totus sericeo-lanatus, caule erecto folioso, foliis lanceolatis acuminatis integerrimis, infer. obovato-oblongis obtusis. Ligula terminali, invol. campanulatis squamis laxè imbricatis dorsis

lanatis linearibus acuminatis, ligulis 25-30 invol. longioribus, achæniis albo-villosissimis.  $\mathcal{L}$  in Brasiliæ prov. Rio-Grande et fortè Sancti-Pauli (h. Mus. imp. Bras. n. 1041 et fortè 439?). Priori simillimus sed inv. squamis foliisque superioribus valdè acuminatis faciliè differt. In specim. prov. Sancti-Pauli pars inferior plantæ glabrescit. (v. s. in h. Mus. reg. Par.)

21. A. ? *sericeus*, herbaceus totus sericeo-lanatus, caule adscendente toto folioso, foliis oblongis acutis basi attenuatis integerrimis, capitulo solitario terminali, invol. campanulati squamis laxè imbricatis siccis linearibus obtusiusculis, ext. subvillosis, cæt. glabris siccis, ligulis vix invol. longioribus, achæniis elongatis hirsutis.  $\mathcal{L}$  in pascuis sabniosis ad Quintero Chilensium legit cl. Bertero (pl. exs. n. 1411) et ad Talcaguano (Cham.). *Diplopappus sericeus* Less. in linn. 1831. p. 110. Cor. flavæ. Folia pollicem longa, 2-3 lin. lata. Pappus pluriserialis rufescens setis ext. brevioribus. An *Noticastro* affinis? (v. s.)
22. A. ? *canescens*, totus adpressè sericeo-villosus canescens, caulibus fruticulosus è collo plurimis apice subramosis, foliis oblongo-linearibus basi attenuatis integerrimis mucronato-acutis infernè setis rigidis paucis ciliatis, ramulis brevibus 1-cephalis, inv. campanulati squamis linearibus acutis subcanescentibus, ligulis paucis invol. paulò longioribus, achæniis villosis pappo dimidio brevioribus.  $\mathcal{L}$  in Mexicanæ prov. Texas districtibus orientalibus nov. et dec. flor. legit cl. Berlandier (pl. exs. n. 1877). Affinis A. sericeo sed capitula triplò 4-plòve minora, pappus pallidè rufescens, achænia etiam ratione capituli multò breviora etc. (v. s. comm. à cl. Berlandier.)
23. A. ? *diversus*, suffruticosus sparsè pubero-villosus, ramis adscendentibus foliosis, foliis spathulato-sen oblongo-linearibus mucronato-acuminatis subintegerrimisqne scabris, capitulis solitariis, invol. squamis lanceolatis laxis, achænio adpressè villosis.  $\mathcal{Z}$  in Monte Video (Pers.), et Rio-Grande (h. Mus. imp. Bras. n. 1002, 1008, 1010, 1022, 1033!). *Erigeron diffusus* Pers. ench. 2. p. 431. *Onoseris Montevidensis* Spreng. syst. 3. p. 502. Affinis A. sericeo sed pube totius plantæ sparsior, caulibus fruticulosus ramosior, folia scabra margine scaberrima, Less. in linn. 1831. p. 110. (v. s. in h. Mus. reg. Par.)

#### Sect. VI.? PYROCHÆTA.

Capitulum radiatum, ligulis 1-serialibus angustissimè linearibus neutris seu styliferis sterilibus, fl. disci tubulosos brevissimè 5-dentatis herm. fertilibus. Invol. imbricatum, squamis linearibus pluriserialibus intimis disco circ. equalibus. Achænia radii abortiva, disci turbinata sericeo-villosissima. Pappus radii setis paucissimis, disci multisetus scaber rigidulus. — Herba Mexicana annua erecta. An genus proprium?

24. A. ? *Hænkii*, radice exili, caule herbaceo simplici aut parcè ramoso glabro, foliis oblongo-cuneatis apice serraturis paucis dentatis junioribus subtus araneosis demùm glabris.  $\text{\textcircled{U}}$  inter plantas Regiomontanas herb. Hænkeani ab ill. de Sternberg ad studium missi conservatur. Herba pedalis. Folia remotiuscula pollicem longa, 3 lin. lata. (v. s. in h. suprà dicto.)

#### Sect. V.? INULOPSIS.

Achænia villosa brevia teretiuscula. Ligulæ lineares numerosissimæ. — Herba Mexicana et Brasiliensis, habitu Inulæ. Fortè genus proprium? imò species duæ in genera duo fortè divellendæ?

25. A. ? *stoloniferus*, caulibus è collo plurimis debilibus parcè foliosis basi hispidulis, aliis sterilibus decumbentibus, aliis 1-cephalis apice villosis-tomentosis et glanduloso-pilosis, foliis radicalibus obovatis hirsutissimis subdentatis, secus caules floridos lanceolatis acutis, secus caules steriles obovato-oblongis subobtusis, invol. squamis linearibus acuminatissimis disco longioribus hispidis.  $\mathcal{L}$  in Mexici summo monte San-Felipe supra locum ubi ex structurâ glaciei apotheca legit cl. Andrienx. Inula stolonifera Gay! in Andr. coll. exs. u. 322. Ligulæ numerosissimæ 1-seriales lineares. Cor. tubulosæ  $\infty$ . Anthere basi nudæ. Invol. subpurpureus. Flores intimi abortu masculi. Achænia juniora villosa, adulta ign. (v. s.)

\* *Herbaceæ.*

1. *L. VULGARIS* (Cass. ex Less. syn. 195), foliis linearibus glabris impunctatis, capitulis corymbosis, invol. squamis appendiculatis apice patulis squarrosis.  $\mathcal{L}$  in Europæ mediæ et australis et imò Tauriæ et Oelandiæ apriciæ rupestribus et argilosis. *Chrysocoma Linosyris* Linn. sp. 1178. DC. fl. fr. 4. p. 141. All. ped. t. 11. f. 2. Schkuhr handb. t. 238. *Linosyris foliosa* Cass. ex Desf. cat. h. Par. p. 196. *Crinitaria Linosyris* Less. syn. 195. *Linosyris nuperorum* Lob. hist. 223. *Aster Linosyris* Bernh. ex Steud. nom. phan. 193. *Erigeron Linosyris* Clairv. man. herb. val. 243. (v. v. et s.)

$\beta$ . *minor* (Wallroth! in litt.), caule 3-5 pollicari.  $\mathcal{L}$  in siccis Thuringiæ, Galliæ, C. *Linosyris tripollicaris* Vill. dauph. 3. p. 188. DC. fl. fr. n. 3130. Culta dimensiones majores mox acquirit (Savi. pug. p. 11). (v. s.)

$\gamma$ . *patula*, foliis patentibus vel reflexis.  $\mathcal{L}$  in collibus calcareis Alsatiæ super. (Nestl.!) et Jurassi Neocomensis (Chaill.!). — An in paludosis Etruriæ circa Stagno et Liburnum si *Chrys. palustris* Savi pug. p. 11. et *Chrys. Liburnica* Spreng. syst. 3. p. 475 huc referendæ? (v. s.)

$\delta$ ? *verticalis*, foliis lanceolatis verticalibus.  $\mathcal{L}$  ad rupium fissuras in Hispaniæ regnis Murcico, Granatensi et Valentino. *Chrysocoma verticalis* Lag. nov. gen. p. 15. n. 330.

2. *L. PUNCTATA* (Cass. dict. 37. p. 476. sub *Crinitaria*), foliis linearibus aut lineari-lanceolatis trinerviis punctatis subscabris, capitulis corymbosis, invol. laxo.  $\mathcal{L}$  in campis apricis siccis aut subsalsis Sibirici ad Irtim (Patr.!), Ural (Fisch.), in Phrygiâ (Parol.!), in campis ad Wolgam (Bieb. Stev.). *Aster acris* Gmel. itin. 1. p. 172. t. 38. f. 2. *Chrysocoma biflora* Linn. sp. 1178 (nomen omn. decipiens). *Chrysocoma dracunculoides* Lam. dict. 2. p. 192. Bieb. taur. 2. p. 288. *Crinita punctata* Moench. meth. 578. *Crinitaria biflora* Less. syn. 195. — Gmel. Sib. 2. t. 82. f. 1. Variat foliis angustissimis 1-nerviis et latioribus 3-nerviis. Cum *Astere* quando confudit Linnæus quando variare cum et sine radio. (v. v. c. et sp.)

3. *L. VILLOSA*, foliis lanceolatis adpressè villosis, capitulis paucis subcorymbosis. involucri squamis adpressis margine scariosis.  $\mathcal{L}$  in apriciis Hungariæ, Caucasi, Sibirici. *Chrysocoma villosa* Linn. sp. 1178. Waldst. et Kit. pl. hung. 1. t. 54. *Conyza oleaefolia* Lam. dict. 2. p. 86. — Gmel. Sib. 2. t. 82. f. 2. (v. s.)

4. *L. DIVARICATA*, herbacea incana, foliis infer. lanceolatis trinerviis, super. lineari-subulatis diminutis, ramis 1-cephalis squamatis virgatis, invol. laxo incanis squamis acutis.  $\mathcal{L}$  in campis siccis circa Astracan ex Gmel. *Chrysocoma divaricata* Fisch. cat. gor. 1816. p. 37. Bieb. suppl. 563. — Gmel. Sib. 2. p. 193. t. 82. f. 3. Minus pubescenti-cana quam *L. villosa*.

\*\* *Fruticosa*, recept. sæpius punctato nec alveolato. (Plerumque minus nota.)

5. *L. CANDICANS*, caule suffrutescente diffuso, foliis linearibus sericeo-pubescentibus margine replicatis, capitulis paniculis confertis 5-9-floris, invol. squamis adpressis apice coloratis.  $\mathcal{L}$  in Ægypto propè Alexandriam. *Chrysocoma candicans* Delil. fl. eg. 127. t. 46. f. 2. Invol. cylindricum nec punctatum nec in prioribus foveolatum.

6. *L. HASTATA*, fruticosa, foliis oblongis sessilibus ad medium lobo horizontalibus hastatis incanis, capitulis solitariis, invol. imbricati squamis acutis.  $\mathcal{L}$  in Arabiâ felici. *Chrysocoma spathulata* Forsk. descr. æg.-ar. 147. *Stachelina hastata* Vahl symb. 1. p. 70. *Pappus biserialis* ex Less. syn. 195.

7. *L. MONTANA*, fruticosa, foliis oblongis villosis integerrimis, capitulis solitariis, invol. oblongis squamis ext. obtusis, int. acutis apice patulis  $\mathcal{L}$  in mont. Horeb Arabiæ et in Palestinâ si pl. n. 440 herb. Novè huc rite referenda *Chrysocoma montana* Vahl. symb. 1. p. 70. *Pappus rufus biserialis* ex Less. l. c. (v. s.?)

8. *L. UNIFLORA*, fruticosa, ramis glabris erectis, foliis linearibus complicatis

utis, pedunculis 1-cephalis, invol. imbricato.  $\delta$  in Egypto. *Chrysocoma afflora* Spreng. nov. prov. 1818. p. 11. Recept. nudum.

1. *GLABRATA* (Lindl. obs. mss. 1835, sub *Crinitaria*), foliis linearibus obtusis punctatis glabris, capitulis corymbosis, invol. obconici squamis ext. pubescentibus, inter. dilatatis obovatis glabris.  $\frac{1}{2}$  in Sibiria Altaica deserto Sonoro-Kirghisico. *Aster dracunculoides discoideus minor* Ledeb. fl. alt. 4. p. 98. species mihi ignota et ex cl. Lindley admissa.

2. *CHRYSOCOMA* Cass. dict. 37. p. 477. Less. syn. 195. — *Chrysocoma* sp. Linn. Thunb. — *Chrysocoma* et *Crinita* sp. Manch.

capitulum homogamum discoideum multiflorum. Recept. nudum alveolatum. Invol. campanulati floribus brevioris squamæ oblongo-ovolatæ imbricatæ. Achænia lateraliter plano-compressa erostria vix hispidula. Pappus setaceo-pilosus 1-serialis. — Frutices Capenses. a linearia sparsa sæpius integra. Rami apice plus minus denudati. itula subglobosa. Flores lutei. — Species recognoscendæ!

### §. 1. *Fruticosa, foliis linearibus, capitulis subsolitariis.*

1. *COMA-AUREA* (Linn. sp. 1178), foliis sparsis linearibus patulis rectis aut brecurvis ramisque glabris aut junioribus subpubescentibus.  $\delta$  ad Cap. Bonæ spei frequens. C. patula Berg. cap. 23? *Crinita linearifolia* Maench suppl. 17. Pluk. alm. t. 327. f. 2. Volk. fl. norib. t. 148. C. aurea Thunb. fl. cap. 626. vixat ramis plus minus divergentibus, omnino glabris aut apice summo pubescentibus; foliis erectiusculis patulive, glabris aut jun. pubescentibus; ramis sic longe aut breviter denudatis; capitulis erectis aut cernuis 30-40-floris, d hæ variationes adeo inter se commixtæ apparent ut vix in statu scientiæ resenti seq. var. sint distinctæ. (v. v. c. et sp.)

2. *patula*, ramis divaricatis.  $\delta$  Chr. patula Linn. mant. 28. Thunb. fl. cap. 626. (v. s.)

3. *cernua*, capitulis nubilibus cernuis  $\delta$  Chr. cernua Linn. sp. 1177. Thunb. fl. cap. 626. — Comm. hort. Amst. 2. p. 89. t. 45. (v. s.)

4. *ciliata* (Linn. sp. 1177), foliis sparsis linearibus patulis rectis serrulato-ciliatis, ramis pubescentibus vix apice nudis  $\delta$  ad Cap. Bonæ-Spei. Chr. ciliaris Willd. sp. 3. p. 1789. non Link. — Icon Comm. hort. 2. t. 48 habitum exhibit, sed cilia foliorum nec in descr. nec in icone indicantur, et folia (in icone nigra) in descr. apice trifida dicuntur. (v. s.)

5. *tenusifolia* (Berg. cap. 235), foliis sparsis lineari-subulatis patulis brevissimis ramisque glabris aut subpuberulis, capitulis breviter pedicellatis circ. 10-floris.  $\delta$  ad Cap. Bonæ-Spei in montibus ad Los-Tafelberg (Drege!); in orcester, Swellendam, Uitenhage (Eckl. et Zeyh.). Chr. microphylla Thunb. cap. 626? Olim recepi sine nom. à cl. Thunb. Folia 1-3 lin. longa, juniora reticulata dein patula, subæcero. Rami folia minima fere usque ad capitulum gerentes numerosi corymbosi. (v. s. sp.)

6. *microcephala*, glabra, foliis linearibus subpatulis acutis, ramis floriferis nigatis ad capitulum usque foliatis, capitulis circ. 20-floris.  $\frac{1}{2}$  ad Cap. Bonæ-Spei (Boiss. herb.), in distr. Uitenhage, Worcester (Eckl. et Zeyh.), ad Isowewelde, Breederivier, Zwarkoprivier (Drege!). Similis C. cernuæ sed ptila fastigiata erecta dimidiò minora; differt à C. ciliatâ et C. tenuifoliâ foliis liberis et à C. pedunculatâ ramis apice non denudatis. (v. s.)

7. *paniculata*, glabra, foliis linearibus erectis sparsis oppositivis, ramis vix vix basi foliosis elongatis nudis, capitulis circ. 20-floris.  $\delta$  ad Cap. Bonæ-Spei in Saldanhabay (Ecklon in h. Dunant), in Namaqualand (Eckl. et Zeyh.), ad Stormberg et Camlsberge (Drege!). Species distinctissima. Folia 4 lin. longa. Pedunculi uni- vel bipollicares. Capitula dimidiò minora quam iorum. (v. s.)

6. *C. LONGIFOLIA*, glabra, foliis angustè linearibus elongatis erectis integerrimis, capitulis pedicellatis corymbosis circ. 20-floris. § in Africa australi ad Olifant-rivier Camisberge legit cl. Drege! Folia 9-15 lin. longa semilineam lata. Pedicelli circ. foliorum longitudine. (v. s.)

*ß. patula*, foliis patulis, corymbis minis confertis. § ad Kaus et Clan-William legit cl. Drege! (v. s.)

7. *C. OBLONGIFOLIA*, fruticosa nana, foliis elliptico-oblongis rigidè ciliatis utrinquè ramisque glabris, ramis florif. apice aphyllis. invol. squamis linearibus acuminiatis margine subserratis. § in Africa Capensi legit cl. Drege! Chr. ciliata E. Mey. ! in Dreg. coll. Melius quàm vera *C. ciliata* nomen meretur sed nec descr. nec icon respondent. (v. s.)

8. *C. TRIDENTATA*, glabra, foliis crassiusculis obtusè trilobis seu grossè tridentatis, paucissimis linearibus integris, capitulis pedicellatis 30-40-floris. § in Africa Capensis deserto Carro et ad Zwartberge legit cl. Drege! Species distinctissima. Capitula *C. Commæ*-aurea ferè majora. Achenia nigra minutissime puberula. Folia variant 3-5 lin. longa. (v. s.)

9. *C. PINNATIFIDA*, fruticosa glabra, foliis sessilibus oblongo-linearibus crassiusculis pinnatifidis, lobis utrinquè 2-3 integerrimis obtusis, marginibus subrevolutis, ramulis apice subnudis 1-cephalis. § in Africa australi ad Zwartberge legit cl. Drege! Caulis circ. bipedalis simplex apice breviter subpaniculatus. Invol. subpurpurascens. Flores pallide flavi. Achenia plano-compressa parva et minutissime puberula (v. s. comm. à cl. inv.)

§. 2. *Herbaceæ, foliis non linearibus, capitulis corymboso-paniculatis.*

10. *C. UNDULATA* (Thunb. fl. cap. 627), herbacea, caule pubescente, foliis cordatis lanceolatis undulatis, capitulis paniculato-fastigiatis. — Ad Cap. Boer-Spei. Chrysocomæ congener ex herb. Thunb. teste Less. syn. 195.

11. *C. DECURRENS*, herbacea subscabrida, foliis elongatis pinnatifidis longè secus caulem decurrentibus, corymbo paniculato-fastigiato, capitulis ad apicem ramulorum congestis. — in Africa australi ad Willbergen alt. 4-500 ped. legit cl. Drege! Herba bipedalis à genere habitu diversissima. Folia ferè *Cony. pinnatilobata*. Flores flavi. Capitula circ. 20-flora. Achenia compressa puberula. Pappus 1-serialis multisetus. Genitalia generis. (v. s.)

12. *C. PTEROCAULA*, herbacea erecta subsimplex hirtello-pubescent, foliis elongatis petiolatis in alas integras secus caulem longè productis, infer. lyncis lobis lateralibus 3-4-jugis ovatis obtusis, terminali lanceolato basi lobato serrato, super. lineari-lanceolatis acuminatis plus minus pinnatilobatis sensim integerrimis, corymbo paniculato laxo. — in Africa australi legit cl. Drege! Prioribus valdè affinis et cum eis sect. naturalem (à Chrysocomia geminis habitu distinctissimam) conficiens. (v. s.)

† *Non satis notæ.*

13. *C. NIVEA* (Willd. sp. 3. p. 1790), fruticosa, foliis lineari-lanceolatis tomentosis, corymbis terminalibus sessilibus. § ad Cap. Boer-Spei. An fortè *Senecionis* sp.?

14. *C. TOMENTOSA* (Linn. syst. veg. 615. Willd. sp. 3. p. 1790), suffruticosa, foliis linearibus utrinquè tomentosis, ramis tomentosis 1-cephalis. § par. et et car. ign. †.

15. *C. ? PURPUREA* (Forst. prodr. n. 286. Willd. sp. 3. p. 1791), herbacea, foliis elliptico-lanceolatis subserratis pubescentibus, paniculâ terminali corymbosa — in insulâ Tannâ. †.

†† *Species exclusæ.*

- C. ACAULIS* Walt. = *Vernonia oligophylla*.

- C. *ADPRESSA* Fl. Flum. = *Vernonia monticola*.  
 C. *AMARA* Schum. = *Decaneurum Senegalense*.  
 C. *ANGUSTIFOLIA* Walt. = *Vernonia angustifolia*.  
 C. *AUREA* Bojer. = *Nidorella Chrysocoma*.  
 C. *BIFLORA* Linn. = *Linosyris punctata*.  
 C. *BIFLORA* Lam. = *Galatella dracunculoides*  $\beta$ .  
 C. *CAMPHORATA* Rob. et Cast. = *Jasonia glutinosa*.  
 C. *CANDICANS* Delil. = *Linosyris candicans*.  
 C. *CAPILLACRA* Michx. = *Eupatorium forniculaceum*.  
 C. *CILIARIS* Link. = *Agathæa Ecklonis*.  
 C. *CINEREA* Labill. = *Ozothamnus cinereus*.  
 C. *CORONOPIFOLIA* Michx. = *Eupatorium coronopifolium*.  
 C. *DENTICULATA* Jacq. = *Plagus virgatus*.  
 C. *DICHOTOMA* Jacq. = *Jasonia dichotoma*.  
 C. *DIVARICATA* Fisch. = *Linosyris divaricata*.  
 C. *DRACUNCULOIDES* Pursh. = *Bigelovia dracunculoides*.  
 C. *DRACUNCULOIDES* Lam. = *Linosyris punctata*.  
 C. *FERRUGINEA* Spreng. = *Ozothamnus ferrugineus*.  
 C. *FOETIDA* Lam. = *Erigeron foetidum*.  
 C. *FRUTICOSA* Stend. = *Vernonia fruticosa*.  
 C. *GIGANTEA* Walt. = *Vernonia angustifolia*.  
 C. *GRAMINIFOLIA* Linn. = *Solidago graminifolia*.  
 C. *GRAVEOLENS* Nutt. = *Bigelovia dracunculoides*.  
 C. *HETEROPHYLLA* Raf. = ? *Baccharidea* ex auct.  
 C. *LAPPULIFOLIA* Bojer. = *Nidorella mucronata*.  
 C. *LEBERNICA* Spreng. = *Linosyris vulgaris*  $\gamma$  *patula*.  
 C. *LINOSYRIS* Linn. = *Linosyris vulgaris*.  
 C. *LINOSYRIS* Aso. = *Galatella rigida*.  
 C. *OLIGOPHYLLA* Fl. Flum. = *Vernonia cephalotes*.  
 C. *MONTANA* Vahl. = *Linosyris montana*.  
 C. *MUCRONATA* Forst. = *Iphyona spinosa*.  
 C. *HAUSEKOSA* Pursh. = *Bigelovia Missouriensis*.  
 C. *NOVEBORACENSIS* Walt. = *Vernonia noveboracensis*.  
 C. *NUDATA* Michx. = *Bigelovia nudata*.  
 C. *ODORATISSIMA* Rensch. = *Liatris odoratissima*.  
 C. *OPPOSITIFOLIA* Linn. = *Pteronia divaricata*.  
 C. *PANICULATA* Gmel. = *Liatris paniculata*.  
 C. *PAUCIFLORA* Fl. Flum. = *Ageratum melissæfolium*.  
 C. *PEDUNCULATA* Fl. Flum. = *Vernonia pedunculata*.  
 C. *PILOSA* Gmel. = *Liatris hirsutissima*.  
 C. *PINIFOLIA* Spr. = *Ozothamnus pinifolius*.  
 C. *PRÆALTA* Desf. = *Vernonia præalta*.  
 C. *PUMILA* Fl. Flum. = *Vernonia grandiflora*.  
 C. *PUNDEKA* Raf. non Forst. = *Liatris* seu *Vernonia* sp.  
 C. *RETICULATA* Labill. = *Ozothamnus reticulatus*.  
 C. *ROSMARINIFOLIA* Spr. = *Ozothamnus*.  
 C. *SAXATILIS* DC. = *Jasonia glutinosa*.



Sack-rivier legit cl. Burchell et mecum comm. sub n. 1405. cat. geogr. Fol. 6-7 lin. longa, 2½-3 lata. Ramuli crebri oppositi ultimi dichotomi. Capitula 16-flora, 8 lin. longa, 4 lin. lata. Stigmata longè exserta. (v. s.)

ß. *rotundifolia*, foliis ovato-subrotundis glabrisculis. — ad Nieuwewede alt. 2-300 ped. legit cl. Drege! (v. s.)

9. *P. oppositifolia* (Lin. syst. veg. 614), ramulis folisque pube subpulverulenti minimà velutinis, foliis oppositis ovato-lanceolatis erectis internodio subæqualibus nervo medio notatis, capitulis terminalibus solitariis sessilibus ovatis glabris apice subviscosis, invol. squamis ovatis obtusis adpressis margine suberoso-ciliolatis. ß ad Cap. Bonæ-Spei. Thunb. fl. cap. 632. Goertt. fr. 2. t. 167. f. 2. *Scopinia dichotoma* Cass. dict. 48. p. 45. Folia 3-4 lin. longa, sesquial. lata. Capit. multiflora 9-10 lin. longa, 6-7 lin. lata. (v. s. comm. à cl. Burchell cat. geogr. n. 6837.)

10. *P. cinerea* (Lin. f. suppl. 356), ramulis subpubescentibus mox glabris, foliis elliptico-oblongis lanceolatisve oppositis cinereo-tomentosis patentibus in ramulis approximatis, capitulis terminalibus sessilibus ovatis 15-20-floris, invol. squamis ovatis obtusis adpressis subciliatis dorso tenuè tomentosis subtristis ß ad Cap. Bonæ-Spei. Thunb. fl. cap. 632. Folia 5-6 lin. longa, juncea can. adulta cinerea. Capit. 8-9 lin. longa, 6-7 lin. lata. Pappus rufus. (v. s. ex herb. Burm. sub nom. *Stahelina*.)

11. *P. xantholepis*, ramulis ultimis folisque pube subtomentosa cano-villosa, foliis oppositis oblongo-linearibus secus ramulos confertissimis, capitulis terminalibus solitariis sessilibus oblongis 6-9-floris, invol. glaberrimi flavi squamis oblongo-linearibus obtusiusculis integris intimis margine angustè membranaceis. ß ad Cap. Bonæ-Spei legit cl. Krebs n. 1901, ad Uitenhage, Swellendam (Ekl. et Zeyh.), et Kaus (Drege!). Folia 3-4 lin. longa, lineam lata. Internodia ramulorum vix semilin. longa. Capit. 6-7-florum. Squamæ serialim imbricatæ exter. nempe suprâ interiores oppositæ nec alternè sitæ. Proximè accedit ad *P. incanæ*. Huc fortè *Eupatorium cinereum* Lin. f. suppl. 354. refer. Pappus albus. Squamæ flavae. (v. s.)

12. *P. incana*, ramulis incanis demum glabris, foliis oppositis linearibus suboblongisve obtusis cano-tomentosis secus ramulos confertissimis, capitulis terminalibus solitariis sessilibus oblongis 10-12-floris, invol. glaberrimi squamis oblongis obtusissimis margine scarioso angustissimo cinctis. ß ad Cap. Bonæ-Spei. *Pteronia rigida* Berg. cap. 231 ex Less. syn. 158. *Chrysocoma incana* N. L. Burm. prod. fl. cap. 24. *Eupatorium cinereum* Thunb. fl. cap. 627? et ideo *Pteronia rigida* Berg. cap. 231? Similis *P. xantholepidi* sed invol. squamis pallidè subfusca nec flavae. Capit. 10-12 nec 6-8-florum. Pappus intense rufus nec albus. Folia multò magis incana nec cinerea. (v. s.)

13. *P. canescens*, ramis glabris vix nascentibus araneosis, foliis oppositis linearibus oblongisve subtrigonis utrinquè canescenti-tomentosis integerrimis ad apices ramulorum confertis, capitulis terminalibus solitariis ovatis, invol. squamis subadpressis ovato-oblongis subacutis dorso araneoso-tomentosis margine angustè scarioso-marginatis. ß in Africâ Capensi ad Onderbokkewald et Zeederberge legit cl. Drege! *Achras sericeo-villosissima* teretiuscula erosius (v. s.)

14. *P. lucilioides*, demissa tota hirtello-velutina canescens, foliis oppositis linearibus obtusis basi cuneato-attenuatis integerrimis enerviis, capitulis terminalibus solitariis sessilibus 7-9-floris, invol. cylindraceis subconicis glaberrimis ß in Africâ Capensi regione Gariepina et Klein-Namaqualand legit cl. Drege! Folia 5-6 lin. longa, 1 lin. lata. Capit. semipoll. longa. Achænis albis hispidis. (v. s.)

15. *P. xanthocarpa*, glabra, junior velutina, foliis oppositis ovatis obtusis subglaucis superioribus internodio longioribus, capitulis terminalibus sessilibus oblongis 4-floris, invol. squamis oblongis glabris acumminatis margine subacutis dorso lineatis, pappo rubro. ß ad Cap. Bonæ-Spei prope Sack-rivier legit Burchell (n. 1403 et 1546 cat. geogr.), et ad Zwart-Ruggens cl. Drege! Ramis

folia novella subvelutina. Rami oppositi adulti rigidi subpinescentes. Folia in. longa, 2 lin. lata. Capit. 5 lin. longum pappo rubro insignitum. (v. s.)

\* \* *Glaberrima.*

*P. MICROPHYLLA*, glabræ ramosissima, foliis oppositis trigonis ad apices ramulorum confertis integerrimis glaucescentibus minimis, capitulis terminalibus litaris plurifloris oblongis basi obconicis, invol. squamis adpressis ovatis, ter. subacutis, inter. subciliatis.  $\frac{1}{2}$  in Africâ Capensi ad Stellenbosch leg. cl. Don! et Zeyher. Affinis *P. ciliatæ*. Folia vix ultra lineam longa. Invol. flamma apice non contractum. Achenium teres sericeo-villosum. Pappus longus et rufescens. (v. s.)

*P. GLOMERATA* (Lin. f. suppl. 356), glabra, foliis oppositis sessilibus ovato-gonis in ramulis præsertim ad apices et axillis glomeratis, capitulis terminalibus sessilibus solitariis obovato-cylindraceis viscosis, inv. squamis adpressis, ter. brevibus ovatis dorso viridibus, cæteris apice flavescentibus ciliatis.  $\frac{1}{2}$  ad p. Bonæ-Spei in Carro (Th. Drege!), ad Zwarteberge et Zwarte-Ruggens (rege!), ad Camisberge et Onder-Bokkewelde (Drege!). Thunb. fl. cap. 63a. web. cat. geogr. n. 1359 et 1435. *Scopinia lepidophylla* Cass. dict. 48. p. 45. Illa 1-2 lin. longa subglauca, summa sub capitulis concava obtusa. Capitulum  $\frac{1}{2}$  lin. longum 4 lin. diam. (v. s.)

*P. PALLENS* (Lin. f. suppl. 357), tota glabra ramosissima, foliis oppositis triangularibus elongatis subfiliformibus obtusis, capitulis solitariis brevè pedunculatis terminalibus turbinateis, invol. squamis ciliatis, inter. margine scariosa subsericeo-ciliatis.  $\frac{1}{2}$  ad Cap. Bonæ-Spei in Carro (Eckl. et Zeyh.), et Gannaland. web. fl. cap. 63o. non Poir. Folia unguicularia ex Th., 10 lin. longa in sem. meo Thunbergiano. Capit. 5-6 lin. longum. Achen. villis albis sericeis nudum onustum. Pappus rufus. (v. s.)

*P. TRICEPHALA*, glabrâ subviscosa ramosissima, foliis oppositis linearibus applicato-trigonis integerrimis apice uncinulato-recurvis, capitulis ad apices ramorum ternis in axillis supremis 4-5-floris, invol. oblongi squamis acutis serribus exter. subtrigonis.  $\frac{1}{2}$  in Africâ Capensi ad Nieuwswyde alt. 2-300 d. legit cl. Drege! Suffrutex vix semipedalis. Folia et invol. 4 lin. longa vix ne vix subœonnata. Achenia teretiuscula erostria villosissima. An fortè fastigiata Thunb. fl. cap. 629? sed in nostrâ folia semilineam lata nec iga. (v. s.)

*P. BRICKROIDES*, ramis glabris gracilibus, foliis oppositis sessilibus linearisæcolatis complicatis integerrimis pulverulento-velutinis, capitulis terminalibus litaris sessilibus oblongo-turbinateis basi attenuatis glabris, invol. squamis æcolatis acuminatis, inf. brevibus apice callosis, aliis apice membranaceis alino-marginatis.  $\frac{1}{2}$  ad Cap. Bonæ-Spei in distr. Swellendam collibus Carro allibus legit cl. Drege! Folia  $1\frac{1}{2}$  lin. longa. Capit. flavicantia 8-10 lin. longa. Achenia sericeo-villosissima teretiuscula. (v. s.)

*P. GYMNOCLINE*, glabra, ramis virgatis striatis, foliis inf. oppositis, super. ternis sessilibus oblongo-linearibus obtusiusculis 1-nerviis cinereis minutissimè punctulatis, capitulis terminalibus solitariis sessilibus oblongo-cylindraceis circ. 15-floris, invol. squamis flavidis adpressis, exter. obtusis, intimis agrioribus acuminatis.  $\frac{1}{2}$  in Africâ Capensis reg. Gariëpinâ, Carro, Klein-maqualand, legit cl. Drege! Folia 5 lin. longa, lineam lata. Capit. flavum 10 lin. longum. Pedicelli apex ex squamarum cicatricibus areolatus. Recept. mura omninò nudum. Achen. villosissimum. (v. s.)

*P. CILIATA*, glabra ramosissima, foliis oppositis ad apices ramorum confertis ipticis rigidis nervo medio prouinulo subcarinatis integerrimis glaucis, capitulis terminalibus solitariis oblongis utrinquè attenuatis, squamis infimis subliaceis carinatis, cæteris adpressis obtusissimis densè et tenuiter ciliatis.  $\frac{1}{2}$  in Africâ Capensi ad Namaqualand leg. cl. Ecklon! et Zeyh. Folia  $1\frac{1}{2}$ -2 lin. longa s.)

- β. subtrigona*, foliis oblongis evidentius subtrigonis.  $\zeta$  ad Olifant-rivier Dregel P. ciliata Thunb. fl. cap. 632? Folia 2-3 lin. longa. (v. s.)
23. *P. succulenta* (Thunb.! prod. 143. fl. cap. 630), tota glabra ramosissima, foliis oppositis trigonis carnosius obtusis patentibus, capitulis terminalibus campanulato-ovatis, invol. squamis ovatis obtusis integerrimis.  $\zeta$  ad Cap. Bonæ-Spei in Karro et Hantum (Th.), ad Clan-William (Eckl.), ad Olifant-rivier (Dreg.). Achænia villis albis sericis erectis densè vestita turbinata sub pappo attenuata. Pappus rufus. Folia in specim. meis 2-6 lin. longa, ex Thunb. angustiora usque pollicaria. Fl. flavi. Fimbrillæ alveolares achæniæ ferè longitudine. (v. s.)
24. *P. sesuvifolia*, glabra ramosissima, foliis oppositis carnosius linearibus acutis planis basi uninerviis patentibus integerrimis, capitulis terminalibus solitariis multifloris, invol. campanulatis squamis ovatis obtusissimis integerrimis.  $\zeta$  ad Cap. Bonæ-Spei, ad Kaus et Olifant-rivier legit cl. Dreg. Affinis *P. succulenta* et fortè eam eâ à Thunb. confusa, sed folia plana nec trigona, acuta nec obtusa, pollicem longa, 2-3 lin. lata. Flores flavi. Achænia sericeo-villosissima. Recept. vix alveolatum. (v. s.)
- §. 3. *Foliis oppositis vaginato-connatis, capitulis multi- pauci-im-*  
*unifloris cylindraceis, invol. squamis adpressis luteis exter. ovatis.*
25. *P. connata*, glabra ad apices viscosa, foliis oppositis basi connatis linearibus filiformibus supernè sulcatis, capitulis ad apices ramorum solitariis ternis ovato-cylindraceis 15-25 floris, invol. squamis adpressis obtusis, ext. ovatis. inter. oblongis.  $\zeta$  in Africâ Capensi in Karro (Dreg.! Eckl.! et Zeyh.), in Swellendam (Eckl.! et Zeyh.). An *P. flexicaulis* Thunb. fl. cap. 629 nec Lin. f.? Folia squamipollicem longa, vaginâ rarissimè ruptâ. Capitula pollicem longa nunc solitaria sessilia, nunc ternâ lateralibus pedicellatis. Flores flavi. (v. s.)
26. *P. flexicaulis* (Lin. f. suppl. 355), glabra ad apicem viscosa, foliis junioribus basi connatis demum ruptis margine membranaceis linearibus filiformibus, capitulis 3-floris densè corymbosis in quoque ramulo subternis pedicellatis, invol. cylindrici glabri squamis oblongis obtusiusculis.  $\zeta$  ad Cap. Bonæ-Spei in Karro (Thunb. fl. cap. 629? Eckl.! et Zeyh.), et inter fluv. Bushman et Gauritz (Roch. cat. geogr. n. 4209), ad Zw. Onder-Bokkeweld et Zeederbergen (Dreg.), in Swellendam et Uitenbage (Eckl.! et Zeyh.). *P. fastigiata* Zeyh.! pl. cap. ex n. 20, non Thunb. qui de eâ dixit folia semilineam longa. Caulis flexuosus cinereus glaber. Folia pollicem longa. Capit. 5 lin. longa. (v. s.)
27. *P. fasciculata* (Lin. f. suppl. 357), tota glabra, foliis oppositis sessilibus confertis lanceolatis acutis viscosius supra concavis subtus carinatis, capitulis plurimis aggregatis fastigiatis oblongis, invol. glabri squamis lanceolatis.  $\zeta$  ad Cap. Bonæ-Spei in Karro prope Hex-rivier (Th.), Gauritz-rivier (Eckl.), in collibus distr. Swellendam (Eckl.), ad Zeederbergen (Dreg.). Thunb. fl. cap. 630. Henanthus fasciculatus Less. syn. 195. Folia pollicem longa, 2 lin. lata, junioribus vaginato-connatis, vaginis demum obliquè ruptis et ligulas triangulares stipulaformes secus ramos persistentes et oppositas relinquentibus. Invol. later.

Sect. II. *PACHYDERIS*. — *Pachyderis* Cass. dict. 56. p. 170. Less. syn. 196.

Capitulum multiflorum. Recept. alveolatum. Achænia compressa (exter. interdum subtrigona) apice brevissimè rostrata, nempe sub pappo in collum crassius glabrum sæpè villis occultatum attenuata. — Folia opposita rarius alterna

§. 1. *Achæniis plus minus villosis hirsutiusc.*

\* *Corollis purpureis.*

28. *P. urtica* (Lin. suppl. 356), ramis hispida, foliis oppositis lanceolatis

acutis suprà glabris subtùs hirsutis margine ciliato-scabris, capitulis terminalibus sessilibus solitariis ovatis multifloris, invol. glabri viscosi squamis ovatis integris.  $\zeta$  ad Cap. Bonæ-Spei, in Karro (Thunb. fl. cap. 631.), in coloniæ distr. orient. (Burch. ! cat. geogr. n. 6380 et 6208), ad Zwartland, Paarlberg (Dreg.!), ad Clan-William, Swellendam, Worcester (Eckl. ! et Zeyh.). An *P. uniflora* Poir. dict. 5. p. 733? Folia 5 lin. longa, 1; lin. lata sub capitulis conferta falso-verticillata. Capit. pollicem longum, 9 lin. diam. Cor. purpurea. Achænia villosa in collum glabrum crassum brevem sed aliis ferè longiorem attenuata. Pappus rufus. (v. s.)

29. *P. CEPHALOTES* (Linn. f. suppl. 338), ramis hirtis subfastigiatis, foliis oppositis lanceolato-trigonis rigidè ciliatis subsabris remotiusculis, capitulis terminalibus sessilibus ovatis solitariis, invol. squamis imbricatis adpressis apice subpuberulis viscosisque, exter. brevibus ovatis, mediis oblongis apice ad margines membranaceis ciliatis.  $\zeta$  ad Cap. Bonæ-Spei prope urbem Burch. ! cat. n. 61. Eckl. ! et Zeyh. Thunb. fl. cap. 631. *Jacea incana*, etc. Burm. ! herb. et ideò *Centaurea incana* Burm. prod. 28 videtur. Capitulum 9 lin. longum, 5 lin. diam. Folia 4 lin. longa erecta patula aut apice recurva. Achænia compressa villosa-hispida breviter et crassè rostrata. Cor. purpurea. Priori valdè affinis. Ramuli interdum parè hirtelli et folia etiam parè ciliata, cæterum glabra (Eckl. !) nundè ad var.  $\beta$  transit. (v. s.)

$\beta$ . *glabra*, ramis glabris, foliis eciliatis in disco utroque glabris complicatis aut planis.  $\zeta$  in Karro legit cl. Ecklon. (v. s.)

\*\* *Corollis flavis.*

30. *P. UNDULATA*, ramis glabris ad apicem tenuissimè folisque glanduloso-pulverulentis, foliis oppositis in petiolum attenuatis ovatis undulato-crispatis, capitulis ex axillis super. pedicellatis in corymbum 3-7-cephalum dispositis oblongis 5-floris, invol. squamis erectis oblongis glabris integris obtusiusculis.  $\zeta$  in Africa Capensi ad Zilverfontain in Klein-Namaqualand legit cl. Dreg. ! Folia cum petiolo sesquialin. 7-8 lin. longa 4 lin. lata. Invol. flavescens pappo maturo dimidiò brevius. Flores flavi. Achæniū compressum subrotellatum parè villosum. (v. s.)
31. *P. OVALIFOLIA*, ramis glabris vix apice summo subcanescentibus, foliis oppositis elliptico-oblongis ovalibusque integerrimis pubè adpressà subtomentosà canis, adultis glabratiss, capitulis ovato-oblongis terminalibus solitariis, invol. squamis adpressis, exter. ovatis, inter. magis oblongis margine scariosis, omnibus mollè et tenniter ciliatis.  $\zeta$  in Africa Capensi ad Olifantivier (Dreg.!), in Worcester (Eckl.!). Achæniū villosissimū brevissimè rostratum. Habitus ferè *P. canescentis*. Folia majora 8 lin. longa. 4 lin. lata, plurima minora. (v. s.)
32. *P. SCARIOSA* (Linn. f. suppl. 356), ramis glabris demum apice rigidis spinescentibus, foliis suboppositis sessilibus ovali-oblongis glabris, capitulis terminalibus solitariis ovatis, invol. glabri squamis ovatis margine latè scarioso-membranaceis mucrone fusco-subpungenti-acuminatis.  $\zeta$  ad Cap. Bonæ-Spei ad Hantam (Thunb. fl. cap. 633), ad Kouss (Dreg.). Achænia compressa villosa breviter rostrata. (v. s.)
33. *P. MEMBRANACEA* (Linn. f. suppl. 357), ramis glabriusculis, foliis oppositis sessilibus oblongis acutis pulverulento-tomentosis, capitulis terminalibus solitariis oblongis, invol. glabri squamis lanceolatis margine latè membranaceis dorso fusco-virescentibus.  $\zeta$  ad Cap. Bonæ-Spei in Karro (Thunb. fl. cap. 633) Ecklon ! et Zeyh. in Bosmanivier et Zneureeberge (Dreg.!). Frutex pedalis, ramis demum induratis subspinescentibus, adultis obtusè tetragonis. Folia 5-8 lin. longa 1-nervia interdum basi trinervia. Invol. ferè *P. scariosa* sed mucro ultra partem scariosam non productus. Achæniū villosissimū teretè angulatum brevissimè rostratum. (v. s.)
34. *P. ACCUMINATA*, tota glabra, foliis alternis lanceolatis utrinquè attenuatis mu-

§. 2. *Acheniis omnibus aut saltem centralibus glabris.*

49. *P. HETEROCARPA*, glabra decumbens?, foliis oppositis breviter connatis linearibus obtusis integerrimis crassiusculis, capitulis terminalibus solitariis sessilibus ovato-campulanatis, invol. foliis proximis superato glabro, squamis ovali-oblongis obtusis margine hyalinis, acheniis exter. villosis-hirsutis, centralibus glaberrimis.  $\frac{3}{4}$  in Africa Capensi ad Olifant-rivier legit cl. Dreg! Folia 12-15 lin. longa, 1  $\frac{1}{2}$  lata. Capit. 8-9 lin. longum latumque. Pappus rigidus. (v. s.)
50. *P. STORRELINOIDES*, glabra, foliis oppositis sessilibus rigidulis glaucis linearilanceolatis complicatis ad margines et sæpè subtus in carinâ rigidè ciliatis, capitulis terminalibus solitariis sessilibus ovato-subcylindraceis, invol. squamis glabris adpressis margine scariosis subfimbriatis, acheniis glaberrimis glandulosis.  $\frac{3}{4}$  in Africa Capensi ad Graaf-Reynet (Dreg! Eckl! et Zeyh.), in Carro et Namaqualand (Eckl! et Zeyh.). Affinis *P. onobromoidi*. Folia 4-5 lin. longa, 1  $\frac{1}{2}$  lata. Capit. pollicem longa. (v. s.)
51. *P. VISCOSA* (Thunb. fl. cap. 632), ramis glabris, foliis oppositis sessilibus ovatis callosis-subacutis margine scabridis patulis, capitulis terminalibus solitariis sessilibus ovatis obtusis integris acheniis glanduliferis.  $\frac{3}{4}$  ad Cap. Bone-Spei in Hantum (Th. Zeyh! in h. Moric. n. 98), ad Zwart-Ruggens in Carro (Dreg!). Capitula crassa pollicem longa. Achenia basi vix brevibus erectis (fortè fimbriis) cincta compressa oblonga utrinquè attenuata glanduloso-scabrida. Pappus rufescens. (v. s.)
52. *P. ONOBROMOIDES*, glabra, foliis alternis sessilibus elongatis lanceolato-linearibus obtusis rigidulis subconvolutis margine rigidè ciliatis, capitulis terminalibus solitariis sessilibus ovatis multifloris, invol. squamis integerrimis adpressis obtusis ext. latè ovatis, int. oblongis, acheniis glanduliferis.  $\frac{3}{4}$  in Africa Capensi ad Olifant-rivier (Dreg!) et in Namaqualand (Eckl!). Rami albi. Folia pallida 12-18 lin. longa, 2-3 lata. Capit. 15 lin. longum, 10-12 lin. latum. Cor. flavæ. Achenia compressa glabra apice breviter rostrata. (v. s.)

Sect. III. *PTEROPHORUS*. — *Pterophorus* Vaill. act. ac. Par. Cass. dict. 37. p. 474. Less. syn. 195. — *Pterophora* Neck. elem. 1. p. 139.

Capitulum multiflorum. Recept. fimbriiferum. Achenium erecto plano-compressum glaberrimum. Pappus subuniserialis, setis basi in anulum subconcreta. — Folia sparsa linearia integra capitula terminalia solitaria, ramis ad apicem ferè foliosis. — Fortè genus inter *Pteronium* et *Chrysocoma* medium.

53. *P. CAMPHORATA* (Lin. sp. 1176. mant. 464), ramis hirsutis, foliis sparsis fasciculatisque linearifiliformibus ciliatis, capitulis terminalibus solitariis sessilibus, invol. squamis serrulato-ciliatis disco brevioribus.  $\frac{3}{4}$  ad Cap. Bone-Spei frequens (Burni! Eckl! et Zeyh. Dreg!). *Pteronia* camphorata N. L. Barm! prod. fl. cap. p. 24. Thunb. fl. cap. 629. Lam. ill. t. 667. f. 1. *Pterophorus* camphoratus Cass. dict. 37. p. 474. — Pluk. mant. t. 345. fol. 56. pl. 2. Florum interdum 6-fidi. (v. s.)
54. *P. STRICTA* (Ait kew. 3. p. 162), ramis apice villosis-scabris, foliis sparsis fasciculatisque linearifiliformibus basi ciliatis, invol. squamis integerrimis apice in folium subacerosum productis disco æqualibus longioribusve.  $\frac{3}{4}$  ad Cap. Bone-Spei ad Zwartberge (Dreg!). *Pterophorus* strictus Less. syn. 195. *P. leucicna* Houtt! in h. Delessert. (v. s.)
55. *P. ASPERA* (Thunb. fl. cap. 631), ramis villosis-scabris, foliis fasciculatis linearibus glabris, invol. squamis glabris lanceolatis acutis apice reflexis.  $\frac{3}{4}$  ad Cap. Bone-Spei, in Stellenbosch, Swellendam, Langekloof, etc. (Eckl! et Zeyh. in Paarlberg et Giftberg (Dreg!). *Pterophorus* asper Less. syn. 195.

*non satis nota (achæniis nempe indescriptis). Plures fortè inter priores jam recensita:*

\* *Alternifoliae.*

**FOSEA** (Poir. dict. 5. p. 731); ramis.... foliis sparsis linearibus sessilibus basi spinuloso-ciliatis, capitulis terminalibus solitariis sessilibus, invol. glabri squamis ovali-lanceolatis obtusis apice subfimbriatis. Bonæ-Spei. Invol. squamæ flayæ.

**SOCOMIFOLIA** (Poir. dict. 5. p. 732), ramis pubescentibus, foliis filiformibus acutis glabris, capitulis confertis, invol. glabri squamis obtusis. ♀ ad Cap. Bonæ-Spei. Cor. in sioco corulescens.

\*\* *Oppositifoliae.*

**ALATA** (Thunb. prod. 143. fl. cap. 629), tota glabra, foliis oppositis ligonibus imbricatis apice recurvis, capitulis terminalibus sessilibus, invol. squamis ovatis. ♀ ad Cap. Bonæ-Spei. Caulis ramosissimus, fragilis flexuosis subfastigiatis. Folia semilineam longa.

**ALATA** (Thunb. fl. cap. 629), tota glabra, foliis oppositis filiformibus, capitulis in paniculam terminalem dispositis, invol. oblongis squamis ovatis obtusis. ♀ ad Cap. Bonæ-Spei. Achæn. ign.

**LA** (Linn. f. suppl. 356), ramis scabridis foliis oppositis linearibus sessilibus, capitulis terminalibus solitariis ovatis, invol. squamis marginibus membranaceis. ♀ ad Cap. Bonæ-Spei Thunb. fl.

**A** (Thunb. fl. cap. 631), ramis glabris, foliis oppositis frequentibus glauco-tomentosis subserratis obtusis, capitulis terminalibus sessilibus, invol. glabri viscosi squamis ovatis integris, infimis carinatis. Bonæ-Spei. Folia semilineam longa.

**TA** (Linn. f. suppl. 356), ramis pilosis, foliis oppositis sessilibus piloso-scabris reflexis, capitulis terminalibus solitariis ovatis, invol. glabri squamis ovato-oblongis integris apice membranaceis. ♀ ad Cap. Bonæ-Spei. Thunb. fl.

**LA** (Linn. f. suppl. 356), ramis glabris, foliis oppositis subsessilibus sessimè pubescentibus (pilosise), capitulis terminalibus solitariis ovatis, invol. glabri squamis ovato-oblongis integris apice membranaceis. Bonæ-Spei in Hantum. Thunb. fl. cap. 633.

†† *Species exclusæ.*

**ica** Spreng. = *Chamæpence mutica*.

**ica** Ten. = *Chamæpence gnaphalodes*.

**ica** f. = *Nestleria minuta*.

**A** Sims. = *Helychrysium*.

**ica** Cav. = *Dysodia porophylla*.

**A** Lour. fl. coch. ed. W. v. 2. p. 597, certè è genere excludenda sed ignotus.

Div. IV. *SOLENOGYNEÆ.*

discoideum heterogatum, fl. radii tubulosis aut vix ac ne vix subsistens sub-3-dentatis femineis pauciseriatis, disci hermaphroditis. Subdivisio ambigua inter *Astereas* et *Baccharideas*.

\* Subtrib. II. *BACCHARIDÆ* Less. *syn.* 200 non 204.

Capitula heterogama aut dioica nunquam radiata, corollæ omnes tubulosæ, femineæ filiformes sæpius multiseriales in ambitu. Antheræ ecandate. Recept. epaleaceum. — Folia alterna.

Div. I. *CONTYÆ* Less. *syn.* 200.

Capitula heterogama monoica.

Subdiv. I. *SPHÆRANTHÆÆ*.

Capitula in glomerulum sphericum congesta ad bracteatum axillas semilia. — Species gerontogææ.

CC. *BLEPHARISPERMUM* Wight herb. DC. in *Wight contrib. bot. ind.* p. 11. et diss. cum icon.

Capitula innumera in glomerulum globosum conservata, biflora, fl. tubulosis, altero femineo apice breviter 3-dentato, altero masculo 5-dentato. Recept. capituli angustissimum paleâ lineari inter flores interjectâ. Invol. squamæ ext. paucæ (2?) breves ovatæ, int. (seu paleæ rec.) duæ flosculis æquales, altera sub fl. femineo plana, altera sub fl. masculum conduplicata. Stylus brevis non exsertus apice vix bifidus, infra emargin. subtumidulus puberulus. Antheræ apice appendiculatæ basi subcaudatæ? Achænium fl. fem. obovale-oblongum obcompressissimum subtetragonum, angulis 2 nerviformibus cum lateribus glaberrimis, 2 marginalibus longè et regulariter ciliatis; fl. masc. gracile elongatum glabrum. Pappus achæniî fem. paleis scariosis 3-5 acuminatissimis, masc. nullus. — Suffrutices Indici glaberrimi. Rami teretes graciles. Folia alterna ovata aut ovalia integerrima penninervia. Pedunculi 1-3 ad apicem rami 1-cephali. Glomerulus globosus. — Affinitus ignota. — Car. præcipuè ex B. petiolari.

1. B. PETIOLARE (DC. l. c.), foliis petiolatis ovato-lanceolatis acuminatis, glomerulis ternis longè pedunculatis.  $\zeta$  in Ind. orient. legit cl. Wight sed loco in herbario non indicatus. Glomeruli Piso paulò majores. Bracteolæ involucri univ. loco parvæ membranaceæ paucæ capitulis breviores. Conf. descript. Kleinii editam in Wight contrib. bot. ind. p. 12. (v. s. in h. Wight.)
2. B. SUBSESSILE (DC. l. c.), foliis ellipticis utrinquè attenuatis subsessilibus. glomerulo terminali solitario subsessili, bracteis foliaceis capitula superantibus.  $\zeta$  in agro Maderaspano. Glomerulus ferè Platani capitulo æqualis. Capitula fortè pluriflora. (v. s. in h. Wight et inter pl. Wallichianas in h. bot. lin. Lond. sine nom. sub n. 7524.)

CCI. *ATHROISMA* DC. in *ann. bot. Guill.* 1833. v. 2. p. 516.

Capitula plurima in glomerulum ovatum conferta. Axis glomeruli cylindrica. Bractææ sub capitulis ovatæ acutæ concavæ. Capitula pluriflora heterogama. Recept. capituli paleas plurimas concavas membranaceas inter flores gerens. Inv. oligophyllum à paleis recept. vix distinctum. Fl. ext. 4-5 feminei tubulosi tenues 4-5 dentati. Anth. .... Stylus fl. int. parvis masculi tubulosi apice latiores 5-dentati. Anth. .... Stylus fl. fem. apice bifidus glabriusculus. Achænia nigra obcompressa ovalia hinc planiusculine convexa ad angulum marginalem apice ciliata cæterum glabra. Pappus parvus villum referens subsetosus. — Herba Indica basi suffruticosa erecta ramosa parçè subpuberula. Folia alterna petiolata lyrata.

pinnatipartita, lobis oblongis obtusis dentatis. Glomerulus terminalis subsessilis stramineus. — *Athropsma* græcè acervus ad capitulorum dispositionem alludit. — Genus videtur medium inter *Sphæranthum* et *Nepharispermum*.

1. *A. LACINIATUM*.  $\frac{5}{2}$  in prov. Indiæ or. Martabanq. ad Troglā legit cl. Wallich. *Sphæranthus laciniatus* Wall. cat. n. 3184. comp. 294. cat. hirm. 1872. (v. s. comm. ab ill. Cor. merc. angl. Ind. or.)

CCII. *SPHÆRANTHUS* *Vaill. act. Par. 1719. t. 20. f. 12. Linn. gen. n. 998. Lam. ill. t. 718. Gærtn. fr. 2. t. 164. Cass. dict. 50. p. 208. Less. syn. 201 non Scop.*

Capitula in glomerulum subglobosum densè aggregata, pluriflora, heterogama, fl. omnibus tubulosis, fœmineis pluriserialibus basi subindurato-dilatatis sæpè tridentatis in ambitu, styli abortu masculis, 5-lentatis in centro. Recept. parziale nudum generale bracteolatum. Invol. parziale polyphyllum uni aut 2-3-seriale, generale varium. Anth. caudatæ. Stylus in capit. ambitu bifidus, in centro indivisus. Achæniium crostre epapposum pilosiusculum aut glabrum. — Herbæ Indiæ aut Africanæ. Folia longè secus caulem decurrentia. Pedunculi alati aut apteri 1-cephali. Glomerulus densus Pisi aut Avellanæ magnitudinis. Cor. violacæ.

# Sect. 1. *EUSPHÆRANTHUS*.

Glomeruli axis plena nec lacunosa; bractææ omnes lineares molles capitulis subæquales, exteriores vix ac ne vix involucrium generale constituentes. — Species Indiæ.

1. *S. MICROCEPHALUS* (Willd. sp. 3. p. 2395), foliis lanceolatis aut subovatis serratis vix asperiusculis in alas integerrimas secus caulem decurrentibus, capitulis globosis, pedunculis sæpius apteris capitulo triplò longioribus.  $\frac{2}{2}$  frequens in uliginosis et oryzetis Indiæ orient. nempe in ins. Luzoniâ (Cham.), Javâ (Blum.), Timor (Kiedl.), ad Prome, Penang, Tavoy, Chapedong, etc. (Wall. Wight!), in Chinâ (Staunt.). *Sph. africanus* Wall. cat. et herb. n. 289 excl. litt. B. *Sph. lævigatus* et *Sph. globosus* Wall. herb. *Sph. indicus* Blum. bijdr. p. 891. var.  $\frac{6}{2}$ . An *Sph. indicus* Linn. saltem fl. Zeyl. n. 312? An *Sph. africanus* ex Indiâ Linn. sp. 1314? Capitula semper in meis specim. globosa Pisi circ. magnit. Pedunculus apterus aut hinc indè rarissimè subalatus. (v. s. specim. co.)

2. *S. STRATUS* (Willd. sp. 3. p. 2395), foliis obovatis serratis utrinquè birtellis in alas serratas desinentibus, glomerulis ovato-globosis, pedunculis glomerulo triplò longioribus aliis serrato-incialis ferè semper instructis. — Frequens in humidis et oryzetis peninsulae Indiæ (Wight!), ad Patna, Kuworkot, Travancore, etc. (Wall.), circa Pondichery (Belang.), in ins. Java circa Buitenzorg (Blum.). *Sph. hirtus* Blum. bijdr. p. 891., Wall. cat. n. 288. D. E. F. *Sph. africanus* Wall. cat. n. 289. B. C. *Sph. indicus* Wight! herb. Icones Pluk. alm. t. 108. f. 7 et 312. f. 6. Burm. Zeyl. t. 94. f. 3. hinc fortè pertinent. Glomerulus Pisi duplò ferè major. Pedunculus rarius subapterus. (v. s.)

3. *S. MOLLISS* (Roxb. herb.), foliis obovato-lanceolatis utrinquè caulisque hirsuto-lanuginosis grossè serratis, serraturis aristatis, subserratis, glomerulis ovato-globosis subhirsutis, pedunculis glomerulo duplò longioribus aliis hirsutissimis apteris aliis interruptè alatis hirsutis.  $\frac{2}{2}$  in Indiâ orient. prov. Silhet, Kamaon, Hindostan (Roxb. et Wall.). — Icon Rhœd. mal. 10. t. 43. hinc fortè refer.? — Caulis ad collum induratus ferè fruticosus. Caulis à collo plurimi sesquipedales ramosissimi. An fortè *Sph. hirti* var. hirsutissima? (v. s. comm. ab hon. Anlâ merc. angl. Ind. or. et à cl. Royle.)



## Sect. II. CUSPIDELLA.

Glomeruli axis subplena aut vix subcava; bracteæ apice acuminato-pungentes, exteriores invol. universale distinctè formantes, ceteræ vix ac ne vix capitula superantes. — Species Africanæ.

4. *S. SENEGALENSIS*, foliis lanceolatis acuminatis utrinquè villosis aristato- et fere spinuloso-serratis secus caulem in alas grossè et subspinuloso-serratas desinentibus, glomerulis globosis, pedunculis interruptè et subspinuloso-alatis glomerulo multò longioribus. (1) in Senegalià ad ripas innund. fluminis præsertim prov. Walo legerunt cl. Adanson, Roussillon! Perrottet! Sph. africanus seu paniculatus Cass. dict. 50. p. 211 et 212. Variat à caule digitali simplici 1-cephalo ad caulem sesquipedalem ramosissimum polycephalum. Capitulum vix-ecum Pisi majoris magnit. (v. s. comm. à cl. Perrottet.)
5. *S. ANGUSTIFOLIUS*, foliis linearibus acutis glabriusculis distanter serratis in alas longas subserratas desinentibus, pedunculis glomerulo duplò ferè longioribus aliis integris ad latus inferius instructis, glomerulo depresso bracteis latè ovatis acuminatis cincto. — in arenosis humidis Senegaliæ ad Casamance legit cl. Perrottet. Planta abnormis et fortè novi generis pignus. Capitula parva pauciora in glomerulum densè aggregata. (v. s. comm. à cl. Perrottet.)
6. *S. PEDUNCULARIS*, foliis linearibus acutis glabriusculis serratis in alas longas angustas subserratis productis, pedunculis folia æquantibus aut superantibus, glomerulo globoso bracteis latè ovatis breviter mucronatis cincto. — in Africa australiori ad Zw. Omsanculo et Omcomas legit cl. Drege! Proximè accedat ad *S. angustifolium* sed pedunculi 2 pollices nec semipollicem longi. Folia 2-3, poll. longa, 2-3 lin. lata. (v. s.)

Sect. III. POLYCEPHALOS. *Polycephalos Forsk. ægypt. descr. p. 154. — Oligolepis Cass. dict. 50. p. 212. — Sphæranthus Gartn. fr. 2. p. 413. t. 164.*

Glomeruli axis plena aut sæpiùs cavo-lacunosa; bracteæ omnes ovato-lanceolatæ cuspidatæ rigidæ capitula superantes, exteriores vix ac ne vix invol. gener. constituentes. — Species Indicæ, Africanæ et Australasiæ.

7. *S. GLABER*, totus glaber, foliis lanceolatis vix subserratis in alas subintegerrimas decurrentibus, pedunculis glomerulo paulò longioribus aliis apteris aliis ad basin hinc indè angustissimè alatis. ? ad ripas flum. Alligatorum in sinu Vadiem ad oram Australiæ borealem legit cl. Cunningham. Axis obovata plena Pedunculus gracilis. Basis corollarum callosa albida. Fl. disci pauci. Achænie glanduloso-viscidum. (v. s. comm. à cl. Cunningham.)
8. *S. SUAVEOLENS*, foliis lanceolatis serratis vix asperiusculis in alas parçè serratis decurrentibus, pedunculis glomerulo dimidio brevioribus subtrigonis angulè alatis subdentatis. 2l. in Ægypto ad ripas Nili! et fortè in Indià orientali ad nulla specimina Indica in herbariis reperi. — *Polycephalos suaveolens* Forsk. æg. p. LXXII et 154. Sph. indicus Lam. ill. t. 718. f. 2. Gartn. fr. t. 164. Del. fl. æg. ill. p. 27. Cass. dict. 50. p. 210 et omn. ferè hort. europ. Vulgò arab. dicitur *Habagbag*. Glomerulus ovato-globosus, axi lacunosâ, 6-7 lin. longus totidemque diam. Cor. violaceo-purpureæ. (v. s. sp. et v. c.)
9. *S. AMARANTHOIDES* (Burm. fl. ind. 186), foliis lanceolatis aut lineari-lanceolatis integerrimis serratisve glabris subtus punctatis breviter decurrentibus, capitula ovato-globosis subsessilibus, pedunculis brevissimis apteris, bracteis acuminatissimis. 2l. in Indiæ orientalis humidis, in prov. Tangore ad oppidum Nepatnam (Wight!), ad ripas Irrawaddy inter Kink-Talong et Vandoboo (Wall.) et fortè in Zeylonâ. — DC. in Wight contrib. bot. ind. p. 11. *S. Zeylanicus* Heyne in h. Wall. n. 290. — Pluk. amalth. t. 357. f. 6. Affinis *S. suaveolenti* sed notis indicatis distinctissimus. (v. s. sp. comm. ab hon. Aulâ merc. angl. Ind. et
10. *S. ? COTULOIDES*, reptans hirsutus, caulibus teretibus gracilibus, foliis ovatis

dentatis mucrone cuspidatis secus caulem in alam parvam interruptam productis, capitulis parvis terminalibus brevissimè pedicellatis. — in insulâ Madagascar ad viarum margines prope urbem Tananarivon legit cl. Bojer. Cotula reptans Bojer! in litt. Herba parva habitu ad Grangeas seu Dichrocephalas accedens, sed ex foliis decurrentibus et squamarum dispositione in anthodio equidem nimis juniore observatâ, speciem cum dubio ad Sphæranthum reduxi. (v. s. comm. à cl. Bojer.)

† *Species non satis notæ.*

11. *S. COCHINCHINENSIS* (Lour. fl. coch. 2. p. 623), foliis oblongis integerrimis lanuginosis secus caulem glabrum indè pseudotriquetrum decurrentibus, glomerulis cordato-ovatis breviter pedunculatis. — in segetibus et hortis Cochinchinæ et Chinæ. Flores albicantes parcè purpurâ conspersi.

12. *S. CHINENSIS* (Linn. mant. 119), foliis sessilibus pinnatifidis, pedunculis alas decurrentes crispas gerentibus. — in Indiâ orientali. An verè Sphæranthi congener? An affinis *S. laciniato* Wall.? An eadem? Cur *Chinensis*, cum ipse cel. auctor ex Indiâ ortam dicat? †.

†† *Species exclusæ.*

*S. AFRICANUS* Burm. = *Dichrocephala latifolia*.

*S. AFRICANUS* Linn. videtur ex priore et ex Sphærantho microcephalo conflatus.

*S. LACINIATUS* Wall. = *Athroisma laciniatum*.

Subdiv. 2. GRANGEINÆ.

Capitula heterogama monoica in glomerulum non congesta. Pappus nullus aut subnullus.

CCIII. *DICHOCEPHALA* DC. in ann. bot. Guill. 1833. v. 2. p. 517 et in *Wight contrib. p. 111*. — *Centipeda* Less. syn. 200 non *Lour.* — *Cotula*, *Grangea*, *Ethulia* et *Artemisia* sp. auct. — *Dichrocephala* sp. *L'Hér. mss.*

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, marginalibus multiseriatis vix 3-4-dentatis fœmineis, centralibus hermaproditis abortu masculis paucis 4-dentatis. Recept. nudum conicum. Invol. expansum subuniseriale, squamis ovatis subæqualibus. Cor. fœm. cylindricæ, centrales fauce campanulatâ. Stylus inclusus. Achænium compressum erostre, fœmineis calvum, masculis setulâ 1-2 coronatum. — Herbæ annuæ ex Asiâ aut Africâ ortæ hirsutiusculæ aut subglabræ ramosæ adscendentes aut erectæ. Folia alterna dentata cuneata aut petiolata, petiolo nudo aut appendiculato. Capitula globosa parva, in racemos paniculasve paucas disposita, pedicellis propriis nudis breviora. — *Centipeda* Loureirii quàm cl. Lessing non ipse vidit, differt à nostro genere, ex descr. auct., 1°. Achæniis oblongis subteretibus nec compressis. 2°. Pappo nullo nec in fl. masc. 1-2-setoso. 3°. Capitulis axillaribus solitariis sessilibus nec racemosis. 4°. Floribus purpurascensibus nec albo-flavis. Potius ad *Myriogynem* pertinere videtur.

1. *D. CAPENSIS*, caule demisso aut adscendente pilis sparsis hirtello, foliis ovatis grossè dentatis sparsè puberulis petiolatis, petiolis aliis nudis aliis lobulis 1-2 utrinquè auctis, capitulis longè pedicellatis racemosis. (1) ad Cap. Bonæ-Spæ (Th.), in Coloniz distr. oriental. (Burch. cat. geog. n. 5243). *Ethulia auriculata* Thunb. fl. cap. 623. *Centipeda Capensis* Less. syn. 201. Herba basi repens adscendens vix pedalis. Capitula parva subglobosa pauca racemosa. (v. s. comm. à cl. Burch.)

2. *D. GRACILIS*, glaberrima aut spice summo subpubescens, caule adscendente debili parcè ramoso, foliis exauriculatis pinnatipartitis lobis utrinquè 2-3-

bus fœmineis tenuissimis limbo parvo 3-dentato, centralibus plurimis masculis tubo obconico 5-dentato. Invol. pluriseriale disco serè brevius. Recept. ebracteolatum suprâ invol. basin elevatum cupulæforme. Achænia epapposa glabra oblonga utrinquè attenuata, rostro brevi pelucido. — Herbæ Indicæ puberæ odoratæ viscosæ erectæ ramosæ. Folia pinnatipartita, lobis obovatis linearibus dentatis, dentibus lobulivè secus rachin inæqualiter intermixtis, caulina basi in aurículas dentatas magnitudine varias dilatata. Capitula parva subglobosa albida, in racemos corymbosos nudos terminales disposita.

1. *C. LYRATA* (Cass. l. c.), foliis inferioribus lyratis, lobis folii superioribus majoribus obovatis. (1) in Indiâ orient. provinciis diversis nempe in Pegu (Bolang.), Nepauli et circa Trogla, Oude, Promé, Taong-Dong (Wall.), Moory-Journa (Ham.). — DC. in Wight contrib. bot. ind. p. 12. *Tanacetum viscosum* A. B. C. D. Wall. l. cat. et herb. n. 3232. comp. 342. *Tanacetum gratum* Wall. herb. Tan. purpureum Ham. l. herb. *Artemisia hirsuta* Rottl. in Spreng. syst. l. p. 490 ex specim. Spreng. à cl. Besser comm. *Herba sesquipedalia*. (v. s.)
2. *C. STRICTA*, foliis pinnatipartitis, lobis sublinearibus grossè et regulariter serratis. (1) in Indiâ orient. verisim. Maderaspatanâ. *Tanacetum viscosum* E et Wall. l. cat. n. 3232. comp. 342. *Artemisia stricta* Heyn. l. herb. (v. s.)

#### CCVI. LESTADIA Kunth in Less. syn. 203.

Capitulum multiflorum heterogamum, fl. fœmineis pluriserialibus in ambitu, reliquis masculis. Receptaculum planum epaleaceum. Invol. imbricatum. Corolla disci et ambitus 5-dentata sed fœminea tenuior. Achænium oblongum rostratum calvum. — Fruticulus ramosissimus circ. 4 pollic. Americæ meridionalis. Folia hirsuta linearia sparsa confertissima. Capitula solitaria in ramis supernè aphyllis, æquè magnæ ac in *Bellide* perenni.

1. *L. PINIFOLIA* (Kunth et Less. l. c.) 5 in Amer. merid. Habitus *Asteris hymphifolii*.

#### CCVII. GYMNA RHENA Desf. mem. mus. 4. p. 1. t. 1. Cass. dict. 21. p. 1132. 23. p. 566. Less. syn. 419. — *Gymnarhea* Steud.

Capitulum multiflorum discoideum, fl. radii multiseriis fœmineis tubulosis 3-4-dentatis post anthesin basi incrassatis et accretis, fl. disci 10-12 bisexualibus ovarii abortu masculis fauce dilatatis 4-dentatis. Invol. nullum aut squamis paucis 1-seriis constans, bracteis foliaceis 1-2 suppletum. Recept. convexum in parte exter. paleas florescentes mineas involventes post anthesin accretas gerens, in disco et radiis fimbrillas floribus interpositas ferens. Anth. basi nudæ. Stylus rubifidus, ramis longis gracilibus semiteretibus, disci simplex subclavatus acutus grossè papillosus. Achænia radii villosa oblonga paucis setoso pluriseriali scabro superata, disci teretia glabra inanitia setis uniseriatis basi inter se concretis superata. — Herba Orientalis densa annua ramosa. Folia alterna pauca oblonga circa capitulum bracteacea, cæterum nulla. Capitula flava. — Genus in ordine anomali videtur affine hinc Evaci suad. cl. Desfontaines sed antheræ ecaudatæ illinc Grangeæ suad. cl. Cassini sed pappus saltem elongatus.

1. *G. MICRANTHA* (Desf. l. c.) (1) in Persiâ inter Mossul et Bagdad leg. cl. Olin et Bruguiera.

## Subdiv. 3. EUCONYZÆ.

ala in glomerulum non congesta. Pappus pilosus 1- aut a-serialis.

**II. THESPIIS DC.** in *Guill. ann. bot.* 1833. v. 2. p. 517. — *Cotula* sp. *Wall.*

culum multiflorum heterogamum, flor. marginalibus multiseri-  
or. tenuissimâ subconicâ stylo vix exserto, centralibus paucis  
is cor. limbo subcampanulato tubo brevissimo anth. inclusis  
is. Recept. nudum planum punctulatum. Involucrum pluriseriale,  
s oblongis acutis disco subbrevioribus. Achænia compressa gla-  
ppus 7-8-setus achænio brevior, setis crassiusculis barbellatis.  
us Indicæ annuæ erectæ ramosæ glabræ. Caules teretes. Folia  
oblonga basi attenuata apice dentata. Pedunculi axillares et  
les capitula pauca parva subcorymbosa congesta gerentes. Flo-  
sicco vid. flavidi. Pappus junior albus demum subrufescens.  
affine Conyzæ et Grangææ. — Nomen alludit ad Thespidem  
um heroem mythologicum qui filias, ut receptaculum nostrarum  
a corollas fœmineas, 50 protulit.

**VARICATA**, caule à basi ramosissimo, ramis divaricatis, pedunculis folio-  
ribus. (1) in Indiæ or. provinciis Silhet et Nepaul legit cl. Wallich. Co-  
varicata Wall.! cat. et herb. n. 3238. comp. 348. Herba 4-5 poll. longa.  
comm. ab hon. Cur. merc. angl. Ind. or.)

**RECTA**, caule basi simplici strictè erecto apice corymboso-ramoso, ramis  
è erectis, pedunculis folio longioribus. (1) in Indiæ orient. prov. Silhet.  
sinapifolia var. G. Wall.! cat. et herb. n. 3227. comp. 347, excl. aliis litt.  
acrocephalam latifoliam referendis. Caulis 12-15 poll. longus. (v. s. comm.  
rior.)

**KARELINIA** Less. ined. ex *Fisch et Mey. in litt.* 1833. — *Serratula*  
sp. *Pall. Willd.*

alum multiflorum heterogamum, fl. ambitus multiserialibus  
s filiformibus truncatis, centralibus 8-10 hermaphroditis 5-  
recept. nudum, planum. Inv. cylindrici squamæ ovato-lanceolatæ  
dpressæ. Antheræ basi ecaudatæ. Stylus fl. fœm. gracilis glaber,  
vergentibus elongatis, fl. herm. inclusus longè hispidulus lobis  
s erectis. Achænia erostris cylindrææ. Pappus fl. fœm. 1-serialis,  
uissimis elongatis capillaribus, fl. hermaphr. setis subplurise-  
ice barbellatis. — Herba Orientalis erecta ramosa glabra. Folia  
oblongo-lanceolata integra margine subscabra. Capitula corym-  
lindrica non malè extus Centauræ cerinthefoliæ seu Acroptili  
ferentia. Flores purpurascens. — Genus affine Conyzæ sed  
oto involucre teretiusculo faciliè distinctum.

**ASPA** (Less. ined.). 2/ in depressis salsis prope mare Caspium olim legit  
et nuperius ad ostia fluvii Ural cl. Karolin. *Serratula Caspia* Pall. voy.  
397. vol. 2. t. 13. p. 470. *Serratula Caspia* Willd. sp. 3. p. 1643.  
hipedalis, corymboso-ramosa. (v. s. comm. à cl. Fisch et Mey.)

## CCX. BERTHELOTIA.

ulum multiflorum heterogamum, fl. radii pluriserialibus fœmi-  
nissimè tubulosis 2-3-denticulatis, disci paucis (5-12) majori-  
dentatis hermaphr. sæpè abortivo-sterilibus. Recept. planum

punctatum nudum. Invol. ovalis squamæ pluriseriales imbricatæ adpressæ, ext. latæ breves obtusæ, intimæ lineares disco subæquales acutæ scariosæ. Antheræ ecaudatæ. Stylus radii gracilis glaber exsertus apice bifidus, disci papillosus subinclusus bifidus, ramis papilloso-hispidis. Achænia sessilia erostria teretiuscula glabra. Pappus 1-serialis, radii setis tenuissimis vix scabridis basi subdistinctis, disci setis rigidis basi subpaleaceis concretis apice barbellato-scabris. — Fruticulus India et Africæ æquinoctialis indigena, erectus ramosus, in totâ superficiei pube brevissimâ subadpressâ molliter canescens. Folia alterna sessilia lanceolata mucronata rigidula integerrima aut hinc inde ad apicem acutè serrata nervosa. Capitula ad apices ramorum densè corymbosa oblonga. Flores purpurascens. Pappus albus. Genus affine Conyzæ sed differt habitu et pappo fl. centralium polyadelpho, paleis nempe linearibus basi concretis. — Fruticulum elegantem dicavi cl. Berthelot floræ faunæque Canariensis strenuo illustratori.

1. *B. LANCEOLATA*.  $\frac{3}{4}$  Frutic. pedalis. An species forsitan duas admittenda?

a. *Senegalensis*, capitulis 5-6 lin. longis, pappi disci setis evidenter basi concretis quasi polyadelphis. In arenosis circa Lamaar et Gandiolo mense junio legit cl. Perrottet. Conyza proteiformis Perr. in litt. Folia infus. a Perr. in litt. magis serrata. (v. s. comm. à el. inv.)

ß. *indica*, capitulis paulò minoribus et confertioribus pappi setis brevibus et obscurius polyadelphis. — in Indiâ orientali ad ripam Gengis prope Cownpore et in ripâ fl. Jumma leg. cl. Wallich et Hamilton. Conyza rubra Ham. herb. Conyza lanceolata Wall. cat. et herb. n. 2991. comp. n. 101 Saussurea mucronata Spreng in litt. ad Wall. (v. s. comm. ab hon. C. Merc. angl. Ind. or.)

CCXI. *LÆNNECIA* Cass. dict. 25. p. 91. 37. p. 462 et 482. Less. syn. 203. — Conyzæ sp. Kunth.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus femineis tenuissimis apice 2-3 dentatis, disci hermaphroditis 5-dentatis. Recept. ebracteolatum punctulatum. Inv. campanulatum imbricatum. squamis linearibus apice subulato-acuminatis glabris coloratis. Antheræ ecaudatæ. Achænium erostre oblongo-obovatum compressum villosus Pappus conformis setosus biserialis, serie ext. brevi multisetosâ, int. longâ pilosâ. — Herbarum Americanarum lanatæ gnaphaloides. Folia alterna oblongo-linearia dentata aut incisa. Capitula ad apices ramulorum ætherium sæpius solitaria.

1. *L. GNAPHALIOIDES* (Cass. l. c.), caulibus diffusis à basi ramosis, foliis oblongo-linearibus inciso-dentatis, capitulis hemisphæricis.  $\frac{1}{2}$  in locis altis monte Cocollar Cumanensium. Conyza gnaphalioides H. B. et Kunth nov. gen. sm. 4 p. 73. t. 327.

2. *L. FILAGINOIDES*, caule simplici stricto, foliis oblongo-linearibus parvè dentatis, capitulis ovato-hemisphæricis. (1) circa urbem Mexici legit cl. Berlandier augusto flor. (pl. exs. n. 820). Priori caracteribus evidenter affinis sed prænotas suprâ indicatas diversa. Capitula dimidiò minor, pedunculis magis hænatiss, etc. (v. s.)

3. *L. PARVIFOLIA*, caulibus diffusis-ramosissimis, foliis oblongo-lanceolatis acutè inciso-dentatis, capitulis ex axillis ortis pedicellatis subsessilibusve in racemum subdigastis. (1) in Mexici valle Toluccanâ octob. flor. legit cl. Berlandier pl. exs. n. 1124!). Flor. herm. 2-3 in centro capituli, feminei circ. 12-15. Caulibus diffusis ad *L. gnaphalioidem* accedit sed folia 2-3 lin. longa, capitula dimidiò et ultra minor. (v. s.)

CCXII. CONYZA *Less. syn. 203.* — *Echenbachia Manch. meth. 573.* — *Dimorphanthès et Fimbrillaria Cass.* — *Conyzæ sp. Linn.* — *Conyzæ, Baccharidis et Erigerontis sp. auct.*

Capitulum multiflorum heterogamum, fl. radii multiseriis scæmineis corollâ tenuissimâ filiformi truncatâ aut breviter 2-3-dentatâ, disci paucis masculis tubulosâ 5-dentatis. Recept. ebracteolatum punctulatum seu fimbriiferum planum, aut convexum. Inv. squamæ pluri-seriales. Anthæræ ecaudatæ. Achænium compresso-planum sæpius glabrum basi attenuatum. Pappus 1-serialis; setis filiformibus vix scabris. — Herbæ ferè omnes Gerontogæ. Caules erecti ramosi teretes. Folia varia sæpè variè incisa semiamplexicaulia. Capitula pedunculata corymbosa aut subpaniculata. Flores flavi. Pappus rufus aut rufescens. — Differt ab *Erigeronte* et *Nidorellâ* ligulis plantæ nullis; à *Blumiæ* achæniis compresso-planis et antheris ecaudatis.

## Sect. I. ANDRYALOIDES.

Recept. alveolare, alveol. marginibus integris. Achænium pappi evid. scabri longitudinè aequale aut ferè longius (nec ut in cæter. multò brevius). — Herba Indica subacaulis lanato-tomentosa, scapis 1-cephalis. — An fortè genus?

1. *C. ANDRYALOIDES*, tota tomento lanuginoso cana subacaulis, foliis subradicalibus petiolatis obovatis obtusis basi subcuneatis grossè et parè dentatis, scapis 1-cephalis folia triplò circ. superantibus foliola linearia acuta integra paucâ gerentibus, inv. squamis biserialibus linearibus acutis dorso puberis vix achænio longioribus. 2 $\frac{1}{2}$  in Ind. prov. boreali occident. legit. cl. Royle. Herba 2-3 poll. longa subcæspitosa verisim. alpina aprica. Achænia compressa elongata villiosissima. Pappus 1-serialis scaber achænio ferè brevior, corollâ longior. Fl. feminei tenues pluriseriales, centrales steriles androgyni, anth. ecaudatis, stylo simplice inclusi. (v. s. comm. à cl. Royle.)

Sect. II. DIMORPHANTHES. — *Dimorphanthès Cass. dict. 13. p. 254; 25. p. 93; 37. p. 461 et 481.*

Recept. planum nudum aut brevissimè alveolato-subfimbriiferum. Achænia pappo setiformi multò breviora.

\* Species Americane. (*Pleræque disco hermaphrodito nec masculo donatæ et forsân sectionem propriam generis constituentes.*)

2. *C. TRIPLINERVIA* (Less. in *linnæâ* 1831. p. 137), herbacea glaberrima, caule erecto sulcato-angulato, foliis petiolatis ovato-lanceolatis basi cuneatis et integerrimis apice acuminatis serratis triplinerviis subtus punctatis, paniculâ corymbosâ polycephalâ, capitulis pedicellatis, invol. squamis lineari-lanceolatis acutis, achænio glabro subcompresso pappo brevior. 2 $\frac{1}{2}$  in Brasiliâ merid. (Sellow!), ad fretum Sanctæ-Catharinæ et Rio Janeiro (Less.), prov. Rio-Grande (h. Imp. Bras. n. 785). Folia 3 poll. longa, pollicem lata. Pappus ♂ apice subclavellatus, ♀ filiformis, uterque in specim. meo albus. Fl. ♂ 3-7. (v. s. comm. à Mus. reg. Berol.)

3. *C. ARGUTA* (Less. in *linnæâ* 1831. p. 138), herbacea glaberrima, caule erecto subsulcato, foliis vix petiolatis lineari-lanceolatis acuminatis per totum ambitum argutè et regulariter serratis uninerviis impunctatis, paniculis corymbosis polycephalis, capitulis pedicellatis, invol. squamis lineari-lanceolatis acuminatis. 2 $\frac{1}{2}$  in Brasiliâ meridionali (Sellow!), prov. Rio-Grande (h. Imp. Bras. n. 783). Folia 3 poll. longa, 3 lin. lata. Capitula ferè *C. triplinervia* sed flores masculi 1-3 undè primâ fronte ad *Baccharides* vergit. (v. s. comm. à Mus. reg. Berol.)

4. *C. THESIOFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 75), suffruticosa basi ra-

mosa, ramis erectis simplicibus glabris angulatis, foliis sessilibus angustè linearibus acutis integerrimis 1-nerviis utrinquè adpressè pilosis, corymbis terminalibus simplicibus, capitulis pedicellatis, invol. squamis 2-3-seriatis subæqualibus linearibus apice subulatis, ext. hispidis.  $\zeta$  in Mexico?

5. *C. KRIGEROIDES*, herbacea erecta ramosissima, caule striato villosa, foliis sessilibus lineari-sublancoeolatis utrinquè attenuatis integerrimis basi subtriplicinerviis pubescentibus, paniculâ diffusâ parè foliosâ amplâ, capitulis pedicellatis, invol. squamis 1-2-seriatis linearibus acuminatis puberis post anthesin omnino deflexis. (1) ? in Brasiliâ circa Bahiam. Simillima hinc *Erig. Canadensis*, illinc *Conyza fastigiata*. Recept. foveolatum omnino nudum vix ultra lineam diam. Pili caulis sublati, ramorum adpressi. Folia 2-2  $\frac{1}{2}$  poll. longa, 3 lin. lata, infima hinc indè dente notata. (v. s.)

6. *C. SINUATA* (Ell. sketch 2. p. 323), herbacea pilosa scabrinuscula, foliis infer. lanceolatis profundè sinuatis, lobis ovalibus acutis, super. linearibus integerrimis, capitulis paniculatis, invol. imbricatis squamis lineari-subulatis. (1) frequens circa Charleston Carolinæ. Flores albi centrales hermaphroditi. Achænia oblonga angulata pilosa.

7. *C. CHILENSIS* (Spreng. nov. prov. 1818. p. 14), caule herbaceo erecto subsimplici tereti striato-nervoso hirsuto, foliis radicalibus patulis oblongis basi attenuatis obtusis sinuatis, caulinis paucis semimplexicaulibus subhirsutis lanceolatis acutis subintegerrimis, capitulis paucis corymbosis latissimis.  $\zeta$  ? in Chili et in Brasiliâ (Chamisso!), prope Bahiam (Salzm. Blanch!), et ad portum Mulgrave Amer. bor. occid. (herb. Hænke!). — Link enum. 2. p. 322. Less. in linnæa 1831. p. 136. *Conyza procera* Desf. cat. h. Par. 1829. p. 164. *Dimorphanthus procera* Cass. bull. philom. 1821. p. 175. dict. 25. p. 93. Fl. innumeri pallidè flavi centrales ferè omnes hermaphr. et plerique fertiles. Pappus in specim. spontaneis rufus, in cultis albidus. Recept. punctulatum, 4 lin. diam. (v. s. sp. et v. c.)

$\beta$ . *monocephala*, caule foliisque glabrinusculis, capitulo solitario. — in Chili cum priore legit cl. de Chamisso. An fortè sp. propria? (v. s.)

8. *C. LINEARIS*, herbacea ramosa, caule pilis patulis hispido ad apicem ramorum steriliùm confertè hirsuto, foliis linearibus integerrimis acutis sæpè spirèliè tortis utrinquè pube brevè confertâ cinereis basi plus minùs villosis, paniculâ racemosâ, capitulis longè pedicellatis, invol. squamis biserialibus linearibus dorso puberis apice glabrinusculis.  $\zeta$  ? in pratis et arvis sabulosis ad Quintero Chilensium legit cl. Bertero januar. flor.; etiam video sine loci design. in h. Dombey. Capit 3 lin. longa, matra 6-7 lin. lata. Recept. subalveolare nudum 3 lin. ferè diam. Affinis *C. diversifolia* ex foliis super. et *C. Chilensis* capit. majusculis. (v. s.)

9. *C. ALBIDA* (Willd. herb. ex Spreng. syst. 3. p. 514), herbacea erecta, caule villosa striato supernè ramoso-paniculato, foliis sessilibus lineari-lanceolatis acuminatis apice remotè et grossè serratis canescenti-pubescentibus, paniculâ foliosâ laxâ, capitulis pedicellatis, inv. squamis 1-2 seriatis linearibus acuminatis dorso pubescentibus. (1) ex Salzm. in Brasiliâ (Cham.), ad Bahiam (Salzm.), Rio-Janeiro (Beyr. Blanch?) et Sanctam-Catharinam (Mac-Rae in h. Lindl.). — Less. in linnæa 1831. p. 136. Habitus *Erig. Canadensis* seu *Boussierensis*. Affinis *C. diversifolia*. Recept. lineam diam., alveolatum, ferè fimbriatum. (v. s.)

10. *C. DIVERSIFOLIA* (Weinm. in florâ 1820. p. 611), herbacea ad collum suffruticosa erecta tota villosa-cinerea, caule simplici pilis confertissimis hirsuto, foliis pube breviori velutino-villosis subtus hirsutis elongato-linearibus acutis. Infer. hinc indè grossè serratis, cæteris integerrimis, paniculâ oligocephalâ pubescente, capitulis pedicellatis, invol. squamis linearibus pube adpressâ parè vix cinerea.  $\zeta$  in pascuis Chilensibus ad S. Fernando legit cl. Bertero. C. ambigua Hook. et Arn. in bot. Beech. p. 57? excl. syn. Plantam descr. ex specim. spontaneis 1-2-pedalibus et foliis 3 poll. longis, 4 lin. latis; sed ut vid

descr. Weinm. in plantâ cultâ caulis 5-6-pedalis, folia 7-9 poll. longa, radic. pinnatifida. (v. s.)

11. *C. ? caryopodioides*, rhizomate suffruticoso; caule herbaceo ramosissimo basi pilosiusculo, ramis virgatis glabresculis, foliis lineari-lanceolatis utrinque acuminatis sparse villosis membranaceis, infer. apice trilobis, mediis remotis et grosse serratis, summis integerrimis, paniculis elongatis apice corymbosis, capitulis longè pedicellatis, invol. squamis linearibus.  $\frac{1}{2}$  in Brasiliæ prov. Rio de Janeiro in Serra dos Orgaos legit cl. Vauthier (pl. exs. n. 324). Recept. angustum nudum. Flor. ambitus numerosi teretes crassiusculi stylo non exserto, masculi pauci centrales vix dissimiles. Achænium compressum. Papp. 1-ser. (v. s.)
12. *C. coronopifolia* (H. B. et Kunth nov. gen. am. 4. p. 70), herbacea erecta canescenti-hispida, foliis sessilibus pinnatipartitis, lobis linearibus acutis basi subinciso-lobatis, capitulis pedicellatis in corymbos terminales ramosos dispositis, invol. squamis subriserialibus aequalibus linearibus apice subulatis.  $\frac{1}{2}$  in altâ planitie Mexicanâ ad littus lacus Texcucensis. Herba bipedalis. Cap. 2  $\frac{1}{2}$  lin. longa. Flor. disci 30-35.
13. *C. hispida* (H. B. et Kunth nov. gen. am. 4. p. 71), herbacea erecta sparse hispida, foliis sessilibus lineari-lanceolatis mucronatis integerrimis, capitulis longè pedicellatis laxè corymbosis hemisphæricis, invol. squamis vix imbricatis linearibus acuminatis dorso hispidulis.  $\frac{1}{2}$  in altâ planitie Mexicanâ ad littus lacus Texcucensis (H. B. et K.), circa urbem (Berl. ! pl. exs. n. 537). Folia sesquipoll. longa, 3-4 lin. lata. Capitula 3-4 lin. diam. Discus circ. 10-florus. (v. s.)
14. *C. obtusa* (H. B. et Kunth nov. gen. am. 4. p. 71), herbacea subcæspitosa canescenti-hispida, caulibus sterilibus subdiffusis, floridis erectis, foliis basi petioliformibus integris apice pinnatipartitis lobis oblongis integerrimis obtusis mucronatis, capitulis racemoso-corymbosis vix pedicellatis, invol. squamis vix imbricatis linearibus acutis dorso hispidulis.  $\frac{1}{2}$  inter urbem Mexici et Tolucam (Berl. ! pl. exs. n. 537). An *Conyza gnaphalioides*? Andrieux pl. exs. mex. n. 274 non Kunth ab hac satis differt? Pappus enim 1-serialis. Folia præsertim ad basin conferta. Capitula ferè *C. sophiæfoliæ*. (v. s.)
15. *C. cardaminifolia* (H. B. et Kunth nov. gen. am. 4. p. 71), caulibus herbaceis cæspitosis procumbentibus pilosis, foliis petiolatis profundè pinnatifidis pubescenti-pilosis, laciniis spathulato-lanceolatis acutis integerrimis, capitulis pedicellatis corymbosis, invol. semiglobosi squamis lanceolato-linearibus apice subulatis.  $\frac{1}{2}$  in ruderatis frigidis ad pagum Mulato Quitensium in radicibus montis Cotopaxi. Flores albi centrales 25-50 hermaphroditi. Affinis *C. coronopifoliæ*.
16. *C. subdecurrens*, herbacea erecta stricta puberula, canle simplici tereti, foliis adnato-sessilibus breviter decurrentibus ligulato-linearibus hinc inde irregulariter et acutè dentatis confertis demùm sæpè inflexis, capitulis brevissimè pedicellatis in paniculam angustam foliis intermixtam digestis, invol. squamis biserialis linearibus acuminatis dorso pubero-hirtellis.  $\frac{1}{2}$  ? in Mexico (Hænke!) ad Cordilleras Guchilapanas (Berl. ! pl. exs. n. 1044). Caulis circ. pedalis. Folia 10-12 lin. longa, 2-3 lin. lata, sæpè crispato-undulata. Capitula 2-3 lin. lata, sæpè crispato-undulata. Capitula 2-3 lin. diam. Flores disci circ. 10. (v. s.)
17. *C. kunthiana*, herbacea pusilla cæspitosa repens, foliis confertis sessilibus lineari-spathulatis obtusis integerrimis albo-lanatis, capitulis terminalibus solitariis sessilibus, invol. squamis subscariosis obtusis, inter. oblongo-linearibus.  $\frac{1}{2}$  in regione alpinâ regni Quitensis? *C. pusilla* H. B. et Kunth nov. gen. am. 4. p. 69 non Houtt. Flores centrales hermaphr. Folia 3-4 lin. longa.
18. *C. myosotifolia* (H. B. et Kunth nov. gen. am. 4. p. 69), caule herbaceo supernè corymboso polycephalo hispido-hirto, foliis sessilibus spathulato-oblongis obtusis hispidulis, capitulis corymbosis pedicellatis, invol. semiglobosi squamis linearibus subulatis.  $\frac{1}{2}$  in temperatis juxta La Sequia Novo-Granaten-



nium ad ripam Smittæ. Flores albi centrales hermaphr. Caulis bipedalis superne angulatus.

19. *C. LYRATA* (H. B. et Kunth nov. gen. am. 4. p. 70), herbacea ramosissima viscosa glanduloso-hirta, caule sulcato, foliis sessilibus ovato-oblongis basi lyratis dentatis, pedunculis terminalibus et axillaribus 3-5-cephalis, invol. subcampanulatis squamis linearibus acuminatis apice hispidis disco longioribus.  $\frac{1}{2}$  in humidis prope Guayaquil Quitensium. Flores albi in invol. 220-250, centrales hermaphroditi. Herba 2-3 pedalis.
  20. *C. SORNIIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 72. t. 326), herbacea stricta virgata hispidula, foliis pinnatipartitis, lobis linearibus apice 2-3-fidis obtusiusculis, capitulis glomerato-spicatis, fasciculis in paniculam dispositis, invol. squamis vix imbricatis linearibus acuminatis hispidulis.  $\frac{1}{2}$  ex H. B. et K., fortè (1) ex spec. sicco. — in valle Mexicanà (Berl. pl. exs. n. 694) inter Chalchicomula et lacum Texcoccensem (H. B. K.). Radix gracilis. Caulis 1-2-pedalis. Capitulum 2 lin. diam. Discus circ. 5-florus. (v. s.)
  21. *C. PULCHELLA* (H. B. et Kunth nov. gen. am. 4. p. 72), herbacea paniculata ramosa hirtella, foliis sessilibus pinnatipartitis, lobis linearibus integerrimis obtusis, capitulis pedicellatis corymboso-subpaniculatis, invol. squamis biaristatis lanceolato-linearibus apice subulatis.  $\frac{1}{2}$  in regione temperatâ circa Moran Mexicanorum. Herba semipedalis. Folia 4 lin. longa. Flores in invol. circ. 100, quorum 6 centrales hermaphroditi.
  22. *C. APURENSIS* (H. B. et Kunth nov. gen. am. 4. p. 73), herbacea erecta superne ramosa hispidula, foliis sessilibus spathulato-oblongis acutis argutè serratis, corymbo fastigiato polycephalo, invol. semiglobosi squamis linearibus apice subulatis.  $\frac{1}{2}$  in humidis prov. Variensis ad ripas flum. Apures. Herba bipedalis. Affinis *C. coronopifoliz*. Flores albi centrales hermaphroditi.
  23. *C. FLORIBUNDA* (H. B. et Kunth nov. gen. am. 4. p. 73), herbacea erecta ramosissima glabriuscula, foliis petiolatis lanceolatis acutè remotè et argutè dentatis utrinque sparè et adpressè puberis, corymbo foliato, invol. subsemi-globosi squamis linearibus acutis glabris.  $\frac{1}{2}$  frequens circa urbem Quito. Flores albi 30-35 quorum 7-8 centrales hermaphr. Herba 3-pedalis gratè odora.  
 $\beta$ . caule hispidulo superne paniculato, foliis subintegerrimis.  $\frac{1}{2}$  circa Guancabamba Peruvianorum.
  24. *C. GRAHAMI*, caule imò basi suffruticuloso herbaceo subsimplici patens et sparè piloso apice confertè hispidò, foliis semialexicaulibus oblongo-linearibus basi integerrimis apice parè et grossè serratis glabriusculis secus nervum et ad margines villosò-ciliatis, capitulis 3-5 subsessilibus congestis, invol. squamis linearibus acuminatis dorso hispidulis.  $\frac{1}{2}$  in Bonariâ ad Mendoza. Affinis *C. Gouani*. Flores innumeri, centrales circ. 20 masculi? Pappus albidus invol. paulò longior. (v. s. cult. in horto Edimb. à cl. Grahami comm.)
- Species Gerontogæ, nempe in *Europâ, Africâ continentali aut Indiâ crescentes.*
25. *C. ALTAICA*, caule herbaceo ramosissimo paniculato parè piloso-setoso, foliis sessilibus oblongo-linearibus setaceo-ciliatis, capitulis solitariis ramosis terminantibus, bracteis foliaceis invol. squamis referentibus sed capitulo longioribus evid. ciliatis invol. cingentibus, invol. squamis linearibus disco æqualibus apice subrecurvis. (2) in Sibiricâ altaicæ planitiebus prope Loktewak. *Erigonum ciliatum* Ledeb. ic. alt. ill. 1. t. 100. fl. alt. 4. p. 92. Cor. radii tenues, albi. ex cl. Ledebour brevissimè ligulatæ, ex specim. observato breves truncatæ, undè ad Conyzam species meo sensu referenda. (v. s. comm. à cl. C. A. Meyer.)
  26. *C. GOUANI* (Willd. sp. 3. p. 1928), rhizomate lignoso, caule glabriusculi herbaceo erecto, foliis lanceolatis apice serratis margine ciliato-scabris, lobis obovatis, summis integerrimis, capitulis ad apices caulium et ramorum congestis, invol. squamis lanceolatis margine membranaceis.  $\frac{1}{2}$  in insulis Canariæ

sibus Palmā et Teneriffā. *Erigeron Gouani* Linn. mant. 469. Jacq. hort. Vind. t. 79. Gouan illust. 66? *Dimorphantes Gouani* Cass. dict. 13. p. 255. *Baccharis hieracifolia* Lam. *Baccharis arunca* Schrank. *Erigeron scariosum* Moench. (v. s. et v.)

β. *diffusa*, caule demissiore ramosiore subdiffuso. ½ *Conyza diffusa* Poir. suppl. 2. p. 338. (v. s. comm. à cl. Broussonet ex insulis Canariis.)

27. *C. ROYLEI*, caulibus herbaceis adscendentibus simplicibus teretibus velutino-subhirsutis, foliis radicalibus petiolatis ovalibus, caulinis basi attenuatis obovatis, omnibus grosse dentato-incisis villosis-subhispidis, capitulis racemosis pedicellatis subsecundis, invol. squamis oblongo-lanceolatis acutis, achæniis compressis in disco glabris margine subciliatis. ½ In Indiæ or. prov. boreali-occid. legit cl. Royle. Herba 6-8 poll. Cor. masculæ flavidæ. (v. s. comm. à cl. Royle sub n. 135.)

28. *C. RUBASCENS*, caulibus herbaceis simplicibus adscendentibus cum foliis et involacris pube minimā brevissimā velutino-pubescentibus, foliis infer. in petiolum longum attenuatis obovato-oblongis, super. brevè basi attenuatis ovali-oblongis acutis, omnibus parvè et minutè dentatis, capitulis racemosis breviter pedicellatis, invol. squamis oblongis mucrone subacutis, achæniis compressis utrinque hispidulis. ½ In Indiæ or. provinc. boreali-occident. legit cl. Royle. Affinis sed diversa videtur à *C. Roylei* quæcum in herb. conjuncta sub n. 135 et ni fallor sub n. 133. (v. s. comm. à cl. inv.)

29. *C. BOVÉI* (DC. in ann. sc. nat. 1834. bot. p. 261), glabra, caule erecto apice corymboso-ramoso tereti, foliis sessilibus semiaplexicaulibus oblongo-linearibus distantè dentatis obtusiusculis, corymbo laxo, invol. squamis linearibus acuminatis pedicellisque puberulis. — in humidis montis Sinai majorē florentem legit cl. Bové. Arabis dicitur *Gazani*. *Conyza Bovéi* pl. exs. n. 118. Herba fortè basi suffruticosa, circiter pedalis. Folia sesquipoll. longa, 2 lin. lata. Capitula 3-4 lin. lata. Fl. feminei innumeri tenuissimi stylo exserto bifido, cor. truncata, centrales 5-dentati pauci stylo simplici verisim. abortivo. (v. s. comm. à cl. invent. cum pl. aliis Arabicis et ideo nomine suo insign.)

30. *C. FASTIGIATA* (Willd. enum. suppl. 57.), caule herbaceo erecto tereti ramoso subcorymboso hinc inde piloso, foliis sessilibus subamplexicaulibus lanceolato-linearibus suberrulatis integerrimisve basi ciliatis, corymbo terminali oligocephalo, capitulis pedicellatis, invol. squamis subniseriis linearibus acutis disci longitudine. ① ex Senegal. habeo spec., etiam in Mauritaniā (ex Spreng. ! syst. 3. p. 514). *C. paniculata* Desf. ! cat. h. Par. 1815. p. 114. non Willd. An *Dimorphantes angustifolia* Cass. dict. 25. p. 95? Valdè affinis hinc *C. ambigua* illinc *C. Gouani*. (v. s.)

31. *C. AMBIGUA* (DC. fl. fr. suppl. p. 468), caule herbaceo erecto ramoso paniculato tereti hirtio, foliis linearibus sessilibus acutiusculis integerrimis aut parvè dentatis scabro-puberulis, paniculā polycephalā, capitulis pedicellatis invol. squamis subniseriis linearibus acuminatis disci longitudine hispidulis. ① in pratis et cultis Galliæ occitanicæ non infrequens, in Italiā circa Romam (Sebast. !), Neapolim (Guss. ! ) et Pisas (Savi !), in Hispaniā Gaditanā (Pic !), Græciā (h. ac. Mon. !). *Erigeron crispum* Poir. chl. narb. n. 470 ex Dub. ! bot. gall. 1. p. 266. *Erigeron hinfolium* Willd. sp. 3. p. 1955. *Erigeron contortum* Desf. h. Par. ex Pers. ench. 2. p. 40. *Erigeron Canadense* Ten. prod. ex syll. p. 428. Habitus *Eriger. Canadensis*. Sola species Europæa nunc in *Conyze* genere remanens! Recept. planum nudum. Flor. fœm. innum., masc. 5-6. Achænia teretiuscula glabra. (v. s.)

32. *C. CAFFRA*, caule herbaceo erecto glabro aut pilosiusculo, foliis semiaplexicaulibus oblongo-lingulatis obtusiusculis remotè serrulatis utrinquè glabris, pedunculis velutinis in corymbum laxum dispositis, invol. squamis linearibus subulatis velutino-canescentibus disci longitudine, cor. masc. circ. 25 apice hirtis, fœm. innumeris. — in Africæ austr. terra Caffrorum, circa Key, legit cl. Dregé ! Herba sesquipedalis. Folia 30-33 lin. longa, 7 lin. lata. Recept. planum

punctulatum, punctis fl. central. majoribus distantibus, fl. form. confertissimis minimis. (v. s.)

33. *C. ÆGYPTIACA* (Ait. kew. 3. p. 183), caule herbaceo erecto tereti hispido, foliis oblongo-spathulatis utrinque pilosis per totam longitudinem grossè et obtusè dentatis, capitulis subpaniculatis globosis, invol. squamis linearibus acuminatis. ① in *Ægypti desertis* Delil. ill. fl. æg. p. 25. *Erigeron ægyptiacum* Linn. mant. 112 (excl. Bocc. syn. et ideò patr. Sicula). Jacq. hort. Vind. 3. t. 19. *Conyza ægyptiaca* et *Baccharis Dioscoridis* Bôv! pl. exa. n. 303 et 304. *Erigeron serratum* Forsk. æg. 148. *Eschenbachia globosa* Moench meth. 573. *Dimorphanthes ægyptiaca* Cass. dict. 13. p. 256. An *C. spathulata* Horn. cat. h. hafn. 2. p. 805 et Link enum. h. berol. 2. p. 322. huc, jam snad. Horn., ut varietas addenda? For. flavi. Recept. punctulatum. *Achnenia compressa*. (v. s. sp. et v. c.)

34. *C. JAPONICA* (Less. syn. 204), caule herbaceo erecto simplici piloso, foliis sessilibus obovatis basi attenuatis apice obtusis serratis villosis, paniculâ terminali. ① in Japoniæ insulâ Nipon. Vulgò dicta *Foso*. Pappus ferrugineus. *Erigeron Japonicum* Thunb. fl. jap. 312.

35. *C. DENTATA* (Willd. sp. 3. p. 1928?), caulibus herbaceis erectis teretibus apice subramosis undique hispidissimis foliosis, foliis utrinque hirsutis oblongis, infimis basi attenuatis parçè inciso-dentatis, super. sessilibus semiamplexicaulibus à basi ad medium inciso-dentatis ferè semipinnatifidis, corymbo terminali composito polycephalo, invol. squamis linearibus acuminatis disco sublongioribus. ② in demissis humidis ad ripas fl. Senegal in regione Walo legit cl. Perrottet. Herba 12-15 poll. longa. Affinis *Ægyptiacæ*. Pappus rufescens. Rec. alveolato-subfimbriatiferum. (v. s. comm. à cl. Perrottet.)

♀. *subnuda*, caule multò minùs præsertim supernè folioso, foliis angustioribus minùs dentatis, corymbo minore confertiore, pappo albido. ② in demissis humidis et arenosis Senegambiæ ad Khorò legit cl. Perrottet. *Erigeron exstipulatum* Schum. gulan. p. 387? An species propria? Flor. form. ad apicem obliquè truncati. (v. s. comm. cum priorè.)

36. *C. AMOENA* (Link enum. h. Berol. 2. p. 322), caule herbaceo hirtò, foliis sessilibus ovalibus obtusis basi attenuatis denticulatis hirtis, paniculâ terminali arctâ. ② ad fluvium Congo. Herba 2-3-pedalis pilis albis glandulosis. Invol. squamæ hirsutissimæ æquales lineares. Cor. purpurascens. An *Blumen* sp.? An prioris var.?

37. *C. ASTEROIDES* (Wall. cat. et herb. n. 3052. comp. 162. non Linn.), caule ascendenti erecto piloso ramoso tereti, foliis oblongis pilosis, infimis basi attenuatis obtusis grossè paucidentatis, superioribus semiamplexicaulibus acutis subintegerrimis, ramis laxis subpaniculatis, capitulis solitariis aut paucis aggregatis, invol. squamis linearibus cuspidato-acuminatis flores æquantibus, recept. in ambitu annuliformi in centro convexo. ② in Indiâ orient. ad Monghir, Dhuuari, Assufgur, Hurduar, etc. Pappus pallidè rufus. *Achnenia compressa*. Flor. form. innumeri filiformes, masc. 15-20. Affinis *C. veronicifoliae*. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

38. *C. SEMIPINNATIFIDA* (Wall. cat. n. 3058. comp. 168), hirsuta, caule herbaceo erecto tereti apice ramoso-corymboso, foliis obovato-oblongis obtusis basi attenuatis grossè et obtusè dentatis aut infernè incisis, summis oblongo hinc inde acutè dentatis, corymbo laxo, capitulis pedicellatis, invol. squamis linearibus acutis disco brevioribus ciliatis demùm reflexis, recept. valdè prominulo convexo-depresso. — in Indiâ or. prov. Martaban ad ripas flum. Saluen legit cl. Wallich. Fol. 2 poll. longa, 8-9 lin. lata. Flor. flavi, form. innumeri teretes obliquè truncati, stigm. non exsertis, masc. circ. 12, anth. inclusis. *Achnenia compressa*. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

39. *C. VERONICIFOLIA* (Wall. cat. et herb. n. 3005. comp. n. 115), caulibus ascendenti-erectis teretibus parçè ramosis pilosis, foliis pilosis infer. oblongis subacutis basi in petiolum attenuatis grossè serratis, superioribus semiamplexi-

caulibus lineari-lanceolatis acuminatis apice serratis, corymbis terminalibus congestis 12-20-cephalis, invol. squamis linearibus acuminatis dorso scabris demum reflexis disco aequalibus. — in Nepaliâ legit indef. Wallich. Conyza congesta Spreng. in litt. ad Wallich. Collum lignosum. Radix gracilis. Caulis 6-12 poll. Cor. fem. innumera truncatæ albide stylo bifido, masc. purpureæ circ. 10. Achænia compressa ut in Erigeronte. Pappus rufus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

40. *C. PETIOLARIS* (Wall. cat. n. 3282. comp. 292), sparsè pilosa, caule herbaceo erecto tereti vix apice ramoso, foliis infer. petiolo longo villosissimo donatis ovalibus obtusis grossè dentatis basi cuneatis, summis sessilibus oblongis serrulatis acutis, capitulis ad apices caulis et ramorum congestis subcorymbosis, invol. squamis linearibus acuminatis discum sequantibus. (1) in Nepaliâ? Recept. nudum medio pilosiusculum. Flor. fem. innumeri filiformes stylo exserto bifido, masc. pauci. Achænia compressa obovalia. Pappus rufus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
41. *C. MULTICAULIS*, caulibus herbaceis è collo plurimis strictè erectis simplicibus teretibus basi pilosis apice puberulis scabridis, infimis basi breviter attenuatis, summis sessilibus subamplexicaulibus, corymbo racemoso conferto, invol. squamis linearibus acuminatis flores subæquantibus, recept. convexo ferè hemisphærico. — in Nepaliâ legit cl. Wallich. Conyza stricta Wall. cat. et herb. n. 3044. comp. 154. non Willd. Pappus rufescens. Achænia glabra compressa. Flor. fem. innumeri tenuissimi truncati, masc. in centro pauci. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
42. *C. VISCIDULA* (Wall. cat. et herb. n. 3006. comp. n. 116), caule herbaceo erecto tereti infernè glabro in ramulis breviter puberulo viscidulo, foliis ramis lineari-lanceolatis utrinquè acuminatis serratis puberulo-viscidulis, paniculâ laxâ foliis intermixtâ, capitulis pedicellatis, invol. 1-seriali 10-phyllò squamis linearibus acuminatis disco aequalibus dorso puberulis. — in Indiâ or. ad Noakote legit cl. Wallich. Herba videtur elata sed pars inf. mihi ignota. Caulis pennæ anserinæ crassitie et ultrâ. Flor. fem. innum. supra modum tenuissimæ, masc. purpurascens circ. 5. Pappus rufescens. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
43. *C. STRIATA* (Wall. cat. et herb. n. 3065. comp. n. 175), glabriuscula aut supernè minutissimè pubens, caule herbaceo erecto striato-nervoso ramoso, foliis lanceolatis aut oblongo-linearibus utrinquè acuminatis integerrimis aut vix suberratis, ramis foliosis in paniculam dispositis corymbo conferto polycephalo terminatis, invol. squamis dorso puberulis lanceolato-linearibus acuminatis disco brevioribus. — in Indiâ orient. ad Prome et in prov. Silhet legit cl. Wallich. Folia 1-2 poll. longa, 3-4 lin. lata. Flores fem. innumeri tenues truncati, masc. 4-5. Pappus in spec. Promenai pallidè rufescens, in Silhetenai rufus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
44. *C. ANGUSTIFOLIA* (Ham. in Wall. cat. et herb. n. 3071. comp. n. 181. non Lam.), caule herbaceo erecto simplici tereti substriato patentem piloso, foliis lineari-lanceolatis elongatis acuminatis basi attenuatis ad medium grossè paucidentatis secus nervum medium subtus villosis cæterum pubescentibus, summis linearibus integerrimis, paniculâ elongatâ laxâ, ramis foliosis, invol. squamis linearibus acuminatis florum longitudine. — in Indiâ or. sed locus ign. Herba elata subcineræa. Fol. 5 poll. longa, 4-5 lin. lata. Flor. flavi, ext. fem. filiformes, inter. circ. 10-12. Achænia compressa, Pappus albidus subflavidus. (v. s. in h. hon. Cur. Ind. or. nunc soc. lin. Lond.)
45. *C. ABSINTHIIFOLIA* (DC. in Wight contrib. p. 16), pilosiusculo-pubescentia erecta ramosissima, foliis obovato-oblongis mucronatis basi cuneato-attenuatis, inf. grossè pauci-dentatis, sup. integerrimis, paniculâ corymbosâ ramosissimâ polycephalâ, inv. squamis linearibus acuminatis florum brevioribus, fl. fem. numerosissimis vix ligulatis. — (1) in Indiâ or. montanis in Nepaliâ, Kamaon, Sirmore, Dendigul, Neelgerry (ubi dic. ex Lessch. *Ouvoua-Guadâ*), peninsulâ Indiæ or. (Wight! herb. 217), in prov. Indiæ bor.-occid. (Royle!

pl. exs. n. 132 et verisim. 129, 130 et 131). *Conyza micrantha* Spreng. in litt. ad Wall. *Baccharis trifurcata* Trevir. in nov. act. acad. nat. ear. 13. p. 201. *Conyza cuneata* Ham! herb. *Conyza montana* et *C. stricta* Heyn! herb. *Conyza pinnatifida* var. Roxb. cat. calc. 61. *Conyza trifida* Ham. ex Dou. *Erigeron pinnatifidus* var. Don. prod. 172 ex Wight! herb. Ambigit inter *Erigeron* et *Conyzam*. *Habitus* *Erig.* acris sed ligulæ nullæ. *Achenium* parvum glabrum lenticulare compressum. *Capitula* parva. *Pappus* rufescens. *Herba* variat foliis angustis et latioribus, integris dentatis aut subpinnatifidis, obtusis aut mucronatis. An fortè plures hic latent commixtæ? (v. s.)

46. *C. WALLICHI*, caule herbaceo erecto tereti supernè ramoso-corymboso glabro, ramulis ultimis pulverulento-puberis, foliis lanceolatis acuminatis basi attenuatis serratis glabris, paniculæ corymbosæ polycephalæ ramis asperè hirsutis, capitulis ad apices ramulorum sessilibus confertis, invol. squamis linearibus acuminatis disco paulò longioribus. — in Indiâ orient. *Conyza disticha* Wall! cat. et herb. n. 3024. comp. n. 134 non Spreng. *Flor. fem.* innumeri, mihi vix ex statu provento specim. noti, masc. 5-7. *Achenia* parva ovalia compressa glabra. *Pappus* rufus.

*a. serrata*, foliis ovali-lanceolatis grossè serratis. — *Conyza disticha* A. Wall! l. c. *Cacalia disticha* Heyn! herb. (v. s. comm. ab hon. Cur. Ind. or. et à cl. Royle n. 128.)

*β. serrulata*, foliis elongato-linear-lanceolatis tenuissimè serrulatis. — circa Onde legit cl. Wallich cat. l. c. litt. B. (v. s. comm. cum a.)

47. *C. TRAVANCORENSIS* (Wall! cat. n. 3067. comp. 177). caule herbaceo erecto tereti subtilissimè striato et puberulo, foliis subpetiolatis lanceolatis utrinque acuminatis subserrulatis glabris cum nervo medio subtus puberulo, ramis corymbosis oligocephalis, capitulis pedicellatis, invol. 1-2-serialis squamis lanceolatis acuminatis flores subæquantibus. — in Indiâ or. ad Travancore. Folia 3-4 poll. longa, 9-10 lin. lata. Recept. convexum subhemisphericum. *Flor. fem.* innum. filiformes, masc. 4-5. *Pappus* rufescens. *Achenia* ex immaturo statu non satis nota. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or.)

*β. brevifolia*, ramulis teretibus vix puberulis, foliis subsessilibus ovali-lanceolatis acutis, capitulis magis corymboso-confertis, invol. squamis magis acuminatis. — in Indiâ or. prov. boreali-occid. legit cl. Royle. Folia sesquipedalia longa, 5 lin. lata. *Achenia* minima compressa glabra. An species propria? (v. s. comm. à cl. Royle sub n. 120.)

48. *C. RUFA* (Wall! cat. et herb. n. 3079. comp. n. 189), caule herbaceo erecto tereti simplici à basi multiplici piloso, foliis villosis oblongis obtusè et inæqualiter dentatis obtusis, infer. basi in petiolum attenuatis, supremis basi cuneatis sessilibus, paniculâ corymbosâ, ramis apice corymbulosis, capitulis pedicellatis, invol. squamis ovatis latè albo-marginatis biserialibus cuspidatis floribus æqualibus. — in Ind. orient. ad Assuighur legit cl. Wallich. *Herba* pedalis. Folia sesquipedalia longa, 4-5 lin. lata. *Pappus* rufus. *Flor. fem.* innumeri, masc. 4-5. *Achenia* ex statu imperf. ignota. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or.)

\*\*\* *Species Madagascarienses aut Mauritanæ.*

49. *C. BOJERII*, suffruticosa erecta ramosa glabra, ramis teretibus, foliis oblongo-lanceolatis sessilibus basi attenuatis integerrimis aut hinc inde subdentatis, pedunculis axillaribus terminalibusve nudis minutissimè pubescentibus capitula 7-8 congesta gerentibus, invol. squamis ovatis obtusiusculis disco dimidiò brevioribus. *β* in sylvia Befourou dictis prov. Ré-tani-mena ina. *Madagasc.* legit cl. Bojer. Folia 2 poll. longa, 7 lin. lata. Flores flavi. *Achenia* glabra. (v. s. comm. à cl. Bojer.)

50. *C. ATTENUATA* (DC. non Wall.), fruticosa glabra, ramis teretibus apice sub herbaceis flexilibus, foliis sessilibus elongato-lanceolatis basi obtusis apice acuminatissimis serratis, corymbo composito polycephalo laxiusculo, invol. squamis linearibus acuminatis nitidulis. *β* in sylvia umbrosis Befourou dictis

- prov. Bé-tani-mena ins. Madagascar legit cl. Bojer. *Baccharis attenuata* Bojer! in litt. 1835. Folia  $2\frac{1}{2}$  poll. longa, 5 lin. lata. Pappus rufescens. Invol. squamæ nervo medio virescente, marginibus pallidis. (v. s. comm. à cl. Bojer.)
1. *C. VIRGATA* (DC. non Linn.), fruticosa erecta, ramis strictis virgatis teretibus pubescenti-hirtellis, foliis sessilibus lineari-lanceolatis acutis serratis pubescentibus subtus punctulatis, corymbo irregulari composito, capitulis ad apices pedunculorum congestis subsessilibus, invol. squamis rufo-scariosis acutis, fl. disci circ. 5. —  $\S$  in Madagascar detexit cl. Bojer. *Baccharis virgata* Bojer! in litt. 1835. Frutex 4-5-pedalis. Folia circ. 20 lin. longa, 4 lin. lata. Achænia compressa glabra. Flores mihi filiformes truncati, ex Bojer flavi marginales sæpè ligulati. An ideo *Nidorellæ* fortè species? (v. s. comm. à cl. inv.)
2. *C. FLEXILIS*, fruticosa glabra subscandens, ramis teretibus flexilibus, foliis sessilibus oblongo-lanceolatis basi attenuatis apice acuminatis integerrimis, paniculâ elongatâ, capitulis ad apices ramulorum congestis subsessilibus parvis, invol. squamis glabris scariosis rufescentibus acutis.  $\S$  in sylvia Béfouroun dictis ins. Madagascar prov. Bé-tani-mena super arbores subscandens. *Baccharis* Bojer! in litt. 1835. Flores flavi. Capitula parva in spec. juniora nec ideo satis ritè observata. Pappus rufus brevis. Folia 3 poll. longa, 8 lin. lata.
3. *C. HIRTILLA*, herbacea subscandens, ramis densè hirtellis, foliis petiolulatis lanceolatis utrinquè acuminatis remotè serratis utrinquè minutissimè scabridis, paniculis parvis axillaribus folio brevioribus, invol. squamis lanceolatis acuminatis scariosis rufescentibus glabris. (1)? in sylvia ins. Madagascar cum *C. flexili* legit cl. Bojer. Folia (petiolo 2-3 lin. incluso) 3 poll. longa, 8-9 lin. lata. Capitula parva. Flores juniores non ritè exam. (v. s. comm. à cl. Bojer.)
- 54? *C. AGERATOIDES*, suffruticosa erecta ramosa infernè glabrata, ramis densè hispidis cylindricis, foliis petiolatis ovatis acutis grossè dentatis utrinquè hirtellis, petiollis densè hispidis, capitulis ad apices pedunculorum 7-8 congestis, invol. squamis glabris linearibus acuminatis, flor. femininis pauciseriatis.  $\S$   $\mathcal{L}$  in campis et agris cultis prov. Emirnæ insul. Madagascar legit cl. Bojer. *Baccharis ageratum* refereas Bojer! An flores (mihi in specim. juniore non satis notis) fortè cœrulei? An à genere removenda et fortè ad *Plucheam* refer.? (v. s. comm. à cl. Bojer.)
55. *C. LINEARILoba*, caule erecto basi fruticuloso, ramis longè et densè hispidis, foliis parvis hispidis linearibus pinnatipartitis lobis 2-3-jugis distantibus linearibus integerrimis, summis indivisis, capitulis racemoso-glomeratis brevissimè pedicellatis, invol. squamis linearibus acuminatis dorso puberulis, fl. disci paucis.  $\S$  in insula Madagascar, Mauritiæ et Bourboniæ, præcipuè in agris cultis legit cl. Bojer. Flor. form. innumeri ut centrales pappo 1-seriali longo subrufescente donati. Recept. foveolatum. Achænia parva compressa glabra. (v. s. ex Madag. comm. à cl. Bojer.)
56. *C. CALOCEPHALA* (Bory! voy. isl. afr. 2. p. 395), suffruticosa tota hirsuto-hispida, foliis sessilibus oblongis integerrimis aut hinc indè subserratis, pedunculis folia superantibus 1-cephalis, invol. squamis biserialibus acuminatis dorso hirsutis, fructibus compressis obovatis, pappo rufo-colorato flocculis longiore.  $\S$  in insula Bourboniæ (Comm.! Bout!), in montanis prope locum dict. Bras-de-Pontesa (Bory!). *Erigeron rutilum* var. Poir. suppl. 5. p. 494. *Dimorphanthus bidentata* Cass. dict. 13. p. 256, et 37. p. 481. Suffrutex vix palmaris. Folia erecta 8-10 lin. longa, 2-3 lata, basi subattenuata utrinquè hirsuto-hispida. Achænia glabrinsculis compressa. Rec. brevissimè fimbriiliferum. (v. s.)
57. *C. ASPERA* (Bory! voy. isl. afr. 3. 174), suffruticosa tota hirsuto-hispida, foliis sessilibus semiamplexicaulibus oblongo-lanceolatis grossè serratis trinerviis, capitulis ad apices ramorum 4-5 aggregatis breviter pedicellatis, invol. squamis acuminatis dorso hirsutis, achæniis compressis oblongis glabris, pappo rufo flocculis longiore.  $\S$  in insula Bourboniæ montanis (Comm.! Bory!). *Conyza melastomoides* Lam. dict. 2. p. 90. *Erigeron scabrum* Pers. ench. 2. p. 432. n. 35 nec n. 23. *Erigeron* Jussæi Spreng. syst. 3. p. 519. Valdè affinis *C. ca-*

locephalæ et *C. lithospermifoliæ*. Achenia compressa. Recept. brevissimè fimbriiferum. (v. s.)

58. *C. LITHOSPERMIFOLIA* (Lam. dict. 2. p. 90), suffruticulosa pubescenti-hirsuta, foliis sessilibus semi-amplexicaulibus oblongis apice serratis basi trinerviis, ramis subvirgatis apice aphyllis 1-cephalis, invol. squamis acuminatis dorso puberulis, acheniis compressis glabriusculis, pappo rufo corollis longiore.  $\frac{1}{2}$  in summis montibus insulæ Bourboniæ (Bory! Bout.). *Elphægea hirta* Cass. dict. 14. p. 364 ex syn. Valdè affinis *C. asperæ* et *C. calocephalæ* sed distincta videtur. Recept. brevissimè fimbriiferum. (v. s.)

59. *C. SERICHA* (Bory! voy. isl. afr. 2. p. 376 non Ait.), suffruticulosa tota villis adpressis sericeis cinerea, foliis sessilibus oblongis basi ferè angustioribus mucronatis integerrimis, pedicellis folio paulò longioribus 1-cephalis ad apicem circ. 3 approximatis, invol. squamis dorso sericeo-villosis subacuminatis, pappo rufo corollis vix longiore.  $\frac{1}{2}$  in insulâ Bourboniâ legit cl. Bory et necnè comm. Hæc et seq. commixtæ adsunt in h. Desf. sub nom. autographo *Erigeron rutilans* Poir.! suppl. Ferè media inter *C. calocephalam* et *C. argenteam* (v. s.)

60. *C. ARGENTEA* (Lam. dict. 2. p. 87), suffruticosa tota villis adpressis sericeis argentea, foliis ovatis semiamplexicaulibus integerrimis macronulatis, capitulis ad apices ramorum subglomeratis breviter pedicellatis, invol. squamis sericeo-villosis acuminatis, pappo rufo flosculis subæquali.  $\frac{1}{2}$  in insulâ Bourboniâ collibus aridis. Folia obscurè trinervia 9 lin. longa, 6 lin. lata. Achenia compressa glabra. (v. s.)

61. *C. ANCHUSÆFOLIA* (Poir. suppl. 2. p. 339), suffruticosa procera hirsuta, foliis ovali-oblongis utrinquè acuminatis in petiolum breviter attenuatis apice serratis penninerviis, corymbo paniculato terminali polycephalo foliis breviorè, pedicellis capitulo æqualibus, invol. squamis extus glabriusculis, acheniis oblongis glabris, pappo corollæ longitudine subrufescente.  $\frac{1}{2}$  in ins. Bourboniâ (Bory! Bout.). *Conyza verbascifolia* Bory! voy. isl. afr. 2. p. 377 non Willd. Folia 5 poll. longa, 15 lin. lata. Flores rubescentes. Achenia compressa glabra. (v. s.)

62. *C. AMYGDALINA* (Lam. dict. 2. p. 90), fruticosa villis longis adpressis sericeis subcanescentibus, foliis petiolatis oblongo-lanceolatis utrinquè acuminatis densato-serratis penninerviis suprà subviridibus subtus sericeo-argenteis, pedunculis axillaribus folio multò brevioribus capitula 3-4 congesta gerentibus, invol. squamis dorso subhirsutis acuminatis, pappo subrufescente corollis longiorè.  $\frac{1}{2}$  in insulâ Bourboniâ prope Castrorum planitiem. Folia cum petiolo semi-pollicari 4 poll. longa, pollicem lata. Achenium ignotum et ideo genus dubium. (v. s. comm. à cl. Bory.)

### Sect. III. TUBIFERA.

Recept. fimbriiferum, fimbriis elongatis. Flores form. tubulosi cylindrici apice brevissimè 3-dentati discum superantes. — Species Bourbonica.

3. *C. LAURIFOLIA* (Lam. dict. 2. p. 89), fruticosa pube brevissimâ subserpiceâ pubescens, foliis oblongo-lanceolatis in petiolum attenuatis subintegerrimis penninerviis, corymbo paniculato amplo, ramis elongatis, pedicellis ultimis vix capitulo longioribus, invol. squamis extus pubescentibus margine membranaceis, pappo rufescente corollis paulò breviorè.  $\frac{1}{2}$  in insulâ Bourboniâ sylva (Bory! Bout.). Fimbriaria tubifera Cass. dict. 17. p. 55. Folia 4-5 poll. longa, pollicem lata. Flor. fœm. pluriseriales tubulosi elongati 3-dentati stylo non exserto, masc. pauci anth. semiexsertis basi muticis. Achenia ex statu immat. sericeo-villosa compressa. Pappus rufus. Recept. fimbriis longis aratis ornatum. (v. s.)

Sect. IV. FIMBRILLARIA. — *Fimbrillaria* Cass. dict. 17. p. 54.

Recept. totum fimbrilliferum, fimbrillis in centro longioribus sæpè setiformibus. — Species Capenses.

64. *C. PODOCETRALA*, caule imò basi suffruticoso multicauli subadscendente, ramis hispidis-scabris, foliis obovatis basi cuneatis integris apice grossè dentatis sparsè scabro-hirtellis, summis oblongis acutis, ramis apice undis 1-cephalis parçè hispidis, invol. squamis lineari-subulatis glabrinusculis disci longitudine.  $\frac{1}{2}$   $\frac{1}{2}$  in Africâ australi ad Zw. Stoorm et Wittbergen alt. 4-500 ped. legit cl. Drege! Recept. alveolatum brevissimè fimbrilliferum. Cor. mascula glabrâ numero fœm. subæquales. (v. s.)

*f. demissa*, caulibus pluribus demissis folisque magis hispidis. — ad Nieuwewelde legit cl. Drege. (v. s.)

65. *C. PINNATILORATA*, caule herbaceo erecto vix puberulo, foliis radicalibus petiolatis, caulinis sessilibus semiamplexicaulibus, omnibus glabrinusculis pinnatipartitis, lobis lanceolatis subdentatis in apice caulis integerrimis margine et secus nervos scabris, corymbo paniculato laxo terminali, pedunculis pubescentibus, achæniis glabris.  $\frac{1}{2}$  ad Cap. Bonæ-Spei post Thunb. legerunt cl. Burckell (cat. geog. n. 433!), Belanger (n. 80), Drege, Ecklon et Zeyh. *Erigeron pinnatum* Linn. f. suppl. 368. Thunb. prod. 153. fl. cap. 666. *Raccharis leucanthemifolia* Burm. f. herb. et ideò ex apha. *Xeranthemum leucanthemifolium* Burm. prod. 26? *Herba sesquipedalis*. Fol. rad. lobi ferè ovati sinuati obtusi. *Achænia* compressa ut in *Erigeronte*. (v. s.)

*f. hispida*, caule magis hispido, foliis vix ad medium incisais, capitulis paucioribus majoribus.  $\frac{1}{2}$  in Wittbergen alt. 3-400 ped. legit cl. Drege. (v. s.)

66. *C. PINNATIFIDA* (Less. syn. 204 non Roxb.), herbacea tota pilis patulis strigosa hirta, foliis caulinis pinnatipartitis lobis lanceolato-linearibus hinc indè dentibus lobulatis acutis distantibus incisais, capitulis 1-2 ad apices ramorum, achæniis puberulis.  $\frac{1}{2}$  ad Cap. Bonæ-Spei (Drege!) in collibus extra urbem (Th.), in planitie prope Zeekuhwallei (Ecklon! in h. Dunant), in Stellenbosch (Eckl.!) *Erigeron pinnatifidum* Thunb. fl. cap. 666. (v. s.)

67. *C. OSCURA*, herbacea erecta villis confertis subrigidis hirsuta, foliis obovato-oblongis sessilibus basi attenuatis subintegris semiamplexicaulibus apice pinnatifidis, lobis obtusis aut in parte caulis super. acutis, corymbis laxiusculis, achæniis subpilosisusculis.  $\frac{1}{2}$ ? in Africâ australi. Drege! Eckl. et Zeyh. (v. s.)

*f. calvescens*, foliis infer. hirsutis, super. cauleque scabris, lobis acutis margine subrevolutis.  $\frac{1}{2}$ ? in Lange-Kloof legit cl. Drege. An spec. propr.? (v. s.)

68. *C. INCISA* (Ait. kew. 3. p. 184), caule herbaceo? erecto puberulo, foliis petiolatis ovatis aut subcordatis subpiloso-viseosis dentato-incisais basi subauriculatis, corymbis terminalibus laxiusculis, achæniis glabris.  $\frac{1}{2}$   $\frac{1}{2}$  ad Cap. Bonæ-Spei. *Erigeron incisum* Th. fl. cap. 666. *Conyza incisa* et *Erigeron incisum* Willd. sp.-3. p. 1945 et 1962. *Raccharis ulmifolia* Burm. f. prod. fl. cap. 26. *Achænia* glabra, compressa ut in *Erigeronte*. Recept. breviter fimbrilliferum.

*a. stipulacea*, petiolo supra medium appendice utrinquè aucto, limbo terminali subcordato. — ad Zneeburgen Drege! *Conyza stipulacea* Desf. cat. h. Paris. 1829. p. 164. excl. patr. *Dimorphanthès stipulacea* Cass. dict. 25. p. 94. excl. patr. (v. s.)

*f. hirta* (Less. syn. 204), caulibus folisque magis hirsutis petiolo exappendiculato, limbo ovali. — ad Cap. Bonæ-Spei (Boiss. herb.), in Tulbagh, Uitenhage, Clan-William (Eckl. et Zeyh.) in Paarl (Drege!) *Erigeron hirtus* Th. fl. cap. 665. (v. s.)

69. *C. SCABRIDA*, tota pubescenti-scabra, caule subherbaceo erecto, foliis penninerviis lanceolatis acuminatis dentatis, inf. petiolatis, super. sessilibus brevissimè



auriculatis semiamplexicaulis, corymbo terminali composito.  $\frac{3}{4}$  ad Cap. Bonæ-Spei (Drege!) in distr. Uitenhage (Eckl. et Zeyh.). An *Erigeron* dentatum Burm.! prod. 26? Media inter *C. incisam* à quâ differt foliis lanceolatis  $1\frac{1}{2}$ -2 poll. longis,  $\frac{4}{5}$  lin. latis, et *C. ivæfolia* à quâ distinguitur foliis pubescenti-scabris nec glabris penninerviis nec trinerviis. *Achænia* compressa. Rec. in abruptu breviter alveolatum medio setis elongatis onustum. (v. s.)

*β. angustifolia*, foliis linearibus remotius dentatis. — Burch.! n. 382. Eckl. n. 771. (v. s.)

70. *C. IVÆFOLIA* (Less. in Linnæa 1831. p. 138), fruticosa, ramis teretibus pubescentibus, foliis lanceolatis acuminatis in petiolum attenuatis grosse serratis basi trinerviis glabris, corymbo terminali composito.  $\frac{3}{4}$  ad Cap. Bonæ-Spei in collibus ad occident. montis Tabularis (Th.), ad Dutois-Kloof (Drege!). *Baccharis ivæfolia* Linn. sp. 1204. excl. syn. et patria. Berg. cap. 275. Thunb. fl. cap. 666. Burm.! prod. 26. *Fimbrillaria baccharoides* Cass. dict. 17. p. 54. Pluk. amalth. t. 454. f. 10. pessima et dubia. Barth. act. hafn. 2. t. 27. Linnæus, qui citat phrasin Pluken. ubi dicitur *Africana*, asserit suam stirpem ex Virginia et Peruv. ortam, sed fortè sub eo nom. *Baccharidem angustifoliam* designavit. Nostra certè Capensis ex spec. spontaneo Drege à me suprâ citato et altero Eckloniano in herb. Dnnant servato, hortensibus simillimis. Recept. fimbriis in setis ferè desinentibus onustum. (v. s. sp. et v. c.)

71. *C. PUBIGILA* (Houttuyn! hist. nat. x. bl. 618. t. 69. f. 1), hispidulo-villosa cinerea herbacea ramosa, foliis superis semilibus ovalibus mucrone apicalatis integerrimis aut ad apicem mucronibus 1-2 auctis basi trinerviis, pedicellis 1-floris folii longitudine in racemum laxum dispositis, involucris linearibus acuminatis puberis demum reflexis. — ad Cap. Bonæ-Spei. Frustulum tantum vidi sed meo sensu distinctissimum. (v. s. in h. Delessert.)

† *Conyze auctorum non satis notæ pleræque verisimiliter è genere excludendæ.*

\* *Gerantogea*.

72. *C. ? ARABICA* (Willd. sp. 3. p. 1949), fruticosa stricta pubescens superne tomentosa, foliis decurrentibus linearibus integerrimis tomentosis, capitulis pedicellatis paniculatis, invol. squamis setaceis.  $\frac{5}{8}$  in Arabia felici. *Erigeron decurrens* Vahl symb. 1. p. 72. An *Pluchea*, an *Blumea* spec?
73. *C. ? DONGOLÆNSIS* (in Caill. Delil. pl. afr. voy. mer. p. 43), ramis villosis, foliis sessilibus oblongis dentatis sublyratis basi incisâ vel pinnatifidis, segmentis acutis. — circa Dongolâh in Nubiâ. Flor. fructusque ignoti et ideo genus dubium.
74. *C. ? CRISPATA* (Vahl symb. 1. p. 71), fruticosa glabra, foliis decurrentibus lanceolatis serrato-dentatis, alis dentato-sinuatis, pedunculis 1-cephalis.  $\frac{5}{8}$  in Arabia. *C. caule alato* Forsk. cat. pl. ar. p. 119. An *Pluchea*? An *Blumea* sp?
75. *C. ? FOLIOLOSA* (Willd. sp. 3. p. 1919, non Wall.), pubescens, caule herbaceo ramosissimo diffuso, foliis oblongis subintegerrimis subamplexicaulis, pedunculis alaribus 1-cephalis demum elongatis folio sextuplo longioribus. — in Indiâ or. Con. foliosa Pers. ench. n. 3. An *Conyza*? an *Blumea*? an *Pluchea*?
76. *C. ? HETEROPHYLLA* (Lam. dict. 2. p. 84), caule striato herbaceo ramoso, foliis petiolatis cinereo-villosis aliis subcordatis aliis rotundioribus, paniculis terminalibus. — In Indiâ or. (Sonn.), in agro Malabarico (Rheed. mal. 10. t. 61) An *Vernonia* species cinerea affinis?
77. *C. INCANA* (Willd. sp. 3. p. 1937), fruticosa tomentosa, foliis linearibus basi attenuatis subdentatis tomentosis, corymbo terminali.  $\frac{3}{8}$  in Arabia felici. *Erigeron incanum* Vahl symb. 1. p. 72. Invol. squamæ lineares. Pappus ferrugineus. An *Blumea*?
78. *C. ? LINIFOLIA* (Burm. prod. fl. cap. 26 non Linn.), vid. *Blumea* sp. ex capitulo à me olim dissecto, sed vix nota!
79. *C. ? LOUREIRI* (Poir. suppl. 2. p. 339), caule suffruticoso erecto simpliciter folio

lanceolatis integerrimis breviter petiolatis subtus tomentosis, pedunculis axillaribus polycephalis, invol. cylindrico, recept. nudo.  $\frac{1}{2}$  prope Cantonem Sinarum. Baccharis Sinensis Lour. coch. 2. p. 604. Conyza Chinensis Pers. ench. n. 16 nec n. 21. An Conyza? an Blumea?

80. *C. MOLLESCANA* (Gaud. voy. p. 465), glabra, ramis suffruticosis sulcatis, foliis oblongis acuminatis basi obtusis argutè et remotè denticulatis subtus pallidioribus, paniculis terminalibus foliatis, involucribus pubescentibus.  $\frac{1}{2}$  in ina. Moluccis ad Rawak. An Blumea sp.?
81. *C. ORIENTALIS* (Willd. sp. 3. p. 1931), herbacea hispido-scabra supernè ramosa, foliis serratis, infer. obovatis petiolatis, superioribus lanceolatis sessilibus, capitulis congesto-corymbosis terminalibus, invol. squamis subulatis.  $\frac{1}{2}$  in Armeniâ. *C. orient. Asteris attici folio flore luteo* Tourn. cor. 35.
82. *C. PUMIGERA* (Linn. mant. 113), caule herbaceo sparsè piloso, foliis oblongis subdentatis subpetiolatis, pedunculis lanatis subdichcephalis. — in Ind. orient. *Sonchus volubilis* Rumph. amboin. 5. p. 299. t. 103. f. 2, è Moluccis ortus et ad suam speciem à Linnæo relatus vix descriptioni respondet et ex iconè videtur potius Senecionea?
83. *C. STRICTA* (Willd. sp. 3. p. 1922), caule herbaceo erecto paniculato, ramis erectis, foliis lineari-lanceolatis basi attenuatis integerrimis pilosis obtusis cum acumine, corymbis fastigiatis. (1) in Indiâ orientali. Pappus rufescens. An Conyza? an Blumea?
84. *C. THYRSOIDRA* (Pers. ench. 2. p. 426. n. 8), herbacea, caule villosa, foliis pubescentibus obovatis sinuato-serratis infernè sublyratis, racemis foliosis terminalibus. — in Senegaliâ. Cæt. ign.

\*\* *Americanae.*

85. *C. CAYENNENSIS* (Spreng. syst. 3. p. 512), herbacea, foliis longè petiolatis subcordato-ovatis acuminatis glabriasculis venosis obtusè serratis, paniculâ terminali, capitulis subsessilibus. — in Cayennâ ex Martin in herb. Willd. Cæt. ign. †.
86. *C. CONICA* (Poir. suppl. 2. p. 341), fruticosa glabra, foliis subsessilibus coriaceis ovato-subrotundatis integris suprâ lucidis subtus pallidis, capitulis cymosis subsessilibus, invol. conici squamis acutis, pappo intensè rufo.  $\frac{1}{2}$  patr. et cæt. car. ign.
87. *C. DISTICHA* (Spreng. syst. 3. p. 509), fruticosa, foliis sessilibus oblongis dentatis subcoriaceis ramisque villosiusculis, spicis distichis foliatis terminalibus.  $\frac{1}{2}$  ad flumen magnum Brasilie. Cæt. ign. An Vernoniæ spec.? †.
88. *C. GRAMINIFOLIA* (Spreng. syst. 3. p. 515), herbacea, caule glaberrimo simpliciter, foliis basi vaginantibus linearibus acuminatis elongatis strictis, pedunculis terminalibus corymbosis squamatis, invol. squamis lanceolatis cuspidatis. — in Monte-Video. Cæt. ign.
89. *C. LAXA* (Spreng. syst. 3. p. 513), herbacea, foliis glabris petiolatis oppositis ovato-lanceolatis utrinquè attenuatis serratis margine scabris triplinerviis, corymbis axillaribus terminalibusque oligocephalis, invol. squamis laxis linearibus. — in Brasiliâ. An Baccharidea? an Eupatoriacea?
90. *C. LUCENS* (Poir. suppl. 2. p. 342), fruticosa tortuosa, ramis teretibus tomentosis, foliis petiolatis ovatis integris suprâ lucidis subtus incano-tomentosis, capitulis terminalibus agglomeratis, invol. squamis linearibus.  $\frac{1}{2}$  in Sancto-Domingo. *C. Domingensis* Willd. herb. ex Spreng. syst. 3. p. 508. Cæt. ign. †.
91. *C. MACROPHYLLA* (Spreng. syst. 3. p. 513), herbacea, foliis amplexicaulibus longissimis lanceolatis glabris sinuato-dentatis, paniculâ corymbosâ, invol. squamis linearibus lanceolatis. — in Monte-Video. Cæt. ign. †.
92. *C. MONTEVIDENSIS* (Spreng. syst. 3. p. 510), suffruticosa glabra, foliis linearibus acuminatis elongatis serratis, capitulis corymbosis terminalibus.  $\frac{1}{2}$  in Monte Video. An Baccharidis spec.? †.
93. *C. OBLONGIFOLIA* (Spreng. syst. 3. p. 512), herbacea, caule simplici stricto

superne aphyllis, foliis glaberrimis subintegerrimis oblongis obtusis sessilibus triplinerviis, inf. oppositis, paniculâ corymbosâ pubescente, invol. squamis oblongis obtusis. — in Brasiliâ ad Rio-Grande. Cæ. ign. †.

94. *C. PANAMENSIS* (Willd. sp. 3. p. 1942), fruticosa, ramis glabris striatis, foliis sessilibus ovato-lanceolatis scabris integerrimis triplinerviis reticulato-venosis, paniculâ terminali, invol. squamis oblongis. ♀ in Panamâ. Ex descr. valde manca videtur potius *Baccharis*?

95. *C. PUNCTATA* (Willd. sp. 3. p. 1937), fruticosa, ramis striatis glabris, foliis linearibus acuminatis basi attenuatis impresso-punctatis, pedunculis longissimi 1-cephalis. ♀ in Chili. Invol. squamæ lanceolata acutæ. Pappus rufus. Cæ. ign. An fortè *Baccharidis* spec.?

96. *C. RUVIDULA* (Spreng. nov. entd. 2. p. 141. syst. 3. p. 513), caule herbaceo erecto piloso, foliis brevissimè petiolatis oblongo-lanceolatis acutis utrinque pubescentibus integerrimis, paniculâ pyramidalî. — in Brasiliâ. Folia sub lente denticulato-ciliata. Involucrum imbricatum glabrum squamis inter. elongatis coloratis. Pappus rufescens. Flores non observ. †.

97. *C. SQUMATA* (Spreng. syst. 3. p. 515), herbacea, caule ramosissimo scabro. foliis sessilibus linearibus acuminatis integerrimis, pedunculis squamatis 1-cephalis, invol. squamis lineari-lanceolatis. — in Monte-Video. Cæ. ign.

98. *C. ? TRIDENTATA* (Poir. suppl. 2. p. 342), fruticosa glaberrima, foliis confertis linearibus basi attenuatis subpetiolatis apice subtridentatis obtusis, capitulis parvis sessilibus. ♀ in Sancto-Domingo. Cæ. ign. An *Baccharidis* species? †.

\*\*\* *Patriæ non satis notæ.*

99. *C. CILIARIS* (Spreng. nov. prov. 14. syst. 3. p. 509), fruticosa, foliis spatulato-lanceolatis subsessilibus glabris ciliatis, pedunculis terminalibus solitariis elongatis 1-cephalis villosis. ♀ Patria et cæ. car. ign.

100. *C. ? GLOMERATA* (Link. enum. h. Berol. 2. p. 322), caule herbaceo alato, foliis late lanceolatis obtusis basi attenuatis serrulatis pubescentibus scabris, capitulis glomeratis bracteâ obvallatis. — Patr. ign. An *Blumea*, *Pluchea* seu *Monstrelis* spec.?

101. *C. LAXIFLORA* (Desf. cat. h. Par. 1830. p. 400), caule herbaceo villosio paniculato, foliis lanceolatis ciliatis, infer. remotè serratis rameis sup. lineari-lanceolatis integerrimis, capitulis paniculatis hemisphæricis, invol. squamis subulatis. (†) Patr. et cæ. car. ign. †.

102. *C. PROCUMBENS* (Balb. cat. h. Taur. p. 28), caule procumbente ramoso, foliis spatulatis subtus tomentoso-niveis margine revolutis, pedunculis nudis monocephalis. — Patr. ign. *Conyza chrysocomoides* Ten. non Willd. ex Balbo. *Conyza prostrata* Ten. cat. h. Neap. 1819. p. 75?

103. *C. ? TORTUOSA* (Linn. hort. Cliff. 405), caule fruticoso tortuoso, foliis ovato-oblongis integerrimis, racemis reflexis. ♀ in Verâ-Cruce (Linn.), Brasiliâ (Spreng.), Madagascar (Vaill. act. par. 1719. p. 396). Species mihi omnino obscura! An fortè *Vernonia*? †.

†† *Conyze exclusæ.*

*C. ABBREVIATA* Wall. = *Vernonia abbreviata*.

*C. ACUMINATA* Wall. = *Vernonia acuminata*.

*C. ADNATA* H. B. K. = *Pluchea subdecurrens*.

*C. ALATA* Roxb. = *Blumea alata*.

*C. ALATA* Bening. = *Inula bifrons*.

*C. ALATA* var.  $\beta$ . Wall. = *Blumea pterodonta*.

*C. AFRICANA* Wall. = *Vernonia albicans*.

*C. ALOPECUROIDES* Lam. = *Pterocanlon alopecuroides*.

- C. AMPLECTENS Wall. = *Blumea amplexans*.
- C. AMPLEXICAULIS Lam. = *Blumea amplexans*.
- C. AMPLEXICAULIS Wall. = *Blumea oligocephala*.
- C. AMPLEXICAULIS Michx. = *Pluchea bifrons*.
- C. ANAGALLIDIFOLIA Wall. = *Blumea anagallidifolia*.
- C. ANGUSTIFOLIA hort. Par. = *Baccharis Pingraea*.
- C. ANGUSTIFOLIA Lam. = *Monarctus salicifolia* β.
- C. ANTHELMINTHICA L. = *Vernonia anthelmintica*.
- C. APPENDICULATA Blum. = *Pluchea appendiculata*.
- C. APPENDICULATA Lam. = *Senecio appendiculatus*.
- C. ARBOREA Wall. = *Vernonia arborea*.
- C. ARBORESCENS L. = *Vernonia arborescens*.
- C. ARBUTIFOLIA Lam. = *Baccharis arbutifolia*.
- C. ARGENTEA Wall. = *Inula argentea*.
- C. AROMATICA Wall. = *Blumea aromatica*.
- C. ARTICULATA Lam. = *Baccharis articulata*.
- C. ASPERA Wall. = *Decaenium asperum*.
- C. ASTEROIDES Desf. = *Pluchea subdecurrens*.
- C. ASTEROIDES Lin. = *Sericocarpus conyzoides*.
- C. ASTEROIDES Wall. = *Chrysopsis gossypina*.
- C. ATTENUATA Wall. = *Vernonia attenuata*.
- C. AURICULATA Moench. = *Phagnalon rupestre*?
- C. AURITA L. F. = *Blumea aurita*.
- C. AXILLARIS Lam. = *Blumea axillaris*.
- C. BALSAMICA Wall. = *Psiadia balsamica*.
- C. BALSAMIFERA L. = *Pluchea balsamifera*.
- C. BARBATA Wall. = *Blumea barbata*.
- C. BELLIDIFOLIA Wall. = *Vernonia bellidifolia*.
- C. BIFOLIATA Linn. = *Blumea bifoliata*.
- C. BIFOLIATA Cham. = *Blumea Chamissoniana*.
- C. BIFOLIATA Wall. = *Blumea Wightiana* β.
- C. BIFOLIATA Walt. = *Sericocarpus tortifolius*.
- C. BIFRONS Linn. = *Pluchea bifrons*.
- C. BLANDA Wall. = *Vernonia blanda*.
- C. BRACTEOLATA Wall. = *Vernonia bracteolata*.
- C. BRYOIDES Lam. = *Baccharis bryoides*.
- C. BUXIFOLIA Lam. = *Baccharis buxifolia*.
- C. CALYCINA Cav. = *Phagnalon calycinum*.
- C. CAMPHORATA Ell. = *Pluchea camphorata*.
- C. CANDIDA Linn. = *Inula candida* α.
- C. CANDIDA Willd. = *Inula candida* β.
- C. CANESCENS Bory. = *Schizogyne sericea*.
- C. CANESCENS Linn. f. = *Webbia pinifolia*.
- C. CAROLINENSIS Jacq. = *Pluchea odorata*.
- C. CERNUA Wall. = *Blumea alata* β.
- C. CENOPODIFOLIA Lam. = *Gynura Mauritanica*.
- C. CHINENSIS Linn. = *Blumea Chinensis*.

- C. *CHINENSIS* Lam. = *Cyanopsis pubescens*.  
 C. *CHRYSOCCOMOIDES* Desf. atl. = *Nolletia chrysoccomoides*.  
 C. *CHRYSOCCOMOIDES* Desf. arb. = *Felicia angustifolia*.  
 C. *CINEREA* Linn. = *Vernonia cinerea*.  
 C. *CINEREA* var. 2 Wall. = *Vernonia physalifolia*.  
 C. *CINERARIOIDES* Bojer. = *Rochonia cinerarioides*.  
 C. *CINERASCENS* Wall. = *Vernonia cinerea* s.  
 C. *CLETHRIFOLIA* Willd. herb. = *Vernonia odoratissima*.  
 C. *COMMUNIS* Wall. = *Blumea hieracifolia*.  
 C. *CORONOPUS* Lam. = *Sarcanthemum Coronopus*.  
 C. *CORYMBOSA* Roxb. = *Pluchea Indica*.  
 C. *CORTESI* H. B. K. = *Pluchea Cortesi*.  
 C. *CUNEATA* Bojer. = *Rochonia cuneata*.  
 C. *CUNEIFOLIA* Lam. = *Baccharis cuneifolia*.  
 C. *CUNEIFOLIA* Wall. = *Blumea cuneifolia*.  
 C. *CUNEIFOLIA* Rensch. = *Solidago spuria*.  
 C. *CUPRESSIFORMIS* Lam. = *Lepidophyllum cupressiforme*.  
 C. *DAMPIERI* Cunn. = *Eurybia Dampieri*.  
 C. *DENSIFLORA* Wall. = *Plumea densiflora*.  
 C. *DICHOTOMA* Linn. = *Jasonia dichotoma*.  
 C. *DIOSCORIDIS* Rauh. = *Bluchea Dioscoridis*.  
 C. *DISCOLOR* Bory. = *Senecio appendiculatus*.  
 C. *DIVARICATA* Spr. = *Vernonia diffusa*.  
 C. *DIVERGENS* Wall. = *Vernonia subsessilis*.  
 C. *ELEGANTULA* Wall. = *Vernonia bellidifolia*.  
 C. *ELÆAGNIFOLIA* Wall. = *Vernonia elæagnifolia*.  
 C. *ELONGATA* Wall. = *Blumea elongata*.  
 C. *ERICOIDES* Lam. = *Vernonia ericoides*.  
 C. *EXCISA* Wall. = *Blumea excisa*.  
 C. *EXTENSA* Wall. = *Vernonia extensa*.  
 C. *FASCICULATA* Wall. n. 3017. = *Blumea flavida*.  
 C. *FASCICULATA* Wall. n. 3094. = *Blumea fasciculata*.  
 C. *FÆTIDA* Willd. = *Erigeron fœtidum*.  
 C. *FÆTIDA* Lam. = *Pluchea Marylandica*.  
 C. *FOLIOLOSA* Wall. n. 3030. = *Pluchea Martabanica*.  
 C. *FOLIOLOSA* Wall. n. 3281. = *Pulicaria foliolosa*.  
 C. *FRUTICOSA* Linn. = *Vernonia fruticosa*.  
 C. *GEMINIFLORA* Ten. = *Phagnalon rupestre*.  
 C. *GENISTELLOIDES* Lam. = *Baccharis genistelloides*.  
 C. *GLOMERATA* Wall. = *Blumea glomerata*.  
 C. *GLOMERATA* C. Wall. = *Blumea leptoclada*.  
 C. *GLUTINOSA* Lam. = *Psadia glutinosa*.  
 C. *GNAPHALONES* H. B. K. = *Lænnecia gnaphalodes*.  
 C. *GRACILIS* Wall. = *Blumea gracilis*.  
 C. *GRANDIS* Wall. = *Pluchea grandis*.  
 C. *GRAVEOLENS* Steud. = *Inula graveolens*.  
 C. *GRAVEOLENS* Wall. = *Erigeron hispidum*.

- C. GUINEENSIS Willd. = *Blumea Guineensis*.
- C. GUMMIFERA Roxb. = *Commidendron gummiferum*.
- C. HALIMIFOLIA Desf. = *Baccharis halimifolia*.
- C. HAMILTONI Wall. = *Blumea Hamiltoni*.
- C. HELIOTROPIFOLIA Lam. = *Monartheus rufescens*.
- C. HEYNEANA Wall. = *Blumea Heyneana*.
- C. HIERACIFOLIA Spreng. = *Blumea hieracifolia*.
- C. HIRSUTA Linn. f. = *Pluchea hirsuta*.
- C. HIRSUTA Comm. = *Cylindrocline Commersonii*.
- C. HIRSUTA Wall. = *Erigeron sublyratum*.
- C. HOLOSERICA Wall. = *Blumea holosericea*.
- C. HYMENOPHYLLA Wall. = *Blumea hymenophylla*.
- C. JAVANICA Blum. = *Blumea lacera* f.
- C. INCANA Heyn. = *Vernonia physalifolia*.
- C. INCANA Wall. = *Vernonia abbreviata* var. ?
- C. INDICA Blum. = *Pluchea indica*.
- C. INDICA Wall. = *Decaneurum Dendigulense*.
- C. INTERMEDIA Lag. = *Phagnalon rupestre* β.
- C. INULOIDES Ait. = *Jasonia dichotoma*.
- C. LACERA Burm. = *Blumea lacera*.
- C. LACINIATA Roxb. = *Blumea laciniata*.
- C. LACINIATA Ham. = *Blumea crepidifolia*.
- C. LACTUCIFOLIA Wall. = *Blumea lactucifolia*.
- C. LANCEOLATA Wall. n. 3059. = *Vernonia divergens*.
- C. LANCEOLATA Wall. n. 2991. = *Berthelotia lanceolata* β.
- C. LIMONIFOLIA Sibth. = *Inula candida* β.
- C. LINEARIFOLIA Lam. = *Psadia linearifolia*.
- C. LINIFOLIA Linn. = *Sericocarpus solidaginoides*.
- C. LINIFOLIA Wall. = *Vernonia leptophylla*.
- C. LOBATA Linn. = *Neurolana lobata*.
- C. LOBATA Juss. = *Ditrichum macrophyllum*.
- C. LONGICAULIS Wall. = *Eupatorium longicaule*.
- C. LONGIFOLIA Heyn. = *Blumea longifolia*.
- C. LUCIDA Spreng. = *Vernonia splendens*.
- C. LYCOPODIOIDES Lam. = *Eriothrix lycopodioides*.
- C. MACROPHYLLA Wall. = *Blumea macrophylla*.
- C. MACROSTACHYA Wall. = *Blumea macrostachya*.
- C. MADAGASCARIENSIS Lam. = *Psadia Madagascariensis*.
- C. MAGELLANICA Lam. = *Baccharis Magellanica*.
- C. MANILLENSIS Less. = *Blumea Manillensis*.
- C. MARYLANDICA Michx. = *Pluchea Marylandica*.
- C. MARYLANDICA Pursh. = *Pluchea foetida*.
- C. MEMBRANACEA Wall. = *Blumea membranacea*.
- C. MOLLIS Willd. = *Vernonia cinerea*.
- C. MULTIFLORA Wall. = *Vernonia multiflora*.
- C. MURALIS Ham. = *Blumea muralis*.

- C. VIRGATA Linn. = *Pterocaulon virgatum*.  
 C. VISCOSA Hortul. = *Inula viscosa*.  
 C. VISCOSULA Wall. = *Blumea viscosula*.  
 C. VOLKAMERIEFOLIA Wall. = *Vernonia volkameriæfolia*.  
 C. VOLUMINIS Wall. = *Microglossum proliferum*.  
 C. VULGARIS Lam. = *Inula conyza*.  
 C. WIGHTIANA Wall. n. 3028. = *Monosis Wightiana*.  
 C. WIGHTIANA Wall. n. 3093. = *Blumea Wightiana*.

CCXIII. PHAGNALON Cass. bull. philom. 1819. p. 174. dict. 19. p. 118. 13. p. 561. et 39. p. 400. Less. syn. 204. — *Conyza* et *Gnaphalii* sp. *emct.*

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, marginalibus pluriserialibus tenuissimis filiformibus foemineis sæpè abortientibus, centralibus 5-dentatis hermaphroditis fertilibus. Recept. nudum planum. Involucrum imbricatum, squamis plus minus scariosis. Achænium teretiusculum erostre adpressè villosum. Pappus 1-serialis paucisetus conformis piloso-setaceus subscaber. — Fruticuli regiones Mediterraneas aut Canarienses habitantes, ramis tomentosis teretibus. Folia sparsa linearia aut lanceolata integra aut subdentata subtus tomentosa sæpius margine revoluta. Capitula ramos sæpius apice aphyllis terminantia ebracteata sæpè solitaria. Flores lutei aut lutescentes.

1. P. SORDIDUM, caule basi fruticoso, ramis tomentosis ultimis aphyllis apice capitula 1-3 sessilia gerentibus, foliis linearibus integerrimis margine revolutis utrinquè tomentosis, invol. ovato-cylindracei squamis ovatis scariosis obtusiusculis. 5 ad rupes et muros Gallie mediterraneæ, Italiae, Mauritanie, Hispanie. *Conyza sordida* Linn. mant. 468. DC. fl. fr. 4. p. 140. *Gnaphalium sordidum* Linn. sp. 1193. *Gnaphalium conyzoides* Lam. fl. fr. 2. p. 63. *Phagnalon tricephalon* Cass. dict. 39. p. 401. — Barr. ic. t. 277 et 368. Invol. rufescens glaberrimum. Cor. flavidæ. (v. v. et s.)
2. P. UMBELLIFORME, caule basi fruticoso, ramis tomentosis apice ramosos phres 1-cephalos subumbellatos gracillimos interdum umbellato-ramulosos gerentibus, foliis integerrimis radicalibus lineari-oblongis planis caulinis linearibus margine revolutis subtomentosis, invol. oblongi squamis angustis acuminatis apice rufo-scariosis. 5 in rupestribus insulæ Palmæ arch. Canar. legerunt d Webb et Berthelot. Affine P. sordido sed evidenter distinctum! (v. s. in b Webb et Berth.)
3. P. SAXATILE (Cass. bull. philom. 1819. p. 174), caule basi suffruticoso ramosissimo, ramis tomentosis apice productis in pedunculos elongatos aphyllis 1-cephalos, foliis infer. oblongo-lanceolatis subdentatis super. linearibus margine revolutis suprà araneosis subtus tomentosis, invol. campanulati glabri squamis sublinearibus acuminatis apice scariosis. 5 in rupibus et muris regionis Mediterraneæ à Cretâ ad insul. Canarienses, à Græciâ ad Lusitaniam. *Conyza saxatilis* Linn. sp. 1206. DC. fl. fr. 4. p. 140. Phagn. subdentatum Cass. dict. 39. p. 400. — Barr. ic. t. 426. Variat foliis caulinis planis aut marginè revolutis, subdentatis aut integerrimis plus minus tomentosis. (v. v. et s.)  
     β. *intermedium*, foliis subadnato-amplexicaulibus, invol. squamis angustioribus minus acutis sed non obtusis. 5 in ins. Scio (Oliv.!), in muris in Cretæ (Sieb.!), in Catalonia (Pourr.!) et in Aragoniâ (Aso!). *Conyza saxatilis* Sieb.! Aso! *Conyza intermedia* Lag. clench. p. 28. n. 33 *Phagnalon* Lagascæ Cass. dict. 39. p. 401. Au species propria? (v. v.)
4. P. RUPESTRE, caule basi suffruticoso procumbente, ramis tomentosis, pediculis elongatis solitariis geminisve monocephalis aphyllis floccosis glabris foliis basi adnato-amplexicaulibus oblongo-linearibus integris subdentatis

pè undulatis suprâ araneosis subtùs tomentosis, invol. subcampanulati glabris squamis rigidis obtusis planiusculis.  $\zeta$  in rupibus calcareis Syriæ, Mauri-  
niæ, Arabiæ, Ægypti, Siciliæ, Capræ, littoris Neapolitani, Sardinie,  
ad Olibiensium, Hispaniæ ad Malagam et in ins. Canariensium. *Conyza*  
*nentosa* Forst. æg. p. 148. *Conyza rupestris* Linn. mant. 113. ex syn. *Co-*  
*za tomentosa* Shaw. sp. n. 176. ic. pess. ? *Conyza saxatilis* Ten. fl. neap. 2.  
213. Bové ! pl. exs. n. 437 non Willd. *Conyza geminiflora* Ten. cat. h. Neap.  
19. p. 75. syl. p. 427. Link enum. h. Berol. 2. p. 323. *Conyza Tenorii*  
reng. ex Ten. Phagnalon Tenorii Presl fl. sic p. xxx. Phagnalon spathu-  
um Cass. dict. 39. p. 401 ? Species à plurimis nunc cum *Ph. saxatili* nunc  
n° *Ph. calycino* mixta sed distinctissima foliis basi subdecurrentes amplexi-  
libus subhastatis, caule basi decumbente, invol. squamis nec denticulatis,  
undulatis nec membranaceo-scariosis. (v. s.)

**CALYCINUM**, caule basi suffruticoso, ramis tomentosis, pedunculis elonga-  
subtomentosis aphyllis 1-cephalis, foliis sessilibus oblongo-linearibus inte-  
s serratis, super. margine revolutis, subtùs plus minus tomentosis, invol.  
spanulati glabri squamis obtusis subserratis membranaceo-scariosis subun-  
natis.  $\zeta$  in rupibus insulæ Maderæ legit cl. Lowe et circa Mogador cl. Brous-  
set. *Conyza calycina* Cav. annal. scienc. nat. v. 4. (1801) p. 87, ex Mogador  
a. Phagnalon Bennetti Low. ! in litt. *Conyza rupestris* Buch in hist. Mad.  
a.)

**MITIDUM** (Fresen. pl. eg. p. 81. t. 4. f. 2), caule vix basi suffrutescente  
etiussculo ramoso tomentoso, foliis linearibus basi subauritis semiamplexi-  
libus margine revolutis acutis suprâ araneosis subtùs cano-tomentosis,  
ol. lati campanulati squamis scariosis glabris nervo medio fusco donatis, ext.  
tis obtusis intimis linearibus acutis.  $\zeta$  in valle Arbain (Rupp. ex Fres.),  
Monte Sinai (Bové !). Arabis dicitur *Honer-el-Naghi* *Conyza* n. 120 Bové !  
exs. Flor. junio. Species ex invol. distinctissima. Pappus 4-5-setus albissi-  
mus. Capitula majores quàm pler. sp. generis. (v. s. comm. à cl. Bové.)

**PUMILUM**, caule ad collum subterroso fruticoso ceterum decumbente  
viscido, foliis obovatis in petiolum attenuatis integerrimis subtùs tomen-  
to, pedunculo vix foliis longiore aphylo 1-cephalo subtomentoso, invol.  
scierialis squamis acuminatis exterioribus dorso tomentosis intimis elongatis  
scariosis.  $\zeta$  in cacumine montium Sphacioticorum Cretæ. *Conyza pumila*  
th. et Sm. prod. fl. Gr. p. 173. fl. Græc. (ined.) t. 863. *Conyza pygmæa*  
! pl. cret. exs. Suffrutex certè omnium minimus, vix pollicem longus.  
a.)

## CCXIV. CHIONOLÆNA.

**PITULUM** multiflorum heterogamum, fl. omnibus tubulosis, radii  
simis 2-3-dentatis vel truncatis fœmineis pluriserialibus, disci  
5 bisexualibus abortu sterilibus 5-dentatis. Invol. squamæ pluri-  
es lineares siccæ albæ glabræ demùm deciduæ. Recept. nudum  
tatum persistens. Anthera ecaudata. Stylus radii exsertus bifidus  
s, disci indivisus filiformis minutè hispidulus. Achænium gracile  
usculum villosum. Pappus 1-serialis setosus. — Suffrutex Brasi-  
s arborescens minutissimam referens, trunco nudo, ramis foliosis  
sericeo-canis mox glabris. Folia alterna conferta patula linearia  
suprà glabra subtùs sericeo-cana enervia. Capitula ramos termi-  
a sessilia solitaria. Invol. ferè Helichrysi. Flores circ. 35 flavi ? —  
tract. accedit ad Conyzam et Phagnalon sed habitus valdè pecu-

**ARBUSCULA**.  $\zeta$  in Brasiliæ provinciâ Minarum-Generalium ad Mariannam  
t cl. Vanthier (pl. exs. n. 298 !). Frutex vix pedalis dendromorphus. Folia  
acerosa, 5-6 lin. longa semilineam lata. (v. s.)



## CCXV. FLACHOTHAMNOS.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus filiformibus fœmineis (fortè sterilibus), disci paucioribus tubulosis 5-dentatis hermaphroditis (fortè abortu masculis). Invol. imbricatum, squamis linearibus acutis 3-4-seriatis. Recept. nudum. Anth..... Styl..... Achænia teretiuscula. Pappus radii pluriserialis setiformis, setis inter se conformibus; disci biserialis, squamellis exter. paleaceis acutis, inter. paucis setiformibus. — Frutex Novo-Hollandicus glaber ramosus erectus. Folia sparsa teretiuscula acutissima carnosæ erecta integerrima. Rami apice squamellati 1-cephali. — Ex capitulis junioribus characterem florum non ritè eruere quivi, sed pappo heterogeneo genus satis distinctum videtur. Nomen ex *ελαχες*, parvus, et *δαιμος*, frutex.

1. E. CUNNINGHAMI. ♂ in humidis depressis ad flumen Lachlan Nov.-Hollandiæ flor. legit cl. All. Cunningham. Senecio othoanoides Cann. in litt. 1835  
Frutex bipedalis Mesembryanthemi facie. Caulis albicans. Folia 7-10 lin. longæ seminilem lata. (v. s. comm. à cl. inv.)

DIV. IV. BACCHARIDÆ. — Baccharidæ veræ *Cass. tabl. op. phyt.* 3. p. 68. — Baccharidæ *Less. syn.* 204 non p. 200.

Capitula (ut in *Eutarchonantho*) dioica.

CCXVI. POLYPAPPUS *Less. in linnæa* 1819. p. 314. 1831. p. 149. *syn.* p. 204

Capitula homogama dioica. Cor. masculæ et fœminæ tubulosæ. ♂ 5-dentatæ, ♀ filiformes. Rachis plana ebracteolata. Invol. campanulatum imbricatum. Achænia teretiuscula. Pappus cum maribus tum fœmineis pluriserialis, setis plurimis apice clavellatis. — Fruticuli Americani ramosissimi. Ramuli pubescentes adulti glabri. Folia alterna sessilia oblonga subcuneata triplinervia. Capitula ramos terminantia solitaria in subdiv. majuscula. Cor. ex sicco purpurascens.

1. P. TRIPLINERVIS (*Less. syn.* 204), foliis apice serratis 3-5 notatis, achæniisque glabris. ♂ in Brasiliâ legit cl. Sellow. *Vernonia triplinervis Spreng. syst.* 3. p. 435. Folia pollicem longa, 3 lin. lata. Capitulum 6-7 lin. diam. Invol. squamæ lanceolatæ acutæ adpressæ minutè ciliolatæ. Pappus rubescens (v. s. ♂ comm. à Mus. reg. Berol.)  
2. P. ? PARVIFOLIUS, foliis linear-oblongis integerrimis junioribus subtus achæniisque puberis. ♂ in Mexico ex h. Hænke. Frustula muscula imperfecta tantum video, sed habitus omninò prioris. Fol. sup. 4 lin. longa, 1 lin. lata. Capitula 4-5 lin. longa. Cor. fœm. paulò crassiores et evidentius dentatæ. Indiv. ♂ ign. (v. s. ♀ in h. Hænke.)

CCXVII. BACCHARIS *Linn. gen. n.* 949. *excl. spec. Michx. fl. bor. Am.* 1. p. 125. *Brown obs. comp. in trans. Linn. soc.* 12. p. 115. — *Molina Ruiz et Pav. prod.* p. 111. t. 24. *syst.* p. 200. — Baccharis et Molina *Less. syn.* p. 204 et 205 — *Sergilus* et *Baccharis Gært.* — *Baccharis*, *Sergilus*, *Pingrea*, *Ternstroemia* et *Arrbenachne Cass.* — *Stephananthus Lehm.*

Capitula multiflora dioica, cor. homogamis tubulosis. Recept. nudum aut in paucis speciebus subpaleaceum. Invol. subhemisphericum aut oblongum pluriseriale imbricatum. ♂ Cor. ad faucem dilatatur 5-fida. Anthæræ exsertæ ecaudatæ. Stylus plus minus abortivus. ♀ Cor. filiformes subtruncatæ, stylus bifidus exsertus. Anth. omninò nullæ. Achæniæ sæpius sulcato-costatum. Pappus pilosus ♂ uniserialis &

pius tortuosus aut subplumoso-penicillatus involucri longitudinem subæquans, ♀ uni- aut pluriserialis apice gracilis invol. longitudinem sæpius superans. — Frutices rariùs herbæ. Species genuinæ omnes in America spontaneæ sæpius glabræ et resinoso-viscosæ interdum variè villosæ. Folia alterna in paucissimis opposita, sæpius ex utroque latere in nervulum aut interdum in alam decurrentia integra aut dentata. Capitula variè disposita. Cor. sæpius albæ. — Species inter se valde affines nec ullo modo in genera plura divellendæ.

§. 1. *Trinervatæ, nempe foliis tri- aut triplinerviis non imbricatis, nec cuneatis, ramis apteris.*

1. *B. SPICIOSA*, suffruticosa glabra, ramis teretibus, foliis in petiolum attenuatis ellipticis acuminatis integerrimis trinerviis et reticulato-venosis, corymbo terminali amplo fastigiato composito polycephalo, invol. utriusque sexus campanulati squamis ovatis submucronatis, achæniis adpressè phloisculis § in montania Guadalupe jnnio flor. legit cl. Perrottet. Frutex 6-8-pedalis, ramis crassis medullosis semilignosis. Folia 3-4 poll. longa, 15-18 lin. lata. Corymbi rami angulati. Pappus rufus, ♀ nniserialis. (v. s. comm. à cl. Perrottet.)
2. *B. NERVOSA*, fruticosa glabra, ramis virgatis teretibus, foliis petiolatis ovalilanceolatis acuminatissimis trinerviis integerrimis membranaceis, corymbis inf. axillaribus super. aphyllis in paniculam terminalem dispositis, invol. ♂ campanulati squamis lanceolatis acutis apice ciliatis § in insul. Trinitatis (Sieb.), et Guadalupe (Krauss!). Eupatorium nervosum Sieb. ! fl. trin. exs. 76. non Swartz. Folia 3 poll. longa, vix pollicem lata. Flor. scem. ign. (v. s.)
3. *B. TRINERVIS* (Pers. ench. 2. p. 423. n. 7), suffruticulosa glabra ramosissima, ramis teretibus, ramulis striato-sulcatis, foliis petiolatis ovato-lanceolatis acutis set acuminatis integerrimis 3-nerviis, caulinis alternis, rameis oppositis, corymbis terminalibus laxis, invol. subcampanulati utriusque sexus squamis ovatis obtusiusculis, achæniis adpressè puberis. § in Brasiliâ (Comm. ! Lhostky!), prope Rio-Janeiro (Lund!). Conyza trinervis Lam. dict. 2. p. 85. Pappus rufus. Folia 1-2 poll. longa, 6-12 lin. lata. (v. s.)
4. *B. QUINQUENRVIS*, suffruticosa glabra, ramis teretibus, foliis alternis petiolatis ovatis acutis quinquenerviis à medio ad apicem crenato-serrulatis, capitulis ♂ ad apices ramulorum 5-7-congestis, invol. squamis lineari-oblongis obtusis disco æqualibus. § in fruticetis prov. Sancti-Pauli Brasiliæ legit cl. Lund. A trinervi et ab omnibus mihi notis foliis quinque seu ferè quintuplinerviis diversa. (v. s. ♂ comm. à cl. Lund.)
5. *B. ANXIORDES* (H. B. et Kunth nov. gen. am. 4. p. 66), fruticosa volubilis, ramis sulcatis pubescentibus, foliis petiolatis elliptico-oblongis brevissimè acuminatis basi acutis integerrimis trinerviis subtilis secus nervos et in petiolo pubescentibus, corymbis axillaribus et terminalibus in paniculam dispositis, invol. campanulati squamis oblongis in ♂ subobtusis in ♀ subacutis, achæniis striatis puberulis. § in Andibus Peruvianorum (H. B. K.), ad Caracas (Vargas!), in Brasiliâ (h. Deless.), ad fluvior. margines circa Rio-Janeiro (Lund ! Gand.) et ni fallor in prov. Matto-Grosso. Affinis *B. trinervi* et cineræ. Frutex ex Kunth volubilis, ex Lund 2-5-pedalis. An plures hic species confusæ ? (v. s.)
6. *B. CAMFORUM*, collo fruticoso, caulibus erectis suffrutescentibus simplicissimis teretibus vix sub lente pubescentibus, foliis sessilibus elliptico-oblongis brevissimè acuminatis trinerviis à medio ad apicem argutè serratis parè reticulatis, capitulis ♀ in corymbum congestum terminalem congestis oblongis, invol. squamis oblongo-linearibus obtusis, interioribus submembranaceis disco æqualibus. § in campis editis prov. Sancti-Pauli Brasiliæ legit cl. Lund. Chrysocoma decussata fl. flum. 8. t. 9? quæ foliis alternis gaudet. Habitus hujus stirpis et plurimarum in campis Brasiliæ obviis præcipuè ex cl. Lund tribuendus more

vulgarissimo igne destruendi herbas camporum, undè rhizoma induratur et quotannis ampliatur (v. s. *Q* comm. à cl. Lund.)

7. *B. CINEREA*, suffruticulosa? totà pube breviter velutino-cinerascente, caule tereti, ramis divergentibus striatis, foliis petiolatis ovato-lanceolatis acuminatis integerrimis trinerviis suprâ glabriusculis alternis ramis hinc inde suboppositis, corymbo terminali laxo, invol. ovati utriusque sexûs squamis lanceolatis acutis, acheniis adpressis puberis. *♂*? in Brasiliâ circa Bahiam leger. cl. Seemann et Blanchet. Affinis *B. trinervi* sed canescenti-cinerea nec glabra, invol. squamæ acutæ nec obtusæ, etc. (v. s.)
8. *B. TRICHOCLADA*, fruticosa, ramis pediculisque teretibus hirsutissimis, foliis vix petiolatis ovato-ellipticis subacuminatis mucrone callosio apiculatis integerrimis grossè trinerviis suprâ glabris subtilis sparsè villosis, pedunculis secus ramulos axillaribus capitula 3-5 subsessilia gerentibus, invol. *♂* squamis lanceolatis acuminatis subciliatis glabriusculis. *♂* in Mexico ex herb. Hænk. Affinis *B. cinerea* et rhexioidi sed ramis saltem junioribus villis longis densè vestitis faciliè differt. Flor. *♀* ign., *♂* non satis not. (v. s. in h. Hænk. à cl. de Sternberg miss.)
9. *B. CINNAMOMIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 65), fruticosa glabra, ramulis angulatis, foliis petiolatis oblongis breviter acuminatis basi angustatis integerrimis subcoriaceis trinerviis suprâ vitidis, paniculis terminalibus ramosissimis, capitulis *♀* pedicellatis, invol. hemisphærici squamis scariosis acutis *♂* in convallibus Quitensium? Folia 4 poll. longa, 18 lin. lata, petiolo 7-8 lin. Flores in invol. crebri, masculi ign. Affinis *B. trinervi* sed folia basi angustata nec rotundata.
10. *B. VENUSTA* (H. B. et Kunth nov. gen. am. 4. p. 66), fruticosa glabra, ramulis angulatis, foliis petiolatis oblongis breviter acuminatis basi acutis integerrimis membranaceis triplinerviis, paniculis axillaribus terminalibusque ramosis, capitulis *♀* pedicellatis, invol. hemisphærici squamis scariosis acutiusculis, achenio sulcato glabro. *♂* in Novâ Andalusâ juxta Caripe et in rivis Orinocensibus. Folia 3 poll. longa, pollicem lata, petiolo 3-4 lin. Flor. *♂* ign. Affinis *B. trinervi*.
11. *B. OXYPHYLLA*, suffruticulosa? glabra resinoso-subviscosa, ramis virgatis sulcato-angulatis, foliis petiolatis elongato-lanceolatis basi acutis apice acuminatissimis integerrimis triplinerviis, paniculâ ramosâ laxâ terminali, capitulis *♀* pedicellatis, invol. squamis lanceolatis acutis intim. linearibus, achenio subpuberulo. *♂*? in Peruvâ legit cl. Poeppig (pl. exs. n. 24). Petioli 4 linear. Limbus fol. 3-3  $\frac{1}{2}$  poll. longus, 7 lin. latus. Nervi laterales duo suboppositi ad quartam nervi medii partem orti, cæterum nulli. Flor. *♂* ign. (v. s.)
12. *B. DIVERGENS*, fruticosa apice viscosa, ramis teretibus floridis divergentibus, foliis brevissimè petiolatis lanceolatis utrinquè acutis subtriplinerviis integerrimis glabris summis pedicellisque subpubescentibus, corymbis compositis confertis, capitulis subfasciculatis, invol. *♀* squamis lanceolatis acutiusculis apice summo subbarbatis. *♂* in Mexico ex h. Hænk. Folia pollicem long. 4-5 lin. lata suprâ viscosa-nitida. Achenia parva glabriuscula. Pappus 1-2 lin. pallidè rufescens (v. s. *♀* in h. Hænk. à cl. de Sternberg comm.)
13. *B. DOUGLASSII*, herbacea erecta glaberrima viscosa, caule simplici apice angulato, foliis breviter petiolatis lanceolatis acutis subtriplinerviis integerrimis utrinquè punctatis, corymbo *♂* composito fastigiato polyccephalo subobuso, invol. campanulati squamis lanceolatis margine membranaceis acutiusculis ciliatis. *♂* *♂* in Californiâ legit cl. Douglas. Folia 2 poll. longa 5-6 lin. lat. super. angustiora. Pappus *♂* invol. æqualis albidus. (v. s. *♂* comm. à hort. Lond.)
14. *B. VIMINEA*, fruticosa glabra, ramis teretibus, ramulis striato-nervosis, foliis breviter petiolatis oblongo-lanceolatis utrinquè acutis basi tenuiter trinerviis, aliis integerrimis, aliis ad apicem minutè denticulatis, corymbis racemosis confertis ramulos terminantibus, capitulis *♂* pedicellatis, invol.

aquamis ovali-lanceolatis acutis membranaceis ciliatis.  $\beta$  in Californiâ legit cl. Douglas. Folia 12-14 lin. longa, 2-2  $\frac{1}{2}$  lata, primâ fronte uninervis imò ferè enervis, ramul. infer. breviora ferè squamis gemmaceis refractis. Pappus  $\sigma^7$  albidus crispus vix invol. longior. Fl.  $\sigma$  ign. (v. s. cumm. ab hunc. soc. hort. Lond.)

15. *B. HÄNKEI*, herbacea? erecta ramosa glabra viscosa, ramis etiam floridâ teretibus, foliis breviter petiolatis ovali- seu oblongo-lanceolatis acutis integerrimis trinerviis utrinquè valdè punctatis, corymbo composito fastigiato polycephalo, invol. campanulati squamis lanceolatis margine membranaceis acutiusculis ciliatis, acheniis glabris minimis.  $\gamma$   $\beta$  in Mexico circa Real del monte legit cl. Hænke. Valdè affinis *B. Douglasii* sed caulis ramosior apice teretiusculus, folia latiora magis punctata pappus rufus (v. s.  $\sigma^7$  et  $\sigma$  in herb. Hænke à cl. de Sternberg misso.)

16. *B. PUNHAUZA*, caule herbaceo? erecto tereti simpliciter pubero, foliis oblongo-linearibus acutis sensibus basi trinerviis integerrimis aut remotè subdentatis pube sparâ subacabridis, paniculâ  $\sigma$  elongatâ angustâ subracemosa, ramulis axillaribus distantibus oligocephalis, pedicellis 1-3-cephalis, invol. squamis oblongo-lanceolatis subacutis apice membranaceis, acheniis pube minimâ pubescentibus.  $\gamma$ ? in Brasiliâ prov. Sancti-Pauli. Caulis bipedalis. Pappus trifidus invol. duplò longior pluriserialis. Achen. et invol. ferè *B. cordifolia* sed cat. diversa. Fl.  $\sigma^7$  ign. (v. s.  $\sigma$  in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 515 misso.)

17. *B. LONGIPES* (Kunze! in Poepp. coll. 2. n. 104), suffruticosa glabra resinoso-viscosa, foliis lineari-lanceolatis utrinquè attenuatis mucronato-acutis integerrimis aut parè serratis punctatis trinerviis, nervis lateralibus margini adproximatis, corymbo composito exserto fastigiato polycephalo aphylo, capitulis  $\sigma$  campanulatis, invol. squamis lanceolatis acuminate glabris margine scariosis, acheniis glabris nervoso-striatis.  $\beta$  in lapidosâ inundatâ Mîo de Chili legit cl. Poeppig. Proximè accedit ad *B. glutinosam* et præsertim ad *B. parvifloram* et si hybridæ facillè admittendæ essent, ut utriusque proles mixta ferè videtur (v. s. ex coll. Poepp.)

18. *B. SALICIFOLIA* (Pers. ench. 2. p. 425. n. 35), fruticosa, foliis lanceolato-linearibus triplinerviis serratis, serraturis remotis, corymbis terminalibus.  $\beta$  in Peruvîæ montibus altis ad Pillao et Acomayo tractus. Molina salicifolia *B.* et Pav. syst. p. 210. An, suad. Spreng., eadem ac *B. gnidiifolia* ? sed capet, non suffic. †.

19. *B. ZARZIMORNA* (H. B. et Kunth nov. gen. am. 4. p. 63), fruticosa ramosissima glabra, ramulis sulcatis viscosis, foliis petiolatis lineari-lanceolatis acutis basi angustatis dentatis trinerviis membranaceis, paniculis terminalibus, capitulis  $\sigma^7$  pedicellatis 20-floris, invol. hemisphærici squamis scariosis acutis.  $\beta$  in calidis prov. Bracamorensis juxta Passo de Matara. Fol. 2  $\frac{1}{2}$  poll. longa, 4 lin. lata, petiolo 3 lin. Flor.  $\sigma$  ign.

20. *B. GNIDIIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 61), fruticosa glabra, ramulis sulcatis, foliis subsessilibus lineari-lanceolatis subacuminatis apicem versus argutè serratis trinerviis subcoriaceis subhis glanduloso-punctatis, corymbis  $\sigma^7$  terminalibus, pedicellis bracteis subulatis gerentibus, invol. hemisphærici squamis lanceolatis acutis scariosis.  $\beta$  in scopulis montis Chimborazo. Folia Daphne Gnidi, 13-15 lin. longa, sesquilineam lata.

21. *B. LANCEOLATA* (H. B. et Kunth nov. gen. am. 4. p. 63), arborea glabra junior viscosa, ramulis sulcatis, foliis petiolatis lanceolatis acutis basi angustatis apicem remotè serratis trinerviis utrinquè glanduloso-punctatis, corymbis terminalibus ramosissimis dichotomis, capitulis  $\sigma^7$  pedicellatis 15-20-floris, invol. hemisphærici squamis scariosis glabris acutis.  $\beta$  in littore Oceani Pacifici juxta Santa Peruvianorum. Folia bipollicaria, 5 lin. lata. Flor.  $\sigma$  ign.

22. *B. RACEMOSA*, fruticosa, foliis ovato-lanceolatis trinerviis acutè serratis, racemis terminalibus.  $\beta$  in sylvis et nemoribus Chili ad Nonquen, Carcamo,

Palomares, Hualpen et Hálqui tractus. Molina racemosa R. et Pav. syst. 209  
Vulgò cum pluribus aliis *Chilca* nuncupatur. Cst. ign. †.

23. *B. COERULESCENS*, suffruticosa erecta glabra, ramis teretibus striatulis, foliis petiolatis elongato-lanceolatis utrinquè attenuatis trinerviis utrinquè punctulatis, infer. ferè à basi grossè serratis, super. minùs serratis imò subintegerrimis, corymbo paniculato laxo, capitulis longè pedicellatis, invol. campanulati squamis lanceolatis acuminatis glaberrimis margine albis scariosis vix subfimbriatis, acheniis glabris.  $\zeta$  in Mexico circa Matamoros legit cl. Berlandier (pl. exs. n. 2343). Cor. ex albo coerulescentes. Capitula 2-3 lin. diam. pallide rufa Pappus fœmin. 1-serialis. (v. s. comm. à cl. invent.)
24. *B. SERRULATA* (Pers. ench. 2. p. 423. n. 87), herbacea erecta glabra, caule basi tereti apice angulato, foliis petiolatis ovato-lanceolatis acuminatis acutè serratis trinerviis minutissimè punctulatis, corymbo composito terminali fastigiato. invol. ♂ campanulati squamis lanceolatis acutiusculis. — in Brasiliæ prov. Rio-Grande. *Conyza serrulata* Lam. dict. 2. p. 85? Specim. mihi obvium à descr. Lamarkii tantum recedit in eo quod sit glabrum nec subglabrum, nec nec fœm. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 798 miss.)
25. *B. BREVIFLORA*, herbacea glabra erecta ramosa, caule ramisque teretibus. foliis subsessilibus elongato-lineari-lanceolatis utrinquè acuminatis acutè serratis triplinerviis, rameis sub 1-nerviis, corymbo composito terminali subpaniculato, invol. ♀ squamis oblongo-linearibus margine subcariosis, ext. obtusis, int. angustioribus acutis pappi parvis glabrisculis — in Brasiliæ prov. Rio-Grande. Affinis *B. ptarmicefolia* sed folia imposita et flores in invol. numerosiores circ. 80. Pappus ♀ albidus 1-serialis vix invol. longior undè primà fronte capitula ♀ videntur ♂. Masc. ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 781 miss.)
26. *B. PHELLOPAPPA*, fruticosa glabra, ramis teretibus apice angulatis, foliis lineari-lanceolatis utrinquè attenuatis basi tenèter trinerviis margine integro apice 1-nerviis serratis subtus punctatis suprà subpunctulatis, corymbo composito terminali, capitulis ♀ pedicellatis, invol. squamis lineari-lanceolatis subacutis, acheniis elongatis striatis, pappo intensè rufè colorato.  $\zeta$  in Americâ olim Hispanicâ sed loc. ign. Folia et rami obscuri in siceo subrafi. Fl. ♀ in invol. 20-25. Achenia ferè 2 lin. longa. Pappus rigidus multisetus. Flor. ♂ ign. (v. s.)
27. *B. ALAMANI*, suffruticosa glabra viscosa, ramis teretibus apice subangulatis. foliis brevissimè petiolatis angustè lineari-lanceolatis triplinerviis utrinquè attenuatis apice subserratis ad utramque faciem punctatis, corymbo paniculato conferto, capitulis pedicellatis ovatis, inv. ♂ squamis lanceolatis acuminatis densè ciliatis laxè subcariosis pappo longioribus.  $\zeta$  in Mexico legit cl. Alamani. Capitula ferè *B. longifolia*, folia ferè *B. coerulescentis* sed angustiora et magis punctata. (v. s. comm. à cl. inv.)
28. *B. LONGIFOLIA*, herbacea erecta glabra ad apicem viscosa, caule tereti, ramis subangulatis, foliis elongatis lineari-lanceolatis breviter petiolatis utrinquè attenuatis à medio ad apicem serratis basi triplinerviis, corymbo composito subpaniculato ferè nudo, inv. campanulati squamis utrinque sexus lanceolatis acuminatis membranaceis densè ciliatis, dorso acheniisque glabris. 2  $\zeta$  in Mexico circa urbem legit cl. Berlandier flor. angusto (pl. exs. n. 624 et 633). Folia impunctata interdum supernè sublepidota, 3-5 poll. longa, semipubescentia. (v. s. ♂ et ♀ à cl. inv. comm.)
29. *B. MARGINALIS*, suffruticosa resinoso-subviscosa glabra, foliis lineari-lanceolatis utrinquè acutis integerrimis aut parè serratis trinerviis, nervis lateralibus margini adproximatis, corymbis compositis foliosis polycephalis, invol. campanulati squamis lanceolatis ciliato-erosis ♂ latioribus magis fimbriatis. ♀ angustioribus, acheniis glabris.  $\zeta$  in versuris et ripis Chaucay et Ceraç prov. Peruviz (R. et Pav.), ad Valparaiso Chilenisium (Gaud.!) Molina parviflora R. et Pav. syst. 209? *Baccharis parviflora* Pers. ench. 2. p. 425. n. 47? Poir. Folia 2 poll. longa, 1  $\zeta$  - 2 lin. lata. Capitul. utriusque sexus 2 lin. diam.

An stirps Chilensis certè eadem ac Peruviana? (v. s. ♂ et ♀ comm. cl. Gaudichand.)

30. *B. FAVILLEI*, frutescens pobe minutissimâ subpulverulenta, ramis teretibus subtriatis, foliis breviter petiolatis lanceolatis utrinquè attenuatis grossè serratis triplinerviis, capitulis ♂ in corymbos compositos terminales subaphyllos digestis 18-20-floris, invol. ovati squamis lanceolatis acutis stramineis. ♀ in Peruviz (Gaud.!), ripis (R. et Pav.), ad sinum Chorillo (Mac-Rae!), et circa Limam (Abadia!). *Conyza frutescens*, etc. Kenill. per. 2. p. 750. t. 37. *B. striata* R. et Pav. syst. 207 excl. syn. ? *B. ivæfolia* Peruviana (nec Capensis) auct. Habitus omninò *Conyze ivæfoliæ* sed etiam genericè distincta. Flor. ♀ ign. (v. s.)
31. *B. SERRIFOLIA*, suffruticosa, ramis striato-angulatis villosulo-hirtis, foliis breviter petiolatis ellipticis triplinerviis conferte et argutissimè serratis secus nervos utrinquè hirtellis, corymbo subpaniculato laxo polycephalo, invol. fl. ♂ glabro, squamis linearibus margine angustè hyalinis apice subciliatis. ♀ in Mexico? Species ex foliis distinctissima. Folia 14 lin. longa, 7 lin. lata, petiolo a lin. hispidulo. Accedit ad *B. multifloram* sed foliis triplinerviis differt. (v. s.)
32. *B. RIPARIA* (H. B. et Kunth nov. gen. am. 4. p. 65), arborea glabra, ramulis striatis, foliis petiolatis oblongo-lanceolatis subacuminatis basi in petiolum angustatis dentato-serratis triplinerviis suprâ nitidis, paniculis terminalibus ramosis corymbosis divaricatis, capitulis ♂ pedicellatis circ. 25-floris, invol. hemisphærici squamis scariosis acutis. ♀ in convallibus Quitensibus prope Guamote et Rio-Chambo (H. et B.) fortè circa Limam? (Née herb.). An var. *B. floribundæ* ut vult Spreng. sed petioli 3-4 nec 6-7 lin. longi? An ♂ *B. polyantha* sed flores in invol. 25 nec 120! Fl. ♀ ign. (v. s.)
33. *B. POLYANTHA* (H. B. et Kunth nov. gen. am. 4. p. 64), fruticosa glabra, ramulis angulatis, foliis petiolatis lanceolatis acutis basi angustatis crenato-serratis triplinerviis suprâ nitidis, paniculis terminalibus ramosissimis corymbosis diffusis, capitulis ♀ breviter pedicellatis 120-130-floris, invol. hemisphærici squamis scariosis acutis. ♀ in Peruviz? (H. B.), in Brasiliâ prope Monte-Video? (Spreng). Folia 4 poll. longa, 10 lin. lata, petiolo 3-4 lin. Flor. ♂ ign.
34. *B. FLORIBUNDA* (H. B. et Kunth nov. gen. am. 4. p. 64. t. 325), arborea glabra, ramulis sulcatis, foliis petiolatis subovato-oblongis subacuminatis basi in petiolum angustatis dentato-serratis triplinerviis suprâ nitidis, paniculis terminalibus ramosissimis corymbosis, capitulis ♂ pedicellatis 40-floris, invol. hemisphærici squamis scariosis acutis. ♀ in Andibus Novo-Granatensium prope Ancalguar et ad flum. Yacanatacu. Affinis *B. polyanthæ* sed folia 2½ poll. longa et petioli 6-7 lin. Arbor 18-24 pedal. Flor. ♀ ign. (v. s. in h. Hænk. à cl. de Sternberg miss.)
35. *B. CONYZOIDES*, herbacea glaberrima, ramis striatis, foliis breviter petiolatis adproximatis rotundo-ellipticis triplinerviis basi integerrimis apice crenato-serratis, paniculis terminalibus subcymosis, invol. ♀ latè campanulati squamis exter. subrotundis obtusis, inter. linearibus apice subciliatis, æcheniis cylindraceis coæstatis hispidiusculis pappo subæqualibus. ♀ — ad fretum Sanctæ-Catharinæ Brasiliæ. Molina conyzoides Less. in Linnæ 1831. p. 139. Fl. ♂ ign.
36. *B. ANOMALA*, suffruticosa ramosa, caule tereti, ramis pubescentibus, foliis petiolatis ovatis basi obtusis apice acutè dentato-serratis trinerviis suprâ sparset puberia subtus densè villosis, paniculis ramos terminantibus laxis, invol. ♂ squamis oblongo-linearibus acutis, fl. masc. stylum exsertum gerentibus. — in Brasiliæ prov. Rio-Grande. Fortè scandens et primâ fronte Clematidem ferè referens. Folia alterna, rarissimè opposita, cum petiolo sesquilineari 7-9 lin. longa, 4 lin. lata. Stigmata maris perfecta sed ovarium abortivum. Pappus rufus scaber. Ind. ♀ ign. (v. s. ♂ in li. Mus. reg. Par. à Mus. imp. Bras. sub n. 856 miss.)
37. *B. PRUNIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 63), fruticosa ramosissima glabra viscosa, ramulis subangulatis, foliis petiolatis oblongis acutis basi an-

gustatis serratis subeoriaceis triplinerviis suprà nitidis, corymbis terminalibus et axillaribus, capitulis form. pedicellatis circ. 250-floris, invol. hemisphaerici squamis scariosis acutis. ♂ in Peruvîâ? Folia 15-16 lin. longa, 6 lin. lata. Flor. masculi ign.

38. *B. SEMISERRATA*, fruticosa, ramis teretibus glabris, ramulis subangulatis apice hirsuto-canescens, foliis lineari-lanceolatis utrinquè attenuatis trinerviis à medio ad apicem serratis, junioribus utrinquè pube sericeâ villosis demùm omninò glabris, pedunculis axillaribus folio brevioribus 1-5-cephalis, invol. ♀ squamis lanceolatis acuminatis, achæniis striatis glabris. ♂ in Brasiliæ prov. Rio-de-Janeiro ad Serra dos Orgaos legit cl. Vauthier (pl. exs. n. 338). Folia 1½ poll. longa, 2 lin. lata, pubem sericeam mox deponentia. Pappus vix rufescens invol. duplò longior. (v. s.)

♂? *glabrata*, ramulis hirsutiusculis, foliis etiam junioribus subglabris. ♂ in Brasiliæ prov. Sancti-Pauli. Nervi fol. laterales validiores paulò alius ori. An spec. propria? (v. s. in h. Mus. reg. Par. à Mus. imp. Brax. mis.)

39. *B. MONTANA*, arborea, ramis teretibus glabris junioribus araneoso-puberulis, foliis oblongo-lanceolatis utrinquè acuminatis margine præsertim ad basin revolutis dentes acutos 2-3 vix apice gerentibus triplinerviis subtus junioribus subcanescentibus demùm subglabris, paniculâ masculâ subracemosa parvi terminali foliis brevior, inv. squamis ovato-lanceolatis margine membranaceis. ♂ in montosis prov. Rio-de-Janeiro vulgaris. Flores albidus octob. proferi ex Lund. mas. Affinis *B. semiserrata* sed folia subintegerrima verè triplinerviis. Ind. form. mihi ign. (v. s.)

40. *B. OXYDONTA*, herbacea? glabra erecta, caule ramisque teretibus, foliis petiolatis lanceolatis basi acutis integris apice acuminatis acutè serratis triplinerviis reticulato-venosis, racemâ compositis axillaribus in paniculam foliosam dispositis, invol. ♂ campanulati squamis lanceolatis acutiusculis. — in Brasiliæ prov. Minas-Geraes ad Villa-Ricca legit cl. Vauthier. Folia ferè 1½ 3 poll. longa. 9-11 lin. lata. Capit. ♂ 3 lin. diam., ♀ ign. An fortè mas *B. triplinervia* sed folia angustiora subulata reticulata, capitula majora. (v. s. ex coll. Vauth. n. 260.)

41. *B. TRIPLINERVIA*, herbacea glaberrima, caule tereti, foliis petiolatis ovato-lanceolatis basi acutis apice longè acuminatis argutè serratis triplinerviis membranaceis, paniculis axillaribus terminalibusque laxis, capitulis ♀ pedicellatis, invol. squamis linearibus subacutis margine membranaceis pappum subæquantibus. — in Brasiliæ prov. Minarum generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 271). Simillima *Conyza triplinervia* sed dioica, folia subtus impunctata argutis serrata, achænia teretiuscula, caulis vix striatus nec sulcato-angulatus. Ind. ♂ ign. (v. s.)

42. *B. PUNCTIGERA*, herbacea pubero-pilosa subviscosa, foliis petiolatis ovato-lanceolatis basi acutis apice longè acuminatis argutè serratis triplinerviis subtus glanduloso-punctatis, corymbo composito fastigiato ramoso polycephalo, capitulis ♀ pedicellatis, invol. squamis linearibus acuminatis glabris. — in Brasiliæ prov. Minarum generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 261). Habitus *Conyze triplinervie* et *Baccharis triplinervie*, ab utroque primâ fronte differt pube pilosâ caulis ramorum petiolorum imò foliorum, et notis indic. Ind. ♂ ign. (v. s.)

43. *B. LUNDII*, fruticosa glabra, ramis teretibus ramulis striatis, foliis longè petiolatis lanceolatis basi cuneato-attenuatis integerrimis cæterum serratis acuminatis triplinerviis membranaceis, corymbo composito terminali fastigiato polycephalo, invol. ♂ et ♀ squamis lanceolatis acutis in ♀ paulò majoribus, achæniis glabris, pappo 1-seriali. ♂ in Brasiliâ circa Rio-de-Janeiro legit cl. Lund. An *B. polyantha* à Spreng. in Brasiliâ citata sed non Kunthiana? *Chrysocoma sancta* fl. flum. 8. t. 3? Petioli 9-10 lin. longi. Limbus folior. 4-5 poll. longus, 1½ latus. Capitula parva 15-20-flora. Affinis *B. oxydonta* sed capit. ♂ dimidiò minora. (v. s.)

44. *B. OLUTINOSA* (Pera. ench. 2. p. 425), suffruticosa glaberrima viscosa, foliis

lanceolatis coriaceis grossè serratis trinerviis et penninerviis basi attenuatis apice acutis, corymbo breviter pedunculato, capitulis ♂ campanulatis, invol. squamis ovali-lanceolatis margine eroso-fimbriatis. ♀ in Chlii prope Talcahuano (Cham. l.), Valparaiso (Bridg.), in aridis et in rudertis ad Concept. et Puchacay prov. (R. et P.). Molina viscosa R. et Pav. syst. 207. Baccharis glutinosa Hook. et Arn. bot. Beech. t. p. 31. Spreng. syst. 3. p. 463 excl. syn. Less. in linnæa 1831. p. 148 et 506. Chilca Fenill. per. 2. t. 37 excl. Tourn. syn. Habitus Conyza ivæfoliæ. Nervi fol. laterales æqualiter à nervo medio et margine distantes. Fl. fœm. ign. undè genus dubium. (v. s. ♂ ex itin. Née et comm. à cl. de Chamisso.)

45. *B. MEDULLOSA*, suffruticosa glabra, ramulis striatis, foliis elongato-linearilanceolatis sessilibus utrinquè attenuatis serratis basi tenuè trinerviis, corymbo æmulo composito fastigiato polycephalo, invol. squamis lanceolato-linearibus acuminiatis. ♀ in Brasiliæ prov. Rio-Grande. Pl. fœm. ign. Canlis digiti crassitie medullâ amplâ donatus. Fol. 5 poll. longa, 5-6 lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 850 miss.)

46. *B. PUNCTULATA*, suffruticosa glabra, ramis teretibus apice angulatis, foliis breviter petiolatis oblongis basi cuneatis integris cæterùm regulariter serratis acutis trinerviis subtili minutè et opacè punctatis, racemis compositis parvis ramulos axillares terminantibus et paniculam foliosam constituentibus, invol. ♀ ovati squamis lineari-oblongis obtusis margine submembranaceis ciliolatis, achænio glabris. ♀ in Brasiliæ prov. Rio-Grande. Fol. 2 poll. longa, 5 lin. lata. Pappus vix subrufescens. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 970 miss.)

47. *B. SORDASCEA*, suffruticosa, ramis sulcato-angulatis puberalis, foliis sessilibus oblongis basi attenuatis apice vix acutis trinerviis velutinis margine tenuissimè serrato-ciliatis, ramulis in paniculam dispositis capitula 3-7 subsessilia gerentibus, invol. ♂ squamis pluriserialibus oblongis subobtusis ad margines densè et longè molliter ciliatis. ♀ in Mexico legit cl. Hænke. Species *B. hirtæ* subaffinis sed ab eâ et ab omnibus ferè involucri distincta. Flor. ♀ ign. ♂ parùm noti. (v. s. in h. Hænke à cl. de Sternberg miss.)

48. *B. HIRTA*, herbacea erecta subsimplex tota strigoso-hispida, caule tereti striato, foliis sessilibus cuneatis apice grossè 3-7-dentatis basi trinerviis nervoso-reticulatis, corymbo composito terminali subpaniculato, capitulis 1-3 ad apices pedicellorum, invol. ♀ squamis lanceolatis vix acutis margine membranaceis subciliatis dorso glabriusculis, achænio jun. subpuberulo adulto glabro. — in Brasiliæ prov. Rio-Grande. Herba bipedalis. Folia 2 poll. longa, 5-8 lin. lata. Dentes sen lobali fol. inf. obtusi, sup. acuti, omnium mucronati. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 833 et 967 miss.)

49. *B. MYRICÆFOLIA*, fruticosa, ramis teretibus glabris, ramulis teretiusculis pube molli cano-hirsutulis, foliis oblongis basi longè attenuatis subpetiolatis apice dentibus utrinquè 1-2 incisus aut integerrimis trinerviis, nervis lateralibus tenuissimis, junioribus floccoso-pubescentibus adultis glabris, pedunculis axillaribus 1-2-cephalis folio brevioribus, invol. ♀ squamis lanceolatis acutis, achænio striato glabro pappi longitudinè ferè æquali. ♀ in Brasiliæ prov. Sancti-Pauli. Affluis *B. semiserratæ* sed faciliè distincta. Capitula dimidiò minorâ longius pediculata. Pappus brevis intensè rufus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 484 miss.)

50. *B. SCANDENS* (Pers. ench. 2. p. 424. n. 33), fruticosa scandens, foliis ovato-lanceolatis dentatis bidentatisque triplinerviis, corymbis terminalibus. ♀ in Peruviz versuris et aridis per Chancay et Cercado provincias ubi vern. *Chilca*. Molina scandens R. et Pav. syst. p. 205. Triorgyalis. Cæt. ign.

§. 2. Cuneifoliæ, nempe foliis obovatis cuneatisve uninerviis aut triplinerviis non imbricatis, ramis apteris.

51. *B. MACELLANICA* (Pers. ench. 2. p. 425. n. 23), fruticosa demisso-cæspitosa glabra viscosa, ramulis angulatis, foliis sessilibus confertis coriaceis obovato-



- cuneatis obtusis alii integerrimis alii apice obtusè tridentatis, capitulis solitariis ad apices ramulorum sessilibus, invol. ovati squamis margine ciliato-fimbriatis, ♂ ovali-lanceolatis, ♀ angustius lanceolatis acuminatis, acheniis striatis glabris. ♂ ad fretum Magellanicum (Lam.), ad portum Egmont et Desendo (Née!), in insulis Maclovianis (Gaud. ! D'Urv. !). *Conyza Magellanica* Lam. dict. 2. p. 91. *Baccharis tridentata* Gaud. ! fl. mal. p. 15. D'Urv. ! fl. mal. non Vahl Differt à spec. Vahl ramis glabris nec subvillosis, foliis 2-3 lin. longis nec pollicaribus, etc. (v. s. ♂ et ♀.)
52. *B. CUNEIFOLIA*, fruticulosa demissa glabra viscosa ramosissima, ramulis subangulatis, foliis sessilibus confertis coriaceis obovato-cuneatis obtusis ad apicem obtusè repando-subdentatis, capitulis ad apices ramulorum sessilibus confertis. ♂ ad fretum Magellanicum (Lam.), ad portum Egmont (Née!) et in Brasiliâ (Vahl), ad Monte-Video (Lam.). *Conyza cuneifolia* Lam. dict. 2. (1780) p. 91. *Baccharis sessiliflora* Vahl symb. 3. (1794) p. 97. An stirps Monte-Videos eadem ac Magellanica? An mera *B. Magellanica* varietas? (v. s. ♂.)
53. *B. ALPINA* (H. B. et Kunth nov. gen. am. 4. p. 49), fruticosa procumbens glabra, foliis obovato-spathulatis obtusis integerrimis coriaceis aveniis, capitulis ♂ subterminalibus solitariis sessilibus, invol. hemisphærici squamis obtusis, intimis oblongo-linearibus, recept. convexo nudo. ♂ ad radices montis Pinchinchæ et Antisanæ Quitensium. Folia 3 lin. et paulò longiora. Pappus ♂ apice incrassatus. Flor. in invol. circ. 7. An à *B. buxifolia* jam innuente Spreng satis differt? sed cl. Kunth qui herb. Juss. sub oculis habuit ab eâ sejunct.
54. *B. ? HUMIFUSA* (H. B. et Kunth nov. gen. am. 4. p. 48. t. 322), fruticosa prostrata repens glabra, foliis obovato-spathulatis obtusis integerrimis coriaceis aveniis, capitulis subterminalibus solitariis subsessilibus, invol. ovati squamis lanceolatis scarionis acutis, recept. conico paleis subulatis obsito, achenio striato. ♂ in radicebus montis Pinchinchæ et Antisanæ Quitensium. Peruvia sp. Cass. dict. 37. p. 480. Affinis *B. Magellanica*. Pappus ♂ apice clavellatus. Flores in inv. circ. 35. — *B. cæspitosa* Pers. ench. 2. p. 428. n. 38, quæ Molina cæspitosa R. et Pav. syst. p. 203 ex Peruviz montanis orta sed vix descripta. ab hac fortè non differt? (v. s.)
55. *B. PROSTRATA* (Pers. ench. 2. p. 424. n. 37), fruticosa prostrata, foliis obovatis denticulatis integrisque, pedunculis axillaribus 1-cephalis. ♂ in Peruviz Andibus alpinis ad Bombom, Huamalies, Canta et Pillau. Molina prostrata R. et Pav. syst. 204. Vernaculè *Pachatacya*. Decoctum in dysuriâ prodest. Cæc. iqa
56. *B. FRIGIDA* (H. B. et Kunth nov. gen. am. 4. p. 49), fruticosa ramosissima glabra, foliis sessilibus lanceolato-spathulatis acutis integerrimis coriaceis uninerviis suprâ viscoso-nitidis subtus punctatis, capitulis ♂ terminalibus solitariis sessilibus, invol. hemisphærici squamis membranaceis, exter. ovato-oblongis, int. linearibus. ♂ in frigidis Peruviz. Flor. in invol. ♂ 60-65 albi, ♀ iqa
57. *B. MICROPHYLLA*, fruticosa erecta? ramosissima glabra viscosa, ramis terribus, foliis petiolatis ovatis aut obovato-cuneatis denticulatis, capitulis numerosis breviter pedicellatis 5-floris, invol. squamis lineari-oblongis obtusis interioribus duplò longioribus, achenio glabriusculo. ♂ in Peruviz legit cl. Hænke. Folia petiolo incluso 4 lin. longa, 3 lin. circ. lata, dentibus parvis denticulata. (v. s. in H. Hænke à cl. de Sternberg comm.)
58. *B. POLYGAMIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 49), fruticosa glabra subviscosa, ramulis subangulatis, foliis breviter petiolatis lanceolato-spathulatis acutis integerrimis coriaceis uninerviis suprâ nitidis subtus subpunctatis, capitulis ♀ axillaribus sessilibus subspicatis, invol. hemisphærici squamis acutis culis scarionis, achenio striato glabro. ♂ juxta Moran Mexicanorum et Paramo et decl. Andium Popayauensium (H. et B.), in Monte-Video Spreng. An plures hic confusa?
59. *B. OBTUSIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 51), fruticosa glabra, ramulis angulatis subviscosis, foliis sessilibus obovatis apice rotundatis et subpandis basi cuneatis coriaceis triplinerviis, capitulis ♀ terminalibus cap-

congestis sessilibus, invol. squamis scariosis acutiusculis, acheniis costatis glabris.  $\frac{3}{4}$  in frigidis prope Gonzanama Peruvianorum. Folia 8-10 lin. longa, 5-6 lata. Pappus fuscescens. Fl. masc. ign.

60. *B. ODORATA* (H. B. et Kunth nov. gen. am. 4. p. 52), fruticosa ramosissima glabra viscosa, ramulis subangulatis, foliis brevissimè petiolatis ellipticis obtusis basi subcuneatis integerrimis coriaceis aveniis viscido-nitidis, capitulis  $\sigma^7$  terminalibus conglomeratis sessilibus, invol. hemisphærici squamis acutiusculis.  $\frac{3}{4}$  in montosis prope Pasto Novo-Granatensium (H. et B.) et fortè in Brasiliâ? (Spr.). Folia ferè Buxi semipoll. longa, odorem balsameum redolentia, Fl. fœm. ign.

61. *B. BERNARDIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 57), fruticosa ramosissima glabra viscosa, ramis teretiusculis, foliis subsessilibus obovatis obtusis mucronatis basi cuneatis integerrimis coriaceis reticulato-venosis nitidis, corymbis  $\varnothing$  terminalibus densis, invol. hemisphærici squamis acuminatis, acheniis costatis glabris.  $\frac{3}{4}$  in altâ planitie Quitensi ad radices montis Rucu-Pichincha. Folia 15 lin. longa, 6 lin. lata. Pl.  $\sigma^7$  ign.

62. *B. SQUARROSA* (H. B. et Kunth nov. gen. am. 4. p. 67), fruticosa glabra, ramis (caulibusve?) sulcato-angulatis pauculato-virgatis apice 1-cephalia, foliis breviter petiolatis spatulato-lanceolatis acutis integerrimis subcarnosis subaveniis viscosis, inv.  $\varnothing$  squamis lineari-oblongis squarrosè imbricatis apice subfimbriatis.  $\frac{3}{4}$  in aridis montium Mexicanorum prope Guanajuato, ad radices montis La Buffa. Folia 10-11 lin. longa, 2 lin. lata.

63. *B. PILULARIS*, fruticulosa ramosissima glabra viscosa, ramulis angulatis, foliis sessilibus obovato-cuneatis obtusis integris aut sæpius subdentatis 1-nerviis coriaceis, capitulis ad apices ramorum et ramulorum axillarum solitariis globosis foliis parvis bracteatis, invol.  $\sigma^7$  squamis oblongis subcatis margine membranaceis apice subfimbriatis.  $\frac{3}{4}$  in Californiâ legit cl. Douglas. Folia 4 lin. longa, 2-3 lin. lata, summa capitulo breviora. Capit. 2 lin. diam. Fl.  $\varnothing$  ign. (v. s.  $\sigma^7$  ex h. soc. hort. Lond. comm. à cl. Benth.)

64. *B. AXILLARIS*, fruticulosa glabra, ramis teretibus junioribus striatis, foliis sessilibus obovatis obtusis basi cuneatis subnerviis integerrimis aut hinc inde subdentatis patulis coriaceis, capitulis ad axillas foliorum solitariis pedicellatis et idè racemos foliatis constituentibus, invol.  $\sigma^7$  ovati squamis lanceolatis acutis, flor. circ. 6 in capit.  $\frac{3}{4}$  in Brasiliæ prov. Sancti-Pauli. Folia vix 3 lin. longa et 1  $\frac{1}{2}$  lata. Pedicelli folio paulò breviores. Ind.  $\varnothing$  ign. (v. s.  $\sigma^7$  in h. Mus. reg. Par. à Mns. imp. Bras. sub n. 481 mis.)

$\beta$ . *dentata*, foliis cuneatis omnibus apice 3-5-dentatis.  $\frac{3}{4}$  in prov. Rio-Grande. Capit.  $\sigma^7$  var.  $\alpha$  conformia. Capit.  $\varnothing$  brevissimè pedicellate oblonga cylindracea var.  $\gamma$  omninò conformia. (v. s. cum var.  $\alpha$ ,  $\sigma^7$  sub n. 956,  $\varnothing$  sub n. 953.)

$\gamma$ . *acutiloba*, foliis dentibus profundis acutis 5-7-incisis.  $\frac{3}{4}$  in Bras. prov. Sancti-Pauli. Similis var.  $\beta$ , sed foliis acutè incisis non apice tantùm sed etiam ad margines dentato-lobatis. Capit.  $\sigma^7$  similia in n. 417 et 419 herb. Mus. imp. Bras. Capitula  $\varnothing$  in n. 418 obvia sunt cylindracea duplò longiora brevius pedicellate, squamis lanceolato-linearibus elongatis acutis. Pappus rufescens pluriserialis. Achenium striatum nitidum. An certè aliarum (quarum specim.  $\sigma^7$  tantùm vidi) conspecifica? (v. s. cum  $\alpha$  et  $\beta$  sub n. 417, 418, 419.)

65. *B. MICROPHYLLA* (H. B. et Kunth nov. gen. am. 4. p. 53), fruticosa ramosissima glabra viscidula, ramulis angulatis, foliis subsessilibus cuneatis apice tridentatis coriaceis uninerviis, capitulis  $\varnothing$  axillaribus breviter pedunculatis, invol. squamis subsquarrosis acutiusculis.  $\frac{3}{4}$  prope Santa-Fe de Bogotá. Bacch. tridentatæ proxima. Pl.  $\sigma^7$  ign.

66. *B. PEDICELLATA*, fruticosa ramosissima glabra viscosa, ramis teretiusculis, foliis obovato-cuneatis sessilibus apicem versus paucidentatis coriaceis sub-3-nerviis, venis lateraliter tenuibus aut subnullis, pedicellis axillaribus subnudi.

- striato-sulcatis 1-cephalis, invol. ♀ squamis ovali-lanceolatis acutis apice subciliatis. ♀ in Chili ex h. Haenke. Folia ferè B. cuneifoliae aut R. concava viscoso-nitida 8-9 lin. longa 4 lin. lata. Pedicelli bracteolâ 1-2 instructi 7-9 lin. longi, involucrique pallida. (v. s. in h. Haenk. à cl. de Sternberg miss.)
67. B. CUNEATA, fruticosa glabra, ramulis angulatis, foliis cuneiformibus apice obtusè 3-dentatis basi in petiolum brevissimum attenuatis, capitulis ad apicem ramulorum congestis subsessilibus ovatis, invol. squamis obtusis marginè scarioso albo subfimbriato, æchæniis glabris. ♀ in Mexico. *Erigeron tricusnatum*. Lin. f. suppl. 368? et ideò forsan *Conyza tricusnata* Willd. sp. 3. p. 1916 excl. Lam. syn. *Baccharis tricusnata* Pers. enob. 2. p. 424. n. 26. Folia accedit ad B. glomerulifloram sed glomeruli terminales nec axillares. (v. s. comm. à cl. Alaman et Mairet.)
68. B. GLOMERULIFLORA (Pers. ench. 2. p. 423. n. 11), fruticosa glaberrima, ramulis angulatis, foliis lævibus cuneato-obovatis in petiolum brevem attenuatis apice obtusis grossè dentatis summis obovato-oblongis integerrimis, capitulis ad axillas foliorum sessilibus congestis, invol. ♂ campanulatis squamis oblongis obtusis. ♀ in sylvis Amer. boreal. littoralis à Virginia ad Carolinam. (Fras! Bosc!) *Baccharis sessiliflora* (Pursh) Michx. fl. 2. p. 135. Ell sketch 2. p. 320 non Vahl. Pappus ♂ brevis, ♀ multò longior. (v. s. ♂.)
69. B. CONNANGUINEA, fruticosa ramosissima glabra junior subviscosa, ramulis decurrenti-angulatis, foliis sessilibus obovato-cuneatis obtusis dentibus acutis utrinquè 2-4 notatis rigidis, capitulis ad apices ramorum et ramulorum sessilibus rarò solitariis sæpiùs 2-4 aggregatis, invol. ♀ squamis oblongo-linearibus obtusis marginè membranaceis subfimbriatis, intimis acutiusculis. ♀ in California legit cl. Douglas. Valdè accedit ad B. confertam, cuneifoliam, etc. Folia 6 lin. longa, 3 lin. lata in sicco suprâ sublepidota. Ind. ♂ ign. (v. s. ♀ comm. ex h. soc. hortic. Lond. à cl. Bentham.)
70. B. SUBDENTATA, fruticulosa glabra, ramulis subangulatis junioribus foliisque subviscosis, foliis sessilibus obovatis ovalibusve coriaceis integerrimis aut hac indè apice subdentatis imâ basi subtrinerviis, capitulis ♀ oblongis in apice breves foliis intermixtis confertis, squamis oblongis acutis inter. longioribus pappum subæquantibus. ♀ in Brasiliæ prov. Minas-geraes (Vauth! pl. ex. v. 339), et Sancti-Pauli (h. imp. Bras. n. 474). Folia 6-9 lin. longa, 2-4 lata. Ind. ♂ ign. (v. s. ♀.)
71. B. OBOVATA (DC. non Hook. et Arn.), fruticosa ramosissima glabra viscosa, ramulis subangulatis, foliis sessilibus obovatis obtusis basi cuneatis integerrimis apice dentibus 5-7 notatis uninerviis, capitulis ♂ ad apices ramorum 3-5 congestis sessilibus, invol. squamis oblongo-lanceolatis acutiusculis. ♀ in Peruviae collibus aridis affatim ad Tarmæ oppidum, vernac. *Tacramacho* apell. Molina obovata R. et Pav. syst. p. 206 (v. s. ex h. Dombeý comm. à Mus reg. Par.)
72. B. SINUATA (H. B. et Kunth nov. gen. am. 4. p. 53), fruticosa ramosissima glabra viscosa, ramulis angulatis, foliis sessilibus subrhomboides basi cuneatis apicem versus sinuatis subcoriaceis, corymbo terminali subtrichotomo, invol. ♂ hemisphærici squamis obtusiusculis, recept. paleas subulatas gerente. ♀ in regione temperatâ Peruviae inter Guancabauha et Sondorillo. Tarseniz p. Cass. dict. 37. p. 478. Folia 1; poll. longa, 15 lin. lata. Cor. ♂ albæ 35-45 in invol., fem. ign.
73. B. PHILIPPENSIS (H. B. et Kunth nov. gen. am. 4. p. 55), fruticosa ramosissima glabra junior viscosa, foliis brevissimè petiolatis obovato-cuneatis apice 3-5-dentatis coriaceis, capitulis terminalibus conglomeratis sessilibus, invol. ♂ squamis scariosis acutis. ♀ in Venezuela prope San-Felipe Affinis dicitur B. tridentatæ. Planta ♀ ign. Cor. ♂ albæ 30 in invol.
74. B. HAMBATENSIS (H. B. et Kunth nov. gen. am. 4. p. 62), fruticosa glabra ramulis sulcato-angulatis, foliis petiolatis lanceolato-oblongis acutis basi cuneatis supernè dentatis subcoriaceis uninerviis reticulato-venosis, corymbo terminalibus subpaniculatis, invol. ♂ hemisphærici squamis scariosis obtu-

- aisculis.  $\frac{5}{8}$  in arida temperatis juxta Hambato Qutensium. Cor.  $\sigma$  albæ circ. 30 in invol.,  $\varnothing$  ign. Folia pollicaria, 4 lin. lata.
75. *B. VAUTHIERI*, fruticosa glabra subviscosa, ramulis decurrenti-angulatis, foliis oblongo-cuneatis basi attenuato-subpetiolatis apice obtusis subtridentatis nitidulis venulosis; capitulis  $\sigma$  ad axillas solitariis breviter pedicellatis folio dimidiò brevioribus ovato-globosis, invol. squamis exter. minimis, inter. oblongis subobtusis disci longitudine.  $\frac{5}{8}$  in Brasiliæ prov. Minarum general. ad Tejucco legit cl. Vauthier (pl. exs. n. 328). Accedit hinc ad *B. tridentatam* illinc ad pilularem. Fœm. ign. Folia pollicem longa. Capit. pedicello incluso 3-4 lin. longa. (v. s. comm. à cl. coll.)
76. *B. TRIDENTATA* (Vahl symb. 3. p. 98), fruticosa glabra, ramulis decurrenti-angulatis, foliis oblongis basi attenuatis apice tridentatis rariùs integris uninerviis, capitulis ad axillas summas et ad apices ramorum congestis subsessilibus oblongis, invol. utriusque sexus oblongis pallidis subacutiusculis, achænio striato glabro.  $\frac{5}{8}$  in collibus aridis Brasiliæ (Vahl), circa Bahiam (Salzm.!), in prov. Minarum general. prope Tejucco (Vanth! sine num.), et Rio-Grande (h. imp. Bras. sub n. 963). *B. tridentata* Pers. et Spreng. excl. syn. Deser. Vahl! non satis completa hæc potius quàm cæteris convenit, sed specim. non vidi. (v. s.)
- $\frac{2}{3}$  *pluridentata*, foliis ad apicem dentibus 5-7 acutis instructis.  $\frac{5}{8}$  circa Bahiam in dumetis legit cl. Blanchet. An fortè spec. propria ad *B. oblongifolias* referenda? (v. s.)
77. *B. BRAVIFOLIA*, fruticosa ramosissima glabra, ramulis subangulatis, foliis obovato-cuneatis obtusis apice 3-dentatis, summis integris, capitulis ad apices ramulorum spicato-congestis numerosissimis, invol.  $\frac{1}{2}$   $\varnothing$  squamis oblongis pallidis subacutiusculis, achænio glabro striatulo.  $\frac{5}{8}$  in campis editis prov. Sancti-Pauli Brasiliæ legit cl. Lund. fl. nov. An var. *B. tridentatæ*, sed folia 3-4 nec 12 lin. longa, obovato-cuneata nec oblonga, capitula paulò minorà multò magis conferta et numerosa (v. s. comm. à cl. Lund.)
78. *B. RETICULARIA*, fruticosa glabra subviscosa ramosa, ramulis striato-subangulatis, foliis obovatis basi cuneato-attenuatis subsessilibus apice obtusis et obtusè 3-5-dentatis utrinquè reticulo tenui nervulorum ornatis, capitulis ad apices ramulorum circ. 3 in axillis ultimis solitariis subsessilibus, invol.  $\varnothing$  5-flori squamis pallidis oblongis vix acutis, achænio striato glabro.  $\frac{5}{8}$  in Brasiliæ prov. Minarum generalium ad Villa Rica legit cl. Vauthier (pl. exs. n. 329). Folia ferè Buxi sed obtusiora magis obovata et dentata. Species affinis *B. pauciflorescens*. (v. s. comm. à cl. coll.)
79. *B. SALZMANNI*, fruticosa ramosissima glabra viscosa, ramulis striato-angulatis, foliis obovato-cuneatis subsessilibus dentibus grossis 7-9 instructis suprâ nitidis nervoso-reticulatis penniveniis, capitulis ad apices ramorum inter folia subcongesto-corymbosis subsessilibus, inv.  $\sigma$  squamis acutis pappo æqualibus,  $\varnothing$  oblongi squamis elongatis acuminatis pappo paulò brevioribus, achænio striato glabro.  $\frac{5}{8}$  in collibus aridis circa Bahiam legit Salzman. Pappus albidus. Folia 12-15 lin. longa, 6-7 lata. (v. s. comm. à cl. coll.)
80. *C. FLATYPODA*, fruticosa glabra resinoso-viscosa, ramis striato-angulatis, floridiis ancipiti-compressis, foliis petiolatis obovato-cuneatis obtusis grossè dentatis coriaceis uninerviis venis hinc inde innumeris pennatis subreticulatis, capitulis  $\sigma$  ad apices ramorum florid. densè aggregato-corymbosis, invol. squamis ovato-oblongis obtusis.  $\frac{5}{8}$  in Brasiliâ (herb. L'Hér.!) prov. Minarum General. ad Mariannam (Vanth! pl. exs. n. 283). Folia 3 poll. longa, 1-1  $\frac{1}{2}$  poll. lata. Capitulorum glomeruli cincti foliis floralibus nunc brevibus oblongis, nunc oblongis oblongo-spathulatis. (v. s.)
81. *B. CONFERTA* (H. B. et Kunth nov. gen. am. 4. p. 55), fruticosa ramosissima glabra apice subviscosa, ramulis decurrenti-angulatis, foliis sessilibus obovato-oblongis acutis basi cuneatis apicem versus utroque latere dente unico instructis subaveniis, capitulis  $\varnothing$  terminalibus confertis subsessilibus, invol. squamis lanceolatis, inter. acuminatis demum squarrosis, achæniis striatis glabris.  $\frac{5}{8}$  in

Mexici temperatis prope Cuernavaca (H. et B.) et ad Real del Monte (Berl. pl. exs. n. 397). Fl. masc. ign. Folia 5-7 lin. longa. Affinis videtur *B. tridentata* et *cuneata*. (v. s.)

82. *B. PTERONIOIDES*, fruticosa ramosissima glabra ad apices viscosa, ramis teretibus, foliis breviter petiolatis oblongo-cuneatis obtusis mucronatis dentibus utrinque 2-3 acutis instructis 1-nerviis utrinque punctatis, capitulis ad apices ramulorum solitariis sessilibus, inv. ♀ squamis lanceolatis acuminatis 5 in Mexico inter Tampico et Real del Monte maio flor. legit. cl. Berlandier (pl. exs. n. 398). Pappus ♀ corollam superans invol. duplò longior pallide rufescens. Folia 3-5 lin. longa. (v. s.)

83. *B. ALATERNOIDES* (H. B. et Kunth nov. gen. am. 4. p. 51), fruticosa ramosissima, ramulis angulatis pulverulento-viscosis, foliis petiolatis obovatis basi cuneatis obsolete et remotè dentatis coriaceis subvenosis glabris brevissimis, capitulis ♂ terminalibus congestis pedunculatis, invol. hemisphaerici squamis scariosis acutis subfimbriatis. 5 in altà planitie Peruvianà ad Caxamarea. Cor. in invol. ♂ 45-50, fœm. ign. Folia 6-7 lin. longa, 4-4 1/2 lata.

84. *B. POEPPIGIANA*, fruticosa glabra viscosa ramosissima, foliis obovatis basi cuneatis subsessilibus apice obtusis repando-dentatis, capitulis pedicellatis ad apices ramorum paucis umbellatis, invol. ♂ campanulati squamis lanceolatis vix apice subciliatis. 5 inter fragmenta rupium graniticarum inter la Guardia et los Ojos de Agua Chilensium. *Baccharis alaternoides* Poepp.! pl. chil. exs. 2. n. 102. non Kunth. Differt à spec. Kunthianà pedicellis folio longioribus, petiolo æqualibus. Affinis *B. cuneifolia* sed magis demissa et viscosa, dentes obtusiores, etc. (v. s. ♂ ex coll. Poepp.)

85. *B. UMBELLIFORMIS*, fruticosa glabra, foliis obovatis obtusis cuneatis et sessilibus planis dentatis uninerviis epunctatis, capitulis pedicellatis ad apices ramorum umbellatis, invol. ♂ campanulati squamis lanceolatis margine scarioso densè ciliatis. 5 in Chili? Poepp. pl. exs. 860. An *Baccharis obovata* Hook. et Arn. bot. Beech. 1. p. 30? sed non Molina obovata R. et Pav. — (v. s. ♂ ex coll. Poepp.)

ℓ. invol. ♀ squamis minus ciliatis — Poepp.! exs. n. 695. An *formosa* var. α? An spec. alt.? (v. s. ex coll. Poepp.)

86. *B. RUPICOLA* (H. B. et Kunth nov. gen. am. 4. p. 52), subarbores glabra ramulis subangulatis, foliis subsessilibus confertis ellipticis obtusiusculis basi subcuneatis apicem versus remotè denticulatis reticulato-venosis subhis viscoso-punctulatis, capitulis ♂ terminalibus congesto-corymbosis, invol. hemisphaerici squamis coriaceis obtusis. 5 in monte altissimo Purace juxta artem Popayan. Arbor 12-18 ped. Folia 6 lin. longa 3 1/2 lata. Fl. ♀ ign.

87. *B. RESINOSA* (H. B. et Kunth nov. gen. am. 4. p. 52. t. 323), fruticosa ramosissima glabra, ramulis striato-angulatis resinoso-viscosis, foliis breviter petiolatis obovato-oblongis acutiusculis basi subcuneatis 3-5-dentatis coriaceis subaveniis viscoso-nitidis subtus punctatis, capitulis terminalibus conglomeratis, ♂ sessilibus, ♂ brevissimè pedunculatis, invol. ovati squamis acutiusculis, achenio sulcato glabro. 5 in Americà meridionali. Folia 8-9 lin. longa. 4-4 1/2 lata. (v. s. ?)

88. *B. ROTUNDIFOLIA* (Spreng. syst. 3. p. 465?), fruticosa, ramis teretibus, ramulis striato-angulatis subviscosis, foliis sessilibus obovato-subrotundis, apice dentibus 3-5 repandis trinerviis coriaceis subtus præcipuè albido-furfuraceis, capitulis ♂ ad apices ramorum congestis sessilibus, ♀ minus congestis subspicatis, bracteis ♂ parvis obovatis, ♀ invol. subæqualibus tridentatis, invol. squamis ♂ ovato-lanceolatis acutis, ♀ magis elongatis vix acutis, achenio striatis glabris. 5 in Brasiliæ prov. Rio-Grande. Phrasis Sprengelii optime indiv. masculo convenit. Folia 6-9 lin. diam. Pappus ♀ rufus longus. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. miss. ♂ sub. n. 964. ♀ sub. n. 965.)

89. *B. CONGESTA*, fruticosa ramosissima glabra viscosa, ramulis striato-angulatis foliis sessilibus obovatis basi breviter cuneatis apice obtusis margine dentibus 3-5 obtusis utrinque notatis penninerviis rigidulis, capitulis ad apices ramorum

sessilibus congestis, invol. ♂ hemisphaerici squamis ovali-oblongis obtusiusculis margine fimbriato-ciliatis. — ♂ ad Mexico et Real del Monte legit cl. Hænke. Accedit ad B. confertam et cuneifoliam. Fl. foem. ign. (v. s. in h. Hænke à cl. de Sternberg miss.)

90. B. XALAPENSIS (H. B. et Kunth nov. gen. am. 4. p. 56), fruticosa glabra, ramulis sulcato-angulatis, foliis subsessilibus obovatis apicem versus acutè 5-dentatis basi cuneatis coriaceis subvenosis, junioribus subviscosis, capitulis ♂ terminalibus capitato-congestis sessilibus, invol. subhemisphaerici squamis ext. ovatis, int. oblongo-linearibus. ♂ in montibus Mexicanis prope Xalapa ubi Escovilla dicitur. Aff. B. cuneifoliae. Ind. ♀ ign.

91. B. HETEROPHYLLA (H. B. et Kunth nov. gen. am. 4. p. 62), fruticosa ramosissima glabra viscosa, ramulis sulcato-angulatis, foliis petiolatis oblongo-lanceolatis apicem versus sinuato-dentatis basi cuneatis membranaceis, pedunculis axillaribus et terminalibus 2-4-cephalis, capitulis ♂ subsessilibus hemisphaericis, inv. squamis ext. ovatis, int. oblongis obtusiusculis. ♂ in convallibus Mexicanis prope Guanajuato et fodinam Valencianae (H. B.). An ad Montevideo, ut vult Spreng. syst. 3. p. 464?

α. foliis obtusis apicem versus sinuato-dentatis.

β. foliis acutis apicem versus dente 1-2 instructis. An spec. dist.?

92. B. HALIMIFORMIS, fruticosa ramosissima apice tenuissimè subpubescens, ramulis angulatis, foliis sessilibus obovato-cuneatis subacutis apice dentibus 5-7 subrepandis penninerviis nervis suprà elevatis reticulatis, capitulis ♂ ad apices ramorum et axillas fol. pedicellatis subcorymbosis dispositis, invol. squamis pallidis subnerviis intimis acuminatis margine hyalinis. ♂ in Brasiliâ circa Bahiam. Foem. ign. Folia accedunt ad B. halimifoliam. (v. s.)

93. B. NERI, fruticosa glabra viscosa, ramulis subangulatis, foliis sessilibus oblongo-subcuneatis obtusis uninerviis dentibus 3-7 notatis, capitulis ♂ ad apices ramorum 3-5 sessilibus congestis circ. 15-floris, invol. squamis ovali-oblongis obtusis, int. linearibus vix acutis. ♂ in Amer. merid. Cordillera del Portillo legit cl. Née. Folia 3½ lin. longa, 1-1½ lata, sæpè deflexa coriacea. Fl. ♀ ign. (v. s. ex itin. Née.)

94. B. CONCAVA, fruticosa ramosissima, ramulis angulatis pubero-subvelutinis, foliis latè obovato-cuneatis apice obtusè tri-rariùs quinque-dentatis sessilibus glabris crasso-coriaceis uninerviis summis circa capitula subconcavis, capitulis ad apices ramulorum 3-5 congestis subsessilibus, invol. campanulati squamis margine scariosis oblongis in ♂ obtusiusculis, in ♀ inter. linearibus subacutis, acheniis compressis glabris. ♂ in fruticetis collium Chilensium ad Conceptionem (R. et Pav. Mac-Rae), Valparaiso (Bert. ! Gaud.), Topocalma (C. Gay.!). Vulgò *Gaultro* ex R. et Pav. *Guanchu* ex Bert. Molina concava R. et Pav. syst. 206. Baccharis tridentata Poepp. ! coll. chil. n. 211. Baccharis resinosa Hook. et Arn. bot. Beech. 1. p. 31. excl. in utr. syn. (v. s. ♂ et ♀.)

95. B. INTERMEDIA, fruticosa glabrescula resinoso-subviscosa, foliis lineari-cuneatis basi attenuatis apice repando-dentatis margine subrevolutis, capitulis ad apices ramulorum congesto-corymbosis brevissimè pedicellatis, invol. ♀ squamis ovali-lanceolatis vix acutis margine scariosis, achenio glabro striato. ♂ in Chili circa Valparaiso (Gaud.!), in fruticetis collium (Bert.!). Media ferè inter B. rosmarinifoliam et concavam. (v. s.)

96. B. VAHLII, fruticosa glabra, ramis teretibus, ramulis angulatis, foliis obovato-cuneiformibus retusis integerrimis penninerviis, capitulis plurimis sessilibus ad apices ramorum congestis ovatis, inv. ♂ squamis ovali-oblongis exter. obtusis inter. vix acutis. ♂ in insulis Caribaeis nempe Montserrat (Vahl), Santo-Domingo (Poit.!), Sancta-Cruce (Vest.). Baccharis dioica Vahl symb. 3. p. 93. t. 74. Spreng. syst. 3. p. 459 (excl. syn. Kunth). Lam. ill. t. 698. t. 3. Cl. Vahl dixit folia trinervia sed in meis specim. (verisim. authenticis) potius penninervia video. Capitula juniora oblonga subviscosa. Nomen sperit. mutavi cum ex gener. caract. Baccharides omnes sint dioice. (v. s.)

97. *B. HALIMIFOLIA* (Linn. sp. 1204), arborea glabra sublespidoto-glaucescens, ramulis angulatis, foliis obovatis supernè grossè angulato-dentatis basi cuneatis in petiolum subattenuatis, summis lanceolatis integerrimis, capitalis ♂ subglobosis pedicellatis solitariis adgregatis, ♀ fasciculatis laxè subpaniculatis ovato-oblongis, invol. squamis, ♂ ovatis ♀ inter. elongatis. 5 in maritimis Amer. bor. à Marylandiâ ad Floridam (Pursh), in Carolinâ (Michx. l.), Georgia (h. DuRoi.), Baccharis halimifolia Duham. arbr. ed. 1. v. s. t. 35. ed. 2. v. 1. t. 60. *B. cuneifolia* Moench. *Conyza halimifolia* Desf. — Plak. t. 27. f. 2. Hérn. parad. t. 225. (v. v. c. et s. sp.)

É. *angustior*, foliis cuneato-lanceolatis apice 3-5-dentatis, summis lanceolatis integerrimis. 5 in insulâ Cubâ circa Havanam. (v. s. comm. à cl. Ramon de la Sagra.)

98. *B. AERUSA*, fruticosa glabra apice viscosa, ramis teretibus, ramulis subangulatis, foliis obovatis subcuneatis in petiolum attenuatis obtusissimis dentibus obtusis utrinquè 7-8 subrepandis basi trinerviis, capitula ♂ ad axillas foliorum super. oblongorum solitariis pedicellatis folio brevioribus racemos foliaceos constituentibus, invol. ovati squamis lanceolatis acutis. 5 in Brasiliâ prov. Sancti-Pauli. Rami flor. laterales breves. Folia caulina cum petiolo 4-5 lin. 2 poll. longa, 9-10 lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 479 miss.)
99. *B. ILLINITA*, fruticosa glabra viscosa, ramis apice striatis, foliis breviter petiolatis latè obovato-cuneatis obtusissimis dentibus 7-9 distantibus grossè notatis suprâ asperè glutine viscoso illinitis et lucidis trinerviis, nervis lateralibus ferè æquidistantibus inter nervum medium et margines, racemo subpaniculato terminali, bracteis linearis-filiformibus pedicello brevioribus, invol. ♀ squamis lanceolato-linearibus acutis, achæniis striatis glabris. 5 in Brasiliâ prov. Sancti-Pauli. Affinis *B. mesoneura* sed bractearum multò angustiorum et breviores, etc. Achænia parva. Pappus rufus invol. paulò longior 1-serialis (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 502 miss.)
100. *B. MESONEURA*, fruticosa ramosa viscosa, ramulis angulatis, foliis obovato-cuneatis in petiolum attenuatis à medio ad apicem grossè dentatis trinerviis, nervis lateralibus æquidistantibus à nervo longitudinali et margine, capitula racemoso-subpaniculatis, foliis floralibus oblongis integris pedicello longioribus, invol. ♀ squamis lanceolatis subacutis, achænio striato ovato glabro. 5 in Brasiliâ prov. Sancti-Pauli. Folia 20 lin. longa, 10 lin. lata. Pappus rufus invol. paulò longior. Fl. masc. ign. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 501 miss.)
101. *B. CASSINOIDES*, fruticosa glabra, ramis teretibus striatis junioribus folisque supernè viscosis, foliis in petiolum attenuatis obovatis obtusis mucronatis integerrimis coriaceis penninerviis subtus minutissimè punctulatis suprâ nitidis, corymbis terminalibus foliis intermixtis, invol. ♂ ovati squamis linearibus acutiusculis. 5 in Brasiliâ prov. Sancti-Pauli. Valdè similis *B. cassinefolia*. Ind. sem. ign. (v. s. ♂ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 477 miss.)
102. *B. CASSINEFOLIA*, fruticosa glabra, ramis teretibus, ramulis striatis, foliis in petiolum attenuatis obovatis obtusis mucronatis integerrimis coriaceis penninerviis subtus minutissimè punctulatis, corymbis terminalibus foliis intermixtis, pedicellis bracteolatis, invol. ♂ ovati, ♀ oblongi, squamis ovatis oblongis, ext. subacutis inter. longioribus obtusis, achæniis glabris. 5 in Brasiliâ circa Rio-Janeiro (Lund!), Bahia (Blanch.). Pappus ♂ invol. cor. æqualis. ♀ duplò longior albus pluriserialis. Folia 1  $\frac{1}{2}$ -2 poll. longa, 6-10 lin. lata (v. s.)
103. *B. VISMIOIDES*, fruticosa, ramis teretiusculis junioribus pedunculisque retrahentibus, foliis in petiolum brevem attenuatis obovatis basi cuneatis integerrimis ciliatis subpenninerviis brevissimè mucronatis, pauculâ terminali foliolis parvis ovali-oblongis instructâ polycephalâ, invol. ♂ squamis ovalibus obtusis glabris margine submembranaceis pappum subæquantibus. 5 in Brasiliâ prov.

- Sancti-Pauli. Fl. fem. ign. Capitula ♂ parva. Folia rufescentia 15-16 lin. longa, 6 lin. lata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 478 miss.)
104. *B. CALVASCHENS*, fruticosa, ramis teretibus junioribus dense villosis adultis glabris, foliis in petiolum attenuatis elliptico-oblongis submucronatis integerrimis uninerviis utrinque junioribus sericeo-villosis demum supra glabris subtus canescenti-velutinis subtuberculatis, racemis ramulos terminantibus simplicibus, invol. squamis oblongis acutis subrufo-membranaceis, acheniis striatis glabris. ♂ in Brasiliæ prov. Minas-Geraes ad Mariannam legit cl. Vauthier (pl. em. n. 33a) et circa Bahiam cl. Blanchet (n. 1972 et 1973!). Pappus rufus x-serialis deciduus Ind. ♂ ign. (v. s. ♂ et ♀.)
105. *B. PAUCIFLOSCULOSA*, fruticosa glabra viscosa ramosa, ramulis decurrenti-angulatis, foliis obovatis basi cuneatis sessilibus apice obtusis obtusè dentatis tenuiter triplinerviis, capitulis ad axillas super. sessilibus solitariis, summis confertis, in ♂ magis glomeratis, in ♀ magis spicatis, invol. 5-8-flori squamis oblongo-lanceolatis, ♂ acutis, in ♀ acuminatis, acheniis glabris striatis. ♂ in Brasiliæ prov. Sancti-Pauli et Miner. gener. Folia 4-6 lin. longa, 3 circ. lata, subtus vix ac ne vix subpunctata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 480 et 486 miss.)
- ♂. *punctulata*, foliis subtus punctulatis. ♂ in prov. Rio-Grande. Indiv. masc. ignota. An fortè species propria? (v. s. cum var. α sub n. 959.)
106. *B. LEPTOCERPHALA*, fruticosa glabra, ramulis decurrenti-angulatis, foliis oblongo-linearibus seu obovato-oblongis integerrimis seu paucidentatis, obtusis basi attenuatis uninerviis coriaceis, capitulis axillaribus sessilibus ad apices ramorum et ramulorum spicato-congestis oblongis, ♂ 8-floris, ♀ 4-floris, invol. squamis oblongo-lanceolatis acutis membranaceis, acheniis glabris, pappo etiam ♀ involucrium vix superante. ♂ in Brasiliæ prov. Rio-Grande (herb. imp. Bras. n. 950, 977 et 948) et Rio-Janeiro (Blanch!). Folia 10-20 lin. longa, 2-4 lata. (v. s.)
107. *B. COGNATA*, fruticosa glabra viscosa ramosa, ramulis decurrenti-angulatis, foliis obovatis basi cuneatis sessilibus apice obtusis obtusè dentatis tenuiter triplinerviis, capitulis ad axillas super. sessilibus congestis subterminalibus, invol. ovatis squamis utriusque sexus ovato-lanceolatis subacutis, floribus in invol. 14-18, acheniis striatis glabris. ♂ in Brasiliæ prov. Rio-Grande. Nimis *B. pauciflosculosa* affinis, sed ex numero florum et squamarum ferè duplo videtur diversa. Folia in specim. cæt. simillimis subpunctata aut epunctata. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 985, 946 et 975 miss.)
108. *B. PLATENSIS* (Spreng. syst. 3. p. 465), fruticosa glabra viscosa, ramis oppositis junioribus striato-angulatis, foliis oppositis oblongo-cuneatis apice dentatis coriaceis, infer. obovato-oblongis, mediis oblongis, summis linearibus subintegerrimis, capitulis sessilibus glomeratis secus ramos et ramulos interruptè spicatis, invol. ovatis squamis ovalibus subacutis pappi longit. subæquantibus. ♂ in Brasiliâ merid. ad Montevideo (Spr.), in littore dicto Bandâ orientale (Racle!), ad Rio-Grande (h. Mus. Bras.). Folia infer. 2 poll. longa, 6 lin. lata. Variat foliis latioribus trinerviis aut angustioribus uninerviis. An duæ species hic fortè confusæ? (v. s.)
109. *B. SUBOPPOSITA*, fruticosa glabra subviscosa, ramulis striato-angulatis foliisque suboppositis cuneatis apice obtusè 5-7 dentatis basi integerrimis subpetiolatis trinerviis, nervis lateralibus margini parallelis, capitulis ad apices ramorum paniculato-congestis breviter pedicellatis, invol. ♀ ovatis squamis ext. ovatis int. oblongis acutis pappo æqualibus. ♂ in Brasiliæ prov. Rio-Grande. Habitus nonnull. Eupatorii specierum ob folia sæpius opposita aut subopposita rarius alterna, sed flores Baccharidis. Folia 10-12 lin. longa, 5-6 lata. Capitula parva. (v. s. ♀ in h. Mus. Par. à Mus. imp. Bras. sub n. 978 et verisim. 949 miss.)
110. *B. AFFINIS*, fruticosa ramosa glabra, ramulis striato-angulatis, foliis cuneatis obtusis basi breviter petiolatis integris apice obtusè 5-7-dentatis basi triner-



viis nervis lateralibus margini parallelis medio cæt. penninervio, summis obovato-oblongis integris, capitulis racemosis subpaniculatis pedicellatis ovatis, squamis utriusque sexus ovato-oblongis subacutis, acheniis striatis glabris, pappo invol. superante. § in Brasiliæ prov. Sancti-Pauli et Rio-Grande. Affinis B. suboppositæ sed folia de generis more alterna obtusiora, capitula duplè majora, etc. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. ♂ sub n. 487 et 391. ♀ sub n. 969 miss.)

111. B. CRENULATA (Spreng. syst. 3. p. 465), fruticosa, foliis oppositis alternisque longè petiolatis oblongis obtusiusculis basi attenuatis crenulatis coccineis reticulato-venosis 3-nerviis glabris, capitulis interruptè spicatis subsessilibus, spicis paniculatis. § in Brasiliâ. †.

112. B. HOOKERIANA, fruticosa, foliis fasciculatis cuneatis basi attenuatis sessilibus dentato-spinosis rigidis, capitulis solitariis ramulos versus apicem cæcis foliosos breves terminantibus et ità quasi spicato-racemosis. § ad Coquimb. Chilensium. Baccharis macronota Hook. et Arn. bot. Beech. p. 30, non Kunth Cæt. ign.

§. 3. Discolores, nempe foliis subcuneatis obovatis oblongis linearibusve uninerviis rarius penninerviis suprâ virescentibus subtus densè cano-tomentosis aut villosis, ramis apteris.

113. B. AVICENNIFOLIA, fruticosa, ramis striatis glabris junioribus antrato-angularis, foliis petiolatis obovato-oblongis obtusis basi attenuatis integerrimis penninerviis suprâ glabris subtus cano-tomentosis, racemis strictis subaphyllis, capitulis breviter pedicellatis ovatis, invol. squamis ovatis acutis. § in Brasiliæ prov. Minarum Generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 266). Conyza racemosa Spreng. syst. 3. p. 508? Petioli 6 lin. Limbus fol. 4 poll. longus, 15-16 lin. latus. Capitula immatura mascula non satis nota, sed certe Baccharidis potius quàm Conyze. (v. s.)

114. B. LESSINGIANA, fruticosa, ramis teretibus, foliis petiolatis ellipticis basi obtusis apice acutis integerrimis suprâ junioribus araneosis adultis glaberrimis subtus cum petiolis ramis pedunculisque densè cano-hirsutis, paniculâ terminali, invol. ♂ squamis ovato-lanceolatis ciliatis, ext. dorso subhirsutis, int. sublinearibus pappo brevioribus. § in Brasiliâ. Bacch. nov. sp. Lem.! mas. in Lund. pl. miss. Flores ♂ in invol. circ. 15, majusculi. Achenium immat. pilosiusculum. Pappus antheris rufescenti-subflavicans. Petioli semipoll. Limbus fol. 3-3 ½ poll. longus, 15 lin. latus. Fl. ♀ ign. (v. s. comm. à cl. Lund.)

115. B. TARCHONANTHOIDES, fruticosa, ramis teretibus pedunculisque molliter tomentosis, foliis subpetiolatis lanceolatis obtusis basi longè cuneatis integris à medio ad apicem grossè serratis coriaceis suprâ glaberrimis subtus tomentosis penninerviis reticulatis, paniculâ terminali hirsutâ ramosisssimâ polycephalâ subaphyllâ, invol. ♂ campanulati squamis hirtellis oblongo-linearibus acutis § seu § in Brasiliæ prov. Minarum General. legit cl. Vauthier (pl. exs. n. 175). Folia 2 ½-3 poll. longa, 8-10 lin. lata. Cor. ♂ et pappus crassiusculis invol. æqualis. Ind. fœm. ign. (v. s. comm. à cl. inv.)

116. B. GRANDIFLORA (H. B. et Kunth nov. gen. am. 4. p. 58), fruticosa robusta, ramulis teretibus pubescenti-tomentosis, foliis petiolatis oblongis acuminatis basi acutis argutè dentatis subrugosis suprâ glabris subtus fuscotomentosis, corymbis terminalibus dichotomis, invol. ♀ subhemisphæricis squamis lineari-lanceolatis acuminatis subfimbriatis, achenio sulcato scaberrimis § in planitie Quitensi juxta Mulato. Folia 3 poll. longa, 10-11 lin. lata. Fl. ♂ ign.

117. B. MOLLIS (H. B. et Kunth nov. gen. am. 4. p. 58), fruticosa, ramis teretibus fusciscenti-villosis, foliis subsessilibus ovatis obtusis basi rotundatis crenatis suprâ pilosiusculis subtus molliter villosis-pubescentibus, dentibus mucronatis, capitulis in apicis axillares terminalesque digestis sessilibus conglomeratis.

invol. ♂ hemisphaerici squamis lanceolatis acutis. ♀ circa urbem Quito. Folia sesquipollicaria. Pl. ♀ ign.

118. *B. UNCINELLA*, fruticosa ramosissima, ramis teretibus glabris junioribus hirsutis, foliis sessilibus confertis patulis coriaceis integerrimis obovato-oblongis supra glaberrimis subtus cano-tomentosis junioribus apice mucrone recurvo uncinulatis adultis obtusis, capitulis ♂ paucis ad apices ramulorum axillaribus subcongestis, invol. ovati squamis exter. dorso subhirsutis, inter. linearibus glabriusculis. ♀ in Brasiliæ prov. Sancti-Pauli. Rami conferti corymposi breves. Folia 3 lin. longa, 1  $\frac{1}{2}$  lata. Flor. form. ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 470 miss.)
119. *B. GNAPHALIOIDES* (Spreng. syst. 3. p. 461), fruticosa, ramis teretibus albo-tomentosis, foliis linearibus utrinque obtusis patulis reflexive uninerviis integerrimis margine subrevolutis supra araneosis subtus albo-tomentosis, capitulis ♀ racemoso-capitatis breviter pedicellatis, invol. squamis ovatis subacutis, acheniis glabris muricatis ♀ in Brasiliæ prov. Rio-Grande. Folia 5 lin. longa, lineam lata. Flor. in invol. ♀ circ. 20. Pappus rufus plariserialis. Species affinis *B. velutina* sed in genere suo distinctissima. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 824 miss.)
120. *B. HELICHRYSOIDES*, fruticosa, ramis teretibus lanato-hirsutis, foliis sessilibus basi truncato-subcordatis lanceolatis acuminatis, summis ovato-lanceolatis, omnibus integerrimis planis supra pubescentibus subtus deusè albo-tomentosis, capitulis laxè corymbosis pedicellatis, invol. campanulati squamis oblongo-lanceolatis hirsutis, in ♀ longioribus acuminatis, achenio subangulato glabro. ♀ in Brasiliæ prov. Rio-Grande et Sancti-Pauli. Pappus albus nitens. Folia 12-15 lin. longa, 3-4 lata. Spec. speciosa distinctissima. (v. s. ♂ et ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 811 et 492 miss.)
121. *B. VELUTINA*, fruticosa ramosa tota molliter velutina, ramulis teretibus, foliis sessilibus linearibus integerrimis 1-nerviis margine subrevolutis, capitulis ♂ ex axillis breviter pedicellatis racemosis pedicellis nudis, invol. squamis oblongis ♀ ex axillis longius pedicellatis, pedicellis bracteolatis, invol. squamis ovatis obtusissimis laxis rufo-velutinis floribus in ind. ♀ 5-6, acheniis glabris, corollâ rigida pappo plariseriali. ♀ in Brasiliæ prov. Rio-Grande. An fortè *B. ochracea* Spreng. syst. 3. p. 460? Habitus ferè *Ericæ*. Folia 3-6 lin. longa, in ♂ acutâ, in ♀ obtusâ. Non satia constat an ind. ♂ et ♀ hic descripta ad unam eandemque speciem pertinet. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. ♂ sub n. 995, ♀ sub 819 miss.)
122. *B. ENIOCLADA*, fruticosa, ramis teretibus præsertim junioribus villosio-hirsutis, foliis sessilibus ovatis basi obtusis apice mucrone acutis integerrimis aut hinc inde 1-2 dentatis coriaceis supra glabris subtus villosis subhirsutis, capitulis ♀ sessilibus axillaribus ad apices ramorum et ramulorum spicato-congestis, invol. squamis lanceolatis rufo-membranaceis glabris, achenio striato glabro. ♀ in Brasiliæ prov. Sancti-Pauli. Folia 6-7 lin. longa, 3 lin. lata. Pappus rufus invol. paulò longior. Fl. masc. ign. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 469 miss.)
123. *B. PHYLLICIFOLIA*, fruticosa, ramis teretibus junioribus canescentibus velutino-hirsutis, foliis sessilibus approximato-patentibus ovato-oblongis basi obtusis subcordatis, apice obtusiusculis submucronulatis margine vix anbrevolutis supra glabris subtus cano-tomentosis, paniculâ subnudâ ramosissimâ apice subcorymbosâ, capitulis pedicellatis, invol. ♂ squamis oblongo-linearibus obtusis dorso lanato-hirsutis. ♀ in Brasiliæ prov. Sancti-Pauli. Accedit ad descr. *B. phyllicoidis*. Folia uninervia 6 lin. longa, 2  $\frac{1}{2}$  lata. Pappus ♂ crispus albidus vix invol. longior. Fl. ♀ ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 491 miss.)
124. *B. LUCOPAPPA*, caule fruticoso ramoso juniore canescente demùm glabrato, ramis teretibus, foliis sessilibus alternis ovatis acuminatis integerrimis supra glabris subtus cano-tomentosis, marginibus subrevolutis, ramis floridis subpaniculatis capitula corymboso-conferta gerentibus, invol. squamis oblongo-

linearibus acutis velutinis sublanatis.  $\frac{3}{4}$  in prov. Rio-Grande Brazil. (h. Mus. imp. Bras. n. 330). Folia coriacea 6-8 lin. longa, 3-4 lata, adolta reflexa. Pappus albusissimus. (v. s. in h. Mus. reg. Par.)

125. *B. PHYLLICORDEA* (H. B. et Kunth nov. gen. am. 4 p. 49), fruticosa ramo aissima, ramulis villosis-pubescentibus, foliis sessilibus approximato-patentibus ovato-oblongis obtusiusculis margine revolutis integerrimis coriaceis enerviâ. suprà nitidis glabris, subtùs villosis-pubescentibus, capitulis  $\sigma$  terminalibus capitato-conglomeratis sessilibus, invol. hemisphærici squamis obtusiusculis  $\frac{3}{4}$  in Peruvia devezis Parami de Guarami. Cor. 13-14 alba,  $\varnothing$  ign.

126. *B. RADICANS*, caulibus suffrutescentibus teretibus virentibus adscendentibus albidis-subtomentosis basi radicanibus, foliis sessilibus linearibus patulis integerrimis acutis margine revolutis suprà glabris subtùs albidis-tomentosis adultis subglabris, corymbo, terminali subcomposito conferto 8-12-cephalo, invol.  $\varnothing$  squamis ovatis exter. obtusiusculis dorso villosis, inter. mucronato-acutis  $\frac{3}{4}$  in Brazilia prov. Rio-Grande. Species distinctissima. Folia 6-8 lin. longa, 1-1  $\frac{1}{2}$  lata, nervo medio suprà exarato. (v. s. in h. Mus. reg. Par. & Mus. imp. Bras. sub n. 906 miss.)

$\S$  4. Oblongifoliae, nempe foliis oblongis ovalibus linearibusve glabris aut rariùs villosis uni- aut penninerviis non distichè imbricatis, ramis apteris.

127. *B. CAPRARIAEFOLIA*, fruticosa, ramis teretibus junioribus laxè et molliùs villosis-hirsutis, foliis sessilibus ovalibus aut obovato-oblongis apice serratis utrinquè 3-5 acutis notatis junioribus laxè villosis adultis glabrisculis, capitulis axillaribus sessilibus ad apices ramorum in spicam foliaceam subconnum digestis, invol. squamis lanceolatis margine membranaceis subacutis, achenio glabro.  $\frac{3}{4}$  in Brazilia prov. Rio-Grande (h. Mus. imp. Bras. n. 966 et 968). Capitula masc. parva, form. duplò ferè majora. Pappus  $\varnothing$  rufus invol. paulò longior. Folia 7-10 lin. longa, 3-lin. lata. (v. s.)

128. *B. ORYZALIS*, fruticosa glabrescens, ramulis subangulatis adultis glabris, foliis petiolatis elliptico-lanceolatis utrinquè acuminatis penninerviis subis pallidioribus et punctulatis à medio ad apicem remotè denticulatis, corymbis axillaribus 5-7-cephalis in paniculam foliosam dispositis, capitulis  $\varnothing$  pedicellatis, invol. squamis int. elongatis acuminatis, acheniis glabris.  $\frac{3}{4}$  in Brazilia Serra dos Orgaos legit cl. Lhostky septemb. flor. Fl. masc. ign. Pappus  $\varnothing$  albidus elongatus. Folia 3 poll. longa, pollicem lata. (v. s.)

129. *B. MICRODONTA*, fruticosa glabra apice viscosa, ramulis decurrenti-angulatis, foliis oblongo-lanceolatis in petiolum subattenuatis acutis minutè et serratis ad apicem serrulatis uninerviis aut subtrinerviis venis lateralibus tenuissimis, capitulis  $\sigma$  ad axillas foliorum secus ramulos axillares solitariis sessilibus ovato-oblongis, invol. squamis lanceolatis.  $\frac{3}{4}$  in Brazilia prov. Sancti-Pauli. Folia 15 lin. longa, 3 lata. Capitula pallida 2 lin. longa. Form. ign. (v. s. in h. Mus. reg. Par. & Mus. imp. Bras. sub n. 483 miss.)

130. *B. MACRODONTA*, fruticosa glabra viscosa, ramulis subangulatis, foliis petiolatis oblongis ovalibusve utrinquè acuminatis hinc inde à medio ad apicem grossè serratis penninerviis, venis inf. submajoribus et inde falsè triplinerviis, pedunculis axillaribus folio brevioribus capitula 5-7 pedicellata corymbosa virentibus, invol.  $\varnothing$  squamis ovali-lanceolatis, int. linearibus, margine membranaceis, acheniis striatis glabris.  $\frac{3}{4}$  in Brazilia prov. Sancti-Pauli. Vidi accedit ad *B. microdonta* sed diversa satis videtur. Flor. masc. ign. (v. s. in h. Mus. reg. Par. & Mus. imp. Bras. sub n. 485 miss.)

131. *B. PENTLANDII*, fruticosa glabra viscosa, ramulis sulcatis, foliis petiolatis lanceolatis utrinquè acutis immò basi integris cæterum argutè serratis penninerviis reticulato-venosis, capitulis  $\sigma$  ad apices ramorum densè congestis inter folia suprema sessilibus, invol. hemisphærici squamis lanceolatis scariosis acutis subimbricatis.  $\frac{3}{4}$  in Republ. Boliviana Amer. austr. legit cl. Pentland et m-

n. comm. Folia cum petiolo 4 lin. bipollicaria, 6-8 lin. lata. Flor. ♀ ign. s.)

**B. SUDDETICORDES** (H. B. et Kunth nov. gen. am. 4. p. 64), fruticosa glabra, ramulis subangulatis, foliis petiolatis lanceolatis acuminatis basi angustatis reticulatis subcoriaceis reticulato-venosis, corymbis terminalibus diffusis, capitulis ♂ pedicellatis 70-80-floris, invol. hemisphaerici squamis lanceolatis subcoriaceis. ♀ in Peruvia? Folia 3 ½ poll. longa, 8 lin. lata, petiolo 1 lin. Fl. ♀ ign.

**B. BOGOTENSIS** (H. B. et Kunth nov. gen. am. 4. p. 61), fruticosa glabra, ramulis virgatis sulcatis, foliis breviter petiolatis lanceolato-oblongis acutis serrulatis coriaceis reticulato-venosis supra nitidis, paniculis ♂ terminibus ramosis foliatis, invol. subhemisphaerici squamis lanceolatis acutis sciliatis. ♀ in altâ planitie Bogotensi juxta Santa-Fé, et inter Chuapa et po. Folia 7-8 lin. longa, 2-2 ½ lata. Fl. ♀ ign.

**B. VOLUMINIS** (H. B. et Kunth nov. gen. am. 4. p. 57), fruticosa glabra voluminosa, ramis striatis, foliis petiolatis ovato-oblongis breviter acuminatis basi rotundatis argutè serrulatis rigidis nitidis, pedunculis axillaribus terminibus oligocephalis, invol. ♀ subhemisphaerici squamis lanceolatis linearibus, achenio costato glabro. ♀ in frigidissimis Andium Peruv. adices Parani de Guarano. Cor. 25-30 albæ, ♂ ign. Folia sesquipollic., 8 lin.

**B. ASCENDENS** (Pers. ench. 2. p. 424. n. 13), caule adscendente, foliis acuminatis profundè dentato-serratis. ♀? in Peruvia ubi dicitur *Chilco* herb. Juss. Cum B. volubili conjunxit Spreng. syst. 3. p. 463. sed distincti servo quia cl. Kunth qui herb. Juss. vidit cum eâ non imò confert. Cæter. †.

**B. ARBUTIFOLIA** (H. B. et Kunth nov. gen. am. 4. p. 54), fruticosa glabra, ramulis angulatis, foliis sessilibus oblongis utrinquè acutis argutè serratis coracis nitidis, capitulis terminalibus corymboso-conglomeratis breviter pedicellatis, invol. squamis ovato-lanceolatis. ♀ in Peruvia (J. Juss.), in deis montis Pichincha Quitensium (H. B.). B. arbutifolia Vahl symb. 3. p. 97? Conyza arbutifolia Lam. dict. 2. (1786) p. 92.

**B. TEINDALENSIS** (H. B. et Kunth nov. gen. am. 4. p. 54), fruticosa ramosissima glabra apice viscosa, ramulis angulatis, foliis breviter petiolatis elliptico-oblongis utrinquè acutis grossè dentatis subcoriaceis, capitulis terminalibus glomeratis breviter pedunculatis, invol. hemisphaerici squamis scariosis tussculis. ♀ in altissimis inter Teindala et Pasto Novo-Granatensium. Cor. e in utroque sexu 30-40. Folia 14-15 lin. longa, 7-8 lata.

**B. MACRANTHA** (H. B. et Kunth nov. gen. am. 4. p. 54), fruticosa ramosissima glabra viscosa, ramis teretibus, foliis brevissimè petiolatis oblongis argutè acutis remotè et obsoletè dentatis subcoriaceis, capitulis ♀ terminibus capitato-congestis breviter pedunculatis, invol. squamis scariosis aculeis. ♀ propè Mulado Quitensium ad radices montis Cotopaxi. Capitula flores circ. 75 invol. longiores gerentia. Folia 12-14 lin. longa, 4-5 lata.

**B. CAPITATA** (H. B. et Kunth nov. gen. am. 4. p. 53), fruticosa ramosissima glabra, ramulis subangulatis viscosis, foliis petiolatis subelliptico-obtusculis basi acutis sinuato-dentatis subcoriaceis glabris viscosis, capitulis terminalibus capitato-congestis sessilibus, invol. subhemisphaerici squamis obtusiusculis, achenio sulcato glabro. ♀ in America meridionali. 10-11 lin. longa. Fl. ♂ ign.

**B. ELEGANS** (H. B. et Kunth nov. gen. am. p. 60. t. 324), fruticosa glabra, ramulis sulcato-angulatis, foliis petiolatis ellipticis acutis basi rotundatis apicè versis argutè dentatis supra nitidis reticulato-venosis, corymbis terminibus ramosissimis divaricatis foliatis, capitulis ♀ ovali-oblongis, invol. squamis lanceolatis acutis, acheniis glabris. ♀ in Mexico? Pedicelli parè bractæolati. r. masc. ign. Affinis videtur B. petiolar.

141. *B. MICRANTHA* (H. B. et Kunth nov. gen. am. 4. p. 61), herbacea? caule tereti striato hirtio, foliis petiolatis oblongis acutis basi subrotundatis argute dentatis subcoriaceis dense hirtis, paniculâ terminali ramosissimâ foliatâ patulâ, invol. ♂ hemisphaerici squamis lanceolatis acutis subciliatis. — Prop. Guanaxuato Mexicanorum. Cor. alba. Fl. form. ign.
142. *B.?* *BRACTEOLATA*, suffruticosa subviscosa, ramis junioribus subhirsutis, foliis linearibus obtusiusculis basi attenuatis confertis punctulatis puberulo-glandulosis integerrimis aut sæpius denticulatis, paniculâ brevi, squamis acutis secus ramulos bracteolatâ, invol. squamis linearibus acuminatis puberulo-ciliatis. ♂ patria ign. Prov. juxta Poir. ex itia. Commers. Specimen habeo olim sub nom. *Chrysocome* in hort. Mousseaux lectum. — *Baccharis* Lax ill. t. 698. f. 2. *Bacch. parviflora* Poir. suppl. t. p. 556. non Pers. Capit. specim. mei ut Poirerii casa et ideo genus dubium. (v. s.)
143. *B. MUCRONATA* (H. B. et Kunth nov. gen. am. 4. p. 60 non Hook.), herbacea?, caule tereti densissimè hispido-hirto, foliis petiolatis oblongis utrinque acutis mucronatis remotè et argutè denticulatis rigidis hirtis, paniculâ terminali ramosâ patulâ bracteâtâ, invol. ♂ squamis lanceolatis acutis ciliatis — in Mexici montibus Guanaxuatensibus. Fl. form. ign.
144. *B. SERRATIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 59), fruticosa, ramulis subangulatis hirto-pubescentibus, foliis brevissimè petiolatis ovato-ellipticè brevissimè acuminatis basi angustatis argutè serratis rigidis suprà hispida pilosis subtus molliter pilosis, paniculis terminalibus ramosis patulis bracteatis, invol. ♂ squamis lanceolatis acutis apice ciliatis. ♂ in montibus Mexicanis propè Guanaxuato (H. B.). An etiam in Martinicâ ut vult Spreng. Syst. 3. p. 463? Bractearum lanceolarum hispida. Fl. form. ign.
145. *B. HIRTELLA*, fruticosa ramosa, ramis teretibus junioribus pubescentibus hirtellis, foliis breviter petiolatis ellipticis acuminatis penninerviis membranaceis suprà sparse setulosis subtus secus nervos hispidulis margine remote callosè serratis, paniculis ramulos terminantibus, capitulis ♂ parvis, inv. squamis lanceolato-linearibus subciliatis vix acutis. ♂ inter Mexicanas h. Hænkeani vidi. Fl. form. ign. Fortè affinis *B. serratifolia* H. B. K. ? sed nec rigida nec brevissimè acuminata nec subtus molliter pilosa. (v. s. in Hænke à cl. de Sternberg miss.)
146. *B. SEMILIFOLIA*, fruticosa, ramis striatis sæpius divaricatis, foliis sessilibus rigidis ovatis acutis basi obtusis triplinerviis reticulato-venosis serratis suprà glabris, paniculâ pyramidato-corymbosâ, invol. campanulati squamis marginè scariosis glabris in ♂ exter. obtusis, in ♀ exter. ellipticis in utroque inter. linearibus, achæniis puberulis. ♂ in Chili.  
*β. hebeclada*, ramis foliisque subtus pubescenti-hirtellis. ♂ ad Tagaen (Cham.!). *M. sessilifolia* Less. ! in linnæâ 1831. p. 140. *Baccharis* n. Papp. l coll. chil. n. 209. (v. s. ♂.)
147. *B. PETIOLATA*, glabra herbacea erecta ramosa, ramis teretibus apice nigro-glandulosis, foliis petiolatis ellipticis grossè dentatis corymbis pedunculatis compositis, inv. ♂ campanulati squamis ovato lanceolatis subhirsutis — in Chili ex h. Hænke. Species distinctissima. Petioli sequenti aut binae Limbus folii 10 lin. longus, 4-5 latus. Fl. form. ign. (v. s. in h. Hænke ad cl. Sternberg comm.)
148. *B. ERIGEROIDES*, suffruticosa erecta glabra, ramis ad collum ortis subtus pilosisculis, foliis sessilibus lineari-sublanceolatis integerrimis sinuato-caulinis glabris, ramis pilosisculis, capitulis in racemum laxum subpaniculatum dispositis, invol. fl. ♀ squamis ovali-oblongis glabris subobtusis pappo dimidio brevioribus, achæniis pubescentibus et globulis glandulosis ovatis. ♂ in campis editis prov. Sancti-Pauli legit cl. Lund. Pappus rigidulus brunescens. Cor. ♀ teretes quasi truncatæ minutissimè 4-dentatæ. Stylus breves parum exserti. Flor. ♂ ign. et ideo genus dubium. (v. s. comm. a cl. Lund.)

149. *B. SEMISERRATA*, fruticosa erecta ramosa glabra, ramis junioribus angulatis adultis glabris, foliis sessilibus lineari-lanceolatis acutis uninerviis inferne integerrimis superne argutè serratis, capitulis 3-5 corymbosis pedicellatis, invol. ♂ hemisphaerici squamis lanceolatis acuminatis margine eroso-scariosis, fl. circ. 30. ♀ in Peruviae Andium vallibus legit cl. Hænke. Folia 1-1½ poll. longa, 2 lin. lata. (v. s. comm. in h. Hænke à cl. de Sternberg miss.)
150. *B. PTARMICIFOLIA*, suffruticosa glabra, ramis striato-sulcatis, foliis subsessilibus lineari-lanceolatis utrinque acuminatis argutè serratis subtrinerviis subtus punctatis, paniculâ terminali laxâ polycephalâ foliosâ, invol. ♀ squamis oblongo-linearibus subobtusis margine membranaceis ciliatis. ♀ in Panamaide (Née! ex h. Thib.). In hortis Mexicanis legit. cl. Berlandier julio flor. (pl. exs. n. 572). Valdè affinis *B. thesioidi* sed folia 2 poll. ferè longa, 3 lin. lata, subtus punctata basi subtrinervia. (v. s.)
151. *B. SULCATA*, fruticulosa ramosissima gracilis glabra, ramulis sulcato-angulatis, foliis sessilibus linearibus acuminatis à medio ad apicem serratis utrinque impresso-punctatis, paniculâ terminali laxâ foliis intermixtâ, capitulis ♂ ad apices pedicellorum solitariis, invol. squamis lanceolatis acutis margine scariosis. ♀ in Mexico circa Villalpando legit cl. Mendèz. Species inter eas quas vidi distinctissima, fortè *B. thesioidi* affinis. Fl. form. ign. (v. s. ♂ comm. à cl. Alaman.)
152. *B. THESIODES* (H. B. et Kunth nov. gen. am. p. 61), fruticulosa glabra, ramulis sulcato-angulatis, foliis brevissimè petiolatis lineari-lanceolatis basi acutis apice acuminatis argutè serrulatis 1-nerviis rigidulis, paniculâ terminali ramosâ foliatâ, invol. squamis lanceolato-oblongis subobtusis sinubriato-ciliatis, pappo ♂ setiformi, achæniis glabris. ♀ in Mexico (Mair.!), circa urbem (Berl. pl. exs. n. 529), juxta Santa-Rosa in montanis (H. et B.). Fol. 10-11 lin. longa, 1-2 lata. Affinis *B. multifloræ*. (v. s. ♂ et ♀.)
153. *B. MULTIFLORA* (H. B. et Kunth nov. gen. am. 4. p. 59), suffruticosa, caulibus simplicibus striatis puberulo-hirtis, foliis brevissimè petiolatis oblongis acutis basi rotundatis argutè serratis suprâ glabris subtus præcipuè secus nervum et petiolum puberulo-hirtis, paniculâ terminali ramosissimâ subfoliatâ, invol. utriusque sexus squamis lanceolatis acutis margine scariosis sinubriato-scariosis, pappo ♂ apice penicillato-plumoso. ♀ in Mexico (Mair.!), circa urbem (Berl. pl. exs. n. 900), et ad Tianguillo et Toluca in frigidis montanis (H. B.). Spec. ♀ descr. cl. Kunth et mea sunt omnia ♂. (v. s.)
154. *B. MYRSINITES* (Pers. ench. 2. p. 424. n. 25), fruticosa ramosissima, ramulis angulatis, foliis lanceolatis confertis apice sæpius bidentatis, capitulis terminalibus sessilibus congestis globosis ovatisve, invol. squamis acutis apice subciliatis. ♀ in Santo-Domingo. *Conyza myrsinites* Lam. dict. 2. p. 92. Spreng. syst. 3. p. 465 excl. syn. Poir. Habitus *B. Magellanicæ* et cuneifoliæ ex Lam.
155. *B. CHILCO* (H. B. et Kunth nov. gen. am. 4. p. 56), fruticosa ramosissima glabra viscosa, ramulis angulatis, foliis brevissimè petiolatis lanceolato-oblongis obtusiusculis basi cuneatis apicem versus utroque latere dente 1-2 instructis subcoriaceis subaveniis, capitulis ♀ terminalibus congestis subsessilibus, invol. squamis scariosis acutiusculis. ♀ prope Santa-Anna Novo-Granatenisium, ubi dicitur *Chilco*. Plures aliæ species ex R. et Pav. in Peruvâ nuncupantur *Chilca*. Fol. 7 lin. longa. Fl. ♂ ign.
156. *B. CHILQUILLA*, fruticosa junior pulverulento-puberula dein glabra subviscosa, ramulis teretiusculis, foliis linearibus elongatis sessilibus acutis hinc inde grosse dentato-serratis aut integerrimis subuninerviis, corymbo terminali polycephalo, invol. ♂ ovati squamis scariosis stramineis glaberrimis ovali-oblongis subobtusis. ♀ in dumetis et sylvaticis secus flumina ad Quillota Chilenisium octob. flor. legit cl. Bertero. Vulgò *Chilquilla del Rio*. Folia 3-4 poll. longa sesquilinea lata. Affinis *B. marginali*. (v. s.)
157. *B. ROSMARINIFOLIA* (Hook. et Arn. bot. Beech. 1. p. 30), caule fruticoso ra-

moço dense folioso, foliis linearibus obtusiusculis crasso-coriaceis integerrimis margine subrevolutis ramulisque resinoso-glandulosis, corymbis terminalibus dense oligocephalis, invol. ovalis squamis elliptico-oblongis ext. obtusis, int. subacutis, acheniis glabris.  $\frac{5}{3}$  in collibus fruticetis sylvaticis et arvis ad Valparaíso (Bert.!), Talcahuano (Cham.!), Conceon (Poepp.!), Rancagua (Bert.!), Concepcionem (d'Urv.!), Colchagua (C. Gay!) Chilensium. Vulgò *Romero*, *Romerito*, *Romero de la tierra*. Baccharis lingulata Kunze! in Poepp. coll. 1. n. 212. Less. in Linnæ 1831. p. 147 et 506. An ex nom. vulg. B. linearis R. et Pav. syst. p. 205 potius hic quam sequenti referenda videtur? (v. s.)

*B. subsinuata*, foliis plurimis obtusè sinuato-repandis. — Verisim. in Chl. legit cl. Dombey. Capitula pedicellata umbellata omnino ut in var. s. (v. s.  $\sigma$  ex h. itin. Dombey comm. à Mus. reg. Par.)

158. B. OBLONGIFOLIA (Spreng. syst. 3. p. 460 excl. syn.), fruticosa, foliis oblongis integerrimis, corymbis terminalibus.  $\frac{5}{3}$  in sylvis circa Concepcionem Chilensium et ad Hualpen tractus. Molina oblongifolia R. et Pav. syst. 203. Cart. ign. An fortè var. latifolia et integrifolia B. Pingræ?  $\frac{5}{3}$ .

*B. glabriuscula*, ramis foliisque subtus glabris aut glabriusculis.  $\frac{5}{3}$  in fruticetis collium circa Valparaíso. Baccharis rigida Hook. et Arn. bot. Beech. 1. p. 57. (v. s.  $\sigma$  et  $\frac{5}{3}$ .)

159. B. PINGRÆA, caule suffruticoso ramoso laxè folioso, foliis linearibus aut lineari-lanceolatis integerrimis serratisve acuminatis ramisque punctulatis, corymbis laxis polycephalis, invol. ovato-subglobosi squamis in  $\sigma$  oblongo-lanceolatis mucronatis, in  $\frac{5}{3}$  inter. acuminatis, acheniis glabris.  $\frac{5}{3}$  in Chili (R. et Pav.?), (Cham.!), frequens ad ripas arenosas fluviorum prope Conceon (Poepp.!), San-Yago (Bert.!), Coquimbo (Gaud.!) et Valparaíso (Macr.!), et in California (Cham. ex Less.), in prov. Rio-Grande (h. imp. Bras.!) et ad Buenos-Ayres (Bacle!). Molina linearis var. R. et Pav. syst. p. 205? Hook. et Arn. in Beech. bot. 1. p. 57. Molina linearis Less. in Linnæ 1831. p. 139 et 505.

*a. latifolia*, foliis lineari-lanceolatis serratis rariùs subintegris imò integerrimis. B. linearis Poepp. coll. chil. II. n. 103. Conyza angustifolia Bert. Par. 1828. Baccharis angustifolia Desf. cat. ed. 3. p. 163 non Michx. excl. patriâ. Pingræa angustifolia Cass. dict. 41. p. 58. (v. s.  $\sigma$  ex Poepp. et hort. Par.  $\frac{5}{3}$  ex Don et Cham.)

*B. angustissima*, foliis angustissimè linearibus integerrimis aut rariùs denticulatis. (v. s.  $\sigma$  ex Gaud.  $\frac{5}{3}$  ex Née et Macrae.)

160. B. LINIFOLIA, herbacea? erecta simplex glabra, caule tereti, foliis linearibus elongatis acutis margine sub lente scabro-serrulatis primâ fronte 1-nerviis, sub lente 3-nerviis, venis lateralibus tenuissimis, paniculâ subcorymbosâ confertâ. invol.  $\sigma$  squamis lanceolatis acutis vix apice subciliatis. — in Mexico ad Real del Monte ex h. Hænke. Habitus ferè Bigelowiæ. Flor. masc. ? in invol. 15-16. Pappus rufus invol. longit. æqualis. Fl. sem. ign. (v. s. in h. Hænke à cl. & Sternberg miss.)

161. B. PANICULATA, fruticosa subviscosa glabriuscula, foliis linearibus acuminatis integerrimis subtrinerviis acumine reflexo, paniculâ elongatâ foliosi. invol.  $\frac{5}{3}$  cylindraceis squamis glabris margine scariosis oblongo-lanceolatis. intimis acutis, acheniis glaberrimis.  $\frac{5}{3}$  in Chili circa Valparaíso (Macrae!) & Coquimbo (Gaud.!). Affinis B. Pingræ sed dispositione capitalorum et foliis apice incurvis facile differt. Fl.  $\sigma$  ign. Cor.  $\frac{5}{3}$  truncatæ apice subpuberula. (v. s. comm. ab hon. Soc. hort. Lond. et ab amic. Gaudichand.)

162. B. PAUCIDENTATA, fruticosa ramosissima glabra, ramulis striato-angulatis. foliis sessilibus linearibus aut lineari-oblongis utrinquè acutis uninerviis aut imâ basi subtrinerviis integerrimis aut dense 1-2 utrinquè notatis, capitula  $\frac{5}{3}$  ad axillas superiores subsessilibus et ideo in spicas breves digestis, invol. squamis lanceolatis acutis margine membranaceis, floribus in invol. circ 10. acheniis striatis glabris.  $\frac{5}{3}$  in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras sub n. 823 et 840), et ad Buenos-Ayres? Affinis B. tenuifoliæ. Ind. masc. ign. Folia 9-12 lin. longa, 1-1' lata. (v. s.)

**ACUNCULIFOLIA**, fruticosa ramosa ad apicem subpubescens, ramulis sessilibus linearibus oblongis acutis utrinque punctatis uninerviis, petiolis apice dentibus acutis parvis utrinque 2-3 notatis, capitulis im- per. breviter pedicellatis in racemum foliosum digestis, invol. squamis acutis margine membranaceis, floribus in invol. 25-30, acheniis 5 in Brasiliæ prov. Rio-Grande, ad Sanctam-Catharinam et Rio-Affinis B. paucidentata. Capitula ♂ pauciora. Variat foliis integerrimisve. Folia 9-11 lin. longa, 2-2½ lin. lata. (v. s. ♂ et ♀ comm. à et in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 817 et 819 miss.)

**HALOTES**, fruticosa glabra, ramulis teretibus, foliis sessilibus ellipticis acutis basi integerrimis apice parce et minute serratis uninerviis erectis subimbricatis, capitulis ad apices ramorum et ramulorum breviter-issimis, foliis summis bracteatis, invol. ♂ squamis oblongis ciliatis, ♀ in Brasiliæ prov. Sancti-Pauli. Folia 7 lin. longa, 2-2½ lata. 4-5 lin. longa, ♀ ign. Pappus ♂ rufus invol. paulo longior. (v. s. in reg. Par. à Mus. imp. Bras. sub n. 476 miss.)

**PROSPERMOIDES**, fruticosa glabra, ramis teretibus junioribus decur- culatis, foliis linearibus sessilibus utrinque attenuatis integerrimis aut subdentatis penninerviis venis subreticulatis inter venas subtus, capitulis ♂ axillaribus solitariis breviter pedicellatis folio breviori- ol. squamis oblongis subobtusis pappum æquantibus. 5 in Brasiliæ neti-Pauli. Affinis B. axillari. Capitula in medio ramorum sita; rami flosi nec floriferi. Folia 9-10 lin. longa, sesquilineam lata. Fl. form. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 482 miss.)

**GRANDES** (H. B. et Kunth nov. gen. am. 4. p. 57), fruticosa ramosissima leosa, ramis sulcato-angulatis, foliis petiolatis oblongis acuminatis integerrimis reticulato-venosis coriaceis nitidis, corymbis ♀ termi- nularibusque, invol. hemisphærici squamis scariosis acutis. 5 in altâ Quilensis ad radices montis Ruca-Pichincha. Folia 2-2½ poll. longa, lata. Pl. ♂ ign.

**ACHYLÆNOIDES**, fruticosa glaberrima nitida, ramis teretibus, foliis petiolatis lanceolatis utrinque acuminatis integerrimis penninerviis, glandis subtus pallidioribus, paniculâ subcorymbosâ laxâ ferè aphyllâ isâ, invol. ♂ squamis oblongis acutis brevibus, ♀ squamis linearibus tibus elongatis, acheniis apice sparse puberis. 5 in Brasiliæ monte Cer- magato flor. legit cl. Lund. Species pulchra habita Brachylæna ne- referens. Fol. 4-5 poll. longa, 9-12 lin. lata. Pappus albus 2-seriatis. mm. à cl. Lund.)

**RUFOSA**, fruticosa ramosa erecta glabra, ramis teretibus, foliis in pe- revem attenuatis subcoriaceis lanceolatis utrinque acuminatis inter- blucidis utrinque venuloso-subreticulatis medio 2-serviis, panicula is corymbosa laxâ ramis ramulisque puberulis, invol. ♂ squamis is acutis, flor. in inv. circ. 15-20. 5 in Peraviæ montibus Orohocen- it cl. Hænke. Affinis B. brachylænoidi. Folia 3-4 poll. longa, 8-10 lin. pus rufescens. Fl. ♀ ign. (v. s. in h. Hænke à cl. de Sternberg eorum.)

**EXA** (Pers. ench. 2. p. 425. n. 41), fruticosa, foliis oblongo-lanceolatis mis denticulatisque, capitulis axillaribus corymboso-racemosis. 5 in sylvaticis altis versus Pillao et Panao tractat. Molina nitida R. et Pav. 104. Cst. ign.

**GUERRINA**, fruticosa glabra, ramis teretibus apice angulatis, foliis sublanceolatis sessilibus integerrimis apice mucrone calloso et denti- pe 1-2 minimis instructis uninerviis venosis, paniculâ terminali sub- tycephalâ, capitulis ♂ pedicellatis, invol. squamis paucis lanceolatis argine membranaceis pappum æquantibus. 5 in Brasiliæ prov. Mine- seralinum ad Serra do Frio legit cl. Vanthier (pl. exs. n. 285). Fl. form. is sesquipoll. longa, 3 lin. lata. (v. s.)



171. *B. vatrónius* (H. B. et Kunth nov. gen. am. 4. p. 50), fruticosa ramosissima glabra viscosa, ramulis striatis, foliis petiolatis oblongis acutis basi angustatis integerrimis coriaceis nitidis, pedunculis  $\sigma$  axillaribus 5-7-nerviis, invol. squamis scariosis acutiusculis, acheniis sulcatis glabris.  $\zeta$  in caulis pro. Juss. de Bracamoros, in ripa flum. Amazonum. Folia 2 poll. longa, 2 lin. lata, petiolo 4-5 lin. Cor. 14 inv. 30-35,  $\sigma$  iga.

172. *B. vranoniorum*, fruticosa, ramis teretibus densè hirsuto-hispidis, foliis sessilibus elliptico-oblongis basi obtusis apice subacutis integerrimis pinninerviis supra puberulis subtus villosis-hirsutis, panicula amplis foliis oblongis internodiis hirsutis, capitulis  $\sigma$  pedicellatis, invol. squamis ext. ovatis dorso hirsutis inter. longioribus membranaceis glabris, acheniis pubentibus, pappo 1-coriell.  $\zeta$  in Brasilia prov. Minas General. ad Marianum h. d. Vauthier (pl. exs. p. 265). Habitus forè Vernoniae puberulae. Hirsutioribus. Folia 2 poll. longa, 6-8 lin. lata. Pappus albus. Fl. mase. iga. (v. s.)

173. *B. Oronotaria*, fruticosa erecta glabra, ramis teretibus vix apice angustis, foliis petiolatis ellipticis utrinque acuminatis subintegerrimis subcoriaceis pinninerviis, racemis axillaribus vix petiolo longioribus, capitulis  $\sigma$  pedicellatis 10-12-floris, invol. squamis ovato-oblongis subacutis.  $\zeta$  in Peruvia montis Oronotensibus legit cl. Hanks. Folia cum petiolo 6 lin. 3 poll. longa, 13-15 lin. lata; nervi laterales 8-9; denticuli 1-2 ad apices foliorum cuspi videntur. Flor.  $\sigma$  iga. (v. s. in h. Hanks. ab ill. de Sternberg const.)

174. *B. vaccinoides* (H. B. et Kunth nov. gen. am. 4. p. 50), fruticosa glabra viscosa, ramulis angulatis, foliis sessilibus oblongis utrinque acutis subintegerrimis coriaceis avenis, capitulis  $\sigma$  terminalibus conglomeratis sessilibus, invol. hemisphaericis squamis scariosis ext. ovatis, inter. oblongis hirsutis. acheniis sulcatis glabris.  $\zeta$  in Mexico temperato prope Moram. Folia ad apicem utrinque subidentata, 5-6 lin. longa, 2  $\frac{1}{2}$  lata. Ind. mase. iga.

175. *B. buxifolia* (Pera. ench. 2. p. 424, n. 24), fruticosa glabra, ramis vix angulatis tuberculatis, foliis subsessilibus confertis oblongo-ovatis integerrimis subretusis, capitulis lateralibus axillaribus solitariis sessilibus, invol. squamis ovali-oblongis apice subciliatis.  $\zeta$  in Peruvia ubi ex Jos. Juss. mss. *Alnusquillo* dicitur. Conyza buxifolia Lam. dict. 2. p. 92. Folia 6-7 lin. longa. Pappus albidus.

176. *B. coridifolia*, fruticosa ramosa glabra, ramulis teretibus, foliis sessilibus linearibus mucrone acutis margine ciliis rigidulis brevibus serrulato-sabris, caulibus elongatis, ramis brevioribus, panicula elongata terminali, capitulis pedicellatis 5-8-floris, invol. squamis ovali-oblongis margine membranaceis obtusis, acheniis pubescentibus.  $\zeta$  in Brasilia prov. Rio-Grande et for. Sancti-Pauli. Caulis basi repens. Panicula maris magis ramosa, capitulis parvis numerosioribus, ferè Artemisiae. Inv. sem. duplò majus, squamis magis scariosis. Pappus rufus pluriserialis. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras sub n. 810, 826, 837, 855 et fortè 510 miss.)

$\beta$ . invol. feminei squamis intimis magis oblongis subacutis.  $\zeta$  in prov. Rio-Grande. (v. s. cum var.  $\alpha$  sub n. 994, 793, 858, 838 et 839.)

177. *B. MEGAPOTAMICA* (Spreng. syst. 3. p. 461), fruticosa glabra, ramulis angulatis, foliis linearibus acutis integerrimis margine revolutis supra nitidis uninerviis ciliatis, capitulis ad axillas fol. sup. solitariis sessilibus in spica dispositis, invol. ovatis squamis lanceolatis acutis, acheniis glabris, pappo  $\sigma$  invol. non superante.  $\zeta$  in Brasilia prov. Rio-Grande et Sancti-Pauli. Spica sessilibus capitulis distantibus interrupta nunc subcontinua. (v. s.  $\sigma$  et  $\zeta$  in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 511, 816, 818, 847 miss.)

178. *B. REVOLUTA* (H. B. et Kunth nov. gen. am. 4. p. 50), fruticosa ramosissima ramulis villosis-pubescentibus, foliis brevissimis petiolatis lanceolato-linearibus integerrimis uninerviis margine revolutis subcoriaceis rigidis villosis-pubescentibus, capitulis  $\sigma$  in corymbum terminalem dispositis, inv. hemisphaerici qu

nis acuminatis scariosis. ♀ in Peruvia? Cor. ♂ 50-60 in inv. fœm., ign. Folia 8-9 lin. longa.

79. *B. LEDIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 50), fruticosa, ramulis cano-tomentosis, foliis sessilibus lanceolato-linearibus acutis integerrimis uninerviis suprâ incauo-pubescentibus subtus cano-tomentosis, capitulis ♂ terminalibus congestis sessilibus, invol. squamis scariosis acuminatis. ♀ in Peruvia? Folia 14-15 lin. longa, 2-3 lata. Cor. masculæ albæ 35-40 in inv., fœm. ign.

80. *B. TENUIFOLIA*, fruticosa ramosissima glabra, ramulis substriatis, foliis alternis et suboppositis linearibus obtusis basi subattenuatis sessilibus integerrimis margine (in sicco) subrevolutis, capitulis ad axillas summas sessilibus spicatis folio suo longioribus oblongis 5-floris, invol. squamis lanceolato-linearibus acutis, achæniis glabris. ♀ in Brasiliæ prov. Sancti-Pauli. Valdè accedit ad *B. leptophyllum*, sed foliis apice obtusis et subtilioribus viscositate ramisque infidis et capitulis spicatis differt. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 512 miss.)

81. *B. LEPTOPHYLLA*, fruticosa ramosissima glabra, ramulis tenuibus striato-angulatis, foliis alternis et suboppositis sessilibus linearibus acutiusculis integerrimis 1-nerviis, capitulis ♀ ad apicem pediculorum 1-3 subracemoso-paniculatis, invol. oblongi squamis lanceolato-linearibus inter. elongatis, floribus in inv. 5-6, achæniis striatis glabris. ♀ in Brasiliæ prov. Rio-Grande. Proximè accedit ad *B. angustifoliam* et ad *tenuifoliam*. Folia 9-10 lin. longa, semilineam lata. Inv. 3 lin. longum. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 993 miss.)

82. *B. GRACILIS*, herbacea glabra, caulibus è collo plurimis gracillimis subnudis striato-sulcatis, foliis sessilibus linearibus acutis tristriatis integerrimis remotis, capitulis paucis longè pedicellatis subcorymbosis, invol. squamis 2-3-seriatis lanceolatis acutis margine membranaceis subciliatis, achæniis glabris ferè invol. equalibus, pappo achæniis longitud. paulò superante scabro. ♀ in Brasiliæ prov. Sancti-Pauli. Valdè accedit ad *B. junceam* sed caulis duplò tenuior et valdè sulcato-angulatus, capitula dimidio minora, pappus intense rufus ferè duplò brevior, achænia duplò longiora, etc. (v. s. ♀ in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 516 miss.)

83. *B. JUNCEA* (Desf. ! cat. h. Par. 1829. p. 163 excl. patr.), herbacea glabra subglaucescens, caulibus è collo plurimis teretibus junceis, foliis sessilibus linearibus subnerviis integerrimis distantibus, capitulis terminalibus sessilibus aut paucis corymbosis pedicellatis, invol. hemisphærici squamis lanceolatis rigidis acutis, intimis linearibus minoribus, achæniis glabris. ♀ in Brasiliâ meridionali ex Less. in linnæâ 1831. p. 149, nec in Ægypto ut vult Lehmann, aut in Senegaliâ ut hab. hort. Par. *Stephananthus junceus* Lehm. cat. sem. h. Hamb. 1826. p. 18. ind. schol. 1828. p. 34. pug. pl. nov. p. 35. act. nov. nat. cur. v. 14. p. 2. p. 821 (descr. ind. masc.). *Arrhenachne juncea* Cass. ! dict. 52. p. 253 (descr. fœm. et masc.). Ad *Eupatorineas* ref. Lehmann et ad *Vernoniaceas* Cassini sed certè (ut jam rectè dixit Less.). *Baccharidis* species. Pappus ♀ pluriserialis pallidè rufescens angustè subpaleaceo-pilosus vix ac ne vix serrulatus. (v. v. ♀ cult. in hort. Par.)

84. *B. ANGUSTIFOLIA* (Michx. fl. bor. am. 2. p. 125), fruticosa glaberrima ramosissima, ramis teretiusculis, foliis angusto-linearibus integerrimis vix acutis, paniculâ ♂ laxâ oligocephalâ, ♀ corymbosâ compositâ polycephalâ, inv. ♂ ovatis squamis lanceolatis subacutis, ♀ oblongi squamis acuminatis, achæniis striatis glabris. ♀ in maritimis Carolinæ et Floridæ (Michx. ! Fras. ! Ell. ! ) et ad ripas fl. Mississipi (Pursh). Frutex 6-10-pedalis. *Baccharis angustifolia* Ell. ! sketch. 2. p. 318 non Desf. Capitula circ. 20-flora. (v. s.)

85. *B. GENISTIFOLIA*, fruticosa ramosissima glabra, ramulis striatis, foliis distantibus paucis sessilibus linearibus obtusis integerrimis subnerviis, capitulis ♂ ad apicem ramulorum 3-5 sessilibus spicatis-congestis ebracteatis, invol. obovatis squamis ovatis acutis margine membranaceis. ♀ in Brasiliæ prov. Rio-Grande. Habitus ferè *Genistæ tinctoriæ*. Folia 5 lin. longa,  $\frac{1}{2}$  lin. lata. Ind.

foem. ign. Pappus ♂ invol. non superans. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 835 miss.)

§. 5. *Sergilæ, foliis minimis aut subnullis, ramis non alatis.*

186. *B. scoparia* (Pers. ench. 2. p. 425. n. 44), fruticosa ramosissima glaberrima viscosa, ramulis gracilibus tetraquetris, foliis distantibus minimis linearibus oblongis obtusis integerrimis, capitulis ♂ terminalibus solitariis ovatis, invol. squamis ovatis. § in Jamaica australis montibus Liguansensibus. Calca scoparia Linn. syst. nat. 3. p. 234. Swartz! obs. 302. *Chrysocoma scoparia* Linn. amœn. 5. p. 381 et 404. *Sergilus scoparius* Gärtn. fr. 2. p. 409. t. 174. — Browne jam. 3: t. 34. f. 4. Vulgò *Mountain Broom tree* Anglis Jam. Spec. foem. non vidi et erg. genus dubium. (v. s. comm. à cl. Swartz et Bertero.)

187. *B. aphylla*, fruticosa ramosissima erecta glabra, foliis nullis aut ad squamulas minimas reductis, capitulis secus ramos sessilibus spicatis ebracteatis. invol. fl. ♀ squamis linearibus oblongis obtusis, pappo demum elongato apice crispo. § in campis editis prov. Sancti-Pauli Brasiliz legit cl. Lund. *Chrysocoma aphylla* fl. flum. 8. t. 1. *B. scoparia* affinis sed nullo modo viscosa, capitula multo majora et spicata, etc. Stirpem huic valde affinem ad Coquimbo Chiliensium lectam video in h. Lindl. An eadem? An species in posterum stabilienda? (v. s. comm. à cl. Lund.)

188. *B. nuda*, fruticosa? dichotomè ramosa erecta glabra, foliis nullis, capitulis 4-5 secus ramos sessilibus spicatis ebracteatis, invol. ovali, pappo cor. brevioris. § in Brasiliz prov. Fluminensi. *Chrysocoma nuda* fl. flum. 8. t. 5. Priori ex icone valde affinis. †.

§. 6. *Caulopteræ, nempe foliis oblongis linearibus aut subnullis, ramis alas foliaceas è foliorum basi utrinquè decurrentes gerentibus.*

189. *B. gaudichaudiana*, fruticosa interruptè bialata, alis planis semi-ellipticis, foliis minimis obtusis, capitulis secus ramos ramulosve sessilibus spicato-paniculatis, invol. squamis ext. parvis acutiusculis, inter. acuminatis, achenis brevibus ellipticis glaberrimis. § in Brasiliâ ad Sanctam-Catharinam legit cl. Gaudichaud. An *Cacalia sessilis* fl. flum. 8. f. 73? Simillima *B. articulata* sed nuda glauca, articuli elliptici latiores nempe 7-8 lin. longi 3 lin. lati. (v. s. comm. à cl. Gaudichaud et postea ♂ à cl. Lund.)

190. *B. articulata* (Pers. ench. 2. p. 425), fruticosa glauca interruptè bialata, alis planis angustis, foliis minimis obtusis, capitulis secus ramorum ramulorumve apices sessilibus spicatis, invol. campanulati squamis ext. obtusis, intimis in foem. acuminatis, acheniis brevibus ellipticis glaberrimis. § in Brasilia meridion. propè Montevideo (Comm. Sellow!) et in prov. Rio-Grande (h. Mus. reg. Par. ex h. Mus. imp. Bras.). *Conyza articulata* Lam. dict. 2. p. 94. Molina *articulata* Less! in linneâ 1831. p. 140. (v. s. ♂ et ♀.)

191. *B. parvifolia*, fruticosa interruptè bialata aut trialata, alis angustis planis foliis ellipticis obtusis, capitulis distantibus lateralibus sessilibus ad apices ramorum interruptè spicatis, invol. ♂ squamis ovali-oblongis vix acutis. § in Peruvia ad Avebrada de Cobrechillo ex h. Henke. Affinis *B. articulata* sed tunc distincta. Folia non squamiformia sed ovalia 2-2 ½ lin. longa. (v. s. in h. Henke à cl. de Sternberg miss.)

192. *B. crispa* (Spreng. syst. 3. p. 466), fruticulosa trialata, alis crispis coriaceis cartilagineo-marginatis, foliis ad squamulas minimas reductis, capitulis approximatis, invol. in ♂ cylindraceis squamis acuminatis. § in Brasiliâ meridionali (Sellow!), prov. Rio-Grande (h. Mus. imp. Bras.). Molina *crispa* Less! in linneâ 1831. p. 141. Alæ angustæ breves. Capit. 2 - lin. longa. Ind. foem. ign. (v. s.)

193. *B. genistelloides* Pers. ench. 2. p. 425 excl. syn. Martegr. ad Cereæ

- phyllanthum potius referend.), suffruticosa trialata, alis planis enerviis brevibus seu crebre interruptis, foliis minimis ad squamulas acutiusculas subternatas reductis, capitulis paucis (1-2) interruptè spicatis, invol. turbinati squamis omnibus acuminatis, rachide ex Less. nudâ et achæniis glabris cylindraceis sulcatis.  $\zeta$  in prov. Tacongâ Peruvîæ (Jos. de Jussieu!), propè Luxam et in Andibus Novo-Granatensium (H. et B.). *Conyza genistelloides* Lam.! dict. 2. p. 93. *Molina reticulata* Less. in linnæâ 1831. p. 143 non R. et P. (v. s.  $\sigma$  comm. à cl. Jussien.)
194. *B. venosa*, suffruticosa trialata, alis angustis venoso-reticulatis remotè interruptis?, foliis ad squamulas minimas reductis, capitulis ad apices ramorum apicato-glomeratis, invol. campanulati squamis omnibus in  $\sigma$  acuminatis.  $\zeta$  in Peruvîa argillosis collibus (R. et Pav.). *Molina venosa* R. et Pav. syst. 212. *A. B. genistelloide* differt articulis longioribus et ferè angustioribus, alis venosis, superficie totâ viscosâ resinosa, et capitulis magis approximatis. (v. s. ex h. Dombey comm. à Mus. reg. Par.)
195. *B. microcephala*, fruticulosa trialata, alis planis angustissimis remotè interruptis, foliis minimis oblongis, capitulis interruptè spicatis solitariis aut adgregatis, invol. campanulati squamis in  $\zeta$  interioribus acutis, in  $\sigma$  omnibus obtusis, achæniis obovatis, costis minutè puberulis.  $\zeta$  in Brasiliâ (Sellow!) prov. Rio-Grande (h. Mus. reg. Par. ex h. imp. Bras.). *Molina microcephala* Less.! in linnæâ 1831. p. 142. Alæ caulinae punctulatæ sæpè 2-3 poll. longæ, 2-3 lin. latæ. Folia 1-2 lin. longa, ad apices ramorum omninò abortiva. Capit. parva. (v. s. comm. à Mus. reg. Berol.)
196. *B. sagittalis*, fruticulosa trialata, alis planis remotissimè interruptis, foliis lineari-lanceolatis acutis internodio multò brevioribus, capitulis in spicam terminalem continuam aut subinterruptam dispositis, invol. campanulato-ovatis squamis in  $\sigma$  omnibus obtusis, in  $\zeta$  extimis acutis intimis acuminatis, achæniis glabris.  $\zeta$  in Chili (Née!), propè Talcaguanho (Cham!), Valparaíso (Macrae!), Coquimbo (Gaud.), ad Conceptionem (R. et Pav.?) et in Cordilleris excelis (cl. Gay!). *Baccharis genistelloides* Poepp. coll. Chil. 1. n. 213 non Pers. ex Less. *Molina sagittalis* Less.! in linnæâ 1831. p. 142. *Molina reticulata* R. et Pav. syst. p. 212? Folia 7-9 lin. longa. (v. s.)
- ß. Pappigii*, foliis infer. ovatis ovalibusve mucronatis, super. lanceolatis.  $\zeta$  in Chili. *Baccharis venosa*? seu *Baccharis tripterix* Poepp.! coll. Chil. 1. n. 210 diar. 46. (v. s.)
197. *B. phytumoides*, herbacea trialata glaberrima simplex seu ramosa, alis planis caule angustioribus, foliis internodio longioribus penninerviis oblongis basi attenuatis, capitulis glomeratis folio suo multò brevioribus in spicam interruptam dispositis, invol. campanulati in utroque sexu conformis squamis intimis acuminatis, achæniis obovatis secus costas minutè puberulis.  $\zeta$  in Brasiliâ meridionali (Sellow!). *Molina phytumoides* Less.! in linnæâ 1831. p. 146. Folia 2-3 poll. longa, 4-9 lin. lata. (v. s.)
198. *B. pentaptera*, fruticulosa quinque-alata, alis planis remotè interruptis, foliis ellipticis obtusis internodio multò brevioribus, ramis apice subcorymbosis, capitulis spicatis adgregatis, invol. cylindracei squamis in  $\sigma$  ext. subacutis, inter. longioribus acutioribus.  $\zeta$  in Brasiliâ (Sellow!), prov. Sancti-Pauli (h. Mus. imp. Bras.). *Molina pentaptera* Less.! in linnæâ 1831. p. 505. Fœm. non novi. (v. s.  $\sigma$  comm. à Mus. imp. Bras.)
199. *B. trimera*, suffruticosa trialata, alis planis reticulato-venosis hinc inde interruptis, foliis ad squamulas minimas acutas adpressas reductis, capitulis glomeratis interruptè spicatis, invol. campanulati squamis exter. acutis, intimis in  $\zeta$  acuminatis, in  $\sigma$  obtusiusculis aut acutis, achæniis oblongis glaberrimis.  $\zeta$  in Brasiliâ littorali (Sellow!), ad fretum Sanctæ-Catharinæ (Cham.), propè Rio-Grande (h. imp. Bras.), pr. Rio-Janeiro (Gaud.), Bahiam (Blanch!). *Molina trimera* Less.! in linnæâ 1831. p. 141. An *Cacalia decurrens* fl. flum. 8. t. 72? Alæ 3-4 lin. latæ 1-3 poll. longæ. (v. s.)

200. *B. CYLINDRICA*, herbacea trialata, alis planis venosis interruptis, foliis ad squamulas minimas reductis, capitulis spicato-glomeratis, invol. ♂ campanulato, in ♀ oblongo et cylindraceo, squamis acutis, ovaris glaberrimis. ♀ in Brasiliâ (Sellow!), prov. Rio-Grande (ex h. imp. Bras.). Molina cylindrica Less. in Linnaea 1831. p. 141. Differt à *B. trimerâ* capitulis valde approximatis, ♀ oblongo-cylindraceis meianthis squamis ciliatis acutis nec intimis acuminatis. (v. s.)

201. *B. MILLEFLORA*, fruticosa ramosissima trialata, alis planis latius crebre interruptis subreticulatis, foliis minimis vix ullis, capitulis spicato-glomeratis, invol. campanulati squamis in ♂ omnibus obtusis, in ♀ interioribus acutis, schœniis oblongo-ellipticis pube costarum minutâ. § in Brasiliâ legit cl. Sellow. Molina milleflora Less. in Linnaea 1831. p. 143. Alæ 4 lin. latæ, 1 1/2 poll. longæ. Recept. subnudum. Capitula numerosa. (v. s. comm. à Mus. reg. Berol.)

202. *B. MYRIOCEPHALA*, fruticosa ramosissima trialata, alis planis subreticulatis hinc inde interruptis, foliis ad squamulas minimas reductis, capitulis parvis glomeratis interrupte spicatis, spiciis numerosis paniculatis, invol. oblongo squamis in ♀ ext. brevibus obtusis, inter. elongatis apice mucronatis membranaceis, schœniis glabris. § in Brasiliâ prov. Minas Geraes legit cl. Vauthier (pl. exa. n. 265!) et in Bandâ orientali (Bacle!). Affinis *B. millefloræ*. Ind. masc. non vidi. (v. s.)

203. *B. JUNCIFORMIS*, herbacea? quinque-alata, alis planis latè linearibus reticulato-venosis, ramis apice angustissimè alatis, foliis ovali-acuminatis petiolulatis internodio multò brevioribus, capitulis spicato-glomeratis, invol. ♀ campanulati squamis ext. obtusissimis intimis acutiusculis, ovaris oblongo-minutè puberulis ♀? in Brasiliâ. Molina juncea Less. in Linnaea 1831. p. 145 et p. 505, sed non est *B. juncea* Desf. cat. h. Par. 1829. Folia 4-5 lin. longa. Masc. ign.

204. *B. POLYPTERA*, caule simplicissimo 7-8-alato, alis angustis valde elongatis, foliis elongato-lanceolatis integerrimis acutis suprâ glaberrimis subius cauleque inter alas densè et candidè tomentosis, spicâ interruptâ terminali valde elongatâ, capitulis sessilibus fasciculatis, invol. ♂ tomentosi squamis ovalibus mucronatis. § in Brasiliâ prov. Rio-Grande. Species distinctissima! Inv. fem. ign. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 746, 750 et 1066 mis.)

§. 7. Distichæ, nempe foliis minimis distichè imbricatis carinatis ramos apteros obtegentibus.

205. *B. FERRUGINEA* (Pers. ench. 2. p. 425. n. 48), fruticosa, ramis junioribus tomentosis foliis obtectis, foliis sessilibus semiamplexicaulibus carinatis approximatis bifariam imbricatis extus glabriusculis intus tomentosis, capitulis (masculis) solitariis terminalibus solitariis. § in Peruvia montibus excelsis vulgò Cordilleras et Punas, ubi vern. *Matara* et *Palmio* appellatur ex R. et P. Molina ferruginea R. et Pav. syst. 211. Rami breves subcorymbosi. Folia angusta extus subrugulosa ferè linearia. Invol. squamæ apice membranaceæ stramineæ subpatulæ. Pappus apice clavellatus. Valde affinis *B. thyoidi*. An hujus fortè indiv. masculum? (v. s. ex h. Dombey à Mus. reg. Par. comm.)

℞. *thyopsis*, foliis latioribus extus glaberrimis lævibus, invol. squamis apice scariosis sed non patulis. § in Amer. merid. cl. Bonpland olim comm. sub nomine *Conyza* thyoidis sed in opere Kunthiano ommissa. Folia omnino *B. thyoidis* sed capitula terminalia mascula. (v. s.)

206. *B. THYOIDES* (Pers. ench. 2. p. 425. n. 49), fruticosa, ramis junioribus tomentosis foliis densè obtectis, foliis sessilibus semiamplexicaulibus carinatis approximatis bifariam imbricatis extus glaberrimis intus lanatis, capitulis axillaribus sessilibus solitariis. § in Peruvia montibus primis legit Jos. d. Jussieu! *Conyza thyoides* Lam. dict. 2. p. 90. ill. t. 697. f. 3. Molina mucra

R. et Pav. syst. p. 211. Vulgò *Palmito* ex R. et Pav. Fl. masc. non vidi. (v. s. specim. Q ex J. Juss. ! Domb. ! Née ! Abadia !)

207. B. CATAPHRACTA (Spreng. syst. 3. p. 462), ramis ? distichè imbricatis conglutinatius glabris, foliis ovatis obtusiusculis rigidis, capitulis terminalibus sessilibus. — in Pern. Conyza cataphracta Willd. herb. Cæt. ign. †.

§. 8. *Lepidophyllæ*, foliis imbricatis squamæformibus, capitulis minimis adgregatis.

208. B. ? LEPIDOPHYLLA, fruticosa ramosissima, ramis teretibus junioribus tomentosis, foliis confertissimis sessilibus ovatis integerrimis concavis glutinosis coriaceis, capitulis ♂ ? circ. 5 sessilibus ad apices ramorum ovatis glutinosis 1-3 ?-floris, invol. squamis ovalibus flavescentibus. § in terrâ Van-Diemen prope Hobart-Town ad summitatem frigidam montis Wellington alt. 4100 ped. angl. supra mare legit cl. A. Cunningham. Genus novum Baccharideum Cunn. ! in Litt. 1835. Habitus *Lepidophylli*, sed frutex dioicus et masculus tantum mihi cognitus. Videtur genus proprium sed characteres mihi non satis clari. (v. s. comm. à cl. Cunn.)

† Non satis notæ.

\* *Gerontogea*. (A genere alienæ.)

209. B. ? ARBOREA (Linn. mant. 284), fruticosa, foliis alternis petiolatis elliptico-lanceolatis integerrimis scabridis, corymbo terminali paniculato, invol. subglobosi squamis imbricatis subulatis, acheniis striatis. § in Ind. or. insulæ Johanne sylvia (Koen. ex Linn.), in ins. Johanna canalisi Mozambici nec Ind. or. (Spreng.). B. arborescens Decaisn. nonv. Ann. Mus. h. n. Par. 3. p. 412. dicitur tripedalis et maxima ordinis ! Car. flor. et fruct. valde incompleti si *Baccharis vera*, sola esset species *Gerontogea*. †.

210. B. ? VILLOSA (Vahl symb. 3. p. 98), foliis lanceolatis petiolatis extrorsum serratis subtus ramisque villosis subtomentosis. — in monte Chadrâ Arabiæ felicis. Ex patriâ et ex invol. squamis lineari-setaceis potius ad *Conyzam*, suadente Persoon, aut ad *Blumeam* referenda.

\*\* *Americana*.

211. B. ? ASSUENSIS (H. B. et Kunth nov. gen. am. 4. p. 67), arborea, ramulis angulatis tomentoso-pubescentibus, foliis petiolatis oblongis acutis basi rotundatis grosse dentatis membranaceis supra glabris nitidis subtus incano-tomentosis, corymbis terminalibus ramosis foliatis, capitulis masculis ? circ. 13-floris longè pedicellatis, invol. squamis lanceolatis acutis, recept. paleis paucis inter flores onusto. § in frigidis montanis Quitensium in Paramo de Assuaye. Cor. flavæ ex Bonpl. albidæ ex Kunth. An stylus et ovarium certe abortiunt ?

212. B. ? BRYOIDES (Pers. ench. 2. p. 425. n. 43), fruticosa ramosa prostrata, ramulis confertis erectis, foliis minimis confertis linearibus subtus incanis, capitulis 6-8-floris terminalibus solitariis sessilibus, invol. cylindrici squamis oblongis. § ad fretum Magellanicum seu littus Patagonicum legit cl. Commerçon. Conyza bryoides Lam. dict. 2. p. 91. Cæt. ign. Genus dubium. †.

213. B. ? CANESCENS (Spreng. syst. 3. p. 466), herbacea, foliis fasciculatis petiolatis lineari-lanceolatis acutis subdentatis incano-tomentosis, paniculâ terminali. — in Monteideo. †.

214. B. CORYMBOSA (Pers. ench. 2. p. 425. n. 40), fruticosa, ramis corymbosis, foliis oppositis lanceolatis integerrimis margine revolutis capitulis axillaribus solitariis. § in Peruvia alpinis versas Pillao. Molina corymbosa R. et Pav. syst. 210. Ex Spreng. phrasis falsa et species eadem ac B. revoluta. †.

215. B. DAPENDENS (Pers. ench. 2. p. 424. n. 28), fruticosa scandens, foliis ovatis cordatisque cum acumine integerrimis subquiquenerviis, capitulis racemosis.

- § § in Andium nemoribus ad Cuchero, Casape et Campillo tractus. Molina dependens R. et Pav. syst. p. 202. Cæt. ign.
216. B. EMARGINATA (Pers. ench. 2. p. 424), fruticosa, foliis ovalibus emarginatis, capitulis subcorymbosis. § in Peruviz collibus copiose in Tarmæ prov. Molina emarginata R. et Pav. syst. p. 202. Cæt. ign.
217. B. ? FULIGINEA (H. B. et Kunth nov. gen. am. 4. p. 68), fruticosa, ramis angulatis pulverulento-pubescentibus, foliis petiolatis oblongis acutis basi rotundatis margine integerrimis et revolutis coriaceis supra glabris nitidis subis fuligineo-tomentosis, corymbis terminalibus ramosis foliatis, capitulis ♂ longi pedicellatis 8-floris, invol. squamis oblongis acutis, recept. paleas paucissimas gerente. § in montanis frigidis Quitensium cum B. Assuensi cui simillima.
218. B. GLABRA (Spreng. syst. 3. p. 466), herbacea, foliis oblongis acuminatis profunde serratis membranaceis utrinque glaberrimis, corymbo terminali. — in Montevideo. †.
219. B. LATIFOLIA (Pers. ench. 2. p. 424. n. 32), fruticosa, foliis lanceolatis triplinerviis argutè serratis, corymbis terminalibus nudis. § in Peruvia rueratis et campis prope Muna et Rondosubi vern. Chilca et *Anchochus dicta*. Molina latifolia R. et Pav. syst. 208. Cæt. ign. †.
220. B. ? QUADRINERVIA (Pers. ench. 2. p. 423. n. 6), foliis subovalibus obtusis mucronatis hinc inde serratis quadrinerviis subviscosis. — in Amer. merid. ex herb. Juss. Numerus nervorum par videtur à genere alienus. †.
221. B. RUPESCENS (Spreng. syst. 3. p. 464), fruticosa, foliis confertis spatulato-lanceolatis linearibusque acutis utrinque mucronato-bidentatis glabris opacis glanduloso-punctatis, racemis foliatis oligocephalis axillaribus. § in Montevideo. †.
222. B. SCABRA (Pers. ench. 2. p. 424. n. 34), ramis scabris, foliis oppositis ovato-lanceolatis acutis trinerviis serratis, corymbis terminalibus. § in Peruv Tarmæ prov. montibus. Molina scabra R. et Pav. syst. 210. Cæt. ign. †.
223. B. ? TOMENTOSA (Pers. ench. 2. p. 424. n. 30 non Thunb.), fruticosa, foliis oblongo-lanceolatis integerrimis tomentosis, capitulis subcorymbosis. § in Peruviz altis frigidis in Caxatambo prov. versus Chenchin. Molina tomentosa R. et Pav. syst. p. 203. Cæt. ign.
224. B. UNIFLORA (Pers. ench. 2. p. 425. n. 46), fruticosa, foliis lanceolatis serrato-dentatis confertis sessilibus, pedunculis terminalibus squamosis 1-cephalis. § in Peruviz collibus aridis ad Tarma. Molina uniflora R. et Pav. syst. 203 Cæt. ign. †.
225. B. ? VIRETA (H. B. et Kunth nov. gen. am. 4. p. 68), fruticosa, ramis pubescenti-hirtellis, foliis subsessilibus oblongo-lanceolatis acutis basi valdè angustatis apicem versus utroque latere 2-3-dentatis crassiusculis glabris glaucrescentibus, corymbis terminalibus simplicibus, capitulis pedicellatis, floribus hermaphroditis, acheniis albo-sericeis. § in declivitate occid. montium prope Cuernavaca Mexicanorum. An fortè Chrysocomea?

N. B. Præter species hinc inde citatas ad Baccharidem referendæ videntur species sequentes siue descriptione in Floræ fluminensis vol. octavo delineatæ. nempe: Chrysocoma cruciata t. 10, Chrysocoma indivisa t. 12, Chrysocoma maritima t. 22, Chrysocoma serrata t. 23, Chrysocoma dentata t. 27, Chrysocoma punctulata t. 49, Cacalia frutescens t. 64, Cacalia trinervia t. 65, Cacalia rosarinus t. 67, etc.

† Species exclusæ.

- B. ABSENTHIOIDES Hook et Arn. = Tessaria absinthioides.  
 B. ADNATA H. B. K. = Pluchea subdecurrens.  
 B. ÆGYPTIACA Forsk. = Pluchea Dioscoridis.  
 B. ARGUTA Pers. = Psidium balsamica.

- B. ARUNCA* Schrank. = *Conyza Gouani*.  
*B. ASTRIFORMIS* Bert. = *Triplolium conspicuum*.  
*B. BARSILAEFOLIA* Bert. = *Tessaria absinthioides*.  
*B. BRASILIENSIS* Linn. = *Vernonia scabra*.  
*B. CHILENSIS* Weinm. = *Euxenia grata*.  
*B. CUPRESSIFORMIS* Pêrs. = *Lepidophyllum cupressiforme*.  
*B. DIOSCORIDIS* Linn. = *Pluchea Dioscoridis*.  
*B. ERICOIDES* Pers. = *Vernonia ericoides*.  
*B. FETIDA* Linn. = *Pluchea fetida*.  
*B. FETIDA* Willd. = *Pluchea Marylandica*.  
*B. HALIMIFOLIA* Moench. = *Conyza ivæfolia*.  
*B. HIERACIFOLIA* Lam. = *Conyza Gouani*.  
*B. INDICA* Linn. = *Pluchea Indica*.  
*B. ILICIFOLIA* Lam. = *Brachylæna dentata*.  
*B. IVÆFOLIA* Linn. = *Conyza ivæfolia*.  
*B. LINEARIFOLIA* Pers. = *Psiadia linearifolia*.  
*B. LYCOPODIODES* Pers. = *Eriothrix lyco podioides*.  
*B. MADAGASCARIENSIS* Pers. = *Psiadia Madagascariensis*.  
*B. NERIIFOLIA* Linn. = *Brachylæna neriifolia*.  
*B. NITIDA* Wall. = *Blumea Chinensis*.  
*B. OVALIS* Pers. = *Pluchea ovalis*.  
*B. OVATA* Sieb. = *Pluchea ovalis*.  
*B. PTEROCAULON* Bojer. = *Monenteles Pterocaulon*.  
*B. RUGOSA* Willd. h. = *Vernonia patens*.  
*B. SALVIA* Lour. = *Pluchea balsamifera*.  
*B. SENEGALENSIS* Juss. = *Decaneurum Senegalense*.  
*B. TRIFURCATA* Trev. = *Conyza absinthifolia*.  
*B. VISCOSA* Lam. = *Psiadia integrifolia*.  
*B. VISCOSA* Walt. = *Pluchea bifrons*.

Subtrib. III. *TARCHONANTHÆ* Less. syn. p. 205 non 208.

Capitula aut dioica aut heterogama nunquam radiata, floribus ambis formineis pluriserialibus tenuissimis, disci hermaphroditis aut masculis paucioribus majoribus, antheræ caudatæ. — Folia alterna.

Div. 1. *EUTARCHONANTHÆ*. — *Tarchonanthæ* Less. syn. 208 non 205. — *Vernoniæ* *Tarchonanthæ* pleræque Cass. tabl. in op. phyt. 3. p. 76.

Capitula (ut in *Baccharis*) dioica.

CCXVIII. *BRACHYLÆNA* Brown. trans. Linn. soc. 12. p. 115. Less. syn. 208. — *Oligocarpus* Cass. dict. 36. p. 21. — *Tarchonanthi* et *Baccharidis* sp. auct.

Capitula multiflora dioica. Recept. angustum nudum (aut ex Cass. sparse bracteolatum sed bracteolas in specim. meis non vidi). Invol. squamæ imbricatæ siccæ corollis breviores. Corollæ tubulosæ inæqualiter 5-dentatæ. ♂ antheræ basi bisetæ concretæ. Stylus filiformis simplex. Ovarium hispidum abortivum parçè papposum. ♀ antheræ liberæ abortivæ. Stylus apice in ramulos breves latos divisus. Achænium



glanduloso-pubescent. Pappus setis biserialibus scabridus. — Frutices aut arbusculæ Capenses. Folia alterna coriacea breviter petiolata integra aut dentata suprâ glabra subtus glabra aut tomentosa. Capitula in racemis subpaniculatos disposita. Cor. luteæ.

1. *B. NERIIFOLIA* (Brown. et Less. ll. cc.), foliis lanceolatis integerrimis utrinque glaberrimis latitudine sextâ circ. longioribus brevissimè petiolatis, paniculis terminalibus polycephalis. ♂ ad Cap. Bonæ-Spei non infrequens. Burch! n. 7260, 4350. Drege! Eckl! et Zeyh. *Baccharis neriifolia* Lin. sp. 1204. Bern! prod. *Oligocarpa neriifolia* Cass. dict. 36. p. 22. *Tarchonanthus lanceolatus* Willd. Spreng. *Tarchonanthus dentatus* Ecklon! coll. in h. Donati. *Corymbus neriifolia* L'Her. ex Steud. Burch! cat. geog. n. 7260 ♂ et 4350 ♀. Panicula pube brevissimâ cinnamomeâ junior velutina, demum glabrata. Folia 3 polli longa, 6-7 lin. lata. (v. s.)
2. *B. GRANDIFOLIA*, foliis petiolatis lanceolatis longè cuneatis apice sinuatis obtusis utrinque glabris latitudine triplò longioribus subtus subferrugineis, paniculæ terminalis ramis divaricatis laxis. ♂ in Africâ Capensi ad Zneurebergen alt. 200-300-ped. legit cl. Drege! Petioli semipollic. Limbus fol. 4-4; polli longus, ferè sesquipollic. latus. Invol. *B. neriifoliae* sed ferrugineo-subvelutina. (v. s. ♂.)
3. *B. ELLIPTICA* (Less. syn. 208), foliis ellipticis denticulatis subtus tomentosis breviter petiolatis, racemis axillaribus glabris. ♂ ad Cap. Bonæ-Spei (Burch! n. 3267 et 4450), ad Klipplat rivier et Fisch rivier (Drege!). *Tarchonanthus ellipticus* Thunb. fl. cap. 638. *Tarch. camphoratus* Houtt! in h. Deless. (v. s.)
4. *B. RACEMOSA* (Less. syn. 208), foliis ellipticis oblongisve mucronatis integerrimis aut subdentatis suprâ glaberrimis subtus tomentosis brevissimè petiolatis, racemis ramulos laterales terminantibus 4-6-cephalis, capitulis distinctè pedicellatis. ♂ ad Cap. Bonæ-Spei in sylvâ Kukâ (Th.), Uitenhage (Eckl! n. 232, 126, 1414, 1326, 8 et 1597). *Tarchonanthus racemosus* Thunb. fl. cap. 638. *Baccharis cuneata* Spreng. in Zeyh! pl. exs. cap. n. 194 ex h. Moric. hæc verisim. referenda? — Burch! cat. n. 4450 ♂ et fortè 3267 ♀. An certè à prius diversa? Folia 7-11 lin. longa, 3-4 lin. lata. Invol. subtrilobis squamæ acutiserialibus, exter. brevissimæ subarctæ, inter. obtusæ dorso puberulæ. (v. s.)
5. *B. DENTATA*, foliis cuneiformibus indivisis aut subtrilobis grossè dentatis denticulatis suprâ glabris subtus adpressè tomentosis brevissimè petiolatis, margines revolutis, racemis ramos terminantibus subspicatis oligocephalis. ♂ ad Cap. Bonæ-Spei in colon. orient. district. Albany (Burchell! n. 3400). *Zneurebergen* (Drege!) et Cafferland (Eckl!). *Tarchonanthus dentatus* Thunb. fl. cap. 638? *Baccharis ilicifolia* Lam. dict. 1. p. 345? Folia pollicem long. 4-7 lin. lata, inter se formis sat varia sed semper basi cuneata. Invol. squamæ acuminatæ (v. s. ♀.)
6. *B. salicina*, foliis lanceolatis per totam ferè longitudinem serratis, paniculis (masculâ) terminali polycephalâ. ♂ in territorio cesso (Eckl!) ad Omsamwulo et Omsamwulo (Drege! n. 5054 et 3659). Folia basi nunc cuneata nunc obtusa, apice indivisa aut fortè trifida, ad margines vix subrevoluta. An var. sexui masculo propria? An species propr.? (v. s. ♂.)
7. *B. DISCOLOR*, foliis petiolatis oblongo-obovatis obtusis repando-dentatis breviter petiolatis suprâ glabris subtus petiolis ramulisque tomentosis, racemis simplicibus subramosisve axillaribus folio vix longioribus oligocephalis, invol. squamis pluriserialibus imbricatis dorso glabris margine araneoso-lanatis. ♂ ad Cap. Bonæ-Spei in Coloniz orient. distr. Uitenhage (Burchell! cat. n. 3751 Eckl! ad Omsamculo et Omtendo (Drege!), ad sinum Algoa (Forb.). Folia 3 polli longa (incl. petiolo 4 lin.), pollicem lata. Achenia matura teretiuscula stricte ferè glabra. (v. s.)

**CCKIX. TARCHONANTHUS** Linn. gen. n. 301. Gærtn. fr. 2. p. 392. t. 166. f. 12. Lam. ill. t. 671. Cass. dict. 52. p. 245. Less. syn. 208 non Vaill. — *Tarchonanthi* sp. Thunb. Willd.

Capitula sæpius multiflora dioica. Invol. marium squamis 5 uniseri-  
libus ad medium coalitis, fœmineorum squamis plurimis distinctis  
biseriatis. Recept. ebracteolatum hirsutum. Cor. tubuloso-campa-  
nulatæ 5-dentatæ extus hirsutæ viscosæ, intus glabræ. ♂ antheræ in  
tubum coalitæ exsertæ breviter appendiculatæ et longè caudatæ, caudis  
setosis, filam. glabris. Ovarium abortivum. Nectarium amplum callo-  
sum apice concavum intra corollam ovarium mentiens (1); stylus fili-  
formis vix apice subbilobus. ♀ Stam. abortiva. Nectarium nullum.  
Stylus exsertus apice bilobus, lobis arcuato-revolutis. Achænium lanâ  
hirsutissimum epapposum. — Arbusculæ Capenses. Folia alterna petio-  
lata coriacea, supra reticulato-areolata demum glabra subtus tomentosa  
aut rariùs ad axillas subsolitaria. — Species omnes ab Hottentotis *Mo-  
hanka* dicuntur ex Burch. trag.

#### Sect. I. EUTARCHONANTHUS.

Capitula fœminea 3-5-flora, invol. squamis latè ovatis donata. Achænia hirsu-  
tissima. — Folia indivisa.

1. **T. CAMPHORATUS** (Linn. sp. 1179), foliis oblongis obtusiusculis integerrimis  
aut subdenticulatis, adultis suprà glabris reticulato-venosis subtus tomentosis.  
paniculâ terminali polycephalâ. ♂ ad Cap. Bonæ-Spei frequens Drege! Eckl.  
Thunb. fl. cap. 637. Berg. cap. 236. — Herm. lugd. b. 229 ic. Pluk. alm. t. 174.  
f. 1. Lam. ill. t. 671. f. 1. Folia 2-3 poll. longa, 6-10 lin. lata, basi plus minus  
cuneata. Frutex camphoræ odorem redolens. (v. v. c. ♂ et s. sp. ♀ et ♂.)
2. **T. LITAKUNENSIS**, foliis oblongis integerrimis acutiusculis adultis suprà glabris  
tenuissimè reticulatis areolis impresso-porosis subtus tomentosis, paniculâ ter-  
minali polycephalâ. ♂ in Africâ australi extratropicâ inter flumen Makkwa-  
rin et urbem Litakun legit cl. Burchell et mecum sp. fœm. fructiferum sub  
n. 2202 cat. geog. comm. Conf. Burch. tr. 2. p. 340. Idem ni fallor ad Omsam-  
culo et Omtendo legit cl. Drege! *T. camphorato* similis sed folia 15 lin. longa,  
3 lata et diversè reticulata. Paniculæ rami et ramuli multò tenuiores, albedo  
nec rufescenti-velutini. Specim. masc. mihi ign. (v. s.)
3. **T. OBOVATUS**, foliis obovatis obtusis basi cuneatis suprà glabris reticulato-ve-  
nosis subtus tomentosis, capitulis ad axillas foliorum solitariis pedicellatis. ♂ ad  
Cap. Bonæ-Spei in regione Trans-Gariepinâ propè Sensavan. Folia 8 lin. longa,  
3-4 lin. lata. Pedicelli cum capitulo circ. folii longitudine. Capitula fructifera  
priori similia sed ad axillas solitaria. (v. s. comm. ♀ à cl. Burchell sub  
n. 2155.)
4. **T. ANGUSTISSIMUS**, foliis oblongo-linearibus integerrimis suprà glabris vix  
reticulatis subtus tomentosis, capitulis solitariis. ♂ ad Cap. Bonæ-Spei in re-  
gionibus interioribus propè Karra-Poort (Burchell! n. 2691), ad Zneenreber-  
gen (Drege!), in Albany et Tambukiland (Eckl.). Folia 7-8 lin. longa, 1 ½-2  
lin. lata. Capitula parca et nimis juniora ut structuram discernere possim et  
ideò mihi ut jam cl. inventori genus omninò dubium. (v. s.)

*N. B. T. minor* Less. syn. 208 (de quo dicuntur : Folia brevissima, capitulorum  
maseolorum squamæ spathulato-obovatæ acutæ foliaceæ impellucidæ extus densæ

(1) Quâ insolitâ nectarii grossificatione decepti fuerunt Bergius et Linnæus. Ipse  
olim hunc errorem, sed dubitans et theoriam veram simul indicans, admisi in dis-  
compos. ann. Mus. h. nat. Par. 16. p. 138.

tomentosæ et circiter sesquilineam longæ), fortè ad quandam ex duabus ultimis speciebus referendus, sed cum capit. fœminea sola viderim, mihi obscurus remanet.

#### Sect. II. HENOTOGYNA.

Capitula fœminea uniflora, invol. squamis linearibus donata. Achenia parva hirsuta. — Folia apice grossè trifida. — An genus?

5. *T. TRILOBUS*, foliis cuneatis margine integris aut subinnotatis apice obtusè trifidis, lobo medio minore, suprà glabris minutissimè reticulatis, subtus hirsutis subtomentosis penninerviis reticulatisque, racemis in paniculam subdispositis. In asperis Africæ austr.-orient ad Omtendo legit cl. Drege! Species distinctissima, ferè inter *Brachylænas* et *Tarchonanthos* veros ex habitu media. Folia rugosa petiolata 3-4 pollic. longa, 1-1½ lata, lobis exceptis *Buddleia* aut *Salvia* folia referentia. Flor. masculi omninò *Eutarchonanthi*, fœminei 1-flori; achenia epapposa ovata per anthesin saltem parè hirsuta. (v. a.)

#### † Species exclusæ.

- T. DENTATUS* Thunb. = *Brachylæna dentata*.
- T. ELLIPTICUS* Thunb. = *Brachylæna serrulata*.
- T. ERICOIDES* Linn. f. = *Eriocephalus glaber*.
- T. GLABER* Linn. f. = *Brachylæna dentata*.
- T. LANCEOLATUS* Thunb. = *Brachylæna neriifolia*.
- T. RACEMOSUS* Thunb. = *Brachylæna racemosa*.

#### Div. II. PLUCHEINEÆ Cass. tabl. sub *Vernoniaceis* (1). — *Epaltides* Less. syn. 205.

Capitula heterogama seu monoica.

CCXX. *BLUMEA* DC. in ann. bot. Guill. 1833. v. 2. p. 514 (2). non Spreng. nec Nees (3). — *Erigeron* Don prod. fl. nep. p. 171. non Linn. — *Conyze* species pleræquæ Blum. bijdr. Wall. cat.

Capitulum multiflorum heterogamum, fl. radii multiserialibus feminis tenuissimè filiformibus truncatis aut sub-2-5-dentatis, disci paucis (5-25) masculis cylindricis 5-dentatis, fauce non aut vix dilatati. Recept. planum sæpius omninò nudum interdum tenuissimè fimbriiferum hirsutum. Invol. pauciseriale subimbricatum, squamis linearibus acuminatis. Anthere basi tenuissimè caudatæ. Achenium teretiusculum. Pappus 1-serialis, setis capillaribus vix subscabris. — Herba aut suffrutices. Species *Gerontogæ*, ferè omnes ex Indiâ orientali seu insulari seu præsertim continentali, paucissimæ Africanæ. Folia alterna sæpius cum caulibus variè villosa. Capitula paniculata aut laxè subcorymbosa. Flores flavi aut purpurascens. — Genus vastissimum à *Conyze* et

(1) Stylus enim florum disci non malè refert stylos *Vernoniacearum*, sed affinitatè potius juxta *Conyzeas* hæc genera trahunt.

(2) Caractere emendato ex observationibus per litteras à cl. Decaisne communicatis.

(3) *Blumia* Nees est *Talauma*. *Blumia* Spreng. est *Reinwardtia* Blum., nomen verdentissimè à Sprengelio injustè mutatum et sanctè servandum cum *Reinwardtia* Spreng. sit. *Dufourea* Kunth. = *Prevostea* Choisy. = *Calyrobolus* Willd., et *Reinwardtia* Dumort. = *Lini* species.

agnalo achæniis teretiusculis nec plano-compressis, ab Erigeronte alis nullis et achæniis teretiusculis, à Laenneciâ pappo 1-seriali distinctum, à Pluchea tandem corollis masculis cylindricis, invol. squamis linearibus vix imbricatis, et inflorescentiâ plus minus paniculatâ versum, dicavi cl. C. L. Blume qui plures hujus species ipse detexit qui Botanicam Indicam strenuè promovit.

**Ser. I. APTERÆ, nempe foliis caulinis non decurrentibus.**

*Distitifloræ, nempe foliis integris dentatis, pedunculis solitariis monocephalis.*

**B. AMPLECTENS** (DC. in Wight contrib. bot. ind. p. 13.), caule herbaceo muscoso tereti divaricato-ramoso, foliis junioribus villosis adultis subglabris, obois caulinis ovatis semiamplexicaulibus acutè dentatis, infimis hinc inde basi necis, pedunculis paucis subterminalibus folio paulò longioribus patentim villosis 1-cephalis, invol. squamis linearibus aristatis disco paulò longioribus. ① in ruderalis Indiæ Maderaspatanæ frequens, circa Negapatam (Wight!). Erigeron obliquum Linn. mant. 572? ex Russ! herb. et idè Conyza obliqua Willd. sp. 3. p. 1930? C. amplexicaulis Lam. dict. 2. p. 85. Willd. l. c.? Folia rar. subrotunda obtusa, ovata, ovali-oblonga acuta, plus minus serrata, semper semiamplexicaulia ut dicitur de C. amplexicauli et ex sicco obliqua ut de C. obliqua. Flor. fœm. pluriser. tenues stigm. exsertis rubris, masc. circ. 15-20 anth. exsertis. Conyza pumila Pers. ench. 2. p. 427, huc ut varietas citata est fortè diversa sed vix nota. (v. s. in h. Cur. Ind. or. nunc soc. Linn. Lond. et in h. Wight.)

**B. ARENARIA** (DC. in Wight contrib. bot. ind. p. 13), basi glabriuscula apice undiquè sericeo-villosissima, caule à basi ramoso diffuso tereti, foliis oblongis sessilibus mucrone calloso acutis integerrimis aut acutè denticulatis, pedunculis solitariis 1-cephalis hirsuto-lanatis axillaribus terminalibusque, invol. squamis linearibus acuminatis dorso hirsutis apice ciliatis, corollis masc. apice barbatis. ① in arenosis Indiæ Maderaspatanæ propè Negapatam (Wight!). Conyza villosa herb. Mair! ex Wall. cat. n. 3105 non 3083 nec Willd. Radix longa gracilis. Flor. fœm. tenues stylo longè exserto bifido, masc. pauci. Rec. nudum. (v. s. in h. Wight et in h. hon. Cur. Ind. or. nunc Soc. Linn. Lond.)

**B. INTERIFOLIA**, herbacea erecta ramosissima infernè pilis confertis anglu-ducosis subhirtella, supernè pilis longioribus sparsis pilosula, foliis lineari-lanceolatis acutis integerrimis, summis linearibus, ramulis elongatis apice 1-cephalis nudis, invol. squamis dorso ad basin pilosiusculis ceterum glabriusculis linearibus demum reflexis. ① in sylvis aridis Novæ-Hollandiæ bor.-occid. ad Port-Keals lat. 14° legit cl. A. Cunningham septemb. flor. Herba semipedalis. Radix gracilis simplex. Flores flavi, masc. in specim. incompleto destructi et idè genus subdubium. (v. s. comu. à cl. Cunn.)

**B. TENELLA** (DC. in Wight contrib. bot. ind. p. 13), caule herbaceo multispiriti puberulo tereti parè ramoso, foliis oblongis basi obtusè cordato-semiamplexicaulibus acutis distanter acutè et calloso-dentatis pube parvâ scabridis, ramis acuminatis subintegerrimis, pedunculis terminalibus et oppositifoliis 1-cephalis puberis folio triplò longioribus, invol. squamis linearibus acuminatis disco fere sublongioribus. ① in insulâ Timor leger. Botanici exped. Baudianæ, et in arenosis versus Negapatam (cl. Wight!). Decaisn! nouv. ann. Mus. h. Par. n. 3. p. 411. Fl. flavi, fœm. innumeris tenuissimis masc. 8-9, anth. inclusis. Valdè affinis B. oligocephalæ et B. amplexicenti. (v. s. comm. à Mus. vg. Par.)

**B. ANAGALLIDIFOLIA** (DC. l. c. p. 13), caulibus à collo multiplicibus diffusis ramosis teretibus apice puberulis, foliis caulinis ovali-oblongis sessilibus hinc inde acutè dentatis, ramis lanceolatis integerrimis puberulis, capitulis longè pedicellatis solitariis axillaribus terminalibus in paniculam foliosam laxam umbocorymbosam dispositis, invol. squamis linearibus acutis glabriusculis disco paulò longioribus. ② in Indiâ or. Maderaspatanâ ad Negapatam in humidis

- frequens (Wight!), in Pegu (Belong.). *Conyza anagallidifolia* Wall.! n. 309. comp. n. 208. Recept. nudum punctulatum. Fl. foem. innumeri tenuissimi masc. 10-12 cor. apice subbarbata. Achæmium glabrum. An forte *Conyza esbra* Linn. mant. 113? quæ vix nota!
6. *B. GLUTINOSA*, herbacea erecta paniculato-ramosa parçè pilosa tota glauce viscosa, ramis teretibus, foliis oblongis grossè et acutè dentatis basi attenuatis, capitulis pedicellatis subsolitariis, invol. squamis linearibus glabrinaculis, floribus masculis circ. 3. — in aridis insulæ Madagascæ circa pagum Majangy in sinu Bombatæ legit cl. Bojer. Cor. glabræ, foemineæ innumerae. Achænia glabra. (v. s. comm. à cl. inv.)
7. *B. OLIGOCEPHALA* (DC. in Wight contr. bot. ind. p. 13), parçè puberula herbacea multicaulis subdiffusa, foliis infer. obovato-oblongis basi cuneatis, super. oblongis ovalibus sessilibus subsemiauriculatis omnibus apice remotè et acutè dentatis, pedunculis 1-3 ex apice caulium 1-cephalis villosis folio longioribus, invol. squamis linearibus acutis disco æquantibus. ① circa Madras (C. amplexicaulis Wight! in Wall.! cat. n. 3104. comp. 214 non Lam. nec Michx. Herba vix semipedalis, affinis *B. bifoliata*, sed præter ear. indic. fl. mac. 15-20, antheris exsertis. (v. s. in h. soc. Linn. Lond. et comm. à cl. Wight.)
8. *B. PUBIFLORA*, caule herbaceo erecto ramoso pilis mollibus petalis plus minus hispidis foliis sagittato-semiauriculatis ovali-oblongis acutis subintegerrimis aut acutè dentatis, ramulis floridis 1-cephalis subnullifoliis pedicelliformibus, invol. squamis linearibus acuminatis flores æquantibus, cor. disci apice hirsutulis. — in Indiâ Maderaspatanâ. *Erigeron asteroides* Roxb. ex Wall.! cat. a herb. n. 2975. comp. n. 85. Styli fl. foem. longè exserti! Variat ramis fere glabris aut molliter apice villosissimis foliorum dentibus nullis aut strigatis 7-8. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
9. *B. BIFOLIATA* (DC. l. c. p. 14 sub *Conyza*), caule herbaceo adscendente tereti ramoso apice præsertim puberulo, foliis acutè dentatis, infer. ovatis basi longè attenuatis glabrinaculis, super. sessilibus oblongis cuneatis puberis, pedunculis axillaribus supra basin sæpius folia duo subopposita gerentibus, pedicellis elongatis 1-cephalis, invol. squamis linearibus acutis disco subæqualibus. ① in Indiæ or. peninsulâ frequens in nemorosis humidis ex Wight! C. foliolosa Wight! in Wall.! cat. n. 3092. comp. 202 non n. 3030 nec Willd. C. bracteolata Rottl.! in h. Puer. Pluk. alm. t. 177. f. 1. Habitus varius et fortè spec. 2-3 sub hac confusæ. In specim. diversis eodem loco collectis video folia nunc longè nunc breviter petiolata, super. et involucra plus minus villosa, fl. nec indiv. 8-16, etc., in omnibus recept. nudum Cor. masc. flavidæ vix ac ne vix apice barbellatæ antheris subinclusis, achænia glabra. (v. s.)
- \*\* *Paniculatæ, nempe foliis dentatis laciniatisve, pedunculis nunc axillaribus polycephalis nunc verò in paniculam dispositis, invol. cinereo-pubescentibus*
10. *B. BARBATA* (DC. l. c. p. 14. sub *Conyza*. Wall.! cat. n. 3099. comp. 209). caule herbaceo erecto simplici tereti pilosiusculo, foliis lanceolatis basi attenuatis acutè et distanter serratis acutis suprà sparè puberis subtus densè et adpressè sericeo-villosis, ramis axillaribus foliosis folio brevioribus apice nemosis, summis corymbosis, invol. squamis linearibus acutis glabrinaculis disci longitudine. ① in Indiæ peninsulâ legit cl. Wight. Herba pedalis. Folia 1-2 poll. longa, 5-6 lin. lata. Recept. nudum. Fl. flavidæ, foem. innumeri styli exserto bifido, masc. 5 cor. apice glabrinacula. Achæn. glabra. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et in herb. Wight.)
11. *B. CHAMISSONIANA*, caule herbaceo erecto tereti apice pilis mollibus plus minus hispidis, foliis obovatis obtusis basi cuneatis subsessilibus denticulatis atropæ molliter pubescenti-villosis pedunculis axillaribus folio brevioribus oligocephalis, invol. squamis linearibus disco æqualibus pubescentibus. ① in Laniâ legit cl. de Chamisso. C. bifoliata Cham! et Less. in Linnæa 1831. p. 135 (v. s. comm. à cl. Chamisso.)
12. *B. AXILLARIS*, caule tereti erecto herbaceo pubescenti-villoso, foliis petalis

obovatis ovalibusve membranaceis argutè denticulatis penninerviis, petiunculis folio brevioribus foliola 1-2 gerentibus 7-8-cephalis, ultimis in paniculam dispositis, invol. squamis dorso hirsutis linearibus acuminatissimis. (C) in insula Mauritii ad ripas rivorum (Bont. ! Bojer !). Conyza axillaris Lam. ! dict. 2. p. 84. Petioli pollicares. Limbi fol. 2  $\frac{1}{2}$  poll. longi. 15-18 lin. lati basi subconnati. Pappus albidus. Cor. purpurascens. Affinis B. Wightianæ et Chamissonianæ, sed foliis petiolatis ab utraq. facile differt. C. axillaris Willd. enum. 8-1. Mura. cat. suppl. 12, est certe diversa sed mihi ignota. (v. s.)

13. B. MANILLENSIS, caule herbaceo erecto tereti striato hirsuto supra parè folioso, foliis semiamplexicaulibus sparè hirsutis obovatis sinuato-dentatis subtus pallidioribus, racemo terminali simplici, capitulis jun. subsessilibus, invol. squamis linearibus ciliatis squarrosis. (C) in Loxoniâ legit cl. de Chamisso. Conyza Manillensis Less. in Linnæâ 1831. p. 135. Ovaria hirsutiuscula. Affinis dicitur B. Wightianæ var.  $\gamma$ .

14. B. WIGHTIANA (DC. l. c. p. 14), caule herbaceo erecto tereti striato puberulo, foliis inf. petiolatis ovalibus basi cuneatis acutè et inæqualiter dentatis subobtusis, super. cuneatis ovali-oblongis, capitulis in paniculam racemosam interruptam terminalem digestis, invol. squamis linearibus acutis inter. glabrimaculis disco paulò longioribus demùm reflexis, cor. masculis extus tuberculoso-glandulosis. (C) in Indiâ Maderaspatanâ legit cl. Wight, ad Pondichéry (Belanger !), et in regione Birmanorum circa Prome (cl. Wallich). Conyza Wightiana Wall. ! cat. n. 3093. comp. 203. Recept. nudum. Fl. form. innumeri tenues, masc. 8-12 ex sicco subrubentes. Achænia glabra. (v. s. comm. ab hon. Cur. Ind. or.)

*P. thureiformis*, foliis fere omnibus oblongo-cuneatis subacutis, capitulis thyrsoideo-confertis. In Indiâ Maderaspatanâ. C. bifoliolata Wall. ! cat. n. 3091. comp. 201 non Linn. (v. s. comm. cum a.)

$\gamma$ . *Madagascariensis*, floribus masc. 5-6. — in agris cultis ins. Madagascar legit cl. Bojer. Similis var. a. (v. s.)

15. B. CUNNINGHAMI, caule herbaceo erecto tereti paniculato-ramoso basi foliisque glabris apice molliter villosis, foliis sessilibus oblongo-lanceolatis grossè et irregulariter inciso-dentatis lobulis dentibusve acutis rigidulis, capitulis paniculatis, invol. squamis linearibus acuminatis dorso puberis apice ciliato-barbatis demùm reflexis. — in umbrosis ripis abruptis Novæ-Hollandiæ bor.-occid. ad Regent's-River latit. 15° legit cl. A. Cunningham. Herba circ. bipedalis. Flores flavidi, masc. pauci. (v. s. comm. à cl. inv.)

16. B. MURRA, tota cineres molliter villosa, caule herbaceo erecto tereti ramoso, foliis ovali-oblongis aristato-serratis basi attenuatis, inf. petiolatis, infernè ly-rato-appendiculis lobatis angustis acutis nunc à limbo distinctis nunc cum eo confluentibus, panicula ramis erectis apice confertè corymbosis, invol. squamis linearibus vix acutis villosis disci longitud. cor. masc. apice barbatis. — in Indiâ orient. prope Holsagung legit cl. Hamilton. Conyza Murra Ham. ! in Wall. ! cat. n. 3095. Rec. nudum. Fl. flavidi, masc. 20-25. Achænia glabra. (v. s. in h. hon. Cur. Ind. or. nunc soc. Lin. Lond.)

17. B. LESCHENAUTIANA, caule erecto herbaceo tereti piloso, foliis ovalibus serratis puberulis, inf. obtusis basi attenuatis, sup. sessilibus acutis, ramulis floridis hirsutissimis oligocephalis infer. axillaribus folio brevioribus, super. in paniculam brevem digestis, invol. squamis linearibus hirsutis discum æquantibus. — in Zeylonâ et in Indicâ peninsulâ legit cl. Leschenault. Rec. subalveolare nudum. Fl. form. innumeri stylo exserto bifido purpureo, masc. 10-11 glabri anih. semixsertis. Achænia glabra. Pappus panicetis albus. Valdè affinis B. molli. (v. s. comm. à cl. Leschenault.)

18. B. LACTUCIFOLIA, infernè sparè pilis rigidulis scabra, supernè confertius et mollius velutina, caule herbaceo erecto tereti apice ramoso, foliis infer. ovatis acuminatis grossè dentatis in petiolum longè attenuatis, mediis ad basin auriculato-cordatis, summis sessilibus oblongo-lanceolatis, paniculâ laxâ patente,

ramis spicato-racemosis, capitulis pedicellatis dissitis, invol. squamis linearibus acutis disci longitudine. (1) in Ind. or. regione Birmannorum ad fl. Salem supra Troglā legit cl. Wallich et nī fallor ad Sornogon cl. Hæntke. Conyza betucefolia Wall.! cat. n. 3088. comp. n. 198. Fl. fem. innumeri, stylo exserto bifido. masc. circ. 25-floris, lobis ciliato-hirtellis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

19. *B. LACERA* (DC. in Wight contr. p. 14), tota tomentosa aut pubescenti-hirsuta albida, caule herbaceo erecto tereti, foliis obovato-oblongis grossè dentatis sæpè basi incisīs aut pinnatifidis, paniculā oblongā aut corymbosā confertā aut laxā, invol. squamis linearibus acuminatis flores aequantibus. — ad oryzeta Ind. or. continent, et insularis nec non in Chinā. Flor. flavi, fem. innumeri, stylo bifido, masc. circ. 20. Achenia subangulata glaberrima. Planta multiformis fortè dividenda, sed caract. non satis perspicui.

a. *Burmanni*, foliis lyrato-pinnatifidis, ramis paniculatis. — in Javā. C. lacera Burm. fl. ind. 180. t. 59. f. 1. Blum.! bijdr 897. (v. s.)

β. *Commersonii*, foliis obovatis grossè dentatis, capitulis confertis. — in Javā. Con. lacera Lam.! dict. 2. p. 83. (v. s.)

γ. *Wallichii*, foliis obovatis profundè et acutè dentatis, paniculā laxā. — in Silhet, Chinā, etc. Con. lacera Wall.! cat. et herb. n. 3045 et 3082. comp. n. 155 et 192. C. villosa Wall. II. cc. n. 3083 et 193, ab hac specie non differre videtur. (v. s.)

δ. *Blumei*, foliis obovatis acutè et exsertè serratis, paniculā laxā. — in Javā Conyza Javanica Blum.! bijdr 897. (v. s.)

20. *B. LACINIATA*, caule herbaceo erecto ramoso tereti puberulo apice mollior et parcè piloso, foliis infer. pinnatipartitis utrinquè subscabris lobis utriusquè 4 oblongis ovalibus summis cum impari subconfluentibus, omnibus subrepundis acutè et inæqualiter dentatis, summis oblongis sessilibus hinc inde serratis, paniculā amplā polycephalā, ramis elongatis apice thyrsoides, capitulis pedicellatis, invol. squamis linearibus acuminatis disco longioribus. — In Ind. orient. Conyza laciniata Wall. cat. n. 3100. comp. n. 210. Recept. nudum. Fl. masc. 15-16 lobis apice puberulis. Achenia pilis sparsis scabrida oblongo-tertia. (v. s. in herb. hon. Cur. Ind. or. nunc Soc. Lin. Lond.)

21. *B. CERNUA*, caule herbaceo erecto tereti subsimplici apice puberulo apice velutino-villoso, foliis obovatis basi attenuatis grossè et acutè dentatis suprà puberulis subtus villosis, infer. longè basi productis, summis oblongis utrinquè villosis, paniculæ elongatæ ramis folio longioribus apice capitula glomerata subsessilia cernua gerentibus, invol. squamis linearibus acutis flores aequantibus. — in Indiā orientali ad Bhiratghur februario flor. legit cl. Hamblus. Conyza cernua Hamilt.! in Wall.! cat. n. 3080. comp. n. 190, non Blum. Folia infer. 6 poll. longa, 1 ½ lata. Fl. fem. innum. tennes, stylo exserto bifido. masc. circ. 7-8. Pappus albus. Achenia puberula. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

22. *B. ? LEUCANTHEMA*, cinereo-pubescenti herliacea, foliis elliptico-oblongis acuminatis sessilibus denticulatis, corymbis compositis patulis polycephalis in paniculam amplam digestis, invol. tomentosi squamis linearilanceolatis aristatis discum subaequantibus. 2l. ad Bessaria pagum Nepaliæ. Erigeron leucanthum Don prod. fl. nep. p. 171. Folia inf. spithamea, 2 poll. lata. Fl. masc. passim.

23. *B. TRICHOPHORA*, tota villis sericis confertis hirsuto-villosa; caule herbaceo stricto erecto purè ramoso, foliis obovatis acutè dentatis, infimis breviter petiolatis obtusis, caulinis basi cuneatis, ramulis floridis folio longioribus in paniculam angustam elongatam dispositis, invol. squamis linearibus disco longioribus. (1) in Nepaliā legit cl. Wallich. Erigeron molle Don prod. fl. nep. p. 172. sed. div. à Conyza molli Willd. An fortè Bl. Wightianæ var.? Recept. nudum Fl. purpurei fem. innum. stylo exserto bifido, masc. 10-12 glabris. (v. s. comm. à cl. Wallich anno 1821.)

24. *B. DAGRENA*, caule herbaceo erecto apice præsertim mollior villosa. (v. s.)

sessilibus obovatis dentatis, infer. basi longè attenuatis in parte attenuatè acutè incisio-dentatis suprà pubescentibus subtus villosis, super. utrinque adpressè sericeo-villosis, paniculâ contractâ sericeo-villosâ, invol. squamis linearibus disci longitudine. -- in Africâ austr.-orient. ad Zw. Omlata et Omsamwahu legit cl. Drege n. 5055. Habitu ferè *B. trichophoræ*. Invol. pubens post anthesin reflexum. Squamæ et styli purpurascens. Recept. punctulatum. Fl. masc. 8-10. (v. s.)

*ß. gracilis*, caule gracili simplici pubescente, foliis infer. basi integerrimis. — euna var. a legit cl. Drege. (v. s.)

25. *B. PARVIFOLIA*, tota et præsertim ad apicem cinerea pubescenti-villosa, caulibus è collo plurimis erectis ramosis, foliis infer. obovatis basi attenuatis breviter petiolatis acutè dentatis, superioribus oblongis sessilibus vix dentatis, paniculâ ramosâ polyccephalâ, invol. squamis exter. brevibus dorso subbirsutis inter. linearibus acuminatis flores subsequantibus demùm reflexis. (1) in Nepaliâ legit cl. Wallich. *Conyza parvifolia* Wall! cat. n. 3004. comp. n. 114. *Erigeron glomeratum* Spreng. in litt. ad Wall. Radix dura subsimplex. Caulis 8-10 poll. Folia pollicem longa, 6 lin. lata. Cor. purpureæ, fœm. innumere, stylo bifido, masc. 8-10. An *Bl. oxyodontæ* var.? (v. s. comm. ab hon. Cor. merc. angl. Ind. or.)

26. *B. TIMORENSIS* (DC. in Decaisne! nouv. ann. Mus. h. n. Par. 3. p. 411), caule erecto tereti apice puberulo basi subglabrato, foliis oblongis acutis puberulis irregulariter serratis, infimis basi attenuatis, mediis basi iuncis auriculâ utrinque exsertâ sublineari semiamplexicaulibus, summis sessilibus subintegris, paniculâ corymbosâ ramis erectis, capitulis pedicellatis, invol. squamis linearibus acuminatis dorso puberis disco ferè superantibus. — in insulâ Timor leg. Baudiniani collectores. Recept. nudum. Cor. rubentes, fœm. innumere, masc. paucæ subpubescentes sub oculo armato. Affinis *B. lactuæfoliæ*. (v. s. comm. à Mus. reg. Par.)

27. *B. TRIGONA*, tota glanduloso-pubera, caule herbaceo erecto infernè trigono superne tereti parçè ramosa, foliis petiolatis ovalibus acutis irreg. dentatis basi inciso-lyratis, summis subintegris acuminatis, paniculâ subramosâ foliosâ strictâ, invol. squamis linearibus acutis glabriusculis, recept. glabro, acheniis teretiusculis glabris. — in Indiâ or. prov. boreali-occid. legit cl. Royle. Valdè accedit ad *Bl. Timorensis* sed diversa videtur. Herba in siccis nigricans. An caulis cava tantùm trigonus? Fl. fœm. innum., masc. circ. 5 breves. (v. s. comm. à cl. Royle.)

28. *B. CICHORIIFOLIA* (DC. in Decaisne! nouv. ann. Mus. h. n. Par. 3. p. 410), caule herbaceo erecto ramoso tereti striato puberulo in ramis floridis villos glandulis intermixtis gerentibus; foliis glabriusculis, inf. lyrato-pinnatilobatis infernè attenuatis lobis inf. parvis oblongis super. majoribus subtriangularibus confluentibus, omnibus irreg. mucronato-dentatis, ful. summis oblongis sessilibus subintegris, paniculâ amplâ divaricatâ, ramis apice racemiferis, invol. squamis linearibus ciliatis disco subæqualibus. — in insulâ Timor colleg. exp. Baudiniana collect. Recept. alveolare fimbriifero-velutinum. Fl. flavidi, fœm. innum. masc. 8-10 glabri. (v. s. comm. à Mus. reg. Par.)

29. *B. SESSILEFLORA* (Decaisne nouv. ann. Mus. h. n. Par. 3. p. 410), ramis herbaceis erectis ramulisque virgatis striatis puberulis, foliis subtruncatis v. lanceolato-oblongis irregulariter serratis sessilibus subauriculatis puberulis, capitulis axillaribus plerumque sessilibus 3-5-glomeratis, invol. squamis linearibus puberulis, recept. glabro subtuberculato. — in insulâ Timor. Proximè ad *B. cichoriifoliam* accedit.

30. *B. CRUPINIFOLIA*, tota tenuiter pubescenti-villosa subcinerea caule erecto striato-sulcato subramoso, foliis infer. basi attenuatis lyratis pinnatipartitis v. lobis inf. ovali-oblongis, summo majore obovato, foliis omnibus acutè et irregulariter dentatis, summis oblongis acuminatis, ramis apice capitula dense thyroidea gerentibus in paniclem dispositis, invol. squamis linearibus acutis dorso pubescentibus disco paulò longioribus. — in Indiâ or. ad Nakis-Wari



februar. florentem legit cl. Hamilton. C. laciniata Ham. l in Wall. cat. n. 30 non Roxb. Recept. densè alveolare. Flores fœm. tenues glabri, stylo brev exserto, masc. circ. 18-20 lobis ciliato-villosis. Achenia subpubescentia. (v in h. Cur. Ind. or. nunc Soc. Liu. Lond.)

31. *B. GLANDULOSA* (DC. in Wight contrib. p. 14), tota brevissimè piloso-glandulosa, caule herbaceo erecto subtereti striato-nervoso, foliis caulinis brev petiolatis ovatis subacuminatis irregulariter dentatis basi lyrato-incis, meis basi cuneatis obovato-oblongis subdentatis, capitulis ad apices ramorum paniculam foliosam constituentibus, pedicellis ramosis glandulosis folio brevibus, invol. squamis linearibus acutis disco sequantibus. (1)? in Indiâ M. respat. propè Negapatam legit cl. Wight. Herba supernè verisim. vici Recept. convexum. Achenia glabra. (v. s. in h. Wight.)

32. *B. PHYLLOSTACHYA*, caule herbaceo erecto tereti basi glabrimaculo a cum ramis petiolisque pilis mollibus patulis hispidis, foliis petiolatis ly pubescentibus, lobis infer. paucis parvis oblongo-linearibus acutis aut tridentis, terminali maximo ovato-obtusobasi sublobato acutè dentato, panicula fertà, ramis brevibus folia obovata ovaliave acutè dentata capitulis intern gerentibus, invol. squamis linearibus acutis glabris disco longioribus. — Ind. orient. verisim. Maderaspatanâ. C. paniculata herb. Madr. ex Wall. n. 3090. comp. n. 200 non Willd. Flor. fœm. innumeri, masc. 5. (v. s. cor ab hon. Cur. merc. angl. Ind. or.)

33. *B. CINERASCENS*, tota cinereo-pubens, pube alterâ brevi confertâ et longis mollibus patulis intermixtis apice confertioribus, caule herbaceo erecto tereti ramoso, foliis ovali-oblongis lanceolatisve, infer. irregulariter dent petiolatis, mediis basi attenuatis, summis integerrimis, omnibus exaristis, paniculâ brevibus polycephalâ, ramis elongatis basi interruptè spicatas congesto-corymbosis, invol. squamis linearibus vix acutis dorso villosis d longioribus. (1) in Nepaliâ legit cl. Wallich. C. runcinata var. B. Wall. n. 3087. comp. 197. Affinis *B. runcinata* ex florum structurâ et numero foliis pilisque certè distincta. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

34. *B. RUNCINATA*, caule herbaceo erecto tereti ramoso pubescenti-villo foliis inf. in petiolum basi auriculato-dilatatum productis oblongis ram irreg. dentatis suprà puberulis subtus et novellis villosis, paniculâ el ramis capitula congesta interruptè spicata apice gerentibus, invol. n linearibus acutis dorso villosis disco longioribus. — in Ind. or. ad Hardwa Delhi legit cl. Wallich. *Conyza runcinata* Wall. cat. n. 3087. A. comp. A. Recept. alveolatum. Fl. fœm. tenues, stylo vix exserto, masc. 12-13 uia subpuberula. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

35. *B. ACURATA* (DC. in Decaisn. ! nouv. ann. Mus. h. n. Par. 3. p. 4) (supernè imprimis) tenuiter velutina pubescens, caule herbaceo ap plici nervulis striato-angulato, foliis petiolatis pinnatis irregulari lobatis, lobis oblongis acuminatis terminali majore ovali, omnibus profundis acutis inæqualibus incis, summis oblongis basi attenuatis, paniculâ contractâ terminali, invol. squamis linearibus pal acutis disco ferè longioribus. (1)? in insulâ Timor leger. Bandiniani tores. Recept. alveolare subhirsutum. Affinis *C. crepidifolia*. (v. Mus. reg. Par.)

β. foliis infimis lyrato-lobatis, cæteris ovatis aut ovato-lance et acutè dentatis. — in Timor cum priore. (v. s. comm. Par.)

36. *B. SONCHIFOLIA*, tota pubescenti-villosa subviscida, caule erecto striato, foliis infer. petiolatis lyratis, lobis inf. paucis oblongis nali maximo triangulari, omnibus grossè et inæqualiter dentatis dentatisve, paniculâ terminali elongatâ, pedunculis late longioribus invol. squamis linearibus acutis disco paulò longior diâ orient. ad Patna legit cl. Hamilton. *Conyza sonchifolia* H cat. n. 3085. comp. n. 195. Affinis *B. laciniata* sed recept. la

flor. fœm. stylo longius exserto prof. bifido, masc. 3-4 cor. brevius ciliatâ. Achænium hirtellum striatum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

37. *B. subcapitata*, tota cinereo-pubens apice et in circulis novellis magis villosa, caule herbaceo erecto tereti simplici unico aut ad basin submultiplici, foliis petiolatis ovatis inæqualiter et acutè dentatis ad basin incisâ fere lyrtatis, capitulis ad apicem caulis in spicam ovato-oblongam congestis, invol. squamis linearibus acutis florum longitudine. ① in Nepaliâ legit cl. Wallich! *Conyza capitata* Wall. cat. n. 3056. comp. n. 166. Herba vix pedalis. Folia æquipoll. longa 8-9 lin. lata. Flores pallidè flaventes, fœm. innumeri graciles stylis exsertis bifidis, masc. circ. 12 antheris non exsertis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

38. *B. Hamiltoni*, caulibus pluribus solitariisve herbaceis erectis simplicibus lamiginosis parce foliosis, foliis in parte caulium infer. approximatis sessilibus oblongis cartilagineo-dentatis suprâ glabriusculis subtus lanatis vix acutis, super. parvis ramosis, capitulis in apicem terminalem ovatam basi subinterruptam adgregatis, invol. squamis linearibus acutiusculis apice ciliatis subcariosis pallidè flavidis disco æqualibus. ② in Indiâ or. ad Kamaon, ad Sukonaghur, ad Koomari legit cl. Hamilton. *Conyza necessaria* Ham.! herb. Gnaphalium Hamiltoni Wall.! cat. n. 3038 comp. n. 48. Fl. fœm. tenues unum. stylo indiviso longè exserto, masc. 8-10. Achæn. glabra. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et à cl. Royle.)

\*\*\* *Senecionifloræ, nempe foliis dentatis incisive membranaceis subglabris, capitulis paniculatis, invol. glabris aristatis.*

39. *B. senecioides*, caule herbaceo erecto tereti striatulo brevissimè subpuberulo, foliis petiolatis ovalibus subacutis basi longè cuneatis distanter callosos-dentati-membranaceis in petiolo et nervo subhirsutis, paniculâ laxâ terminali subnuda, capitulis longè pedicellatis, invol. squamis linearibus acuminatis ciliatis disco subæqualibus. ① in Indiâ orient. loco dicto Nawabgunj legit cl. Hamilton. *Conyza senecioides* Hamilton in Wall.! n. 3102. comp. n. 212. Fl. flavi, fœm. tenues stylo bifido, masc. cylindrici circ. 15. Achænia glabra (v. s. in herb. hor. Cur. Ind. or. nunc Soc. Lin. Lond.)

40. *B. FLAVA*, tota glabrâ herbacea erectâ, caule tereti lævi ramoso, foliis infimis obovatis basi attenuatis subserratis obtusis, caulinis cordato-semiamplexicaulibus acuminatis grossè callososerratis, ramis floridis axillaribus racemum corymbosum confertum gerentibus in paniculam laxè dispositis, invol. squamis linearibus cuspidato-acuminatis disco circ. æqualibus. ② ? in Indiæ or. prov. Nepal, Silhet, Ava, Tavoy, ins. Penang, etc. *Conyza fasciculata* Wall.! cat. n. 3017. comp. n. 127. non n. 3094. comp. 204. *Senecio fasciculatus* et *Baccharis amplexicaulis* Wall.! herb. non Lam. *Erigeron obovatus* Spreng.! in litt. ad Wall.! *Conyza obliqua* Ham.! herb. *Erigeron falcatus* Don prod. 172? sed quid falcatus? Flores flavi, fœm. multiseriales tenuissimè truncati, stylo exserto bifido, masc. 10-12. Pappus albus. Ovarium glabrum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et à cl. Royle.)

♂. *subnuda*, caule simplici subaphyllo. Hab. cum var. α. et cum eâ comm. (v. s.)

41. *B. VIRENS* (DC. in Wight contrib. p. 14), caule herbaceo erecto tereti glabro ramoso, foliis rigidulis sessilibus utrinquè sparsè puberulis, caulinis elongatis basi cuneatis sessilibus trifidis, lobis lanceolatis mucronatis regulariter serratis, terminali longiore, ramis obovatis basi cuneatis parce serratis, paniculâ laxâ, pedicellis axillaribus folio longioribus 1-5-cephalis capitulis pedicellatis, invol. squamis int. linearibus disco longioribus mucronatis. — in Indiâ orient. legit cl. Russel, in arenosis maritimis ad Negapatam (Wight!). *Conyza virens* Wall.! cat. n. 3037. comp. n. 147. var. C. *Erigeron virens* Russ.! herb. Fl. purpurascens, fœm. innumeri, masculi 5. Achænia puberula. (v. s. in herb. hon. Cur. merc. angl. nunc Soc. Lin. Lond.)

42. *B. LAPSEMOIDES* (DC. l. c. p. 15), glabra, caule herbaceo erecto tereti ramoso, foliis petiolatis membranaceis, caulinis sinuato-repandis hinc inde mucronato-serrulatis, lobis mucronatis lateralibus subtriangularibus terminali ovato, rameis obovatis vix subserratis, paniculâ laxâ oligocephalâ, ramis lateralibus foliosis, capitulis pedicellatis, inv. squamis linearibus mucronatis disco paulò longioribus. (1) in Indiæ orient. sylvis secus ripas flum. Salween prov. Martabanis ad Trogia, Kogon, Moslimyue et Amherst leg. cl. Wallich, et Courtallum in Carnaticâ austr. Conyza virens A. B. D. Wall. ! cat. et herb. n. 3037. comp. 147. Flor. flavidi, fœm. innumeri, masc. 10-12. Invol. post anth. omnino reflexa. Rami floridi glabri aut pube glandulosâ minimè subviscosi. (v. s. com. ab hon. Cur. Ind. or. et in herb. Wight.)
43. *B. HYMENOPHYLLA*, grabriuscula, caule herbaceo erecto tereti ramoso, rami floridis glanduloso-puberis, foliis petiolatis membranaceis mucronibus exsertis sparsè subserratis, infimis longius petiolatis basi sublyrato-incisâ, super. ovatis ovalibusve, paniculâ laxâ, pedunculis folio longioribus, capitulis pedicellatis, invol. squamis linearibus flores non superantibus. (1) in Martabanis secus flumen Salween leg. cl. Wallich. Conyza hymenophylla Wall. ! cat. n. 3038. comp. n. 138. Valdè affinis *B. lapsanoidi*, sed folia minus divisa, s. masc. circ. 20 et fœm. pauciores. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
- ß. subpubescens*, foliis inferioribus ovalibus basi cuneatis, vix subrepandis nec basi sublyratis, ramis floridis pube parçâ non glandulosâ subpubescentibus. (1) in Martabanâ ad Kogon. (v. s. comm. com a.)
44. *B. PAUCIFOLIA*, caule herbaceo erecto tereti subsimplici undiquè minutissimè puberulo apice subviscido, foliis petiolatis obovato-oblongis basi cuneatis hinc inde exsertè serratis suprâ adultis glabris subtus et novellis utrinque molliter pubescenti-villosis paniculâ elongatâ subaphyllâ, invol. squamis linearibus acuminatis flores æquantibus. (1) in Indiæ orient. verisim. prov. Silhet. Conyza virens E. Wallich ! cat. et herb. n. 3037. comp. 147. end. A.-D. Affinis *B. lapsanoidi* et *hymenophyllæ*. Invol. demum omnino reflexi. Fl. flavidi, masc. 12-18. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
45. *B. NAPIFOLIA*, caule herbaceo erecto tereti basi subglabro apice pubescente subglanduloso, foliis pubescentibus petiolatis lyratis lobis dentatis lateralibus ovali-oblongis terminali subrotundo, paniculâ parvâ oligophyllâ, ramis parè foliosis, invol. squamis linearibus acuminatis florum longitudine. — in Indiâ orient. ad Tavoy. Conyza napifolia Wall. ! cat. n. 3075. comp. n. 185. Herba pedalis. Folia sesquipoll. longa. Flor. flavi, fœm. pauciseriales truncati, stylo exserto bifido, masc. centrales circ. 30. An fortè genericè distincta? (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
46. *B. MURALIS*, infernè glabra medio puberula apice glanduloso-pilosa, caule herbaceo erecto simplici tereti, foliis in petiolum angustatis lyrato-pinnatifidiis, lobis grossè dentato-serratis inf. utrinquè 2-3 triangularibus oblongioribus super. confluentibus ultimo majore ovato, paniculâ elongatâ laxâ, capitulis pedicellatis, invol. squamis linearibus acuminatis glabris subcoloratis flores paulò superantibus. (1) in Indiâ orient. ad Paingti (an fortè Paingkhandi dicitur Kamaon?). Conyza muralis Ham. ! in Wall. ! cat. n. 3077. comp. n. 111. Fol. infer. 5 poll. longa, 1 ½ poll. lata. Inv. et fl. purpurasc. Fl. fœm. innumeri stylo bifido exserto, masc. circ. 5-7 breviores. Pappus albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
47. *B. TRISULCA*, caule herbaceo erecto levi trianlco, foliis sessilibus hispidis dentatis, radicalibus cuneatis, caulinis oblongis amplexicanlibus, corymbis paniculatis oligocephalis, squamis invol. aristatis disco longioribus. (1) in Nepala. Erigeron trisulcum Don prod. fl. nep. 171. Conyza trisulca Ham. ex Don Pappus niveus. Flor. masc. pauci.
48. *B. MEMBRANACEA*, caule herbaceo erecto simplici tereti infernè glabro et trunco piloso, foliis membranaceis sparsè pilosis grossè serratis, super. obovatis acuminatis basi longè cuneatis in parte cuneatâ nunc integerrima ses-

inciso-lobata sublyrata, summis distantibus linearibus serratis, pedunculis axillaribus glanduloso-pilosis capitula subglomerata sessilia gerentibus interrupte spicatis in paniculam laxam dispositis, invol. squamis linearibus acuminatis disco aequalibus. ① in Ind. orient. ad Promē regni Avenais legit ocul. Wallich. Conyza membranacea Wall.! cat. n. 3019. comp. n. 129. Folia infer. panca, 7-8 poll. longa, 2-2½ poll. lata; summa pedunculis paulò breviora. Flor. fœm. pluriseriales tenuissimi circ. 30-40, masc. circ. 25-30. Pappus albus tenuissimus parvus. (v. s. com. ab hon. Cur. merc. angl. Ind. or.)

49. *B. viscosula*, caule herbaceo erecto tereti subsimplici apice puberulo-viscoso, foliis membranaceis denticulis exsertis callosis serratis obovatis basi cuneatis subpetiolatis sæpe lyrato-incisis tenuissimè utrinquè puberulis, paniculâ laxâ, ramis subnudis apice capitula pauca sessilia subspicata gerentibus, invol. squamis linearibus acuminatis flores æquantibus in lineâ mediâ extus puberulis. ① in Indiæ orient. regno Avenai propè Taong-Dong novembri legit cl. Wallich. Conyza viscosula Wall.! cat. n. 3081. comp. n. 191. Herba pedalis. Fol. 4 poll. longa 1-1½ lata. Capitula parvula 2 lin. lata. Flor. fœm. tenues truncati stylo brevissimo exserto, masc. 15-20. Pappus albus. (v. s. comm. ab hon. Cur. Ind. or.)

β. *hispidula*, caule a basi ad apicem patentim hispido, pilis supernè pluribus glanduloso-viscosis foliis subtus villosis-hirsutis, invol. squamis flores superantibus dorso puberulis apice longè ciliatis. ① cum priori eodem loco et die lecta à cl. Wallich et tamen fortè species propria? (v. s. comm. cum α.)

50. *B. Heyneana* (DC. in Wight contrib. p. 15), caule herbaceo erecto ramulosisimo subtilissimè striatulo et puberulo, foliis oblongis basi attenuatis apice obtusis hinc inde dentes exsertos acutos gerentibus utrinquè glabrisculis, infer. basi lyrato-suritis, paniculâ foliosâ laxâ, ramis folii longitudine apice subcorymbosis, invol. squamis linearibus acuminatis dorso subpuberulis disco paulò longioribus. — in Indiâ orient. legit cl. Heyne. Conyza Heyneana Wall.! cat. n. 3089. comp. n. 199. Con. aurita Heyne! herb. non Lin. f. Folia 2-3 poll. longa, pollicem lata, infima ignota. Fl. flavidi, fœm. innum. tenues stylo exserto, masc. circ. 12. Pappus albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et in herb. Wight.)

51. *B. cuneifolia*, pube brevissimâ subcabrida, caulibus herbaceis erectis teretibus striatis, foliis sessilibus cuneiformibus obtusis serratis exsertis acutis mucroniformibus distantibus sæpè reflexis, paniculâ oligocephalâ ramis erectis capitula pauca congesta apice gerentibus, invol. squamis linearibus acutis disco æquantibus, fl. masc. numero fœmineos superantibus. — in Indiæ prov. Martaban ad ripas flum. Saluen propè Phaoes legit cl. Wallich. Conyza cuneifolia Wall.! cat. n. 3201. comp. n. 211. non Lam. nec Rausch. Folia 12-15 lin. longa, 5-6 lata. Recept. nudum. Flores flavidi, fœm. pauciseriales, masc. circ. 40-50. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

52. *B. subsimplex*, caule imâ basi fruticuloso gracili glabro, ramis herbaceis teretibus elongatis subnudis pubescenti-villosis, foliis infer. obovatis petiolatis basi cuneatis rariter et acutè callosis-dentatis suprâ glabrisculis subtus pallidis pubescentibus, ramis minoribus petiolatis ad petiolum sæpè incisis, capitalis ad axillas supremas 2-3 brevè pedicellatis apicem interruptam fere constitinentibus, invol. squamis linearibus acuminatis dorso puberulis apice glabris arcuato-reflexis. — in Indiæ or. prov. Silhet. Conyza subsimplex Wall.! cat. n. 3003. comp. n. 113. C. paniculata Willd. sp. 3. p. 1920 ex Spreng. in litt. ad Wallich sed descr. non ritè quadrat. Cor. ochroleuca, fœm. tenuissimæ stylo simplici, masc. circ. 20. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

\*\*\*\* Fasciculiforme, nempe foliis dentatis, ramis elongatis capitula ad axillas foliorum floralium aggregata et interruptè spiciformia gerentibus.

53. *B. RACEMOSA*, caule herbaceo tereti villosa aut demum glabrato, foliis ellipticis acutis basi attenuatis petiolatis inaequaliter serratis utrinque operis villosis, capitulis ad axillas foliorum superior. dense congestis sessilibus in apice interruptè formam longè digestis, invol. squamis linearibus acuminatis dorso puberis demum stellato-patulis intus nitidis. ① in Indiâ orient. ad Noakote legit cl. Wallich. *Conyza racemosa* Wall.! cat. n. 3002. comp. n. 112. 808 Spreng. *C. stachyoides* Spreng. in litt. ad Wall. Cor. purpurea, fœm. immære stylo bifido, masc. 5-7. Pappus albidus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
54. *B. MACROSTACHYA*, tota dense hirsuta albida, caule herbaceo erecto tereti subsimplici, foliis lanceolatis utrinque acuminatis acutè serratis, superioribus subsessilibus ex axillis ramos abbreviatis foliolosos dense floridos gerentibus. paniculâ elongatâ interruptè subspicatâ confertâ fasciculatâ, invol. squamis linearibus acutis, ext. hirsutis, interioribus apice coloratis. — in Ind. orient. ad Noakote Nepalensium martio flor. legit cl. Wallich. *Conyza macrostachya* Wall.! cat. n. 3053. comp. n. 143. *Baccharis auriculata* Wall.! olim in herb. Flor. purpurei, fœm. innum. tenues, masc. 10-12. Achænia glabra. Rec. phnum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
55. *B. HIERACIFOLIA* (DC. in Wight contrib. p. 15), tota hirsuta, caule herbaceo erecto tereti simplici, foliis calloso-dentatis, infer. obovatis obtusis in petiolum attenuatis, super. ovalibus oblongisve acutis sessilibus aut semiplexicanibus, capitulis sessilibus confertis in thyrsum ovato-oblongum digestis, invol. squamis linearibus acuminatis glabriusculis disco longioribus. ① in Indiæ orient. prov. Nepaliâ (Ham.), Silhet (Wall.) et penins. Carnaticâ (Wight!). *Conyza hieracifolia* Spreng. syst. 3. p. 514. *Erigeron hieracifolius* Don prod. fl. nep. 172. *Conyza communis* Wall.! cat. et herb. n. 3018 comp. n. 128. Herba pedalis. Fol. subtus magis villosa-hirsuta. Invol. apice subpurpurascens. Flor. fœm. innumeri tenues stylo exserto indiviso, masc. 5. Ovaria pubescentia. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et in herb. Wight.)
56. *B. HOLOSERICA*, tota villis sericeis longis vestita, caule herbaceo erecto tereti apice hirsuto subsimplici, foliis oblongis basi attenuatis acutis profundè parè et acutè serratis adpressè sericeo-villosis, paniculâ elongatâ interruptâ. ramis baseos folia superantibus apice racemosis, sup. brevioribus congestè polycephalis, invol. squamis linearibus acuminatis puberis ciliatis disco fere longioribus. — in Indiæ or. prov. Martabanâ ad ripas flum. Saloen propè Phasor legit cl. Wall. *Conyza holosericea* Wall.! cat. n. 3103. comp. n. 213. *C. sericea* Wall.! cat. burm. n. 1892. Pars plant. inferior ignota. Flor. purpurascens. fœm. innumeri tenues, stylo exserto subsimplici, masc. 10-12 glabri, ant. exsertis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
57. *B. VIMINEA*, caule erecto ramoso, ramis elongatis teretibus pubescentibus. foliis lanceolato-oblongis basi attenuatis acuminatis grossè ad apicem et acutè serratis utrinque pubescentibus, ramulis floridis axillaribus folio brevioribus foliola et capitula 7-8 gerentibus, invol. squamis linearibus pubescentibus disci longitudine. — in insulâ Timor. Recept. alveolare glabrum. Flor. fœm. purpurei innumeri stylo exserto bifido, masc. 5-7 glabri. Achænia glabra. (v. s. comm. à Mus. Reg. Par.)
58. *B. FASCICULATA*, herbacea erecta ramosa tota cinereo-villosa, fol. infer. super. oblongo-lanceolatis basi attenuatis sessilibus superne rariter serratis acutis, capitulis sessilibus interruptè spicatis nempe ad axillas foliorum rami inferiorum solitariis superiorum 2-3 congestis, invol. squamis linearibus acuminatis flores æquantibus, ovario glabro, pappo 6-8-seto. — in Indiæ or. insulâ Timor et regione Burmannorum propè Amherst. *Conyza fasciculata* Wall.! cat. n. 3094. comp. n. 204. *C. glomerata* Wall.! cat. Burm. n. 1409. (v. s. comm. à Mus. Reg. Par.)

59. *B. GRACILIS*, infernè glabra supernè puberula, caule herbaceo striato erecto? ramoso, foliis oblongis basi attenuatis obtusis, inf...., mediis sinuato-incisis, summis suberrulatis acutis, paniculâ ramosisimâ laxâ, invol. squamis subvillosis linearibus acutis flores subaquantibus. — in Indiâ orient. legit cl. Heyne. *Conyza gracilis* Heyne! in Wall.! cat. n. 3064. comp. n. 174. Pars infer. plantæ ignota. Folia sub paniculâ ramis ferè pollicem longa, 3-4 lin. lata. Flores pallidè flaventes, fœm. innum. tennes, masc. circ. 12. Achænia minima teretiuscula. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
60. *B. PURPURA*, tota parçâ villosa, caule erecto herbaceo simplici tereti, foliis ovatis acutis serratis basi integerrimis attenuatis, ramis plurimis subnudis paniculatis dispositis capitula sessilia apicata subconferta gerentibus, invol. squamis linearibus acutis, exter. brevibus cinereo-virescentibus, intimis coloratis flores paulò superantibus. ① in montibus seu collibus Indiæ or. circa Rajmahal legit cl. Hamilton. *Conyza purpurea* Ham.! in Wall.! cat. n. 3076. comp. n. 186. Radix exilis subsimplex. Herba pedalis. Involucra purpurascens. Flor. fœm. innumeri tennes, masc. circ. 20. Pappus albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
61. *B. GLOMERATA* (DC. in Wight contrib. p. 15), tota villosa-subhirsuta cinerea, caule herbaceo erecto tereti ramoso, ramis in paniculam dispositis foliosis apice capitula sessilia glomerata interruptè apicata gerentibus, foliis lanceolatis serratis utrinquè acutis, invol. squamis multiserialibus, ext. minimis, intimis linearibus flores paulò superantibus, fl. masc. 30. ① in Indiâ orient. reg. Burmannorum ad Prome, Taong-Dong, Amherst, Moalmeyne legit cl. Wallich, in arenosis maritimis circa Madras et Negapatam cl. Wight. *Conyza glomerata* Wall.! cat. n. 3055. comp. n. 165 excl. litt. C. *Conyza argentea* Spreng. in litt. ad Wall. non alior. Flor. fœm. vix masc. numerosiores truncati, stigm. bifido purpureo, masc. 5-fidi antheris luteis exsertis. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
62. *B. ZAPFOCLADA*, infernè glabrata supernè cinereo-villosula, caule herbaceo erecto tereti ramoso, ramis virgatis paniculatis capitula sessilia solitaria aut apice glomerata interruptè apicata gerentibus, foliis glabriusculis obovato-oblongis basi longè cuneatis ferè petiolatis apice grossè et acutè serratis, invol. squamis linearibus acuminatis florum circ. longitudine, fl. masc. circ. 12-15. ① in Nepaliâ legit cl. Wallich. *Conyza glomerata* var. C. Wall.! cat. et herb. n. 3055. comp. n. 165. Valdè accedit ad *B. glomeratam* sed pars infer. ferè glabra, invol. squamæ pauciores, fl. masc. dimidiò numerosiores, antheræ incluse. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
63. *B. PERROTETIANA*, tota cinereo-velutina, caulibus herbaceis erectis subsimplicibus teretibus, foliis oblongis acutis serratis basi attenuatis, racemo spiciformi terminali oligocephalo, invol. squamis linearibus puberulis apice subcoloratis acutis disco paulò longioribus. — in Africa occid. arenosis locis inundatis propè flum. Senegal et inter fl. Gambie et peninsulam promontorii Viridis mense martio flor. legit cl. Perrottet. Flor. purpurei, fœm. innumeri, masc. 15-20. Achænia teretiuscula glabra. (v. s. comm. à cl. Perrottet.)
64. *B. SOLIDAGINOIDES*, caule herbaceo erecto tereti subsimplici molliter villosa, foliis lanceolatis sessilibus basi attenuatis suprâ pubescentibus subtus molliter villosa-canescens serratis, capitulis aggregatis in racemum interruptum subfoliosum subspicatum dispositis, squamis lineari-subulatis acutis villosis apice laxiusculis, flor. fœm. innumeris, masc. 4-5. ②? in Sierra-Leonâ. Gnaphalium solidaginoides Poir.! suppl. 2. p. 805. excl. patr. Species à Poiretto descr. in h. Desf. ex specim. quod ipse cl. et amico magistro olim communicavi, ex herb. L'Hér. orto. (v. s.)
- \*\*\*\*\* Oxyodontæ, nempe basi glabrata apice hirsuta, foliis obovatis dentes duros acutos subaristatos gerentibus, capitulis fasciculatis aut subpaniculatis.
65. *B. SPIRÆULOSA*, caule herbaceo tereti basi pilosiusculo apice ramisque sericeo-villosissimo, foliis infer. obovatis acutis longè cuneato-atenuatis sparsè pilosis

inæqualiter et grossè serrato-spinulosis, summis oblongo-linearibus utrinque sericeo-villosissimis æqualiter spinuloso-serratis, ramis floridis paniculatis folia longè superantibus apice capitula sessilia conferta spicata gerentibus. ①? in Indiâ orient. ad Promæ regni Avenais legit cl. Wallich. *Conyza sericea* Wall.! cat. et herb. n. 3021. comp. n. 131 non Ait. nec Bory. Folia inf. 6 poll. longa, 2 poll. lata. Rami florif. inter. 6 pollic. Flor. form. pluriser. terminati stylo indiviso, masc. circ. 20. Pappus albus tenuis deciduus parvus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

66. *B. BELANGERIANA*, caule herbaceo erecto à basi ramoso basi sparè hispido apice molliter villosa, foliis infimis obovatis subobtusis in petiolum brevem attenuatis, summis subsessilibus, omnibus argutè serratis suprè sparè pilosis subtùs densè sericeo-villosis, capitulis ad axillas superiores congestis fere interruptè spicatis, invol. squamis linearibus glabriusculis apice ciliatis. ①? ad limes Malabaricum legit cl. Belanger. Affinis *B. oxyodonta* sed erecta, multò major. Recept. angustum nudum. Flor. fœm. in genere pauci, masc. 8-10. Antheris exsertis. *Achænia teretia glabriuscula*. (v. s. comm. à cl. inv.)

67. *B. BRIANTHA* (DC. in Wight contrib. p. 15), caule herbaceo erecto tereti basi glabro apice ramoso et mollissimè hispido, foliis membranaceis cœulis sessilibus ovalibus acutè et exsertè apiculato-dentatis molliter pubescentibus, summis cordato-amplexicanalibus, pedunculis axill. et terminalibus 1-cephalis solitariis aut paucis aggregatis demum valdè elongatis, invol. squamis linearibus ciliatis acuminatis, cor. fœm. et masc. apice barbatis. ①? in Indiâ versim. Maderaspatanà legit cl. Wight. Herba pallidè viresca. Fol. inf. ignota cœulis 3 poll. longa, 15-16 lin. lata. Invol. demum reflexum. Rec. glabrum. Flor. masc. 25-30. (v. s. in h. Wight.)

68. *B. OXYODONTA* (DC. in Wight contrib. p. 15), herbacea villis longè sericeis laxè villosa, caulibus pluribus tenuibus simplicibus ramosive diffusis, foliis argutè et grossè serratis subtùs magis sericeo-villosis, infimis obovatis petiolatis, superioribus ellipticis subsessilibus, capitulis paucis ad apices ramorum ramoso-corymbosis subsessilibus, invol. squamis linearibus acuminatis disco paulò longioribus. ① in Indiâ orient. ad Promæ regni Avenais legit cl. Wallich, circa Pondichéry (Belanger!) et ad littora maria versus Negapatam frequens (Wight). *Conyza oxyodonta* Wall.! cat. n. 3015. comp. n. 125. *Erigeron graminosus* Russ.! herb. Folia infima 2-2½ poll. longa, 9-10 lin. lata, caulina 9-12 lin. longa, 4-5 lata. Flor. fœm. pluriseriales, masc. circ. 15-20. Pappus albidus parvus tenuis deciduus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

β. *tenera*, tota magis sericeo-villosa, foliis summis vix serratis, panicula laxiore polycephalâ. ① in Indiâ orient. ad Oude, Holasannj et in Nepaliâ leger. cl. Hamilton et Wallich. *Conyza tenera* B. et C. Wall.! cat. et herb. n. 3033. comp. n. 133. *Conyza bovina* Ham.! herb. *Erigeron cuspidatum* Spreng. in litt. ad Wall. (v. s. comm. cum α.)

γ. *glabrata*, tota glabriuscula laxa, capitulis distantibus longius pedicellatis. — in Indiæ prov. boreali-occid. *Conyza tenera* A. Wall.! cat. et herb. n. 3033. comp. n. 133. (comm. cum var. α et β à cl. Royle.)

\*\*\*\*\* Semivestitæ, nempe infernè glabræ supernè velutinæ, foliis oblongis acuminatis subintegris, pappo albo.

a. Receptaculo hirsuto.

69. *B. RIPARIA*, glabra, caule fruticoso scandente tereti, foliis breviter petiolatis ellipticis breviter acuminatis glanduloso-serratis rigidis subtùs subpallidioribus paniculæ oblongæ ramis velutinis folio longioribus 1-3-cephalis, capitalis pedicellatis, invol. squamis lanceolato-linearibus acutis florum disco paulò brevioribus. ½ in insulâ Javâ ad ripas fluminum legit cl. Blume. *Conyza riparia* Blume! Bijdr. 899 non Kunth. Flor. masc. circ. 20 lobis valdè hirsutis. Recept. hirsutum. (v. s. comm. ab ill. inv.)

70. *B. CHINENSIS*, caule herbaceo erecto tereti glabro apice cum pedunculis is-

volucrisque pubescenti-velutino, foliis breviter petiolatis ovalibus apice angustè acuminatis utrinquè glabris densibus distantibus callosis acutis exsertis hinc indè instructis, paniculâ laxâ, invol. squamis linearibus, exter. brevissimis, intimis multò longioribus, recept. villissimis, acheniis teretiusculis subcostatis puberulis. — in Chinâ (Linn.), in Javæ sylvis montosis (Blume!), et insulâ Penang (Wallich!). *Baccharis nitida* Wall. cat. et herb. n. 3043. comp. n. 133. *Conyza Chinensis* Linn. sp. 1208 (excl. Rumph. syn.). Blume! bijdr. 898. Pers. ench. n. 21 nec n. 16. *Eupatorium reflexum* Linn. ex Steud. Affinis *B. semivestitæ* nec ullo modo genericè separanda. Differt foliis etiâ superioribus petiolo donatis, paniculâ minùs velutinâ, invol. squamis acutioribus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

71. *B. SEMIVESTITA*, caule herbaceo tereti erecto apice cum ramis floridis densè velutino-villoso, foliis elliptico-oblongis acuminatis basi attenuatis subsessilibus grossè serratis utrinquè glabris, paniculâ terminalis ramis inferioribus distantibus, capitulis ad apicem ramulorum sessilibus densè congestis, invol. squamis linearibus acutis dorso pubescentibus flores sequantibus, recept. villissimis. in Nepaliâ legit cl. Wallich. *Conyza semivestita* Wall. cat. n. 2996 A. comp. n. 106 A. Folia 6 poll. longa, 2 poll. lata. Fl. fœm. innumeri, masc. pauci. Ovarium puberulum subcylindraceum. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

72. *B. PROCERA*, caule herbaceo erecto tereti paniculato-ramoso infernè glabro apice ramisque lanuginoso-villosis, foliis caulinis glabris sessilibus ovali-lanceolatis acuminatis subserratis, paniculâ amplâ divaricatâ, capitulis ad apicem ramulorum aggregatis, invol. squamis dorso pubescenti-hirsutis linearibus flores paulò superantibus, receptaculo villosis. — in Indiâ or. prov. Silhet. *Conyza procera* Wall. cat. n. 3050. comp. n. 140 non Desf. Folia 5 poll. longa, 2 poll. lata, infima ignota. Fl. purpurei, fœm. innumeri, masc. 12-15. *Achænia* puberula. Pappus albus. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

b. Receptaculo glabro.

73. *B. MYRIOCEPHALA*, caule herbaceo erecto tereti infernè cum foliis glabro supernè cum paniculæ ramis puberulo-hirtis, foliis oblongis acuminatis basi longè in petiolum attenuatis subserrulatis, paniculâ ramosissimâ subaphyllâ, capitulis secus ramulos lateralibus sessilibus et terminalibus, invol. squamis linearibus acuminatis recurvo-patentibus dorso puberulis disco paulò longioribus. — in Indiâ or. prov. Silhet legit amic. Wallich. *Conyza squarrosa* Wall. cat. et herb. n. 3025. comp. 135 non Linn. Herba elata spectabilis. Folia 8-12 poll. longa, 1-2 poll. lata. Rami inf. ex axillis orti villosi oligocephali, super. elongati ramosissimi. Flor. fœm. innumeri truncati stylo bifido, masc. 5-9. *Achænia* teretiuscula. Pappus pallidè rufescens. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

74. *B. SPECTABILIS*, ferè tota glabra ramulis subhirsutis, caule herbaceo erecto paniculato ramoso tereti, foliis utrinquè lævibus ovali-lanceolatis basi longè attenuatis apice acuminatis hinc indè subincisis remotè et calloso-serratis superè lucidis, paniculæ amplissimæ laxæ, ramis velutino-hirtellis lateralibus folio longioribus polyccephalis, invol. squamis linearibus acuminatis dorso glabriusculis apice ciliatis flores ferè superantibus, recept. alveolato glabro. — in peninsulâ Indiæ orient. legit cl. Wight (pl. exs. 2. 16). Media omninò inter *B. myriocephalam* et elongatam, sed paniculâ laxissimâ polyccephalâ ab utraque præter car. indic. diversa. Folia 8-9 poll. longa, 1½-2 lata. An var. *Bl. myriocephala*? sed capitula duplò majora quam *B. myriocephalæ*, invol. squamæ non squarrosæ. (v. s. comm. à cl. Wight.)

75. *B. ELONGATA*, basi glabriuscula supernè pubero-velutina subsericea, caule herbaceo erecto simplici tereti, foliis superè scabridis subtus pubescentibus lanceolatis calloso-serratis, inferioribus basi attenuatis et hinc indè subincisis, superioribus vix basi attenuatis apice acutioribus, paniculæ elongatæ ramis pluribus brevibus oligocephalis, invol. squamis linearibus acuminatis dorso



- puberulis apice ciliatis flores paulò superantibus demum rufescent. recept. alreolato glabro. — in India orient. insula Penang legit cl. Wallich. *Conyza densiflora* Wall. cat. n. 3078. comp. n. 188. Herba bipedalis. Flores albi, innumeri tenues stylo exserto apice bifido, masc. circ. 12. Pappus albus. *Achnania glabra* teretiscula. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
76. *B. OBOVATA*, caule herbaceo erecto tereti simplici molliter hirsuto, foliis obovatis acuminatis basi cuneatis apice serrulatis supra sparse pilosis subtus cinereo-villosis, pedunculis paucis molliter hispidescentibus longioribus, cephalis in paniculam laxam oligocephalam dispositis, invol. squamis linearibus acuminatis disco paulò longioribus, recept. glabro. ① in Nepalià inferiore legit cl. Wallich. *Conyza obovata* Wall. cat. n. 3079. comp. n. 138. Herba sesquipedalis. Folia 5 poll. longa, 2 poll. lata. Capitula diam. semipoll. Fl. parparascentes. fœm. innumeri truncati tenues stylo bifido, masc. circ. 2. Pappus albus deciduus. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)

\*\*\*\*\* *Macrophylla, nempe foliis ovalibus oblongis subnervigerrimis apice, capitulis paniculatis, pappo rufescente.*

77. *B. MACROPHYLLA*, caule suffruticoso, foliis ovato-ant oblongo-lanceolatis acuminatis basi in petiolum angustatis glanduloso-serratis supra puberulis subtus cinereo-tomentosis, paniculâ divaricatâ laxâ, ramis nudis apice subramosis, invol. squamis linearibus acuminatis glabrisculis flores squantibus. ③ in insula Java montanis ad rivulos legit cl. Blume. *Conyza macrophylla* Blume! hgh. Spk. Folia pedem et ultra longa, 5 poll. lata. Recept. alreolatum. Flor. fœm. innumeri stylo exserto bifido, masc. pauci. Pappus pallidè rufescent. *Achnania teretiscula* striata pallida subpubescentia. (v. s. comm. à cl. Blume.)
78. *B. AROMATICA*, caule herbaceo erecto tereti ramoso pube minimâ cæcatis subglandulosis præsertim supernè vestito, foliis ovatis acuminatis basi longè secus nervum petioliformem productis utrinquè puberulis, paniculâ laxâ polycephalâ, invol. squamis exter. brevibus subrecurvis dorso puberulis, inter elongato-linearibus acuminatis glabris flores paulò superantibus. — in Ind. or. inter vallem censu stricto Nepaliæ et Noakote legit cl. Wallich mæris ferè defloratam. *Conyza aromatica* Wall. cat. n. 3054. comp. n. 164. *Erigon Cochinchinense* Spreng. in litt. ad Wallich non Pers. Folia 5-6 pollic. long. 1-2 lata. Pappus rufus. Flor. fœm. innumeri filiformes, masc. circ. 12. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
79. *B. DENSIFLORA*, caule herbaceo erecto ramisque molliter lanuginosis, foliis elliptico-oblongis acuminatis basi attenuatis hinc indè acutè imbrico-dentatis supra scabridis subtus lanuginosis nonnullis basi sublyratis, paniculâ ramosissimâ confertissimâ, invol. squamis linearibus acuminatis glabrisculis corollas subquantibus, flor. fœm. innumeri, masc. circ. 5. — in India orient. ad Pundua legit Fr. de Silva. *Conyza aromatica* Wall. cat. n. 2997. comp. n. 107. *Conyza amygdalina* Spreng. in litt. non Lam. Folia ad paniculam ramosam lineari-lanceolata rariter serrata ramis breviora, canina infer. 6 poll. long. 2 poll. lata. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
80. *B. EXCISA*, caule herbaceo erecto tereti ramoso velutino-pubescente, foliis elliptico-oblongis dentatis basi incisè sublyratis supra puberulis subtus lanuginosis, paniculâ amplâ ramosissimâ, invol. squamis linearibus disco longioribus hirsutis subacutis falcatis recurvis. — in India Malesica insula Penang legit cl. Wallich. *Conyza excisa* Wall. cat. n. 3011. comp. n. 121. Affinis *B. densifloræ* sed panicula magis laxa. Fol. infer. basi longè attenuata. Fl. fœm. innumeri, masc. 5-6. Pappus rufus. (v. s. comm. ab hon. Car. merc. angl. Ind. or.)
81. *B. LONGIVOLIA*, caule herbaceo erecto tereti simplici infernè cum foliis glabro supernè cum panicula ramis pubero aut villosa, foliis oblongis utrinquè acuminatis serratis, paniculâ elongatâ polycephalâ, ramis subsimplicibus, cephalis lateralibus subsensilibus et terminalibus subdivisitis, invol. squamis linearibus acuminatis glabris flores squantibus. — in India orient. legit cl. Heyne. Co-

*nyza densiflora* Heyne in Wall! cat. n. 3026. comp. 126. Folia 8-9 poll. longa, 1½-2 poll. lata. Puncula pedem et ultra longa. Inv. squamis longiores 4 lin. Pappus rufescens. Achænia striato-angulata velutina. Fl. form. innumeri tenuissimi stylo bifido, masc. circ. 5. Cor. extus ad apicem parvè glandulosa. (v. s. comm. ab hon. Curia merc. angl. Ind. or.)

2. *B. SYLVATICA*, caule herbaceo simplici, foliis breviter petiolatis oblongo-lanceolatis utrinquè acuminatis suprà basin argutè serrulatis et ad costam infrà subvelutinis, racemis simplicibus paniculatis nutantibus, invol. squamis linearibus acutis exter. brevibus pubescentibus int. disco paulò longioribus. — in insulæ Javæ montanis Megamendung. Con. sylvatica Blume! bijdr. 898. Pappus rufescens. Flor. form. innumeri tenues stylo exserto bifido, masc. 6-7 lobis hirtellis anth. et interdum stylo exsertis. (v. s. comm. à cl. Blume et Belanger.)

3. *B. sessilifolia*, caule herbaceo erecto apice confertè puberulo, foliis sessilibus oblongo-lanceolatis utrinquè acuminatis supernè serratis discoloribus suprà scabris infrà rugosis et arachnoideo-puberis, racemis compositis laxè paniculatis, invol. squamis linearibus, exter. brevibus puberulis, int. glabris disci longitudine. — in ins. Javæ altis montis Burangrang. Conyza sessilifolia Blume! bijdr. p. 897. Pappus subrufescens. Fl. form. tenues stylo longè exserto bifido, masc. 5 antheris exsertis lobis hispidatis. (v. s. comm. à cl. Blume.)

\*\*\*\*\* Appendiculatæ, nempe foliis petiolatis quasi lyratis sem petiolis appendiceas vel lobos distinctos gerentibus.

4. *B. BALSAMIFERA*, caule basi suffruticoso, ramis teretibus lanato-villosis, foliis oblongo-aut elliptico-lanceolatis duplicato-dentatis rugoso-venosis suprà villosis subtus sericeo-lanatis, petiolis lobis linearì-lanceolatis appendiculatis, corymbo subpaniculato divaricato, invol. squamis linearibus acutis hirsutis. 5 7 in Indiâ orient. continentali et insulari, Luzoniâ, Javâ, Cochinchinâ. Conyza balsamifera Linn. sp. 1208. Blume! bijdr. 895. Wall! cat. et herb. n. 2998. comp. 108. Conyza vestita Wall! herb. 1824. Pluchea balsamifera Less. syn. 1831. p. 150. An Erigeron sericeum Retz obs. 5. p. 28? C. odorata Rumph. amb. 6. t. 24. f. 1. non Linn. Baccharis Salvia Lour. cochinch. 2. p. 603. Variat foliis suprà nunc villosissimis nunc subvillosis ferè glabratiss, dentibus inæqualibus aut æqualibus acutis subobtusisve, petiolis interdum nudis sæpius appendiculatis undè folia stricto sensu lyrata. Pappus rufescens. Antheræ caudatæ. Achæmium maturum glabrum striato-sulcatum (v. s.)

5. *B. GRANDIS*, caule herbaceo erecto angulato tenuiter pubescente, foliis ovalibus acuminatis suprà glabris subtus pubescentibus serratis petiolatis, petiolo lobatos 5-6 distantes lineares acutos gerentibus. — in Indiâ orient. ad Tavoy Burmannorum legit cl. Wallich sinè floribus. Conyza grandis Wall! cat. n. 3051. comp. 161. Conyza falcigera Wall! cat. Borm. n. 2048. Valdè affinis *B. balsamifera* et ex hac analogiâ ad Blumeam relata. Folia cum petiolo y poll. longa, 3 poll. lata. (v. s. sine fl.)

6. *B. APPENDICULATA*, caule suffruticoso, foliis oblongo-lanceolatis argutè serrulatis subtomentosis et subrugosis, petiolis appendiculatis, racemis compositis paniculatis. 5 in insulâ Javâ ad fluviorum ripas circa Buitenzorg. Conyza appendiculata Blume bijdr. p. 895 non Lam. Baccharis gratissima Blume! in litt. 1831? Florum non satis evolutorum struct. non examinare quivi.

## Ser. II. CAULOPTERÆ, nempe foliis secus caulem ramosve decurrentibus.

\* Alatæ, nempe alis integerrimis.

7. *B. VERONOIDES*, suffruticulosa tota densè velutino-hirsuta, foliis ovato-lanceolatis acutè et callosè denticulatis in alas integerrimas desinentibus, paniculæ terminalis ramis axillaribus 2-3-cephalis folium paulò superantibus, invol. squamis imbricatis linearibus acuminatis subseparatis, exter. dorso

hirsutulis, intimis glabris scariosis longioribus.  $\zeta$  in Indiâ or. *Vernonia Wightiana* Wall. cat. n. 2923. comp. 33 sed in h. Wight non reperi et ideo errorem quemdam timeo. Diversissima à *Conyzâ Wightianâ* quæ *Blomea nostra* Wightiana. Fl. purpurei, margin. innumeri feminei, centrales 10-12 masculi stylo incluso purisim. ovario abortivo. Achenium teres tenue. (v. s. in h. Car. Ind. or. nunc soc. Linn. Lond.)

88. B. *OLOPTERA*, tota pubescenti-subglandulosa, caule basi suffruticuloso, foliis oblongis rariter denticulatis acutis basi productis in alas longas latas integerrimas, capitulis ad apices ramulorum solitariis, invol. squamis multiserialibus basi erectis apice subulatis squarroso-reflexis.  $\zeta$  in Africâ ad flumen Combe mense martio legit cl. Perrottet, in campis recenter igne devastatis. Fl. fem. innumeri tenuissimè truncati, masc. exactè cylindrici. Anth. basi caudata ut vix ac ne vix subdentata. Achenium teres villosum. Recept. glabrum. Pappi flosculis longior. (v. s. comm. ab indef. Perrottet.)

89. B. *SALVIFOLIA*, herbacea simplex molliter villosa-tomentosa, foliis lanceolato-oblongis subintegerrimis acutis basi semiamplexicaulibus decurrentibus, ramis lateralibus 2-5-cephalis in paniculam elongatam dispositis, capitulis multiserialibus, invol. squamis ext. brevibus foliaceis squarrosis, intimis elongatis acuminatis glabris.  $\zeta$  in insulâ Bourboniâ ubi ab incolis dicitur *Saage*. *Conyza salvifolia* Bory voy. ial. afr. 3. p. 176. Folia trita odorata, loco Tabaci adhibetur. Rec. nudum. Fl. fem. innumeri filiform. stylo exs. masc. circ. 12-15 Achenia teretiuscula glabra. (v. s.)

90. B. *ALATA*, caule herbaceo erecto ramoso foliisque pube brevi subrufâ pubescenti-velutinis, foliis elliptico-oblongis dentatis secus caulem in alas integerrimas longè productis, pedunculis axillaribus mono-aut oligocephalis racemoso-paniculatis, capitulis suberectis, squamis ext. lanceolatis foliaceis squarrosis subpuberis, inter. linearibus scariosis flores æquantibus. — in Indiâ or. legit cl. Heyne. *Conyza alata* Roxb. cat. h. calc. p. 61. *Conyza alata* var. A. Wall. cat. n. 3039. comp. 149. *Erigeron alatum* Don prod. 171. *Vernonia alata* Heyne! herb. Fl. purpurei, masc. 10-12. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

$\beta$ . *cernua* (Wall. l cat. et herb. n. 3012. comp. n. 122 sub *Conyzâ*), caule herbaceo erecto ramoso foliisque pube brevi subrufâ pubescenti-velutinis, foliis oblongis acuminatis denticulatis secus caulem in alas integerrimas longè productis, pedunculis axillaribus mono-aut oligocephalis racemoso-paniculatis recurvis, capitulis cernuis, invol. squamis ext. lanceolatis foliaceis puberis, inter. longis nitidis scariosis apice recurvis demum persistentibus. — in Nepaliâ, ad Kamaon, Sirmore. Pubes fortè glandulosa. Invol. radicans Carliæ more. Fl. fem. innumeri stylo bifido brevè exserta, masc. 12-18. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

$\gamma$  *Natalensis*, alis integerrimis in parte caulis infer. subundulatis, capitulis cernuis, invol. multiserialibus magis squarrosis. — in Africâ austr.-orient. ad Port-Natal legit cl. Drege, u. 4996. An species propr.? Fl. masc. circ. 11 anth. ecaudatis. Cor. apice summo subscabris. (v. s.)

91. B. *GARIEPINA*, suffruticulosa ramosissima tota canescenti-villosa, foliis oblongo-linearibus integris subacutis in alas integras productis, capitulis pedicellatis subpaniculatis, invol. squamis lineari-subulatis rectis disco subaequalibus, intimis non scariosis.  $\zeta$  in Africâ austr. regione Gariepina legit. cl. Drege, n. 2722. Folia 8-9 lin. longa, 1'-2 lata. Fl. masc. 10. Anth. tenuissimæ candatæ? Achenia tenuia vix pubescentia. (v. s.)

•• Auritæ, nempe alis incisîs aut interruptis.

92. B. *PTARODONTA* (DC. in Wight contrib. p. 15), caule herbaceo tereti ramoso vix subpuberulo apice viscoso, foliis elliptico-oblongis glabris subserratis in alam profundè et acutè dentatam ferè incisam longè productis, ramis foliolis subpaniculatis apice mono-aut oligocephalis, pedicellis nudis, invol. squamis ext. oblongis foliaceis brevibus, intimis scariosis linearibus acutis flores pube superantibus. — in Indiâ or. littore arenoso maris Coromandeliani frequens

ex Wight! herb. nec ab eâ differre videntur specimina à cl. Bojer in ins. Madagascar ad urbem Tananarivou lecta; alæ caulina in iis variant nunc grossæ nunc minutissimè serratæ. *Conyza alata* B. Wall.! cat. n. 3039. comp. 149. *Ethulia divergens* et *Inula indica* Russ.! herb. Valdè accedit ad *B. alatum* sed decurientiis fol. inciso-dentatis faciliè differt. (v. s.)

3. *B. aurita* (DC. in Wight contrib. p. 16), hirsuto-villosa, canle erecto tereti ramoso, foliis lyrato-pinnatifidis, lobis oblongis acutis dentatis, terminali majore, infer. basi decurrentibus secus caulem 1-2-auritis, paniculâ terminali laxâ, capitulis pedicellatis erectis, invol. squamis lanceolatis acuminatis, ext. dorso hirsutis, inter. subscariosis disco paulò longioribus. ① in Indiæ or. peninsulâ (Lessch.!), frequens in agro Maderaspatano et Coromandeliano (Wight!), ad Surjaghora et Tukanaghur (Ham.!), in Pegu (Belang.!), etc. Con. aurita Lfou. suppl. 367? Roxb.! Ham.! in Wall. cat. n. 3069 et 3086. Ex Linn. pars infima planta glabra, sed mihi ignota. Recept. nudum. Flor. masc. 5-8. Achania glabriusc. Styli fl. fœm. exserti. (v. s. comm. ab bon. Car. merc. angl. Ind. or.)

β. *Ægyptiaca*, canle folisque hirsutis subsericeis, capitulis subminoribus. — in *Ægypto* ad Hermonthim legit cl. Sieber, et in Sennaar cl. Acerbi. *Erigeron Chinensis* Sieb.! fl. eg. exs. non Jacq. An *Conyza villosa* Willd. sp. 3. p. 1929? (v. s.)

γ? *Berteriana*, tota piloso-subhirsuta vix subcinerea, foliis infimis latè obovatis. ① Hanc stirpem ex insulâ Americanâ Sanctæ-Marthæ ad Balbium olim misit Bertero, sed ibi fortè culta aut introducta? An *B. auritæ* aut *B. Senegalensis* var.? (v. s.)

4. *B. Guineensis*, tota puberula subglandulosa, foliis oblongis acutis dentatis basi incisâ infernè secus caulem 1-2-auritis, paniculâ terminali laxâ, capitulis breviter pedicellatis erectis, invol. squamis lanceolatis acuminatis, ext. dorso puberulis, intim. subscariosis disco paulò longioribus. ① in Guineâ (Willd. Thonn.), in Senegaliæ arenosis apricis humidis (Perr.!). *Conyza Guineensis* Willd. sp. 3. p. 1930. *Erigeron stipulatum* Schum. guin. p. 385. *Conyza argentea* Perr.! et Lepr.! ined. Nimis *B. auritæ* affinis et fortè varietas? (v. s.)

β. *foliolosa*, squamis foliaceis linearibus innumeris secus pedicellos. — *Conyza Senegalensis* Sieb. fl. exs. n. 42. non Willd. Potius monstrum quàm varietas vera. (v. s.)

5. *B. Senegalensis*, tota puberula subglandulosa, foliis obovatis obtusis dentatis basi attenuatis et incisâ subdecurrenti-auritis, paniculâ confertâ terminali, capitulis pedicellatis, invol. squamis lanceolatis acuminatis, exter. brevibus puberulis, int. subscariosis disco vix longioribus. ① in Senegaliâ olim legit Roussillon, ad Gambiam recentius Don (h. Lindl.!). *Conyza Senegalensis* Willd. sp. 3. p. 1928. (v. s.)

CKXI. *PLUCHEA* Cass. bull. philom. 1817. p. 31. non Arrab. fl. flum. (1835). — *Pluchea* et *Chlenobolus* Cass. dict. 42. p. 1. — *Conyza* et *Baccharidis* sp. auct. — *Pluchea* sp. Less. syn. 207.

Capitulum multiflorum heterogamum, fl. ambitus multiserialibus imineis filiformibus truncatis aut 2-3-dentatis, centralibus paucis asculis aut subhermaphrodito-sterilibus 5-dentatis glabris. Recept. anum epaleaceum nudum aut hirsuto-fimbriiferum. Invol. pluririale imbricatum. Anthere basi bicaudatæ. Achenium erostre cylindricum aut sulcato-angulatum. Pappus uniserialis, setis filiformibus et scabris. — Herbæ aut suffrutices in Americâ, Asiâ imò Africâ ritibus calidis dispersi. Folia alterna, varia in speciebus diversis, sæpè bescenti-glandulosa. Capitula corymbosa pedicellata. — Genus multiforme, in posterum fortè dividendum, à *Conyza* et *Blumea* antheris udatis diversum. — Placus Lour. huc fortè revocandus, sed non iis notus.

Sect. 1. *STYLIMNUS*. — *Stylimnus* et *Gynema Rafin. journ. phys.* 1819. p. 100 et *ann. nat.* 1820. p. 15. — *Leptogyne Ell. sketch.* 2. (1824). p. 322. non *Less.*

Recept. nudum.

§. 1. *Foliis decurrentibus*. — *Species Americana et Gerontogea.*

1. *P. MACROCEPHALA*, herbacea erecta simplex pilosa, caule striato, foliis longè decurrentibus, infer. ovali-oblongis serratis basi longè attenuatis, super. lanceolatis linearibusve integerrimis, capitulis 7-8 in corymbum terminalem dispositis, pedicellis patentim hirsutis, involucris tomentoso-lanatis. — in Brasiliæ prov. Rio-Grande (h. imp. Bras. n. 747). Herba 1½-2-pedalis. Capit. 9-10 lin diam. Pappus copiosus sabrufescens. Anth. longè caudata. (v. s. in h. Mus. reg. Par.)
2. *P. QUIROC*, herbacea, foliis sessilibus longè in alas foliaceas decurrentibus oblongis lanceolatisve calloso-denticulatis, corymbo composito subfasciculato, invol. squamis lanceolatis subacutis. (1) in Brasiliæ campis arenoso-pulvis (Lund!), circa Bahiam (Blanch. Salzm.), Rio-Janeiro (Gaud.), in littore dicto Banda orientale (Bacle!), in prov. Rio-Grande, Mato-Grosso. Dicitur var *Quitoc* (Lund), *Quitocus* (Blanch.), *Quitoco* (h. Mus. Bras.). Flor. nov. dec. Gnaphalium suaveolens Arrab. fl. flum. 8. t. 100. Herba odore suavi aromatico, sapore amaro subcamphorato. Fl. invol. innumeri, marginales pallidi flavescentes, centrales violacei. Variat superficie glabrâ aut plus minùs pubescente, imò in vaginâ foliorum infer. canescenti-subvelutina; foliis oblongis lanceolatis aut subovatis acutis, rariùs obtusis, plus minùs dentatis; alis caulinis integrè aut dentatis. Ex alia caulinis accedit ad Pterocaulon sed ex habitu toto et caract. capituli certè Pluchea pertinet. (v. s.)
3. *P. SUBDECURVENS* (Cass. dict. 42. p. 4), caule fruticoso erecto tereti superius præsertim puberulo, foliis in alam subintegram obtusam brevem decurrentibus lanceolato-linearibus acutis à medio ad apicem serratis suprè scabrimulis subtus viscoso-punctulatis, corymbo terminali conferto polycerphalo, capitulis pedicellatis, invol. squamis lanceolatis acuminatis disco subæqualibus glaberrimis. § in Mexici salis inter Ixtla et Acamiatla (H. B. K.). *Conyza atrovirens* Desf. cat. h. par. 1815. p. 114. non Linn. nec Walt. *Conyza adnata* H. B. et Kunth nov. gen. am. 4. p. 74. *Baccharis adnata* Willd. enum. 870. Flor. lat. foem. pluriserialis truncati stylo exserto, masc. 5-10. Achenia glabra. (v. s. in h. Par.)
4. *P. OVALIS*, tota cinereo-puberula, caule suffruticoso ramoso tereti, foliis ovalibus ovatiave dentatis basi productis in alam inæqualiter dentatam, capitulis corymbosis. § in Senegaliâ olim leger. Geoffroy de Villeneuve, Roméille et Sieber, in siccis ad Promont. Viride recentius Perrottet. *Baccharis ovalis* Pan. ench. 2. p. 424. n. 14. *Baccharis ovata* Sieb. fl. seneg. exs. n. 61. Invol. pallidè squamæ inter. apice subobtusæ ciliatæ. (v. s.)

§. 2. *Antheris brevius caudatis, foliis sessilibus non decurrentibus aut petiolatis*. — *Species Gerontogea.*

5. *P. DIOSCORIDIS*, tota cinereo-puberula, caule herbaceo basi suffruticosius tereti apice ramoso, foliis sessilibus ovalibus acutis serratis basi attenuatis inciso-sublobulatis, lobulis stipulæ-formibus, capitulis corymbosis, invol. squamis acuminatis. § 2 in Egypto ad Alexandriam, Kahirem, Damiettam; in Nubiâ (Delil.) et in Syriâ? (J. Bauh.). *Baccharis Dioscoridis* Linn. amma. sc. 4. p. 289. *Conyza Dioscoridis* Rauw. it. t. 54. *Baccharis Egyptiaca* Forsk. (v. s. et v.)
6. *P. TOMENTOSA* (DC. in Wight contrib. p. 16), fruticosa, ramis teretibus multinerviis et foliis utrinquè pube brevi vix tomentosa griseo-velutina, foliis ovalibusve cordato-semiamplexicaulibus dentatis, pedunculis ex axillis

superioribus folio longioribus, capitula pedicellata conferta gerantibus, in paniculam confertam brevem dispositis, invol. squamis oblongo-linearibus acutis disco paulò brevioribus. § in Indiâ or. circa Mysore legit cl. Heyne. Conyza tomentosa Wall.! cat. n. 3031. comp. n. 141. Wight! herb. non Mill. nec Forst. Eupatorium Mysorense Heyne! herb. Folia 12-15 lin. longa, 9-10 lin. lata. Flor. fœm. innum. stylo vix exserto, masc. 5. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

7. *P. INDICA* (Less.! in Linnæâ 1831. p. 150), caule fruticoso tereti supernè corymboso ramoso, infernè glabro, ramulis pedicellis folisque novellis minutè pulverulentò-puberulis, foliis obovatis breviter petiolatis distanter et acutè serratis, corymbis ramos terminantibus confertis, invol. ovatis squamis ext. ovatis dorso puberulis, intimis linearibus acuminatis glabris. § in Ind. or. ad ripas Gangis, prope Chitagong et Pansang (Wall.!), in Luzoniâ (Cham.!), Javâ (Bl.!); Timor (h. Mus. Par.), Borneo? (h. Merat). Baccharis indica Linn. sp. 1205. Willd. sp. 3. p. 1916. Conyza indica Blume! herb. Conyza corymbosa Roxb. mss. Wall.! cat. et herb. n. 3009. comp. n. 119 non Mill. — Breyne. cent. t. 70. Flor. fœm. innumeri, masc. 3-5. Antheræ breviter caudatæ. (v. s.)

8. *P. WALLICHIANA*, caule fruticoso tereti ramoso foliis pedunculisque pube brevissimâ subpulverulentâ scabridis, foliis ovalibus obovatisve acutiusculis obtusiusve serrulatis integerrimisve sessilibus, corymbis terminalibus trichotomis, capitalis breviter pedicellatis ovali-oblongis, invol. squamis lanceolatis acutis florum longitudine. § in Indiâ orient. ad Srinaghur. Conyza sessilifolia Wall.! cat. et herb. n. 3029. comp. n. 139 non Blume. Folia 1-2 poll. longa, 6-10 lin. lata. Fl. fœm. innumeri tenues truncati stylo bifido, masc. circ. 5. Antheræ basi brevissimè caudatæ. Valdè habitu accedit ad Pluch. Indicam. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

9. *P. FOLIOLOSA*, glabra, caule fruticoso tereti ramoso, foliis ellipticis subacutis grossè et acutè paucidentatis breviter petiolatis, summis basi cuneatis sessilibus, corymbis terminalibus compositis confertis, invol. squamis inf. latè ovatis, super. linearibus acuminatis florum longitudine. § ad ripas fluminum Salween et Caluen prov. Martabanæ Ind. or. (Wall.!) et in Pegu (Belang.). Conyza foliolosa Wall.! cat. n. 3030. comp. n. 140 non Willd. Eupatorium foliolosum Wall. cat. Burm. n. 1565. Habitus Pluchæ Indicæ et Wallichianæ. Anth. brevissimè caudatæ. Flor. fœm. innumeri tenues stylo bifido, masc. circ. 5. Ovaria glaberrima. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

§. 3. *Antheris longius caudatis, foliis non decurrentibus sessilibus petiolatisve. — Species Americanæ.*

10. *P. OBLONGIFOLIA*, herbacea subfusco-pubescentia subglutinosa, foliis sessilibus basi obtusè truncatis oblongis acutis serratis venoso-reticulatis, corymbo terminali subcomposito, invol. squamis dorso puberis glandulosis lanceolatis acuminatis disco longioribus. ½ in Brasiliâ paludosis ad Rio-Janeiro et Bahia legit cl. Blanchet (n. 71. 72 et 1035!) et in prov. Sancti-Pauli cl. Lund! Affinis *P. bifronti*, sed notis indicatis faciliè distincta. (v. s.)

11. *P. BIFRONS*, herbacea subglutinosa, foliis amplexicaulibus cordatis ovali-lanceolatis serratis nervoso-reticulatis, corymbo composito conferto, invol. squamis lanceolatis acutis dorso puberis subglandulosis. ½ in humidis uliginosis et aggeribus Carolinæ. Conyza bifrons Linn. sp. 1207. Ell. sketch 2. p. 322. Conyza amplexicaulis Michx. fl. bor. am. 2. p. 126 non Willd. Baccharis viscosa Walt. fl. car. p. 202. Conyza uliginosa Pers. ench. 2. p. 427. — Pluken. alm. 1. 87. t. 4. Cor. sordidè purpurascens ut congeneres, nec pallidè flavæ ut vult Pursh. Per hiemem in horis matutinis, si cœlum serenum, fibræ cristallinæ pollicem longæ è basi caulis oriuntur! (Ell. l. c.). (v. s.)

12. *P. MARYLANDICA* (DC. non Cass.), herbacea minutè pubescens subglutinosa, foliis sessilibus lanceolatis acutis subserratis membranaceis uninerviis, corymbo composito fastigiato polycephalo subaphyllo, capitulis pedicellatis, invol.

- squamis lanceolatis acutis florum disco brevioribus. ① in humidis et innatis Amer. bor. à Noveboraco ad Carolinam (Pursh fl. 523), Nova-Cæsaræ (Torr.). Conyza Marylandica Michx. fl. 2. p. 126. Ell. sketch 2. p. 321. Baccharis foetida Walt. car. 202? Stylimus maritimus Raf. in litt. 1819. Sæpe in horti vagat sub nom. Erig. camphorati. (v. s.)
13. *P. PORTIDA*, herbacea glabriuscula, foliis petiolatis ovali-lanceolatis utrinque acuminatis grossè serratis penninerviis membranaceis, corymbo fastigiato subpaniculato, capitulis pedicellatis, invol. squamis lanceolatis acutis dorso subpubescentibus ciliatis. ② in Virginia (Linn.), in Kentucky (Raf.). Baccharis foetida Linn. sp. 1205. Gynema dentata Raf. in litt. 1819. Gynema vicina Raf. ann. nat. 1820. p. 15. Conyza marylandica Pursh fl. 523? Pluchea marylandica Cass. dict. 42. p. 2 non DC. Dill. elth. t. 89. f. 105. (v. s.)
14. *P. CAMPHORATA*, herbacea subpubescens, foliis petiolatis ovatis seu ovato-lanceolatis acutis subrepando-dentatis membranaceis penninerviis subtis granula resinosa gerentibus, corymbo composito fastigiato, capitulis pedicellatis, invol. squamis membranaceis ciliatis ovato-lanceolatis acutis dorso pubescenti-glandulosis. ① circa Bostoniam (Greene!), in Pennsylvania nec in Carolina et Georgia (Ell.). Erigeron camphoratum Linn. sp. 1212. Willd. sp. 3. p. 1960 Conyza camphorata Ell. sketch 2. p. 321 non Pursh? Pluchea petiolata Cass. dict. 42. p. 2. Gynema balsamica Raf. (v. s.)
15. *P. PURPURASCENS*, herbacea pubescens, foliis petiolatis lanceolato-ovatis acutis serratis subtomentosis, corymbo terminali composito, capitulis pedicellatis ovato-globosis, invol. squamis ext. ovatis ferrugineo-pubescentibus, int. ovato-lanceolatis patentibus. ① in subhumidis et depressis Jamaicae. Conyza purpurascens Swartz fl. ind. occ. 1342. — Sloan. hist. 1. p. 152. f. 1 mak. Brown. jam. 318. Affinis dicitur *P. odorata*, sed herbacea serratifolia, etc.
16. *P. GLABRATA*, herbacea glabra aut vix subpubescens, foliis petiolatis lanceolatis basi longè acuminatis apice acutis serratis penninerviis, corymbis compositis laxis, capitulis pedicellatis subrotundis, invol. squamis ovato-lanceolatis acutis. ① in Guadalupâ legit olim cl. Badier et recentius Perrotet ad Pointe à Pitre. An var. glabra *P. purpurascens*? An Conyza odorata Guadalupensis Spreng. syst. 3. p. 507 ad hanc aut ad *P. odorata* referenda? (v. s.)
17. *P. ODORATA* (Cass. dict. 42. p. 3), fruticosa pubescenti-velutina, foliis petiolatis ovalibus oblongisve utrinque attenuatis integerrimis subtis subcanescentibus, corymbo composito fastigiato polycephalo, capitulis pedicellatis, invol. squamis ext. ovatis dorso pubescentibus, inter. linearibus acuminatis scaros glabriusculis. ③ in America merid. (Linn.), Mexico ad Tampico de Tamasopo (Berl. pl. exs. n. 3) et in distr. Oaxacamo (Andr. pl. exs. n. 275), in Guadalupâ (Badier!). Conyza odorata Linn. sp. 1208. — Plum. ed. Barm. t. 97 (bi sear.), sed nunquam vidi folia inciso-dentata. Folia tacta odorem subnavem exhibent. Ab hac nullo modo differre videtur Conyza Carolinensis Jacq. ic. rar. 3. t. 585. coll. 2. p. 271. Willd. sp. 2. p. 271, in insulis Bahamensis spontanea, sed verisim. excl. habit. in Carolinâ et Floridâ. (v. v. c. et s. sp.)
18. *P. CORTESII*, suffruticosa ramosissima canescenti-hirta, foliis petiolatis oblongis acutis basi angustatis integerrimis canescentibus, corymbis terminalibus ramosis, capitulis breviter pedicellatis, invol. squamis imbricatis oblongo-lanceolatis apice subulatis. ③ prope villam Ferd. Cortesii et urbem Cærteran Mexicanorum. Conyza Cortesii H. B. et Kunth nov. gen. am. 4. p. 25. Folia 3 poll. longa, 10 lin. lata. Flor. disci circ. 20. Affinis dicitur Pluch. odorata: ad Plucheam ideò referenda? quod in specim. Kunthiano confirmavit occ. Decaisne.
19. *P. CHINGOYO*, fruticosa, ramis hirtis pubescentibus, foliis subrotundo-ellipticis utrinque obtusis integerrimis suprâ hirtis pubescentibus subtis mollioribus pubescentibus canescentibus, corymbis terminalibus ramosis, invol. hemisphaerici squamis glabris linearibus apice subulatis. ③ in locis planis prope urbem Santa Peruvianorum ubi dicitur *Chingoyo*. Conyza Chingoyo H. B. et Kunth

nov. gen. am. 4. p. 76. t. 328. Ex habitu ad Plucheam referenda videtur, cl. Decaisne reverà ex specim. Kunthiano hanc suspicionem confirmavit.

N. B. *Placus tomentosus* Lour. fl. coch. 2. p. 606. *Placus laevis* Lour. fl. coch. 2. p. 606. *Gynema balsamica* Raf. fl. lud. 63. *Gynema*? *argentea* Raf. fl. lud. 63 et *Gynema*? *parviflora* Raf. fl. lud. 64 sunt species fortè huic sectioni addendæ sed omninò ignotæ.

### Sect. II. HEBEPHORA.

Recept. hirsutum. Caules herbacei. Folia sessilia non decurrentia. — Species Asiaticæ, mediæ inter Plucheam veram et *Pterocaulon*, à priori recept. hirsuto, poster. invol. persistente et inflor. corymbosâ diversæ.

10. *P. HIRSUTA* (Less. in linnæâ 1831. p. 150), caule herbaceo erecto tereti hirsuto, foliis sessilibus lanceolatis basi acuminatis apice subærratis suprâ lucidis asperis subtus subhirsutis, capitulis spicato-glomerulatis, invol. cylindrici squamis linearibus acuminatis hirsuto-ciliatis, recept. hirsuto-fimbrillifero, corollis glabris, antheris longè caudatis. — in Chinâ (Torr. ex Linn.), in Cochinchinâ? (Lour.), in Luzoniâ (Cham. ex Less.). *Conyza hirsuta* Linn. sp. 1209. An plantæ omnes hic recensitæ omninò eadem?

11. *P. SCABRIDA*, suffruticosa, ramis teretibus confertè pulverulento-scabris, foliis subsessilibus basi attenuatis obovatis grossè et irregulariter dentatis utrinquè sparè pulverulento-scabris, corymbo terminali composito, capitulis ad apices ramorum sessilibus fasciculatis, invol. ovati squamis ext. ovalibus, inter. linearibus glabris apice subciliatis. 3 ad Luzoniam legit cl. Hænke. Proxima *P. hirsuta* Less. sed foliorum formâ et capitulorum dispositione diversæ. Recept. hirsutum. Cor. masc. glabræ apice tuberculatæ. (v. s. in h. Hænke à cl. de Sternberg miss.)

† *Species exclusa.*

12. *PYCNOSTACHYA* Less. = *Pterocaulon pycnostachyum*.

XXXII. *PTEROCAULON* Ell. sketch 2 (1824). p. 323. — *Chlænobolus* Cass. 49. (1827) p. 337. — *Pluchea* sp. Less. — *Conyza* sp. Lam.

Capitulum multitorum, fl. marginalibus fœmineis filiformibus dentatis pluriserialibus. Recept. tenuissimè fimbrilliferum hirsutum, arius nudum. Invol. oblongi squamæ imbricatæ pluriseriales adpressè ut apice summo vix squarrosæ caducissimæ! Antheræ basi caudas ubulatas gerentes subexsertæ. Achænia angulata adpressè pilosa. Pappi etæ filiformes pluriseriales inter se æquales involucri longiores. — Herbæ interdum basi suffruticosæ Americanæ. Rhizoma subtuberosum sæpè lignosum. Caulis tomentosus alis sex foliaceis ornatus. Folia alerna lanceolata, integra aut denticulata, suprâ glabra aut juniora arachnosa, subtus densè tomentosa, utrinquè producta in alas longas integras. Capitula sessilia glomerata in spicas thyrsosve spicatos digesta.

### Sect. I. CHLÆNOBOLUS.

Recept. hirsutum. Folia subtus, caules et pagina alarum cum inferiore folii contigua densè et cano-tomentosa.

1. *P. PYCNOSTACHYUM* (Ell. sketch 2. p. 324), caule herbaceo simplici, foliis lanceolatis subundulatis minutè et remotè denticulatis suprâ glabris, capitulis in spicam densam continuam confertis, invol. squamis extus sericeo-tomentosis apice squarrosis. 2/ in arenosis siccis campis et sylvis Carolinæ et Floridiæ (Pursh, excl. syn. Cav.). *Conyza pycnostachya* Michx. fl. 2. p. 126. *Gnaphalium undulatum* Walt. fl. car. 203. *Chlænobolus pycnostachyos* Cass. dict. 49. p. 338. *Pluchea pycnostachya* Less. Radix tuberosa vulgè *Black-Root* dicta ulcera vetera sanat. (v. s.)



2. *P. ALOPECUROIDES*, basi suffrutescens simplex, foliis ellipticis irregulariter et callosa-crenato-serratis suprâ junioribus arachnoideis adultis glabris, capitulis in thyrsam spiciformem densum aggregatis, infer. hinc inde interruptis, invol. squamis dense tomentosis apice acutis glabris squarrosis. ♂ in subæol. aridis Jamaica australis (Sw.), Guadalupe (Bert.) et Martinicæ (h. Daby!). *Conyza alopecuroides* var. *a.* Lam. dict. 2. p. 93. *Conyza alopecuroides* Swartz fl. ind. occ. 1344. *Chænobolus alopecuroides* Cass. dict. 49. p. 339. (v. s.)
- ♀. *polystachyum*, caule apice ramoso, spicis thyrsoides plurimis. ♂ in insul. Caribæis? (v. s. comm. à cl. Lambert.)
3. *P. SPICATUM*, suffrutescens simplex ramosumve, foliis oblongo-ant elliptico-lanceolatis acutis irregulariter dentatis subrugosis capitulis thyrsoides-spinosis nempe in glomerulos spicatis aggregatis, invol. squamis sericeo-tomentosis squarrosis, inter. subglabratia. ♂ in Brasiliâ circa Rio-Janeiro et Bahia. An forte *Conyza alopecuroides* Kunth prope Camanum lecta huc referenda?
- a. brachystachyum*, thyrsis spicato ovato denso continuo. ♂ frequens circa Rio-Janeiro et in prov. Sancti-Pauli (Lund!). *Conyza rugosa* Vahl synb. x. p. 71. *Chænobolus rugosus* Cass. dict. 49. p. 341. Habitus *P. alopecuroides* et forsan species duas conjungens. (v. s.)
- ♀. *cylindricum*, thyrsis cylindrico elongato continuo. ♂ in Amer. merid. (Lam. Cav.). *Conyza spicata* Lam. dict. 2. p. 93. Cav. ic. 1. t. 12. *Chænobolus spicatus* Cass. l. c. (v. s.)
4. *P. INTERRUPTUM*, suffrutescens, foliis oblongo-lanceolatis irregulariter dentatis, capitulis fasciculatis sessilibus, fasciculis in thyrsam spicatum cylindrico-elongatum interruptum digestis, invol. squamis linearibus, ext. dorso tomentosis subacutis, inter. elongatis acuminatis apice glabris. ♂ ♀ in Brasiliâ circa Bahiam (Salzm.), Rio-Janeiro (Gandich.), prov. Rio-Grande (h. Mus. Bras!)
- a. polystachyum*, ramis plurimis elongatis apice thyrsoides-spinosis. (v. s.)
- ♀. *monostachyum*, caulibus simplicibus apice thyrsoides-spinosis. An species propria ex inv. brevioribus et thyrsis minus interruptis. (v. s.)
5. *P. ANGSTROPHIUM*, herbaceum gracile ramosum, foliis linearibus acutis intergerimis, capitulis sessilibus in spicam subrotundam brevem aphyllam confertissimis, invol. squamis inter. linearibus acuminatis glabris, flor. hermaphrod. 2-3. ♀ in Brasiliæ prov. Sancti-Pauli. Folia sesquipoll. longa, 2 lin. lat. suprâ juniora araneosa, adulta glabra. Capitula 15-20 oblonga in glomerulum sphaericum conferta. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 431 miss.)
6. *P. VIRGATUM*, herbaceum simplex, foliis lineari-lanceolatis vix serratis acutis, capitulis glomeratis, glomerulis spicato-thyrsoides valde remotis, invol. squamis dense tomentosis mucrone acuto glabro. ♂ ♀? in insulis Caribæis nempe Jamaicâ (Sloan.), Portorico et Sancti-Thomasi (Bert.), nec in Carolinâ. *Conyza virgata* Lin. sp. 1206 Lam. dict. 2. p. 93. *Gnaphalium virgatum* Lin. amœn. 5. p. 405. *Chænobolus virgatus* Cass. dict. 49. p. 340. — Plum. ed. Burm. t. 98. f. 2. Sloan. hist. 1. p. 206. t. 152. f. 5. (v. s.)

## Sect. II. PTEROCAULOPSIS.

Recept. nudum nec hirsutum. Invol. squamæ subæquales vix imbricatæ glabræ et folia subtus non cano-tomentosa. — Herba basi suffruticosa foliis Pluchæ. inflorescentiâ Pterocaulis, fortè cum Monentelis sectione primâ genus constituens.

7. *P. POLYSTACHYUM*, suffruticosum erectum puberulo-hirtellum, foliis lanceolatis seu oblongo-lanceolatis acutis dentato-serratis reticulato-nervosis in alas foliosas dentatas aut subintegras secus caulem decurrentibus, ramis floridis plurimis paniculato-corymbois, capitulis oblongis sessilibus fasciculatis secus ramos thyrsoides-spinatis. Invol. squamis linearibus acuminatis glabris. ♂ in Brasiliæ prov. Rio-Grande. Cor. ferè Pluchæ, sed folia decurrentia et apices thy-

oidem Pterocauli. Fl. disci 2-3. fœm. innumeri, omnes longitudine involucri. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub. n. 861 miss.)

**CCKXIII. MONENTELES** Labill. *sert. caled. p. 43. Cass. dict. 53. p. 236.* — *Tessariae sp. Less. syn. 207.*

Capitulum pluriflorum, fl. marginalibus 2-3-seriatis fœmineis filiformibus apice subdentatis, disci 1-5 hermaphroditis seu styli abortu masculis subcylindricis 5-dentatis extus subglandulosis. Recept. nudum planum angustum. Involucri oblongi squamæ biseriatae persistentes subæquales. Antheræ disci caudis subulatis donatæ. Styl. disci et radii apice bifidus. Achænia oblonga subcompressa parçè villosa. Pappus 1-serialis, setis filiformibus imâ basi subconcretis vix invol. longioribus. — Herba Filagineæ ramosæ. Folia oblonga seu lanceolata denticulata, in alas secus caulem longas integerrimas producta. Capitula sessilia, ad axillas foliorum floralium fasciculata, in spicas interruptas disposita. — Genus ad Pterocaulon proximè caracteribus et habitu accedens, sed satis diversum videtur invol. persistente non imbricato, recept. nudo, floribus disci paucioribus imò solitariis, ovario subcompresso et pappo 1-seriali.

### §. 1. Floribus disci 3-5. — An potius Pterocauli sectio?

1. **M. PTEROCAULON**, basi suffruticosus totus cinereo-villosus, caule erecto ramosissimo, foliis oblongo-linearibus distanter denticulatis secus ramos in alas longas foliaceas integerrimas productis, capitulis ad axillas supremas et apices ramorum sessilibus congesto-fasciculatis. ♂ in agris circa urbem Tananarivon metrop. prov. Emirnae insul. Madagascar legit cl. Bojer. Baccharis Pterocaulon Bojer! in litt. 1833 cum icon. Habitus omninò Monentelæ. An fortè Conyza decurrens Lin. sp. 1206 eadem aut affinis? (v. s. comm. à cl. inv.)

### §. 2. Flore disci solitario.

\* *Capitulis in glomerulos spicatum dispositos aggregatis.*

2. **M. SPICATUS** (Labill.! *sert. nov. cal. 43. t. 43*), caulibus herbaceis dense tomentosis alatis, foliis longè decurrentibus oblongis serrulatis suprâ araneosis subtus dense lanuginosis, capitulis in spicas elongatas dispositis, squamis invol. omnibus acuminatis, exter. brevioribus lanatis. ♀ in Novâ-Caledoniâ (Labill.!) et in Indiâ montibus regni Avenais ad Taong-dong, Paghamew, alibi ad ripas fl. Irrawaddy (Wall.!). *M. spicatus* Cassini dict. 53. p. 237. Gnaphalium cylindrostachyum Wall.! cat. n. 3931. comp. n. 41. Gn. polycephalum Wall.! cat. hirm. n. 655. Recept. nudum. (v. s.)
3. **M. REDOLENS**, caule lignoso ramoso toto alato, foliis longè decurrentibus lanceolatis integerrimis suprâ pubescentibus subtus leviter cano-tomentosis, capitulis in spicas terminales glomeratis. ♂ in Novâ-Caledoniâ (Forst.) et in Lonzoniâ (Cham.) Gnaphalium redolens Forst. prod. n. 533. Conyza redolens Willd. sp. 3. p. 1951. Tessaria redolens Less. in linnæ 6. p. 151.

\*\* *Capitulis in glomerulum globosum dispositis.*

4. **M. GLOBIFRUTUS**, caulibus suffruticulosus, foliis lanceolatis acutis reticulato-nervosis utrinquè pubescenti-hirsutis in alas angustissimas productis, glomerulis globosis terminalibus, invol. squamis exter. brevibus hirsutis apice maculatis subtruncatis, inter. elongatis linearibus acutis glabris, flor. fœm. capituli cujusque 10-12. ♂ in paludosis sylvaticis ad fluvium Brisbane in sinu Moreton legit cl. A. Cunningham. Affinis *M. sphacelato*. (v. s. comm. à cl. Cunningham.)
5. **M. INTERMEDIUS**, caulibus suffruticulosus hirsutus, foliis elliptico-lanceolatis

mucronato-acutis reticulato-venosis utrinque hirsutis in alas latiusculas innigerrimas longè productis, glomerulis terminalibus globosis, invol. squamis ext. hirsutis apice ustulatis, inter. elongatis glabris, floribus fœm. circ. 10. 5 in arenosis littoribus ad Caput-Cleveland Novæ-Cambriz aust. inter tropicos. Species media inter *M. globiferum* et *M. sphacelatum*, à priori alis caulinis dupli triplove latioribus et longioribus, superficie magis hirsutâ, foliis bracteis majoribus, à sequente foliis acutis et fl. fœmineis paucioribus diversa. (v. s. comm. à cl. Cunningham.)

6. *M. SPHACELATUS* (Labill. sert. n. cal. 43. t. 44), caulibus herbaceis alatis, foliis longè decurrentibus oblongis obtusis, capitulis in glomerulos laterales terminalesve globosos aggregatis, invol. squamis exterioribus ad apicem ustulatis. 2? in Novâ-Caledoniâ. Monenteles sphacelatus Cass. dict. 53. p. 23. Flos centr. ex. Cass. fertilis, fl. fœm. circ. 25. Recept. pilos lanatos paucos gerens.
7. *M? SPHERANTHOIDES*, caule fruticuloso glabro, foliis lineari-lanceolatis elongatis margine subrevolutis rugosis penninerviis denticulatis in alas dentatis decurrentibus glabrisculis, pedunculis pubescenti-hirsutis subcæbris, glomerulis globosis terminalibus. 3 in insulâ Enderby Archipelagi Dampieriani ad oram bor. occid. Australiæ legit cl. A. Cunningham. *Sphæranthoides* Cunningh. in litt. Frondescentia ferè *Sphæranthi*, capitulum *Monetellis*, sed in specim. equidem malè conservato achænia omninò calva! An pappus deciduus, an verè nullus? Invol. squamæ ext. breves hirsutæ, inter. elongatæ glabræ. Species distinctissima! Genus dubium! (v. s. comm. à cl. peregr.)

CCXXIV. *TESSARIA* Ruiz et Pavon prod. p. 112. t. 24. syst. p. 213 Cass. dict. 53. p. 233. — *Tessariz* sp. et *Gynheteria* Less. syn. 207 et 208. — *Gynheteria* Willd. mag. nat. Berl. 1807. p. 140. — *Gynheteria* Spreng. neu. entd. 2. p. 135.

Capitulum multiflorum, fl. marginalibus pluriserialibus tenuissimis truncatis seu 2-3-dentatis fœmineis, centralibus 1-7 majoribus hermaphroditis aut styli abortu masculis, cor. apice dilatata 5-fida. Recept. tenuiter fimbriatiferum hirsutum. Invol. oblongi squamæ imbricatæ scariosæ, ext. breves persistentes ciliatæ, inter. elongatæ acutæ decedæ subintegerrimæ. Antheræ basi brevissimè caudatæ. Stylus disci indivisus undiquè subhispidulus. Fl. fœm. exsertus glaber bifidus. Achænia brevia teretiuscula glabra. Pappus 1-serialis, setis filiformibus. — Frutices Americæ australis incolæ, foliis ramulisque adpressè cano-villosis. Folia alterna ovali-oblonga aut lanceolata integra aut in iisdem indiv. plus minùs dentata. Corymbi terminales ramosi. Cor disci purpureæ, fœmineæ pallidæ.

### §. 1. *Flore hermaphrodito-sterili unico centrali.*

1. *T. LEGITIMA*, foliis ramulisque hirsuto-canescens ovali- aut subobovatis oblongis vix acutis, invol. squamis ext. ciliis subhirsutis fimbriatis. 3 in Peruvia ubi dicitur *Paxaro-Bobo*. *Tessaria* Cass. dict. 53. p. 233. Variat foliis integerrimis aut sinuato-subdentatis: ergò certè videtur *T. integrifolia* R. et Pav. syst. p. 213 et fortè etiam *T. dentata* R. et Pav. p. 214. *Gynheteria* incana Willd. mag. nat. berl. 1807. p. 140? Spreng. neu. entd. 2. p. 135. t. 1 f. 5-11 ex icone, sed excl. patr. et descr. part. (v. s. ex itin. Dombey, Nec et Gaudichaud.)
2. *T. MUCRONATA*, ramis glabrisculis, foliis adpressè sericeo-incanis ellipticis basi mucronato-acutis apice mucronato-acutis, aliis integerrimis, aliis paucidentatis, invol. squamis glabris margine vix ac ne vix subciliatis. 3 in 5 ad Baranquilla Novo-Granatensium propè ostia fl. Magdalenz legit cl. Bertero. *Gynheteria* incana è Magdalena Spreng. syst. 3. p. 482 (excl. syn. et aliis habit.) ex specim. ab ipso auct. ad Balbisiu misso! An fortè *Cavendishii*?

*riparia* Magdalenensis H. B. et Kunth nov. gen. am. p. 4. 76? (ex quâ cl. Cassini genus suum *Phalacrotesum* dict. 53. p. 235. 57. p. 339. 60. p. 586 constituit) sed certè nostræ flos centralis papposus nec nudus. (v. s. comm. à cl. Balbis et Bertero.)

§. 2. *Floribus disci 3-8 hermaphroditis seu abortu masculis.*

3. T. *AMBIGUA*, ramulis foliisque adpressissimè pulverulento-canis, foliis lanceolatis subobtusis basi in petiolum attenuatis sinuato-paucidentatis, invol. squamis exter. ovato-rotundis ciliatis, inter. linearibus scabris glabris, disco 3-floro. § in Americâ merid. fortè in Peruvîâ. Habeo ex itin. Nezaro sub nom. vernaculo *Pajro-Bovo*, vix à nom. primæ generis speciei distincto.
4. T. *ABSINTHIODES*, ramis foliisque adpressissimè canis subargenteis, foliis lanceolatis utrinquè acuminatis, nunc integerrimis, nunc dentibus grossis acutis hinc inde incisus, subtus nervoso-reticulatis, invol. squamis glabris acutis, disco 7-8 floro. § in Chili ad Valparaíso, Coquimbo, et Conceptionem secus flumina. Vulgò ex Bert. dicitur *Bréa*. Baccharis absinthioides Hook. et Arn. bot. Beech. p. 57. Baccharis banksiaefolia Bert. herb. Gynheteria incana Chilensis Spreng. syst. 3. p. 482. (v. s. comm. à cl. Bertero, D'Urville, Gaudich.)

† *Species exclusa.*

T. *REDOLENS* Less. = *Monenteles redolens*.

CCXXV. *MONARRHENUS* Cass. bull. philom. 1817. febr. dict. 32. p. 433. 53. p. 236. — *Mahometa* DC. 1802. ms. — *Conyzæ* sp. Lam. — *Tessaræ* sp. Less.

Capitulum pluriflorum heterogamum, fl. omnibus tubulosis, ambitus plurimis subbiserialibus filiformibus 3-4-lobis fœmineis, disci solitario 5-dentato hermaphrodito aut ovarii abortione masculo. Recept. nudum planum parvum. Invol. ferè campanulatum floribus paulò brevius imbricatum, squamis oblongis obtusis apice summo subtomentosis. Cor. disci extus subglandulosa. Antheræ breviter caudatæ longè appendiculatæ. Stylus disci ramis inclusis brevibus, fl. fœm. longis exsertis filiformibus. Achænia fl. fœm. oblonga cylindracea supernè subhispidula callo basilari crasso. Pappus setis 1-serialibus scabris. — Frutices Mauritanii. Folia alterna linearia aut sublanceolata integerrima, suprâ scabrida, subtus tomentosa aut sericea. Corymbus compositus. Capitula ad apices pedunculorum conferta subsessilia. Flores latei. — Genus affine Tessariæ et Monenteli, ob flor. ♂ solitarium inter plurimos ♀ olim à me Mahometa dictum.

1. M. *SALICIFOLIUS* (Cass. l. c.), foliis breviter petiolatis lanceolato-linearibus subtus sericeo-tomentosis penninerviis. § in insulis Mauritiî ad montem Pollicis, et Borboniæ ad montem Bernica (Bout.). *Conyza salicifolia* var. *α*. Lam. dict. 2. p. 89. Bory! voy. isl. afr. 2. p. 28. Folia omninò Salicis viminalis, 2-3 poll. longa, 4 lin. lata. Vulgò *Bois de Senil* et *Bois de Chenille* dicitur. (v. s.)
2. M. *PINIFOLIUS* (Cass. ll. cc.), foliis angustè linearibus sessilibus margine subrevolutis uninerviis subtus tomentosis. § in insulâ Borboniæ. *Conyza salicifolia* var. *β*. Lam. dict. 2. p. 89. *Conyza salicifolia angustifolia* Pers. ench. 2. p. 428. Vulgò *Bois de Senil de Bourbon*. Folia 2-3 poll. longa, semi-lin. lata. Priori valde affinis sed distinctus videtur et nulla specimina intermedia observantur. (v. s.)
3. M. *AUFESCENS*, foliis sessilibus lineari-lanceolatis planis penninerviis utrinquè ramulisque pube rufâ minimâ ferrugineis integerrimis sat hinc inde argutè et minutissimè serratis. § in insulis Mauritanis nempe in insulâ Borboniæ (Comm.) in insulâ Mauritiî (Bout. in h. Lindl.). *Conyza heliotropifolia* Lam. dict. 2. p. 90? (v. s. in h. Lindl.)

CCXXVI. CYLINDROCLINE Cass. bull. philom. 1817. jan. dict. 12. p. 318. *Las. syn.* 207. — *Lepidopogon Tausch in florâ 1829. p. 2. p. 37. et 1831. p. 1. p. 224.*

Capitulum pluriflorum heterogamum, flor. omnibus tubulosis, marginalibus in parte recept. spiciformi seu inter involucri squamas sitis pluriserialibus foemineis filiformibus, disci in parte recept. terminali sitis paucis abortu styli subsimplicis masculis. Recept. totum bracteolatum et spiciforme apice subconcauum. Invol. oblongum, squamis secus pedicelli apicem spicæ more dispositis imbricatis oblongis, infer. sterilibus apice ad dorsum tomentosis, superioribus in axilla floriferis fimbriato-ciliatis. Achænium oblongo-cylindraceum basi attenuatum subarcuatum 8-costatum pilosiusculum. Pappus 1-serialis, paleis ex corneis pectinato-barbellatâ. — Fruticulus Mauritanus ramis crassis basi glabris inter folia densè et longè sericeo-hirsutis. Folia ad apices ramorum conferta rosulata integra ovalia basi attenuata et vix idè subspathularia juniora, suprâ et demùm ad basin sericeo-hirsuta, ad apicem breviter strigilloso-scabra, subtus albo-tomentosa ad nervos et basin sericeo-hirsuta. Corymbus compositus confertus, ramis longè sericeo-hirsutis aphyllis folia vix æquantibus. — Est quasi *Evax arborescens*!

1. C. COMMERSONI (Cass. ll. cc.). § in insulâ Mauritiâ montibus propè monte Pollicis et *Piton de la rivière noire* dictis (Boj.!). *Lepidopogon gnaphaloides* Tausch in florâ 1831. *Conyza hirsuta* Commers. herb. *Conyza Sieb.*! fl. maur. exs. 2. n. 334. *Lepidopogon* Ponz. Tausch in florâ 1829. c. 2. p. 37 et 1831. c. 1. p. 224. Flores cœrulei! Species valdè singularis identitatem squamarum involucri et palearum receptaculi optimè demonstrans! (v. s.)

CCXXVII EVAX Gærtn. fr. 2. p. 393. t. 165. *Lam. ill.* t. 694. *Pers. ench.* n. 1842. — *Filago Willd. sp.* 3. p. 2387. *Cass. dict.* 17. p. 2. — *Gnaphalium Vaill. act. ac. Par.* 1719. p. 314. — *Filaginis* sp. *Linn.* — *Micropis* sp. *Desf.*

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, radi multiseriatis foemineis gracilibus, disci paucis masculis 4-crenatis. Receptaculum elongatum subulatum, apice inter fl. masc. nudum, paleolas scarioras acuminatas inter. fl. foem. gerens. Invol. 1-2-seriale, squamis adpressis scariosis ovalibus aristato-acuminatis. Achænia (radii) obcompressa obovalia glabra calva. — Herba Europæ australis et Orientis indigenæ, minimæ, subcaules, tomento plus minus copioso canescentes. Folia alterna circa capitulorum glomerulum rosulata. Capitula plurima conferta subumbellata, medio majore.

#### Sect. 1. EUEVAX.

Invol. squamæ et recept. paleæ planæ scariosæ abruptè aristato-acuminatæ. Capitula densè umbellata, bracteis foliaceis inter se nullis, foliis floralibus circa umbellam radiantibus.

1. E. PYGMÆA (Pers. ench. 2. p. 422), foliis floralibus obovatis obtusis. In apricis et rupestribus maritimis Europæ mediterraneæ. *Lam. ill.* t. 694. f. 1. *Filago pygmæa* Linn. sp. 1311. *Filago acutis* All. ped. n. 620. *Gnaphalium pygmæum* Lam. dict. 2. p. 761. *Evax umbellata* Gærtn. l. c. *Micropis pygmæa* Desf. atl. 2. p. 307. — Barr. in. t. 127. n. 14. Variat caule simplicissimo vel pygmæo subnullo et caule à basi ramoso multicapiti longiore ferè pollicari. (v. et s.)
2. E. EXIGUA, foliis floralibus oblongis acutis vix glomerulo longioribus, inter

- squamis exter. dorso lanatis. ① in apricis et rupestribus Hispaniæ circa Madridum (Lag.), in Siciliâ meridionali (Guss.), in archipelagi insulis (Sibth.), in rupibus vulcanicis novæ insulæ Camini prope Theram (D'Urv.), et in agro Byzantino (Cast. !); etiam habeo inter plantas itin. Brousson. sed locus non indistinctus. *Filago* exigua Sibth. fl. græc. t. 922. ined. ex prod. a. p. 207. *Micropus* exiguus D'Urv. enum. p. 109. n. 802. *Filago pygmaea* Cav. ic. 1. p. 23. t. 36. non Linn. Variat ut prior caule simplici humilimo et à basi multicauli inèd ramosissimo 2-4 poll. longo. Folia valdè cano-lanata. Omnino media inter priorem et sequentem. (v. s.)
3. *E. ASTERICIFLORA* (Pers. ench. 2. p. 422), foliis floralibus oblongis acutis radiantibus capitulorum glomerulo multò longioribus. ① in apricis et rupestribus Hispaniæ (Lam. ex Vahl's specim.), Sardinia (Moris!), Sicilia (Guss.), Calabria (Duf.), et Orientis (Labill.) Lam. ill. t. 694. f. 2. *Gnaphalium astericiflorum* Lam. dict. 2. p. 760. — Moris. ox. s. 7. t. 12 f. 5. Variat etiam caule simplici 2-poll. et ramoso, sæpius ad apicem ramosiore, 3-4 poll. longo. Folia glomerulorum circumdantia 9-10 lin. longa. (v. s.)

Sect. II. PSEUDEVAX.

Invol. squamæ exter. lanatæ, inter. et recept. palææ oblongæ chartaceæ obtusæ. Capitula sparsa aut congesta, bracteis foliaceis intermixtis et foliis floralibus irregulariter circa glomerulum dispositis. — Habitus *Filaginis*.

4. *E. DISCOLOR*, foliis obovato-lanceolatis, floralibus vix involucrium superantibus, capitulis congestis lanâ floccosâ basi circumdatis. ① in Sicilia montibus Nebrodensibus (Guss.). *Filago discolor* Guss. in litt. 1831 et 1834. Herba 1-2-poll. à basi ramosa. (v. s. comm. à cl. Guss.)
5. *E. MAREOTICA*, foliis lineari-oblongis, floralibus vix capitulo longioribus, capitulis axillaribus unilateralibus imbricatis. ① prope Alexandriam ad lacum Mareotidem. *Filago Mareotica* Delil. fl. eg. 130. t. 47. f. 2. *Micropus Mareoticus* Sprung. syst. 3. p. 499. Habitus serè *Filaginis montanae* sed caract. omnino *Evæci*.
6. *E. PROLIFERA* (Nutt. in litt. 1825), caule solitario erecto apice capitula inter folia glomerata gerente, ramis subglomerulo ortis ipsis apice capitula glomerulata gerentibus, foliis lineari-oblongis, floralibus conferto-subverticillatis capitula superantibus ① in America bor. ad Red-River detexit cl. Nuttall. Herba stricta 2  $\frac{1}{2}$ -3 poll. longa tota villosa-canescens. (v. s. comm. à cl. Nutt.)
7. *E. MULTICAULIS*, caulibus à collo plurimis subdiffusis erectisculis ramosis, foliis oblongis obtusis laxè ramisque araneoso-lanatis, summis circa capitula lanata glomerata conferto-verticillatis, invol. squamis obtusis membranaceis. ① in Mexico circa lacum Sancti-Nicolai in sinu Spiritûs-Sancti et mi fallor in campis prov. Texas ad Fernando de Bexar legit cl. Berlandier (pl. exa. n. 1958 et 2109). Herba 4-6 poll. longa. (v. s.)

† *Species dubia.*

- E. VERA* (Raf. herb. p. 24), in Texas et Louisiana. An huic generi adnominanda?

†† *Species exclusæ.*

- E. ERICOIDES* Schrank = *Eolopes tenuifolia* et *Helichrysum teretifolium*.  
*E. INVOLUCRATA* Schrank = *Petalacte coronata*.  
*E. INDICA* Ham. = *Filago pygmaea*.  
*E. SPATHULATA* Lem. = *Gnaphalium bracteatum*.

CCXXVIII. MICROPSIS.

Capitulum pluriflorum heterogamum, fl. omnibus tubulosis, exter. 2-3-serialibus fœmineis, inter. circ. 3 masculis. Invol. squamæ 2-3-se-

riales apice membranaceæ oblongæ subacutæ glabræ. Recept. angustum nudum. Achænia villosa oblonga pappo parvo squamelloso superata. — Herba Chilensis habitu Evacis seu Micropi, undè nomen.

1. *M. NANA*. (1) in pascuis declivibus collium prope Quillota Chilensium legit el. Bertero! (pl. exa. n. 1146), flor. septemb. Herba multicaulis vix pollicem longa. Folia alterna obovata adpressè cano-villosa integerrima obtusa mucronata, super. congesta subverticillata in axillis capitula parva sessilibus tomentosa (v. s.)

CCXXIX. *MICROPUS* Linn. gen. n. 996. Gærtn. fr. 2. p. 389. t. 164. Cat. dict. 31. p. 39. Less. syn. 205. — *Gnaphalodes* Tourn. inst. 439. t. 261. *Adans. fam.* 2. p. 118. — *Micropi* sp. Desf.

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, radii circ. 5-7 uniserialibus fœmineis filiformibus, disci 5-7 masculis 5-dentatis. Invol. biseriale, squamis flores et fructus radii involventibus. Recept. ebracteolatum angustum. Achænia radii lateraliter compressa obovata glabra calva squamis inv. involuta et cum eo decidua. — Herbæ Europæ, lanato-tomentosæ, pusillæ, Filaginis s. *Gnaphalii* facie. Folia alterna. Capitula glomerata.

#### Sect. I. ACANTHOLÆNA.

Invol. squamæ achænia involventes dorso aculeis rigidis echinatæ.

1. *M. SUPINUS* (Linn. sp. 1313), totus adpressè sericeo-lanatus, caulibus diffusis erective, foliis spathulatis obtusis, capitulis axillaribus vix aggregatis, invol. squamis dorso echinatis. (1) in aridis inundatisque desiccatis apricis Europæ Mediterraneæ ab Hispaniâ ad Persiam, nec non in Oriente et Mauritaniâ. Cr. ic. 1. t. 35. Lam. ill. t. 694. f. 1. Filago supina Lam. fl. fr. 2. p. 60. *Gnaphalodes* dentata Mœnch meth. 568. — Pluk. alm. t. 187. f. 6. Variat caule setosius erecto, aut plurimis diffusis imò procumbentibus. Fructus squamis inv. mucronati involuti, forè triangulares, non malè *Melampodii* achænia referunt (v. v. et s.)

#### Sect. II. BOMBYCILÆNA.

Invol. squamæ achænia involventes dorso inermes lanatæ.

2. *M. ERECTUS* (Linn. sp. 1313), totus niveo-lanatus, caulibus erectis aut subdiffusis, foliis oblongo-lanceolatis obtusiusculis, capitulis axillaribus terminalibusque aggregatis, invol. squamis dorso convexis parè lanatis. (1) in spiculis sterilibus rupestribus et inundatis exsiccatis Europæ Mediterraneæ et mediæ Hispaniâ ad Persiam, nec non in Oriente et Mauritaniâ. Lam. ill. t. 694. f. 1. Loefl. itin. 166. t. 1. f. 3. Lag. nov. gen. et sp. p. 32. n. 401. Variat ut pro caule sæpius erecto simplici aut multicauli, rarius multicapiti supino (Filago multicaulis Lam. fl. fr. 2. p. 59). (v. v. et s.)
3. *M. BOMBYCINUS* (Lag. nov. gen. et sp. p. 32. n. 400), totus niveo-lanatus, caulibus erectis aut subdiffusis, foliis lineari-oblongis obtusis, capitulis congestis axillaribus et terminalibus, invol. squamis dorso convexis lanâ floccosissima tumida (nec adpressâ) obductis. (1) in collibus aridis Hispaniæ circa Madridem (Lag.!), Tudelam (Duf.!), et Galliæ australis in Occitaniâ totâ. *Micropus erectus* Guss. pl. rar. p. 345? Cass. dict. 31. p. 40? — Clus. hist. 1. p. 329. f. 1? Variat ut duæ sp. priores caule erecto subsimplici apice subdichotomo. (*M. discolor* Pers. ench. 2. p. 422), aut multicauli supino vel ascendente (*Barr.* t. 296). Prioris fortè varietas? (v. v. et s.)
4. *M. GLOBIFERUS* (Bert. herb. n. 433), totus niveo-tomentosus, caulibus basi ramosis diffusis, foliis oblongo-linearibus, floralibus latioribus ebracteatis capitulis terminalibus lateralibusque, invol. squamis planiusculis inermibus (v. v. et s.)

mentosis obtusissimis. ① in Chili ad Rancagua legit cl. Bertero. Squamæ recept. involutæ obovato-compressæ tomentosæ achænium foventes et globum incanum cum invol. constantes. (v. s.)

5. *M. minimus*, totus niveo-tomentosus, caule brevissimo erectiusculo parçè ramoso, foliis ovali-oblongis obtusis, capitulis ad apices ramorum confertis. ① in Mexico ad Monterey januar. fl. legit cl. Berlandier (pl. exs. n. 1861). Priori affinis sed multò minor. Species quoad caract. fl. et fr. recognoscenda. (v. s.)

† *Species exclusæ.*

*M. exiguus* Spreng. = *Evax exigua*.

*M. marboticus* Spreng. = *Evax Mareotica*.

*M. pygmaeus* Desf. = *Evax pygmaea*.

CCXXX. *EPALTES* Cass. bull. philom. 1818. p. 139. dict. 15. p. 6. 37. p. 339. Less. syn. 206. — *Ethulia* Gertn. fr. 2. p. 389. t. 164. non Cass. — *Ethulia* sp. Linn.

Capitulum multiflorum heterogamum, fl. radii pluriserialibus fœmineis, disci paucioribus androgynis sterilibus. Invol. bi- aut pluriserialia imbricata campanulatum. Recept. nudum convexum. Cor. radii filiformes tenuissimæ vix apice dentatæ, disci obconicæ 5-dentatæ extus parçè glandulosæ. Antheræ inclusæ. Stylus radii exsertus ramis revolutis, disci breviter exsertus indivisus. Achænium radii obovato-subangulatum glabrum epapposum, disci abortivum. — Herbæ ramosæ. Folia alterna oblonga aut obovata integra aut vix dentata sessilia aut longè decurrentia. Pedunculi oppositifolii. Capitula solitaria aut gemina pedicellata subpaniculata. — Genus affine *Denekia* et *Dichrocephala*.

\* *Foliis decurrentibus.*

1. *E. divaricata* (Cass. ll. cc.), caule herbaceo erecto divaricato-ramosissimo, foliis lineari-oblongis basi attenuatis longè decurrentibus remotè denticulatis, pedunculis folio brevioribus subbracteatis, invol. squamis acuminatissimis disco longioribus. ① frequens in orysetis et agris Indiæ orient. *Ethulia divaricata* Linn. mant. 110. Burm. ind. 176. t. 58. f. 1. Wall. cat. et herb. n. 98. A et B. — Pluk. alm. t. 21. f. 4. Variat omninò glabra aut hinc inde versus apicem sublanuginosa. Invol. albida subcariosa rariùs subpurpureascentia. (v. s.)
2. *E. linearifolia*, caule herbaceo erecto apice corymboso-ramoso, foliis linearibus angustè decurrentibus subintegerrimis, pedunculis folio longioribus bracteolatis, invol. squamis acuminatissimis disco longioribus. ① in Ind. orientali circa Promæ regni Avenis. *Ethulia divaricata* var. C. Wall. cat. et herb. n. 98. C. Priori simillima et fortè varietas? Involucrum in omnibus specim. mihi obvis subpurpureascent. Folia infer. sesquipollicaria vix ultra lineam lata. Pedunculi ferè pollicares bracteolas 3-5 acutissimas patulas gerentes. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)
3. *E. pygmaea*, caule herbaceo nano ramoso, foliis lineari-oblongis subdenticulatis angustissimè decurrentibus, pedunculis ebracteatis folia vix superantibus, involucri squamis adpressis vix acutis flores non superantibus. ① in Indiæ orientalis agro Carnatico legit cl. Belanger. Herba sesquipoll. glabra. (v. s. comm. à cl. inv.)
4. *E. brasiliensis*, caule herbaceo erecto ramoso, foliis oblongo-lanceolatis acuminatis dentato-serratis in alas integerrimas latas foliaceas secus caulem et ramos decurrentibus, pedunculis ramosis subulatis, corymbis folio brevioribus, invol. squamis ovalibus obtusis flores non superantibus. ① in humidis circa Bahiam legit cl. Salzmann. Habitus ferè *Pluchæa* Quitoc, sed caract. genericiis



2. *I. ROYLEANA*, tota molliter villosa subtomentosa, caule erecto simplici 1-cephalo, foliis ovatis subdenticulatis, caulinis infer. in petiolum alatum basi obtusè auriculato-dilatatum semisimplexicaulem productis, summis pæcis sessilibus cordato-amplexicaulibus, achæniis teretiusculis?  $\frac{1}{2}$  in Indiæ prov. boreali-occid. detexit cl. Royle. Species insignis Inulæ Helenio habitu, involucri ligulis numerosissimis angustis et pappo rigidiore proxima, sed foliis etiam caulinis plerisque longè petiolatis mediisque subpandæreiformibus distincta. Achænia ex immaturo statu vid. teretiuscula nec tetragona. (v. s. comm. à cl. Royle.)

Sect. II. BUBONIUM. — *Enula Duby bot. gall. 1. p. 267* (sed *Enula* veterum est *Corvisartia*.)

Squamæ lineares aut lanceolatæ, apice appendiculo parvo foliaceo auctæ, extr. subbreiores.

3. *I. CONYZA*, caule erecto apice ramoso subcorymboso pubescenti-subtomentoso, foliis ovato-lanceolatis subacutis subintegerrimis suprà subpubescentibus subtus tenuè tomentosis, radicalibus in petiolum attenuatis, caulinis sessilibus, invol. squamis linearibus apice foliaceis patulo-recurvis, fl. radii femineis trifidis subligulatis.  $\frac{1}{2}$  ad sepes, colles, prata sicca præsertim in calcareis totis ferè Europæ usque ad Byzantium, Tauriam et Persiam. *Conyza squarrosa* Linn. sp. 1205. fl. dan. t. 622. engl. bot. t. 1195. Cass. dict. 10. p. 306. Lam. ill. t. 697. f. 1. *Conyza vulgaris* Lam. fl. fr. p. 73. *Inula squarrosa* Bernh. 208 Linn. *Erigeron squarrosus* Clairv. herb. val. 243. *Inula* sp. Less. syn. 191. — Cam. epit. 612. Herba sordida obscura trita fetida. Invol. è virescente perparascens. Cor. flavæ, marginales ad latus internum profundius fissæ et ideo ad ligulas subcedentes. (v. v.)
4. *I. SUAVIOLENS* (Jacq. hort. vind. 3. p. 29. t. 51), caule erecto simplici apice subcorymboso, foliis ellipticis subvillosis, inferioribus in petiolum longè attenuatis remotè serrulatis, caulinis basi breviter attenuatis integerrimis, capitula 5-pedunculatis, invol. campanulati squamis apice foliaceis subrecurvis, ligulis linearibus involucri longioribus.  $\frac{1}{2}$  in Italiæ littoribus circa Monfalcone (Suffr. ex Reichenb. fl. exc. 2. p. 237), fortè in Gallo-Provincia ex Lam. Marr. syst. p. 766. Willd. sp. 3. p. 2099. DC. fl. fr. n. 3144. Herba trita potius grave quàm suavè olens. Valdè affinis *I. Conyzæ* et hujus varietas ligulata videbatur nisi capitula majora. (v. s. c. in hort. Taurin. et Cels.)
5. *I. CANDIDA* (Cass. dict. 23. p. 554), tota densè candido-tomentosa, foliis ovatis radicalibus longè, caulinis breviter petiolatis, pedunculis terminalibus et axillaribus 1-cephalis, invol. squamis oblongo-linearibus obtusis.  $\frac{1}{2}$  in repæstriis apricis Crætæ, Sami, Dalmatiæ, Istriæ.
- a. *verbascifolia*, foliis subcrenato-dentatis acutiusculis, capitulis radiatis, ligulis elongatis.  $\frac{1}{2}$  *Conyza verbascifolia* Willd. sp. 3. p. 1924 non Bory. Con. candida Linn. sp. 1208. Sibth. fl. græc. t. 861 ex Sm. prod. 2. p. 173 — Bocch. sic. t. 31. f. 2. (v. s. ex Dalmatiâ.)
- ß. *limonifolia*, foliis integerrimis obtusis, capitulis flosculosis aut ligulis involucri breviores gerentibus.  $\frac{1}{2}$  *C. limonifolia* Sibth. fl. græc. t. 865 ex Sm. prod. 2. p. 174. *C. candida* Will. sp. 3. p. 1923 non Lour. (1) Bory. ic. t. 217. (v. s. ex Atticâ, Crêtâ et Samo.)
6. *I. THAPSOIDES*, tota tomentoso-lanata, caule erecto apice corymboso-ramoso, foliis serratis acuminatis radicalibus petiolatis ovali-oblongis, caulinis admodum longè decurrentibus, capitulis corymbosis confertis, invol. squamis linearibus  $\frac{1}{2}$  in herbidiis ad Caucasum. Etiam si nonnulli dixerint nihil esse clarior ac identitas synonymi. infra citatorum, fortè 3 varietates sequentes ut species habende.
- a. *Sprengelii*, tomento denso crasso, foliis caulinis brevius decurrentibus.

(1) *Conyza candida* Lour. coch. 2. p. 605 est verisimiliter species omnino diversa sed nondum satis nota.

Rhantario) opalescentum. Antheræ caudatæ. — Folia alterna. Capitula in glomerulum non aggregata. Ligulæ ubi adsunt disco concolores. — Species omnes Gerontogæ.

## Div. I. EUINULÆ.

Capitula pluri- aut multiflora in glomerulum non concreta.

CCXXXII. RHANTERIUM Desf. *fl. atl.* 2. p. 291. t. 240. *Cass. dict.* 45. p. 312. *Less. syn.* 199.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus foemineis ligulatis, disci tubulosis hermaphr. 5-dentatis. Invol. ovatum, squamis basi concavis imbricatis productis in appendicem rigidam uncinato-reflexam. Recept. bracteolatum, bracteolis involucrique squamis internis flores subamplectantibus. Antheræ apice acutè appendiculatæ, basi caudatæ. Achænia ovali-subpentagona glabra erostris, radii epapposæ, disci coronatæ paleis 5 caducis basi integerrimis utrinquè ad apicem densè barbellatis. — Suffrutex Mauritanicus rigidus ramosus parcè foliosus, ramis tomentosis apice 1-cephalis. Folia sparsa lanceolata argutè serrata glabriuscula. Flores lutei. — Genus recept. paleaceo ad Bupthalmæas accedit.

1. R. SUAVEOLENS (Desf. l. c.). 5 in arenis maritimis prope Sfax regni Tunisani. Folia tria suaveolentia. (v. s. comm. à cl. Desfontaines.)

CCXXXIII. INULA *Gartn. fr.* 2. p. 449. f. 170. *Less. syn.* 198. — Inulæ et Conyzæ sp. *Lin.* — Inula, Limbarda, Corvisartia et Conyza *Cass.* — Conyza *Don prod. fl. nep.* 176 non *Lin.*

Capitulum multiflorum heterogamum, fl. radii 1-serialibus foemineis abortu interdum substerilibus, sæpius ligulatis, rarius subtubulosis 3-fidis, disci hermaphroditis tubulosis 5-dentatis. Invol. imbricatum pluriseriale. Recept. planum aut subconvexum nudum. Antheræ basi bisetæ. Achæmium erostre teretiusculum aut in I. Helenio tetragonum. Pappus conformis 1-serialis, setis capillaribus subscabris. — Herbæ gerontogæ pleræque Europææ aut Asiaticæ, perennes, rarius biennes aut annuæ. Folia caulina alterna, sæpè amplexicaulia, indivisa integerrima aut serrata. Capitula ad apices pedunculorum solitaria sæpè corymbosa flava.

Sect. I. CORVISARTIA *Merat fl. par. ed.* 2. v. 2. p. 261. *Cass. dict.* 10. p. 572.

Invol. squamæ ext. latè ovatæ foliaceæ, inter. obovato-spathulatæ obtusæ. Achæmium glaberrimum, in I. Helenio tetragonum, in I. Royleanâ fortè teretiusculum. — Si in utraqûe specie achæmium tetragonum Corvisartia ut genus habita consentiente admittendum, si achænia diversa I. Royleana optimè jungit I. Helenium cum cæteris Inulis.

1. I. HELENIUM (*Lin.* sp. 1236), caule erecto, foliis dentatis subulis velutino-tomentosis acutis, radicalibus ovatis in petiolum longè attenuatis, caulinis semiamplexicaulibus, pedunculis paucis 1-cephalis ad apicem corymbosis. 4 in Angliâ, Belgio, Galliâ septentr. et orientali, Germaniâ. fl. dan. t. 728. *Gartn. fr.* 2. t. 170. Aster Helenium *Scop. carn.* n. 1078. Aster officinalis *All.* n. 705. Corvisartia Helenium *Merat et Cass. ll. cc.* Helenium *Cam. epit.* 35. *Fuchs hist.* 242. Radix magna amara et aromatica, *Enula Campana* offic. dicta (v. v. et s.)

2. *I. ROYLEANA*, tota molliter villosa subtomentosa, caule erecto simplici 1-cephalo, foliis ovatis subdentulatis, caulinis infer. in petiolum alatum basi obtusè auriculato-dilatatum semimplexicaulem productis, annuatis paucis sessilibus cordato-amplexicaulis, acheniis teretiusculis?  $\frac{1}{2}$  in Indiæ prov. boreali-occid. detexit cl. Royle. Species insignis Inula Helenio habitu, involacro, ligulis numerosissimis angustis et pappo rigidiore proxima, sed foliis etiam caulinis plerisque longè petiolatis mediisque subpanduræformibus distincta. Achenia ex immaturo statu vid. teretiuscula nec tetragona. (v. s. comm. à d. Royle.)

Sect. II. BUBONIUM. — *Enula Duby bot. gall. 1. p. 267* (sed *Enula* veterum est *Corvisartia*.)

Squamæ lineares aut lanceolatae, apice appendiculo parvo foliaceo auctæ, exte. subbreviares.

3. *I. CONYZA*, caule erecto apice ramoso subcorymboso pubescenti-subtomentoso, foliis ovato-lanceolatis subacutis subintegerrimis suprâ subpubescentibus subtus tenuè tomentosis, radicalibus in petiolum attenuatis, caulinis sessilibus, invol. squamis linearibus apice foliaceis patulo-recurvis, fl. radii feminæ trifidis subligulatis.  $\frac{1}{2}$  ad sepes, colles, prata sicca præsertim in calcareis totis ferè Europæ usque ad Byzantium, Tauriam et Persiam. *Conyza squarrosa* Lam. sp. 1205. fl. dan. t. 622. engl. bot. t. 1195. Cass. dict. 10. p. 306. Lam.  $\frac{1}{2}$  t. 607. f. 1. *Conyza vulgaris* Lam. fl. fr. p. 73. *Inula squarrosa* Bernh. non Linn. *Erigeron squarrosus* Clairv. herb. val. 243. *Inula* sp. Less. syn. 191. — Cam. epit. 612. Herba sordida obscura trita fœtida. Invol. è virescente perperascens. Cor. flavæ, marginales ad latus internum profundius fissæ et idcirco ligulas subaccedentes. (v. v.)
4. *I. SUAVIOLENS* (Jacq. hort. vind. 3. p. 29. t. 51), caule erecto simplici apice subcorymboso, foliis ellipticis subvillosis, inferioribus in petiolum longè attenuatis remotè serrulatis, caulinis basi breviter attenuatis integerrimis, capitulis 5-pedunculatis, invol. campanulati squamis apice foliaceis subrecurvis, ligulis linearibus involucri longioribus.  $\frac{1}{2}$  in Italiæ littoribus circa Monsaleon (Suffr. ex Reichenb. fl. exc. 2. p. 237), fortè in Gallo-Provincia ex Lam. Mar. syst. p. 766. Wild. sp. 3. p. 2099. DC. fl. fr. n. 3144. Herba trita potius græquàm suavè olens. Valde affinis *I. Conyza* et hujus varietas ligulata videtur nisi capitula majora. (v. s. c. in hort. Taurin. et Cels.)
5. *I. CANDIDA* (Cass. dict. 23. p. 554), tota densè candido-tomentosa, foliis ovatis radicalibus longè, caulinis breviter petiolatis, pedunculis terminalibus et axillaribus 1-cephalis, invol. squamis oblongo-linearibus obtusis.  $\frac{1}{2}$  in rupestribus apricis Cretæ, Sami, Dalmatiæ, Istriæ.
- a. *verbascifolia*, foliis subcrenato-dentatis acutiusculis, capitulis radialis ligulis elongatis.  $\frac{1}{2}$  *Conyza verbascifolia* Willd. sp. 3. p. 1924 non Bory de Saint-Vincent, candida Linn. sp. 1208. Sibth. fl. græc. t. 861 ex Sm. prod. 2. p. 171. — Boeck. sic. t. 31. f. 2. (v. s. ex Dalmatiâ.)
- ß. *limonifolia*, foliis integerrimis obtusis, capitulis flosculosis aut ligulis involucri breviores gerentibus.  $\frac{1}{2}$  *C. limonifolia* Sibth. fl. græc. t. 851 ex Sm. prod. 2. p. 174. *C. candida* Will. sp. 3. p. 1923 non Lour. (t. Mart. t. 217. (v. s. ex Atticâ, Cretâ et Samo.)
6. *I. THAPSOIDES*, tota tomentoso-lanata, caule erecto apice corymboso-ramoso, foliis serratis acuminate radicalibus petiolatis ovali-oblongis, caulinis adaxialibus longè decurrentibus, capitulis corymbosis confertis, invol. squamis linearibus.  $\frac{1}{2}$  in herbis ad Caucasum. Etiam si nonnulli dixerint nihil esse clarius ac idcirco titulus synonym. infra citatorum, fortè 3 varietates sequentes ut species habendas.
- a. *Sprengelii*, tomento denso crasso, foliis caulinis brevius decurrentibus.

(1) *Conyza candida* Lour. coch. 2. p. 605 est verisimiliter species omnino diversa sed nondum satis nota.

capitulis paucioribus latioribus, involucri tomentoso-lanati squamis obtusis, ligulis involucrum discumque longè superantibus. — *Inula thapsoides* Spreng. pug. 1. p. 58. Cat. sem. h. hal. 16. Media ferè inter *I. candidam* var. *a* et var. seqq. (v. s. comm. à cl. Spreng. et v. c. in h. Monsp.)

β. *Poiretii*, tomento parciore vix albido, foliis caulinis longius decurrentibus, corymbis numerosioribus confertis, invol. ovali squamis linearibus acutis glabrisculis, ligulis vix involucro longioribus. — *Inula verbascifolia* Poir. suppl. 3. p. 154. Horn. h. hafa. supp. 99. *Inula thapsoides* Reichenb. icon. t. 347. (v. s. et v. c. in hort. Par.)

γ. *Biebersteinii*, tomento denso-crasso, foliis caulinis longè decurrentibus, corymbis numerosis confertis, invol. ovato-oblongi squamis linearibus rectis acutis vix dorso villosis, ligulis nullis aut paucissimis brevissimis trifidis. — *Conyza thapsoides* Bieb. fl. taur. et suppl. n. 1716. (v. s. comm. à cl. Steven et v. c. in hort. Monsp.)

7. *I. POLYGONATA*, ramis teretibus villosis apice tomentosis foliorum decurrentiis integerrimis longis alatis, foliis elliptico-oblongis utrinquè attenuatis dentatis supra asperis subtus et summis utrinquè densè tomentosis, corymbo terminali, capitulis subsessilibus paucis bracteis tomentosis circumdati, invol. squamis ext. oblongis densè tomentosis obtusis, intimis lanceolatis acuminatis scariosis glabris, ligulis nullis. 5? in Indià or. ad Promè in regno Avenis legit cl. Wallich. *Conyza polygonata* Wall. cat. u. 3010. comp. n. 120. Caulis fortè basi frutescens? Rami pedales et ultrà crassitie pennæ anserinæ. Foliorum superiorum dentes callosi. Flores ex statu misero specim. non ritè vidi et hic retuli speciem ex analogiâ cum *In. candidâ* et *thapsoides*. (v. s. comm. ab amp. Car. merc. angl. Ind. or.)

8. *I. BIFRONS* (Lin. sp. 1236), caule apice ramoso corymboso-subdichotomo viscoso, foliis glabris ovali-oblongis aliis cordato-amplexicaulibus aliis decurrentibus, basi dentatis apice integerrimis mucronatis, capitulis confertè corymbosis subsessilibus, invol. squamis oblongo-linearibus subsquarrosis, ligulis brevissimis. (2) in vallibus et humidis Pyrenæorum (Herm.), Arvernæ, Occitanie, Galloprovincie, Pedemontii, Liguriæ, Gallicie (Bess.), Italiæ. DC. fl. fr. 4. p. 155. Reichenb. icon. 4. n. 527. t. 346. Inv. glabra Bess. cat. h. Crzm. ex Reich. — Herm. par. t. 127. Bocc. mus. 1. t. 121. Garid. Aix. t. 23. — *Conyza alata* Baumg. ex Reichenb. (sed in fl. trans. non reperi). *Aster bifrons* All. ped. n. 714. *Inula glomeriflora* Lam. fl. fr. 2. p. 150. *Conyza bifrons* Gou. hort. 436 non Linn. (v. s.)

9. *I. OCVLUS-CHRISTI* (Linn. sp. 1237), caule erecto apice parçè ramoso-corymboso molliter villosa aut lanata, foliis oblongo-lanceolatis integerrimis utrinquè villosa-sericeis, radicalibus in petiolum attenuatis, caulinis basi obtusâ amplexicaulibus, capitulis 3-5 longiusculè pedunculatis, invol. squamis oblongo-linearibus dorso hirsutis, ligulis invol. duplò longioribus. 2/ in collium lateribus graminosis apricis Austriæ, Hungariæ, Dalmatiæ. Willd. sp. 3. p. 2090. — Clas. hist. 2. p. 20. f. 2. Plantam ex Austriâ loco Clusiano lectam (quæ typus speciei) à Jacquinò illustratam et à filio suo mihi missam comparavi cum Caucasica nec ullum discrimen specificum vidi. — Inul. montanam fl. taur. n. 1746 huc refert cl. Steven. (v. s.)

β. *campestris*, tota parciùs villosa, foliis superioribus non tantum cordato-amplexicaulibus sed etiâ subdecurrentibus. 2/ in Podoliâ (Bess.), Tauriâ (Bieb. Stev.!) et Persiâ (Szow.!). *I. Oculus-Christi* Bieb. fl. taur. n. 1744. *I. campestris* Bess. enum. Vollh. p. 33 et 108. Recept. in hac et in var. *a* pubè minutissimâ breviter puberulam. (v. s.)

10. *I. WELSHIODES* (DC. fl. fr. suppl. (1815) p. 470), caule erecto hirsuto supernè corymboso-ramoso, foliis ovali-oblongis subhirsutis subdentatis, radicalibus petiolatis, superioribus basi obtusâ amplexicaulibus, capitulis 3-5-foliis floralibus bracteatis, invol. campanulati squamis linearibus hirsutis, ligulis invol. duplò longioribus, recept. ad medium setuloso. 2/ in agris asperis Pyrenæorum

orientalium demissorum legi ad pagos Finestret, Ris, Billoe, etc. et in montibus Corbariensibus. Inula Oculus-Christi Lapeyr. abr. 522 non Linn. (v. v. sp.)

β. *flouculosa*, ligulis nullis. (v. s. c.)

11. I. HYRTA (Linn. sp. 1239), caule erecto subramoso piloso-hirto, foliis sessilibus lanceolato-oblongis rigidis integerrimis secus nervos et margines hirtis, capitulis solitariis aut subternis corymbosis, invol. squamis hirtis ciliatis linearilanceolatis, exterioribus interiores superantibus, acheniis glabris. ½ in pratis asperis et montosis Galliae imò prope Parisios, Pedemontii, Appennini Genensis et Etrusci, Sabaudiae, Helvetiae, Carnioliae, Thuringiae, Austriae, Tauriae, Sibiriae, Dahuriae. Jacq. fl. austr. t. 358. DC. fl. fr. 4. p. 151. Inula montana Poll. pal. Aster hirtus Scop. carn. n. 1082. t. 58. Clas. hist. s. p. 14. f. 2. Variet 1-3-cephala. (v. v. et s.)

12. I. VAILLANTII (Will. dauph. 3. p. 216), caule erecto ramosissimo subcorymboso foliisque subtus pube brevi velutino-cinerea, foliis oblongo-lanceolatis subserratis, canlinis sessilibus, infimis basi subattenuatis, invol. squamis ext. oblongo-linearibus subfoliaceis, intimis linearibus glabris, ligulis invol. duplo longioribus, acheniis glabris. ½ in dumetis et pratis montosis Helvetiae orientalis, Sabaudiae, Delphinatus et Pedemontii. Fl. fr. n. 3152. Gaud. fl. helv. 5. p. 325. Aster Vaillantii All. fl. ped. n. 719. In. cinerea Lam. dict. 3. p. 159. Inula ramosissima Clairv. man. herb. 244. — Hall. helv. n. 73. t. 2. opt. Herba Melissam ferè olet. (v. v. et s.)

13. I. SEMIAMPLEXICAULIS (Renter! mem. soc. Gen. 7. p. 169. lc.), caule erecto apice corymboso pubescenti-subviloso, foliis semiamplexicaulibus oblongo-lanceolatis acutis calloso-serratis supra glabris subtus pubescenti-villosis, invol. squamis ext. oblongo-lanceolatis patulis foliaceis, inter. linearibus glabris, ligulis invol. duplo longioribus, acheniis glabris. ½ olim legi in Pedemontii agro Cuneensi et cum I. Vaillantii praesente Allionio (n. 719?) confundi, recentius in sylva La Batie prope Genevam cum I. Vaillantii et I. salicina legitur ocul. Renter. Utriusque fortè hybrida? Differt ab I. salicina foliis dorso canaliculatis villosis, herbâ Polii odore donatâ, foliis frequentius serratis. Distinguitur ab I. Vaillantii foliis amplexicaulibus minus villosis-tomentosis paulò rigidioribus, odore subdiverso, ligulis paulò longioribus. (v. v.)

14. I. SALICINA (Linn. sp. 1238), glabra, caule erecto apice corymboso-ramoso, foliis semiamplexicaulibus lanceolatis margine ciliato-scabris apice subrecurvis, ramis 1-cephalis, invol. campanulati squamis ovato-lanceolatis serratis-scabris apice subreflexis, acheniis glabris. ½ in pratis asperis dumetis et rivulis Galliae, Helvetiae, Italiae, Germaniae, Daniae, etc. Fl. dan. t. 786. Fl. fr. 4. p. 151. Gaud. fl. helv. 5. p. 324. Aster salicinus All. ped. n. 709. — Cha. hist. 2. p. 15. f. 1. Mor. ox. s. 7. t. 21. f. 10. Herba inodora. (v. v. sp.)

β. *latifolia*, foliis ovato-lanceolatis, rigidioribus margine magis scabrida

½ Inula aspera Poir. suppl. 3. p. 154. (v. v. c.)

γ. ? *lancifolia*, caule hirtio, capitulis minoribus. ½ Inula lancifolia Wenderoth in florâ 1829. erg. p. 1. p. 26. An potius I. squarrosa seu spiraeifolia var.?

15. I. ENSIFOLIA (Linn. sp. 1240), caule erecto glabro apice araneoso simpliciter 1-5-cephalo, foliis sessilibus linearibus calloso-acuminatis multinerviis glabris, capituli squamis lanceolatis erectis foliosis subtus araneoso-tomentosis. ½ in collibus et asperis Pedemontii, Italiae, Carnioliae, Bavariae, Austriae, Tauriae, Caucasi. Jacq. fl. austr. t. 162. Bieberst. fl. taur. n. 1753. D'Urv. Enum. n. 811. Aster ensifolius Scop. carn. n. 1084. — Bocc. mus. 1. p. 31. t. 18. Variet caule 1-cephalo aut polyccephalo sed tres varietates speciesve mihi non satis notè distinguit cl. Wenderoth in florâ 1829. p. 1. p. 26. (v. s.)

16. I. SQUARROSA (Linn. sp. 1240 non Bernh.), caule erecto vix ramoso scaberrimo, foliis oblongis acuminatis serratis utrinquè et margine plus minus scabridis reticulato-venosis, capitulis ad apices ramulorum solitariis vix corymbosis, invol. campanulati squamis ovatis apice patulo-reflexis margine scabris

¶ in sylvis petrosis Gallie, Helvetiæ et Germaniæ australis, Italiæ, etc. Bert. ann. ital. p. 192 et Gaud. fl. helv. 5. p. 321. var.  $\alpha$ . DC. fl. fr. n. 3148. *Aster corymbosus* Moench. non Ait. In. spiræifolia Lam. dict. 3. p. 258. — Pluk. alm. t. 16. f. 1. (v. v. sp.)

¶ *latifolia*, foliis ovato-lanceolatis subcabris. ¶ in Delphinato prope Gratiopolim. Inula germanica Vill. Dauph. 3. p. 219. Lam. dict. 3. p. 258. DC. fl. fr. n. 3149 non Linn. (v. s.)

17. *I. spiræifolia* (Linn. sp. 1238 non Lam.), puberulo-scabra, caule erecto simplici apice summo confertè corymboso, foliis sessilibus oblongis cartilagineo-denticulatis et mucronatis rigidis reticulatis, invol. squamis ovatis glabris obtusiusculis squarrosis. ¶ in Pedemontio, Carniolâ calidiore, Austriâ littorali, Podoliâ, Tauriâ, Caucaso, Persiâ. Inula Bubonium Murr. syst. 767. Jacq. fl. austr. app. t. 19. Bieb. fl. taur. n. 1751. In. salicina Pall. ind. taur. n. 154 ex Bieb. *Aster Bubonium* Scop. carn. n. 1083. t. 58. *Aster squarrosus* All. pedem. n. 708. excl. syn. ex cl. Moretti qui primus synonym. hujus speciei elucidavit. — *Aster* IV Clus. hist. 2. p. 14. Bubonium Tab. hist. 716. Inula spiræifolia et Bubonium Willd. sp. 3. p. 2094 et 2097. (v. s.)

¶ *hybrida* (Baumg. fl. trans. 3. p. 132), capitulis subternis majoribus. ¶ in montosis Podolizæ et Transylvaniæ. Fortè hæc var. *I. spiræifolia* et *I. squarrosam* Linn. præsentibus cl. Lamarck Duby et Gaudin in unicum colligit. (v. s.)

18. *I. micranthos* (Poir. dict. suppl. 3. p. 153), caule erecto simplici pubescente, foliis sessilibus lanceolatis integerrimis suprâ subcabris subtus villosis subsericeis, capitulis corymbosis confertis, invol. squamis linearibus acutis dorso puberulis, acheniis villosis. ¶ in Armeniâ (Tourn.), in collibus Bulgariæ circa Sisepolim (D'Urv.) et in herbidiis apricis Tauriæ (Bieb.). Inula orientalis Willd. sp. 3. (1800) p. 2100, non Lam. (1789). *I. germanica* Bieb. fl. cauc. n. 1752 non Linn. ex D'Urv. enum. 815. *Aster armenius* salicis folio flore luteo minimo umbellato Tourn. cor. 36. Affinis *I. Germanicæ* sed quidquid dix. Spreng. diversa foliis subtus sericeis et præsertim invol. subcampanulato nec oblongo, acheniis villosis, etc. (v. s. comm. à cl. D'Urv.)
19. *I. Germanica* (Linn. sp. 1239), caulibus erectis simplicibus apice corymbosis, foliis cordato-semiamplexicaulibus ovali-lanceolatis integerrimis suprâ nodiusculis subtus pubescentibus, capitulis confertis breviter pedicellatis oblongis, invol. squamis adpressis apiculo subreflexo, ligulis disco parum longioribus, acheniis glabris. ¶ in herbidiis apricis Saxoniæ, Thuringiæ! Beasarabias!, in Tauriâ ad Odessam, etc. fortè in Sibirîâ (si icon Gmel. fl. sib. s. t. 78. f. 2. huc pertinet?). — Jacq. fl. austr. t. 134. opt. *I. Germanica* D'Urv. enum. 11. *I. media* Bieb. suppl. p. 576. Petif enum. pl. pal. 94. — Moris. ox. s. 7. t. 19. f. 26. pess. — *I. Germanica* florist. Gall. est diversa. (v. v.)
20. *I. media* (Bieb. suppl. 576), pilosiuscula, caule apice corymbum compositum glomeratum gerente, foliis sessilibus cordato-lanceolatis, infimis subdentatis, ligulis invol. parum longioribus, invol. submentosis. ¶ in Poloniâ australi legit cl. Besser ex Bieb. *Media* inter *I. salicinam* et *Germanicam*. *Corymbi* rami sub-1-cephali capitulo æquales aut breviores. †
21. *I. Caspica* (Blum. in Ledeb. ind. sem. h. dorp. 1822. p. 10), caule herbaceo tereti stricto simplici glabro, foliis elongatis lineari-lanceolatis sessilibus basi obtusè subauritis glabris integerrimis margine scabris, corymbi compositi ramis villosis, invol. squamis linearibus, ext. subfoliaceis rigidè ciliato-scabris, ligulis numerosis angustis. ¶ ad mare Caspium in salsis prope Astrachan. Folia 4 poll. longa, 3-4 lin. lata. (v. s. comm. à cl. Fischer.)
22. *I. Britannica* (DC. fl. fr. 4. p. 149), caule erecto villosa apice corymboso 3-5-cephalo, foliis longè lanceolatis remotè denticulatis subtus villosis, infimis basi attenuatis, superioribus dilatatis semiamplexicaulibus, invol. squamis linearibus laxis acheniisque hirsutis. ¶ in pratis secus vias et fossas, in Galliâ ferè totâ, Germaniâ, Daniâ, Italiâ, Græciâ (Sibth.), Persiâ (Scow.), Bulgariâ (D'Urv.), Sibirîâ, sed non in insulis Britannicis! Inula Britannica Linn.

sp. 1237. Sebkühr handb. t. 247. Fl. Dan. t. 413. Aster Britannicus All. pedem. n. 712. In. hirta Poll. pal. — Moris ox. a. 7. t. 19. f. 8. — Britanica Dalech. hist. 1083 unde nomen specificum à plurimis orthographiâ erroneè corruptum *Britavica*, Diosc. ex Dalech. Variat foliis integerrimis aut semieratis, caule 1-9-cephalo, capitalis minoribus ubi numerosa, majoribus ubi pauciora adsunt, foliis præsertim superioribus latioribus quæ I. Oestulian Reich. fl. exc. 2. p. 237. (v. v. et s.)

β. *comosa*, invol. squamis in folia disco longiora abeuntibus. I. *comosa* Lam. fl. fr. 2. p. 147. (v. s. olim.)

γ. *dichotoma*, pedunculis bifidis dichotomisve, capitalis minoribus, invol. squamis apice patulo-reflexis. — in Oriente nempe in Ægypto? ex Zucc. in Persiæ prov. Aderbeidschan ad rivalorum margines circa Khoi (Szow ex sp. Fisch.) et ad mare Caspium prope Astrachan et Hinkenberg? ex Ledeb. Inula dichotoma Zucc. obs. bot. n. 88. Inula Britanica anthod. minor. Fisch. et Mey. pl. Szow. ined. (v. s.)

23. I. *MONTANA* (Linn. sp. 1241), caule erecto 1-cephalo villosa, foliis lanceolatis integerrimis villosis, radicalibus in petiolum attenuatis, invol. squamis ext. oblongo-lanceolatis obtusiusculis extus villosis, inter. linearibus acutis, æchæniis hirtis. 2 in sterilibus montanis apricis Hispaniæ, Sicilia, Gallie australis, Pedemontii, Helvetiæ et Germaniæ australis, Istriæ et Tauris. DC. fl. fr. 4. p. 154. — Garid. Aix. t. 10. — Lob. ic. 350. f. 2. — Aster montanus All. ped. n. 706. In. odora D'Urv. enum. n. 812. excl. syn. Pulicaria calycis Presl. del. prag. p. 96 et Inula calycina Presl fl. sic. 1. p. xxxi hæc pen. monente cl. Gasone, Multum refert Pulicariam odorem sed facili differt pappo uno nec biseriali. Variat caule solitario aut multicapiti, 1-cephalo aut raris apice ramoso 1-3-cephalo. (v. v. et s.)

24. I. *GRANDIFLORA* (Willd. sp. 3. p. 2098), caule simplici 1-cephalo apice villosa, foliis sessilibus oblongis parè et sparsè pubero-villosis serratis, serraturis apice glandulosis, invol. squamis lineari-subulatis fulvo-hirsutis. 2 in alpe-tribus Caucasi Iberici (Stev.!), et ad Gurriel Persiæ (Szow.!), Bieb. fl. taur. et suppl. n. 1748. Inula alpina Adam in Web. et Mohr cat. 1. p. 68. In. glandulosa J. Sims. bot. mag. n. 1907. (v. s. comm. à cl. Steven.)

25. I. *GLANDULOSA* (Willd. sp. 3. (1800) p. 2097 non Lam. 1789), caule simplicissimo 1-cephalo patentem et molliter hirta, foliis sessilibus ovali-oblongis utrinquè villosa-hirtis glandulas secus marginem sessiles dentiformes gerentibus, invol. squamis lineari-subulatis nigricantibus hirsutis. 2 in alpe-tribus Caucasi. Bieb. fl. taur. et suppl. n. 1749. Bot. mag. t. 1907. bot. reg. t. 33. I. orientalis Lam. dict. 3. (1789) p. 255. non Willd. (1800). Capitulum raris excepto ferè sesquipoll. diam. Ligulæ lineares angustæ pollicem longæ. Vix affinis I. grandifloræ. Nomen Lamarckii arbitrariò à Willd. mutatum, imò altero confusum, stricto jure servandum esset. (v. s. comm. à cl. Steven.)

26. I. *GRAVEOLENS* (Desf. cat. ed. 2. p. 121), piloso-viscosa ramosissima, foliis inf. oblongo-lanceolatis subdentatis, sup. linearibus integris, ligulis nigris vix invol. æquantibus. 2 in apricis Europæ australis ad Constantinopolem, agrum Trajanum, Græciam, Italiam, Galliam australem et occident. Hispaniam, etc. Cass. dict. 23. p. 553. Erigeron graveolens Linn. sp. 1210. Schleg. graveolens Lam. fl. fr. 2. p. 145. DC. fl. fr. n. 3162. — Barr. ic. t. 3. 3. 2. Ca- lis vetustus sæpè indurato-suffrutescens. Pappi setæ numerosiores quàm in ce-teris. (v. v. et s.)

β. *frutescens*, caule humili fruticoso cæspitoso ramosissimo, foliis linearibus brevibus obtusiusculis integerrimis, ligulis fortè nullis. 3 ad Alexandriam Ægypti. (v. s. comm. à cl. Martins.)

27. I. *LATIFOLIA*, caule simplici basi tereti glabrescente suffruticuloso, sp. striato-sulcato setis hispido, foliis sessilibus latè ovalibus acutis basi longe cæneatis remotè serratis sparsè setosis obliquè penninerviis, capitalis paucis minimalibus, invol. squamis ext. linearibus subfollaceis hirtis interioribus paucioribus, ligulis invol. superantibus, æchæniis villosis. 3 2 in India orient.

- legit cl. Royle (n. 113. 2). Pedalis aut sesquipedalis. Folia circ. 4 poll. longa, a poll. lata, membranacea, serraturis callosis exsertis. (v. s. comm. à cl. Royle.)
28. *I. AFRICANA* (Lam.) dict. 3. p. 256), caule herbaceo tenui vix pilosiusculo ramoso, foliis amplexicaulibus subscaëtidis subtus glanduloso-punctatis, inferioribus oblongis acutis serratis, summis lanceolatis brevioribus subintegerrimis, ramis apice 1-cephalis aliis aphyllis aliis parçè foliolosis, invol. squamis lineari-setaceis. — ad Cap. Bonæ-Spei ex Sonnerat in h. Lam. et ad Galam Africæ (Cass.). *Iphiona punctata* Cass. dict. 23. p. 610. *Inula punctata* Less. syn. 198. Folia 6-7 lin. longa, 2-3 lin. lata. Capitula subglobosa 3-4 lin. diam. pallidè flava. Ligulæ breves. Habitus Pulicariæ sed pappus 1-serialis, Vicox sed fl. radii papposi! (v. s.)

## Sect. III. CAPPA.

Invol. squamæ multiseriales obtusiusculæ regulariter imbricatæ, ext. brevissimis, int. gradatim longioribus. Ligulæ paucæ parvæ aut subnullæ, fl. radii nempe 2-3-dentatis fere tubulosis (sed uniserialibus fœmineis ut in Inulâ). — Suffrutices Indici.

29. *I. EUPATORIODES*, caule fruticoso, ramis teretibus striatis rufo-velutinis, foliis ellipticis acuminatis basi attenuatis breviter petiolatis râriter et minutè serratis suprâ sparse scabris subtus rufo-cinereo-pubescentibus, paniculâ thyrsoidæ polycephala compositâ, ramis inf. axillaribus, invol. squamis oblongis obtusis subhirsutis, ligulis paucis inv. paulò longioribus, achæniis villosis. § in montibus prov. Silhet et fortè in Nepaliâ Ind. or. leger. cl. Bruce et Wallich. *Conyza eupatorioides* Wall. ! cat. et herb. n. 2993. comp. n. 103. Anth. breviter caudatæ. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or.)
30. *I. OBLONGA*, fruticosa, ramis teretibus apice præsertim sericeo-villosis, foliis brevissimè petiolatis oblongis utrinquè attenuatis dentato-serratis suprâ scabris subtus sericeis pubescentibusve, corymbo terminali congesto oligocephalo, capitulis vix pedicellatis, invol. squamis linearibus acutis velutino-hirsutis disco brevioribus, fl. fœm. paucis circ. 2-serialibus. § in Indiæ orient. montanis. Cor. purpureæ. Pappus apice crassius barbellaris subclavellatus. Ovarium teres pubescenti-sericeum. Anth. caudatæ.
- a. argentea*, foliis subtus densè sericeis albis, à medio ad apicem serratis. § ad Taong-Dong, Srinaghur, Kamroop et Charcooli in Deyra-Dhoou legit cl. Wallich. *Conyza argentea* A. B. C. Wall. ! cat. et herb. n. 3014. comp. 124 non Lam. *Conyza oblonga* Wall. ! herb. (v. s. comm. ab ampl. Cur. merc. Angl. Ind. or.)
- β. pubescens*, foliis tenuioribus subtus pubescentibus vix subsericeis fere à basi ad apicem serratis. § in demissis Nepalæ inferioris. C. argentea D. Wall. ! cat. l. c. An species propria? (v. s. comm. cum *a.*)
31. *I. CAPPA*, caule fruticoso, ramis teretibus albedo-velutinis, foliis elliptico-oblongis acuminatis basi attenuatis breviter petiolatis suprâ scabris subtus tomento sericeo niveo lanuginosis, paniculâ compositâ terminali polycephala, ramis niveis, inf. axillaribus folio multò brevioribus, invol. squamis lineari-oblongis subacutis albo-velutinis, ligulis subnullis, achæniis villosis. § in Nepaliâ superiore. *Conyza Cappa* Ham. in Don prod. 176. *Conyza lanuginosa* Wall. ! cat. et herb. n. 299a. comp. n. 102. *Eupatorium lanatum* Spreng. in litt. ad Wall. Frutex elegans. Pappus apice scaber. Fl. radii 1-ser. tubulosi 3-dentati aut subligulati sed cæteros non superantes. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or.)
32. *I. PSEUDO-CAPPA*, caule fruticoso, ramis teretibus scabridis villosis, foliis ellipticis vix acutis basi subobtusis sessilibus subintegerrimis suprâ sparse scabris subtus sericeo-lanato-villosis albidis, paniculâ terminali compositâ polycephala, ramis inf. axillaribus folia subabortiva triplo superantibus, inv. squamis lineari-oblongis vix subhirsutis, ligulis subnullis, achæniis villosis. § in Chinâ legit cl. Staunton. Valdè affinis *I. Cappæ*. Folia infer. non vidi, sed super. sessilia ubi in *I. Cappâ* petiolatâ. (v. s. comm. à cl. Lambert.)



33. *I. xrisophora*, caule suffruticoso erecto tereti densè hirsuto, foliis subsemissibus oblongo-lanceolatis subintegerrimis suprà hispidovulvatinis subitis densalbo-lanuginosis, corymbo terminali conferto 8-9-cephalo, invol. ovatis squamis linearibus acutis multiseriatis dorso tomentosis, ligulis subnullis, acheniis villosis.  $\zeta$  in Nepaliâ inferiore legit cl. Wallich. *Conyza eriophora* Wall: cat. n. 2013. comp. 123. Fl. radii. 1-ser. form. tubulosi tenues, cast. herm. stigmat. longis exsertis. Pappus 1-serialis. Affinis *I. Cappæ*. (v. s. comm. ab hon. Cur. merc. Angl. Ind. or.)

Sect. IV. LIMBARDA. — *Limbarda Adans. fam. 2. p. 125. Cass. dict. 23. p. 565. et 26. p. 437. Reichenbach fl. exc. 2. p. 237.*

Invol. squamæ omninò exappendiculatæ adpressæ acuminatæ pauciserialis, inter. non multò exteriores superantibus. Achenium villosum teretiusculum.

34. *I. CRITHMORHIZA* (Linn. sp. 1240), fruticulosa glaberrima, foliis linearibus obtusis carnosius integerrimis aut apice trifidia.  $\zeta$  ad rupes maritimas secus Oceanum ab Angliâ ad Lusitaniam et secus mare Mediterraneum ferè totum. DC. fl. fr. 4. p. 154. Engl. bot. t. 68. Inula crithmifolia Willd. sp. 3. p. 210. Senecio succulentus Forst. descr. 2. p. 149. Senecio crithmifolia Scop. carn. n. 1066. Limbarda tricuspidata Cass. dict. 26. p. 438. Inula Canariensis Mill. dict. n. 12? excl. patr. — Lob. ic. t. 395. f. 2. (v. v.)

35. *I. AMMOPHILA* (Bunge! enum. mong. ex litt.), glabriuscula, caulibus suffruticosis diffusis gracilibus, foliis lanceolato-linearibus sessilibus, infer. subcordatis integerrimis crassiusculis patentibus, capitulis solitariis terminalibus, invol. squamis adpressis acutis pallidis, fl. radii angustè ligulatis.  $\mathcal{Z}$   $\zeta$  in sabulosis Chinæ borealis legit cl. Bunge. Folia 2-3 lin. longa subvillosa, infer. breviora latiora, sup. angustiora paulò longiora. (v. s. comm. à cl. Bunge.)

*fl. salsoloides*, glanduloso-subviscosa, foliis evidentiùs cordatis, fl. radii tubulosi.  $\mathcal{Z}$   $\zeta$  in sabulosis Mongoliæ Chinensis circa Chadatu detexit cl. Turczaninow. *Conyza salsoloides* Turcz. ! pl. chin. et mong. dec. n. 20. Certè Inula ex antheris basi bisetis et *I. ammophila* var. *discoïdes* videtur etiamvis viscosa et foliis paulò latioribus donata. (v. s. comm. à cl. Turczaninow.)

36. *I. VISCOSA* (Ait. kew. ed. 1. v. 3. p. 223), caule erecto piloso-viscoso apice paniculato-ramoso, foliis lanceolatis serratis utrinquè glanduloso-pilosis sessilibus cordato-auriculatis, auriculis reflexis, ramis lateralibus foliosis oligocephalis, invol. squamis linearibus.  $\mathcal{Z}$  in apricis Teneriffæ (Cham.), ins. Archipelagi (Beaupré!), agri Hierosolymitani (Bové!), Hispaniæ, Galliæ australis. Italiæ mediter., Mauritanie. Desf. fl. atl. 2. p. 274. DC. fl. fr. 4. p. 173. *Ergeron viscosum* Linn. sp. 1209. Jacq. hort. vind. t. 165. *Solidago viscosa* Lam. fl. fr. 2. p. 144. — Clas. hist. 2. p. 20. f. 1. Bocc. sic. t. 7. f. B? Variat totis 3-9 lin. latis. (v. v. et s.)

37. *I. BARBATA* (Wall. ! cat. et herb. n. 71), caulibus erectis subsimplicibus glabris apice sparse pilosis, foliis oblongo-lanceolatis acuminatis basi vix attenuatis semiamplexicaulibus serrulatis ciliato-barbatis, capitulis sessilibus subsolitariis foliis cinctis, invol. squamis lanceolato-linearibus foliaceis long-barbato-ciliatis.  $\mathcal{Z}$  ad Srinaghur legit cl. Blindworth. Herba pedalis et sequiped. Radix fasciculata multicaulis. Capitula ampla ferè euphthalmoides. Ligulæ longissimè lineares stylo brevissimo subabortivè glabræ pollicem longæ sessilin. latæ. Pappus 1-serialis capillaris albus. Achenium parvum subpabernaculæ (v. s. comm. ab hon. Aul. merc. angl. Ind. or. et à cl. Royle.)

38. *I. VESTITA* (Wall. ! cat. et herb. n. 72), caule erecto ramoso apice præsertim cum foliis villosissimo, foliis semiamplexicaulibus lanceolatis caulibus brevibus apice acuminatis irregulariter serratis utrinquè villosis, pedunculis aphyllis molliter hirsutissimis, invol. squamis lineari-subulatis scabridè linearibus barbatis.  $\textcircled{1}$ ? ad Kamaon et Srinaghur legit rev. Blindworth. Herba pedalis à basi ramosa. Ligulæ stylo subabortivo fortè steriles. Pappus al-

capillaris 1-ser. Achænium parvum teretiusculum glabrum. Hieracium hispidum Don prod. fl. nep. 165 ex Spreng. in litt. An rectè? (v. s. comm. ab hon. Anl. merc. angl. Ind. or. et à cl. Royle.)

fl. minor, caule digitali aut subpalmari simplici oligocephalo. — ad Doab Wall. l. c. var. B. (v. s. comm. cum var. a.)

39. I. ? *NERVOSA* (Wall. cat. et herb. n. 70), caule erecto villoso subsimplici, foliis ovali-lanceolatis basi attenuatis apice acuminatis penninerviis utrinquè villosis scabridis serraturis callosis distantibus, capitalis paucis corymbosis terminalibus, pappi setis apice rigidis barbellatis. ½ in Nepaliâ legit cl. Wallich. Herba pedalis, caule tereti aut subcompresso. Invol. campanulati squamæ lanceolato-lineares exteriores subfoliaceæ dorso scabridæ. Ligulæ lineares angustæ glabræ. Achænium villosum teretiusculum. Pappus 1-serialis basi villis achæni quasi calyculatus rigidus basi subinteger, apice barbellis scaber. (v. s. comm. ab hon. Anl. angl. Ind. or.)

40. I. *ACUMINATA*, caule erecto ramoso teretiusculo puberulo, foliis caulinis glabris integerrimis basi dilatatis subauriculatis cordato-semiamplexicaulibus apice lanceolatis acuminatis, capitulis ad apices ramorum solitariis, invol. squamis linearibus foliaceis laxiusculis, achæntis glabris. ½ in Indiæ prov. boreali-occid. legit. cl. Royle. Paulò accedit ad I. spiræifoliam et I. Germanicam sed foliis ad basin dilatatis et ad apicem acuminatis faciliè differt. (v. s. comm. à cl. Royle.)

41. I. *JAPONICA* (Th. fl. jap. 318), caule erecto villoso tereti, foliis sessilibus sub-amplexicaulibus oblongo-lanceolatis serratis villosis, pedunculis axillaribus monocephalis subpaniculatis tomentosis, invol. squamis lineari-subulatis villosis. ½ in Japoniâ. Banks ic. Kæmpf. t. 30. Sen Fuku Kæmpf. amoen. 877. — Affinis videtur I. nervosæ Wall. sed differt foliis caulinis basi nec attenuatis nec subpetiolatis. Pappus indescr.

† *Species non satis notæ.*

42. I. *QUADRIDENTATA* (Lag. nov. gen. et sp. 30. n. 374), herbacea villosa-viscosa, foliis lanceolatis integris, floralibus linearibus, squamis invol. exterioribus laxis, corollulis 4-dentatis. ① in arenosis ruderalis cultis Hispaniæ ab urbe Alonensi ad Murciam usque. An vera Inulæ species? An fortè Francoeuræ sp.?

43. I. *PROVINCIALIS* (Linn. sp. 1241), caule erecto 1-cephalo, foliis subserratis subtus tomentosis, radicalibus ovatis, caulinis paucis lanceolatis sessilibus. ½ in Collendaniæ Galloprovinciæ. I. Provincialis Gon. videtur Senecio uniflorus, sed species Linnæana omninò dubia! An fortè Pulicariæ odoræ var.?

44. I. *MOLLIS* (Bernhardi in Link enum. h. ber. 2. p. 335), caule erecto tenuitomentoso, foliis serrulatis, radicalibus in petiolum attenuatis, caulinis lanceolatis acutis sessilibus hirtis, invol. squamis lanceolatis hirtis extimis reflexis. ½ Patr. ign. †.

45. I. ? *PRIMUMFOLIA* (Lam. dict. 3. p. 261), caule erecto herbaceo polycephalo, foliis spathulatis crenatis subtus incanis. — in pratis montosis Sancti-Domingi. — Plum. et Burm. t. 40. f. 1. Verisim. è genere excludenda, sed hodiernis ignota.

†† *Species exclusæ.*

I. *ACRIS* Bernh. = Erigeron acre.

I. *ÆSTUANS* Linn. = Verbesina? æstuens.

I. *APPENDICULATA* B. Wall. = Vicoa appendiculata.

I. *APPENDICULATA* C. Wall. = Vicoa Indica γ.

I. *ARABICA* Linn. = Pulicaria Arabica.

I. *ARGENTEA* Pers. = Chrysopsis argentea.

I. *AROMATICA* Linn. = Printzia aromatica.

I. *CANADENSIS* Bernh. = Erigeron Canadensis.

- I. *CAPENSIS* Spreng. = *Pulicaria Capensis*.
- I. *CAUCASICA* Pers. = *Telekia cordifolia*.
- I. *CRANUA* Berg. = *Printzia Bergii*.
- I. *CHRYSOCHOMOIDES* Poir. = *Jasonia Sicula*.
- I. *CÆRULÆ* Linn. = *Printzia Bergii*.
- I. *CONTRETA* Lam. = *Pulicaria dysenterica*.
- I. *CONYZOIDES* Desf. = *Pulicaria odora*.
- I. *CRISPA* Desf. = *Francoeuria crispa*.
- I. *CRISPA* Sieb. = *Pulicaria undulata*.
- I. *CYLINDRICA* With. = *Pulicaria vulgaris*.
- I. *DENTATA* Sibth. = *Pulicaria dentata*.
- I. *DIVARICATA* Nutt. = *Heterotheca Lamarckii*.
- I. *DUBIA* Thunb. = *Aster Japonicus*.
- I. *DYSENTERICA* Linn. = *Pulicaria dysenterica*.
- I. *FALCATA* Pursh. = *Chrysopsis falcata*.
- I. *FORTIDA* Linn. = *Erigeron foetidum*.
- I. *GLANDULOSA* Lam. = *Chrysopsis Mariana*.
- I. *GLUTINOSA* Pers. = *Grindelia glutinosa*.
- I. *GNAPHALODES* Vent. = *Strabonia gnaphalodes*.
- I. *GOSYPIA* Michx. = *Chrysopsis gossypina*.
- I. *GRAMINIFOLIA* Michx. = *Chrysopsis graminifolia*.
- I. *INCISA* Lam. = *Pulicaria incisa*.
- I. *INDICA* Linn. = *Vicoa Indica*.
- I. *LANATA* Banks. = *Chrysopsis gossypina*.
- I. *MACROPHYLLA* hortul. = *Telekia cordifolia*.
- I. *MARIANA* Linn. = *Chrysopsis Mariana*.
- I. *MONTANA* Biv. = *Pulicaria odora*.
- I. *ODORA* Linn. = *Pulicaria odora*.
- I. *PALUDOSA* Link. = *Pulicaria dentata*?
- I. *PINIFOLIA* Linn. = *Senecio pinifolius*.
- I. *FUNCTATA* Muhl. = *Heterotheca scabra*.
- I. *PULICARIA* Brot. = *Pulicaria dentata*.
- I. *PULICARIA* Linn. = *Pulicaria vulgaris*.
- I. *PULICARIA* d'Urv. = *Pulicaria dysenterica*.
- I. *QUADRIFIDA* Ham. = *Francoeuria crispa* β.
- I. *ÆREPENS* Fisch. = *Pulicaria uliginosa*.
- I. *SATURNIA* Spreng. = *Lorentea satureioides*.
- I. *SATURNEOIDES* Mill. = *Lorentea satureioides*.
- I. *SAXATILIS* Lam. = *Jasonia glutinosa*.
- I. *SCABRA* Nutt. = *Heterotheca scabra*.
- I. *SCABRA* Wall. = *Vicoa appendiculata*.
- I. *SERRATA* Pers. = *Grindelia inuloides*.
- I. *STOLONIFERA* Gay. = *Aplopappus stoloniferus*.
- I. *SUBAXILLARIS* Lam. = *Heterotheca Lamarckii*.
- I. *TRICHOPHYLLA* Nutt. = *Chrysopsis trichophylla*.
- I. *TRIXIS* Linn. = *Trixis frutescens*.
- J. *TUBEROSA* Lam. = *Jasonia tuberosa*.

1. *ULIGINOSA* Stev. = *Pulicaria uliginosa*.

1. *UNDULATA* Linn. = *Pulicaria undulata*.

1. *VILLOSA* Vahl. = *Pulicaria Arabica*.

CCXXXIV. *SCHIZOGYNE* Cass. dict. 56. p. 23. Less. syn. 123. — *Conyza* et *Chrysocoma* sp. auct. — *Gnaphalium* sp. Poir.

Capitulum multiflorum heterogamum, fl. omnibus tubulosis, marginalibus paucis 1-seriatis tenuibus bidentatis fœmineis, cæteris 5-dentatis hermaphroditis. Recept. nudum angustum. Invol. oblongo-obconicum imbricatum, squamis colorato-scariosis oblongis acutis serrulatis glabris ciliolatis. Antherae basi breviter bisetæ. Achænium oblongo-cylindraceum glabriusculum. Pappus 1-serialis setosus scabridus. — *Frutex* Canariensis ramosissimus. Folia sparsa sessilia linearia obtusa integerrima. Corymbi terminales subpaniculati polycephali. Pedicelli squamulosi. Invol. citrina. Flores intensius lutei. — Genus differt à *Conyza* et *Phagnalo* floribus fœmineis uni- nec multiserialibus, à *Chrysocoma* floribus non omnibus hermaphroditis. Ex antheris bisetosus-caudatis accedere videtur ad *Inulam* sed faciliè differt capitulis non radiatis, involuacro imbricato, etc.

1. *S. SERICEA*, ramis folisque tomento albo subsericeo undiquè obtectis. ♂ in insulis Canar. Teneriffâ, Palmâ, magnâ Canariâ legerunt et mecum comm. cl. Collignon, Ledru, Broussonet, Bory, Chr. Smith, Webb. et Berthelot, fortè etiâ ex h. Puer. in insulis Azoricis sed non in ins. Mauritiî aut ad Cap. Bonæ-Spei quidquid dixerit cl. Cassini. *Conyza sericea* Ait. kew. 3. p. 183. L'Hér. ! descr. mas. Link. Buch! can. 165. *Chrysocoma sericea* Linn. f. suppl. 360. *Conyza tenuifolia* Salisb. prod. 194. *Conyza canescens* Bory ! in litt. Sch. obtusifolia Cass. l. c. ips. test. in h. Merat. *Gnaphalium multiflorum* Poir. ! suppl. 2. p. 803 (ex specim. Ledru) et idèò *Gnaph. Teneriffæ* Spreng. syst. 3. p. 468. *Gnaphalium compressum* Poir. suppl. 2. p. 804 (ex specim. Brousson.) ipso auctore teste et autogr. in h. Desf. ! Variat involucri glaberrimis aut extûs subaraneosis. Cortex sapore acri contra odontalgiam ab incolis usitatus ex Linn. (v. s.)

2. *S. GLABERRIMA*, ramis folisque glaberrimis. ♂ in insul. Canariensisbus verisim. Teneriffâ. *Conyza Chrysocomæ linosyridi* affinis Webb et Berth. ! herb. Priori simillima et fortè varietas. Differt tamen glabritie perfectâ, foliis paulò brevioribus, corymbis oligocephalis. (v. s. in h. Webb. et Berth.)

• CCXXXV. *VARTHEMIA*.

Capitulum multiflorum homogamum, fl. omnibus tubulosis 5-dentatis hermaphroditis. Recept. alveolatum. Invol. obovatum imbricatum, squamis adpressis acutiusculis gradatim longioribus, intimis disco æqualibus. Antherae basi bisetæ. Stigmata brevia inclusa. Achænium compressum oblongum pubescens. Pappus 1-serialis, setis subscabridis. — *Suffrutex* Persicus erectus ramosissimus glaberrimus. Folia sparsa paucalinearum aut oblongo-linearum integerrima patulo-deflexa. Capitula ad apices ramorum solitaria. Flores lutei. — Ex capitulis homogamis ad *Chrysocomam*, ex antheris basi bisetis ad *Inulam* accedit. — Genus dicavi veteri peregrinatori Lud. de Varthemo, qui ineunte seculo sexdecimo Persiam et Orientem ferè totum peragravit.

1. *V. PERSICA*. ♂ in Persiâ prope Teheran leger. cl. Olivier et Bruguière et recentius cl. Belanger. (v. s.)

CCXXXVI. VICOA *Cass. ann. sc. nat.* 1829. p. 33. *dict.* 60. p. 596. *Less. syn.* 199. *Inulæ* sp. *Linn.*

Capitulum multiflorum heterogamum, fl. radii angustè ligulatis fœmineis uniserialibus, disci tubulosis 5-dentatis hermaphr. Invol. imbricatum, squamis lineari-subulatis. Recept. ebracteolatum arcolatam plano-subconvexum. Antheræ basi setoso-appendiculatæ. Achænium erostre. Pappus radii nullus, disci capillaris 1-serialis. — Herbas Indice annuæ ramosæ erectæ. Folia alterna sagittato-auriculata semiamplexicaulia lanceolata aut linearia integra aut dentata. Rami parçè foliosi apice 1-cephali. Capitula flava.

1. *V. AURICULATA* (Cass. Il. cc.), foliis subdentatis oblongis acutis basi auriculatis, ligulis disco subæqualibus. ① in Ceylonâ.

2. *V. INDICA* (DC. in Wight contrib. bot. ind. p. 10), foliis basi auriculatis lanceolatis acuminatis serratis subintegerrimisve utrinquè plûs minûs puberulis, ligulis disco duplò longioribus. ① per totam Indiæ orientalis peninsulam fræuens. Capitula 3 lin. diam. (v. s.)

a. *serrata*, foliis per totam longitudinem serratis. Inula Indica Willd. sp. 3 p. 2092. Wight! herb. Inula Indica var. F. Wall.! cat. n. 74. Folia basi latè dilatato-auriculata in parte dilatata præsertim dentata.

β. *subintegra*, foliis subintegerrimis aut hinc inde serratis. — Inula Indica Linn. sp. 1237. Wall. cat. n. 74. — *Doronicum calcaratum* Roxb. fl. ind 3. p. 434 ex Wight. Burm. ceyl. 124. t. 55. f. 2. Pluk. alm. t. 149. n. 4. f. 3. — *Aster Indicus* Willd. sp. 3. p. 2041 non Linn. ex Wight contrib. bot. ind. p. 10.

γ. *attenuata*, foliis subintegerrimis inferioribus basi angustatis ferè petiolatis nec auriculatis. — Inula appendiculata var. C. Wall.! cat. n. 76 lecta ad Promæ.

δ. *aspera*, foliis subintegerrimis omnibus auriculatis suprâ sparçè et rigide setosis, subtus puberulis. Inula Indica var. G. Wall.! herb.

3. *V. AURITA*, foliis subintegerrimis omnibus basi latè et obtusè auriculatis angustis suprâ densè setoso-echinatis subtus pubero-subtomentosis, ligulis disco paulò longioribus. — Inula auriculata Wall.! herb. n. 75. *Erigeron linearis* Spreng. in litt. Valdè accedit ad *V. Indicam* sed ex pubescentiâ subtomentosi vid. diversa. (v. s. comm. ab hon. Cur. merc. angl. Ind. or.)

4. *V. APPENDICULATA*, foliis linearibus basi auriculatis acuminatis subintegerrimis margine revolutis utrinquè minutissimè et densè puberulis, ligulis disco duplò longioribus. ① in Indiâ orientali ad ripas excelsas flum. Yenangham. Inula scabra Wall.! cat. h. bot. Inula appendiculata Wall.! cat. et herb. n. 76. var. β Folia 3 poll. longa, 2 lin. lata. Capitula radio excepto 6-7 lin. diam. et idè priori dupla. (v. s. comm. ab hon. Aul. angl. Ind. or.)

CCXXXVII. PENTANEMA *Cass. bull. philom.* 1818. p. 75. *dict.* 23. p. 565. β p. 373. *Less. syn.* 198.

Capitulum multiflorum heterogamum, fl. radii 1-ser. fœmineis angustè ligulatis puberis apice 3-dentatis, fl. disci tubulosis hermaphr. 5-dentatis. Recept. ebracteolatum subconvexum. Invol. subglobos squamæ lineares imbricatæ villis longis mollibus hispidæ. Achænium erostre hispidulum. Pappus 1-serialis, setis 5 capillaribus. — Herba Orientalis annua erecta gracilis molliter hispida. Rami divaricati rubentes. Folia alterna obovata aut oblonga obtusa integerrima parçè villosa membranacea. Capitula pedunculata oppositifolia subglobosa flava. Nomen ex πέντε, quinque, et νημα, filum, et ἐργὸ neutri ne firminini generis.

1. *P. DIVARICATUM* (Cass. ll. cc.) (I) in Oriente sine loci design. legit cl. Michaux, et postea inter Alep et Bagdad cl. Olivier et Brugnière. Capitulum omnino *Pallcaris trichocephalæ* sed pappus diversus. (v. a.)

CCXXXVIII. *FRANCOEURIA* Cass. dict. 34. p. 44. v. 38. p. 374. Less. syn. 198. — *Duchesnia* Cass. dict. 13. p. 545 non Smith et excl. car. — *Inula* sp. Pers. — *Asteris* sp. Forsk.

Capitulum multiflorum heterogamum, fl. radii ligulatis breviusculis 1-ser. fœmineis, disci tubulosis 4-dentatis hermaphr. Recept. nudum planum. Invol. campanulatum imbricatum, squamis lineari-subulatis. Anthæræ basi setoso-appendiculatæ. Achænium erostre. Pappus caducus, setis 1-serialibus infernè subscabris supernè barbellato-subplumosis basi conferruminatis in annulum brevissimum setulis paleolisve minimis coronatum. — Suffruticulus Africanus erectus. Rami teretes 1-cephali divergentes. Folia alterna semiamplexicaulia oblonga dentata. Capitula flava.

1. *F. CRISPA* (Cass. ll. cc.), foliis margine undulato-crispatis.  $\frac{5}{8}$  et sæpè (I). In arenosis apricis Senegaliz et Egypti. *Helenium Ægyptiacum*, etc. Vaill. act. par. 1720. p. 304. n. 24. *Aster crispus* Forst. æg. ar. descr. p. 150. I. crispa Pers. ench. 2. 450. excl. syn. Vent. Delille! fl. eg. p. 25. t. 45. f. 2. *Duchesnia crispa* Cass. dict. 13. p. 545. *Conyza undulata* Poir. suppl. 2. p. 339? Variet subherbacea aut fruticosa, tota subglabrata aut densè tomentoso-alba, foliis angustioribus latioribus semper margine crispatis. (v. a.)

$\frac{1}{2}$ . *Indica*, foliis margine planiusculis. — ad Bar in prov. Bahar prope ripas Gangis legit cl. Hamilton. *Inula quadrifida* Ham. herb. Wall. cat. n. 73. (v. a. comm. ab hon. Aulâ merc. angl. Ind. or.)

CCXXXIX. *IPHIONA*. — *Iphionæ* sp. Cass. — *Chrysocomæ* sp. Forsk. — *Conyzæ* sp. Lam. — *Stæbelinæ* sp. Vahl.

Capitulum multiflorum homogamum, fl. omnibus hermaphroditis tubo stricto sensu subnullo, fauce longâ cylindricâ, dentibus 5. Recept. alveolatum angustum. Invol. imbricatum obovato-oblongum, squamis adpressis acuminatis seu ovalibus mucronatis. Anthæræ basi breviter setoso-caudatæ. Filam. imæ cor. insertæ! Achænium subcylindraceum striato-sulcatum parcè hispidum. Pappus rigidus multiserialis subscaber, setis exter. brevioribus, intimis gradatim longioribus corollam æquantibus. — Fruticuli Ægyptiaci glabri ramosissimi. Folia sparsa pauca subulata rigida pungentia basi auriculis 1-2 pungentibus instructa. Capitula ad apices ramorum subnudorum solitaria lutea. Pappus rufescens.

1. *I. JUNIPERIFOLIA* (Cass. dict. 23. p. 610), tota glabra, foliis apice subulato-pungentibus, inv. squamis obtusis mucronatis.  $\frac{5}{8}$  in Ægypto prope Cahirum inter urbem et Soneys, verus Ramalieh, etc. (Delil.! Acerb.! Radd.!), in valle Hebran (Rupp.), in Arabiæ desertis circa montem Sinai (Boré!). Arabicè *Dafry* in Ægypto, *Daffara* in Arabiâ dicta. *Conyza Memphitica*, etc. Vaill. act. par. 1719. p. 301. *Chrysocoma mucronata* Forsk. descr. 147. Fres. beitr. p. 77. *Stæbelina spinosa* Vahl symb. 1. p. 69. *Conyza pungens* Lam. dict. 2. p. 86. *Chrysocoma spinosa* Delil. fl. eg. p. 128. t. 46. f. 3. *Conyza pungens* et *Stæbelina spinosa* Pers. *Serratula spinosa* Poir. Iph. dubia Cass. bull. philom. 1817. oct. Jasonis sp.? Less. syn. 195. Folia variant basi uni- aut bispinosa et rarius in medio adhuc 1-spinosa, summa semper squamiformia integerrima. (v. s. comm. à cl. Acerbi et Raddi.)

2. *I. SCABRA* (DC. in ann. sc. nat. 1834. bot. p. 263), tota pube brevi glandulosa

scabrâ, foliis subulato-pungentibus, invol. squamis lineari-lanceolatis glabris.  $\zeta$  in Ægypti deserto circa Suez et in Arabiâ circa Tor mense junio flor. legi. cl. Bové. Chrysocoma? Bové! pl. exs. n. 104. Arabis dicitur *Saffara* ex Bové. Priori habitu similis sed folia et capitula majora. Caulis ramosissimos. Folia lineari-subulata pungentia utrinquè ad basin dentibus 1-3 spinosis seta. Achænia villosissima. (v. s. comm. à cl. Bové.)

† *Species exclusa.*

I. PUNCTATA Cass. = *Inula Africana*.

CCXL. JASONIA. — Jasonia Chiliadenus et Allagoppus Cass. dict. 24. (1825) 34 (1825) et 56 (1828). — Jasonia Allagoppus et Donizæ sp. Less. syn. 19. — Inulæ et Erigerontis sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœminis uniserialibus aut homogamum fl. omnibus tubulosis 5-dent. hermaphrod. Recept. nudum subalveolatum planum. Invol. imbricatum disco circ. æquale, squamis linearibus acutis. Antheræ basi setis cædatæ. Achænia villosum oblongo-cylindraceum basi attenuatum Pappus biserialis utrinquè setis filiformibus barbellatis, exterioribus brevissimis, inter. corollam ferè æquantibus. — Herbæ aut suffrutescunt humiles in regione Mediterraneâ aut Canariensi indigenæ, sæpius pilosæ aut glandulosæ ramosæ, ramis patentibus apice 1-cephalis. Folia altera sessilia oblonga aut linearia integra aut dentata. Flores lutei. — An genus in quatuor divellendum?

Sect. I. CHILIADENUS Cass. dict. 34. (1825) p. 34. — Myriadenus Cass. dict. 23. p. 565 non Desv. — Orsina Bertol. giorn. h. nat. 1829. v. 1. p. 362.

Capitula discoidea homogama. Achænia basi villosa apice glandulosa. Pappus inter. multi- (10-15) setosus. — Caules herbacei.

1. J. GLUTINOSA, herbacea erecta glutinoso-pubera, foliis lineari-lanceolatis integerrimis sæpè obliquè tortis, ramis ad apicem usquè foliosis.  $\zeta$  in apricis rupestribus Galloprovinciæ, agri Ruscinonensis (Benth.), Catalauniæ (Poir.) Siciliæ, Lopedusæ et Melitæ (Guss.). Erigeron glutinosum Linn. mant. 1. Inula saxatilis Lam. fl. fr. 2. p. 153. DC. fl. fr. ed. 3. v. 4. p. 154. Chrysocoma camphorata Rob. et Cass. in litt. Chrysocoma saxatilis DC. fl. fr. suppl. p. 168. Chiliadenus camphoratus Cass. ! dict. 34. p. 35. Orsina camphorata Bertol. giorn. sc. nat. Bol. v. 1. 1829. p. 362. opt. ex Guss. ! — Bocc. sic. t. 13. f. 4. ideo Inula fetida Zeraph. fl. melit. prod. p. 25. ex Bertol. — Barr. ic. t. 134 (v. s.)

Sect. II. EUJASONIA. — Jasonia Cass. dict. 24 (1822). p. 200.

Capitula radiata aut subdiscoidea. Achænia teretiuscula undiquè villosa. Pappus interior multi- (10-20) setus.

2. J. TUBEROSA, suffrutescens erecta, radice præmorsâ incrassato-tuberosâ lignosâ, foliis linearibus aut sublinearibus integerrimis, ramis parvè foliosis, involucri longioribus.  $\zeta$  in apricis Galliæ mediterraneæ, Hispaniæ, Syriæ, etc. Erigeron tuberosum Linn. sp. 1212. Willd. sp. 3. p. 1961. Inula tuberosa Lam. dict. 3. p. 260. DC. fl. fr. ed. 3. v. 4. p. 153. Jasonia radus Cass. dict. 24. p. 201. Aster punctatus Lapeyr. abr. pyr. p. 513. excl. syn. a Benth. — Lobel ic. t. 250. f. 3. Variat caule vix palmari et culta pedali et quipedali. (v. v.)

3. J. SICULA (DC. in ann. sc. nat. 1834. bot. p. 261), herbacea erecta subulata.

radice gracili, foliis radicalibus lanceolatis dentatis, caulinis linearibus basi auriculatâ semiamplexicaulibus integerrimis ramis parvis foliosis. ☉? ♀ in paludosis maritimis arenosis Cretæ, Siciliæ, Hetruriæ, Galliæ et Hispaniæ mediterraneæ, Mauritanie et ad montem Sinai (Bové! n. 118). Caulis 1-x  $\frac{1}{2}$  pedalis ramosus purpurascens. Rami paniculato-corymbosi. Pappus biserialis, serie extimâ brevissimâ.

*a. discoidea*, floribus marginalibus subtubulosis aut in ligulam involuero brevioribus fissis. — *Erigeron* Siculum Linn. sp. 1210. *Conyza* Sicula Willd. sp. 3. p. 1931. DC. fl. fr. 4. p. 129. Poir. suppl. 2. p. 358. *Jasonia* discoidea Cass. dict. 24. p. 201. et 39. p. 406. *Inula* chrysocomoides Poir. voy. 2. p. 239. Lam. dict. 3. p. 261. ill. t. 680. f. 3. Desf. atl. 2. p. 275. *Tubilium* Siculum Choisy! in h. DC. ined. — Boeck. sic. t. 31. f. 4. Magn. monsp. 76. io. Pluk. alm. t. 168. f. 2. Species ab auctoribus sæpè confusa et characteres radii reverà variantes et fallaces. An *Dimorphanthès* Sicula Cass. dict. 13. p. 255. huc revocanda sed excl. car. gener.? (v. v. et s.)

*β. radiata*, floribus radii in ligulam involuero longiorem productis. — in Galliâ Narbonensi et Mauritaniâ Tingitanâ. DC. fl. fr. 4. p. 129. Salzmann! pl. exs. *Aster* pyrenaicus Pourr.! in h. L'Hér. excl. syn. *Ligulæ* luteæ ferè similis priori. (v. s.)

Sect. III. ALLAGOPAPPUS Cass. dict. 56. p. 21. Less. syn. 197.

Capitulum discoideum homogamum. Achænium angulatum ad angulos villorum. Pappus inter. punci- (5-7) setus, setis filiformis scabris, exter. brevis paucisetus. — Caules fruticosi. Folia glandulosa. An genus proprium?

4. *J. dichotoma*, fruticosa subdichotomè ramosa glabriuscula, foliis linearibus basi attenuatis apice serratis, ramis floriferis squamas plurimas lineari-subnatas gerentibus, ligulis nullis. ♀ in insula Canariis, Palmâ, Gran-Canariâ et frequentior in insulâ Teneriffâ ad rupes Barancorum legerunt Ledru! Chr. Smith! Broussonet! Webb et Berthelot! nec in insulâ Mauritii ut incautè dixit Cass. ex h. Merat. *Conyza* dichotoma Linn. f. suppl. 359. *Chrysocoma* dichotoma Jacq. ic. rar. 1. t. 171. *Conyza* inuloides Ait. kew. 3. p. 182. *Conyza* Canariensis Willd. sp. 3. p. 1937. Link in Buch can. 148 et 165. *Allagopappus* dichotomus Cass. l. c. exo. patr. Antheris basi, jàm teste Aitonio, bisetis à Conyzâ et Chrysocomâ differt. Pappo biseriali extimo minimo pluri-seto ad *Eriasoniam* accedit. Folia utrinquè multiglandulosa. (v. s.)

Sect. IV. DONIOIDES. Doniæ sp. Less.

Capitulum radiatum, ligulis fœmineis. Achænium subangulatum hispidum. Pappus interior 7-10-setus, setis crassis ciliato-scabris, exterior pluri-setus brevissimus.

5. *J. lævigata*, fruticosa glabra lævis, foliis oblongo-lanceolatis subcoriaceis integerrimis aut apice grossè serratis, pedunculis subaphyllis. ♀ in ins. Teneriffâ ad Caput Teno rarior Broussonet! Webb et Berthelot! Folia basi subtriner-via 1-2 poll. longa, 7-12 lin. lata. Inv. squamæ foliaceæ disco ferè longiores. *Rophthalmum* lævigatum Brouss. herb. Willd. mag. nat. ges. Berl. 1807. p. 138. *Donia* Canariensis Less. syn. 199. (v. s.)

CCXLI. PULICARIA. — Inulæ sp. Linn. — Pulicaria et Tubilium Cass. Less.

Capitulum multiflorum heterogamum, fl. radii fœmineis 1-serialibus ferè semper ligulatis, disci tubulosi 5-dent. hermaphr. Recept. nudum areolatum planiusculum. Invol. laxè imbricatum pauciserialè, squamis linearibus. Achænium erostre puberulum teretiusculum nec compressum. Pappus biserialis, serie exteriori coroniformi dentatâ brevissimâ, inter. setis 10-20 scabridis. — Herbæ pleraque Europææ nonnullæ Africanæ, in uliginosis præsertim nascentes, villosæ odoratæ



erectae ramosae. Folia caulina cordato-sagittata integra aut dentata lanceolata. Pedunculi apice 1-cephali. Capitula flava.

Sect. 1. PULICARIA. — *Pulicaria Gært. fr. 2. p. 461. t. 173. Cass. dict. 23. p. 565 et 44. p. 93. Less. syn. 197.*

Capitula radiata, fl. angustè ligulatis.

1. *P. TRICHOCEPHALA*, caule demisso à basi ramosissimo cespitoso, foliis lanceo-sublanceolatis integerrimis acutis sessilibus pubescentibus, capitulis parvis foliis bracteatis lateralibus subsessilibus et terminalibus subglobosis, invol. squamis linearibus villosis, radii ligulis disco paulò longioribus, pappo ext. dentato, inter. 10-seto. (†) in Oriente sed locus propr. ignotus fertè Syria. Valdè accedit ad *P. Arabicam* sed differt præcipuè involucri villis longioribus hirsutissimo fertè ut in *Pentstemonate*. Herba vix digiti longitudo. Radix perpendicularis subsimplex. (v. s.)
2. *P. ARABICA* (Cass. dict. 44. p. 94), caule erecto divaricato-ramoso, foliis oblongis cordato-semiamplexicaulibus mucrone acutis utrinque pilis minus pubescenti-villosis integerrimis, pedunculis axillaribus terminalibusque, invol. squamis lineari-undulatis puberulis, radii ligulis disco paulò longioribus, pappo ext. dentato, inter. 10-seto. (†) in Arabia? (Forsk. ex Linn.), in cæcis montosis ad Sinai (Bové), Egypto (Delil. Coqueb.), Creta (Oliv. & B.), Mauritania (Desf. Seism.), Andalusia (Hasselt), Inula arabica Linn. mant. 114? (excl. syn. Pluk. et ideo petr. in India India.) Desf. fl. et. 1. p. 276. Inula villosa Vahl ex Horn. h. hafa. 2. p. 82a non Nutt. Pul. villosa Link. enum. h. herol. 2. p. 336. Variat caule decumbente aut subdiffuso. Capit. 5 lin. diam. Valdè dubium mihi videtur stirpem in Arabia habitare. Fossilis in florâ suâ Arabicâ hanc non enumerat. Desor. Linnæi est etiam valdè rara. (v. s.)
3. *P. DESERTORUM* (DC. ann. sc. nat. 1834. bot. p. 263), caulibus erectis ramosis pubescenti-villosis, foliis semiamplexicaulibus oblongis grossè et obtusè dentatis undulato-crispis villosis, pedunculis 1-cephalis, capitulis depressis hemisphaericis, invol. squamis linearibus acuminatis, ligulis invol. paulò longioribus, pappo ext. brevissimo subintegerrimo, inter. 10-seto. (†) ? in Arabia desertis circa montem Sinai legit cl. Bové mense junio flor. Arabica dicitur. Inula Arabica Bové! pl. exs. n. 109. Ex loco natali est verè Arabica sed differt à stirpe sub hoc nomine adhuc admittit et fortè in Arabia usquam repertâ. (v. s.)
4. *P. VULGARIS* (Gært. fr. 2. p. 461. t. 173), caule paniculato erecto, foliis semiamplexicaulibus oblongis acutis undulatis subintegerrimis villosis, pedunculis 1-cephalis oppositifoliis, capitulis hemisphaericis, radii ligulis brevissimis, pappo ext. dentato, inter. circ. 15-seto. (†) in uliginosis viis et plateis Europæ temperatæ et australis naque ad Caucasum (Bieb. Stev.), et ad Sibiriam Altham prope Semipalatinsk (C. A. Mey.), in Teneriffâ fortè introducta (Broom). Inula Pulicaria Linn. sp. 1238. Fl. dan. t. 613. Engl. bot. t. 1196. Aster panicarius Scop. carn. n. 1080. Inula cylindrica With. Huds. et Inula uliginosa Sibth. ex Sm. fl. brit. 892. Conyza minima Dod. pempt. 52. Variat capitulis flosculosis ex Sm. fl. brit. Variat præcipuè invol. in region. frigidioribus glabrisculis aut molliter hirsutis in reg. calidioribus. An fortè species hæc adhuc confusa? (v. v. et s.)
5. *P. URGENTOSA* (Stev. in litt.), caulibus erectis apice tantum paniculato-ramosis, foliis caulinis cordato-auriculatis semiamplexicaulibus lanceolatis acutis subserratis subtus præsertim velutino-tomentosis, capitulis pedunculatis terminalibus et axillaribus, invol. squamis lineari-subulatis, radii ligulis duplò fertè longioribus, pappo ext. dentato, int. 15-seto. (†) in Tauri montionall ad fossas aquosas legit cl. Steven. Eandem ni fallor habeo lectam à ripas Jordanis et prope Beyrouth in Bové pl. exs. n. 441? *P. dysenterica* Rab.

suppl. p. 574 ex Stev. *Inula repens* Fisch. ex Trevir. act. nat. cur. 13. p. 203. Ferè media inter *P. vulgarem* et *dysentericam*. (v. s. comm. à cl. Steven.)

6. *P. angustifolia*, tota subcinereo-pubescenti, caule basi suffruticulosæ ramosæ, foliis sessilibus oblongo-linearibus integerrimis aut hinc inde dentatis, pedunculis subnudis 1-cephalis folio triplò longioribus, invol. squamis foliaceis, ligulis invol. longioribus, pappo ext. denticulato, inter. 10-12-seto.  $\frac{3}{4}$  in India prov. boreali-occident. legit cl. Royle. Accedit ad *P. uliginosam* sed foliis sessilibus nec cordato-auriculatis capitulis longius pedunculatis et invol. squamis latioribus faciliè differt. (v. s. comm. à cl. Royle.)

7. *P. dysenterica* (Gærtn. fr. 2. p. 462), caule paniculato erecto, foliis caulinis aurito-amplexicaulibus ovali-oblongis subintegerrimis callosæ-acutis subtus incano-tomentosis, pedunculis axillaribus et terminalibus 1-cephalis, squamis invol. apice setaceo-cuspidatis, radii ligulis disco duplò triplòve longioribus, pappo ext. dentato, inter. circ. 20-seto.  $\frac{1}{2}$  in Europæ ferè totius uliginosis, fossis, humidis. Cass. dict. 44. p. 93. *Inula dysenterica* Linn. sp. 1237. DC. fl. fr. 4. p. 149. Fl. dan. t. 410. Engl. bot. t. 1115. Hayn. arzn. 8. t. 46. Schkuhr hand. t. 247. ic. fr. *Aster dysentericus* All. pedem. n. 711. *Inula conyzæ* Lam. fl. fr. 2. p. 149. *Aster undulus* Mœnch. *Inula pulicaria* D'Urv.! enum. n. 816. excl. syn. Variat foliis longioribus concoloribus. An fn. planifolia Link ab hac satis differt? — Fuchs hist. 436. ic. (v. v.)

$\beta$ . *breviradiata*, ligulis invol. paulò longioribus aut eo subæqualibus.  $\frac{1}{2}$  circa Genevam ad pagum Vezenaz obs. Reuter. (v. s.)

8. *P. undulata*, albo-lanuginosa, caule erecto paniculato, foliis cordato-auriculatis semiamplexicaulibus lanceolatis undulato-crispis dentatis, pedunculis squamiferis terminalibus et axillaribus, invol. squamis linearibus acutis subcahris, radii ligulis disco paulò longioribus, pappo ext. dentato, inter. 15-20-seto. (12) in desertis Egypti. *Inula undulata* Linn. mant. 115. Lam. dict. 3. p. 256. Delil. fl. eg. 126. t. 46. f. 1. excl. var.  $\beta$ . *Helenium Egyptiacum tomentosum*, etc., Vaill. excl. Par. 1720. p. 305. *Inula crispata* Sieb. herb. Egypt. I. odora Forsk. fl. eg. p. 150? (v. s.)

9. *P. incisa*, pubero-villosa, caule erecto paniculato, foliis basi dilatata cordato-semiamplexicaulibus oblongo-linearibus acutis plus minus incisè dentatisve planiusculis, pedunculis apice subaphyllis terminalibus et axillaribus, invol. squamis linearibus acutis glabriusculis, radii ligulis disco paulò longioribus, pappo exter. dentato, inter. 15-seto. (1) in Senegalià propè St.-Louis legerunt cl. Roussillon et Perrottet. *Inula incisa* Lam. dict. 3. p. 256. In. undulata  $\beta$  Pera. Delil. Valdè affinis I. undulatæ sed nec albo-lanuginosa nec undulatifolia. (v. s.)

10. *P. odora* (Reich. fl. exc. 2. n. 1532), caule erecto subsimplici oligocephalo, foliis villosis remotè denticulatis, radicalibus petiolatis ovatis, caulinis auriculato-amplexicaulibus oblongis lanceolatis, pedunculis subternis axillaribus et terminalibus, radii ligulis disco duplò longioribus, pappo ext. 10-dentato, inter. 10-seto.  $\frac{1}{2}$  in apricis agri Ruscinonensis (Duby bot. gall. 1. p. 268). Galloprovinciæ, Liguriæ, Istriæ (Reich.), Græciæ (Sibth.; h. Acad. mon.!). Aleppi, Constantinopoli, Etruriæ, agri Neapolitani (Ten. syll. p. 464), Siciliæ (Guss.), Ischiæ (Schonv.), Mauritaniz, Corsicæ, nec in Burgundiâ ut falsò indicavit Durande. *Inula odora* Linn. sp. 1236. *Aster odoratus* All. ped. n. 713. *Inula dumetorum* Salzm. herb. — Barr. ic. t. 1145. bona. An fortè huc referenda *Pulicaria calycina* Presl. del. 96, quæ *Inula montana* Rivon. rar. cent. 1. p. 34. non Linn. et *Inula calycina* Spreng. syst. 3. p. 521 et I. conyzoides Desf. cor. Tourm. p. 49. t. 38? quæ *Aster Creticus conyzoides* flore magno Asphodeli radice Tourm. cor. 36. (v. v.)

11. *P. capensis*, caule erecto basi simplici apice corymboso foliisque præsertim superioribus utrinquè densè sericeo-villosis, foliis cordato-amplexicaulibus mucrone callosæ apiculatis et callosæ-serrulatis, invol. squamis lineari-acuminatis laxè imbricatis pedunculisque villosissimis, ligulis linearibus involucri paulò longioribus, achæniis villosis, pappo ext. 10-dentato inter. 15-seto.

℥ ad Cap. Bonæ-Spei ad fossas juxta montem Diaboli (Eckl. ! it.), in distr. Worcester (Eckl. !), ad Kleindrakenstein (Drege !), ad Port-Natal (Drege !), et in coloniz distr. orient. (Burch. ! cat. geog. n. 4407). Herba 1-2-pedalia. Capitula 5-6 lin. diam. Variat foliis caulibusve plus minus sericeo-tomentosis et incanis. An fortè duæ species distinguendæ? Species simillima *P. inuloides* radiata. (v. s.)

12. *P. ERIGEROIDES*, caule erecto basi simplici apice subpaniculato pubescente, foliis semiamplexicaulibus oblongo-lanceolatis remotè denticulatis aut subintegerrimis minutissimè puberulis, invol. squamis lineari-subulatis subcabratis, ligulis angustis parvis, pappo exter. 10-dentato, acheniis vix pilosisculis. ℥ ? in Africâ Capensi Burch. ! cat. n. 4408 et 1949. Eckl. ! Drege ! *Erigeron scabrum* Thunb. fl. cap. 665. Affinis *P. vulgari* et *P. Capensi* sed ab utraq. diversa. (v. s.)

13. *P. DENTATA*, caule erecto ramosissimo, foliis amplexicaulibus acuminatis dentatis, capitulis hemisphericis, radii ligulis longitudine involucri. ℥ in insulâ Crêtâ. *Inula dentata* Sibth. fl. græc. t. 874 (ined.) ex Smith prod. 2. p. 11. *Inula pulicaria* ß Linn. sp. 1238. syn. dub. An fortè in. *paludosa* Link, seu la. *Pulicaria* Brot. huc referenda? †.

Sect. II. *TUBILIUM* Cass. bull. philom. 1817. p. 153. dict. 23. p. 565. v. 56. p. 19. Less. syn. 197.

Capitulum discoidem, fl. radii 3-5-fidis stamina efforta sæpè gerentibus. Achenia glabrescens. — Sectio cum *Pulicariâ* conjungenda eodem jure ac *Corva squarrosa* cum *Inulâ*.

14. *P. INULOIDES*, caule herbaceo erecto puberulo, foliis semiamplexicaulibus obtusè auriculatis lineari-oblongis integerrimis aut dentato-subincisis vix pubescentibus, ramis 1-cephalis paucis subcorymbosis, invol. squamis lineari-subulatis villosis, fl. radii irregulariter tubulosi cæteros paulò superantibus. ℥ ad Mogador (nec in Teneriffâ) legit cl. Broussonet ! *Erigeron inuloides* Poir. ! suppl. 5. p. 664. *Tubilium angustifolium* Cass. ! dict. 56. p. 20. Similium *P. Capensi* et etiã ad *Conyzam Gonani* habitu accedens. Fol. 2 poll. long. 2 lin. latu. (v. s. comm. à cl. Broussonet.)

15. *P. FOLIOLOSA*, tota cinereo-villosa, caulibus herbaceis ramosissimis apice molliter hispidis, foliis oblongis acutis utrinquè puberulis, inf. basi attenuatis subdentatis, super. integris basi auriculatis semiamplexicaulibus, invol. squamis lineari-subulatis villosis, fl. radii tubulosi tenuissimis cæteris ferè brevioribus, pappo ext. coroniformi dentato, inter. 5-10-seto. ℥ ? in Indiâ oriet. prope Patna legit cl. Hamilton, et in prov. bor.-occident. cl. Royle. *Corva foliolosa* Ham. ! in Wall. cat. n. 3281. comp. 391 non 3092 nec Willd. An setas ex cor. minutie non vidi sed ex habitu et pappo hic refero speciem. Cor. apice subglandulosa. (v. s. in h. Cor. Ind. unne Soc. Linn. Lond.)

16. *P. HYDRABADENSIS* (Lindl. ! in litt. 1835), caule herbaceo erecto tereti ramoso subpubero, foliis lanceolatis puberulis, infer. obtusis dentatis basi in petiolo attenuatis, super. sessilibus angustis acutis subintegris, ramis parvis foliis apice monocephalis, invol. squamis linearibus acuminatis, exter. foliis apice puberulis, inter. siccis glabris. ① in Indiâ orient., ad Hyderabad (ex nom. ad muros Seringapatam (h. madr. !). *Aster Hydrabadensis* Wall. ! cat. n. 29. ! comp. n. 83. Recept. latum eleganter punctatum aut brevè alveolatum. Pappo interior rufus rigidulus deciduus coronâ albidâ cinctus. Cor. tubulosæ brevissimè 5-dentatæ. Anth. tenuissimè caudatæ. (v. s. comm. ab hon. Aul. angl. Ind. or.)

† *Species exclusæ.*

*P. ANNUA* Gærtn. = *Stenactis annua*.

*P. CALYCINA* Presl. = *Inula montana*.

## CCXLII. STRABONIA.

utulum multiflorum heterogamum, fl. radii subligulatis difformi-  
 nimis 1-serialibus fœmineis, disci tubulosi 5-dentatis herma-  
 itis. Recept. planum nudum. Invol. squamæ lineares pauciseriales  
 limidiò breviores. Antheræ basi bisetæ. Achænium compressum  
 in apice obtusum basi subangustatum. Pappus collo vix notando  
 in apice distinctus duplex, serie exter. brevissimâ dentatâ sub-  
 formi, int. setis pluribus densè et crassè barbellatis 1-serialibus.  
 R. Persica multicaulis erecta tota lanugine albâ molli densè  
 , pedicellis involucrisque subglabratissimis pubescentibusve. Folia  
 o-spathulata obtusissima undulata semiamplexicaulia. Capitula  
 cæca ramulorum solitaria. Flores flavi. — Genus achænio com-  
 mune teretiusculo ab Inuleis plerisque diversum. Dicavi memoriæ  
 Strabonis inter veteres geographi insignis.

*GNAPHALODES*. (1) ex Vent. sed 24 vid. ex radice crassâ tereti subligulata. In  
 1 inter Teheran et Ispahan ex herb. Oliv. et Brug. prope Tegrich ad ra-  
 montium Albourghs ex semin. ad Cels. missis. Inula gnaphalodes Vent.  
 r. jard. Cels n. et t. 75. (v. s. ex h. Oliv. et Brug.)

## CCXLIII. PEGOLETTIA Cass. dict. 38. p. 230. Less. syn. 200.

utulum multiflorum homogamum. Recept. ebracteolatum puncta-  
 tum. Invol. 2-3-seriale imbricatum floribus brevius, squamis  
 subulatis. Cor. tubulosæ 5-fidæ subringentes. Antheræ basi  
 nicaudatæ. Stylus Inulearum. Achænium cylindraceum sulcatum  
 ulatum eroste. Pappus conformis biserialis, paleis seriei ext.  
 as basi paleaceis, inter. elongatis piliformibus barbellatis ferè  
 10. — Fruticuli vel herbæ Africani. Folia alterna punctata ellip-  
 tica linearia. Capitula ad apices ramorum solitaria. Flores flavi.

*WEGALENSIS* (Cass. dict. 38. p. 230), herbacea erecta tota villis brevibus  
 la subglutinosa, foliis sessilibus lineari-sublanceolatis acutis vix apice  
 mutatis, pappo biseriali, inter. subplumoso, exter. brevi paleaceo barbato.  
 arenosis secus flumen Senegal in regione Walo, ad Daganâ? legit indef.  
 itet. Herba pedalis paniculato-ramosa. Radix subsimplex. Flores flavi  
 vultus subpurpurei. (v. s.)

*XYODONTA*, herbacea, basi glabra supernè puberulo-glandulosa viscosa,  
 ovali-lanceolatis utrinquè acuminatis dentibus utrinquè 2-3 grossis acutis  
 de callosis, invol. squamis acuminatis, floribus in invol. 35-40, pappo  
 ali exteriori brevissimo parco obtuso. — in Africâ austr. ad flum. Gariep,  
 sin-Namaqualand, legit cl. Drege. Folia a poll. longa, 7 lin. lata. Achæ-  
 nium sulcatum hinc indè pilosiusculum. (v. s.)

*UMINATA*, suffruticosa versus apices glandulosa, foliis oblongis integris  
 què acuminatis, invol. squamis longè acuminatis apice laxis, pappo bise-  
 riali breviori. 3 in Africâ australi ultrâ flum. Gariep legit cl. Burchell,  
 36. Ramuli et folia puberulo-glandulosa. Achænium glabratum striatum  
 acutum. (v. s.)

*HYGALÆFOLIA* (Less. syn. 200), fruticosa glabra, ramis tortuosis (aut ex  
 b. retrofractis), foliis petiolatis obovatis ovalibusve acutis integris, invol.  
 as acuminatis, pappo biseriali ext. brevi subpaleaceo. 3 ad Cap. Bonæ  
 Th. Licht.), ad Klein-Fischrivier et Zwarteberge (Drege!). Eupatorium  
 fractum Thunb. fl. cap. 628. Vernonia polygalæfolia Licht. mss. ex Less.

*ECCHARIDIFOLIA* (Less. syn. 200), fruticosa, foliis petiolatis ovalibus obo-

vatiave viscidis grossè dentatis, invol. squamis acutis, circ. 2-3-serialibus pappi pluriseriali, exter. vix distincto. ♀ ad Cap. Bonæ-Spei (Krebs! pl. exs. n. 191) ad Fischrivier et Zneenreberge (Drege!) *Pteronia dentata* Spreng. in h. Moric. Valdè priori ex habitu affinis, sed viscosa, folia dentata et pappi series externa vix ab inter. distincta. (v. s.)

6. P. ? *TANKEA*, herbacea erecta tennis sparsè hispida, foliis elliptico-oblongis utrinquè attenuatis integerrimis, ramis apice longè aphyllis pubescentibus r-cephalia, invol. squamis biserialibus acuminatis, pappo biseriali, exterior brevissimo. — in Africa anstr. legit cl. Drege, n. 5159, sine loci design. Flore non vidi undè genus subdubium. Recept. nudum punctatum. Achenium tera substriatum. Pappi setæ crassæ nives. (v. s.)

CXXIV. *GEIGERIA* *Griessolich* in *linnæa* 1830. p. 411. — *Geigera* Less. m. 199. — *Zeyhera* Spreng. f. diss. 1828 non Mart. 1826. — *Disonium* Willd. herb. ex Schlecht.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci tubulosis regulariter 5-dent. hermaphr. Recept. conicum ebracteolatum piloso-fimbrilliferum. Invol. imbricatum, squamis ext. bractæformibus foliaceis laxis, cæteris adpressis. Achenium subtrigonum erostre basi attenuatum pilis vestitum. Pappi paleæ 8-12 biseriales obovatæ apice obtusæ et ex Griess. subincisæ. — Herba humilima Capensis, radice seu rhizomate lignoso simplici, caule brevissimè subdichotomo. Folia linearia conferta integerrima. Capitula fasciculata sessilia. Flores lutei.

1. G. *AFRICANA* (Griess. l. c.). ♀ ad Cap. Bonæ-Spei. Zeyher! coll. exs. (185 n. 147 in h. Moric. Eckl.! pl. exs. n. 650. *Zeyhera acanthis* Spreng. f. l. c. *Dacnium longifolium* Willd. herb. ex Less. (v. s.)

#### DIV. II. CÆSULIÆ.

Capitula 1-flora in glomerulum aggregata, involucri proprio bivalvi dena cum achenio concreto. — Locus in serie naturali dubius.

CXXLV. *CÆSULIA* Roxb. *corom.* 1. p. 64. t. 93. *Cass. dict.* 6. suppl. p. 4. *Burm. trans. lin. soc.* 12. p. 94. Less. *syn.* p. 200. — *Meyeræ* sp. Don. Spreng.

Capitula plurima uniflora adgregata in glomerulum seu capitulum generale. Involucri proprium valvis 2 carinatis membranaceis inter se basi et demum cum ovario concretis tunc pappum *Bidentis* more bipartitum simulantibus. Cor. tubulosa, fauce ovata, lobis 5 parvis Styli rami filiformes divergentes glabri. Antheræ rigidulæ, basi setæ lævibus caudatæ. Achenium compressum epapposum quasi bihilum ex involucri valvis concretis. — Herba Indica glabra adscendens. Folia alterna lanceolato-lineariter utrinquè acuminata distanter serrata, basi ad florum originem dilatata semiamplexicaulia. Ramuli florales axillares brevissimi; folia floralia 2-3 colorata orbiculata capitulum generale adjectis bracteis paucis parvis cingentia. Flores pallide violacei aut rubri à centro ad peripheriam evoluti. — Ad *Heliantheas* et deindè ad *Vernoniaeas* dubitanter refert Cass.; Less. ad *Asteroides* *Inuleas*. *Stylis* enim non vid. *Vernoniaceæ* et antheræ *Inulearum*, sed habitus et flores. valdè diversi. Habitus ferè *Cryphiospermi* sed alternifolia.

1. C. *AXILLARIS* (Roxb.! l. c. et in fl. ind. 3. p. 447). ♀ in locis subhum. Coromandeliam, in oryzetis prope Tanjore (Wight!) et in paludosis Dep. inferioris regione subsylvaticâ Turraye. Wallich! cat. n. 2183. comp.

Andr. bot. rep. t. 431. *Meyera orientalis* Don prod. fl. nep. 180. *Melananthera orientalis* Spreng. in litt. ad Wall. 1830. Folia 3 pollices longa, 3 lin. lata. (v. s. comm. ab hon. Anlā angl. Ind. or.)

*Species exclusæ.*

C. *RADICANS* Willd. = *Cryphospermum repens*.

C. *SCANDENS* Poepp. = *Wulfia scandens*.

Subtrib. V. *BUPHTHALMEÆ*. — Inulæ *Bupthalmæ* §. 1. *Cass. dict.* 23. p. 566. — *Bupthalmæ* Less. *syn.* p. 209 (*adm. Ceruana*). — *Bupthalmi* *Linn. spec. Gerontogææ*.

Capitula multiflora heterogama (aut in Anvilleâ homogama), fl. marginalibus fœmineis sæpius ligulatis, in Ceruanâ tubulosi 3-fidis. Anth. nunc longè nunc breviter caudatæ nunc ecaudatæ (in *Lepidochætâ* et *Ceruana*). Recept. totum paleaceum, paleis floris discum subæquantibus. Achænium erostre aut brevissimè rostratum. Pappus coroniformis lacerus aut piloso-dentatus. — An genera infrâ indicata pro sect. tantum habenda? — Herbæ aut suffrutices. Folia in omnibus ritè cognitis alterna integra aut dentata. Capitula lutea. — Species omnes *Gerontogææ*.

CCXCVI. *BUPHTHALMUM* Neck. *elem.* 1. p. 20. *Cass. dict.* 34. p. 276. *Less. syn.* 209. — *Buatia* sp. *Adans.* — *Asteroidis* sp. *Tourn.*

Radius 1-serialis, ligulis latiusculis; corollarum omnium tubus teres infernè angustatus. Antheræ obsoletissimè caudatæ. Achænium radii triquetrum angustè trialatum, disci plano-compressum margine interiori unialatum. Pappus scariosus coroniformis piloso-dentatus conformis. — Herbæ perennes glabriusculæ. Folia lineari-lanceolata subintegra. Capitula terminalia solitaria. Invol. squamæ pauciseriales longè acuminatæ. — Genus habitu *Inulas* referens.

1. *B. SALICIFOLIUM* (DC. fl. fr. 4. p. 218), caule herbaceo, foliis lineari-sen oblongo-lanceolatis subdentatis, achæniis radii apice obtusis subtruncatis. ♀ in Europâ ferè totâ ad radices montium. Jacq. fl. austr. t. 370. *B. grandiflorum* Schkuhr handb. t. 257. *B. grandiflorum* et *salicifolium* Linn. sp. 1275. Lam. dict. 1. p. 516. Reichenb. fl. exc. 2. p. 236. Gand. fl. helv. 5. p. 380. Variat caule simplici aut ramoso, subvillosa aut glabra, ramis floridis erectis aut subpatulis, foliis acuminatis aut rariùs obtusis integerrimis aut dentatis, subvillosis aut glabris, latioribus et angustioribus, basi subæmiamplexicaulibus aut subangustatis, sed hæ variationes adeò in specim. variis combinantur ut species auctorum vix inter varietates veras recipiendæ videantur. (v. v.)

2. *B. INULOIDES* (Moris! sard. app. 1828), caule basi suffruticuloso, ramis foliis involucriæque pube brevi hirtis viscosis, foliis infer. ovatis grossè dentatis petiolatis summis lanceolato-linearibus integerrimis sessilibus, achæniis radii supernè in cornua duo acuminata productis. ♀ ♂ inter fissuras rupium calcar. insulæ Tavolara prope Sardiniam. Pappus irregulariter coroniformis inæqualis dentato-ciliatus. Paleæ minùs acuminatæ quàm prioris. (v. s. comm. à cl. Moris.)

3. *B. ANGUSTIFOLIUM* (Pursh fl. 2. p. 564), foliis alternis linearibus supernè latioribus integerrimis glabris, invol. squamis acutè lanceolatis. — in *Georgiâ* et *Floridâ* ex h. Banks in quo Pursh descr. Elliotio et aliis bot. amer. ign. An *Mnesithea* album Raf. fl. Ind. 68? ex Spreng.

† *Species non satis notæ.*

4. *B. LONGIPES* (Cass. bull. philom. 1822. p. 144. dict. 34. p. 277), caule herbaceo

viscidis grossè dentatis, invol.  
 triseriali, exter: vix distincto. 5 ad 6  
 Fischriev et Zuccareherge (Drege)  
 aldè priori ex habitu affinis, sed  
 vix ab inter. distincta. (v. s.)  
 herbae erecta tenuis  
 integerrimis, r  
 triserialib

viscidis grossè dentatis, invol.  
 riseriali, exter: vix distincto. 5 ad C  
 Fischriev et Zneareherge (Drege)  
 aldè priori ex habitu affinis, sed v  
 gna vix ab inter. distincta. (v. s.)  
 herbae erecta tenui  
 integerrimis, r  
 liserilib

P. TENELLA, herbacea erecta tenuis, utrinque attenuatis integerrimis, rari-  
t-cephalis, invol. squamis biseriatis  
brevissimis. — In Africa austr. legit et  
non vidi unde genus subdubium. Re-  
substriatum. Pappi setæ crassæ nives.

CCKLIV. GEIGERIA Griesslich in  
 199. — Zeyher Spreng. f. diss. 182  
 herb. ex Schlecht.

Capitulum multiflorum heterogamum  
fœmineis, disci tubulosi regulariter  
cum ebracteolatum piloso-fimbriatis  
ext. bracteaeformibus foliaceis laxis  
trigono-erostre basi attenuatum per  
seriales obovatas apice obtusæ et ex-  
limba Capensis, radice seu rhizomate  
subdichotomo. Folia linearia confertim  
sessilia. Flores lutei.

1. *G. AFRICANA* (Griess. l. c.). 2 ad C.  
n. 147 in h. Moric. Eckl. 1 pl. exs. n. 1  
nium longifolium Willd. herb. ex Les  
Div. II.

Div. II.

Capitula 1-flora in glomerulom aggregatis  
 phario concreto. — Locus in seri

CCXLV. CÆSULIA Roxb. corom. 1. p.  
trans. lin. soc. 12. p. 94. Less. syn. P.

CKLV. CÆSULI. *Roxb. Less. 3. p. 447.*  
trans. lin. soc. 12. p. 94. Less. 3. p. 447.  
Capitula plurima uniflora adgreg  
generale. Involucrum proprium valv  
se basi et demum cum ovario conc  
bipartitum simulantibus. Cor. tubulo  
Styli rami filiformes divergentes glabr  
lævibus caudate. Achanium compress  
ex involucri valvis conerctis. — Herba in  
alterna lanceolato-linearia utrinquæ acut  
ad florum originem dilatata semialexicau  
brevissimi; folia floralia 2-3 colorata orb  
adjectis bracteis paucis parvis cingentia. Flor  
à centro ad peripheriam evoluti. — Ad Helian  
nonieas dubitanter refert Cass.; Less. ad Aste  
enim non vid. Vernoniaceæ et antheræ Inulearu  
floresc. valde diversi. Habitus ferè Cryphiosperm.  
C. ARYZETIS (Roxb.) Less. 3. p. 447.  
C. ARYZETIS (Roxb.) Less. 3. p. 447.  
C. ARYZETIS (Roxb.) Less. 3. p. 447.

1. *C. AXILLARIS* (Roxb.) L. c. et in fl. ind. 3. p. 447). 2.  
Coromandeliae, in orysetis prope Tanjore (Wight!) et in  
inferioris regione subsylvatica Turraye. Wallich! cat. n. 2.

# DICOTYL. SEU EXOGENÆ.

EUPHATHALMÆ.

e, foliis alternis petiolatis cordiformibus obtusis trinerviis inæqualiter  
is, pedunculis elongatis oppositifoliis 1-cephalis, 1-bracteatis. — in im.  
gascar. Certè ex Cass. priorum congener sed habitus diversissimus et  
tactus indescriptus dubitationem suadent.

B. ? MELISSOIDES (Poir. suppl. 1. p. 747), caule herbaceo aspero, foliis oppo-  
sitis longè petiolatis cordato-ovatis obtusè crenatis asperis, pedunculo longis-  
simo 1-cephalo. — Patr. et car. pleriq. ignot. An Helianthea? fortè Wedelia?

PLERACEUM (Lour. fl. coch. 2. p. 618), caule herbaceo erecto lævi, foliis  
linearibus lanceolatis inæqualiter dentatis recurvis glabris, capitulis soli-  
tis, invol. squamis acutis membranulâ laterali connexis. — Cultum in hortis  
— hincinæ et Chinæ. — Discus flavus. Radius albus. Recept. paleæ minime.  
à cælo differt à B. graveolente ad quod Willd. refert et imò à genere om-  
nino alienum videtur.

RAMOSUM (Forsk. ag. ar. descr. 151), caule herbaceo hispido, foliis oppo-  
sit lanceolatis scabris enerviis denticulatis subciliatis, capitulis terminalibus  
— nculatis solitariis, invol. squamis ext. foliaceis radio parum brevioribus  
in Arabiâ felici. Vahl symb. 2. p. 92. Ligulæ circ. 8 apice tridentate per-  
tecto-nervosæ.

SAGITTATUM (Pursh fl. bor. am. 2. p. 564), tomentosum, foliis radicalibus  
gissimè petiolatis oblongis sagittatis integerrimis subtrinerviis, caulibus  
longis in petiolum attenuatis, caule subtricephalo, invol. squamis ext.  
o longioribus. ? in Amer. bor. montibus dictis Rocky-Mountains. Cap-  
itulum magna lutea (Pursh). Non est Bupthalmum (Nutt.), sed quid? †

LITENSE Ræusch. }  
REGÆ Spreng. } nomine tantum nota.

†† Species exclusæ.

- B. AQUATICUM Linn. = Asteriscus aquaticus.
- B. ARABICUM Delil. = Anvillea Garcini.
- B. ARBORESCENS Linn. = Borrichia arborescens.
- B. ASPERRIMUM Spr. = Anomostephium bupthalmoides.
- B. ASPERUM Horn.? = Wedelia hispida.
- B. ASTEROIDEUM Viv. = Pallenis spinosa.
- B. AUSTRALE Spreng. = Wollastonia aristata.
- B. BONARIENSE Pers. = Acicarpha spathulata?
- B. CAPENSE Linn. = Oederia prolifera.
- B. CANUM L'Hér. herb. = Borrichia argentea.
- B. CHRYSANTHEMOIDES Poir. = Asteriscus imbricatus.
- B. CORDIFOLIUM W. et Kit. = Telekia cordifolia.
- B. CRASSIFOLIUM Raf. = Asteriscus maritimus.
- B. DIFFUSUM Vahl mss. = Eclipta procumbens β.
- B. DURUM Linn. = Osteospermum spinosum.
- B. ELEGANS Brouss. = Asteriscus imbricatus.
- B. FLOSCULOSUM Vent. = Anvillea Garcini.
- B. FRUTESCENS Linn. = Borrichia frutescens.
- B. GARCINI Burm. = Anvillea Garcini.
- B. GRAVEOLENS Vahl = Asteriscus graveolens.
- B. HELIANTHOIDES Linn. = Heliopsis lævis.
- B. HETEROPHYLLUM Willd. = Wedelia pedunculosa?



- B. *HELIANTHOIDES* Forst. = *Wollastonia aristata*.  
 B. *IMBRICATUM* Cav. = *Asteriscus imbricatus*.  
 B. *INCANUM* Mill. = *Borrichia frutescens*.  
 B. *LEVIGATUM* Brouss. = *Jasonia lævigata*.  
 B. *LINKERI* Willd. = *Borrichia frutescens*  $\beta$ ?  
 B. *MARITIMUM* Linn. = *Asteriscus maritimus*.  
 B. *ODORUM* Schousb. = *Asteriscus odoratus*.  
 B. *PERRICUM* Garc. = *Anvillea Garcini*.  
 B. *PERUVIANUM* Lam. = *Borrichia Peruviana*.  
 B. *PRATENSE* Vahl = *Cerussa pratensis*.  
 B. *REPENS* Lam. = *Wedelia carnosa*.  
 B. *RANTILLA* Ham. = *Guizotia oleifera*.  
 B. *SCABRUM* Cav. = *Wedelia hispida*.  
 B. *SERICEUM* Linn. f. = *Asteriscus sericeus*.  
 B. *SENSILE* Mill. = *Asteriscus maritimus*.  
 B. *SPECIOSISSIMUM* Linn. = *Telekia speciosissima*.  
 B. *SPECIOSUM* Schreb. = *Telekia cordifolia*.  
 B. *SPINOSUM* Linn. = *Pallenia spinosa*.  
 B. *STENOPHYLLUM* Link. = *Asteriscus sericeus*  $\gamma$ .  
 B. *STRIGOSUM* Spreng. = *Wedelia pedunculosa*?  
 B. *TRIFOLIUM* Poepp. = *Oyedæa bupththalmoides*.  
 B. *UNIFLOSUM* Forst. = *Wollastonia Forsteriana*.

CCXLVII. *TELEKIA* Baumg. *fl. trans.* 3. (1816) p. 149, *malè*. Less. *syn.* 209.  
 — *Molpadia* Cass. *bull. philom.* 1818. *dict.* 32. p. 400. — *Inula* sp. *Bieb.*  
*Pers.* — *Asteroides* sp. *Tourn.*

Radius 1-serialis ligulis longis angustis; corollarum omnium tubus teres infernè sensim angustatus. Anth. basi caudis elongatis. Achænium disci et radii lineare elongatum multicostatum exalatum triquetro-obcompressum. Pappus coroniformis denticulatus conformis subcartilagineus (nec plumosus ut vult Baumg.). — Herbæ proceræ habitu *Inulæ Helenii*.

1. *T. CORDIFOLIA* (Kit. pl. rar. hung. 2. p. 117. t. 113. sub *Buphth.*), caule erecto pubescente, foliis subtus pubescentibus inferioribus petiolatis cordatis duplicato-callosis-serratis, superioribus ovatis sessilibus simpliciter serratis.  $\frac{1}{2}$  in pomariis sylvaticis et pratis montanis Banatûs, Croatiae, Transylvanicae, Tauricae, Cappadociae, Persiae ad Guriel (Szow!). *Bieb.* fl. taur. n. 1796. *Telekia speciosa* Baumg. fl. trans. 3. p. 150. *Molpadia suaveolens* Cass. *dict.* 32. p. 401. *Inula Caucasica* Pers. *ench.* 2. p. 450. *Bupththalmum speciosum* Schreb. *dec.* p. 11. t. 6. *Asteroides orientalis petasitidis folio*, etc. *Tourn.* cor. 51. *inst.* t. 487. Capitulum *Inulæ Helenii*. Invol. squamæ ovatae mucronatae. Pappus brevissimus. Herba trita suaveolens. (v. s.)
2. *T. SPECIOSISSIMA*, caule erecto sparsè piloso, foliis cordato-amplexicaulibus ovalibus acutis serratis rigidulis glabris nervo medio subtus subvillosis.  $\frac{1}{2}$  in vallibus montium Brixienensium (Ard.), ad rupes circa lacum Comensem (Com!), et in provinciis vicinis ad Corni di Canzo (Duby!), in valle Susanâ (Thom!), etc. *Bupththalmum speciosissimum* Ard. *specim.* 1. p. 26. t. 1. *Linn.* *mant.* 117. nec 517. *Reichenb.* *icon.* 4. p. 37. t. 349. *Gaud.* fl. helv. 5. p. 382. Invol. squamæ elongatae acuminatae. Pappus paulò magis elongatus quàm prioris. (v. s.)

CCXLVIII. *ASTERISCUS* Moench meth. 59a. Less. syn. 209. — *Nauplius* Cass. dict. 34. p. 272. — *Asterisci* sp. Tourn.. — *Asteroidis* sp. Faill. — *Odonatosperma* Nees? ex Less.

Radius 1-serialis, ligulis cuneiformibus apice tridentatis, tubo brevi basi biauriculato, cor. disci exalata tubo inferne incrassato. Anth. basi longe caudatæ. Achænium triquetro-obcompressum exalatum conforme. Pappus conformis coroniformis irregulariter dentatus. — Herba basi sæpe suffruticosa ramosa regiones mediterraneas habitantes plus minus hirsutæ. Folia obovata oblonga aut linearia integerrima. Capitula terminalia sæpius ramis lateraliter ortis et elongatis lateraliter facta. Invol. squamæ ext. foliaceæ inermes obtusæ disco multò longiores.

1. *A. MARITIMUS* (Moench l. c.), caule irregulariter ramoso, ramis apice 1-cephalis nunc foliosis nunc denudatis, foliis oblongo-spathulatis villosis, invol. bracteis spathulatis.  $\frac{1}{2}$  in rupibus maritimis *Hispania*, *Gallia australis*, *Italia*, *Aleppi*, *ins. Mediterran.*, *Mauritanie*. *Bupthalmum maritimum* Linn. sp. 1274. *Nauplius maritimus* Cass. dict. 34. p. 274. *Bupthalmum crassifolium* Raf. speech. 1. p. 191. ex descr. et sp. à cl. Guss. miso. — Barr. ic. t. 261. Booc. mus. t. 129. Clus. hist. 2. p. 13. Variat caule basi suffruticoso et herbaceo, supino et suberecto, ramis brevibus foliosis (*Bupth. sessile* Mill.) aut elongatis parvis foliosis post anthesin nudis. Invol. squamæ ext. spathulatis post anthesin deciduas, int. tunc adpressæ in discum depressum. (v. v. et s.)
2. *A. AQUATICUS* (Moench l. c.), caule erecto di- aut trichotomo, capitulis sessilibus axillaribus et terminalibus, foliis oblongis obtusis villosis aut glaberrimis. (1) in maritimis aquaticis regionibus *Mediterraneæ* à *Canariis* ad *Cretam*, à *Lusitaniâ* ad *Bysantium*. *Bupthalmum maritimum* Linn. sp. 1274. *Gusta* f. t. 169. DC. fl. fr. 4. p. 217. Schkuhr handb. t. 257. *Nauplius aquaticus* Cass. dict. 34. p. 273. — Seba thes. 1. t. 29. f. 7. Zan. hist. t. 24. *Breyn.* cent. t. 77. Interdum caulis induratur. (v. v. et s.)
3. *A. GRAVEOLENS*, caule erecto, ramis sub capitulis ortis elongatis prolixis, foliis oblongis basi cordato-amplexicaulibus incisè dentatis villosis, capitulis axillaribus brevissimè pedicellatis et terminalibus, invol. bracteis oblongo-linearibus acutis.  $\frac{1}{2}$ ? in *Ægypti* desertis circa *Kahiram*, necnon circa *Sinai* (*Bové!*). *Bupthalmum graveolens* Forsk. descr. 151. Vahl synb. 1. p. 74. t. 19. Sieb. herb. *Ægypt.* *Dalil.* fl. Capitulum post anthesin subglobosum. (v. s.)
4. *A. SERICEUS*, caule fruticoso ramoso, foliis cuneatis linearibusve integerrimis villosis-sericeis, capitulis terminalibus et lateralibus sessilibus, bracteis linearibus, rec. bracteolis acuminatis.  $\frac{1}{2}$  in *ins. Canariis*. *Bupthalmum sericeum* Linn. f. suppl. 379. Folia alterna nec opposita.
  - a. *latifolius*, foliis spathulato-oblongis densè villosis-sericeis, bracteis ralis brevioribus lineari-subspathulatis, invol. squamis linearibus sericeis-linearibus.  $\frac{1}{2}$  in insulâ *Fuertaventuræ*. *B. sericeum* L'Hér. herb. etms. Ait. kew. 3. p. 245. Sims. bot. mag. t. 1836. opt. Capit. cum radio disci sesquipollicari. (v. s.)
  - β. *intermedius*, foliis spathulato-oblongis adpressis villosis-sericeis bracteis oblongo-subspathulatis radio subæqualibus, invol. squamis linearibus pubescenti-sericeis.  $\frac{1}{2}$  in *ins. Lancerotte*. *Bupth.* *sericeum* Link in *Buch!* can. 150. (v. s.)
  - γ. *stenophyllus*, foliis linearibus acutis villosis-sericeis, caule ramosissimo depresso, bracteis folio conformibus, invol. squamis adpressissimis.  $\frac{1}{2}$  in campis maritimis *ins. Magnæ-Canariæ*. *Bupth.* *stenophyllum* Link in *Buch!* can. 150. Gravè olet ex Chr. Smith. Capit. cum radio 9-10 lin. diam. (v. s.)
5. *A. ODOREUS*, caule suffruticoso erecto trichotomo, foliis lineari-lanceolatis

acutis integerrimis pubescenti-subsericeis, capitulis terminalibus et axillaribus subsessilibus, bracteis plurimis folio conformibus, rec. bracteolis obtusè truncatis apice ciliato-villosis.  $\zeta$  in arenosis imperii Maroccani (Schousb.), prope Mogador (Brouss.)<sup>1</sup>, Bupthalmum odororum Schousb. mar. 2. p. 199. t. 7. Prioris var.  $\gamma$  affinis sed bracteolis obtusis faciliè differt. (v. s.)

6. *A. IMBRICATUS*, glaber atrovirens, caule fruticoso ramoso, foliis in petiolum attenuatis spatulatis obtusis argutè et grossè serratis, capitulis terminalibus solitariis, bracteis cuneiformibus apice 3-dentatis.  $\zeta$  in agro Mogadoriensi detexit cl. Broussonet. Bupthalmum imbricatum Cav. annal. scienc. nat. 1801. v. 4. p. 94. Bupth. elegans Willd. mag. naturl. ges. Berl. 1807. p. 138. Buph. chrysanthemoides Poir. suppl. v. 1. (1810) p. 747 excl. patr. Nomina 3 certè synonyma cum tres descr. à specim. Broussonetianis concinnatæ! (v. s. comm. à cl. Brouss.)

CCXLIX. *PALLENIS* Cass. dict. 37. p. 275. Less. syn. 210. — *Athalmum* Neck. elem. 1. p. 20? — *Asterisci* sp. Tourn. — *Asteroidis* sp. Vail.

Radius biserialis. Cor. radii angustè ligulatè apice tridentata tubo crasso utrinquè alato, disci tubo crasso hinc marginato-alato. Achænium obcompressum orbiculatum alatum; disci subtriquetrum villosiusculum. Pappus denticulato-ciliaris, radii dimidiatus externus, disci coroniformis. — Inv. squamæ ext. foliaceæ patentès apice spinosæ.

1. *P. SPINOSA* (Cass. l. c.), caule erecto hirsuto, foliis obovato-sut. oblongo-lanceolatis, invol. squamis ext. patulis spinescentibus. ① in asperis et subhumidis apricis, ad margines agrorum Europæ australis et Africæ borealis, à Teneriffa ad Ægyptum, ab Armorica Nannetensi ad Constantinopolim. Bupthalmum spinosum Linn. sp. 1274. DC. fl. fr. 4. p. 217. Schkuhr handb. t. 258. Bupth. asterioidum Viv. fl. lyb. sp. p. 57. t. 25. f. 2. — Blackw. herb. t. 272. Cles. hist. 2. p. 13. f. 1. Variat caule plus minus hirsuto simplici 1-céphalo aut corymboso, capitulis centralibus sessilibus cæteris longè pedunculatis, foliis integerrimis aut denticulatis. B. aureum Salz. pl. exs. circa Tanger lectum videtur var. foliis latioribus caule et involucri basi hirsutissimis. (v. v. sp.)

# CCL. ANVILLEA.

Capitulum multiflorum homogamum, fl. omnibus tubulosis 5-dentatis hermaphroditis. Recept. convexi paleæ basi oblongæ convexæ, apice ext. in acumen, interiores in setam productæ. Involucri campanulati squamæ ext. seu bracteæ foliaceæ spatulatæ patulæ, inter. biseriales adpressæ imbricatæ. Antheræ basi ecaudatæ (Vent.). Achænium tetragonum durum pappo coroniformi brevi integro coronatum. Herba, erecta aut multicaulis diffusa præsertim ad summities velutino-canescens, caule ad basin aut rhizomate fruticoso. Folia obovata spatulatave basi attenuata grossè et obtusè dentata. Capitula sessilia terminalia, ramis sæpè sub capitulis ortis quasi lateralialia. Flores aurei. — Genus à cæteris Bupthalmis distinctum floribus omnibus tubulosis hermaphr. et ideo ut ex antheris ecaudatis ad Cernuam vergens. — Dicevi cl. d'Anville (Joh. Bapt. Bourguignon d'Anville), summo geographo Parisiensi qui Orientalium regionum situs labore indefesso sagacissimè exploravit.

1. *A. GARCINI*.  $\zeta$  in Oriente non longè à Bagdad leger. cl. Olivier et Bruguère! in Arabia-Petræa (ex syn. Delil.) et in Persiâ olim legit Garcin (ex Barm.). Bupthalmum Persicum Garc. ex Barm. Bupthalmum Garcini Barm. fl. ind. t. 60. f. 1. Anthemis Garcini Barm. fl. ind. p. 183. Bupthalmum Arabicum

Delil. fragm. fl. arab. petr. p. 14. t. 4. ex Guillemin qui specim. vidit. Buphthalmum flosculosum Vent. h. Cels n. et t. 25. (v. s. ex herb. Oliv. et Brug.)

CCLI. CERUANA Forsk. ag. descr. 153. Cass. dict. 8. p. 22. Less. syn. 202. — Buphthalmi sp. Vahl.

Capitulum multiflorum heterogamum discoideum; flor. radii pluri-seriales fœminei graciles 3-dentati, disci hermaphr. 5-dentati. Recept. convexum totum bracteolatum. Invol. campanulato-hemisphaericum bracteis foliaceis extus cinctum erectum subbiserialia. Antheræ ecaudatæ. Achænium oblongo-obovatum bicostatum compressum, callo apicali rostrum brevissimum et annuliforme sistente. Pappus brevissimus coroniformis piloso-dentatus. — Herbæ rariùs fruticuli hirsuti Africani. Folia alterna subsemialexicaulia grossè dentata aut incisopinnatifida. Capitula cymosa, nempe pedunculi 1-3 infra capitulum terminale quodque nascentes et elongati in pseudo-laterales conversi. Flores lutei. — Genus habitu Buphthalmo affine sed discoideum, Grangeæ sed recept. bracteolatum.

1. C. PRATENSIS (Forak. fl. æg. p. LXXIV), caule herbaceo erecto hispido, foliis obovato-cuneatis dentatis aut pinnatifidis, bracteis capitulo longioribus lanceolato-linearibus. ① in Ægypto ad ripas Nili. Buphthalmum pratense Vahl symb. 1. p. 75. Delil! fl. eg. p. 132. t. 48. f. 2. Sieb. herb. eg. exs. (v. s.)
2. C. SENEGALENSIS, caulibus herbaceis erectis hispidissimis, foliis obovato-cuneatis grossè dentatis, bracteis ovatis obtusis capituli longitudine. ①? in Senegaliâ ad Podor legit indef. Perrottet mensæ 7<sup>br</sup> flor. Caulis à basi multiceps. Radix simplex crassitie pennæ columbinæ. Herba tota hispidissima. An forte ex patriâ C. rotundifolia Cass. dict. 41. p. 123? sed forma foliorum non rîc cum nostrâ congruit. (v. s. comm. à cl. Perrottet.)
3. C. FRUTICOSA (Less. syn. 202), caule procumbente lignoso ramosissimo, foliis oblongo-ellipticis subsemialexicaulibus integerrimis seu pinnatifido-dentatis, ramulis confertis 1-cephalis. 5 in Ægypto prope Alexandriam. Fides Graugeæ Maderaspatanæ.

Subtrib. VI. ECLIPTEÆ Less. syn. 210. — Helianthearum gen. Cass.

Capitula multiflora heterogama, fl. radii fœmineis (excl. Chrysospermo) ligulatis, disci hermaphroditis. Antheræ ecaudatæ. Recept. paleaceum. Pappus nullus seu aristatus nunquàm setosus. Folia opposita. — Subtribus Heliantheis proxima sed stigmati structurâ ad Asteroides potius referenda.

CCLII. BORRICHIA Adans. fam. 2. (1763) p. 130. — Diomedea Cass. bull. philom. 1815. p. 175 et 1817. p. 70. dict. 13. p. 283. Kunth nov. gen. 4. p. 212. Less. syn. 240. non Zoolog. — Odontospermum Neck. elem. 1. (1791) p. 20? — Buphthalmi sp. Linn.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis 1-serialibus, disci hermaphrod. tubulosi 5-fidis. Involucrum hemisphaericum, squamis imbricatis, ext. foliaceis acutis, int. obtusissimis margine membranaceis. Recept. planum paleis lanceolatis onustum. Cor. faux parçè dilatata. Styli rami longi per totam ferè longitudinem hispidi acuti. Antheræ nigricantes. Achænia cuneata compresso-angulata corticata apice pappo brevi coroniformi dentato coronata, disci subdifformia subcalva. — Frutices suffruticesve Americani maritimi. Folia opposita oblonga integerrima coriacea mediantibus utrinquè li-

gula triangulari adpressâ (stipulâ auct.) subconnata. Capitula terminalia pedicellata. Flores lutei. — *Borrichia* est nomen vetustius, caractere pro tempore rectè stabilitum, dicatum cl. Borrich. *Diomedea* est nomen multò recentius et jam veteribus et nunc zoologis avem designat.

1. *B. ARBORESCENS*, glabra, foliis lanceolatis mucronato-acutis basi angustatis integerrimis, invol. squamis ovatis ext. subacutis adpressis, paleis spathulatis obtusis.  $\zeta$   $\zeta$  in insulis Cayo, Flamingo et Bonito, ad littus australe insulæ Cubæ (H. B.), in insulis Bahamensibus (Linn.), et Bermudianis (Pluk.). *Buphthalmum arborescens* Linn. sp. 1273. *Diomedea indentata* Cass. dict. 13. p. 284. *Diomedea glabrata* H. B. et Kunth nov. gen. am. p. 213. — Dill. elth. 1. t. 38. f. 43. Plum. amer. ed. Burm. t. 16. f. 2. (v. s.)
2. *B. FRUTESCENS*, adpressè sericeo-canescens, foliis lanceolatis obovatisve obtusis basi attenuatis utrinquè grossè et acutè ad basin 1-2-dentatis, ramis sæpius edentulis, invol. squamis ext. apice recurvo-squarrosis, paleis acuminatis indurto-spinosis.  $\zeta$   $\zeta$  in Jamaicâ (Sloane), in maritimis Americæ borealis à Virginiâ ad Floridam (Pursh!), in Mexico (Berl. Pl. exs. n. 1963). *Buphthalmum frutescens* Linn. sp. 1273. *Buphthalmum incanum* Mill. *Diomedea bidentata* Cass. dict. 13. p. 283. — Dill. elth. 1. t. 38. f. 44. Catesb. car. 1. t. 93. Plum. et Burm. t. 107. f. 1. (v. s.)  
 $\beta$ . *angustifolia*, foliis lineari-lanceolatis medio aut basi obtusè et acutè dentatis.  $\zeta$  in Mexico ad Tampico (Berl. Pl. exs. n. 126). An *Buphthalmum lineare* Willd. sp. 3. p. 2260 ad hanc aut ad sequentem referendum? (v. s.)
3. *B. PERUVIANA*, foliis suprà glabriusculis subtùs sericeo-canescens foliis lineari-lanceolatis submucronatis medio denticulis subobtusis 1-2 sæpius notatis.  $\zeta$  in Peruvîâ (Domb. ex h. L'Hér.). *Buphthalmum Peruvianum* Lam. dict. 1. p. 515. (v. s. sine capit.)
4. *B. ARGENTEA*, sericeo-lanata argentea, foliis lanceolatis mucronatis integerrimis, involucri squamis extimis ovatis sericeis vix mucronatis adpressis, int. glabris ciliatis amplis, paleis acutis flosculis ferè brevioribus.  $\zeta$  in Peruvîâ (Domb. ex h. L'Hér.), in maritimis insulæ Cubæ prope Havanam (La Sagra) et Guanavacoa (H. et B.), in ins. Magni-Kaimanis prope Jamaicam (Don ex Lindl.). *Diomedea argentea* H. B. et Kunth nov. gen. am. 4. p. 213. Lindl. trans. hort. soc. Lond. 7. p. 50. report 1827. p. 7. *Buphthalmum canum* L'Hér. herb. et descr. mas. Cum priore in hortis interdum confusa sed involucri diversissima. (v. s.)
5. *B. DEMISSA*, caule demisso ramoso glabro, foliis strigilloso-pubescentibus confertis subspathulatis basi longè attenuatis obtusis integerrimis, pedunculis terminalibus, invol. squamis ovatis adpressis post anthesin conniventi-clausis, paleis acutis flosculo brevioribus confertis et alveolas ferè constituentibus.  $\zeta$  circa Carthagenam Amer. austr. legit cl. Gaudichaud. (v. s.)

**CCLIII ECLIPTA** Linn. mant. (1767) p. 157. Gært. fr. 2. p. 441. t. 169. Cass. dict. 14. p. 231. 46. p. 399. et 410. Less. syn. 212. — *Micrelium* Forsk. descr. p. 152. — *Eupatoriophalacron* Vaill. act. Par. 1720. excl. sp. Adans. fam. 2. p. 130.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus fœmineis angustissimè et breviter ligulatis, disci tubulosi 4-dentatis hermaphr. Recept. è plano convexum, bracteolis lineari-filiformibus apice ciliatis achænia æquantibus instructum. Inv. biseriale, squamis 10-12 ovato-lanceolatis acuminatis. Achænia radii triquetra, disci compressa ad latera muricato-tuberculosa, apice epapposa mutica aut denticulis 1-3 minimis apiculata, juniora ad apicem puberula. — Herbæ æquinociales plùs minùs scabræ aut hirsutæ ramosæ erectæ aut prostratæ. Folia opposita integra aut serrata penninervia. Pedunculi axillares soli-

tarii gemini aut terni 1-cephali. Flores albi. Succus ad colorem nigrum inficiendum adhibendus. — Genus ex Cass. affine hinc Buphthalmum illinc Melantheræ.

Sect. 1. EUECLIPTA. — Ecliptæ sp. auct.

Achenia omnia muricato-tuberculosa calva aut vix ac ne vix aristolata, rari triquetra, disci modicè compressa oblonga.

1. *E. erecta* (Linn. mant. 286? Willd. sp. 3. p. 2217), caule erecto adpressè strigoso, foliis oblongo-lanceolatis utrinquè acuminatis, pedicellis solitariis geminisve capitulo quintuplò longioribus. ① per orbis totius regiones calidas in graminosis humidis argillois inundatis ferè vulgata; habeo ex China, Cochinchina, Ind. or., Ægypto, Senegaliâ, agro Owarienti, Africâ austr.-orient. prope Port-Natal (Drege!) et ex insulis Cuba, Trinitatis, Mauritii, etc.; habet etiam in Moluccis, Javâ, Asiâ minore, Surinamo, Virginia, Florida, Nova-Granatâ, Brasiliâ, Sancto-Domingo, sinu Mullgrave Amer. bor.-occ., etc. Verbesina alba Linn. sp. 1272. Cotula alba Linn. syst. nat. 2. p. 564. Micrelidium asteroides Forst. descr. 152. Ecl. adpressa Mench. meth. — Dill. elth. f. 137. Rumph. amb. 6. p. 18. f. 1. Pluk. alm. t. 109. f. 1. Folia utrinquè adpressè strigosa, integra parè aut intensè serrata. Fortè species Linnaeus et sequens sed cum omnes auctores hanc sub nomine *E. erectæ* habeant, etiâ admisi. (v. v. et s.)

β. *diffusa*, caule à basi ramosissimo diffuso aut ascendente pedunculis vix capitulo duplò longioribus. ① in Senegaliæ regione Walo cum var. erecti in argillois inundatis legit cl. Perrottet. (v. s. comm. à cl. inv.)

2. *E. longifolia* (Schrad! in litt. 1832), caule erecto adpressè strigoso, foliis oblongo-lanceolatis acuminatis basi obtusis semisimplexicaulibus ad latera reflexis pedunculis 1-3 capitulo quintuplò longioribus. ① in insulis Caribæ, Guadalupe; fortè in Porto-Ricco, Domingo et Martinicâ si Bellis ramosa Jacq. amer. 216. t. 129 et ideò *E. punctata* Linn. mant. 2. p. 286 huc ritè referendæ? *E. erecta* Linn. mant. 286? (v. s.)

3. *E. Zippeliana* (Blum! bijdr. 914), caule erecto apice patentim hirsuto, foliis ovali-lanceolatis undulatis utrinquè attenuatis, pedunculis 1-4 capitulo vix duplò longioribus. ① in graminosis paludosis Bataviæ. Pedunculi hirsuti. Folia utrinquè adpressè strigosa serrata. (v. s. comm. à cl. Blume.)

4. *E. prostrata* (Linn. mant. 286), caule prostrato aut ascendente adpressè strigoso ramulis subhirsutis, foliis ovali- aut oblongo-lanceolatis basi attenuatis subserratis subundulatis scabris, pedunculis 1-2 capitulo duplò triplòve longioribus. ① in humidis Indiæ orientalis ut vid. frequens et in insulâ Madag. ad rivulos indigena facta ex rev. Lowe Verbesina prostrata Linn. sp. 2. p. 127. Cotula prostrata Linn. syst. 2. p. 564. Micrelidium Tolak Forst. descr. 153. Dill. elth. f. 138. Pluk. alm. t. 118. f. 5. (v. s.)

β. *undulata*, ramis apice strigosis patentibus hirsutis, foliis ovatis, pedunculis vix capitulo longioribus. ① in Indiâ orientali. Ecl. undulata Willd. sp. 3. p. 2219. Hanc et veram prostratam conjungo ex pluribus specimenibus à cl. Wallichio sub n. 319 et 320, et à cl. Royle comm. nec meo meo sensu separatu dignis. (v. s.)

5. *E. parviflora* (Wall! cat. et herb. n. 323), caule prostrato-repente adpressè etiam in ramulis strigoso, foliis oblongo-lanceolatis utrinquè acuminatis subintegerrimis subscabridis, pedunculis 1-2 capitulo vix duplò longioribus. ① in Indiæ orientalis humidis legerunt cl. Heyne et Wight. Hic etiam pertinere videtur *E. prostrata* D et *E. Wall!* cat. et herb. comp. n. 319 et Ecl. prostrata cl. Chamisso in Luzoniâ lecta. Folia ferè *E. erectæ*, habet *E. prostrata*, caulibus longè radicanibus ab utrâque distincta. (v. s.)

6. *E. thermalis* (Bunge! enum. pl. chin. bor. n. 224), caule erecto simplici subdivaricato-ramoso adpressè strigoso apice subhispido, foliis oblongis strig-

attenuatis subintegerrimis adpressè strigosis, pedunculis 1-2 capitulo vix longioribus, achæniis angulis crassioribus granulatis. ① ad thermas prope Tanschan Chinæ borealis legit cl. Bunge fl. maio. (v. s. comm. à cl. inv.)

7. *E. PROCUMBENS* (Michx. fl. bor. am. 2. p. 129), caule procumbente adpressè strigoso, foliis longo-lanceolatis utrinquè attenuatis rariter serratis utrinquè adpressè scabris, pedunculis 1-2 capitulo duplò circ. longioribus, inv. squamis oblongo-lanceolatis acuminatis. ① in humidis Carolinæ (Bosc!). Ell. sketch 2. p. 403. Grangea lanceolata Poir. suppl. 2. p. 826. Rami ab Ell. dicuntur radicales. Invol. squamæ radio longiores. (v. s. comm. à cl. Bosc.)

*ß. patula*, foliis latioribus ovali-lanceolatis subtriplinerviis. ① Patr. ign. Ecl. patula Schrad. l in litt. 1832. Bupthalmum diffusum Vahl herb. ex Puer. l herb. (v. s.)

8. *E. BRACHYPODA* (Michx. fl. bor. am. 2. p. 130?), caule divaricato-prostrato adpressè strigoso, ramis apice pedunculisque hirsutis, foliis longo-lanceolatis utrinquè acuminatis rariter serratis utrinquè adpressè scabris, pedunculis 1-2 capitulo paulò longioribus, inv. squamis ovatis cuspidato-acuminatis. ① in Louisiana prope Novam-Aureliam et in Carolina (Michx.) fortè in Mexico littorali ad Tampico (Berl. l n. 23?). Amellus Carolinianus Walt. fl. car. p. 213 ex Michx. Flores dixit 5-fidos cl. Michaux et in nostrà 4-fidi! sed ipse Elliot nunquam 5-fidos vidit. An species dum hic confusæ? an caract. varians? (v. s.)

## Sect. II. PLATECLIPTA.

*Achænia lævia aut partim muricata aristis 2-3 coronata, lateraliter in alam latam complanata, radii subtrigona, disci compressissima subrotunda. — Herbæ glabriusculæ, foliis basi triplinerviis.*

9. *E. LANCEOLATA*, erecta glabra, foliis oblongo-lanceolatis integerrimis, pedicellis terminalibus et è dichotomiis ortis folio longioribus apice adpressè pubescentibus, invol. squamis 10 lanceolatis inter se subæqualibus, achæniis medio dorso muriculatis. ②? in Brasiliæ prov. Rio-Grande. Caulis purpurascens pedalis sesquipedalisve. Folia 12-15 lin. longa, 4-5 lata internodiis breviora. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1007.)
10. *E. ELLIPTICA*, ascendens glabra, foliis elliptico-oblongis basi attenuatis integerrimis, pedicellis terminalibus folio paulò longioribus adpressè pubescentibus, invol. squamis 5 ext. majoribus oblongis, 5 inter. submembranaceis dimidiò minoribus, achæniis non aut vix tuberculosis. ②? in Brasiliæ prov. Rio-Grande. Folia vix pollicem longa 4 lin. lata internodiis sæpè longiora. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. n. 1024.)

## † Species vix notæ.

11. *E. ? PALUSTRIS* (Forst. prodr.), erectiuscula glabra foliis lanceolatis subintegerrimis, capitulis pedunculatis. — in insulis maris Pacifici. An Meyeræ seu Enhydræ sp. ? †.
12. *E. ? RUSILLA*, basi repens, caule simplici foliis petiolatis ovalibus acutis glabris subdenticulatis 3-nerviis, capitulis axillaribus pedicellatis solitariis. — in Porto-Ricco. Cor. flavidæ seu purpurascens. Verbesina pusilla Poir. dict. 8. p. 459. †.
13. *E. CILIATA* (Raf. lud. fl. p. 69), ex foliis alternis à genere videtur aliena nec satis nota.

## †† Species exclusæ.

- E. HUMILIS* Kunth. = Sabazia humilis.  
*E. INTEGRIFOLIA* Spreng. = Bellis integrifolia.  
*E. LATIFOLIA* Linn. f. = Blainvillea latifolia.  
*E. SCABRIUSCULA* Wall. = Wollastonia scabriuscula.  
*E. SENSILIS* Sw. = Enhydra sessilis.

CCLIV. BLAINVILLEA *Cass. journ. phys.* 1823. p. 216. *dict.* 29. p. 493. 47. p. 99. 59. p. 138. *Less. syn.* 212. — *Ucacea Cass. dict.* 27. p. 9 *excl. syn. ad.* — *Verbesinæ et Ecliptæ sp. auct.*

Capitulum pluriflorum heterogamum, fl. radii paucis 1-seriatis femineis in ligulam latam brevem 5-dentatam irregularem expansis, disci tubulosi 5-dentatis hermaphr. Invol. ovato-cylindraceum 1-2-seriale. squamis erectis exter. ovali-oblongis foliaceis, int. paleiformibus. Recept. planum angustum, paleis flores amplectentibus concavis submembranaceis apice dentatis. Stylus disci inclusus, ramis brevibus fert Bupthalmi. Achænia radii triquetra glabriuscula in angulis hispidula aristas 3 persistentes rigidas ciliatas apice gerentibus, disci laterali compressa biaristata rarius 3-4-arist. — *Herbæ Æquinoctiales Gerontogæ erectæ scabræ aut hirsutæ. Folia opposita petiolata triplinervia ovato-rhombea aut lanceolata dentata. Pedunculi ex axillis seu alis ramorum 1-3, monocephali nudi. Flores albi.*

1. B. RHOMBOIDEA (Cass.! *dict.* 29. p. 493), foliis rhomboideis acuminatis irregulariter dentatis utrinque tomentoso-hirsutis, ramis inf. oppositis summis dichotomis, pedicellis petiolo multò longioribus, achæniis aristis crassis triquetris basi ad apicem barbato-plumosis. (1) in Brasiliâ (Comm.! ) circa Rio-Janeiro (Lund.! Gand.! ) et Bahiâ (Blanch.! Salzmann.! ). *Verbesina dichotoma* Murr. *comm.* Gætt. 1779. p. 15. t. 4 opt. sed excl. patr. Stirps in hortis botanicis non infrequens, sequenti similis sed certè diversa. (v. a. sp. et v. c.)

β. *lanceolata*, foliis angustioribus oblongo-lanceolatis. — *Verbesina lanceolata* Poir.! *dict.* 8. p. 461. (v. s. c.)

2. B. LATIFOLIA (DC. in Wight *contrib.* p. 17), foliis rhomboideis acuminatis grossè dentatis utrinque sparsè pubescenti-hirtellis, ramis petiolisque hirtellis oppositis summis dichotomis, pedunculis petiolo plerumque brevioribus, achæniis aristis basi barbatis apice nudis sæpè subabortivis. (1) in Indiâ orient. ad montes Dendigul Malabaricæ legit cl. Wight et ad Munghir. *Eclipta latifolia* Linn. f. *suppl.* 378 et Linn.! *herb.* ex Wight! *herb.* *Verbesina dichotoma* Cef. Wall.! *cat. et herb.* n. 314. *Verbesina Lavenia* Roxb. fl. ind. 3. p. 443 non Linn. ex Wight. Ad priorem valdè accedit sed pubero-hispidula nec velatè hirsuta, folia latiora duplò longiùs petiolata papyraceo-membranacea, superiora ferè lanceolata. (v. s.)

β. *angustifolia*, foliis angustioribus oblongo aut vix ovato-lanceolatis. — in Indiâ orient., Ceylonâ, etc. *Verbesina dichotoma* Wall.! *cat. et herb.* n. 314 A. C. D. (v. s.)

3. B. GAYANA (Cass.! *dict.* 47. p. 90), foliis oppositis et alternis longè petiolatis ovato-lanceolatis dentatis suprâ scabris subtus pubescentibus, pedunculis dichotomiis ramorum ortis aut oppositifoliis petiolo duplò longioribus, invol. squamis int. striatis, vix exter. subvillosis. (1) in Senegaliâ ex semine. ad cl. Gay missis. Ligulæ dicuntur biligulatæ, in specim. meo deciduæ. (v. s. *comm.* à cl. Gay.)

4. B. PRURIKANA, foliis oppositis breviter petiolatis oblongo-lanceolatis acuminatis integerrimis aut subserratis utrinque strigoso-hispidis subtus pallidioribus, pedunculis axillaribus et alaribus petiolo triplò et ultra longioribus. invol. squamis linearibus dorso strigosis intus lævibus non striatis. (1) in Senegaliâ superiore ad montes Bakel legit cl. Leprieur. Flores non vidi. Achænia omnia in specim. meo oblonga subcuneata utrinque convexa biaristata apice biaristata juniora præsertim villosa submaculata. An fortè *Wedelia Africana* Pers. *ench.*? p. 490 huc refer.? (v. s. *comm.* à cl. inv.)

5. B. BIARISTATA, foliis oppositis longè petiolatis ovatis acutis serratis utrinque sparsè et adpressè pilosis subtus pallidioribus, pedunculis alaribus petiolo duplò longioribus, invol. squamis 5 oblongis acutis, achæniis omnibus biaristatis. (1) c.



Brasiliz prov. Rio-Grande. Ligulæ parvæ fœminæ. Capitula parva pauciflora. Petioli 8-9 lin. longi. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 873 miss.)

CCLV. SALMEA DC. cat. h. Monsp. (1813) p. 140. Brown. trans. linn. soc. 13. p. 112. Cass. dict. 47. p. 87. Less. syn. 212 non Cav. nec Willd. (1). — Hopkirkia Spreng. nov. prov. (1818) p. 23. — Bidentis sp. Linn.

Capitulum homogamum. Inv. imbricatum. Recept. conicum paleaceum, paleis convolutis flores amplectentibus. Cor. tubulosæ, fauce vix dilatata, lobis 5 subrevolutis. Antheræ sagittatæ (Br.). Styli basi subbulbosi rami breves crassi extus subhispidi supra papuloso-puberi obtusi. Achænia obcompressa linearia aptera, marginibus ciliatis apice in aristis 2 persistentes productis. — Frutices erecti aut subscondentes Americani et præes. Caribæi. Rami teretes. Folia opposita petiolata ovata acuta penninervia dentata aut subintegerrima. Capitula paniculato-corymbosa. Flores albidii. — Genus affine Dahliz ex stigmate, sed valdè affine Spilantho ex receptaculo achænio et paleis. Valdè dubius hæreo de loco huic generi in ordine naturali tribuendo, aut inter Eclipteas aut inter Verbesineas.

1. *S. EUPATORIA* (DC. cat. h. monsp. p. 140 cum ic. ined.), caulibus plurimis suberectis glabris apice pubescentibus, foliis ovatis acuminatis hinc indè dentatis glabris, aristis æqualibus, corollis rectis.  $\frac{3}{4}$  in ins. Caribæis. Bidens scandens Swartz obs. 927? Salmea grandiceps Cass. dict. 47. p. 88? Hopkirkia eupatoria Spreng. syst. 3. p. 443. Jacq. f. ecl. v. 2. t. 139. ined. Eupatorium Houstonis et Eup. Gouani hortul. non Linn. Rami divaricati. Pedunculi puberulo-hirsuti divaricati, inf. sæpius alterni paniculati, sup. oppositi corymbosi. (v. s. sp. et v. c.)

$\beta$ . *intermedia*, ramis ramisque glaberrimis, foliis basi cuneatis repandentatis glabris.  $\frac{3}{4}$  in Jamaica! An fortè species propria exactè inter *S. eupatorium* et scandentem media. (v. s.)

2. *S. SCANDENS* (DC. l. c.), caule scandenti ramisque etiàm ad apicem lævibus, foliis ovatis acuminatis subintegerrimis glabris, aristis æqualibus, corollis rectis.  $\frac{3}{4}$  in ins. Caribæis et sp. in Jamaica. Bidens scandens Linn. sp. 1167. hort. cliff. 399. Sinis bot. mag. t. 2062. Salmea scandens Brown l. c. Bidens frutescens Mill. dict. n. 7. Hopkirkia scandens Spreng. syst. 3. p. 443. Calea Amellus Linn. sp. 1179? ex Spreng. et suad. Willd. sp. 3. p. 1795. Salmia parviceps Cass. dict. 47. p. 88? Isocarpha Amellus Spreng. in h. Balb. (v. s. sp.)

3. *S. HIRSUTA* (DC. l. c.), caule scandente striato, ramis hirsutis, foliis ovatis basi subcordatis apice acuminatis integerrimis supra pubescentibus subtus villosis hirsutis, aristis inæqualibus, corollis rectis.  $\frac{3}{4}$  in Jamaica. Bidens hirsuta Swartz fl. ind. occ. 3. p. 1295. Salmea hirsuta Brown l. c. Hopkirkia hirsuta et Eupatorium vestitum Spreng. syst. 3. p. 444 et 416. Salmea oppositiceps Cass. dict. 47. p. 89? (v. s. comm. à cl. Bertero.)

4. *S. ? CURVIFLORA* (Brown trans. soc. lin. 13. p. 112), tubo corollæ insigniter extrorsum curvo, aristis inæqualibus, interiori alata, stigmate revoluta.  $\frac{5}{6}$  Patr. et cœt. car. ign.

N. B. Hopkirkia fruticulosa Spreng. syst. 3. p. 444, capitulo radiato et pluribus aliis caracteribus à genere diversa, est Armania fruticulosa Bert.

(1) Salmea Cav. est Sansevieria. Salmia Willd. est Carludovica.

lete bicornæ. — Herbaria mexicanæ grandes. Roma opposita pariter bipinnatipartita, segmentis ovatis acutis serratis. Racemulæ, aliis cylindricis, aliis oblongo-tuberculatis. Rami apertissimi 1-cephali. Capitula versicolora disco nempe luteo, radiis albo aut flavo. — Herbas hortenses autumnales speciatim in Europam introduxit cl. Sessè Moçino et Cervantès 1790. Innumerae varietates et variationes in hortis enatae sunt ad 1500 enumerat cl. Walner in cat. Dahl. Genevæ 8° 1833 in hortis quotidie creantur sed species primitivæ videntur c

1. *D. VARIABILIS* (Desf. cat. h. par. ed. 3. p. 182), caule non pruinoso, organa feminea gerentibus fertilibus sterilibusve.  $\mathcal{U}$  in Mexico. *D. rosea* Cav. ic. 1. p. 33. t. 80 et 3. t. 265. Thonin! ann. mus. 3. f. 1 et 3. *D. sambucifolia* Salisb. parad. t. 16. *D. purpurea* P. p. 444. *Georgina variabilis* Willd. hort. berol. 2. t. 93-95. Georg. DC. ann. mus. Par. 15. p. 310. *Dahlia superflua* Ait. Kew. ed. 1. *Georgina variabilis* & Kunth. nov. gen. am. 4. p. 243. *Coreopsis nuda* Cass. dict. 18. p. 442. Variat caule elato orgyali aut 2-pedunculato (hortul.), rachi plus minus alata, foliis puberulis aut glabriusculis, flosculis ligulatis 1-seriatis (*D. simpl. hort.*) pleriseriatis (*D. fl. & omnibus ligulatis* (*D. plenis*), sæpius expansis aut in globum (*D. en boule, globe flowered*), ligulis purpureis, rubris, roseis, & stramineis flavisque, nunc præsertim in ligulis 1-seriatis formis nunc præsertim in pluri- aut multiseriatis femineis abortivis. At differt *D. pusilla* Zucc. in herb. Acad. Monac. distincta statura & squamis angustioribus, inter. subacutis? (v. v. c. et s. sp.)
2. *D. CERVANTESII* (Lag. mant. ined.), caule non pruinoso solide destitutus sen verè neutris.  $\mathcal{U}$  in Mexico. *Georgina Cervantesii* Lag. gard. 2. t. 22. *Dahlia acutiflora* fl. mex. ic. ined. Media inter puerilem & adultam. Caulis orgyalis. Fol. rachis aptera. Invol. inter. squamis angustæ elongatæ reflexæ, inter. etiã angustiores. Ligulæ purpureæ aut coccineæ. Achæmium ferè spatulatum. (v. s.)
3. *D. COCCINEA* (Cav. ic. 3. p. 33. t. 266), caule pruinoso cavo, ligulis neutris.  $\mathcal{U}$  in Mexico. *Georgina variabilis* & Kunth nov. gen. 2.

## CCLVII. LEPTOCARPHA.

**Capitulum** multiflorum heterogamum, fl. radii ligulatis neutris 1-ser., disci hermaphr. 5-dentatis. Invol. squamæ biserialis æquales lanceolato-lineares. Recept. subconvexum, paleis paucis linearibus angustissimis deciduis. Cor. disci tubo cylindrico extus pubero, fauce vix dilatata. Styli rami breves lineari-lingulati dorso puberi. Achænia compressa (ext. obcompressa?) obovata membranacea pubescenti-subglandulosa, aristis 2 brevibus tenuibus ad angulos coronata. — Suffrutex Austr.-Americanus glabriusculus erectus. Folia petiolata ovato-lanceolata dentato-serrata basi obtusa apice acuminata membranacea penninervia inf. opposita, sup. subalterna. Capitula pedicellata circ. 5 subcorymbosa. Corollæ luteæ. Anth. fuscae. — Differt ab *Heliantho* paleis recept. stigmatibus et corollæ disci structurâ.

1. *L. RIVULARIS*. 5 in America australi (Chili aut Peruvia?) legit cl. Pœppig. *Helianthus rivularis* Pœpp. pl. exs. n. 716. Pedicelli petioli ramique juniores pubescentes. Fol. limbus glaber. (v. s.)

CCLVIII. *SIEGESBECKIA* Linn. hort. cliff. 412. *L'Hér. stirp.* 37. *Gärtn. fr.* 2. p. 424. t. 168. *Cass. dict.* 49. p. 114. *Less. syn.* 211. — *Siegesbeckia* sp. Linn. gen. n. 973. — *Sigesbeckia* et *Trimeranthes* *Cass. dict.* 59. p. 237.

**Capitulum** multiflorum heterogamum, fl. radii 1-serialibus fœmineis ligulatis aut difformibus, disci tubulosis hermaphr. 3-5-dentatis. Invol. biseriale, squamis ext. 5 lineari-spathulatis patentibus pilis glanduloso-capitatis ornatis, inter. achænia radii semi-involventibus erectis dorso piloso-glandulosis. Recept. planum paleis ovali-oblongis achænia involventibus. Achænia obovato-oblonga subtrigona intus subarcuata epapposa. — Herbæ dichotomæ in partibus superioribus glanduloso-pilosæ. Folia opposita sæpius in petiolum angustata ovato-triangularia grossè dentata subvillosa aut pilosa. Pedunculi ex alis ramorum orti, 1-cephali, subcorymbosi. Flores lutei.

Sect. 1. PENTAMERANTHES. — *Siegesbeckia Mœnch meth. Cass. dict.* 59. p. 237. 49. p. 114.

Flores fœminei ligulâ planâ donati. Fl. disci 5-dentati 5-andri. Achænia areolâ seu collo brevissimo integerrimo asperata aut subcalva.

1. *S. ORIENTALIS* (Linn. sp. 1269), foliis ovatis basi cuneatis apice acuminatis grossè dentatis, summis oblongo-lanceolatis, invol. exter. squamis interiore duplò longioribus. (1) in Chinâ, Indiâ orientali frequens. Cl. Wight legit in montibus Dendigul ad altit. 2000 ad 2500 ped. Etiâ recepi ex ins. Mauritii sed verisim. introductam, in ins. Societatis, imò ex cultis Chilensibus. Linn. hort. cliff. t. 23. — Icon Pluk. amalth. p. 58. t. 380. f. 2. non malè specim. mea Chinesisia refert nec meo sensu excludenda. (v. v. c. et s. sp.)
2. *S. JORULLENSIS* (H. B. et Kunth nov. gen. am. 4. p. 284), foliis ovato-triangularibus basi cuneatis apice acuminatis grossè dentatis, summis ovato-triangularibus, invol. exter. squamis interiore triplò quadruplòve longioribus. (1) in Mexico ad montem Jorullo (H. et B.) Lindl. bot. reg. t. 1061. *S. droseroides* Sweet br. fl. gard. t. 203. *S. triangularis* Cav. ic. 3. p. 27. t. 253? Priori valdè affinis sed videtur diversa. (v. s.?)
3. *S. CORDIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 283), foliis deltoideo-ovatis in petiolum decurrentibus acutis cordatis irregulariter dentatis summis pilis articlatis hinc indè ciliatis oblongo-lanceolatis, invol. ext. squamis interiore

triplo longioribus. (1) in planitie Bogotensi ad urbem Santa-Fé (H. et B.), foris in Mexico ad Zacuatlipam (Berl.). (v. s.)

4. *S. serrata*, foliis ovatis acutis regulariter dentato-serratis breviter petiolatis, summis sessilibus ovato-lanceolatis ciliatis, invol. ext. squamis interiore duplo longioribus. (1) in Chili legit cl. Chamisso et in Americâ austr. cl. Poepp. *S. cordifolia* Poepp.! diar. 463. herb. n. 55. Differre videtur à priore potius semipollic. nec 3-pollic. limbo ovato nec cordato, inv. ext. squamis brevioribus. (v. s.)
5. *S. IBERICA* (Willd. sp. 3. p. 2220), foliis deltoideo-triangularibus grossè et irregulariter dentatis in petiolo cuneato-decurrentibus, summis ovato-lanceolatis, invol. ext. squamis interiores paulò superantibus. (1) in Caucasi Iberici sylvis (Stev.), Tauris merid. vineis (Bieb. Mey. enum. n. 595) et circa pagos Mædæ (Buxb.). *S. Orientalis* Bieb. fl. taur. n. 1791. — Buxb. cent. 3. p. 29. t. 52. (v. s.)
6. *S. MICROCEPHALA*, foliis breviter petiolatis oblongo-lanceolatis hinc inde serratis utrinque sparse puberulis, involucri ext. squamis linearibus subspathulatis glanduloso-ciliatis vix capitulo longioribus. — in Novâ-Hollandiâ ex herb. Lambert. Caulis dichotomus gracilis sparse puberulus. Petioli 2 lin. Folia fere sesquipollic., 5-6 lin. lata. Capit. 3-4 lin. diam. Ligulæ paucæ obovatæ 3-dentatæ. (v. s. comm. à cl. Lambert.)
7. *S. GRACILIS*, foliis breviter petiolatis oblongo-lanceolatis subintegerim utrinque sparse puberulis, invol. ext. squamis linearibus apice subspathulatis glanduloso-ciliatis capitulo duplo longioribus. — in Novâ-Hollandiâ ex herb. Lambert. Similis *S. microcephalæ* et fortè mera varietas. (v. s. comm. à cl. Lambert.)

Seet. II. *TRIMERANTHES* Cass. dict. 49. p. 114. 59. p. 237. — Schkuhria Mæch meth. 566, non Roth.

Fl. radii difformes breviter et irregulariter ligulati 2-3-dentati, disci 3-dentati 3-andri. Achænia collo omninò destituta.

8. *S. PUSCULOSA* (L'Hér.! stirp. nov. p. 37. t. 19), foliis sessilibus ovatis dentatis invol. ext. squamis interiore duplo longiore. (1) in Peruvîâ. Schkuhria dichotoma Mæch meth. 566. *Trimeranthes dichotoma* Cass. dict. 49. p. 115. (v. v. cl.)

CCLIX. *SABAZIA* Cass. dict. 46. p. 480. v. 55. p. 264 et 273. Less. syn. 211. — Ecliptæ sp. Kunth.

Capitulum multiflorum heterogamum, fl. radii fœmineis 1-seriatis ligulatis, ligulis cuneatis grossè tridentatis 6-nerviis invol. longioribus. disci hermaphr. tubulosi 5-dentatis. Recept. conicum, paleis linearibus seu setiformibus glabris achænia ferè superantibus, intimis sæpi post anthesin ad apicem recept. persistentibus. Invol. semiglobosum biseriale 7-8-phyllum. Cor. extus pilosiuscula. Achænia conformia pentagona striata cuneata apice omninò calva nec apiculata. — Herba Mexicanæ ramosæ. Folia opposita petiolata oblonga dentata glabra. Pedunculi alares aut plerique terminales elongati 1-cephali. Ligule subidæ extus purpurascens. Genus ab Ecliptâ differt ligulis late cuneatis, tubulis 5-dentatis, fructibus striatis nec tuberculatis 5-gonis omnibus nec 2-3-gonis dissimilibus apice omninò calvis et muticis. Habitus Galinsogæ sed achænium calvum.

§. 1. *Recept. conico, achæniis glabris.*

1. *S. HUMILIS* (Cass. dict. 46. p. 481), erectiuscula ramis pedunculisque pubescentibus, ligulis profundè trifidis. (1) in Novâ-Hispaniâ prope Arrio et Pucucaro. Eclipta humilis H. B. et Kunth nov. gen. am. 4. p. 264. t. 39.
2. *S. SARMENTOSA* (Less. in linnæa 1830 p. 148), caule sarmentoso, foliis ovatis

acuminatis teneris triplinerviis pilosiusculis anticè remotè serratis, invol. squamis obtusis, inter. ciliatis majoribus. — in Mexico ad Serro-Colorado.

3. *S. MICROGLOSA*, caule erecto pedunculisque glabris, foliis ovato-lanceolatis acuminatis teneris ferè à basi ad apicem remotè serratis, invol. squamis obtusis, inter. vix ciliatis, ligulis minimis. (1) in montanis circa Mexico ad S.-Augustinum legit Berlandier (pl. exa. n. 733). (v. s.)

ß. *puberula*, caule ad apicem pedicellisque pubescentibus. (1) circa Mexico in montanis. (Berl. ! pl. exa. n. 910). (v. s.)

### §. 2. *Recept. convexo, achæniis hispidulis.*

4. *S. ? URTICIFOLIA*, hispida, foliis ovatis acuminatis grossè dentatis, invol. squamis exter. piloso-glandulosis, recept. convexo, achæniis hispidulis. ½ in agro Quitensi inter Mulalo et Pensache. *Viborgia urticæfolia* H. B. et Kunth nov. gen. am. 4. p. 267. t. 389. An *Calydermis* spec. Less. syn. 245? *Habitus* *Galinsogæ* sed achænium calvum. Differt à *Calyderme* capitulis radiatis.

### † *Species exclusæ.*

*S. OCCIDENTALIS* Lin. = *Verbesina phætusa*.

*S. OCCIDENTALIS* Walt. = *Verbesina Siegesbeckia*.

*S. PORTORICCENSIS* Bert. = *Ogiera Portoriccensis*.

CCLX. *CRYPTIOSPERMUM* Beauv. fl. owar. 2. p. 24. t. 74. (excl. descr. ferè tota et icon. stigm.). Cass. dict. 12. p. 78. Less. syn. 210. — *Wahlenbergia* Schum. pl. guin. 387. non Schrad. — *Enhydra* sp. Cass. — *Meyera* sp. Brown. — *Casulie* sp. Willd.

Capitulum multiflorum heterogamum, fl. omnibus intra rec. paleas complicatas omninò involutis, ext. pluriserialibus femineis, centr. hermaphroditis stigmatis alteratione subabortivis. Invol. 4-phyllum, squamis foliaceis, 2 oppositis majoribus. Recept. hemisphæricum paleaceum, paleis coriaceis subglandulosis apice pilos articulos paucos gerentibus. Cor. basi latiores corneæ apice attenuatæ, limbo brevissimo, radii subligulato 3-4 dentato, disci 5-dentato campanulato. Stigmata breviter extus arcuata obtusa. Achænium obcompressum basi attenuatum erostre omninò calvum paleâ inclusum. — Herba Africana aquatica radicans glabra oppositifolia Enhydram omninò referens, dichotoma procumbens. Folia integerrima oblonga remota subserrata. Capitula axillaria solitaria sessilia.

1. *C. REPENS* (Beauv. ! l. c.). — in Africa æquinoctiali ad ripas flumin. Formosæ (Beauv. !), in Guineâ (Thoun.). Less. in Linzæ 1831. p. 154. *Casulia radicans* Willd. sp. 3. p. 1797. *Wahlenbergia glomerata* Schum. guin. 387. *Enhydra casuloides* Cass. dict. 14. p. 554. *Meyera Guineensis* Spreng. syst. 3. p. 602. (v. s.)

### Trib. IV. *SENECIONIDEÆ* Less. syn. p. 218.

Stylus apice cylindraceus, in flor. hermaphroditis supernè bifidus, ramis longiusculis linearibus apice penicillatis, nunc truncatis, nunc ultra penicillum in conum brevem aut appendiculum elongatum angustum et hispidulum productis. Stigmatis series latiusculæ et prominulæ usque ad penicillum seu originem coni seu appendicis extensæ. Corolla disci pellucida regularis. Pollen globosum echinulatum.

## CONSPECTUS TRIBUS.

Subtrib. I. MELAMPODINÆ. Flores omnes unisexuales nec ulli hermaphroditi, masculi et feminei in diversis plantis (dioici). aut in eorundem plantarum capitulis diversis (heterocephali). aut in iisdem capitulis (monoici). Antheræ ecaudatæ. Recept. plerumquæ paleaceum. Pappus nunquam setosus.

Div. I. EUXENIÆ. Capitula dioica multiflora discoidea. Achænia ecorticata.

Div. II. MILLERIÆ. Capitula monoica pauciflora, fl. radii pauciligulatis aut trifidis. Achænia ecorticata. Recept. sæpius paleaceum.

Div. III. SILPHIÆ. Capitula monoica multiflora, fl. radii pluribiligulatis. Achænia ecorticata. Recept. totum aut saltém margine paleaceum.

Div. IV. MELAMPODIÆ. Capitula monoica multiflora radiata Achænia corticata.

Div. V. AMBROSIÆ. Capitula heterocephala, nempe alia ♂, alia ♀. Antheræ marium liberæ. Cor. tubulosæ, interdum in ♂ nullæ.

Div. VI. IVEÆ. Capitula monoica. Cor. tubulosæ aut rariùs in ♀ nullæ. Antheræ liberæ.

Div. VII. PARTHENIÆ. Capitula monoica radiata. Achænia rariùs obcompressa aut rariùs trigona. Ligulæ tardè deciduæ.

Subtrib. II. HELIANTHÆ. Capitula sæpius heterogama radiata aut homogama discoidea. Recept. totum aut margine paleaceum. Cor. fl. herm. lobi crassi. Pappus nullus coroniformis aut aristatus. Antheræ nigricantes ecaudatæ. — Folia sæpius opposita.

Div. I. HELIOPSIDIÆ. Capitula heterogama ligulis femininis et discoidea. Achænia grossa cortice crasso sæpius adhaerens instructa, nunquam obcompressa. Pappus nullus, coroniformis, aut breviter aristulatus.

Div. II. RUDBECKIÆ. Capitula heterogama ligulis neutris et sterilibus. Achænia erostria. Pappus nullus aut coroniformis.

Div. III. CORROPSIDIÆ. Capitula heterogama, ligulis neutris et simè discoidea. Achænia erostria. Pappus aristis 2-4 et interdum etiã squamellis constans.

Div. IV. BIDENTIDÆ. Capitula heterogama ligulis neutris, rariùs discoidea. Achænia rostrata, rariùs calva, sæpius aristis 1-4 superata.

Div. V. VERBENIÆ. Capitula heterogama ligulis femininis, rariùs discoidea. Achænia cortice tenui donata, compressa aut obcompressa, calva, aut aristas tot quot anguli, aut setas rariùs fragiles gerentia.

**Subtrib. III. FLAVERIÆÆ.** Capitula uni- aut pauciflora densè aggregata heterogama, fl. fœmineis nunc solitariis nunc pluriserialibus, aut heterogama cum prioribus mixta. Achænia calva. Recept. in capit. paucifloris epaleaceum, in multifloris paleaceum. — Folia opposita.

**Subtrib. IV. TAGETINÆÆ.** Capitula heterogama fl. radii fœmineis aut discoidea homogama. Invol. sæpiùs 1-seriale aut gamophyllum. Recept. epaleaceum. Achæmium erostre basi attenuatum. Pappus aristatus, rariùs setosus. — Folia opposita rariùs alterna et involucra grossè pellucido-glandulosa.

**Div. 1. TAGETÆÆ.** Capitula sæpiùs radiata. Cor. disci regulares. Styli rami cono superati. Pappus aristatus.

**Div. II. POROPHYLLÆÆ.** Capitula discoidea. Cor. disci inæqualiter fissæ. Styli rami in appendicem subulatam producti. Pappus setosus pluriserialis.

**Subtrib. V. HELENIÆÆ.** Capitula sæpiùs heterogama. Achæmium erostre. Pappus squamellis pluribus 1-serialibus scariosis ferè semper liberis constans, rarissimè calvus. — Folia alterna rariùs opposita.

**Div. 1. GAILLARDIÆÆ.** Receptaculum epaleaceum.

**Subdiv. 1. EUGAILLARDIÆÆ.** Ligulis neutris.

**Subdiv. 2. EUHELENIÆÆ.** Ligulis fœmineis aut nullis.

**Div. II. GALINSOGÆÆ.** Receptaculum paleaceum. Involucri squamæ achænia radii non involventes.

**Subdiv. 1. EUGALINSOGÆÆ.** Ligulis fœmineis seu nullis.

**Subdiv. 2. SPHENOGYNEÆÆ.** Ligulis neutris.

**Div. III? MADIÆÆ.** Recept. margine aut ferè totum paleaceum. Invol. squamæ 1-2-seriales convolutæ aut complicatæ achænia radii semper calva includentes.

**Subtrib. VI. ANTHEMIDÆÆ.** Capitula sæpissimè heterogama, ligulis fœmineis seu neutris. Antheræ ecaudatæ. Styli rami truncati barbati rarissimè cono superati. Achæmium angulatum teres aut in radio obcompressum. Pappus nullus aut coroniformis, rariùs squamellis aut pilis capillaribus constans. — Folia sæpissimè alterna.

**Div. 1. EUANTHEMIDÆÆ.** Receptaculum paleaceum. Capitula radiata.

**Div. II. CHRYSANTHEMÆÆ.** Recept. epaleaceum. Capitula radiata.

**Div. III. COTULÆÆ.** Recept. epaleaceum. Capitula discoidea homogama aut heterogama.

**Div. IV. ATHANASIÆÆ.** Receptaculum paleaceum. Capitula homogama, cor. omnibus teretibus.

**Div. V. ARTEMISIÆÆ.** Recept. epaleaceum. Capitula discoidea homo- aut heterogama. Cor. disci teretes stylum bifidum gerentes.

Div. VI. HIPPIÆ. Capitula monoica discoidea. Recept. epaleaceum. Cor. teretes. Stylus disci simplex clavatus.

Div. VII. BRIOCEPHALÆ. Capitula monoica, fl. formineis pauciligulatis tubulosisve. Stylus fl. disci simplex clavato-truncatus. Receptaculum paleaceum.

Subtrib. VII. GNAPHALIÆ. Capitula homogama aut heterogama. discoidea rarissimè radiata, interdum 1-flora. Antheræ caudatæ. Achænia pappo piloso seu setaceo coronata, rarissimè calva. — Folia sæpius alterna.

Div. I. ANGIANTHÆ. Capitula pauciflora in glomerulum invol. generali cinctum densè aggregata.

Div. II. CASSINIÆ. Capitula non glomerata. Recept. totum aut margine paleaceum.

Div. III. HELICHRYSÆ. Capitula non glomerata. Recept. epaleaceum.

Div. IV. SERIPHIÆ. Capitula 1-flora, nunc libera, nunc glomerata et invol. generali cincta.

Div. V. ANTENNARIÆ. Capitula dioica aut subdioica non glomerata. Styli fl. masc. simplices. Recept. epaleaceum aut margine tantum paleaceum.

Div. VI. LEYSSEÆ. Capitula multiflora radiata, ligulis formineis. Recept. epaleaceum. Pappus squamellis setisve constans.

Div. VII. RELHANIÆ. Capitula multi- rariùs pauciflora homogama aut radiata. Recept. epaleaceum. Pappus nullus aut coraciformis.

Subtrib. VIII. SENECTIONIÆ. Capitula homo- aut heterogama, discoidea aut radiata. Antheræ ecaudatæ. Achænia pappo piloso seu setaceo coronata, exteriora rarissimè calva. — Folia alterna.

Div. I. NEUROLÆNÆ. Receptaculum paleaceum.

Div. II. ERECHTITÆ. Recept. epaleaceum. Capitulum discoidum heterogamum, flor. radii tubulosis formineis.

Div. III. EUSENECTIONIÆ. Recept. epaleaceum. Capitulum aut homogamum discoideum, aut radiatum. Cotyledones planæ.

Div. IV. BALBISIÆ. Recept. epaleaceum. Capitulum discoidum homogamum aut radiatum subdioicum. Cotyledones involutæ.

Subtrib. I. MELAMPODINÆ. — Melampodiæ et Ambrosiæ *Linn.* 213 et 218. — Ambrosiæ, Milleriæ et Balumoreæ *Cass. opusc. 3. p. 61.* — Polygamia necessaria *Linn.*

Flores omnes unisexuales, nec ulli hermaphroditi, in iisdem aut diversis plantis, et in eadem stirpe in iisdem aut diversis capitulis siti, semper monoici aut dioici. Antheræ ecaudatæ. Recept. ferè semper paleaceum. Pappus nullus, subcoroniformis aut aristatus, nunquam setosus. Styli nulli pilis collectoribus sæpius deficientibus sæpius subdeformati, aut *Senecionidearum* apice barbati.



1. EUXENIÆ. — Asteroideæ Melampodieæ Euxeniæ Less.  
*syn.* 216.

*titula* dioica, multiflora, discoidea. Achænia ecorticata.

1. EUXENIA *Cham. hort. ber.* (1820). p. 75. t. 16. (*excl. syn.*) *Cass. dict.* 35. 16. 50. p. 472. *Less. syn.* 216 non *Nees et Mart.* — *Ogiera Spreng. syst.* 3. 74 non *Cass.* — *Podanthus Lag. nov. gen.* (1816). p. 24. *Cass. dict.* 46. 14. *Spreng. syst.* 3. p. 456. *Less. syn.* 419 non *Haw.* (1812).

*titula* dioica globosa homogama discoidea. Invol. 1-seriale patulum reflexum, squamis oblongo-lanceolatis 10 liberis æqualibus oppositis sæpe majoribus. Recept. convexum paleis oblongis sub deciduis onustum. Cor. tubo angusto, fauce campanulatâ extus â et glandulosâ, limbo 5-dentato. ♂ anth. ecaudatæ parùm cohæ- (nec liberæ) nigricantes. Stylus inclusus, ramis brevibus crassis s dorso hispidulis. Ovar. abortivum. ♀ anth. abortivæ liberæ. ami crassi obtusi exserti. Achænium obpyramidatum basi attenuatum tetragonum calvum aut rarius pappo subauriculiformi coronabulum apice pilis minutissimis circa areolam subinstructum. — es Chilenses dichotomi ramosissimi, ramis teretibus junioribus sæpe pubescentibus. Folia opposita breviter petiolata ovata aut lanceolata acuta serrata triplinervia utrinquè subscabrida. Pedunculus 3 è dichotomiis orti 1-cephali. Flores flavi.

1820 (Cham. hort. ber. p. 75. t. 16), foliis latè ovatis in petiolum non rentibus. 5 in rupestribus apricis Chilensibus (Domb.!), nempe ad aguan (Cham.), Conception (Beech.), in Brasiliâ? (Nees et Mart. beitr. 7. act. nat. cur. 12. p. 7). *Cass.* 50. p. 472. *Hook. et Arn. bot. Beech.* 1. 7. *Ogiera triplinervia Spreng. syst.* 3. p. 674 non *Cass.* *Baccharis Chilensis Weinm. syll. pl. nov.* 2. p. 21 ex *Less.* *Podanthus ovatifolius Lag. nov.* 24. n. 322. Vulgò Chilensibus dicitur *Palo-Negro* ex *Lag.* (v. a.)

1821 (Lag. nov. gen. p. 24. n. 322), foliis ovali-lanceolatis basi longè cuneatis apice acuminatis medio serratis. 5 in Chili circa Valparaiso (Macr.! Gand.!), in sylvaticis colad Quillots et Rancagua (Bert.!), *Podanthus Mitiqui Lindl.!* in litt. Loud. hort. brit. p. 488. *Ogiera triplinervia Bert. herb. non Cass.* Vulgò meibus dicitur *Mitirin* ex Bert. et verosim. *Mitiqui* ex nom. spec. Lindl. Priori affinis sed foliorum formâ sat distincta. (v. a.)

1822 (Lag. nov. gen. p. 24. n. 322), foliis subintegerrima, 5 in Chili. *Græmia aromatica Pæpp.!* pl. exa. n. 208. diar. n. 30 non *Hook.* Folia minora ferè omnia integerrima, hinc inde subdenticulata. An fortè species propria? (v. a.)

† *Species exclusa.*

1823 *Nees.* = *Gymnopais euzenoides.*

1824 L. PETROBIUM *R. Brown trans. Linn. soc.* 12. p. 113. — *Laxmannia nov. gen. n. et t.* 47 non *Schreb.* nec *R. Br.* nec *Fisch.* (r). — *Spilanthi Forst.* — *Drimphyllum Burch. ms.*

*titula* pluri- (10-12) flora dioica. Recept. paleis membranaceis squamas referentibus inter flores sitis. Invol. subcampanulatum de-oligophyllum floribus brevius, squamis ovali-oblongis glabris. ibus teres vix aristis achæniis longior, limbo 4-partito, lobis ovali-

*Laxmannia* *Schreb.* (1791) est *Cymnosma*. *Laxmannia* *Fisch.* (1812) est *Stictisect.* *Gei.* *Laxmannia* *Brown* (1810) est genus inter *Asphodeless.* Stricto jure orsteri servandum et novum genus ad *Brownii* nomen fingendum esset, sed ioritatem hic neglexi ne augeatur confusio.

oblongis. ♂ antheræ exsertæ appendiculatæ. Stigmata acuta hispidula. ♀ stam. distincta. Stigmata acuta recurva. Achænium lineare plano-compressum, ad latus utrinquè medio nervigerum, sparsè hirtellum, apice productum in aristas 2 rigidulas ex angulis ortas margine strigillis erectis subciliatis. Interdum arista tertia longior adest. — Arbor trioryzi Heleniana. Ramuli alterni. Folia opposita petiolata latè obovata ses ovalia, anticè serrata, suprà glabra, subtus sparsè subpuberula. Panicula (ex Burch.) terminalis oligocephala ramulis oppositis. Capitula longè pedicellata erecta albida.

1. *P. ARBORÆUM* (Brown l. c.), 5 in insulâ Ste-Helenæ ubi Angli dicitur *White-Wood*, *Cabbage-Tree*. *Bidens* arborea (fem.) et *Spilanthes* tetradactyla (mascul.) Roxb. in Beata. app. p. 301 et 325. *Spilanthes* arborum G. Forst. comm. goett. p. p. 66. *Laxmannia* arborea Forst. nov. gen. t. 47. *Spilanthes* pseudogummifera Forst. in h. Mus. Par. Drimiphyllum Helenianum Burch. in litt. (v. s. specim. form. comm. à cl. Burch.)

CCLXIII. *ASTEMMA* Less. syn. 216. — *Monactidis* spec. Kunth.

Capitula 10-15-flora homogama discoidea dioica. Invol. tubulosum campanulatum, squamis linearibus obtusis subimbricatis. Recept. paleis squamas referentibus membranaceis ciliatis. Cor. tubulosa dentibus 5 revolutis, fœmineæ stam. sterilia gerentes. Achænium lineare calve breviter rostratum stipitatumque. — Arbor Americæ sequinoct. Ram. angulati puberuli. Folia alterna subdenticulata subtriplinervia. Panicula fastigiata terminalis ramosissima foliata. Corollæ albæ ex Kunth, flavæ ex Less. — Genus valdè dubium.

1. *A. DUBIUM* (Less. l. c.), 5 in valle Tarquensi regni Quitensis. Monactidia H. B. et Kunth nov. gen. am. 4. p. 187.

Div. II. MILLERIÆ. — Helianthæ Milleriæ veræ et Baltimore Cass. opusc. 60. — Asteroideæ Melampodiæ Milleriæ et Scaevionideæ Ambrosiæ Less. syn. 2-15 et 219.

Capitula monoica, fl. nempe in iisdem capitulis fœmineis sæpè paucis ligulis aut tubulosis subtrifidis, masculis centralibus tubulosis. Achænia ecoriatis non cum squamis inv. seu paleis rec. non concoreta exalata, calva aut raris pappi coroniformi parvo superata. Recept. interdum epaleaceum. Antheræ cœcæ.

- CCLXIV. *ELVIRA*. — *Elvira* Cass. dict. (1824) 30. p. 67 et *Meratia* Cass. l. c. p. 65 non Lous. — *Delilia* Spreng. bull. philom. 1823. apr. non Kunth. — *Emelia* fl. mexic. ic. ined. Milleria sp. L.

Capitulum monoicum biflorum, flore altero fœmineo fertè ligulato, altero masculo tubuloso 5-fido. Invol. 2-4-phyllum squamis oppositis, altero majore orbiculato basi cordatâ subcrenato, altero minore ovali. 5-4-ve ubi adsunt oblongæ acutæ. Recept. minimum epaleaceum. Cor. fœminea longè tubulosa: ligulâ brevi integerrimâ? Anth. ♂ vix exsertæ Styl. abortivus breviter bifidus. Styl. ♀ apice bifidus. Achænium planobcompressum ovale involucri immutato tectum nec involutum, calva. — Herba Americana annua trichotoma. Folia opposita petiolata ovata aut oblonga acuminata serrulata triplinervia pubescenti-subscabrâ. Pedicelli in axillis supremis aggregati 1-cephali subumbellati. — Descr. Sprengelii pessima; ex eâ sagacius. Cassinius finxit Meratiam suam cujus descriptio multò præcellit descr. autoptæ sed adhuc tamen subinstructa. Nomen et descr. fl. Mex. optima, sed inedita et ideo *Elvira* se-

men admittendum, etiamsi sensûs saltem cogniti expers. An capit. interdum 3-florum, ut patet ex descr. Linassi et Spreng.?

1. *E. biflora*. (Y) in agro Campechiano (Mart.), in umbrosis circa Bahiam (Salzm.), Novâ-Andalusâ prope Camanacoa (H. et B), insulâ Cubâ juxta Havanam (Lamagr.), ad Santa-Martha (Bert.) et in Mexicâ ditione Oaxacâ circa prædium Tapanala (Alam.). *Milleria* Mart. cent. t. 47. f. 1. Linn. hort. Cliff. t. 25 *Milleria biflora* Linn. sp. 1301 Gærtn. fr. 2. p. 423. t. 168. Lam. ill. t. 710 f. 1. *Delilia Berterii* Spreng. bull. phil. 1823 apr. syst. 3. p. 674. *Elvira Martyni* et *Meratia Sprengelii* Cass. ll. ec. (v. s.)

CCLXV. *MILLERIA* Cass. dict. 3o (1824). p. 67. fig. p. 235. Less. syn. 215. *Milleria* sp. Linn. Mart. Gærtn.

Capitulum monoicum pauciflorum, fl. fœmineo ligulato, cæteris masculis tubulosis 5-dent. Invol. 3-5-phyllosum, altero majore cucullato. Recept. epaleaceum. Ligulæ profundè trifidæ. Stylus ♂ integerrimus, ♀ bifidus. Achæmium nitidum plano-obcompressum calvum involucreo demum carnosio indurato atque irregulariter gibbo omninò involutum, areolâ laterali. — Herba Americana annua trichotoma subvillosa. Folia opposita petiolata ovato-rhomboidæ tenerè membranacea triplinervia dentata. Stipula ovata sæpè reflexa utrinquè inter folia ut in Rubiaceis. Pedunculi terminales dichotomi, pedicelli è dichotomiis et apice ramorum. Cor. luteæ.

1. *M. quinqueflora* (Linn. sp. 1301), canle subtetragono pubescente, foliis latè ovatis subcordatis abruptè acuminatis serratis infernè in petiolum longè decurrentibus. (Y) in Panamá, Verâ-Cruce, Mexico (Linn. Cav.), in valle Sameti-Jacobi (H. et B.), in Chilapæ circuitibus (fl. mex. ic. ined.). Gærtn. fr. 2. p. 423. t. 168. f. 5. Lam. ill. t. 710. f. 2. *M. dichotoma* Cav. ic. 1. p. 58. t. 82. — Mart. cent. t. 41. Caulis obtusè tetragonus variat glaber pubescens aut villosus. Florum capituli 5-9 (v. s.)

β. *maculata*, foliis maculatis profundius serratis Mart. cent. t. 47. f. 2. — *M. maculata* Mill. dict. n. 2.

2. *M. glandulosa*, canle tereti pilis glandulosis asperulo, foliis ovalibus acuminatis serraturis callosis minimis donatis, in petiolum latè alatum attenuatis. (Y) in Mexico legit el. Berlandier (pl. exa. n. 9551). Petiolus seu nervus medius infra folii dilatationem latus, superne glaber, ad marginem utrinquè serie pilorum quasi sursum ciliatus. Pedicelli fructu ferè breviores. Capitulum minimum. Fructus grossus obtusus irregulariter rugosus constans squamis 2 coriaceis subinaequalibus, achæmio obovato laevi longitudinaliter striatulo apice mucronulato calvo. Cæt. ut in ic. cit. Gærtn. (v. s.)

#### † *Species exclusæ.*

*M. alba* Hortul. — *Fougeronxia alba*.

*M. angustifolia* Cav. = *Flaveria angustifolia*.

*M. biflora* Linn. = *Elvira biflora*.

*M. contrayerva* Lam. = *Flaveria contrayerva*.

CCLXVI. *RIENCOURTIA* Cass. bull. philom. 1818. p. 76. dict. 45. p. 466. Less. syn. 215. — *Petrantha* Poit. ms.

Capitulum monoicum 4-7-florum discoideum heterogamum, fl. disci 3-6 masculis, radii unico fœmineo tubuloso. Invol. oblongum disco brevius, squamis biserialibus in utrâque serie 4 æqualibus adpressis coriaceis. Recept. parvum nudum. ♂ Cor. disci fauce amplâ, limbo 4-5-dentato, lobis fasciculo pilorum superatis. Anth. nigricantes. Sty-

lus symplex exsertus. ♀. Cor. radii cylindrica 3-dentata. Styl. uni-  
due. Achanium obcompressum ovale s. orbiculatum calvum. — Herba  
erecta ramosa Australi-Americana. Folia opposita petiolulata lan-  
olata 3-nervia pilosa subdentata. Capitula spicata vel glomerata ad spi-  
ces ramorum, superne aphyllorum. — Conjug. cl. Cassini orta à gen-  
te Riencourt.

1. *R. stricteolivera* (Cass. fl. ex.), capitulis in spicam dispositis distichis et  
bracteorum axillis brevibus, rachis hispida, bracteis ellipticis. — in Annali  
austr. ? Corollae macula 5-dentata. (v. s. specim. incomp. in h. Jan.)
2. *R. glomerata* (Cass. dist. 43. p. 376. 45. p. 468), capitulis plurimis in glo-  
merulos terminales solitarios subglobosis collectis bracteis, bracteis qui-  
bus piloso-hispidis. — in Galia gallica. Tetraena suaveolens Poit. inel.

## CCLXVII. LATRILIZA.

Capitulum monoicum multiflorum, fl. radii 3-4 feminis, flus  
masculis. Recept. subspiciforme. Invol. squame 1-3-seriales, ciliatis  
breves potius bracteis dicendis, inter. majores concavas. Rec. plus  
late ovales suborbiculatis ciliatis inter flores sitis. Cor. radii breviter  
bulosus subtruncatus pilos articulates extis gerentes, disci 5-fidis  
glabris. Antherae concretis eadentis. Stylus radii bifidus, ramis 2 cili-  
atis glabris acuminatis intus sulcatis; disci simplex inclusus abortiva.  
Achania radii grossa glabra obcompressa, disci abortiva. — Herba  
Brasilensis habitu paululum Chromolaena. Rhizoma lignosum. Culi  
herbacei basi striatis glabris apice puberulus aut hispidus. Folia op-  
posita sessilia oblonga, infer. glabra, super. hirtella. Corymbes breves  
trifidis hirtis; capitulis circ. 3-5 ovalis lupulaceis. — Dixi in me-  
moriam cl. Petr. Andr. Latreille entomologi peritissimi, de geographia  
etiā bene meriti et omnibus naturae curiosis morum amantibus cari.

1. *L. serrata*, caule dichotomo basi glabro apice puberulo, foliis oblongis  
coarctatis à medio ad apicem grossè et remotè callosis-serratis. ♀. in campis  
prov. Sancti-Pauli novemb. flor. legit cl. entomologus Lund. Folia 3 pol-  
longa, 4-6 lin. lata subtriplinervia. Capitula nivea. Invol. veri squame lim-  
bo ovate apice acuminatae. Rec. paleae apice late fimbriatae basi angustatae (v. s.  
comm. à cl. Lund.)
2. *L. integrifolia*, caule simplici basi glabro apice hispido, foliis lanceo-  
latis integerrimis. ♀. in Brasiliis prov. Sancti-Pauli ex h. Mus. imp. Bras. a 44.  
Invol. squame obtusissimae dorso hispidae. (v. s. in h. Mus. reg. Par.)

## CCLXVIII. ICTYOTHERE Mart. arzn. pfl. p. 27.

Capitulum monoicum, floribus omnibus tubulosis, radii feminae  
subquinis fertilibus, corollulae limbo in fimbriis tenuissimis lacinis.  
disci plurimis 5-fidis. Involuorum minimum cupuliforme persistens.  
Recept. paleaceum, paleis singulis florem singulum foventibus. Antherae  
basi intus denticulo auctae. Stylus radii bifidus, disci steriles. Achania  
intra paleas extimas concavas grandefactas obovata absque pappo. —  
Herba perennis, foliis oppositis, petiolis breviter vaginantibus limbo  
argutè serratis. Capitula florum pedunculata terminalia aggregata,  
rachis post grossificationem per dichotomiam excrecente, involucri  
persistentibus tunc grandefactis. — Affinis Bailleriae Cacosmiae et Ma-  
lampodio ex Mart.

1. *I. CUNANI* (Mart. l. c.). — in Brasiliis prov. Rio-Negro. Pileos involucri  
nom. gener.

CCLXIX. CLIBADIUM *Linn. mant.* 161 *non Neck.* — Clibadium, *Oswalda et Baillieria Cass.* 59. p. 324. — *Baillieria et Oswalda Less. syn.* 215. — *Baillieria Aubl.* — *Trixia Swartz, Pers. non P. Brown.*

Capitulum monoicum pluriflorum discoideum, fl. radii paucis tubulosis 5-fidis hermaphroditis sterilibus. Invol. ovatum, squamis conniventibus ovatis concavis, interioribus achænia radii in parte concava semifloventibus. Recept. planum in disco omnino epaleaceum, paleas seu invol. squamas inter. in radio interdum biseriali gerens. Stylus radii bifidus, ramis divaricatis subrevolutis glabriusculis subobtusis, disci ex *Cass.* indivisus vix subbifidus apice clavatus hispidulus. Achænia radii grossa obovata intus obcompressa extus convexa epapposa apice pilis articulis villosa, disci abortiva gracilia linearia apice villosissima. — *Herbæ seu suffrutices Americani.* Rami teretes sæpius subvillosi. Folia opposita petiolata ovata aut lanceolata serrata plerumque aspera. Capitula ovata albida parva in paniculam sæpius corymboso-fastigiatam disposita.

Sect. I. EUCLIBADIUM. — Clibadium *Linn. l. c. Cass. dict.* 9. p. 395. 29. p. 176 et 181. 59. p. 324.

Achænia drupacea succo viscoso scatentia, radii uniserialia. Paleæ recept. nullæ.

1. C. SURINAMENSIS (*Linn. mant.* 294), foliis petiolatis ovatis acuminatis acutè crenatis scabris, pedunculis oppositis. — Surinami. C. fœtidum *Alam. ex Linn.* Invol. violaceum. Cor. alba. — Species ab hodiernis ign. Habui ex Amer. sub hoc nom. fragmentum sine flore, cum descriptione non malè conveniens. Rami et petioli hirsutuli. Petioli 10-12 lin. longi. Folia 4 poll. longa, 2  $\frac{1}{2}$  poll. lata, argutè serrata, suprâ scabrida, juniora villosa demum scabrida. (v. s.?)

Sect. II. OSWALDA *Cass. dict.* 59. p. 319 et 322. *Less. syn.* 215. — *Baillieria Aubl.* — *Baillieria sp. Spreng.* — *Trixidis sp. Sw. Pers.*

Achænia sicca villosa, radii uniserialia obcompressa obovata dorso convexa, disci gracilia linearia apice hispida. Paleæ recept. nullæ.

2. C. ROTUNDIFOLIUM, foliis brevissimè petiolatis ovato-subrotundis basi triplinerviis utrinque subscabridis, paniculâ corymboso-fastigiata.  $\S$  in Brasiliâ circa Bahiam legerunt cl. Salzmann, Blanchet, etc., in prov. Sancti-Pauli (h. *Mss. Bras.* n. 466). Petioli 3-4 lin. longi. Limb. fol. 3 poll. longus, circit. 2 poll. latus. Capitula ovato-globosa. Pedunculi hirsuti. Inv. squamæ pubescentes, inter. glabræ multistriatæ acutæ. (v. s.)

$\beta$ . Foliis ovatis acutis. — in prov. Sancti-Pauli (Lund!), et prov. Minarum Generalium (Gaudichaud!). (v. s.)

3. C. PERUVIANUM (Poepp.! pl. exs. n. 35. diar. n. 1611), foliis petiolatis ovali-lanceolatis longè acuminatis basi cuneatis adpressè serrulatis triplinerviis suprâ scabris subtus ramis petiolisque hirsutis, paniculâ corymboso-fastigiata.  $\S$  in Peruvia legit cl. Poeppig. Petioli 4-5 lin. longi. Limb. fol. 4 poll. longus, 10-12 lin. latus. Capitula ovato-globosa. Inv. squamæ ext. dorso puberæ, intus glabræ striatæ apice puberæ acutæ. (v. s.)

4. C. TRINITATIS, foliis petiolatis ovali-lanceolatis basi longè cuneatis apice acuminatis acutè serrulatis subtrinerviis, suprâ secus nervos et vix inter nervulos scabridis, subtus ramisque hirtello-subhirsutis, paniculâ ovata.  $\S$  in insulâ Trinitatis legit cl. Sieber fl. trin. pl. exs. n. 71! Petioli 6-7 lin. longi. Limb. fol. 5 poll. longus, 1  $\frac{1}{2}$  latus. Capitula ovato-globosa parva. Panicula non fastigiata, foliis sub brevior. Inv. squamæ pauca extus subpuberæ apice eroso subdenticulatæ. (v. s.)

flava. — Differt ab *Unia* acheniis nec compressis nec areolâ laterali depictis; à *Villanova* et *Chlamispermate* disco masculino; ab omnibus acheniis glanduloso-puberis et foliis alternis. — Nomen ex *areola*, tener.

1. *A. ANTHEMIFOLIUS*. ① in paucis sterilibus circa *Ranegua*, *Valparaiso* et al. montem *La Laguna Chilensis* legit cl. *Bertero*. *Unia anthemifolia* *Bert.* *lib.* n. 726.

CCLXXII. *FRONACRON* *Cass. dict.* 43. p. 390. *Less. syn.* 214.

Capitulum monoicum multiflorum, fl. radii 5 uniseriatis feminis ligulatis, disci circ. 12 masculis tubulosis. Invol. subglobosum limbi disco aequale, squamis 7, duabus ext. rotundatis villosis subaequalibus, 5 int. glabris membranaceis. Recept. planiusculum, paleis paucis inaequalibus subulatis. Cor. radii tubo glanduloso, ligulâ ellipticâ integerrâ. Achenia glabra, disci gracilia, radii lateraliter plano-campum subenticularia crassa laviâ truncata calva. — Herba ramosissima *Guianensis* villosa. Folia opposita lanceolata subdentata. Capitula pedunculata terminalia inter ramulos novellos orta. Flores flavi.

1. *P. RAMOSISSIMUM* (*Cass. l. c.*). ①? in *Guianâ* gallicâ.

CCLXXIII. *ATLOTHECA*.

Capitulum monoicum multiflorum, fl. radii circ. 5-6 ligulatis feminis, disci tubulosis 5-dent. masculis. Invol. biseriale, squamis ext. oblongis erectis 8-10 disco brevioribus, inter. paleas referentibus et acheniis radii semi-includentibus. Recept. parvum convexum, paleis membranaceis cuneatis apice dentatis acutis achenia involv. Ligulae breviter obtusae 3-dentatae. Achenia radii villosa subtrigona epapposa villis subcoronata inter squamas int. et paleas ext. inclusa, disci gracilia gibberima calva. — Herba *Mexicana* incana apice corymboso-ramosa. Caulis teres basi glabratus. Folia alterna sessilia oblonga obtusa sinuata. Capitula corymboso-fastigiata. Cor. lutea. — Habitus herbae affinis *Parthenio* incano sed fructus diversus et flores lutei. — Nomen à *ατλος*, varius, et *θηκα*, theca.

1. *A. PARTHENIOIDES*. — in Mexici ditione Sancti-Ludovici de Potosi decembris flor. legit cl. *Berlandier* (pl. exs. n. 1341). Herba circ. pedalis. (v. a.)

CCLXXIV. *TRIGONOSPERMUM* *Less. syn.* 214. et in *linnaea* 1834. p. 26.

Capitulum monoicum minutum pluriflorum, fl. radii 3 foemineis ligulatis profundè 3-4-lobis parvis, disci crebrioribus masculis tubulosis. Invol. 6-8-phyllum, squamis linearibus acutis 1-nerviis, alternis majoribus rotundis. Recept. paleis scariosis difformibus, ext. majoribus erosis, int. linearibus. Stylus ♂ integerrimus inclusus abortivus, ♀ bifidus. Achenia disci 0, radii triangularia squamâ int. et paleâ cincta lucida calva. — Herba annua *Mexicana* erecta dichotoma. Folia opposita triplinervia rhombeo-ovata breviter acuminata usque ad basin petioli cuneato-decurrentia utrinque hirsutiuscula. Paniculae cymosae. Rami glanduloso-hirsuti. Capitula lutea.

1. *T. ADENOSTEMOIDES* (*Less. l. c.*), capitulis minimis subdiscoideis, ligulis nequidiz discum superantibus. ① in Mexico circa *Jalapam*. Herba ultra-organica. Capitula parva. Cor. masc. 4-dentata. Folia 6 poll. longa, 3 lata.

12. C. ? PEDUNCULOSUM, herbaceum, ramis gracilibus apice summmitate promissâ nudis 2-3-cephalis, foliis lanceolatis serratis 5-nerviis hispidis. — in Cayennâ. Trixis pedunculosa Rich. act. soc. h. n. Par. 1792. p. 112. Pers. ench. 2. p. 491. Baillieria pedunculosa Spreng. syst. 3. p. 625.
13. C. ? BARBASCO, fruticosa, foliis oppositis acutis basi valdè angustatis remotè et obsoletè serrulatis suprà pilosiusculis subtus adpresso-pubescentibus, corymbis sessilibus triobotomis, recept. paleis oblongis singulis flore masculo duplò brevioribus, ovario compresso? dentibus 2 minutis apice instructo. 5 in ripâ flum. Tuamini et Tima prope Javita in missionibus Orinoci. Baillieria Barbasco H. B. et Kunth nov. gen. am. 4. p. 289. Vulgò dicitur *Barbasco*. An hujus generis?
14. C. ? NERITIFOLIUM, arboreum, ramis albo-lanatis, foliis sparsis verticillato-congestis lanceolatis integerrimis coriaceis suprà glabris subtus albo-lanatis, corymbis terminalibus villosissimis, invol. squamis interioribus lineari-lanceolatis, recept. paleaceo convexo-conico, fl. femineis ligulatis, achænio margine villosa coronato. 5 in monte Sillâ de Caracas ubi dicitur *Incienso*. Baillieria neritifolia H. B. et Kunth nov. gen. am. 4. p. 289. Trixis neritifolia Humb. rel. hist. 1. p. 645. A genere certè removenda! An *Melampodii* sp.?

CCLXX. UNXIA Linn. f. suppl. 56. Juss. gen. 186. Less. syn. 220 non Kunth.

Capitulum monoicum pluriflorum, fl. radii 5 ligulatis femineis parvis ovatis, disci 5 masculis tubulosis 5-dentatis. Invol. subrotundum, squamis 5 ovatis. Recept. nudum? Achænia radii ovata dura (Linn. f.) compressa lævissima calva, areolâ laterali (Less.). — Herbas dichotome Guianenses. Folia opposita hirsuta oblongo-ovata acuminata basi rotundata 5-nervia villosa. Capitula axillaria breviter pedunculata. — An *Clibadio* affinis? †.

1. U. CAMPHORATA (Linn. f. suppl. 368), foliis lanceolatis, capitulis 10-floris. — in locis arenosis Surinami. Odor gravis camphoratus. *Camphortplant Surinam* Holland.
2. U. HIRSUTA (Rich. act. soc. Par. 1. p. 105), foliis oblongis subcordatis, capitulis multifloris. (1) Cayennæ. Pers. ench. 2. p. 492.

### †† Species exclusæ.

- U. ACHILLEOIDES Spr. = Villanova achilleoides.  
 U. ALTERNIFOLIA Spr. = Villanova alternifolia.  
 U. ANEMONEFOLIA H. B. K. = Villanova anemonefolia.  
 U. ANTHEMIFOLIA Bert. = Apalus anthemifolius.  
 U. DISSECTA Hook. = Villanova dissecta.  
 U. PRATENSIS H. B. K. = Chlemisperma pratensis.

### CCLXXI. APALUS.

Capitulum monoicum, fl. radii circ. 5-7 uniseriatis femineis ligulatis ovati-oblongis integris, disci circ. 8-10 masculis glabris tubo tereti limbo campanulato 5-dentato. Invol. 5-7-phyllum uniseriale, squamis ovatis. Recept. nudum angustum. Styl. radii bifidus, ramis teretibus extrorsum arcuatis. Achænia radii oblonga erostris calva sulcato-angulata pube glandulosâ brevi crassâ velutino-hirtellâ, disco epigyno. — Herba Chilensis tenella annua erecta glabra versus apicem subvillosa. Folia alterna semiamplexicaulia, basi linearia petioliformia integra apice pinnatipartita, lobis linearibus oppositis integris obtusis utrinquè 2-4. Rami apice pedicelliformes 1-cephali. Cor. disci et radii

flavae. — Differt ab *Uxia* acheniis nec compressis nec areolâ laterali depictis; à *Villanovâ* et *Chlamispermate* disco masculino; ab omnibus acheniis glanduloso-puberis et foliis alternis. — Nomen ex *arab.* tener.

1. *A. ANTHEMIFOLIUS*. (1) in paucis sterilibus circa *Ramagum*, *Valpurga* et al. montem *La Laguna Chilianam* legit cl. *Bertero*. *Uxia anthemifolia* *Benth.* *h.* n. 736.

CCLXXII. *PRONACRON* *Cass. dict.* 43. p. 370. *Less. syn.* 214.

Capitulum monoicum multiflorum, fl. radii 5 uniseriatis feminis ligulatis, disci circ. 12 masculis tubulosis. Invol. subglobosum limbi disco aequale, squamis 7, duabus ext. rotundatis villosis subsejunctis, 5 int. glabris membranaceis. Recept. planiusculum, paleis parvis inaequalibus subulatis. Cor. radii tubo glanduloso, ligula elliptica integra. Achenia glabra, disci gracilia, radii lateraliter plano-concavo sublenticularia crassa laevia truncata calva. — Herba *ramagum* *Guianensis* villosa. Folia opposita lanceolata subdentata. Capitula pedunculata terminalia inter ramulos novellos orta. Flores flavi.

1. *P. RAMOSESIMUM* (*Cass. l. c.*). (1) in *Guianâ* gallicâ.

CCLXXIII. *ANLOTHECA*.

Capitulum monoicum multiflorum, fl. radii circ. 5-6 ligulatis feminis, disci tubulosis 5-dent. masculis. Invol. biseriali, squamis ext. oblongis erectis 8-10 disco brevioribus, inter. paleas referentibus et acheniis radii semi-includentibus. Recept. parvum convexum, paleis membranaceis cuneatis apice dentatis acutis achenia involv. Ligula breviter obtuse 3-dentata. Achenia radii villosa subtrigona epapposa villis subcoronata inter squamas int. et paleas ext. inclusa, disci gracilia gibberima calva. — Herba *Mexicana* incana apice corymboso-ramosa. Caulis teres basi glabratus. Folia alterna sessilia oblonga obtusa sinuata. Capitula corymboso-fastigiata. Cor. lutea. — Habitus herbae affinis *Parthenio* incano sed fructus diversus et flores lutei. — Nomen à *αἰολος*, varius, et *θησαυρος*, theca.

1. *A. PARTHENOIDEA*, — in *Mexici* ditione *Sancti-Ludovici de Potosi* decembris flor. legit cl. *Berlandier* (*pl. exs.* n. 1341). Herba circ. pedalis. (v. a.)

CCLXXIV. *TRIGONOSPERMUM* *Less. syn.* 214. et in *linnaea* 1834. p. 36.

Capitulum monoicum minutum pluriflorum, fl. radii 3 foemineis ligulatis profundè 3-4-lobis parvis, disci crebrioribus masculis tubulosis. Invol. 6-8-phyllum, squamis linearibus acutis 1-nerviis, alternis majoribus rotundis. Recept. paleis scariosis difformibus, ext. majoribus erosio, int. linearibus. Stylus ♂ integerrimus inclusus abortivus, bifidus. Achenia disci 6, radii triangularia squamâ int. et paleâ et cincta lucida calva. — Herba annua *Mexicana* erecta dichotoma. Folia opposita triplinervia rhombeo-ovata breviter acuminata usque ad basin petioli cuneato-decurrentia utrinquè hirsutiuscula. Paniculae cymosae. Rami glanduloso-hirsuti. Capitula lutea.

1. *T. AENEOSTEMOIDEA* (*Less. l. c.*), capitulis minimis subdiscoideis, ligulis nervis discum superantibus. (1) in *Mexico* circa *Jalapam*. Herba ultra-organa. Capitula parva. Cor. masc. 4-dentata. Folia 6 poll. longa, 3 lata.



2. *T. MELAMPODIOIDES*, capitulis evidenter radiatis, ligulis nempe disco triplo circ. longioribus. — in Mexico ad montem San-Felipe prope Oaxacam legit cl. Andrieux (pl. ex. n. 320). Caulis subhirsutus. Folia 3 poll. longa, sesquipoll. lata. Paniculae rami pilos parcos et glandulas turbinatas gerentes. Capitula magna. ferè Siegesbeckiæ. Ligulæ 3-5 elongatæ apice irregulariter lobato-dentatæ, discique cor. aurantio-flavæ. Achænia trigona nigra. (v. s. ex coll. Andr.)

## CCLXXV. XENISMIA.

Capitulum monoicum, fl. radii 5-7 fœmineis ligulatis, disci 10-12 tubulosi 5-dentatis masculis. Invol. 1-serialis squamæ oblongæ. Recept. epaleaceum. Ligulæ obtusæ basi ciliatæ. Anth..... Styl..... Achænia disci 0, radii crassa glabra aculeis rigidis crassis undiquè echinata calva. Semen crassiusculum oblongum basi attenuatum. — Herba Australi-Africana annua subpuberula multicaulis erecta. Radix exilis. Folia alterna oblongo-cuneata apice dentata basi attenuata. Capitula ad ramorum apices pedicellata solitaria parvula. — Nomen (ex ξενισμός, peregrinitas) alludit ad habitationem plantæ quæ sola totius divisionis in Africâ versatur.

1. *X. AGANTHOSPERMA*. ① in Africâ austr.-orient. legit cl. Drege! Herba 5-6 poll. Habitu Acanthospermum humile paulò referens. (v. s.)

CCLXXVI. SCOLOSPERMUM Less. in *linnæa* 1830. p. 152. t. 2. f. 19-31. 577. p. 219.

Capitulum pluriflorum monoicum, fl. radii fœmineis ligulatis, disci masculis tubulosi obconici 5-dentatis. Invol. triserialis squamæ in seriebus 2 ext. septenæ subæquales. Recept. paleaceum. Stylus fl. ♂ indivisus exsertus teres obtusiusculus apice hirsutiusculus. Achænia calva triquetra versus apicem tuberculata in squamis invol. intimis capsulatum inclusa. Embryo triqueter. Cotyledones conduplicate. — Herba Mexicana erecta dichotoma Fougereuxiæ facie. Folia opposita petiolata ovato-lanceolata acuminata serrata triplinervia. Capitula in dichotomiis axillaria et terminalia pedunculata lutea.

1. *S. FOUGEROUXIÆ*. — in Mexico ad Santa-Fé (Schied.) et ad Panamam (Hænk.!). 8. *Baltimoroides* Less. ll. cc. Petioli breves hispidi. Folia utrinquè adpressè villosa. (v. s. in h. Hænk. à cl. de Sternberg miss.)

CCLXXVII. FOUGEROUXIA. — *Baltimora* et *Fougeria* Moench *meth.* 592 et suppl. 243. *Cass. dict.* 17. p. 283. 46. p. 399 et 412. — *Baltimora* Linn. *mant.* 288. *gen. n.* 1323. *Gærtn. fr.* 2. p. 443. t. 169. *Lam. ill.* t. 709. — Niebuhr *Scop. non.* DC.

Capitulum pluriflorum monoicum, fl. radii 5 ligulatis fœmineis, disci masculis tubulosi 5-dent. Invol. 2-serialis squamæ adpressæ lanceolatæ, ext. foliaceæ. Recept. planum, paleis amplectentibus membranaceis apice ciliato-fimbriatis. Cor. disci extus hirsutiusculæ. Achænia radii juniora subtrigona obcompressa pappo coroniformi dentato deciduo (1), adulta trigona omninò calva lævia, disci abortiva elongata compressa papulosa. — Herbæ annuæ erectæ Americanæ. Caulis obtusè tetra-

---

(1) Quo pappi delapsu plurimi decepti fuerunt, et præ aliis Moenchius qui plantam adhuc pappiferam sub nom. *Baltimoræ* et pappo dejecto sub nom. *Fougeriæ* descripsit.

genus subscaber. Folia opposita petiolata ovata acuta serrata trinervi subscabrida. Capitula pedicellata è dichotomiis orta aut rarissime in iisdem indiv. oppositifolia. Flores pallidè flavi aut albi? — Nomen Baltimore rejiciendum! cum planta ad Baltimorem non crescat, admisi cum mutatione parçâ etymologiæ conformi nomen Mœnchii in honorem cl. Fougereux de Bondaroy Gaillardæ monographi botan. et agronomi periti sæc. XVIII.

1. F. RECTA, foliis pubescentibus serratis, pedicellis elongatis. (X) in Mexico (Lam.) fortè circa Veram-Crucem nec certè circa Baltimorem (Pursh. Nutt.). *Baltimore recta* Linn. mant. 288 excl. Pluk. syn. et ideò patriâ. Schkuhr. handb. 3. a. 261 b. opt. Baltimore trinnervata Morach meth. 59a, et Fougeria tetragona Mœnch suppl. 243. Moris. oxon. 5. 7. t. 23. f. 16. (v. v. cult.)
2. F. ALBA, foliis subtomentosis subserratis, capitulis subsensilibus. — cùm in horto Trianonesi culta ex specim. herb. Rich. sub nom. Milleris alba. Baltimore alba Pers. ench. 2. p. 489.

CCLXXVIII. CHRYSOGONUM Linn. gen. n. 988. Gartn. fr. 2. p. 436. t. 174. Lam. ill. t. 713. Less. syn. 220. — Chrysogonum et Diotostephus Cass.

Capitulum multiflorum monoicum, fl. radii circ. 5 uniseriatis ligulatis fœmineis, disci masculis tubulosis 5-dentatis obtusis. Recept. plenum, squamis linearibus obtusis. Cor. disci glabrinsculas, tubo brevissimo, fauce cylindricâ. Achænia radii obcompressa vix subtrigona glabra pappo coroniformi persistenti coronata, disci abortiva linearia elongata pappo coroniformi dentato. — Herbæ perennes Boreali-Americanæ plus minùs hirsutæ humiles. Caules è collo plurimi, alii floridi, alii stoloniformes. Folia alia radicalia, alia secus caules opposita petiolata ovata crenata. Capitula solitaria breviter pedicellata. Flores lutei.

1. C. VIRGINIANUM (Linn. sp. 1303), petiolis limbo longioribus, foliis demum glabratâ, pappo radii coroniformi ad latas interius fissas apice crenato. ? in Virginia? (Linn.), in Carolinâ superiore (B. Greene!). Gartn. fr. 2. p. 433. t. 174. f. 8. Pluk. alm. 100. t. 83. f. 4 (excl. altero Pluk. syn. à Linn. et Willd citato). Lam. ill. t. 713. f. B. (v. a.)
2. C. DIOTOSTEPHUS, hirsutissimum, petiolis limbo subbrevioribus, pappo radii biarrito. ? in siccis pinguibus Carolinæ. Diotostephus repens Cass. dict. 41. p. 543. Chrys. Virginianum Michx. amer. 2. p. 472. Ell. sketch 2. p. 471. — Lam. ill. t. 713. f. A. Achænia radii ante squamas invol. interiores sita et cum floribus disci circit. 4. 5. à recept. cum squamâ ipsâ separata. (v. a.)

† Species vix nota.

C. DICHOTOMUM (Vahl in Vest St.-Cr. p. 303. sine descr.). — in insulâ Sandæ-Crucis.

†† Species exclusa.

C. PERUVIANUM Linn. = Zinnia pauciflora.

DIV. III. SILPHIÆ. — Asteroidæ Melampodiæ Silphiæ Less. syn. 213.

Capitula monoica multiflora, fl. fœmineis pluribus (3-10) ligulatis. Achænia eecorticata calva aut biaristata. Recept. paleaceum, paleis squamisve nunquam cum achæniis concretis.

CCLXXIX. *GUARDIOLA* *H. Bonpl. pl. eq. 1. p. 144. t. 41. H. B. et Kunth nov. gen. am. 4. p. 247. Cass. dict. 20. p. 12. 59. p. 319. Less. syn. p. 419. — Guardi-  
ola Steud. nom. 304.*

Capitulum pluriflorum heterogamum, fl. radii 3-5 ligulatis fœmineis, disci 10-11 masculis tubulosis 5-dentatis. Invol. 3-phyllum, squamis ovalibus erectis. Recept. planum angustum paleaceum. Ligularum tubus longus basi hispidus, cor. radii tubus longior glaber. Achænia disci compressa (Kunth) seu obcompressa (Cass.) lævia calva, disci abortiva. — Herba Mexicana glabra trichotoma. Folia opposita longè petiolata ovato-lanceolata acuminata serrata basi subcuneata. Capitula terna pedicellata corymbosa terminalia. Discus flavus, antheris nigricantibus, radius albus brevis 3-dentatus. — Certè *Helianthea* sensu Cassiniano. An *Millerieis*, an *Silphieis* adjungenda?

t. *G. MEXICANA* (H. et B. l. c. t. 41). ① in aridis Mexici inter Ario-Aguasarco et montem Jorullo.

CCLXXX. *HIDALGOA* *Less. in linnæa 1831. p. 406. syn. 213. non Lallav.*

Capitulum monoicum multiflorum, fl. radii 5 uniseriatis fœmineis ligulatis, radii tubulosis abortu masculis. Inv. cylindraceo-campanulatum. biseriale disco brevius, squamis integerrimis obtusis. Recept. planum, paleis membranaceis linearibus. Stylus disci simplicissimus clavatus, radii bifidus ramis semi-lanceolatis. Achæmium disci lineare, radii valdè compressum læve ellipticum angustè bialatum apice continuatum in pappum brevissimum bicornem. — Herba scandens Mexicana. Folia alterna petiolata pedati- seu ternati-secta, segmentis lateralibus plus minus bilobis. Capitula solitaria longè pedunculata aurantiaca.

. *H. LESSINGII*. — in Mexico temperato, in sylvia Misanthe, in Cuesta-Grande de Yala-Cingo. *H. ternata* Less. ll. cc. non Lallav. Ex descr. auctorum differt à stirpe Lallavei (mibi ut Sprengelio *Melampodio* adjungenda), genericè fructu nec drupaceo nec apice cucullato; specificè foliis alternis nec oppositis, pedatis nec strictè ternatis-sectis, segmentis lateralibus bilobis.

#### † *Species exclusa.*

*H. ternata* Lallav. = *Melampodium Hidalgoi*.

CCLXXXI. *SILPHIUM* *Linn. gen. n. 1334. Cass. dict. 59. p. 319 et 34. Less. syn. 213.*

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis, ligulis 1-serialibus, fructibus pluriserialibus, fl. disci tubulosis 5-dentatis abortu styli masculis. Invol. campanulatum imbricatum, squamis basi adpressis strictis apice foliaceis laxis. Recept. subconvexum paleaceum. Cor. disci tubo brevissimo, fauce cylindracea lobis, superne puberulo-hirtis. Stylus indivisus apice hispidus. Achænia radii plano-obcompressa bialata emarginata dentibus aristivè 2 nunc liberis nunc cum alâ utrinquè concretis coronata. — Herba Boreali-Americana sæpius elata interdum gummifera, caule tereti aut tetragono, foliis alternis verticillatis aut oppositis sæpè connatis. Capitula pedicellata. Flores lutei. — *Silphium* veterum erat etiam herba gummifera inter Umbelliferas querenda.

§. 1. *Tereticaulia, caulibus teretibus seu obtusis subterangulis; foliis alternis aut tornato-verticillatis, achantis apice biseriatis et bidentatis.*

\* *Foliis pinnatifidis alternis.*

1. *S. LACINATUM* (Linn. sp. 1301), caule tereti superne hispido, foliis alternis petiolatis pinnatifidis scabris amplexicaulibus, lobis acutis indivisis aut dentato-sinatis, invol. squamis hispidulis basi ciliatis late ovatis in apicem lanceolatum acuminatum productis.  $\frac{1}{2}$  ad ripas flum. Amer. bor. perennans ad Mississipi (Pursh), in Georgia occident. et ad montes Alleghany (Ell.), Linn. f. dec. p. 3. t. 3. Ell. sketch 2. p. 461. Jacq. f. col. 1. t. 60. *S. apiculatum* Poir. suppl. 3. p. 157. Herba 2-3-origalis. Capitulum amplum. Achenia radiis plicatula, aristis 2 cum alâ condensibus. (v. s.)
2. *S. COMPLANATUM* (Ell. sketch 2. p. 460. in bdn.), caule valde hispido et atro, foliis sinuato-pinnatifidis subscabris subhispidis, lobis acutis indivisis, invol. squamis ovatis acuminatis margine hispidis.  $\frac{1}{2}$  in pratis Alabamæ prope Georgiam occid. Herba 2-3-pedalis valde *S. laciniatum* affinis. Caulis gemmis subinthesum copiosum fundit. Capitula majora quam in omnibus aliis speciebus. Ariste subulata.
3. *S. PINNATIFIDUM* (Ell. sketch 2. p. 462), caule etiam ad apicem glaberrimo, foliis sinuato-pinnatifidis subscabris subhispidis, lobis ovatis indivisis, invol. squamis glabris obtusis, ext. rotundis, int. ovalibus, ligulis invol. circa squamulas.  $\frac{1}{2}$  in pratis Alabamæ et Georgiæ occid. Herba 4-6-pedalis. Fol. lobi supris acutis *S. pinnatifidum* Nees et. sem. h. Bonn. 1831 omnino idem?
4. *S. COMPOSITUM* (Michx. fl. bor. am. 2. p. 145), caule tereti laevi, foliis alternis petiolatis sinuato-pinnatifidis lobis obtusis secus petiolum et marginem lobi vix ac tæ vix pilosiusculis, invol. squamis ovalibus obtusis glabris.  $\frac{1}{2}$  in quibus glaberrimis maritimis à Virginia ad Carolinam (Pursh), in pinetis adiacens *Ceanothus mollis* (Wres! Rose!). *S. laciniatum* Walt. cor. 217. *S. sinuatum* Banks et ex Pursh. *S. pseudo-laciniatum* Fras. in herb. L'Hér. Radix ex Fraserum venenata. Capitula parva paniculata. (v. s.)
5. *S. SUBCAULE* (Nutt. in Sillim. am. journ. 5. 1822. p. 301.), scaberrime, caule brevissimo, pedunculis longissimis, foliis alternis sessilibus subimbricatis pinnatifidis dentatis, invol. squamis dilatatis rhomboidis. — in Florida orient. Caulis vix 2-3-poll. Folia 2-3 poll. longa exacte ut in *Senecione alba* sinuata. Pedunc. palmares.

\*\* *Foliis indivisis alternis.*

6. *S. TERREBINTHINACEUM* (Linn. suppl. 383), caule tereti striato glabro, his alternis, infer. petiolatis ovatis subcordatis serratis scabridis, sup. ovatis, panicula laxa nudiuscula, invol. squamis glabris ovatis obtusis.  $\frac{1}{2}$  in montibus Carolinæ (Ell.) et Louisiana (Pursh). Jacq. hort. vind. 1. t. 43. Garta. k. 1. p. 445. t. 171. Herba 4-5-pedalis fere tota glabra. Denticuli 2 ad apicem achenii breves ab alâ disjuncti. Variat, ex Ell. sketch 2. p. 463, foliis radicalibus reniformibus subhispidis scaberrimis et foliis lobato-angulatis subhispidis fere glabris. Prima var. est verisimil. *S. reniforme* Raf. med. fl. 2. p. 283 et ultima h. *S. elatum* Pursh fl. bor. am. 2. p. 579 ex h. Banks in Carolina indicatum.
7. *S. PUMILUM* (Michx. fl. bor. am. 2. p. 146), caule adscendente tereti tomentoso, foliis alternis petiolatis cordato-ovatis dentatis supra pubescentibus subtus petiolique subtomentosis, capitulis paucis longe pedicellatis, invol. squamis late ovatis obtusis velutinis.  $\frac{1}{2}$  in Florida (Michx.).  
 $\beta$ . *tomentosum* (Nutt. gen. am. 2. p. 183), caule petiolis foliisque sub dense albo-tomentosis.  $\frac{1}{2}$  in Georgia (Bartr.), pinetis siccis Carolinæ (Fras! Ell.). *S. tomentosum* Pursh fl. 2. p. 579. *S. pumilum* Ell. sketch 2. p. 469. Achenia adpressè villosa, vix alata, juniora setas 2 deciduas feruntia demum mutica apice hispidula. (v. s.)
8. *S. ASTERISCUS* (Linn. sp. 1302), caule tereti hispido, foliis inf. oppositis

grossè serratis, super. alternis sessilibus oblongis seu ovali-lanceolatis hirtis serratis integrive, capitula paucis solitariisve, invol. squamis ovatis ciliatis, exter. acutis.  $\frac{1}{2}$  in pratis montanis (Pursh) et arenosis siccis (Ell.). Virginie et Carolinæ. Dill. elth. 42. t. 37. f. 42. S. reticulatum Mönch. Achænia radii latè alata apice callis 2 obtusis ab alâ disjunctis coronata. (v. s. sp. et v. c.)

$\beta$ . *lævicaulis*, caule lævi, invol. squamis dorso glabris margine ciliatis apice subobtusis, achæniis dentibus paulò magis conspicuis.  $\frac{1}{2}$  in Georgia occident. Silph. Asteriscus  $\beta$  Nutt. gen. 2. p. 183. S. scabrum Walt. car. 217. S. dentatum Ell. sketch 2. p. 468. (v. s.)

9. S. QUERCIFOLIUM, caule tereti hispido densè folioso, foliis sparsis petiolatis elliptico-sublanceolatis obtusiusculis obtusè sinuato-repandis penninerviis supra strigilloso-scaberrimis subtus puberis, petiolis ciliatis, capitulis corymbosis, invol. squamis ovali-oblongis obtusis glabris subciliatis apice patentis-squarrosis.  $\frac{1}{2}$  patr. ign. Vidi vivum anno 1807, in horto Burdigalensi nec unquam in hortis inveni. Differt à S. Asterico foliis petiolatis; à S. trifoliato et ternato caule hispido et foliis alternis; ab omnibus foliis repando-sinuatis.

10. S. RETICULATUM (Pursh fl. bor. am. 2. p. 579, caule . . . , foliis alternis ovato-lanceolatis serratis obtusiusculis villosiusculis.  $\frac{1}{2}$  in Florida. Cæter. ign. Species Nuttalis dubia. †.

\*\*\* *Folius indivisis verticillatis.*

11. S. TRIFOLIATUM (Linn. spec. 1302), caule tereti subhexagono lævi, foliis ternato-verticillatis ovatis dentatis scabris, infer. longè, super. brevissimè petiolatis, petiolis subnudis, paniculâ 3-chotomâ, invol. squamis ovali-oblongis subacutis vix ciliatis, ligulis 15-18.  $\frac{1}{2}$  in montosis Virginie, Carolinæ (Pursh) et Georgiæ (Ell. sketch 2. p. 466). S. ternatum Desf.! cat. h. Par. ed. 3. p. 403 non Retz. — Moris. ox. s. 6. t. 3. f. 68. Achænia glabra latè alata apice emarginata dente callosa intra marginatorem notata. Hæui sub nom. admissio ex hort. Hafn. et ex Lugd. Bat. tempore Royeni collectum. (v. v. c.)

12. S. TERNATUM (Retz in Willd. sp. 3. p. 2333), caule tereti lævi, foliis ternato-verticillatis breviter petiolatis lanceolatis acuminatis serratis scabris, petiolis ciliatis, paniculâ subdichotomâ, invol. squamis ovali-oblongis obtusiusculis ciliatis, ligulis 8-9.  $\frac{1}{2}$  in Amer. bor. (Retz.), montanis Carolinæ et Georgiæ (Ell. sketch 2. p. 467). S. ternifolium Michx. fl. bor. am. 2. p. 146. S. trifoliatum Desf.! hort. Par. ed. 3. p. 182. Hook. bot. mag. t. 3355. non Retz. Linnæus Michaux et plures alii S. ternatum et trifoliatum verisim. confuderunt. S. ternatum præter alias notas differt achænio apice subciliato, aristas 2 obtusas cum aliis concretas gerente. (v. v. c.)

$\beta$ . *atropurpureum* (Nutt. gen. am. 2. p. 183), caule tereti lævi atropurpureo, foliis quaterno-verticillatis sparsive lanceolatis scabris, petiolis ciliatis.  $\frac{1}{2}$  in Amer. bor. Carolinæ et Georgiæ (Pursh fl. 2. p. 575). S. atropurpureum Retz in Willd. sp. 3. p. 2334.

13. S. 2. Tetragona, caule tetragono apice interdium teretiusculo, foliis oppositis, superioribus sæpè connatis, achæniis exaristatis, aristis nempe cum alâ utrinquè omninò concretis.

13. S. FENFOLIATUM (Linn. sp. 1301), caule tetragono lævi, foliis oppositis ovato-deltoides grossè serratis glabriusculis, inf. petiolatis, sup. connato-perfoliatis, invol. squamis latè ovatis obtusis tenuè ciliatis, ext. subquarrosis, ligulis 15-20.  $\frac{1}{2}$  in Amer. bor. à Pennsylvania ad Carolinam in montibus Alleghany (Pursh). Cass. dict. 59. p. 324. Hook. bot. mag. t. 3354. S. tetragonum Mönch. Achænia glaberrima latè alata emarginata exaristata, lobis obtusis, aristis nempe cum alâ omninò et strictissimè sub nervi marginalis specie concretis. (v. v. c.)

$\beta$ . *conjunctum*, invol. squamis ext. 4 expansis et cætera superantibus. S. conjunctum Willd. enum. p. 633 ex Pursh.

14. *S. HORNEMANNI* (Schrad. ! hort. Gœtt. sem. 1829 et in litt. 1832), caule tetragono infernè hispido, foliis oppositis subtus subecabris latè ovatis acutis grossè serratis, infer. petiolatis petiolis aspius alatis, superioribus connato-perfoliatis, invol. squamis ovatis subobtusis tenuè ciliatis, ligulis circ. 20.  $\frac{1}{2}$  in Amer. bor. *S. Asteriscus* Hornem. h. Hafn. ex Schrad. *S. connatum* Desf. hort. Par. cat. ed. 3. p. 182 non Linn. *S. perfoliatum* Poir. suppl. 5. p. 157? *Achnia* glaberrima alata apice acutè biloba, more *S. perfoliati* exaristata. Species media inter *S. perfoliatum* et *S. connatum* et cum utroque aspè confusa. (v. s. comit. à cl Schrad. et v. c.)
15. *S. CONNATUM* (Linn. mant. 574), caule basi tetragono apice tereti hispido-simo, foliis oppositis latè connato-perfoliatis ovato-oblongis serratis subis, invol. squamis ovatis obtusiusculis subsquarrosis, ligulis circ. 12.  $\frac{1}{2}$  in Amer. boreali. *S. scabrum* Moench. Ligulæ 7-8-nerviæ latiores quàm *S. perfoliati*. *Achnia* non vidi. (v. s. cult. in hort. Hafn.)
16. *S. SCABERRIMUM* (Ell. sketch 2. p. 466), caule subangulato, angulis superè scabris, foliis breviter petiolatis interdum connatis ovatis acuminatis serratis rigidis utrinquè scaberrimis, capitulis subcorymbosis, invol. squamis ovatis ciliatis, ligulis 12-14.  $\frac{1}{2}$  in Amer. bor. Georgiâ. *Achnia* orbiculata alata profundè emarginata.
17. *S. INTEGRIFOLIUM* (Michx. fl. bor. am. 2. p. 146), caule tetragono aspero, foliis oppositis sessilibus oblongis integerrimis scabris, capitalis paucis breviter pedunculatis, invol. squamis oblongo-ovatis glabris ciliatis, ligulis circ. 14.  $\frac{1}{2}$  in regione Illinoensium (Michx.), ad montes Alleghany (Pursh), in montanis Carolinæ (Ell. sketch 2. p. 465.)
18. *S. LÆVIGATUM* (Pursh fl. bor. am. 2. p. 578), caule tetragono sulcato glabro simplici, foliis oppositis sessilibus ovatis acuminatis tenuissimè serratis lœvis subcordatis utrinquè glabris, capitulis densè corymbosis, invol. squamis ovatis ciliatis.  $\frac{1}{2}$  in Georgiâ. inter Savannah. et Louis-Ville (Ensl.) et in distr. occident. (Ell. sketch 2. p. 466.)
19. *S. EXTEROCAULON* (Bernh. ex Spreng. syst. 3. p. 630), caule tetragono glabro, foliis oppositis cordato-lanceolatis acutis inæqualiter dentatis scaberrimis, petiolis alatis perfoliatis, invol. squamis oblongis acutiusculis.  $\frac{1}{2}$  in America boreali.

† *Species vix nota.*

*S. TRACOPUS* (Raf. herb. p. 24). — In America boreali.

†† *Species exclusæ.*

*S. SOLIDAGINOIDES* Linn. = *Heliopsis lævis*.

*S. TETRAGONOTHECA* Gært. = *Tetragonotheca helianthoides*.

*S. TRILOBATUM* Linn. = *Wedelia carnea*.

CCLXXXII. *POLYMNIA* Linn. gen. n. 987. Gært. fr. 2. p. 429 t. 174. f. 2. *Lev. syn.* 213. — *Polymnia* et *Alymnia* Neck. elem. 1. p. 31. — *Polymnia* et *Polymniastrum* Lam. ill. t. 711 et 712. Cass. dict. 59. p. 246 et 247.

Capitulum multiflorum monoicum, fl. radii fœmineis ligulatis 1-seriis, disci tubulosis 5-dentatis abortu styli masculis. Invol. duplex. ext. squamis 5 patulis magnis foliaceis ovato-lanceolatis, int. squamis pluribus brevioribus (sed disco longioribus) adpressis achnia amplectentibus. Recept. planum paleaceum. Stylus disci supernè pubescenti apice bifidus. *Achnia* glabra, disci abortiva teretia, radii obovato-crassa subcompressa lævia exalata calva. — Herbar. Americanæ erecta. Folia alterna seu opposita aspera semiamplexicaulia subcorymbosa. Flores lutei, disco sæpè fusco-purpurascente.

Sect. I. UVEDALIA. — *Polymnia Neck. Lam. Cass. ll. cc.*

*Ligulæ planæ elongatæ involucri superantes.*

. *P. UVEDALIA* (Linn. spec. 1303), caule sulcato glabriusculo, foliis oppositis utrinque sparè puberis sinuato-lobatis subtrilobis in petiolum decurrentibus, ligulis 10 lineari-lanceolatis tridentatis.  $\frac{1}{2}$  in pascuis montanis Amer. bor. à Virginiâ ad Carolinam. Michx. fl. 2. p. 147. Ell. sketch 2. p. 471. Mill. dict. n. 1. Lam. ill. t. 711. f. 2. bona. Willd. sp. 3. p. 2335. excl. syn. Cav. — Pluk. t. 83. f. 3. Moris. ox. s. 6. t. 7. f. 55. (v. s.)

. *P. MACULATA* (Cav. ic. 3. p. 14. t. 227), caule sulcato plûs minùs villosò sæptis maculato, foliis oppositis junioribus molliter villosis, inf. deltoideo-cuneatis sinuato-angulatis, supremis rhombeis subrepandis, ligulis circ. 20 lanceolatis tridentatis, invol. squamis exter. latè ovatis.  $\frac{1}{2}$  in Mexico (Cav.), in valle Mexicanâ (Berl.! pl. exs. n. 719), in montibus Aishualtampec (fl. Mexic. ined.). *P. Uvedalia* hort. Paris. 1818 et hort. Lugd. B. tempore Royeni. (v. v. c. et s. sp.)

. *P. ASPERA* (Mart.! herb.), caule teretiusculo scabro apice breviter hirsuto, foliis oppositis suprâ aspero-scabris subtùs molliter velutinis subcanescentibus hastato-trifidis, lobis acuminatis suberoso-dentatis, lateralibus subbifidis ad basin acutè auriculatis, invol. squamis exter. lineari-oblongis disco duplò ferè longioribus.  $\frac{1}{2}$  in Mexico. Enata anno 1829 in hort. Monac. seminibus ab hortul. Kerl. adportatis. Species aff. *P. maculatæ* sed folia basi amplè et acutè auriculata. Ex capitulis junioribus caract. non eruere potui. An *P. scabra* Zaccar. pl. nov. 1. p. 313 eadem? Ex cl. auct. capitula dicuntur discoidea nec radiata! (v. s. in h. Mart.)

. *P. RIPARIA* (H. B. et Kunth nov. gen. am. 4. p. 282), caule compresso fistuloso glabro, foliis oppositis ovatis sinuato-angulatis subcordatis basi in petiolum decurrentibus dentatis hispidulis, ligulis 12-15 elongatis 3-dentatis.  $\frac{1}{2}$  ad ripam fl. Magdalensæ inter Cascajo et Angustura de Carare.

. *P. GLABRATA*, caule tereti subangulato glabro ad apices scabriusculo, foliis oppositis subconnatis ovato-lanceolatis acuminatissimis basi in petiolum alatum decurrentibus subsinuato-dentatis sen integerrimis, adultis glabris, junioribus subasperulis, corymbo composito laxo, invol. squamis exterioribus latè ovalibus.  $\frac{1}{2}$  in Chili legit cl. Hænke. Folia 9 poll. longa, 3 poll. lata. (v. s. in h. Hænke à cl. de Sternberg comm.)

*ß. angustifolia*, foliis oblongo-lanceolatis subintegerrimis.  $\frac{1}{2}$  in Cordilleris Chilensium et fortè Peruvianorum legit Hænke. Folia 4 poll. longa, pollicem lata. (v. s. comm. cum  $\alpha$ .)

Sect. II. ALYMNIA. — *Alymnia Neck. elem. 1. p. 31. — Polymniastrium Lam. ill. t. 712.*

*Ligulæ involucri breviores obovatæ aut lineares.*

. *P. CANADENSIS* (Linn. sp. 1303), caule tereti puberulo-glanduloso, foliis oppositis alternisve ovato-triangularis sublobatis acutis petiolatis sæpè basi auriculatis, infimis sublyratis, invol. squamis ovatis acuminatis ciliatis, ext. paulò majoribus, acheniis obovatis subcompressis, ligulis minutissimis trifidis.  $\frac{1}{2}$  in montanis Americæ borealis à Canadâ ad Virginiam (Pursh), in Carolinâ (Ell. sketch 2. p. 471), in Kentucky (Raf.!), ad saltus Niagaræ (B. Greene!), Linn. smœn. ac. 5. p. 15. t. 1. f. 5. Lam. ill. t. 711. f. 1. Michx. fl. bor. am. 2. p. 147. (v. s.)

. *P. VARIABILIS* (Poir. dict. 5. p. 505), caule tereti puberulo-glanduloso, foliis oppositis ovato-triangularis sublobatis in petiolum cuneato-decurrentibus subpuberis, invol. squamis subæqualibus subacutis dorso puberis eciliatis, ligulis minutis subintegris. — patr. ign. *Polymniastrium* Lam. ill. t. 712. Capitulum

primâ fronte discoides sed varè radiatum ligulis minutissimis. An à primâ certè distincta? (v. s. c. in hort. Par.)

8. *P. SIMONNEA*, caule tereti trichotomo apice puberulo, foliis oppositis sessilibus ovato-deltoides basi latè cuneatis apice acuminatis subdentatis induratis, invol. squamis ext. 5 oblongo-linearibus subpuberulis, int. summo dno pilos apice glandulosos gerentibus, ligulis parvis subintegris. — in Brasiliâ circa Rio-Janeiro legit cl. Gandichaud. Species distinctissima et primâ fronte i genere diversa, sed à prioribus non separanda. Stylus disci minimus subnullus. Recept. post. antihesi elevatum supernè plano-subconcaevum et Reclinatus recept. ferè referens (v. s. comm. à cl. inv.)

9. *P. ATLANTICUS*, caule tereti puberulo-glanduloso, foliis oppositis alternis membranaceis puberulis triplinerviis grossè dentatis apice subtrilobis, basi petiolum alatum irregulariter dentatum attenuatis, prope basin auriculato-dentatis, inv. squamis ext. ovatis acutis dorso villosis, acheniis obovatis compressis, ligulis linearibus integris. — in Brasiliâ prov. Rio-Grande (h. Ma. imp. Bras. n. 8851). Affinis priori sed folia basi dilatata, capitula duplè majora.

† *Species non satis nota aut dubia.*

10. *P. CAROLINIANA* (Poir. diet. 5. p. 505), caule tereti pubescente, foliis alternis indivisis crenatis subtus albidis, invol. squamis ext. majoribus, int. 4 minutibus. — in Carolinâ. †.

†† *Species exclusa.*

- P. AFRICANA* LINN. — *Gnossia oleifera*.  
*P. CARNOSA* LINN. — *Didelta carnosa*.  
*P. FRONDOSA* BRAC. = *Gnossia oleifera*.  
*P. PERFOLIATA* POIR. = *Melampodium perfoliatum*.  
*P. REVOLUTA* QUORUMD. ex Steud. = *Wedelia perfoliata*.  
*P. SPINOSA* LINN. = *Didelta spinosa*.  
*P. TETRAGONOTHECA* LINN. = *Tetragonotheca helianthoides*.  
*P. WEDELIA* LINN. = *Wedelia frutescens*.

CCLXXXIII. *ESPELETIA* Mutis in Humb. et Bonpl. pl. equin. 2. p. 11. t. 797.  
H. B. et Kunth nov. gen. am. 4. p. 280.

Capitulum multiflorum monoicum, fl. radii numerosissimis 1-seriis ligulatis fœmineis, disci tubulosi 5-dentati bisexualibus abortu masculis. Invol. campanulatum imbricatum polyphyllum. Recept. planiusculum paleaceum, paleis membranaceis obtusis. Antheræ erectæ. Stylus radii bifidus gracilis, disci simplex. Achenia obovata subangulata calva, disci verisim. abortiva. — Herbaræ æquinoct. Americane dense lanato-tomentosæ resinosæ perennes. Folia opposita integra. Capitula ramosa terminantia corymbosa lutea. — Genus caract. videtur affinis *Silphio* sed habitus diversus. Antheræ ex icone breviter caudatæ, sed de hoc caract. cum habitu discrepante silent descriptiones.

1. *E. GRANDIFLORA* (H. et Bonpl. l. c. t. 70), rufescenti-lanata, foliis radicalibus lanceolatis subtus lineatis, corymbo simplici laxo. ♀ in Amer. locis subfrigidis prope Santa-Fé de Bogota et in Andibus Quinduanensibus Bogotensibus dicitur *Fraxylejon*. Fundit resinam *tremenithina* dictam. Achenia omnino glabra. Ligulæ circ. 60 lineares apice 3-dentatæ.  
2. *E. ARGENTEA* (H. et Bonpl. l. c. t. 71), caule densissime argenteo-piloso, foliis supra strigoso-strigosis subtus molliter tomentosis, corymbo ramoso subfastigiato. ♀ in Novæ-Granatæ frigidis juxta urbem Zypaquira. Capitula dicuntur radiata, sed ligulæ indescriptæ nec in icone obviæ. Stylus radii et



basin more insolito moniliformi-incrassatus. Achænia omninò glabra. Cor. ad tubi apicem pilosiusculæ.

3. *E. corymbosa* (H. et Bonpl. l. c. p. 16. t. 72), incano-tomentosa, foliis coriaceis, radicalibus supernè lato-lanceolatis subtùs venosis tomentosis, capitulis subcorymbosis confertis. 2½ in frigidis Andium Novo-Granatensium juxta Almaguer. Esp. rigida H. et B. pl. equin. t. 72. Ligulæ circ. 20 invol. paulò longiores. Cor. in tubo glabræ. Achænium apice barbatum.

## CCLXXXIV. BERLANDIERA.

Capitulum multiflorum monoicum, fl. radii 1-serialibus ligulatis femineis ad axillas squamarum invol. interiorum sitis, fl. disci masculis tubulosis 5-dentatis ad axillas palearum recept. exteriorum sitis, centralibus abortivis. Recept. turbinatum obconicum supernè truncatum. Invol. 3-seriale foliaceum, squamis ext. circ. 3 oblongis, mediis obovatis majoribus acutiusculis circ. 4-5, intimis maximis subrhombicis subdentatis circ. 6-8. Paleæ recept. foliæ, exteriores binæ floriferae ad basin ejusque squamæ recept. intimæ ortæ cucullatæ apice dilatata obtusæ, aliæ ex margine et ex disco terminali recept. ortæ minores cuculliformes florem ad filum reductum aut nullum gerentes. Achænia radii obcompressa obovato-orbiculata exalata coriacea pappo brevissimo subaristato coronata. Cotyledones subcarnosæ invol. squamis parallelæ. — Suffrutex Texano-Mexicanus. Rami teretes hirsuti. Folia alterna sessilia cordata ovato-sublanceolata crenata, suprâ strigilloso-villosa, subtùs canescenti-villosa. Capitula longè pedicellata ad apices ramorum 3-5 corymbosa. Invol. et recept. paleæ pubescentes. Flores lutei. — Aff. videtur hinc Melampodio, illinc Polymniæ. — Genus dixi in honorem cl. Berlandier qui Mexicanas plagas peragravit et plantas plurimas his regionibus proprias collegit.

2. *B. TEXANA*. 3 in sylvis Mexicanæ provincie Texas legit cl. Berlandier. (Pl. Mex. D. 2044.) (V. 2.)

**Div. IV. MELAMPODIEÆ.** — Asteroideæ Melampodieæ veræ. Less. syn. 217.

Capitula monoica multiflora, fl. femineis pluribus ligulatis. Achænia corticata (1) calva. Recept. paleaceum, paleis plùs minùs cum achæniis concretis. — Summi hanc et priorem divisionem sub nom. *Helianthæ Milleriæ* Melampodijunxit. (Opusc. 3. p. 60.)

CCLXXXV. MELAMPODIUM Linn. gen. n. 989. Brown trans. hort. soc. 12. p. 104. H. B. et Kunth nov. gen. am. 4. p. 272. Less. syn. 217. — Melampodium, Dysodium et Alcina Pers. — Melampodium, Zarabella et Alcina Cass.

Capitulum multiflorum monoicum, fl. radii 5-10 uniseriatis femineis ligulatis, disci tubulosis masculis. Invol. duplex, squamis ext. rariùs 3, planis patulis, int. tot quot flores radii et eos involventibus. Recept. convexo- aut subulato-conicum paleaceum, paleis diaphanodiscidis. Achænia disci abortiva, radii obovata subcurvata lævia intra squamas inv. interiores plùs minùs inclusa et ideò sæpiùs

(1) Corticata dixi achænia squamis inv. paleisve recept. involuta et cum iis adœquata ut massa unica ex achænio et squamâ coadunatis evadat. Sed involuta dixi a squamis paleisve sunt inclusa nec concreta. Ergò achænia *Chrysogoni* et plur. eorum generum à cl. Lessingio corticata dicta hoc nomine non designavi.

externè tuberculosa aut rugosa et apice supè coronata. — Herba ut suffruticuli Americani. Caules dichotomi. Folia opposita. Pedicelli inter dichotomias orti 1-cephali. Flores flavi, ligulis in unica spec. albis.

Sect. 1. EUMELAMPIDIUM. — *Melampodium* Linn. l. c. Gartn. fr. 2. p. 436. t. 169. Lag. nov. gen. et sp. p. 32. Cass. dict. 59. p. 237.

Invol. squamis interiores achenia involventes, extus tuberculosas cum mucato nervoso-costatis, superius productas in cucullum nunc truncatam nunc aristam uncinatam desinentem.

1. *M. diffusum* (Cass. dict. 59 (1829). p. 238), caule erecto-diffuso di-trichotomo, foliis oblongis utrinque acutis integerrimis supra pubescentibus subitè longe sericeo-villosis, pedunculis è dichotomiis ortis 1-cephalis, invol. squamis 5, ext. obovato-eminentis mucronatis achenia involv., basi variegatis seriem tuberculosam ad latum internum gerentibus, cucullatis in acumen uncinatum productis. ① in insulâ Luzoniâ ad Manillam (Champ.). *M. Manillense* Less. in Linnæ 1830. p. 155. t. 2. f. G. (Achenia). *Alia M. Americano*. Discus oblongus. Folia ex Less. interdum variant pinnatis (v. s.)
2. *M. strictum* (Lag. elench. h. madr. 1805. nov. gen. (1816) p. 32. n. 464 in Kunth), caule erecto diffuso di-trichotomo, foliis supra acutulis subitè sericeo-villosis, infer. trifidis, sup. integris linear-lanceolatis, capitatis è dichotomiis et apicibus breviter pedicellatis, invol. squamis ext. oblongis villosissimis, int. achenia involv. undique muricatis apice cucullatis in acumen extensum productis. ① in Mexico (Lag.) prov. Leonine ad occid. m. G. Guanaxtato (Méndez). *M. diffuso* valde affinis, sed pedicellis viz invol. longius bracteis foliaceis binis angustis sub involucri et acheniis valde muricatis differt. (v. s.)
3. *M. ramosissimum*, caule suffruticuloso ramosissimo glabro, foliis linearibus subadpressè pubescentibus, aliis integerrimis, aliis hinc indè obtusè dentatolobatis sinuatis, pedicellis folio longioribus 1-cephalis, invol. squamis ext. ovalibus extus pubescentibus, inter. achenia involventibus basi tuberculosis apice expansis in cucullum latum dorso in acumen uncinatum desinentem. ① in Mexici prov. Texas legit cl. Berlandier (pl. exs. n. 2017). Ligulæ oblongæ lineares parvæ. (v. s.)
4. *M. linearilobum*, caule erecto diffusè ramoso, foliis pinnatipartitis lobis utrinque 2-4 linearibus supra strigoso-pilosis subitè sericeo-villosis, nunc paucis integerrimis linearibus, pedicellis elongatis 1-cephalis, invol. squamis ext. latè ovalibus basi villosissimis apice membranâ marginatis inter. achenia involv. basi rugoso-tuberculatis apice cucullatis, cucullo margine intus revolutò dorso brevissimè apiculato. ① in Mexico ad Guilo-tepec et Las Bajas. Ligulæ latè obovatæ. Cucullus interdum exapiculatus interdum apicalo recurvo. Facile lobis fol. linearibus ab icone Houston. typo *M. Americano* differt. (v. s. comm. à cl. Alaman.)
5. *M. Americanum* (Linn. sp. 1303), caule procumbente, foliis integris seu lobis lato-lanceolatis utrinque 2-3-fidis supra strigoso-hispidis subitè junioribus sericeo-hirtis, pedicellis elongatis hispidis 1-cephalis, ligulis 10 oblongis invol. duplò longioribus integris, invol. squamis ext. latè ovatis, inter. achenia obtegentibus costatis auriculato-tuberculatis, apice cucullatis, cucullo ext. aspero marginibus inflexo exappendiculato. — in Mexico ad Vera-Cruz (Houston reliq. p. g. t. 21) et ad Santa-Fé in savanis (Less. in Linnæ 1830. p. 144 Gartn. fr. 2. p. 436. t. 169. f. 8.
6. *M. cinereum*, caule diffuso villosò ramoso, foliis oblongo-linearibus obtusè utrinque cinereo-pubescentibus paucis integris plerisque obtusè sinuato-repandis, pedicellis 1-cephalis folio paulò brevioribus, ligulis invol. longioribus apice emarginatis, invol. squamis ext. ovatis subacutis dorso villosis, int. achenia invol. basi muricato-tuberculatis apice cucullatis, cucullo ample

ariolato extus lævi, margine integerrimo intus revoluto. (1) in Mexico ad San-Fernando oct. fl. legit Berlandier (pl. exs. n. 2243). Flores ex eo albi, in sicco albidii. (v. s.)

7. *M. KUNTIIANUM*, caule fruticoso, ramis junioribus villosa-pubescentibus, foliis lanceolato-linearibus integerrimis supra sericeo-pubescentibus, subtus lanuginoso-sericeis argenteis, pedicellis 1-cephalis folio longioribus, invol. squamis ext. obovato-cuneatis acutis extus sericeis, int. achænia involv. basi costatis muricato-tuberculatis apice cucullatis, cucullo vix apiculato marginibus vix inflexis. 5 in Mexico inter, ad Tehuillotepec et Tasco. Mel. sericetum H. B. et Kunth nov. gen. am. (1820), p. 272. t. 398 non Lag. Flores aurantisco-flavi; ligulæ invol. triplò longiores. Capitulum 8-9 lin. diam. Caules et pedunculi ex icone hispiduli (v. s. sine fl.)
8. *M. LANCEOLATUM*, caule fruticuloso basi glabro ad apices adpressè villosa, foliis lanceolatis acuminatis integerrimis supra parcè strigillosis demùm glabris subtus sericeo-villosis argenteis, pedicellis 1-cephalis vix folio longioribus, invol. squamis ovatis dorso villosis, int. achænia involv. nervuloso-costatis et tuberculatis apice cucullo exaristato instructis. 5 in itiner. suis legit Née sed locus ignotus. An Manillense? an potiùs Americanum? Species facilè distincta. Capitula 4 lin. diam. Discus elevatus ut in *Spilantho*. Fl. flavi. Ligulæ inv. duplò longiores ovals. (v. s. ex herb. Thib.)

Sect. II. *ZARABELLIA*. — *Dysodium* Rich. in *Pers. ench.* 2. p. 489 non Cav. — *Zarabellia* Cass. dict. 59. p. 240 non Neck.

Invol. squamæ inter. achænia radii involventes, rugosæ aut tuberculatæ, supernè aut truncatæ aut productæ in mucrones dentesve 1-3 anticè subhiantes et achæniæ apicem denudatum monstrantes. — Sect. inter priorem et tertiam media, ad *Eumelampodium* M. heterophyllo, ad *Alcinam* M. divaricato ex habitu accedens.

9. *M. HETEROPHYLLUM* (Lag. nov. gen. et sp. p. 33. n. 405), caule fruticuloso erecto piloso, foliis lanceolatis in iisdem aut diversis individuis indivisiis trifidisque, supra puberulo-scabridis, subtus sericeo-villosis argenteis, pedicellis 1-cephalis folio longioribus, invol. squamis exter. ovato-rhombeis dorso villosis, int. achænia involv. nervuloso-subcostatis parcè tuberculatis in mucrones 1-3 minutos subproductis. 5 in Mexico (Lag.), in pratis Papantlæ (Schied.), ad colles prope Tantoyma et Tampico de Tamaulipas (Berl. pl. exs. n. 2161 et 1281). Less. in linnæa 1831. p. 407. Herba ferè *M. sericei* seu lanceolati sed cucullus ad apicem achæniæ nullus. (v. s.)

10. *M. ANGUSTIFOLIUM*, caule fruticuloso erecto puberulo-hirtello, foliis linearibus integerrimis acutis supra puberulo-hirtellis subtus sericeo-villosis argenteis, pedicellis folio duplò longioribus, invol. squamis ext. ovatis acutis dorso villosis, inter. achænia involv. apice truncatis vix mucronatis. 5 in Peruviae montanis, ad Oronocum (Hænk. herb.). Hæc et præcedens species habitu sect. primæ optimè exhibent sed cucullus nullus. Fortè caract. paulò immutato ad *Eumelampodium* referendæ. (v. s. in h. Hænke à cl. Sternberg miss.)

11. *M. GRACILE* (Less. in linnæa 1831. p. 407), caule herbaceo erecto trichotomo, foliis semiaimplexicanibus supra asperimis junioribusque strigosohispidis subtrès asperis pallidioribus et glabriscaulis elliptico- seu obovato-rhombeis integerrimis seu hastato-panduratis posticè sinuato-dentatis, sup. triangularibus integerrimis acuminatis basi obtusissimis, invol. squamis ext. 3. ligulis 5, fructibus trapezoideo-obovatis apice nudis glabris laevibus obsolete reticulato-venosis. (1) in pratis Papantlæ Mexicanorum. Simile *M. paludoso* sed multò tenerius.

12. *M. OBLONGIFOLIUM*, caule herbaceo erecto di- et trichotomo sparè puberulo-hirtello, foliis sessilibus oblongis acuminatis integerrimis aut basi interfûm subdilatato-denticulatis utrinquè sparè et rariter strigillosis, pedicellis folio longioribus, invol. squamis ext. 3 lanceolatis, ligulis 5 bifidis, fructibus obo-

- vatis apice nudis dorso reticulato-rugulosis. (1) in Mexico ad Tantoyuca legit Berlandier (pl. exs. n. 2153). Capitula parva lutea. Affine *M. gracili*, sed foliis nec rhombea nec formis varia nec ullâ basi obtusissima. (v. s.)
13. *M. MICROCEPHALUM* (Less. in *linnæa* 1834. p. 268), herbaceum gracile hirsutiusculum, foliis oblongo-rhomboides basi perangustâ sessilibus, pedicellis folio longioribus, invol. squamis ext. 5 curvilineo-acuminatis, achæniis curvato-obpyramidalis apice truncatis et nudis dorso tuberculatis. (1) in Mexici culis prope Hacienda de la Laguna. Proximum dicitur *M. divaricatum*.
14. *M. HISPIDUM* (H. B. et Kunth nov. gen. am. 4. p. 273 t. 399), herbaceum erectum, ramis oppositis hispidis, foliis sessilibus subconnatis linearilanceolatis integerrimis suprâ hispido-scabris subtus pubescentibus, pedicellis folio duplò triplòve brevioribus, invol. squamis ext. 5 ellipticis obtusis dorso villosis, int. achænia involv. costatis spinuloso-tuberculatis apice truncatis, ligulis bifidis brevibus. (1) in Mexico inter Pascuaro et Valladolid de Mechoacan. Ligula flavidæ.
15. *M. LONGIFOLIUM* (Cav. hort. madr.), caule herbaceo erecto sparsè puberulo-hirtello, foliis oblongis subrhombicis semiamplexicaulibus obtusè cordato-auritis apice subacutis integerrimis utrinquè sparsè puberulis, pedicellis folio brevioribus, invol. squamis ext. ovato-lanceolatis subacuminatis dorso puberis, int. achænia involv. compressis rugosis apice muticis aut dorso brevissimè mucronatis. (1) in Mexico (Cav.). Willd. enum. p. 934. Brouss. hort. mœss. 1806. (v. v. c.)
16. *M. RHOMBOIDEUM*, caule herbaceo erecto bifariam piloso, foliis ovato-oblongis basi aurito-semiamplexicaulibus à medio ad basin angustatis apice subacutis vix dentatis utrinquè puberulis, pedicellis è dichotomiis ortis folio brevioribus gracilibus, invol. squamis ext. 5 lanceolatis acuminatis, inter. achænia involv. rugosis dorso ad apicem in spinam erectam productis. (1) in Amer. calid. ad locum ign. Zarabellia rhomboidea Cass. dict. 59. p. 240. Dysodium divaricatum hort. par. 1819! An *M. Berterianum* Spreng. syst. 3. p. 619? sed inter plantas Berterianas non reperio. Affine *M. longifolio*, sed differt caulibus bifariam nec sparsè pilosis fructibusque longè spinosis. (v. v. c.)
17. *M. DIVARICATUM*, caule herbaceo dichotomo bifariam piloso, foliis ovatis acuminatis basi cuneatis grossè dentatis suprâ scabris subtus hispidulis, pedicellis è dichotomiis ortis glabresculis folio sæpius longioribus, invol. squamis ext. 5 obovatis obtusis basi subconcretis, int. achænia obtegentibus compressis rugosis apice truncatis extus subdentatis, ligulis brevibus 3-dentatis. (1) in Amer. paludosis ad sinum Darien (H. et B.), circa Sanctam-Martham (Rich. Bert.), in Mexico (Lag. Karv.). Dysodium divaricatum Rich. (1807), in Pen. ench. 2. p. 489. Colla in Ac. Taur. 1827. p. 323 cum icon. Melampodium peludosum H. B. et Kunth nov. gen. am. (1830) 4. p. 215. Less. in *linnæa* 1830. p. 149. Alcina ovatifolia Lag. nov. gen. et sp. (1816) n. 403. Jacq. f. eclog. 1. p. 115. t. 78. Melamp. ovatifolium Reich. exot. (1827) p. 30. t. 42, sed oculis undiquè pilosis! Wedelia ovatifolia Willd. enum. suppl. 61. Wedelia minor Hornem. Hafn. hort. p. 835. Alcina minor Cass. dict. 59. (1829) p. 243. Dysodium radiatum Desf. cat. h. par. 1830. p. 182. (v. v. c. et sp.)

Sect. III. ALCINA. — Alcina Cav. ic. p. 10. t. 15. Cass. dict. 59. p. 242. — Wedeliæ sp. Willd. — Alcina Steud. — Camutia Bonst.

Invol. squamæ int. achænia radii involventes ovali-oblongæ laeves apice tubercula minima 2-4 gerentes et inter eos foramine minimo vix apertæ. Achænium striatum.

18. *M. PERFOLIATUM* (H. B. et Kunth nov. gen. am. 4. p. 274), herbaceum erectum dichotomum glaberrimum, foliis basi appendiculatis in paxidem connatis in petiolum alatum angustatis latè ovato-rhombicis triplinerviis subdentatis, pedicellis è dichotomiis ortis folio paulò brevioribus, invol. squamis ext. ovali-oblongis capitulo duplò longioribus, int. achænia involv. ovatis laevibus

apice summo tubercula 3-5 minima gerentibus. ① in Mexico (Cav. Alam.!), inter Valladolid et Pasouaro (H. B), in valle Tolucañā (Berl.!) pl. exs. n. 1134). *Aleina perfoliata* Cav. ic. 1. p. 10. t. 15. Lag. nov. gen. p. 32. *Wedelia perfoliata* Willd. sp. 3. p. 2335. *Polymnia perfoliata* Poir. dict. 5. p. 506. *Melamp. connatam* fl. mex. ic. ined. *Camutia perfoliata* Bonat. ex Steud. (v. s.)

† *Species non satis notæ.*

19. *M.?* *DOMBEYANUM*, caule herbaceo erecto basi glabrato medio sparse apice conferte hirsuto, foliis sessilibus basi breviter attenuatis ovatis vix acutis grosse dentatis glabriusculis, junioribus subtus villosis et glandulosis, capitulis in dichotomiis summis subsessilibus, invol. squamis ext. ovali-oblongis obtusiusculis..... ①? in Amer. austr. legit Dombey sed locus propr. ign. Ob capitula nimia juniora caract. gener. non eruere quivi. Habitus ferè *M. paludos.* (v. s. comm. à Mus. reg. Par.)
20. *M.?* *RUDERALE* (Swartz fl. ind. occ. 1372), caule herbaceo erecto tetragono pubescente, foliis petiolatis ovato-lanceolatis acutis serratis trinerviis hispidis, pedicellis axillaribus petiolo brevioribus, capitulis subnatis discoidis. ① in ruderatis Jamaica australioris. A genere videtur removenda. Inv. squamæ 5 oblongæ erectæ. Fl. 10-11 hermaphr., exteriores tantum fertiles. Fructus ovato-oblongi tetragoni, ad angulos obtusè muricati, calyculo minuto cylindrico superati. Paleæ acuminatæ flosculos cingentes. (Sw.)
21. *M.?* *HIDALGOA*, caule volubili, foliis oppositis longè petiolatis ternatisectis, segmentis ovatis crassiusculis glabris serratis, invol. biseriali, ligulis 5, achænio drupaceo ovato, initio apice auriculato, maturo forcipato. — in densissimarum sylvarum marginibus arborescens scandens prope flumen Blanco non longè à San-José del Corral Mexicanorum. *Hidalgoa ternata* Lallav. et Lex. nov. veg. mex. 1. p. 15 non Less. Ligulæ coccineo-aurantiæ. Discus flavus. Conf. *Hidalgoa* Lessingii.

†† *Species exclusæ.*

- M. AUSTRALE* Linn. = *Acanthospermum xanthioides*.  
*M. BARANQUILLÆ* Spreng. = *Sclerocarpus Africanus*.  
*M. HUMILE* Sw. = *Acanthospermum humile*.

CCLXXXVI. *ACANTHOSPERMUM* Schrank pl. rar. hort. monac. v. 2. (1819) t. 53. — *Centrospermum* Kunth nov. gen. am. (1820) p. 271. t. 397 (non Spreng.). Cass. dict. 59. p. 235 et 244 (excl. syn. Spreng.). Less. syn. 217. — *Echinodium* Poir. mss. ex Cass. — *Melampodii* sp. Læft. Linn. Brown.

Capitulum pluriflorum heterogamum, fl. radii 1-serialibus foemineis ligulato-cucullatis, disci masculis tubulosis 5-dentatis. Invol. 1-seriale, squamis 5 ellipticis concavis. Recept. planum paleaceum, paleis exter. ovaria involventibus extus aculeis herbaceis uncinatis onustis, inter. concavis apice dentatis fl. masculos foventibus. Antheræ ecaudatæ appendiculatæ. Styli rami recurvati apice glabri. Achænia paleis aculeatis erectè involuta compressa obtusa basi subattenuata epapposa. — Herbae Americanæ ramosæ demissæ aut diffusæ trichotomæ. Folia opposita subtus punctata. Capitula solitaria ad apices aut dichotomias ramorum per anthesin parva. Cor. luteæ.

☞

Sect. I. XANTHIODES.

Paleæ exteriores seu achæniorum tegumenta longitudinaliter striato-sulcata, nervulos aculeos herbaceos uncinatos gerentia, apice obtusa nec in cornua producta, fructus *Euxanthii* referentia.

1. *A. XANTHIODES*, caule diffuso, ramis pubescentibus, foliis petiolatis subintegris aut dentatis glabriusculis, fructibus apice inermibus. ① in Amer. austr.

7. *X. PARVIFOLIUM*, spinis solitariis simplicibus, foliis oblongis subintegerrimis mucrone spinuloso terminatis utrinque glabrescentibus. (L.)? Patr. ign. varia. Amer. austr. Spinæ crassæ 3-4 lin. longæ. Folia 4-5 lin. longa, 1  $\frac{1}{2}$ -2 lin. lata (v. s. commun. ab am. Delessert.)

† *Species vix nota.*

8. *X.?* *ROMOTHALAMUM* (Spreng. neu. entd. 1. p. 259), caule inermi, involucri androgynis pentaphyllis, foliis obovato-subrotundis. — in Brasiliâ. Speciei genere videtur aliena, et postea ab auctore ipso omissa! †.

†† *Species exclusa.*

*X. FRUTICOSUM* Linn. = *Franseria artemisioides*.

CCLXXXVIII. *FRANSERIA* Cav. ic. 2. p. 78. Willd. sp. 4. p. 378. Cass. dict. 17. p. 364. Less. syn. 219.

Capitula monoica homogama. ♂ Invol. hemisphæricum 8-12-dentatum seu squamis 8-12 uniserial. concretis. Recept. paleis filiformibus inter flores. Flores 15-20. Cor. 4-fida. Stam. immo corolla inserta, vix huic adnata. Antheræ liberæ. Pistilli rudimentum. ♀ Invol. ovatum extus aculeos uncinatos aut spinescentes gerens et in cornum cum 3-4 apice desinens, 1-florum aut in loculamenta 3-4 uniflora divisum. Cor. nulla aut stylo adpressissima. Anth. o. Stylus filiformis intra coram productus bifidus. Stigm. apice obtusa. Achænium calvum in invol. nidulans. — Suffrutices aut herbæ in Americâ crescentes. Folia altera grossè dentato-lobata uni- aut bipinnatisecta. Capitula spicata, macula superiora, feminea inferiora sæpius aggregato-spiculata. — Genus medium iter Xanthium et Ambrosiam.

#### Sect. I. XANTHIOPSIS.

Invol. frutigeræ aculeis mollibus uncinatis hispida.

1. *F. ARTEMISIOIDES* (Willd. sp. 4. p. 374. hort. berol. 1. t. 2), fruticosa, foliis bipinnatifido-dentatis, petiolis alatis, limbo suprâ virescente subitè albepubescente. ♂ in Peruvîâ. Xanthium fruticosum Linn. f. suppl. 418. Ambrosia arborescens Lam. dict. 1. p. 127. Invol. masc. circ. 12-dentatum. fem. et Willd. biflorum. (v. s.)
2. *F. AMBROSIOIDES* (Cav. ic. 2. p. 79. t. 200. excl. syn.), fruticosa, foliis oblongo-lanceolatis dentatis basi cordatis truncatis inæqualibus utrinque hirsutis viridibus, petiolo sæpè appendiculato cæterum aptero. ♂ in Mexico. fl. mex. ic. ined. Invol. masc. 8-fidum. Cor. more Compositarum 5-nervis! lvs. femineum intus 3-4-los. 3-4-florum. (v. v. c. in hort. Monsp.)

#### Sect. II. ACANTHOLÆNA.

Invol. frutigeræ aculeis spinescentibus echinata.

3. *F. CHAMISSONIS* (Less. in Linnæâ 1831. p. 507), subherbacea flexuoso-decumbens, foliis rotundo-ellipticis utrinque hirsuto-incanis longè petiolatis. — in Californiâ. Huc cl. Lessing. refert var. duas diversissimas in posterum fortè separandas.

*a. malvæfolia*, basi lignosâ, foliis integris plicatis crenatis hirsuto-subarveis.

*β. bipinnatisecta*, canescens, caule herbaceo inani, foliis bipinnatisectis. segmentis divergentibus oppositis, segmentulis minutis obtusis. (v. s. olim specim. in h. Par. culta ex sem. Californicis à cl. Collignon lectum et sp. sicc. ex Dougl. itin. ab hon. soc. hort. Lond. comm.)

**OCLXXXIX. AMEROSIA** Tourn. inst. t. 252. Linn. gen. n. 1057. Gærtn. fr. 2. p. 417. t. 164. Cass. dict. 25. p. 203. Less. syn. 219.

Capitula in eadem plantâ alia feminea ad basin apicis, alia mascula ad apicem, omnia homogama. ♂ Capitula pluriflora. Invol. squamis universalibus ferè omninò in cupulam concretis. Recept. epalesceum. Cor. tubulosa brevis. Stam. corollæ non adhærentia. ♀ Capitula plura 1-flora aggregata, invol. communi cincta, bracteolis interpositis instructa. Corolla nulla. Styli rami elongati è collo involuclli exserti. Achænium ovatum intrâ invol. persistens incurvatum et sæpius in dentes aut cornua excrescens. — Herbæ interdum suffruticosæ, aliæ gerontogæ, aliæ Americane. Folia infer. opposita, super. alterna, pinnatilobata trilobata aut indivisa. Rami desinentes in apicem basi compositam foemineam, apice masculam simplicem. — Genus quoad characteres genericos difficile intelligendum et quoad specierum discrimina, fructibus sæpius indescriptis, non satis elaboratum.

\* *Folii uni- aut sæpius bipinnatifidis.*

1. **A. MARITIMA** (Linn. sp. 1481), tota villosissima cinereo-canescens, caule divaricato villosulo-hispido, foliis bipinnatipartitis, lobis apice obtusis hinc inde utrinquè 1-dentatis, invol. masculo breviter hispido campanulato 15-20-floro, form. fructifero ovato cornubus 4-5 rectis echinato. (X) in herbâs ad ripas maris et fluviorum in Hispaniâ circa Taragonam (Delaroches!), ad Niceam (All.), Venetiam (Moric.), in Hetruriâ, agro Neapolitano (Ten.), Cretâ (Oliv.), Egypto (Coqneb. ! Bové !), Semnaar (Acerb. !), Cappadociâ (Linn.), etc. Schkuhr handb. t. 292, sed cornua inv. fruct. omnia. A. villosissima Forsk. descr. eg. p. 160. Dod. pempt. 35. 10. A. artemisifolia et A. elatior Bess. prim. 21. fl. gal. 2. p. 389 et 390 mihi ignota, videntur ex descript. merè A. maritimæ variet. ? (v. s.)

2. **A. SENEGALENSIS**, tota cinereo-pubescente, summitatibus villosis, caule ramoso subvillosulo, foliis bipinnatipartitis, lobis acutis dentatis inter dentes suberrulatis, invol. masculo campanulatis breviter hispido 15-20-floris, form. ovatis cornubus 4-5 rectis echinatis. (X) secus ripas fl. Senegal locis humidis argillosis legit. cl. Baole et Perrottet. An A. maritima var. ? sed pubes multò minor et lobis fol. longiores acuti denticulati discrimina in genere adeò intricato fortè constituunt. (v. s.)

3. **A. CRITHMIFOLIA**, caule basi suffruticoseo supernè confertè velutino, foliis oppositis petiolatis crassiusculis bipinnatifidis, lobis angustis obtusis utrinquè pube adpressâ puberis, apicis paucis, terminali multò longiore, fructibus junioribus villosis inermibus. 2/ 3 in ins. Cubâ circa Havanam legit cl. Ferrero. Amer. maritima Ferr. in litt. 1821. Inter Americanas magis eadem quàm alteram species A. maritimam refert ? (v. s.)

4. **A. ABROTANIFOLIA** (Wall. ! cat n. 3234. comp. 344), minutissimè pubescens, caule basi suffruticoseo ramosissimo, foliis pinnatifidis basi cuneatis petiolatis, lobis oblongis acutis hinc inde dentatis subtrifidis, summis linearibus integerrimis, racemis brevibus foliosis, invol. mass. viz puberulo 5-7-floro, form. ovato viz corniculato. 3/ culta in h. bot. cal. et ex Wall. viz non Indica. Eandem sine loci design. video inter plantas Lundii Brasilienses, et circa Bahiam à cl. Blanchet (pl. exs. n. 972) lectam. Facile ab omnibus differt capitulis masculis minimis. (v. s.)

5. **A. FRUTICOSA**, caule fruticoso erecto ramoso, foliis bipinnatipartitis, lobis linearibus sublancoatis acutis integris aut lobulis hinc inde incisè suprâ glaberrimis subtrifidis adpressè pubescentibus, racemis terminalibus elongatis inter capitula mascula obrectatis, fructibus echinato-spinosis. 3/ in locis umbrosis Mexici in distr. Tamulipas prope Reynosa april. flor. legit cl. Berlandier (pl. exs. n. 2112 !). Variat caule ramisque setis strigillois sparsè hispido aut viz

hispidula subpubescens, capitulis masculis minoribus aut paulò majoribus. (v. s.)

β. *intermedia*, caule suffruticoso supernè hispido, foliis bipinnatifidis lobis acutis, racemis elongatis confertis. ♂ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 869!). Media inter *A. fruticosam* et *A. crinitifoliam*, priori caule et foliis, posteriori inflorescentiâ similis. An species propria? (v. s. in h. Mus. reg. Par.)

6. *A. ELATIOR* (Linn. sp. 1401), caule virgato vix pubescente, foliis bipinnatifidis glabresculis, petiolis ciliatis, racemis paniculatis, lateralibus abbreviatis, fructibus apice 6-cornutis. (1) in pascuis fertilibus à Canadâ ad Carolinam (Pursh), in Georgiâ circa Savannah (h. Duby!). Willd. sp. 4. p. 376. Ell. sketch 2. p. 477. Affinis *A. artemisiæfolia*, sed foliis caulibusque glabresculis, racemis lateralibus caule brevioribus et petiolis longè ciliatis differt. (v. s.)

7. *A. POLYSTACHYA*, tota pube confertâ velutina, caule sulcato, foliis bipinnatifidis lobis elongatis inciso-dentatis acuminatis, racemis numerosissimis in paniculam amplam dispositis, fructibus inermibus. — in Brasiliæ prov. Sancti-Pauli locis cultis humidis nov. et dec. flor. legit cl. Lund; etiâ video in h. Mus. imp. Bras. sub n. 460! Affinis hinc *Franseriæ artemisioidi*, illinc *Ambrosiæ artemisiæfoliæ*; à priore diversa caule herbaceo, petiolis teretibus nec albis et caract. genericis; à posteriore caule sulcato multò elatiore, foliorum lobis multò magis acuminatis, rachi inter lobos lobulatâ, racemis numerosis, fructibus inermi, etc. An *A. maritima* Arrab. fl. flum. 10. t. 26 non Linn.? (v. s. com. à cl. Lund.)

8. *A. ARTEMISIEFOLIA* (Linn. sp. 1401), caule tereti pubescente, foliis bipinnatifidis subcanescentibus, summis pinnatifidis, ramis fastigiatis, racemis terminalibus ternis, fructibus spinis brevibus acutis. (1) in agris Amer. bor. à Canadâ ad Carolinam Ell. sketch 2. p. 477. Lam. ill. t. 765. f. 1. *A. abrotanifolia* Michx. fl. bor. am. 2. p. 183. Fortè etiâ hab. in Mexico (fl. mex. in ind. circa urbem (Berl.!) pl. exs. n. 573) et ad Havanam (Ferr.!), sed ex similitudine speciem dubius hæreo. — Hern. mex. p. 446. f. 1? (v. s.)

9. *A. PAILOSTACHYA*, caule herbaceo erecto tereti paniculato-ramoso foliisque scabridis, foliis subsessilibus pinnatipartitis lobis lineari-lanceolatis acuminatis, hinc indè pinnatim incisim, racemis plurimis gracilibus inter capitula mascula remota ebracteatis, fructibus infra apicem tuberculis obtusis 5-6 ornatis apice conico elongato. — in Mexico inter San-Fernando et Matamoros legit cl. Berlandier (pl. exs. n. 2280!). Fructus plurimi parvi in invol. sciss. Capitula masc. pubero-scabrida parva. (v. s.)

10. *A. CONFERTIFLORA*, caule herbaceo erecto sulcato paniculato-ramoso, foliis petiolatis pinnatisectis suprâ glabresculis subtus in petiolo puberulo-scabris segmentis pinnatipartitis linearibus elongatis integris aut subincisis, racemis plurimis elongatis confertifloris inter. capitula masc. ebracteatis. — in Mexico ad Matamoros legit cl. Berlandier (pl. exs. n. 2297!). Affinis *A. pailestachya*, sed foliis petiolatis pinnatisectis et capitulis masc. confertis diversa. Fructus ex statu juniore non satis noti. (v. s.)

11. *A. ACANTHOCARPA* (Hook. fl. bor. am. 1. p. 309), valdè ramosa, setis asperis rigidis hispidis, foliis bipinnatifidis lobis linearibus, racemis terminalibus, fructibus spinis rigidis subulatis longis patentibus undiqñ tectis. — in Amer. bor. ad ripas fluminum secus Saskatchewan, Red-River, Columbia.

12. *A. PERUVIANA* (Willd. sp. 4. p. 377), caule erecto virgato glabresculo, foliis infer. bipinnatifidis, interm. pinatifidis, summis oblongo lanceolatis indivisis, racemis terminalibus solitariis. 22 in Peru (Poepp.!) pl. exs. n. 17. ad sinum Chorillo (Macrae!), in Mexico (Alam.!), ad Real del Monte (Hauke!). An stirps Mexicana omninò eadem ac Peruviana? Huc refer. videtur *A. hemilis* Arrab. fl. flum. 10. t. 25? (v. s.)

13. *A. HETEROPHYLLA* (Muhl. in Willd. sp. 4. p. 377), caule pubescente villosulo paniculato, foliis caulinis pinnatifidis subdentatis petiolatis, racemis summis lanceolatis sessilibus, petiolis longè ciliatis, racemis terminalibus



solitaria, fructu sub apice spinis 5-6 acutis instructo. ① ad ripas fluminum circa Philadelphiam (Pursh). (v. s.)

i. *A. MICROCEPHALA*, glabrescens, foliis superioribus alternis longè petiolatis pinnatilibus sinuatis obtusis, lobis inæqualibus integris obtusissimis, foliis inferioribus indivisis subspathulatis obtusis. — in Cayennâ legit olim cl. Patris. Invol. mascula nutantia orbiculata integerrime dimidiò minora quàm in omnibus aliis. Fl. feminae juniores tantum vidi. (v. s.)

l. *A. PANICULATA* (Michx. fl. bor. am. 2. p. 183), caule pubescente ramosissimo superne paniculato, foliis utrinque viridibus bipinnatifidis, lobis lanceolatis, fructibus aggregatis pusillis globoso-obovatis subnervosis. ① in cultis à Canadâ ad Floridam necnon ad Nootka Am. bor. occid. (Hook.). Willd. sp. 4. p. 3-6. Ell. sketch 2. p. 478. Iva monophylla Walt. car. 232. — Pluk. alm. t. 10. f. 5. — Ambr. pusilla Rafin. in Ktt. hujus videtur var. nana. (v. s.)

k. *A. TOMENTOSA* (Nutt. gen. am. 2. p. 186), caule humili, foliis bipinnatifidis subtus albis et tomentosis, spicis solitariis. ② in Louisiana super. ad ripas flum. Missouri, rara, 1-2 pedalis.

j. *A. HISPIDA* (Pursh fl. suppl. p. 743), undique canescenti-hispida, foliis bipinnatifidis, lobis incisus, racemis terminalibus subpaniculatis ① in Carolina meridionali ex herb. Sherard sed Elliottio et Nuttalis ignota.

h. *A. TENUIFOLIA* (Spreng. syst. 3. p. 851), caule hispido, foliis bipinnatifidis subtus canescentibus, lobis remotis linearibus acutis subdentatis, racemis solitariis. — in Montevideo.

\*\* *foliis trilobatis.*

p. *A. CUMANENSIS* (H. B. et Kunth nov. gen. am. 4. p. 276), ramis hirsutis, foliis tripartitis pubescentibus subtus canescentibus, lobis pinnatilibus, intermedio trifido, fructibus 4-5 spinosis. ② in humidis prope Cumanam Novæ Andalusie.

o. *A. TRIFIDA* (Linn. sp. 1401), caule erecto piloso aspero, foliis etiã super. oppositis pilosis et scabris trifidis, lobis ovali-lanceolatis acuminatis serratis, petiolo angustè alato ciliato, fructu infra apicem tuberculis 6 subspinosis instructo apice conico. ① ad ripas flum. Amer. bor. à Canadâ ad Georgiam. Ell. sketch 2. p. 476. — Moris. ox. s. 6. t. 1. f. 4. Herba 4-6-pedalis. Cor. albæ. (v. v. c. et s. sp.)

n. *A. APTERA*, junior sparse setulosa, adulta tuberculis aspera, caule striato, foliis trifidis, lobis ovali-lanceolatis acuminatis serratis, petiolo tereti aptero, fructu infra apicem tuberculis 5-6 subspinosis instructo apice conico. ① in Mexico circa Bejar locis inundatis aug. flor. legit cl. Berlandier (pl. exs. n. 17781). Herba 3-6-pedalis ad *A. trifidam* proximè accedens, sed petiolo foliis distincta. Lobi interd. 5, duobus exter. minoribus. (v. s.)

\*\*\* *foliis indivisis.*

m. *A. EMBRYATA* (Michx. fl. bor. am. 2. p. 182), hirsutissima, foliis arctè sessilibus indivisis oblongis supra basin latecentem utrinque unidentatis, fructibus tetragonis infra apicem quadrispinosis. ① in regione Illinoensi (Michx.), ad partes occident. montium Alleghany (Pursh).

q. *A. INVASIVOLIA* (Muhl. in Willd. sp. 3. p. 275), caule..., foliis oppositis sessilibus ovatis acuminatis serratis basi ciliatis, racemis terminalibus plerumque ternis. ① ad ripas paludum Pennsylvaniae et Virginiae. Pursh fl. 2. p. 580. (v. s.)

† *Species exclusa.*

*A. ARBORESCENS* Linn. = *Franseria artemisioides*.

Div. VI. IVEÆ. — Ambrosiæ IVEÆ *Cass. dict.* 29. p. 175. — Ambrosiæ spuria *Cass. opusc.* 3. p. 61.

Flores ♂ et ♀ in eisdem capitulis. Cor. omnes tubulosæ 5-dentatæ aut inter-

dum in ♀ nullæ. ♂ Anthæræ liberæ approximatæ. Styli apice incrassati. ♀ Styli bipartiti aut simplices elongati ferè subulati hispiduli.

CCXC. PINILLOSIA *Ossa in litt. descr. et ic.*

Flores monoici ♂ et ♀ in iisdem capitulis: foeminei 2 apicali ante invol. squamas 2 majores siti; masculi 2 cum iis alterni corollâ tubulosâ 5-fidâ donati. Invol. 4-phyllo squamis cruciatim oppositis 3-nerviâ imâ basi subconcretis constans, bracteolis 2-linearibus brevibus extis stipatum. Recept. planum fortè omninò epaleaceum. Cor. ♂ tubo tereti, fauce apertâ, lobis 5 ovatis papilloso-puberis. Anth. exsertæ ovato-oblongæ, utrinquè obtusæ approximatæ nec concretæ. Stylus antheras paulò superans apice bilobus incrassatus glaber. Ovarium abortivum. Cor. ♀ nulla. Styli rami à basi ferè liberi elongati graciles apice subhispidi. Achænia obovata subcuneata, juniora præter apicem mucroniformem instructa pappo 4-aristato, aristis crassis retrorsim aculeolatis apice capitellatis, adulta quadricornuta cornubus coarctis nudis patulis et nectario mucroniformi terminata. Embryo crassus, cotyl. obovatis extus convexus. — Herba Caribæa repens glaberrima, Tetranthi facie. Folia opposita orbiculata longè petiolata integra. Pedunculi axillares petiolis longiores nudi 1-cephali. Flores albi. — Plantam ante diem supremum misit b. Ant. de La Ossa rogans ut si genus ex ei exstruendum, dicatum fuerit ill. Claudio Martínez de Pinillos insule Cubæ gubernatori, scientiæ fautori!

1. P. TETRANTHOIDES. ? in insulâ Sancti-Domingi legit cl. Bertero et in Cabi circa Havannam cl. de La Ossa et de La Sagra. Tetranthus Berteri Spreng. syst. 3. p. 459. Herba elegans ferè Linnæam referens sed multò minor. (v. s.)

CCXCI. TETRANTHUS Swartz prod. 116. fl. ind. occid. 3. p. 1885. t. 27. Cass. dict. 57. p. 343. Less. syn. 420.

Flores monoici, ♂ et ♀ in iisdem capitulis; foeminei 2 ante squamas inv. majores tubulosi 5-dentati; masculi 2 ante squamas minores etiâ tubulosi 5-dentati. Invol. 4-phyllo squamis ovalibus obtusis 2 sub ext. paulò majoribus constans, bracteolis 4-5 linearibus ciliatis acutis stipatum. Recept. planum paleis convolutis amplex. Cor. tubo gracili, fauce infundibuliformi, lobis oblongis subpuberulis subaequalibus. ♂ Anthæræ oblongæ apice breviter mucronatæ approximatæ nec concretæ. ♀ Styli rami longi graciles exserti apice divergentes hispiduli. Achæniû oblongum substriatum intra paleam involutum. Pappus brevis coroniformis ciliolatus. — Herba Domingensis repens. Folia opposita petiolata ovata mucronulata. Pedunculi axillares nudi foliis longiores 1-cephali. Flores albidii.

1. T. LITTORALIS (Swartz! l. c.), caule petiolisque glaberrimis, pedunculis adpressè subpubescentibus, foliis ovato-orbiculatis basi hinc inde cordatis. ♂ et ♀ ripas fluviorum Hispaniolæ. (v. s.)  
2. T. HIRsutus (Spreng. neu. entd. 3. p. 40), caule petiolisque hirsutis, pedunculis basi laxè hirsutis cæterum adpressè pubescentibus, foliis ovatis æquâ cordatis. ① in Sancto-Domingo legit cl. Bertero. T. littoralis Spreng. syst. 3. p. 459 non Sw. Paleæ distinctè ciliatæ. Herba tota priore paulò major. (v. s.)

† *Species exclusa.*

T. BERTERI Spreng. = Pinillosia tetranthoides.

ICH. IVA Linn. gen. n. 1059. Gærtn. fr. 2. p. 394. t. 164. Cass. dict. 24. 43. 59. p. 176. Less. syn. 219 non Riv. — Denira Adans. fam. 2. p. 118. — archonanthi sp. Faill. — Parthenii sp. Linn. ed. 1737. p. 265.

Flores monoici, ♂ et ♀ in iisdem capitulis, fœmineis paucis 1-sermibitu tubulosis seu campanulatis, masc. pluribus in disco tubulosis entatis. Invol. sæpius 3-5-phyllo campanulatum, squamis ovatis uniseriatis, rariùs imbricatum squamis 3-4-seriatis. Recept. planum lineari aut lineari-spathulatis. Styli fl. fœm. longè exserti subulati hispidi, fl. masc. breviores apice incrassati. Achænia disci abortiva, li obcompressa calva cornubus nec tuberculis onusta. — Herbæ aut lices ex Americæ ripis orti. Folia opposita aut summa alterna tri- via serrata aut integra. Capitula ad fol. super. axillas solitaria seu 2, breviter pedicellata, deflexa, semiglobosa, racemos terminales aceos constituentia. Cor. albidæ. Anth. luteæ. — Habitus Artemisiæ capitulis monoicis et fructu facillè diversa.

§. 1. *Involucris 3-5-phyllis subuniseriatis.*

I. XANTHIFOLIA (Nutt. gen. am. 2. p. 185), herbacea, foliis oppositis petiolatis ordato-ovatis acuminatis duplò serratis sericeo-villosis subtus canescentibus, pectis paniculatis nudis, invol. squamis 5 ovato-lanceolatis acuminatis. ① in ridis circa Fort Mandan ad ripas flum. Missouri. Herba 5-6-pedalis. Folia 2-3 Xanthii strumarii.

I. CILIATA (Willd. sp. 3. p. 2386), caule herbaceo hirsuto, foliis oppositis etiolatis lanceolato-ovatis rariter serratis, petiolis longè ciliatis, racemo constrictissimo, bracteis ovato-lanceolatis acuminatis longissimè ciliatis, capitulis axillaribus deflexis. ① in Amer. bor. pratis in dition. Kentuckensium et Illinoisiam. Pursh fl. 2. p. 580. Iva annua Michx. fl. bor. am. 2. p. 184 non ign.

♂. latifolia, foliis latè ovatis remotè dentatis trinerviis utrinquè petiolisque pubescentibus, summis alternis lanceolatis basi petiolisque ciliatis, bracteis ovatis acuminatissimis squamisque involucri obtusis ciliatis. — in Mexico circa Bejar legit cl. Berlandier (pl. exs. n. 18641). An à specie Illinoisensi diversa? (v. s.)

I. AXILLARIS (Pursh fl. bor. am. 2. suppl. p. 743), herbacea subpilosa, foliis utrumquè oppositis lineari-lanceolatis integerrimis, capitulis axillaribus solitariis nutantibus, invol. simpliciusculi squamis ovatis. ① in Louisiana super. (Pursh), in aridis et salinis ripis fl. Missouri (Nutt.), ad flum. Saskatchewan et Columbia (Hænk.). — Herba 8-9 poll. Flores albi (Pursh). Folia cuneato-oblonga obtusa ex Nutt. gen. am. 2. p. 185. — An species ab Hookero fl. bor. am. 2. p. 309. t. 107 egregiè descripta et delineata, ipso auct. dubitante, mininò eadem ac Purshiana?

I. ANGUSTIFOLIA (Nutt. in litt. 1825), herba tota adpressè pubera, foliis sessilibus linearibus aut lanceolato-linearibus, infer. subserratis oppositis, super. alternis integerrimis, capitulis axillaribus subsessilibus deflexis, invol. obtusis lœso subpuberis. ① in Amer. bor. territorio Arkansano detexit et mecum comm. cl. Nuttall. Capitula multò minora quàm I. annuæ, magis approximata et spicam foliaceam continuam constituentia. Fruct. mat. non vidi. (v. s.)

I. ASPERIFOLIA (Less. in linnæa 1830. p. 151), caule herbaceo adscendente, foliis lanceolato-linearibus acutis strigosis integerrimis, infer. oppositis, capitulis alternis cernuis, invol. 3-gono denticulato gamophyllo. — in pascuis prope Vera-Cruz.

I. ANNUA (Linn. sp. 1402), caule herbaceo striato petiolisque pubescentibus, foliis oppositis petiolatis ovali-lanceolatis utrinquè acutis grossè serratis trinerviis scabridis, summis integerrimis, capitulis axillaribus cernuis pedicellatis

invol. squamis ovalibus dorso pubescentibus. (1) in America merid. (Linn.), sed locus propr. incert. Linn. hort. ups. 285. Schmid. icon. p. 61. t. 16. Vitis foliis ternato-verticillatis, (v. s.)

7. I. FRUTESCENS (Linn. smcen. ac. 3. p. 25), fruticosa, foliis oppositis lanceolatis subpetiolatis acutis profunde serratis subscabris, summis integerrimis, apicalis axillaribus deflexis vix pedicellatis depresso-globosis, invol. 5-phylo obtusissimo. 3 in saluginosis maritimis à Novâ-Angliâ ad Floridam (Pursh), in Novâ-Cesareâ (Torr. 1). Ell. sketch 2. p. 474. Gaertn. fr. 2. t. 164. Lem. ill. t. 166. f. 2. — Pluk. alm. t. 27. f. 1. vulgò dicitur *Marble-Elder* ex Nutt. Variat foliis ternato-verticillatis. (v. v. c. et s. sp.)

8. I. CHRISANTHIFOLIA (H. B. et Kunth nov. gen. am. 4. p. 276), fruticosa pubescenti-cauescens, foliis oppositis petiolatis lanceolatis integerrimis glauciscentibus tripplinerviis, capitalis ad axillas subternis cernulis, invol. squamis 3-5 orbiculatis, rec. paleis obtusis, ext. obovatis, int. linearis-spathulatis 3 in im. Cubæ humidis juxta Havanam.

### §. 2. *Involucris imbricatis 6-9-phyllis.*

9. I. IMBRICATA (Walt. car. 232), suffrutescens seu herbacea glabra, foliis subulibus linearis-lanceolatis cuneatis carnosis, infer. oppositis trinerviis denticatis, super. alternis integerrimis, invol. imbricatis 3-4-seriatis, recept. paleis ovatis latius. 2; 3 ad ripas arenosas maritimas Carolinæ et Georgiæ. Michx. fl. h. am. 2. p. 184. Pursh fl. 2. p. 580. Nutt. gen. 2. p. 185. Ell. 1 sketch 2. p. 474. dms. opt. Inv. squama 6-9. Cor. fœm. tenues, stigm. simplicibus elongatis. Cal. fl. fœm. interdum ad latus internum fissum et idè forè ad ligulatas et ad div. vergentes. (v. s. sp.)

### + *Species exclusa.*

- I. MONOPHYLLA Walt. = *Ambrosia paniculata*.

## CCXCIII. EUPHROSINE.

Capitula monoica, fl. radii fœmineis apetalis 1-seriatis, disci masculis tubulosus 5-dentatis. Invol. 1-2-seriale patens squamis obovatis margine submembranaceis. Recept. planiusculum, paleis obovatis membranaceis, exter. majoribus, intimis multò minoribus. ♂ Ach. distinctæ. Cor. breves, tubo vix ullo. ♀ Styli rami teretes glabri obtusi. Flos centralis omninò abortivus filiformis apice penicilliformis. Achænia obcompressa, margine calloso cincta, lævia, inter squamas et paleas exter. sita, apice omninò calva. — Herba Mexicana glabra erecta, habitu nunc *Ambrosiam* nunc *Parthenium* referens. Folia alterna petiolata pinnatipartita, lobis oblongis pinnatilobatis sinuatis, lobulis secus rachin intermixtis parvis sinuato-dentatis. Pili pauci subtilius secus nervos primarios rigiduli. Capitula in paniculam ramosissimam disposita breviter pedicellata. Flores albidii. Achænum *Parthenii*, sed cum squamis vicinis non adhærens. — Ex fœminis nudis *Euphrosyne* et alludens ad nuditatem *Gratiarum*.

1. E. PARTHENIFOLIA. 2? (1)? in Mexico circa urbem legit cl. Berlandier (exs. n. 587). *Gymnostyles parthenifolia* Moç. fl. mex. icon. ined. Folia 5 in longa, variegata pinnatilobata. (v. s.)

DIV. VII. PARTHENIÆ. — Senecionidæ Ambrosiæ Partheniæ Linn. syn. 220.

Fl. ♂ et ♀ in iisdem capitulis, fl. radii 1-seriatis fœmineis ligulatis, Epib. subcoriaceis persistentibus aut tardè deciduis, fl. disci masculis tubulosus 5-dentatis. Recept. sæpius paleaceum. Achænia radii obcompressa margine callo cincta aut rarius trigona.

## CCXCIV. CONIOTHELE.

Capitulum multiflorum monoicum, ligulis 7-10 uniserialibus fœmineis, fl. disci pluribus masculis tubulosis fauce ampliata 5-dentata. Invol. subcampanulati squamæ 7-10 subuniseriales oblongæ. Recept. epaleaceum convexum. Achænia.... teretia calva, papillis minutis confertis pulverulento-velutinis, disci omnino nulla. — Herba Californica glaberrima erecta tenuis annua subsimplex. Folia alterna pinnatipartita, rachi lobisque remotis 2-3-jugis linearibus. Capitula terminalia pedicellata solitaria. Flores pallidè flavi. Inter Parthenieas accedit ad *Leptosynen* sed ab hac et à cæteris differt recept. nudo. — Nomen ex *κονία*, pulveris, et *θηλη*, papilla, alludit ad fructum pulverulento-pappilloseum.

1. C. CALIFORNICA. ① in Novâ-Californiâ legit cl. Douglas. Semen hujus stirpis sub microscopio observavit cl. Daby. Papillæ aquâ imbibitæ elongantur, rampuntur et ex iis oriuntur filamenta tenuia non articulata moleculis intus donata quæ seminis quasi integumentum villosum efficiunt. (v. s. comm. ab hon. Soc. hort. Lond.)

## CCXCV. LEPTOSYNE.

Capitulum multiflorum, fl. radii ligulatis fœmineis, disci tubulosi 5-dentati bisexualibus abortu sterilibus? Invol. biseriale, squamis exter. seu bracteis paucis linearibus foliaceis disco sublongioribus, inter. ellipticis vix acutis submembranaceis disco æqualibus. Recept. paleis membranaceis lanceolatis persistentibus onustum. Ligulæ latè ovato-subrotundæ. Cor. tubi fauce obconicâ basi subbarbellatâ. Anth. ecaudatæ. Stylus bifidus, radii vix exsertus obtusus, disci inclusus ramis cono hispido lato brevi capitellatis. Achænia (ex flore) obcompressa, calva seu pappo minimo coronato superata, margine subalata, glabra. — Herba Californica glabra annua tenuis. Radix simplex gracilis. Folia subradicalia erecta angustissimè linearia, alia integerrima, alia lobis parvis linearibus pinnatipartita. Caulis teres gracilis nudus 1-cephalus foliis triplò quadruplo longior. Flores flavi. — Genus fructibus maturis ignotis subdubium. An, floribus disci verè hermaphroditis admissis, potiùs *Anthemideum*? sed *Anthemideæ* paleaceæ nullæ in Americâ hucusque notæ? — Nomen ex *λεπτοσύνη*, tenuitas, gracilitas.

1. L. DOUGLASII. ① in Californiâ legit cl. Douglas. Herba strictè erecta 6-8-pollic. (v. s. comm. ab hon. Soc. hort. Lond.)

CCXCVI. PARTHENIUM *Linn. gen. n. 1058. Gærtn. fr. 2. p. 429. t. 168. Cass. dict. 38. p. 14. Less. syn. 319. — Partheniastrum Nissol. act. ac. Par. 1711. — Hysterophorus Vaill. act. ac. Par. 1720. — ArgYROCHÆTA Cav. ic. 4. p. 54. — Villanova Ort. non Lag. — Trichospermum Beauv. ined. ex Cass.*

Capitulum multiflorum heterogamum, fl. radii 5 uniserialis fœmineis ligulatis, disci tubulosi 5-dentati abortu styli masculis. Invol. hemisphæricum biseriale, squamis ext. ovatis, int. suborbiculatis. Recept. conicum cylindraceumve, paleis semiamplect. membranaceis apice latoribus onustum. Fl. ♂ stam. imo tubo inserta, anth. juniores vix cohærentes. Stylus indivisus. ♀ Ligulæ obcordatæ breves. Styli rami semiteretes obtusi. Achænia obcompressa lævia, cincta margine calloso, utrinquè basi squamis contiguis adhærente et demùm ab ovario secedente. Pappus hisquamellatus, squamellis aristæformibus seu suborbi-

culatis. — Herbæ seu suffrutices Americani hirsuto-canescentes. Folia alterna. Capitula albida fastigiato-paniculata.

### Sect. I. PARTHENIASTRUM.

Pappi squamellæ tennes brevissimæ imò nullæ. — Folia indivisa.

1. *P. FRUTICOSUM* (Less. in Linnæi 1830. p. 152), caule fruticoso, foliis petiolatis triangularibus acutis basi subcordatis aut in petiolum decurrentibus imbris dentatis, invol. squamis sub-3-seriatis obtusissimis.  $\zeta$  in Mexico ad Pta del Rio.
2. *P. TOMENTOSUM*, caule fruticoso, foliis petiolatis ovatis acutiusculis in petiolum non decurrentibus grossè dentatis suprâ molliter velutinis subtus petiolo remisque albo-tomentosis, paniculâ corymbosâ polycephalâ confertâ, invol. squamis biserialibus, inter. obtusissimis, achenio epapposo.  $\zeta$  in Mexico inter Oaxacam et Mitlam legit cl. Andrieux (pl. exs. n. 281). Habitus in sicco paulo refert icon. 44. ser. 2. Sweet br. fl. gard. ad genus omninò diversum referendam. (v. s.)
3. *P. INTEGRIFOLIUM* (Linn. sp. 1402), caule herbaceo, foliis ovalibus asperis inæqualiter dentatis aut basi subincisis, infer. in petiolum decurrentibus, super. amplexicaulibus, invol. squamis exter. subacutis.  $\zeta$  in montosis Virginie (Pursh), Carolinæ (Fras.!) et Georgiæ (Ell.). Willd. hort. ber. t. 4. — Delth. 302. t. 225. f. 292. (v. v. c. et s. sp.)

### Sect. II. PARTHENICHÆTA.

Pappus biaristatus, aristis longis rigidis. — Folia obtusè sinuato-pinnatifida.

4. *P. RAMOSISSIMUM*, caule fruticoso ramosissimo, foliis obovatis obtusè sinuato-incano-lanatis, invol. squami. obtusissimis, acheniis puberis obovatis, aristis a rectis elongatis  $\zeta$  in Mexici ditione Sancti-Ludovici de Potosi legit Berthier (pl. exs. n. 1342!). Achenia fortè juniora non marginata nervo solutè observavi, cæterum *P. incano* valdè simile. An genus diversum. (v. s.)
5. *P. INCANUM* (H. B. et Kunth nov. gen. am. 4. p. 260. t. 391), herbaceum totum albedo-incanum, foliis obtusè sinuato-pinnatifidis, lobis inf. inciso-dentatis, terminali trilobo, invol. squamis omnibus obtusissimis, acheniis aristis recurvo-subdivergentibus ① in montibus circa urbem Mexici. (Berl. pl. ex. n. 632.) (v. s.)

### Sect. III. ARGYROCHÆTA. — *Argyrochæta* Cav.

Pappi squamellæ ovali-oblongæ obtusæ membranacæ. Folia bipinnatifida

6. *P. HYSTEROPHORUS* (Linn. sp. 1402), herbaceum puberulum basi subbipedum apice subincanum, foliis variè bipinnatifidis, invol. squamis acutis ① in glareosis Amer. calid. frequens, nempe circa Mexico (Berl. pl. exs. n. 49), in prov. Texas (n. 1575), prov. Leoninâ (Mendex!), ins. Cubâ (h. Madr.), Trininitatis (Sieb.), Jamaicâ (h. L'Hér.), Sancti-Thomasi (Wydl.), Martini. (Willd.), etc., nunc in ius. Mauritiî introd. (Bout.). *Argyrochæta bipinnatifida* Cav. ic. 4. p. 54. t. 378. Villanova bipinnatifida Ort. dec. 48. t. 6 — Nissol. act. par. 1711. p. 423. t. 13. f. 2. (v. v. c. et s. p.)

† *Species exclusa.*

*P. LUTEUM* Spreng. = *Guzotia oleifera*.

### CXCVII. MENDEZIA.

Capitulum multiflorum radiatum, fl. radii paucis 1-serialibus formis ligulatis, disci tubulosis 5-dentatis hermaphr. abortivis. Invol. squamæ paucæ 2-3-seriales ovales obtusæ imbricatæ. Recept. parvum

conicum, paleis membranaceis oblongis complicatis. Cor. radii tardè deciduæ ellipticæ, radii tenues glaberrimæ. Styli disci rami apice summo hirtelli. Achænia radii obcompresso-plana elliptico-cuneata apice emarginata breviter biaristata utrinquè glabra margine subtuberculata, tuberculis pilos fasciculatos gerentibus; disci compressa abortiva. — Herba Mexicana annua trichotoma, caule tereti pubescente. Folia opposita oblongo-linearia integra puberula. Capitula terminalia solitaria bicolora, disco nempè intensè croceo ligulis pallidis. — Genus affine Verbesinæ ex achænio biaristato, sed multò affinius Parthenio et Tragocerati, priori characteribus, posteriori habitu. Dixi in honorem cl. Mendez botanici Mexicani, qui, mediante amiciss. L. Alaman, hanc et plurimas alias ab ipso detectas humanissimè communicavit.

1. *M. NICOLOR*. (1) in Mexici prov. Leoninâ ad occid. urbis Guanaxato legit cl. Mendez. (v. s.)

CCXCVIII. TRAGOCERAS Less. syn. 220. — *Tragoceros H. B. et Kunth nov. gen. am. 4. p. 248. t. 385.*

Capitulum 11-14 florum heterogamum, fl. radii 6-8-ligulatis foemineis, disci 5-6 tubulosis masculis 5-dentatis. Invol. ovatum, squamis 5-7 imbricatis membranaceis obtusis. Recept. minutum paleaceum. Styli rami capillacei patentes. Achænium disci abortivum, radii oblongo-obovatum, calvum apice truncatum, corollâ persistente magnâ coronatum. — Herbae Mexicanæ ramosissimæ graciles humiles annuæ. Folia opposita integra. Capitula solitaria ramos supernè aphyllis coronantia.

#### Sect. I. EUTRAGOCERAS.

Achænia radii costis extûs crenatis callosis destituta.

1. *T. LINNÆOIDES* (H. B. et Kunth l. c.), ligulis apice in cornua duo ferè subulata recurva fissis, achæniis utrinquè glabris subnucicostatis. (1) in aridis salis Mexici inter Aria et Montem Jorullo. Cor. disci violaceæ, radii albidæ.
2. *T. MICROGLOSSUM*, ligulis indivisis concavis achænio duplò brevioribus extûs villosis, achæniis intûs 3-striatis apice utrinquè villosis. (1) in Mexici prov. Leoninâ ultra Guanaxato legit cl. Mendez. Cor. disci pallidæ, radii virescentes. (v. s.)

#### Sect. II. SKIRROGLYPHIS.

Achænia radii costis 3 callosis, mediâ minore, lateralibus crenato-tuberculatis instructa.

3. *T. FLAVICOMUM*, ligulis indivisis 2-3-dentatis planis achænio vix brevioribus, suprà subtilissimè velutinis, subtûs sparè glandulosis et puberulo-hirtellis, achæniis intûs glabriusculis 1-costatis extûs callosis 3-costatis, costâ mediâ minimâ, lateralibus crenato-tuberculatis. (1) in Mexico circa Tehuantepec, Chigultan et Izaltepec. Fl. jul. aug. Caulis purpurascens. Ligulæ supernè pallidæ lotæ, subtûs subvirescentes. (v. s. comm. à cl. L. Alaman.)
4. *T. SCHRIBNEANUM* (Less. in linnæâ 1834. p. 269<sup>1</sup>), ligulis emarginatis concavis reflexis subtûs pubescentibus achænio multò brevioribus, achæniis intûs pubescentibus costis integerrimis extûs glabris calloso-3-costatis, costâ mediâ angustâ, lateralibus crenato-tuberculatis. (1) in Mexico ad Malpays de Nanligo. Ligulæ albidæ. Ex descr. conjungere videtur ligulas *T. microglossi* et achænia *T. flavicom.*

† *Species exclusa.*

*T. PRUCEDANIFOLIUM* Spreng. = *Isostigma prucedanifolium*.

Subtrib. II. *HELIANTHÆ* Less. *syn.* p. 221. — *Helianthæ* sect. 1, 3 et 4 *Cass. opusc.* 3. p. 57. *excl. gen. paucis.*

Capitula sæpius heterogama, fl. radii ligulatis seu fœmineis, disco hermaphr., rarissimè homogama discoidea, nunquam monoica seu dioica. Recept. paleaceum aut rarissimè ad centrum subnudum. Cor. fl. hermaphr. lobi crasse sæpius supernè papilloso. Achænia sæpius tetragona, nunc teretia aut obcompressa aut compressa, alia calva, alia pappo coroniformi aristato aut partim paleaceo, nunquam nec verè setoso nec omninò paleaceo superato. Anthæræ nigricantes ecaudatæ. Styli fl. herm. rami divergentes nunc truncati apice penicillati, nunc sæpius cono acuto aut appendiculo lineari terminati. — Folia sæpius opposita.

Div. I. *HELIOPSIDÆ*. — *Wedeliæ* et *Heliopsidæ* Less. *syn.* 221 et 223. — *Helianthæ* *Heliopsidæ* et *Rudbeckiarum* pars *Cass. opusc.* 3. p. 59.

Capitula heterogama, fl. radii fœmineis, aut rariùs homogama discoidea. Achænia sæpè cortice crasso donata obovata angulata aut compressa, nunquam obcompressa, calva aut pappo coroniformi aut setis paucis rigidulis irregularibus coronata. — Cortex achæniæ (calycis tubus) nunc à pericarpio vero subseparabilis, nunc arcuè adhærens.

CCXCIX. *PHILACTIS* Schrad. *ind. sem. hort. Gœtt.* 1831 et *diss. mus. cambricæ*, 1832.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis circ. 8-10 uniser., disci tubulosi 5-dentati hermaphroditis sterilibus. Invol. campanulatum imbricatum, squamis biserialibus adpressis. Recept. elongato-conicum scrobiculatum, paleis oblongis acutis complicatis. Cor. radii cum ovario continuæ oblongæ amplæ obtusæ 5-dentatæ, disci extus parèe glandulosæ. Anth. disci ecaudatæ. Styli radii exserti bipartiti, disci sæpius inclusi, ramis cono brevi superatis. Achænia radii trigona obpyramidata fertilia glabra coronata, extus ligulâ persistente, intus aristâ crassâ acutâ subciliatâ, disci compresso-subtetragona ex sicco abortiva subhirtella coronata aristis 4 cruciatim inæqualibus scabris imâ basi subconcretis ex angulis ortis. — Herba basi suffruticosa Mexicana dichotoma. Rami teretes, juniores villosi. Folia opposita petiolata latè ovata acuminata triplinervia serrulata utrinquè adpressè villosa. Pedicelli inter dichotomias orti 1-cephali foliis breviores. Ligulæ flavæ demùm subrufæ. Discus luteus. — Genus inter *Tragoceratem* et *Zinniam* medium.

1. *P. ZINNIODES* (Schrad. l. c.). 2. 5 in Mexico. Petioli 4-lin. Limbus fol. 1 poll. longus, sesquipollic. latus. (v. s. comm. à cl. Schrad.)

CCC. *ZINNIA* Linn. *gen. n.* 974. *Gertn. fr.* 2. p. 459. t. 172. *Cass. dict.* 3. p. 315. *Don trans. lin. soc.* 16. p. 2. p. 155. Less. *syn.* 224. — *Crassina* Scop. — *Lejica* Hill.

Capitulum radiatum, fl. radii fœmineis ligulatis, disci tubulosi 5-dentati hermaphr. Invol. imbricatum, squamis ovato-rotundis nigro marginatis. Recept. conicum cylindraceumve, paleis oblongis complicatis disci flores involventibus. Ligulæ coriaceæ reticulatæ obovatæ vel oblongæ cum achæniæ apice continuæ persistentes. Cor. disci super ovarium articulatæ, supernè pilis corollinis velutinæ. Achænia corticata



membranacea subalata, radii subtrigona calva, disci ext. subtrigona, inter. compresso-plana, apice ex angulo interiore uniaristata aut biaristata (aristâ inter. longiore), aut bidentata seu mutica. Cotyled. in achæniis trigonis involucrio parallelæ, in achæniis compressis invol. contrariæ. — Herbæ Americanæ annuæ, ramis foliisque oppositis. Folia 3-5-nervia integra. Capitula ramos terminantia solitaria, elegantia, diu persistentia. — Cl. Cassini dubitat an ligula sit verus pappus radii et cl. Don hanc sententiam confirmat.

§. 1. *Achæniis disci ad latus interius uniaristatis aut rariùs subbinis valde inæqualibus, paleis receptaculi integerrimis obtusis.*

1. *Z. tenuiflora* (Jacq. ic. rar. 3. t. 590. coll. 5. p. 153), caule erecto vix pubescente, foliis brevissimè subpetiolatis cordato-lanceolatis, pedunculis longissimis cylindricis, invol. oblongi squamis adpressis vix marginatis, ligulis lineari-oblongis sæpius apice revolutis, recept. paleis integerrimis, achæniis disci uniaristatis. ① in Mexico (Cav.), ad Tlapujahua (Keerl.) prope Cinapequara, Valladolid, Guanajuato, etc. (H. et B.). Sims bot. mag. t. 555. *Z. revoluta* Cav. ic. 3. t. 251. Ligulæ suprà coccineæ, 7-8 lin. longæ, vix 2 lin. latæ, integræ ant apice bidentatæ. (v. v. c. et s. sp.)
- a. *Z. acutispis* (ic. fl. mex. ined.), caule erecto hirtello, foliis breviter petiolatis lanceolatis villosis, pedunculis folio brevioribus cylindricis, invol. ovalis squamis adpressis, recept. paleis integerrimis, ligulis lineari-lanceolatis apice acutè bifidis, achæniis disci uniaristatis. ① in Mexico. An var. *Z. tenuifloræ*, sed ligulis suprà sordide albidovirescentibus et notis suprà indicatis satis distincta videtur. (v. ic. pict.)
3. *Z. leptorhoda*, caule erecto confertè pubescente, foliis sessilibus ovali-oblongis, pedunculis cylindricis folio æqualibus longioribusve, invol. obovatis squamis adpressis, ligulis obovatis emarginatis, recept. paleis integerrimis obtusis, achæniis disci 1-aristatis. ① in Mexico, ad Tlacolola in Oaxacanâ ditione (Andr. fl. exs. n. 314), et in montibus circa urb. Mex. (Berl.) pl. exs. n. 686). *Z. pauciflora* in fl. mex. ined. non Linn. Affinis *Z. multifloræ* sed pedunculis nec cavis nec incrassatis nec striatis et capitulo minore differt. Ligulæ suprà purpureæ aut rubræ. (v. s.)
4. *Z. multiflora* (Linn. sp. 1269. Linn. f. dec. 23. t. 12), caule erecto ramoso subhirtello, foliis vix subpetiolatis ovato-lanceolatis, pedunculis folia superantibus apice (præsertim centrali) obconicis cavis striatis, invol. campanulatis squamis adpressis, ligulis obovatis obtusis emarginatisve, recept. paleis integerrimis obtusis, achæniis disci 1-aristatis. ① in Mexici pratis juxta Valladolid et Páezaro (H. et B.), in Louisiana (Willd.), in Brasiliâ circa Bahiam (Blanch.) n. 191, ad littora maritima ins. Mauritiæ (Met.), sed verisim. culta aut introducta. An à *Z. pauciflorâ* satis distincta?
  - a. *rubra*, ligulis supernè rubris coccineisve. Jacq. obs. 3. p. 40. (v. v. c.)
  - β. *flava*, ligulis supernè flavis luteisve. H. B. et Kunth nov. gen. am. 4. p. 250. (v. s. in Carol. introd. ex Bosc.)
5. *Z. verticillata* (Andr. bot. rep. 3. t. 189), caule erecto sparsè hirsuto, foliis oblongo-lanceolatis, nunc irregulariter verticillato-congestis, nunc apicaliter subdigestis, pedunculis brevibus obconicis cavis striatis, invol. campanulatis squamis adpressis, ligulis obovatis apice emarginatis sæpius 2-3-seriatis, paleis recept. integerrimis subacutis, achæniis disci 1-aristatis. ① in Mexico. *Z. hybrida* olim in hort. Par. *Z. verticillata* Schkuhr handb. 3. t. 252 evidenter monastrum sed verisim. speciei nondum in statu naturali descriptæ. Caules angulato-sulcati sæpè quasi fasciati. Capitula fortè è pluribus facta. Fl. disci 5-10-andri, stylf 3-5-fidi. Conf. Don l. c. (v. v. c.)
6. *Z. pauciflora* (Linn. sp. 1269), caule erecto hirsuto, foliis sessilibus cordato-lanceolatis subamplexicaulibus, pedunculo terminali striato obconico cavo

# DICOTYL. SEU EXOGENÆ.

ERINIA.

lateralibus longioribus tenuioribus, invol. squamis adpressis, recept. achæniis subbiaristatis, aristâ ext. brevissimâ aut nullâ. (1) in ill. t. 685. f. 1. Rudbecka fol. oppos. Zinn. h. Gætt. 409. t. 1. Peruvianum Lin. sp. ed. 1. p. 920 excl. syn. Zinnia lutes Gært. t. 172. Bidens Mill. dict. t. 64. Lejica Hill. exot. 29. Crassia, ligulæ flavæ obovatæ apice obtusæ aut 1-2-dentatæ. (v. v.)

*Achæniis disci apice biaristatis bidentatisve aut muticis, recept. paleis integerrimis acutis.*

ERIDA (Sims bot. mag. t. 2123), caule erecto pubescente, foliis cordatis sessilibus amplexicaulibus, pedunculo terminali solitario hemisphaerico invol. squamis adpressis, recept. paleis acutis integris, achæniis subbiaristatis. (1) in Amer. merid. ex h. Pavon. Z. grandiflora hortul. Alor. sed differt caule pubescente nec hirsuto, fl. disci obsoleti, ligulæ obscure-coccineis nec flavis, pedunculo non striato. (1) hybrida ex Z. pauciflora et Z. elegans? (v. v. c.)

ERISTIFOLIA (H. B. et Kunth nov. gen. am. 4. p. 251), caule erecto paucis sparsè hirtello, foliis sessilibus lanceolatis hispido-scabris, pedunculis subaequantibus cylindricis, invol. squamis squarrosis, ligulis ellipticis, paleis integerrimis angustis acutis, achæniis disci emarginato-bidentibus ciliatis. (1) in Mexico prope Guanajuato. Ligula supra flavæ, convexus, ligulis paulò intensius flavus. (v. s. sp. comm. i. d. Jamaica.)

ERISTIMA (H. B. et Kunth nov. gen. am. 4. p. 251), caule procumbente sessilibus laevi apice subhispido, foliis breviter petiolatis oblongis obtusis o-scabris, pedunculis striatis superne incrassatis, invol. hemisphaerico adpressis, recept. paleis lanceolatis integerrimis, achæniis disci apice (1) in arenosis maritimis prope Acapulco.

## §. 3. Achæniis disci muticis aut subbidentatis, recept. paleis acuminate apice serrato-cristatis.

10. Z. ELEGANS (Jacq. coll. 3. p. 152), caule erecto hirtello, foliis sessilibus amplexicaulibus cordato-ovatis, pedunculis solitariis folio longioribus cylindricis, invol. squamis ovatis obtusis, superioribus nigro marginatis, recept. paleis serrato-cristatis, achæniis disci muticis vix subbidentatis, ligulis obovato-subtundis. (1) in Mexico (Cav. Karv.), ad montes Tixtlæ (fl. mex. ined.), cum Cuernavaca (Berl. n. 975). Planta in hortis seminibus varians, ab hac species duas ex involucri laevi aut squarroso divellenda?

A. invol. squamis adpressis aut intimis vix subaequantibus.

a. violacea, palearum apicibus ligulisque supra purpureo-violaceis. Z. violacea Cav. ic. 1. p. 57. t. 81. Andr. bot. rep. 1. t. 55. Z. elegans Jacq. rar. 3. t. 589. Sims bot. mag. t. 527. (v. v. c.)

β. alba, palearum apicibus ligulisque supra albis. (v. v. c.)

B. invol. squamis omnibus squarrosis.

γ. purpurascens, palearum apicibus ligulisque supra purpureo-violaceis. (v. v. c.)

δ. coccinea, palearum apicibus ligulisque supra splendide coccineis. — Z. violacea coccinea Lindl. bot. reg. t. 1294. (v. v. c. et s. sp.)

† Species exclusa.

Z. BIDENS Retz. = Glossogyne pinnatifida.

CCCI. *HELICTA* Cass. *bull. philom. nov.* 1818. *dict.* 20. p. 461. 46. p. 399 et 406 non Less.

Capitulum multiflorum heterogamum, fl. radii ligulatis 1-seriatis fœmineis, disci hermaphr. tubulosi 5-dentatis. Invol. campanulatum biseriale disco longius, squamis ext. 5 spathulatis apice foliaceis, inter. minoribus oblongis. Recept. convexum, paleis amplectentibus. Ligulæ ad basin usque fissæ nec basi tubulosæ. Cor. disci tubo nullo et ideò stam. sublibera. Antheræ nigricantes in appendice terminali tubercula grossa gerentes. Achænia conformia erostria compressa obovata oblonga, supernè hispidula, margine calloso utrinquè cincta. Pappus sessilis coroniformis brevis cartilagineus dentatus. — Frutex (patriâ ignotâ) sarmentosus, ramis teretibus junioribus hispidulis. Folia opposita subconnata ovalia obtusa remotè subdentata crassiuscula lucida. Pedicelli solitarii terminales et axillares. Flores lutei.

1. *H. SARMENTOSA* (Cass. l. c.). 3 Patr. ign. diù in hort. Par. sub nomine *Verbæ sinæ muticæ* culta.

† *Species exclusæ.*

*H. ALTERNIFOLIA* Less. = *Epallage salicifolia*.

*H. MADAGASCARIENSIS* Less. = *Epallage dentata*.

CCCH. *ALARÇONIA*.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis 1-seriatis, disci tubulosi 5-dentatis hermaphr. Invol. 2-3-seriale campanulatum, squamis laxis foliaceis hirsutis oblongis disco æqualibus longioribusve, intimis minoribus paleas referentibus. Recept. planum, paleis flores subæquantibus et semiamplect. Ligulæ amplæ apice 3-dentatæ. Cor. disci tubo brevi coriaceo, fauce cylindricâ longâ 5-nerviâ, dentibus vix extus apice subbarbatis. Styli rami fl. ♀ glabri, fl. ♂ longi densissimè hispidi acuti, exserti revoluti. Achænia prismatica crassa elongata, pappo calyciformi vario. — Herbæ perennes villosæ Californicæ Inulæ Helenii faciem ferentes. Caulis teres simplex hirsutus. Folia alterna oblonga aut elliptico-oblonga, inferiora basi attenuata, super. semiamplexicaulia, integerrima. Capitula ampla solitaria terminalia. Flores flavi. Genus habitu spectabile. An *Wedeliæ*, an *Heleniæ* affine? — Dixi in honorem peregrinatoris Ferdin. Alarçon, qui primus rut inter primos Californiam adiit anno 1540.

1. *A. HELENIOIDES*, caule foliisque molliter velutinis, radicalibus petiolatis amplis elliptico-oblongis utrinquè acutis, invol. squamis disco longioribus, achæniis glabro dentibus 5 lanceolatis acutis sericeo-cillatis crassis coronato. 2½ in Californiâ legit cl. Douglas. Folia radicalia pedem longa, 2½ pollic. lata. Ligulæ ferè sesquipollic. (v. s. comm. ab hon. soc. hortie. Londin.)

2. *A. ANGUSTIFOLIA*, caule hirsuto, foliis oblongo-lanceolatis, infer. subglabris, super. vix subhirsutis, invol. squamis disco æqualibus longè hirsutis, achæniis apice villosis, nunc truncato, nunc in aristam setosam scabram elongatam productis. 2½ in Californiâ legit cl. Douglas. Recepti mixtam cum præcedente cui habitu sat similis, sed caract. carpici adeò dissimilis ut facile ex hac specie genus constituendum esset. (v. s. comm. cum priore.)

CCCH. WEDELIA Jacq. amer. 217. t. 130. Rich. in Pers. ench. 2. p. 490 ad Adams. — Wedelia, Stenmodontia et verisim. Trichostemma Cass. dict. 46. p. 407-409. — Wedelia sp. Less. syn. 222.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligatis fœmineis, disci hermaphr. tubulosis 5-dentatis. Invol. 2-3-seriale, squamis ext. foliaceis, intimis membranaceis. Recept. subconvexum paleaceum. Styli fl. hermaphr. rami cono superati. Achænia obovata aut compressa erostrata, superata pappo coronati seu calyciformi magis subatipitato squamellis concretis dentatis ciliatisque constante. — Sæp. frutices aut herbæ. Species pleræque Americanæ, rarius Asiaticæ et Australasicæ. Folia opposita breviter petiolata serrata rarius trifida. Pedicelli terminales sæpitis solitarii, sæpè ex dichotomiis ramorum, rarius ex axillis foliorum, orti. Corollæ luteæ.

### Sect. I. CYATHOPHORA.

Achænia omnia exalata, apice calyculum dentatam gerentia, nec in tubercula callosa producta.

\* *Herbaceæ, foliis ad medium trilobis aut subtrilobis.*

1. W. CARNOSEA (Rich. in Pers. ench. 2. p. 490), caule herbaceo repente glabro, foliis sessilibus subcarnosis obovatis basi cuneatis subtrilobis repente subdentatis glabris, pedunculis axillaribus 1-cephalis, folio brevioribus aut paulo longioribus adpresso-villosis, invol. squamis hisciatis, ext. foliaceis oblongis obtusis disco aequalibus dorso subpuberis, int. minoribus membranaceis glabris, pappo coroniformi irregul. denticulato. ? in maritimis insul. Cuba-rum, ad Sanctum-Domingum (Poir.), Jamaicam (Sloan.), etc. — Cien. de 46. p. 409. Plum. ed. Barm. t. 107. f. 2. Silphium trilobatum Linn. sp. 136. Bupthalmum repens Lam. dict. 1. p. 515. Sloan. jam. t. 155. f. 1. Ligula 10-12 apice dentatæ. (v. s.)

β. *triloba* (Rich. l. c.), parcè pilosa, foliis manifestè trilobis. — Habet et ins. Cuba fol. lobis ferè acuminatis et ex Guadalupâ fol. lobis obtus. An diversæ? (v. s.)

γ. *aspera* (Rich. l. c.), foliis asperiusculis obsolete trifidis. — in Guadalupe legit Bertero. (v. s.)

2. W. PALUDOSA, caule herbaceo adscendente? glabro subanguloso, foliis subperennulis sessilibus oblongo-lanceolatis, basi angustatis obtusis, apice acuminatis, longè supra basin triplinerviis remotè serratis, serraturis duabus mediis longioribus, pedicellis axillaribus folio triplò longioribus pubescentibus, invol. squamis oblongis subacutis foliaceis disco paulò longioribus, intimis paleaceis brevioribus scariosis, achæniis elongati pappo calyciformi denticulato. (¶) in margines paludum in Brasiliâ circa Rio-de-Janeiro legit cl. Lund. Flora americana Ligula apice acutè 2-3-dentatæ. (v. s.)

β. *vialis*, caule hinc inde pilosiusculo magis demisso, foliis breviter acutè trilobis, pedicellis villosis, invol. squamis paulò latioribus ciliatis — ad viarum margines circa Bahiam legit cl. Salzmann. An species propria, an eadem in solo sicciorè nata? (v. s.)

γ. *latifolia*, caule glabriusculo, foliis latè ovatis profundè trilobis subatipatis. — circa Bahiam legit cl. Blanchet. Huc fortè pertinet *Acmella Brasiliensis* Spreng. syst. 3. p. 592, sive *Bupthalmum strigosum* Spreng. ar. entd. 2. p. 140, sed nihil asserere queo cum *Acmella* Spreng. septem et ciebuis ad septem genera diversa referendis constet. (v. s.)

3. W. GRACILIS (Rich. in Pers. ench. 2. p. 490), caule herbaceo repente subuliformi, foliis anguloso-lobatis pubescentibus membranaceis, achæniis turpè subtrigonis muricatis. — in Antiquâ et Sancto-Domingo, in siccis et montibus. An ex ipso auct. var. W. carnosæ?

\*\* *Herbaceæ, foliis serratis.*

4. *W. HISPIDA* (H. B. et Kunth nov. gen. am. 4. p. 215. t. 71), herbacea, ramis erectis hispidis, foliis lanceolatis aut subovato-lanceolatis basi cuneatis apice acuminatis medio inciso-dentatis subtriplinerviis utrinquè hirtis, pedunculis elongatis 1-cephalis è dichotomiis ortis, invol. squamis ext. oblongo-lanceolatis hirtis disco æqualibus, int. glabriusculis subæqualibus, achænio compresso-tetragono, pappo coronif. potiùs calyculiformi ciliato denticulato. ? in Mexici valle Tenochtitl (H. et B.), Ker bot. reg. t. 543. Desf. cat. h. par. ed. 3. p. 185. *W. calendulacea* Rich. in Pers. ench. 2. p. 490 non Less. *Stemmodontia scaberrima* Cass. dict. 46. p. 407. *Bupthalmum scabrum* Cav. ic. 2. p. 53. t. 167 vix bona, sed sub hoc nomine semina et specim. ex hort. Madr. habui. An fortè *Bupthalmum asperum* Horn. h. Hafn. suppl. p. 102? Flores subroccei ad quod alludit nomen Rich. Achænia sparsè puberula maculata utrinquè ad basin disco rufo calloso ornata. Pappus hinc indè in setulam productus. (v. s. et v.)
5. *W. MACRODONTA*, caule herbaceo ramoso tortuoso hispido, foliis sessilibus ovato-lanceolatis acutis grossè dentatis utrinquè hispidis, pedicellis è dichotomiis et apicibus ortis 1-cephalis hispidissimis aphyllis, invol. squamis foliaceis scabris acutis disco circ. æqualibus, ligulis plurimis vix invol. longioribus. ? in campis editis prov. Sancti-Pauli Brasiliæ ad Taubalé, novemb. flor. legit cl. Lund. Capitula multiflora, ligulis 15-20. (v. s. comm. à cl. Lund.)
6. *W. LUNDII*, caule herbaceo erecto trichotomo gracili setis hispido, foliis sessilibus lanceolatis utrinquè acuminatis utrinquè hispidis serratis paucis minimis apice donatis, pedicellis è dichotomiis ortis elongatis aphyllis 1-cephalis, invol. squamis ext. foliaceis disco longioribus, pappo calyciformi ciliato subaristato. ? in campis editis prov. Sancti-Pauli, ad Mugi flum. legit cl. Lund. Affinis *W. hispida*, sed habitu multò graciliori et foliis vix subserratis præcipue differt. Ligulæ apice dentatæ 8-9 lin. longæ. (v. s. comm. à cl. Lund.)
7. *W. VAUTHIERI*, caule subherbaceo erecto villis hirtello subsimplici, foliis ovalibus sessilibus acutiusculis basi trinerviis antrò reticulatis utrinquè sparsè setuloso-scabris, capitulo terminali solitario pedicello hirsuto instructo, invol. squamis ovalibus obtusis dorso scabris disci circ. longitudine. ? in Brasiliæ prov. Minarum-Generalium legit cl. Vauthier (pl. exs. n. 319!). Affinis hinc *W. Lundii*, illinc *W. radiosæ*. Rhizoma fruticosum. Ligulæ 8-9 lin. longæ, apice dentatæ, invol. duplò et ultrà longiores. (v. s. comm. à cl. Vauthier.)
8. *W. AFRICANA* (Pers. ench. 2. p. 490), herbacea, foliis breviter petiolatis lanceolatis serratis acuminatis trinerviis cauleque pubescentibus, invol. squamis ext. obtusis dorso puberis, int. subacutis glabriusculis paulò longioribus, ligulis circ. 20 apice bidentatis, achæniis calyculo subseasili denticulato. ? in Africæ æquinoctialis regno Owariensi. Beauv. fl. owar. 2. p. 18. t. 69.
9. *W. CALENDULACEA* (Less. syn. p. 222 non Rich.), foliis oblongo-lanceolatis basi attenuatis utrinquè strigoso-pilosis apice rariter serratis, pedunculis 1-cephalis solitariis axillaribus folio triplò longioribus, invol. squamis ext. oblongis subacutis disco longioribus, achæniis calyculo substipitato denticulato. — in Indiâ orient. pascuis humidis frequens, in Silhet, Bengalo, agro Maderapatanò, Zeylonâ, etc. DC. in Wight contrib. p. 17. *Verbesina calendulacea* Linn. sp. 1272. Willd. sp. 3. p. 2226. Wall. cat. et herb. n. 315. Lam. ill. t. 686. f. 1. *W. Benghalensis* Rich. in Pers. ench. 2. p. 10? *Jageria calendulacea* Spreng. syst. 3. p. 500. — Burm. zeyl. 52. t. 22. f. 1. Rheed. mal. 10. t. 42? mal. Variat foliis obtusis aut sæpius acuminatis, integerrimis plùs minùs serratis aut incisissimis, imò rariis, ferè trilobis. Achænia crassa obtusè tetragona rugoso-muricata margine 4-5-dentato crasso obtuso persistente coronata. (v. s.)
10. *W. URTICÆFOLIA* (DC. in Wight contrib. p. 18), herbacea subscandens, foliis petiolatis ovato-lanceolatis grossè et subinæqualiter serratis utrinquè strigoso-hispidis basi brevè apice longè acuminatis et sæpè incurvis, pedunculis solitariis 1-cephalis, invol. squamis 10 biseriatis acuminatis dorso pubero-strigosis, recept.

paletis acuminatissimis, acheniis calyculo breviter denticulato superatis. — in India orient. Fortè species 2 seu 3 distinguendæ, nempe:

- a. *Wallichii*, pedunculis folio brevioribus, foliis utriusque longè acuminatis membranaceis parciat involucrique strigosis. — in Nepaliâ. Wedd. Wallichii Less. in linnæa 1831. p. 162. (v. s. comm. à cl. Wall. anno 1831.)
  - β. *Wightii*, pedunculis folii circ. longitudinem sequentibus, foliis basi breviter acuminatis parciat strigosis. — in collibus Indiæ australis, ad Dugl. *Verbesina urticifolia* Wight! herb. *Verbesina biflora* var. Wall. ex n. 317. litt. B. et fortè D. (v. s.)
  - γ. *scaberrima*, pedunculis folio longioribus, foliis breviter basi acuminatis magis ovatis utriusque caulibus involucrique scaberrimis. — in regno An. ad Prome et in ins. Javâ (Bl.). Culta flore pleno evadit. *Verbesina urticifolia* Blume! hijdr. gr. r. *Verbesina scaberrima* Wall. cat. n. 311. An quis propria? (v. s.)
11. *W. CUNNINGHAMI*, herbacea erecta, caule subaspero, foliis petiolatis oblongo-lanceolatis basi acutis apice acuminatis serrulatis utriusque puberulis, petioli axillaribus terminalibusque 1-cephalis folio brevioribus, invol. squamis 6 linearibus, ext. lanceolatis acutis longioribus puberis, acheniis obcompressis calyculo breviter denticulato superatis. — in rupestribus insularum Goulbura ad em. borealem Australis mart. flor. legit cl. A. Cunningham. Affinis *W. urticifolia* var. sed involucri oligophyllis folisque multo angustioribus faciliè diff. (v. s. comm. à cl. inv.)
  12. *W. TRICHOSTEPHIA*, caule herbaceo erecto tereti hispido, foliis breviter petiolatis ovali-oblongis remotè serratis triplinerviis utriusque hispidis, petioli solitarii terminalibus villosis, invol. squamis ovato-lanceolatis foliaceis disco paulò longioribus, acheniis tetragonis pilorum zona superatis. — in Bruti *Trichostemma hispidum* Cass. dict. 46. p. 409. *Trichostephus* Cass. dict. 6. p. 618. *Trichostephium* Cass. opusc. 3. p. 59.
  13. *W. MOLLIS* (Rich. in Pers ench. 2. p. 490), herbacea, foliis petiolatis oblongo-lanceolatis crenatis membranaceis subtiliter pubescentibus. ? in America merid. Caulis pubescens. Folia obsolete 2?-nervia, sed imprimis ab eis valde venosa. †.
  14. *W. SUBVELUTINA*, caule herbaceo erecto tereti villosa, foliis vix petiolatis angustè lanceolatis basi obtusis apice acuminatis parvè et remotè serrulatis, præ ad basin triplinerviis, subtus reticulato-velutinoso villosis, supra scabridis demum glabris, pedicellis solitariis brevissimis, invol. squamis ovato-lanceolatis, acheniis calyculo minimo denticulato-ciliato. ? in Brasiliæ prov. Sancti-Pauli (h. imp. Bras. n. 4081), in paludosis (Lund!). Folia 3 poll. long. 5 lin. lata, petiolo 2 lin. longo. (v. s.)
  15. *W. PUBERULA*, caule herbaceo erecto tereti striato apice puberulo, foliis sessilibus ovatis vix acutiusculis dentatis utriusque tenuè puberulis subtus propè basin triplinerviis nervoso-reticulatis, pedicellis terminalibus ternis vix capient longitudine, invol. squamis ovatis disco brevioribus, achenio subpuberulo calyculo multidentato coronato. ? in Brasiliæ prov. Minas Generalium ad Barra legit cl. Vauthier (pl. exs. n. 327!). Folia sesquipollicem longa, fere pollicem lata. Ligula ovali-oblongè obtusè. (v. s.)
  16. *W. ERRENBERGII* (Schlecht. in linnæa 1831. p. 729), caule herbaceo tereti teretoso-piloso imò basi repente, foliis petiolulatis triplinerviis oblongo-ellipticis acutis serrato-crenatis hirsuto-hispidis supra asperimis, capitulis pedicellis terminalibus, invol. 10-phylli squamis ext. apiculatis extus hirsutis, paleis ciliatis, ligulis circ. 12 truncatis sub-3-crenatis, achenio obovato-tetragono pappo brevissimo coroniformi. — in Sancto-Domingo ad urbem Cap. Francæ

\*\*\* *Fruticosa aut suffruticosa, foliis serratis dentatisve.*

17. *W. ARTICULATA*, caule suffruticuloso sulcato adpressè puberulo subsæpe foliis breviter petiolatis ovato-lanceolatis basi obtusis apice acuminatis sub-

- rulatâ rigidulis suprà strigilloso-asperis subâs velutino-pubescentibus subtriplinerviis nervoso-reticulatis, pedicellis folio brevioribus solitariis adpressè villosis, invol. squamis ovalibus disco subbrevioribus, ext. dorso puberulis, achæniî calyculo villosulo, radii margine ciliato. ♀ in Porto-Ricco legit cl. Bertero, et verisimiliter in aliis insul. Caribæis. W. Jacquinî Rich. in Pers. ench. 2. p. 490. excl. syn. W. frutescens Spreng. ! in h. Balb. non Jacq. Ligulæ oblongo-lineares obtusæ invol. duplò longiores. (v. s.)
18. W. RADIOSA (Ker bot. reg. t. 610), canle basi suffruticuloso tereti asperulo, foliis petiolatis ovato-sen oblongo-lanceolatis acutis serrulatis utrinquè puberulis asperulis, pedunculis è dichotomiâ et spice ramorum 1-3 monocephalis folio longioribus, invol. squamis 3-seriatis, ext. ovatis puberis ciliatis foliaceis, intimis oblongo-linearibus membranaceis, pappo coroniformi ciliato-serrulato. ♀ in Brasiliâ. Ligulæ flavæ obtusæ vix 3-dentatæ invol. duplò longiores. Achænia juniora obcompressa disci ciliata, radii hirsutula, matura mihi ignota. (v. s. ex hort. Angl.)
19. W. CARACASANA, canle suffruticoso tereti subscabrido, ramulis pedicellis petiolisque hispidulis, foliis petiolatis ovato-lanceolatis basi cuneatis apice longè acuminatis serratis triplinerviis utrinquè sparè strigosis subâs pallidioribus, pedicellis terminalibus 1-cephalis subternis, invol. squamis ext. lanceolatis acuminatis strigillosis radium subæquantibus, ligulis angustis acutè bidentatis, achæniî pubescentis calyculo serrulato-ciliato. ♀ in Amer. austr. propè Caracas legit cl. Vargas. Folia ferè W. acuminatæ, sed distinctiùs triplinervia, pedunculi terni longiores. Cor. intensè luteæ. (v. s.)
20. W. ACUMINATA, canle fruticuloso, ramis teretibus pedunculisque strigososcabris, foliis petiolatis ovato-lanceolatis basi cuneatis apice longè acuminatis serratis utrinquè strigilloso-asperis subtriplinerviis, pedicellis solitariis 1-cephalis folio brevioribus, invol. squamis ext. lanceolatis acuminatis strigillosis radium subæquantibus, ligulis angustis acutè bidentatis, achæniî pubescentis calyculo substipitato denticulato. ♀ in ins. Cubâ juxta Havannam legit cl. Ant. de la Ossa. Folia 3 poll. longa, 12-15 lin. lata, interdum basi inæqualia. Petiolus junior ciliatus. (v. s.)
21. W. FRUTESCENS (Jacq. amer. p. 217. t. 130), canle fruticoso scandente glabro, ramis teretibus, foliis breviter petiolatis ovatis acutis leviter serratis utrinquè scabris, pedicellis terminalibus solitariis vix capituli longitudine, invol. squamis ext. subovatis laxis, ligulis obcordatis semibilobis, achæniî obovati calyculo 10-dentato. ♀ in fruticetis et sylvaticis circa Carthagenam. Polymnia Wedelia Linn. mant. 118.
22. W. AMBIGUA, canle fruticoso tereti glabro, ramulis petiolisque hispidissimis, foliis breviter petiolatis ellipticis utrinquè acuminatis subserratis, suprà sparè strigillosis, subâs secus nervos pubero-scabris, vix triplinerviis, pedicellis 1-cephalis hispidis folio paulò brevioribus, invol. squamis ext. lanceolatis scabris, int. paulò longioribus ovatis obtusissimis glabris, lignis acutè bidentatis, achæniî calyculo subdenticulato. ♀ in Guadalupâ legit cl. Bertero. W. calycina Spreng. ! syst. 3. p. 580. ex ipso in h. Balb. sed excl. syn. etiam Sprengeliano ! Achæniî calyculus nigricans. (v. s.)
23. W. LANCEOLATA, canle fruticoso tereti nervulis elevatis striato, ramis oppositis piloso-scabris, foliis breviter petiolatis lanceolatis basi brevè cuneatis apice acuminatis serrulatis suprà strigilloso-scabris subâs puberulis triplinerviis, pedicellis solitariis hispidulis folio paulò longioribus, invol. squamis ext. foliaceis lanceolatis acutis, int. obtusis subbrevioribus ovali-oblongis glabris, achæniî radii pubescente calyculo substipitato inæqualiter denticulato. ♀ in Porto-Ricco legit cl. Ledru. Ligulæ invol. paulò longiores oblongo-lineares obtusæ vix subdentatæ. Folia supernè ad orig. pilorum albo-punctata. Affinis W. frutescenti, sed invol. squamæ acuminatæ nec obtusæ. (v. s.)
24. W. AFFINIS, canle fruticoso dichotomo tereti glabro, ramulis hispidulis, foliis breviter petiolatis lanceolatis basi cuneatis apice acuminatis subserrulatis

- utrinque parce puberulis vix subtriplinerviis, pedicellis solitariis 1-cephalis hispidulis vix folio longioribus, invol. squamis foliaceis linear-lanceolatis disco paulo longioribus, ligulis acutè bifidis vix invol. longioribus, achenii puberuli calyculo subultrapitato vix subdentato.  $\S$  in insulâ Sancti-Thomasi legit cl. Wydlar (pl. exa. n. 51). W. calycina Rich. in Pers. ench. 2. p. 490? Vix affinis W. lanceolatae. (v. s.)
25. W. *PULCHRELLA* (H. B. et Kunth nov. gen. am. 4. p. 214), caule fruticoso procumbente, ramis teretibus adpressè pilosis, foliis petiolatis ovatis aut basi cuneatis serratis triplinerviis utrinque piloso-strigosis, pedicellis 1-terminalibus elongatis pubescentibus, invol. squamis ext. obovatis aut subultrapitando-ovatis acutis adpressè hispidulis, inter. glabris, ligulis circ. 7 bifidis, achenii pubescentis calyculo fimbriato.  $\S$  in humidis prope Cumana ubi dicitur *Miracul.*
26. W. *ACAPULCENSIS* (H. B. et Kunth nov. gen. am. 4. p. 215), caule fruticoso procumbente, ramis striatis junioribus piloso-hispidis, foliis petiolatis ovato-oblongis acuminatis basi acutis remotè serrulatis triplinerviis apice hispidè-scabris subtus piloso-strigosis, pedicellis solitariis 1-cephalis folio brevioribus pilosisculis, invol. squamis ext. lanceolatis aut ovato-lanceolatis hispidis ligulis apice bifidis, achenio puberulo calyculo subultrapitato fimbriato-ciliato  $\S$  prope Acapulco Mexicanorum ad littus Oceani Pacifici (H. et B.), et in ins. Sancti-Thomasi ex Schlect. in Linnaea 1831. p. 727. An W. calycina Rich. in Pers. 2. p. 490? ex Schlect.
27. W. *LATIFOLIA*, caule fruticuloso tereti, ramis scabridis, foliis petiolatis ovatis basi subobtusis, apice acuminatis remotè subserratis utrinque sparsiè adpressè puberulis triplinerviis, petiolis ciliatis, pedicellis 1-cephalis capiti longitudine, invol. squamis ext. ovato-lanceolatis acuminatis dorso villosis scabè radium subquantibus, ligulis latis acutè bidentatis, achenio calyculo subultrapitato denticulato-ciliato.  $\S$  in Amer. austr. Terrâ firmâ legit cl. Bertoni. *Nympha australis* Spreng. ! in h. Balb., sed in ejus syst. non reperire quivi. *C. flavo-croceæ* ! (v. s.)
28. W. *CRENATA* (Rich. in Pers. ench. 2. p. 490), suffrutescens procumbens. foliis petiolatis oblongis inæqualiter crenatis obtusis membranaceis subrugis. ovarii glandulosis, pappo obsoleto.  $\S$  in Guadalupæ arenosis. Ligula obtusè 3-dentatæ.  $\dagger$ .
29. W. *SERRATA* (Rich. in Pers. ench. 2. p. 490), suffruticosa, foliis longioribus petiolatis subcoriaceis lævibus lucidis, serraturis magnis, ligulis tridentatis. achenii calyculo margine excepto glabro.  $\S$  ad Sanctum-Domingum. Folia ferè llicina.  $\dagger$ .
30. W. *CRUCIANA* (Rich. in Pers. ench. 2. p. 490), frutescens tota mollissimè villosa, foliis petiolatis ovatis acutè dentatis, pedunculis longissimis, achenii puberulis variegati calyculo stipitato multipartito.  $\S$  in insulâ Sanctæ-Crucæ ad sepes arvorum viarumque W. Caribæa Spreng. syst. 3. p. 580.  $\dagger$ .
31. W. *PARVIFLORA* (Rich. in Pers. ench. 2. p. 490), frutescens, foliis sessilibus ovato-lanceolatis acutissimis integris asperimis rugosis, pedunculis brevibus. ligulis semibifidis, achenii puberulis calyculo subintegro seu levisimo crenatis.  $\S$  in campestribus Guadalupæ. Ex foliis sessilibus ab omnibus differre videtur.  $\dagger$ .

#### SECT. II. CALLOPHORA.

Achenia omnia exalata apice producta in tubercula duo callosa obtusa inter quæ pappus brevissimus coroniformis subserrulato-ciliatus sedet.

32. W. *GAUDICHAUDI*, caule suffruticuloso tereti, ramis pedicellisque hispidis. foliis subsessilibus ellipticis utrinque acuminatis vix apice serrulatis subtriplinerviis, supra strigillosis, subtus præsertim secus nervos puberulis, pedicellis elongatis 1-cephalis terminalibus seu è dichotomiis, invol. squamis brevibus oblongis villosis-subhirtis disco ferè longioribus, ligulis obtusis achenii spiræ



obtusè bicallosis et in centro pappum sessilem minimum subserulato-ciliatum gerentibus.  $\frac{5}{2}$  in Peruvâ legit cl. et am. Gaudichaud. Habitus *W. Caracasana* et *W. acuminata* sed achæniū diversissimū. An genus propriū? (v. s.)

## Sect. III. ACTINOPTERA.

Achænia radii alâ angustâ non membranaceâ marginatâ, disci exalata, omnia in pappum cyathiformem membranaceum obliquum dentatum producta.

33. *W. HELIANTHOIDES* (H. B. et Kunth nov. gen. am. 4. p. 216. t. 372), caule herbaceo dichotomo, ramis teretibus piloso-pubescentibus, foliis breviter petiolatis ovato-oblongis acuminatis remotè serratis triplinerviis suprà hispidulo-scabris subtus pubescenti-pilosis, pedicellis terminalibus elongatis solitariis invol. squamis subæqualibus oblongo-lanceolatis hispidis, ligulis 15-25? bi-dentatis, achæniis maculatis calyculo obliquo laciniato-fimbriato coronatis.  $\frac{2}{2}$  in collibus siccis inter Alauai et Ticsan Quitensium. Achænia disci immarginata, radii alâ utrinquè cincta, quâ notâ à genere *Wedelia* differt.

## Sect. IV? AGLOSSA.

Capitulum homogamum discoideum. Involucrum 1-seriale 5-phyllum. Achænia ad angulos tuberculosa. — An certè congener?

34. *W. DISCOIDEA* (Schlect. in linnæâ 1831. p. 728), caule herbaceo prostrato subhirsuto, foliis oppositis petiolulatis triplinerviis ovatis obtusis serrato-crenatis scabris utrinquè hirsutis, capitulis ad apices ramorum 1-3 breviter pedunculatis discoideis, involucri 1-serialis squamis 5-ovatis obtusis, paleis acuminatis apice longè ciliatis, achæniis ad angulos tuberculatis. — in insulâ Sancti-Thomasii.

† *Species exclusæ.*

*W. ARISTATA* Less. = *Wollastonia strigulosa*.

*W. AUREA* Hook. = *Verbesina aurea*.

*W. CHAMISSONIS* Less. = *Wollastonia canescens*.

*W. FORSTERIANA* Endl. = *Wollastonia Forsteriana*.

*W. MINOR* Spreng. = *Melampodium divaricatum*.

*W. OVATIFOLIA* Willd. = *Melampodium divaricatum*.

*W. PARVIFOLIA* Willd. = *Melampodium perfoliatum*.

CCIV. *JÆGERIA* H. B. et Kunth nov. gen. am. 4. p. 277. t. 400. *Cass. dict.* 24. p. 125. *Less. syn.* 223. non *Ræusch.* (1)

Capitulum multiflorum radiatum, ligulis fœmineis 1-serialibus, disci hermaphr. 5-dentatis. Invol. campanulatum oligophyllum æquale, squamis florum fœmineorum basin et demum achænia involventibus. Recept. parvum conicum, paleis carinatis diaphanis persistentibus. Styli rami recurvati. Achænia compressa 3-4-gona oblonga lævia apice calva aut pappo coroniformi minimo superata disco epigyno perexiguuo. — Herbæ Americane annuæ pilosiusculæ erectæ simplices aut oppositè ramosæ. Folia opposita subsessilia aut breviter petiolata ovalia subserrata. Capitula parva pedicellata.

1. *J. MNIODEA* (H. B. et Kunth l. c. t. 400), caule simplici puberulo 1-2-cephalo, invol. squamis ligulisque. 5 (1) in temperatis prope Ario Mexicanorum. Flor. lutei. Achænia calva.

---

(1) *Jægera* *Ræusch* genus monandrum sine descript. omnino ignotum.

# DICOTYL. SEU EXOGENÆ.

280851.

*virgata*, caule subsimplici oligocephalo sparsè pubero apice hirsuto, petalis ciliatis attenuatis subintegris supra sparsè pilosis subtus inflexis ovalibus sublanceolatis, ligulis paucis minimis. (I) in Brasiliæ cti-Pauli. Affinis *J. hirtæ*. Fructus junior refert iconem Kunth. Ligula ad. albæ aut albidæ. (v. s. in h. Mus. Par. à Mus. imp. Bras. sub n. 431

ss. syn. 223), caule oppositè ramoso hirsuto polycephalo, foliis l. squamis ligulisque circ. 10 ligulis obovatis exiguis. (I) in America aquosis prope Jalapam (Less.), in Brasiliâ (h. Mus. Berol.), à Lag. nov. gen. et sp. 3o. n. 383. Less.! in Linnæa 1830. p. 154. (v. s.)

*peruviana* caule ramoso adscendente ad nodos radicante infernè glabrato superius breviter hirsuto, foliis ovatis ovalibusve subacutis trinerviis is parçè puberulis, ligulis circ. 10 invol. longitudine. (I) ? in Brasiliâ de Janeiro ad Serra dos Orgaos legit cl. Vauthier (pl. rar. Brasiliæ, 1. 1. 1. 1.)

! *J. MACROCEPHALA* (Less. in Linnæa 1834 p. 270), caule ramoso oligocephalo, foliis plerisque oblongis, ligulis oblongo ellipticis. (I) prope Jalapam Mexicæ Capitula longè pedunculata solitaria minora quàm in *Erigeronte* alpino.

## † Species exclusæ.

*J. ABYSSINICA* Sm. — *oleifera*.

*J. CALENDULA* L. — *ulacea*.

CCCV. *LIPOTRICHE* R. Brown in Linn. soc. 12. p. 118. in add. 2o L.

Capitulum multiflorum radiatum, radii ligulatis 3-dentatis femininis 1-serialibus, disci hermaphr. tubulosi. Invol. duplici serie imbricatum subæquale. Recept. convexum, paleis foliaceis distinctis. Styl. disci rami appendice acuta hispidula. Achænia subuniformia turbinata, pappo setaceo caduco uniseriali 8-10-seto denticulato coronato. — Herbari in Africâ æquinoctiali spontanæ. Folia opposita indivisa. Pedunculi terminales terni. Flores flavi. Genus à cl. auctore *Melantheræ* proximè accedit, affine etiam *Wedeliæ* *Ecliptæ* et *Diomedæ* dicitur.

1. L. BROWNEI. — in Africâ æquin. ad ripas fluminis Congo legit infelix. Cl. Smith. Cæt. ign.

## † Species exclusæ.

L. AUSTRALIS Less. = *Lipochæta lobata*.

L. GYMNOLOMOIDES Less. = *Perimenium gymnomoloides*.

CCCVI. *MELANTHERA* Rohr in Kiobeh. nat. hist. selsk. 2. (1792) p. 1. p. 223. Cass. journ. phys. 1823. mai. p. 207. dict. 29. p. 483. 59. p. 158. — *Melanthera* Rich. in Michx. flor. bor. am. 2. (1803) p. 106. Brown. trans. soc. Linn. 14. p. 117. Less. syn. 222. — *Amellus* P. Brown. jam. 317 non Linn. — *Bidens* sp. Linn. — *Calex* sp. Jacq.

Capitulum multiflorum homogamum. Invol. disco breviori squamæ biserialibus subæquales. Recept. convexum paleis flores amplectentibus persistentibus onustum. Cor. tubo brevi, limbo 5-fido, lobis supra papillois. Achænia compresso-subtrigona basi incrassata apice truncata. Pappi setæ 2-10 rigidulæ caducæ irregulariter dispositæ. — Herbari rariùs suffrutescentes Americane ramosæ. Caules subtrigoni.

Folia opposita, rarius ternato-verticillata, petiolata, triplinervia, ovato-deltoidæa aut triloba, dentato-serrata, scabra aut subcanescentia. Capitula longè pedunculata 1-3 terminalia. Cor. albæ. Anth. nigricantes. — Nomen *Melantheræ* vetustius et constructum ex typis veteribus consecratis nempe *Melanthium*, *Melanthemum*, etc.

1. *M. HASTATA* (Rich. in Michx. l. c.), foliis oppositis hastato-trilobis scabris, invol. squamis lanceolatis, recept. paleis longè acuminatis rigidis.  $\frac{1}{2}$  in sicca Carolina (Boec! Fras! etc.) et in Cubâ prope Havanam (La Sagra!). Ell. sketch 2. p. 314. *Bidens nivea* Linn. sp. 1167 (excl. primò Dill. syn.)

a. *lobata* (Pursh fl. bor. am. 2. p. 519), foliis profundè trilobatis. — *Melanthera trilobata* Cass. dict. 29. p. 485. — Dill. elth. t. 47. f. 55. *Athanasia hastata* Walt. car. 201? (v. s.)

ß. *pandurata* (Pursh l. c.), foliis leviter lobatis panduræformibus. — *Melanthera panduræformis* Cass. dict. 29. p. 485. *Bidens nivea* Walt. car. 201? — Dill. elth. t. 46. f. 54. (v. s.)

2. *M. DELTOIDEA* (Rich. l. c. in adnot. sub *Melantherâ*), foliis oppositis indivisis ovato-deltoidæis aut subcordatis canescenti-scabris, invol. squamis ovali-oblongis, recept. paleis obtusis breviter mucronatis subfoliaceis.  $\frac{1}{2}$  seu  $\frac{3}{4}$  in loco natali; sæpius ① in hortis Eur. Hab. in insulis Caribæis, Jamaicâ (Br. Sw.!), Martinicâ (h. Duby!), Santo-Domingo (Bert.!), Cubâ (La Sagra!), Novâ-Granatâ ad fl. Magdalensæ (H. et B.). *Melanthera urticifolia* Cass. dict. 29. p. 484. *Melanthera* Linuzi Kunth nov. gen. 4. p. 199. *Bidens nivea* Linn. sp. 1167 var. a excl. syn. Dill. Swartz! obs. bot. 296. *Calea aspera* Jacq. ic. rar. t. 583. *Amellus* P. Browne jâm. 217. — Dill. elth. p. 55. t. 47. f. 55. n. 3. (v. v. c. et s. sp.)

3. *M. OXYLERIS*, foliis oppositis ovato-deltoidæis acuminatis dentatis utrinquè scabris subtus nervoso-reticulatis pallidis, involucri squamis receptaculique paleis acuminatis.  $\frac{1}{2}$  ad Guyaquil legit cl. Hænke. Affinis *M. deltoideæ* sed paleis valde acuminatis et capitulis paulò minoribus præcipuè differt. *Achenia* truncatis setis fragilibus numero variis interdum nullis (aut deciduis) coronata. (v. s. in h. Hænke.)

4. *M. CORYMBOSA* (Spreng. syst. 3. p. 445? excl. syn.), foliis ternato-verticillatis ovato-lanceolatis acuminatis serratis utrinquè scabris subcanescentibus, capitulis pedunculatis laxè corymbosis, invol. squamis ovatis, recept. paleis linearicuneatis vix acutis.  $\frac{3}{4}$ ? in Santo-Domingo legit cl. Bertero, fortè etiâ in Guadalupâ et Antignâ ex Spreng. neu. entd. 2. p. 135 sed ipse in vol. 3. p. 40 ejusd. libri Antignam omisit? An var. prioris? an folia constanter verticillata? Folia cum petiolo 15-16 lin. longa, 5-6 lin. lata. (v. s. ex herb. Bert. comm. à cl. Balb. qui etiâ ad cl. Spreng. misit.)

† *Species exclusæ.*

*M. GLANDULOSA* Wall. = *Wollastonia scalariscula*.

*M. ORIENTALIS* Spreng. = *Cassia axillaris*.

CCCVII. *OGIERA* Cass. dict. 35. p. 445. 43. p. 371. *opusc. phyt.* 3. p. 61. — Wedeliæ sp. Less. syn. 222.

Capitulum circ. 8-florum hæmogamum discoideum. Invol. biserialis circ. 10-phylli squamæ ext. foliæ majores, inter. paleas referentes. Recept. parvum planum paleaceum. Cor. 5-dentata. Styli rami in appendicem hispidam producti. *Achenia* oblonga gracilia subtetragona apice in collum brevissimum attenuata calva. — Herbæ piloso-hispidæ Americanæ. Folia opposita petiolata ovalia subdentata triplinervia subtus plus minus glandulosa. Capitula parva pedicellata. Cor. lutæ. — Cassinius prope Siegesbeckiam, Lessingius inter Wedeliæ collocat. Styli rami, in paucis floribus equidè visi, potius ad *Heliopsis* per-

tinent, sed à Wedeliâ genus facile differt capitulis discoidicis et achænio castrato calvo.

1. *O. TRIPLINERVIS* (Cass. dict. 35. p. 445). foliis utrinque hispidis subtus glandulosis, achæniis tuberculatis, capitulis solitariis. — in Guianâ?

β. *leiocarpa*, adscendens, foliis utrinque pilosiusculis, achæniis levibus, capitulis subadgregatis. — in Guianâ gallicâ. *Ogiera leiocarpa* Cass. dict. 3. p. 371. *Chalarium* Poit. ined. (v. s. specim. à cl. Ad. Brongniart mis.)

γ. *Portoricensis*, prostrata, foliis utrinque pilosiusculis, pedicellis ad ramorum dichotomias 2-4 subadgregatis, achæniis immaturis levibus, maturis tuberculosis. (1) in insulâ Porto-Ricco legit cl. Bertero. *Siegbertii* Portoricensis Bert. herb. (v. s.)

† *Species exclusæ.*

*O. TRIPLINERVIS* Bert. = *Euxenia* Mitqui.

*O. TRIPLINERVIS* Spreng. = *Euxenia* grata.

CCCVIII. *MONACTIS* H. B. et Kunth nov. gen. am. 4. p. 286 (excl. sp. 2). Less. syn. 211.

Capitulum 5-11-florum heterogamum, fl. radii unico ligulato semineo, disci tubulosis hermaphrod. 5-dentatis. Invol. cylindraceum subimbricatum, squamis lanceolato-linearibus. Recept. planum, paleis squamis referentibus sed carinatis. Stigma disci exsertum lobis revolutis. Achænia compressa, breviter rostrata et stipitata, calva glabra. — Arbor Aust.-Americana glabriuscula. Rami angulati subflexuosi. Folia alternâ petiolata ovato-oblonga subdenticulata triplinervia. Panicula terminata fastigiata ramosissima foliata. Flores flavi. Squamæ bracteolæque et Less. demum spiraliter tortæ.

1. *M. FLAVERIÆ*. 5 in prov. Bracamorensi inter urbem Jaen et ripam fl. Amazonum. *Monactis Flaverioides* H. B. et Kunth nov. gen. am. 4. p. 286. t. 46. *Phactusa* australis Spreng. syst. 3. p. 591.

OCCIX. *WOLLASTONIA* DC. in *Decaisn. nouv. ann. Mus. h. n. Par.* 3. p. 11. — *Wedeliæ* sp. Less. — *Verbesinæ* sp. Blum.

Capitulum multiflorum heterogamum, fl. radii 1-seriatis ligulatis fœmineis, disci hermaphr. tubulosis 5-dentatis. Invol. 2-3-seriale, squamis ext. oblongis foliaceis, internis minoribus membranaceis. Recept. planum aut subconvexum paleaceum. Cor. super ovarium articulare. Styli fl. herm. rami cono superati. Achænia crassa, obovata turbinate aut compressa, apice umbilicata, nunc nuda, sæpius aristas 1-5 tenues rigidulas fragiles deciduas gerentia. — Suffrutices aut rarius herbe. *Species* ex Indiâ orient., Moluccis aut Australiâ ortæ, sæpius hispide. *Wedeliæ* facie. Folia opposita breviter petiolata ovata acuta triplinervia dentata. Pedicelli sæpius ad apices ramorum terni 1-cephali. Flores lutei. — Genus medium inter *Wedeliæ* et *Heliopsidem*, ab utraque achæniis structurâ et pappo diversum. — Dixi in memoriam cl. chemici anglî Wollaston de scientiis physicis et naturalibus optime meriti et eorum suavitatem semper lugendi.

§. 1. *Wollastonia veræ, gerontogæ, radiatæ.*

1. *W. BIFLORA*, foliis petiolatis ovatis basi breviter apice longè acuminatis serratis supra sparso puberulo-scabris subtus glabriusculis, pedunculis 1-3 nervis

cephalis, uno terminali, 1-2 ex axillis supremis ortis, invol. squamis biserialis elongato-lanceolatis, achenio calvo sen 1-aristato. — in Indiâ orient. ad Malabarum, Martabaniam, Sangore, etc., frequens. *Verbesina biflora* Linn. sp. 1272. Willd. sp. 3. p. 226. Wall. cat. n. 317. excl. specim. paucis. *Wedelia biflora* DC. in Wight contrib. p. 18. *Acmella biflora* Spreng. syst. 3. p. 591. Rheed. mal. 10. t. 40 bona. Invol. squamæ oblongo-lineares acutæ foliaceæ dorso puberulæ. Recept. paleæ similes. Achenia crassa obtusè 3-4-gona basi attenuata 3-4-sulcata tuberculato-rugosa, vertice lato pubero, areolâ minimâ nunc aristâ ad latus inter. instructâ nunc nudâ. Certè ad Wollastoniam, nec ad *Adenostemma* ut vult Lessing (p. 156), nec ad *Acmellam* ut Spreng., pertinet. (v. s. comm. ab. bon. Cæt. merc. angl. Ind. or.)

2. *W. ASPERRIMA* (Decaisne nouv. ann. Mus. h. n. Par. 3. p. 414), herbacea, ramis teretibus subtrigosis, foliis petiolatis lanceolato-attenuatis acutis serratis basi acutis trinerviis utrinquè asperimis, capitulis solitariis, invol. squamis lanceolatis, bracteis scutè acuminatis, acheniis obovatis apice truncatis angularis tenuissimè punctulatis v. tuberculatis, pappo subunisetis. — in insulâ Timor.
3. *W. SCABRUSCULA* (DC. in Decaisn. n. ann. mus. 3. p. 414), caule herbaceo sulcato-angulato glabriusculo, foliis petiolatis ovato-lanceolatis acuminatis basi cuneatis supra basin triplinerviis grossè serratis suprâ sparè strigillosis subtus adpressè puberulis, corymbis terminalibus compositis trichotomis, invol. squamis biserialis ovali-oblongis acutis, acheniis crassis apice umbilicatis exaristatis. — in insulâ Timor leger. itin. Baudiniani botanici, in Indiâ orient. ad Penang, Singapore, Silhet, Rangoon cl. Wallich, et in insulis Mariannis cl. Hænke. *Eclipta scabruscula* Wall. cat. n. 322. comp. 322. *Melananthera glandulosa* Wall. herb. Meyera capitata Spreng. syst. 3. p. 601 ex litt. ad Wall. sed excl. syn. *Verbesina dichotoma* Heyn. herb. *Verbesina biflora* Blum. bijdr. p. 911. *W. aristata* et glabrata proxima, sed folia angustiora basi sensim cuneata supernè pilis delapsis albo punctata, capitula numerosiora compositè corymbosa et achenia calva, sed in a prioribus aristæ adeò variabiles ut vix earum absentiam tutò admittam. (v. s. comm. à Mus. reg. Par. et ab hou. Cur. Ind. or. angl.)
4. *W. MONTANA*, subscandens, foliis breviter petiolatis ovatis acuminatis grossè serratis utrinquè sparè striguloso-scabris, pedunculis 1-3 folio longioribus, invol. squamis 5 uniserialis ovali-lanceolatis acutis, acheniis 1-aristatis vix angulosis. — in montanis insulæ Javæ. *Verbesina montana* Blum. bijdr. 911. Caulis vix puberulus. Paleolæ recept. acuminatæ apice serrulatæ. Achenia apice minutè puberula ferè obovata. (v. s. comm. à cl. Blum.)
5. *W. MOLUCCANA*, suffruticosa di-trichotoma, foliis breviter petiolatis ovatis acuminatis rariter serratis basi sub-5-nerviis utrinquè ramulisque scabriasculis, pedunculis confertis subumbellatis scabrido-subhirsutis folio brevioribus, invol. 4-5-phylo, squamis ovatis paleisque margine ad apicem hirsutis. 3 in insulis Moluccanis. *Verbesina Moluccana* Blum. bijdr. p. 912. (v. s. comm. à cl. Lambert.)
6. *W. ZANZIBARENSIS*, caule herbaceo erecto glabriusculo, foliis petiolatis ovato-subrhombeis apice acuminatis trinerviis grossè serratis utrinquè actulas sparsas adpressas gerentibus, capitalis subternis longiusculè pedicellatis, invol. squamis et recept. paleis oblongo-linearibus subfoliaceis peristentibus cor. disci brevioribus, acheniis obovatis glabris obtusis, junioribus apice hirsutulis, adultis calvis. — in ins. Zanzibar legit cl. Bojer. *Anthemioipsis macrophylla* Bojer. in litt. 1836. Styli rami inclusi apice hispidi cono acutissimo. (v. s. comm. à cl. inv.)
7. *W. CANESCENS*, herbacea scandens? hirsuto-canescens, foliis ovato- seu elliptico-lanceolatis serratis acuminatis hirsuto-canescensibus, paniculis subcorymbosis, invol. biserialis squamis hirsutis acuminatis, acheniis muticis. — in insulâ Guahan. *Verbesina canescens* Gaudich. in Freyc. voy. p. 463. *Wedelia Chamissonis* Less. in Linnæâ 1831. p. 161.

8. *W. ELONGATA*, caule herbaceo erecto tereti ramoso glabrescente, foliis petiolatis oblongo-lanceolatis acutis serratis utrinque setis adpressis subacutis, pedicellis ternis folio duplò longioribus 1-cephalis, invol. squamis ovali-oblongis dorso subacutis disco aequalibus, acheniis obovatis glabris calvis. — in insulâ Madagascar legit cl. Bojer. *Anthemiopsis elongata* Bojer. l in litt. 1835. Foliorum joga remota internodio breviora. Recept. palæ submembranaceæ. Achenia epapposa collo minutissimo terminata nec omnino truncata. (v. s. comm. à cl. Bojer.)
9. *W. STRIGULOSA* (DC. in Decaisn. nouv. ann. mus. 3. p. 414), foliis longè petiolatis trinerviis aut propè basin triplinerviis ovatis acuminatis grossè serratis, suprà sparsè, subtùs in nervis et petiolis striguloso-pubescentibus, pedunculis ad apices ramorum 2-3 subalternis, invol. squamis biserialibus ovato-lanceolatis acuminatis dorso striatulis puberulis. — in insulis Rawak, Waigion, Pisang, et Moluccarum. *Verbesina strigulosa* Gaudich! in Freyc. voy. bot. p. 463. *Wedelia aristata* Less. l in Linnæa 1831. p. 160. *Buphthalmum helianthoides* Forst. prod. 57. non L'Hér. *Buphthalmum australe* Spreng. syst. 3. p. 605? ex Less. Achenia obtusè obpyramidata 3-et 4-gona obtusa subpuberula aristis 1-3 deciduis coronata. Recept. palæ obovato-oblongæ mucronatæ. (v. s. comm. à cl. Gaudichand et Less.)
10. *W. GLABRATA*, caule herbaceo glabro, foliis longè petiolatis altè triplinerviis ovatis subacuminatis remotè dentatis, suprà sparsè, subtùs secus nervos pilis parvis substrigillosis paucis puberis, pedunculis ternis terminalibus, invol. squamis biserialis ovali-oblongis acutis, acheniis 1-aristatis mucronatis. — in insulâ Timor legit. bot. Baudiniana exped. Affinis *W. aristata* sed car. infacile differt. (v. s. comm. à Mus. reg. Par.)
11. *W. INSULARIS*, caule basi suffruticoso adscendente, ramis vix apice pubescentibus, foliis petiolatis ovato-lanceolatis basi acutis trinerviis apice acuminatis subserratis utrinque petiolisque pubescentibus, pedunculis 2 dichotomis apice folia 2 ferè linearia et capitula 3 pedicellata gerentibus, invol. brevibus polyphyllis squamis ovali-oblongis acutis, fructibus exaristatis. Folia virilibus saxosis insularum ad oram borealem Australiæ apr. flor. legit cl. Cunningham. Affinis ex habitu *W. scabrescens*, ex calyce *W. Forsteriana*. (v. s. comm. à cl. inv.)
12. *W. PROCUMBENS*, caule procumbente basi suffruticuloso, ramis adscendentibus scabris, foliis sessilibus ovato-lanceolatis seu lanceolatis acuminatis serratis et grossè dentatis utrinque strigillosis asperimis trinerviis, capitulis pedicellatis ex dichotomis solitariis, invol. biserialis squamis substrigillosis ex foliaceis oblongis disco longioribus, achenio subtrigono glabro coronulâ triè brevissimâ superato.  $\frac{5}{2}$  in littoribus insularum ad oram borealem Australiæ apr. flor. legit cl. A. Cunningham. Folia pilis albis et post eorum delapsu maculis albis ut in *Borragineis* notata. Capitula pauciflora. Lignæ vix itè longiores. (v. s. comm. à cl. inv.)
13. *W. FORSTERIANA*, caule suffruticoso, foliis petiolatis oblongo-lanceolatis acutis grossè serratis suprà strigosis subtùs albido-villosis, pedunculis axillaris terminalibusque 1-cephalis.  $\frac{5}{2}$  in insulâ Norfolk (Forst.), locis apricis saxosis ad oram meridionalem insulæ (Cunn.), *Buphthalmum unilobum* Forst. prod. 541. Spreng. syst. 3. p. 605. *Wedelia Forsteriana* Endl. fl. not. p. n. 102. Ex Spreng. folia quintuplinervia sed in specim. meo evidè triplinervia. Invol. squamis 15-20 bi-seu triseriatis constans. Folia specim. mei ovali-oblonga latè dicenda. Fructum ex specim. juniore non judicare queo. (v. s. comm. à cl. Cunningham.)

## §. 2. *Wollastonia dubia*, *Americana*, *discoidea*

14. *W. LANTANOIDES*, caule fruticuloso, ramis striato-sulcatis subsapideis foliis petiolatis latè ovatis basi obtusis apice vix acutis crenatis, propè basin trinerviis, utrinque subpuberulis, subtùs nervoso-reticulatis, pedunculis terminalibus ternis 1-cephalis, capitulis discoideis, invol. squamis ellipticis axillaris

- disco brevioribus, acheniis obovatis calvis aut setis 1-3 fragiles eaduncas-apice gerentibus.  $\frac{3}{4}$  in insulis Caribæis legit cl. West. *Euphthalmum latifolio* affine Puer. herb. et h. Vahl. An *Wedelia discoides* Less. syn. 222 affinis aut fortè eadem? Cor. flavæ. Petioli pollicares. Limbus fol. 2 poll. longus et latus. (v. s.)
15. W. PROSTRATA, caulibus herbaceis procumbentibus adpressè pubescentibus, foliis oppositis obovatis obtusis basi in petiolum attenuatis integris glabrioribus, pedicellis axillaribus adscendentibus 1-cephalis folio triplò longioribus, invol. sub-1-serialis squamis elliptico-oblongis, acheniis crassis subtrigona compressive calvis aut aristam brevem gerentibus, recept. paleis lineari-subulatis.  $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Herba habitu ferè *Veronica* Allionii, fortè genus proprium constituens, sed floribus omninò in speciem deficientibus hic provisoriè adjecta. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1073 miss.)

## CCCX. TILESIA Meyer esseq. p. 252.

Capitulum multiflorum homogamum seu heterogamum, fl. radii ligulatis fœmineis bifidis, disci tubulosis 5-dentatis hermaphr. Invol. imbricatum 2-3-seriale, squamis coriaceis ovatis apice squarrosis. Recept. hemisphericum, paleis squamas referentibus flores amplectentibus. Styli disci et fortè radii rami setacei plumosi. Achenia obpyramidata tetragona immarginata calva, integumento externo (calycis tubo) separabili. — Planta scandens Guianensis scabra. Folia opposita petiolata ovata serrato-crenata utrinquè scaberrima. Capitula 1-3 ad apices pedunculorum congesta. Cor. crocæ.

- r. T. CAPITATA (Mey. l. c.). — in Guianâ batavâ ad Essequibo. Meyera capitata Spreng. syst. 3. p. 601. ex syn.

## CCCXI. PASCALIA Ort. dec. 4. p. 39. t. 4. Willd. sp. 3. p. 2228. Cass. dict. 46. p. 406. Less. syn. 223.

Capitulum multiflorum radiatum, fl. radii ligulatis fœmineis 1-serialibus, disci hermaphr. tubulosis 5-dentatis. Invol. disco æquale, squamis subbiseriatis linearibus subfoliaceis. Recept. subconvexum paleaceum, paleis lanceolatis semiamplectentibus. Achenium radii triquetrum squamis inv. non inclusum, disci tetraquetrum obovatum paleis semiinclusum (utrumque maturum ex Ort. drupaceum carnosum). Pappus coroniformis brevissimus irregulariter dentatus. — Herba Mexicana scabriuscula parcè ramosa strobilorum Pini odorem spirans (Ort.). Folia opposita lineari-lanceolata subtripplinervia integerrima, inferiora irregulariter dentata. Capitula terminalia solitaria. Cor. luteæ.

- r. P. GLAUCA (Ort. l. c.).  $\frac{1}{2}$  in Chili non longè ab oppido Chillon. (v. v. s.)

† Species vix nota.

- P. GLANDULOSA Fisch. ex Steud. nom. ph. 592.

†† Species exclusa.

- P. BACCATA Spreng. = *Walfia membranifolia*.

## CCCXII. RUMFORDIA.

Capitulum radiatum, ligulis multinerviis fœmineis circ. 10, tubulis hermaphroditis 5-dentatis. Invol. duplex, exter. squamæ 5 patulæ foliaceæ ovali-oblongæ disco breviores, inter. circ. 10 complicatè ovatæ mucronatæ, dorso glanduloso-puberæ, achenia radii involventes. Recept. plano-convexum, paleis squamas inv. inter. referentibus. Cor.

omnes basi puberæ super ovarium articulatae. Stylus radii longus apice bifidus glabriusculus, disci exsertus ramis semiteretibus apice puberulis cono brevissimo subsuperatis. Achænia obovato-compressa glabra calva conformia. — Frutex Mexicanus glaber. Rami teretes. Folia opposita ampla sessilia latè ovalia, basi attenuata, apice breviter acuminata, ad quartam partem longitudinis triplinervia, margine basi integerrima apice serrulata. Panícula oppositè ramosissima polycephala bracteis oblongis intermixta. Capitula pedicellata. Ligulæ pallidè flavæ, discus obscurius lutens. — Caract. ferè Heliopsidis, sed habitus diversissimus. Dicavi in memoriam cl. physici Benj. Thomson comitis de Rumford qui experimentis circa calorem et lucem et etiàm circa lignorum naturam momentosis inter hodiernos spectabilis remanet.

1. *R. FLORIBUNDA*. ♂ in Mexico detexit cl. de Karwinaki. Folia super. 8 poll. longa, 5-6 poll. lata. (v. s. in h. acad. reg. Monacensis.)

CCCXIII. *HELIOPSIS* Pers. ench. 2. p. 473. Dunal mem. Mus. h. n. Par. 5. p. 54. Less. syn. 223. — *Heliopsis* et *Kallias* Cass.

Capitulum multiflorum heterogamum, fl. radii ligulatis foemineis 1-serialibus, disci hermaphr. tubulosis 5-dentatis. Invol. bi-seu pleriseriale, squamis ext. sæpius foliaceis, cæt. subimbricatis. Recept. conicum, paleis oblongo-lanceolatis achænia amplectentibus. Achænia angulato-compressa glabra omninò calva, disco epigyno parvo. — Herbe Americanæ erectæ perennes oppositifoliæ. Folia dentata. Capitula solitaria pedunculata. Cor. flavæ. — An ab hac satis differt *Helepta* R. neogen. 1825. p. 3?

Sect. I. *EUEHELIOPSIS*. — *Heliopsis* Cass. dict. 24. p. 333.

Corollæ omnes glabræ, ligulæ super ovarium articulatae. Achænia lævia Invol. 2-3-seriale. — Herbe Boreali-Americanæ. Folia serrata.

1. *H. LÆVIS* (Pers. ench. 2. p. 473), caulibus glaberrimis, foliis glabris ovato-acuminatis serratis, invol. squamis externis lanceolatis patulis apice subseratis. ♀ in Amer. bor. à Pensylvaniâ ad Carolinam (Pursh), in fluminis ripis umbrosis rupestribus (Dun.). Dun. ! mem. mus. 5. p. 55. Hook. bot. mag. t. 337a. *Helianthus lævis* Linn. f. suppl. 1278. *Euphthalmum helianthoides* Linn. hort. ups. 264. L'Hér. stirp. 93. t. 45. *Silphium solidaginoides* Linn. p. 1302? *Rudbeckia oppositifolia* Linn. sp. 1280. — Moris. ox. 2. 6. t. 3. f. 64 (v. v. c. et s. sp?)

♀. foliis lævissimis, ligulis pallidioribus. *Silphium helianthoides* Gron. virg. 134. ex Dun. l. c.

2. *H. SCABRA* (Dunal ! mem. mus. h. n. Par. 5. p. 56. t. 4), caulibus foliisque scabris, foliis ovato-oblongis acuminatis serratis, squamis invol. ovato-acuminatis integris. ♀ in Amer. ad flumen Missouri et (ex Hook. fl. bor. am. 1. p. 310) ad ripas fl. Rubri (Red-River.). *H. canescens* Don in bot. reg. t. 39: excl. syn. (v. v. c. et s. sp.)

Sect. II. *KALLIAS*. — *Kallias* Cass. dict. 24. p. 326. 46. p. 406.

Corollæ omnes glabræ, ligulæ cum ovario continuæ nec articulatae. Achænia subdrupacea rugosa. Invol. 2-4-seriale imbricatum. — Herbe Austroli-Americanæ. Folia serrata. — An fortè ex ligulis persistentibus *Phylactidi* affiniore?

3. *H. EUPHATHALMOIDES* (Dunal ! mem. mus. Par. 5. p. 57), caulibus puberulis ap. villosis, foliis pilosiusculis ovato-acuminatis serratis, invol. squamis limbo integris imbricatis. ♀ in Peruvîa. *Anthemis ovalifolia* Ort. dec. 6. p. 73 lra



- cat. ed. 2. p. 125. *Verbesina ovata* hort. Par. ex Poir. dict. 8. p. 454. *Anthemis buphthalmoides* Jacq. h. Schomb. 2. p. 13. t. 151. Willd. sp. 3. p. 2185. excl. Cav. syn. *Amella buphthalmoides* Pers. ench. 2 p. 473. *Kallias ovata* Cass. dict. 24. p. 327? ex syn. (v. v. c.)
4. *H. CANESCENS* (H. B. et Kunth nov. gen. am. 4. p. 212), caulis sub foliorum jugis pedunculisque bifariam pilosis osterum glabrescens, foliis latè ovatis infer. subcordatis acuminatis serratis subvillosis, invol. squamis ovalibus imbricatis.  $\mathcal{U}$  in Peruvia. *H. rugosa* hort. berol. *Helianthus buphthalmoides* hortol. *Helenomium canescens* Willd. herb. ex Kunth. *Helianthus cordatus* Lamb. herb. ex Don. *Kallias ovata* Cass. dict. 24. p. 327? ex descr. *Kallias dubia* Cass. dict. 2. p. 333. Recept. intus cavum. Spec. à priori fortè non satis diversa. (v. v. c. et s. sp.)

Sect. III. *BALSAMORHIZA* Hook. fl. bor. am. 1. p. 310.

*Achenia compresso-tetragona calva. Capitula ferè Helianthi. Radix crassa balsamifera. Folia radicalia petiolata pinnatifidata. — An genus proprium ex Hook. ? †.*

5. *H. ? BALSAMORHIZA* (Hook. l. c.), subpubescens, radice crassâ balsamiferâ, foliis radicalibus plurimis petiolatis pinnatisectis, segmentis subpinnatifidis incisis, invol. squamis numerosis lanceolato-acuminatis imbricatis caulem paucifoliatum subsequantibus.  $\mathcal{U}$  in glareosis ripis fl. Columbia prope arcem Vancouver. Radix fusiformis succo terebinthinaceo. Folia 6-8 poll. longa.
6. *H. ? TEREBINTHACEA* (Hook. l. c.), pubescens, radice crassâ balsamiferâ, foliis radicalibus petiolatis ovato-lanceolatis sinuato-pinnatifidis crenato-serratis, invol. squamis numerosis lanceolato-acuminatis imbricatis caulem paucifoliatum subsequantibus.  $\mathcal{U}$ . Hab. cum priore ad quam valdè accedit.

† *Species exclusa.*

*H. PLATYGLOSSA* Cass. = *Guizotia oleifera*.

CCCXIV. *GUIZOTIA* Cass. bull. philom. 1821. p. 187. dict. 59. p. 247. — Ramilla DC. in Wight contrib. bot. ind. p. 18.

Omnia *Heliopsisidis*, sed cor. imâ basi et ad medium tubi pilis crassis subarticulatis annulatim instructæ, omnes super ovarium articulatæ. Achenium læve. Invol. biserialle. — Species *Gerontogea* annua. Genus inter *Heliopsisidem* et *Tetragonothecam* medium. Priori affine achenii disco minimo et invol. squamis liberis, sed distinctum corollis basi barbatis super ovarium omnibus articulatis; ad posterius accedens corollis basi barbatis et invol. biseriali, sed discrepans invol. squamis liberis et disco epigyno majore.

1. *G. OLEIFERA* (DC. mem. soc. hist. nat. Genev. v. 7. cum tab. 2), caule ad apicem pubescente, foliis semiamplexicaulis subcordato-aut ovato-lanceolatis remotè serratis subcabridis, invol. biserialis squamis ext. 5 latè ovatis foliaceis inter. superantibus. ① in Abyssiniâ ex Linn. f. sed fortè culta? in Indiâ orient. frequenter culta et verisim. etiâ spontanea? *Polymnia Abyssinica* Linn. f. suppl. (1781) p. 383. *Polymnia frondosa* Bruce voy. ed. 4<sup>e</sup>. gall. v. 3. p. 575. *Verbesina setifera* Roxb. cat. calc. (1814) p. 62. Sims bot. mag. t. 1017. Ainal. mat. mod. ind. 2. p. 256. *Verinnua Heyn.* tract. on Ind. p. 49. ex Ainal. *Parthenium luteum* Spreng. nov. prov. (1818) p. 31. *Heliopsis platyglossa* Cass. dict. 24. (1822) p. 332. *Tetragonotheca Abyssinica* Ledeb. ind. hort. dorp. suppl. 1824. p. 7. *Jugera Abyssinica* Spreng. syst. 3. (1826) p. 590. *Guizotia Abyssinica* Cass. dict. 59. (1829) p. 248. *Helianthus oleifer* Wall ! cat. et herb. n. 3194. comp. 304. (1830) et *Bidens*? Ramtilla Wall. l. c. n. 3191. comp. 301. *Ramtilla oleifera* DC. diss. mss. in soc. Genev. 1833 et in Wight contrib. bot. p. 18. *Buphthalmum* Ramtilla Ham. herb. et *Anthemis Mysorensis* herb. madr. ex

Wall. Calia in Abyssiniâ et in Indiâ orient. præsertim in agro Abyssinici ad oleum ex seminibus effluendum in lampadibus et condimentis tubulosis et olei acetum succedunt. Verase. in Indiâ dicitur Anti-PIN; Ann-TIL, Kurelloo, Kus-Yelloo, Werinusa, et in Abyssiniâ Noof ex Bratt.

a. *Andra*, foliis elongato-lanceolatis gracili serratis, ramulis elongatis subsymbois. — Hæc plerique synonyma supra citata pertinent. (v. v. c. et in Ind.)

ß. *augustæ*, foliis lanceolatis subdentatis, ramulis axillaribus scissis brevissimis. ① in Bengala inferiore ad marginem aquarum dulcium. Plantæ arboris cæcis sylvestris Wall.! n. 8194. (v. s. ap. in h. bot. Lin. Lond.)

CCCXV. TETRAGONOTHECA Dill. elth. 378. Linn. gen. ed. 1. n. 800. ed. 2. n. 976. L'Hér. carp. 197. Cass. dict. 69. p. 319. — Polymnia sp. Linn. syn. veg. Willd.

Capitulum multiflorum heterogamum, fl. radii ligulatis feminis 1-seriatis, disci hermaphroditis tubulosis 5-dentatis. Invol. duplex, ex squamis 4 basi in cupulam tetragonam concretis latè ovatis acuminatis maximis, int. squamis 8 ovali-lanceolatis parvis achenia radii amplectentibus. Recept. convexo-conicum paleaceum, paleis membranaceis lanceolatis acuminatis. Cor. omnes secus tubum hispidæ, super ovar. articulatis, radii latæ multinerviæ, disci dentibus erectis. Styli disci rami hispidi exserti. Achenia omnia obovata vix subangula crassa lævia calva, areolâ terminali latiusculâ, nectario elevato. — Heri Boreali-Americana oppositè ramosa. Caulis hirtellus teres. Folia opposita dentata parcè villosa. Pedunculi terminales 1-cephali. Capitula ampla. Flores lutei.

1. T. HELIANTHOIDES (Linn. sp. 1273). ① in Virginiâ (Linn.), Carolinâ (Frm.). locis arenosis siccis (Ell. sketch 2. p. 407) L'Hér.! stirp. 177. Polymnia Tetragonotheca Linn. syst. veg. p. 658. Abbot insect. 2. p. 137. t. 69 ex Willd. 3. p. 2116. Silphium Tetragonotheca Gærtn. — Dill. elth. 378 t. 283. f. 35. (v. s. ex herb. L'Hér.)

† Species exclusæ.

T. AFRASINICA Ledeb. = Guizotia oleifera.

CCCXVI. FERDINANDA Lag. nov. gen. et sp. p. 31. Cass. dict. 20. p. 429. ß. p. 404. Less. syn. 224.

Capitulum multiflorum heterogamum, fl. radii ligulatis feminis 1-serialibus, disci hermaphr. tubulosis 5-dentatis. Invol. hemisphericum disco brevius, squamis 2-3-seriatis adpressis. Recept. conicum. paleis oblongis submembranaceis amplectentibus. Anth. subnigris ecaudatis. Styli rami pubero-hirtelli cono brevi superati. Achenia rad. obconica angulata pappo 2-5-squamellato subcoroniformi minimo vel nullo superata, disci oblonga compressa sub-4-costata omnino calva. — Frutices Mexicani. Folia alterna ovata aut oblonga basi attenuata apice obtusa integerrima aut crenata triplinervia, supra pubescentia aut scabra, subtus velutino-argentea aut pubescentia. Corymbus ramosus polycephalus. Folia summa bracteiformia colorata. Flores lutei. — Genus ambigit inter Heliantheas et Anthemideas, prioribus stigmatibus posterioribus habitu toto affinis. — Toto cælo differt à Roseniâ quæ a Sprengel conjungere voluit.

1. F. AUGUSTA (Lag.! l. c.), foliis basi abruptè cuneato-attenuatis supra pube-

centibus subtus velutino-argenteis, floralibus integerrimis suprâ præcipuè aureo-flavicantibus, pappo radii squamellato irregulari, squamellis vix imâ basi concretis.  $\text{\textit{f}}$  in Mexico (Lag. Alam.), in montibus circa urbem (Berl. n. 663 et 932), in prov. Leoninâ ad occid. urbem Guanajuato (Mendez.). Ferdinanda velutina Desf. l. h. par. Cass. dict. 20. p. 430. Variat foliis ovatis oblongis integerrimis aut latè crenatis, semper versus basin abruptè angustatis. (v. v. c. et s. sp. et ex hort. Madr.)

2. *F. LUTESCENS*, foliis oblongis lanceolatisve obtusis integerrimis infernè sensim attenuatis suprâ vix pubescentibus subtus incanis, pappo radii nullo.  $\text{\textit{f}}$  in Mexici collibus aridis de Guadalupe prope urbem (Lallav. et Lex.). Anthemis lutescens Lallav. et Lex. nov. veg. t. p. 30 ex Alam. l. in litt. Priori valdè affinis. Rami ex sicco subscandentes. Folia et flores fragrantissimum odorem Abrotani æmulum spirant (Lallav. et Lex.). (v. s.)
3. *F. MINANS* (Lag. l. c.), foliis suprâ scabris subtus pubescentibus, petioliis foliaceo-alatis.  $\text{\textit{f}}$  in Mexico. Crt. ign. †.

CCCXVII. CHRYSOPHANIA Kunth in Less. syn. p. 224.

Capitulum radiatum, ligulis parvis rotundis fœmineis, cor. disci tubulosis hermaphr. basi obsolete puberulis. Invol. subuniserialia. Recept. conicum, paleis oblongo-ellipticis obtusis multinerviis. Stylus Zaluzaniæ. Achænia quadrangulata quadricostata conformia omninò calva, disco epigyno parvo, sectione transversali quadrata aut compresso-quadrangula. — Fruticulus Mexicanus dichotomus. Folia alterna elliptica, grôsse et inæqualiter crenata, petiolata, secus petiolum obtusè decurrenti-acuminata, suprâ canescentia, subtus incana. Paniculæ terminales fastigiatæ parùm ramosæ. Capitula lutea magnitudine Spilanthis acmellæ.

1. *C. FASTIGIATA* (Kunth l. c.).  $\text{\textit{f}}$  in Mexico prope Actopam. Folia 15 lin. longa, 9 lin. lata.

CCCXVIII. ZALUZANIA Pers. ench. 2 (1807). p. 473. Cass. dict. 56. p. 332 non Comm. nec Neck. (1). — Zaluzaniæ sp. Less. — Anthemidis sp. Ort.

Capitulum multiflorum heterogamum, fl. radii 5-7 ligulatis 1-seriatis ovalibus fœmineis, disci hermaphr. tubulosis 5-dentatis. Invol. biserialia petulum, squamis sericeis, int. brevioribus cuneatis trilobis. Recept. conicum, paleis membranaceis amplexantibus cuneatis apicè trilobis squamas int. referentibus. Cor. disci hispidæ, tubo faucis aequali basi achæniæ obtegente. Styli rami puberuli cono brevi terminati. Achænia calva, radii subtrigona hispida, disci compressa glabra lævia exangulata. — Herba perennis basi suffruticosa Mexicana ramosa villosa-tomentosa. Folia infima opposita, cætera alterna, petiolata pinnetipartita, lobis paucis obtusè sinuato-repandis sæpius trilobis. Capitula pedicellata corymbosa laxè subpaniculata. Flores lutei.

1. *Z. TAILORA* (Pers. ench. 2. p. 473).  $\text{\textit{f}}$  in Mexico (Ort.) circa Real del Monte (Berl. l. pl. exs. n. 396), in collibus de Los Remedios San-Angel (Lall. et Lex.). Anthemis triloba Ort. dec. 6. p. 72. Anthemis sinuata Lallav. et Lex. nov. veg. mex. 1. p. 30. Acnella trilobata Spreng. syst. 3. p. 591. (v. v. c. et s. sp.)

---

(1) Zaluzanskia Neck. = Marsilea. Zaluzania Comm. ined. = Berterizæ sect.

5. *R. vulpina* (Aiton kew. ed. 1. v. 3. p. 251), caule hispidulo ramoso polyphlo, foliis oblongo-lanceolatis hispidis, infer. denticulatis, super. integerrimis, basi angustatis subcordatis apice acuminatis, invol. squamis ovali-oblongis radium sequantibus, paleis recept. lanceolatis, pappo vix ullo.  $\frac{1}{2}$  in pratis sylvis montanis à Pennsylvania ad Carolinam. Sims bot. mag. t. 1948. *R. chrysomela* Michx.-f. bor. am. 2. p. 143. *R. aspera* Desf. h. par. ed. 3. p. 251. *R. fulgida* et *aspera* Pers. each. 2. p. 477. *R. fulgida* et *chrysomela* Spreng. syst. 3. p. 612. Ligulæ circ. 12 intensè luteæ, 3-dentatæ. Discus atro-purpureus (v. v. c.)
6. *R. hirta* (Linn. sp. 1280), caule hirsutissimo parce ramoso oligocephalo, foliis latè ovatis aut obovatis tripli-ant quintuplinerviis serratis hirtis, invol. squamis foliaceis hirsutis radium subequantibus, paleis recept. lanceolatis, pappo vix ullo.  $\frac{1}{2}$  Desf.  $\odot$  Pursh in montanis à Virginia ad Floridam. Sweet bot. fl. gard. 1. t. 82. — Vill. elth. t. 218. f. 285. Pluk. alm. t. 242. f. 2. Mill. z. t. 224. f. 1. Ligulæ flavæ subitè pubescentes. Discus atro-purpureus. (v. v. c. et s. sp.)
7. *R. grandiflora* (Gmel. hort. Carls. ex Steud.), caule ramoso angulato hispidulo, foliis petiolatis acutis basi attenuatis reticulato-venosis, infer. ciliatis 7-nerviis crenato-dentatis, super. lanceolatis scaberrimis oboletè crenatis. invol. squamis radio quintuplo brevioribus, ligulis dorso præcursat ad basin hispidis, recept. paleis pungentibus, pappo brevissimo subintegro.  $\frac{1}{2}$  in America boreali. *Centrocarpha grandiflora* Don in Sweet brit. fl. gard. ser. 2. v. 1. t. 87. Rudb. nudicaulis Nutt. mas. Ligulæ flavæ 2 poll. longæ dependens. Discus convexus atro-purpureus (v. v. in h. Gen.)
8. *R. stricta* (Wiegler. in Flor. 1829. 1. p. 30. Schrad. in litt. 1832), caule hispidulo, foliis puberulis, radicalibus longè petiolatis ovatis acuminatis 5-nerviis, caulinis sessilibus lineari-lanceolatis grosse indico-dentatis, cuneatis integris, invol. squamis lineari-lanceolatis ciliatis radio dimidiò brevioribus, paleis lanceolatis, pappo vix ullo. — patr. ign. Folia radicalia forè Plantaginis similis, caulina 5 poll. longæ, 6 lin. lata. Ligulæ circiter 24 latæ 8-10 lin. longæ apice tridentatæ. Discus atro-purpureus. (v. s. et v. c.)
9. *R. mollis* (Ell. sketch 2. p. 453), caule hispidulo-villoso ramoso, foliis sessilibus semisimplexicaulibus ovali-sublanceolatis dentatis et sup. integris mollibus tomentosis, invol. patulo villosissimo ligulis dimidiò brevioribus, paleis recept. lanceolatis, æcheniis vix 4-dentatis.  $\frac{1}{2}$  in Georgia distr. occidentilibus. Tota subcanescens. Folia caul. 15 lin. longæ, 9 lin. lata. Ligulæ 12-20 basi intrax. apice pallidiùs flavæ, 6-7 lin. longæ. Discus atro-purpureus. (v. s. e. ex bot. Cels. in h. L'Hér.)
10. *R. radula* (Pursh fl. bor. am. 2. p. 456), caule infernè hispidulo super. glabro nudiusculo, pedunculis longissimis 1-cephalis, foliis ovatis attenuatis tuberculatis hispidis, invol. squamis imbricatis ovatis acuminatis ciliatis. — in Georgia.
11. *R. discolor* (Pursh fl. bor. amer. 2. p. 574), caule foliisque strigis rigidis subtuberculatis hispidulo-scabris, foliis sessilibus spathulato-lanceolatis subintegerrimis, invol. squamis ovatis obtusiusculis radio paulò brevioribus, recept. paleis oblongis acuminatis apice fimbriatis, pappo vix ullo. — in Florida (Bart. à Pursh), Georgia distr. occident. (Ell. sketch 2. p. 455), et in territ. Arkansas (ex sem. à cl. de Pourtalès lectis). Ligulæ supernè intensè luteæ, subitè pubescentes pallidæ. Discus valdè convexus atro-purpureus. Nostra melius convenit cum descr. Elliotti quàm Purshii. (v. s. comm. à cl. de Charpentier.)
12. *R. lævigata* (Pursh fl. bor. am. 2. p. 574), glaberrima, caule lævigato punctulato, foliis ovato-lanceolatis utrinquè acuminatis integerrimis aut subintegerrimis triplinerviis, invol. squamis lanceolatis longitudine radii.  $\frac{1}{2}$  in Georgia pinetis. Ell. sketch 2. p. 454. Ligulæ breves pallidæ.
13. *R. glabra*, glaberrima, foliis caulinis distantibus sessilibus elongato-lanceolatis integerrimis utrinquè vix attenuatis apice callosulis, invol. squamis lanceolatis radio dimidiò brevioribus, paleis recept. lanceolatis, pappo 4-dentato. —

in Georgiâ circa Savannah? An fortè var. *R. lævigata*? Caules ramive elongati monocephali. Folia 5 poll. longa, 6 lin. lata, nervulis anastomosantibus notata. Ligulæ 12-15 lin. longæ flavæ apice acutè dentatæ. Discus fusco-purpureus. (v. s. comm. à cl. Doby.)

14. *R. SPATHULATA* (Michx. fl. bor. am. 2. p. 144), gracilis pubescens, caulis 1-cephalis, foliis amplexicaulibus ovato-spathulatis lævibus integerrimis, involucri patulo imbricato (?), ligulis 3-dentatis. — in montanis Carolinæ (Michx.) et Floridæ (Bartr.). Nutt. gen. am. 2. p. 178. De involucri imbricato silet Nuttall.

15. *R. GRACILIS* (Nutt. gen. am. 2. p. 178 non Banks), tenuis pubescens 1-3-cephala, pedunculis longissimis, foliis spathulato-ovatis remotè denticulatis, involucri patulo. — in pinetis à Virginiâ ad Floridam. Discus conicus fuscus.

+ *Species dubia, extra Americæ borealis confæderationem nascentes, fortè omnes à genere excludendæ.*

16. *R.?* *NAPIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 244), caule simplici scapiformi piloso 1-cephalo, foliis radicalibus hastato-lyratis crenato-serratis adpresso-hispidulis, invol. squamis biserialibus lanceolatis acuminatis, paleis recept. lanceolatis scariosis, pappo paleis ambuletis pluribus minutissimis constante. ? in Mexico juxta Santa-Rosa de la Sierra. An *Chrysocentematia* sp.? an genus proprium?

17. *R.?* *IRUCANTHA* (Lag. nov. gen. et sp. p. 3a. n. 399), foliis deltoideo-ovatis acutis serratis triplinerviis. ? in Mexico. Cæter. ign.

18. *R.?* *NUDICAULIS* (Pers. ench. 2. p. 477. n. 11), caule subscapiformi hirsuto, foliis oppositis sublanco-latis obsolete dentatis glabris, pedunculis longissimis strictis nudis. — ad Monte-Video ex herb. Juss.

19. *R.?* *DENSIFOLIA* (Smith in Rees cycl. 30. n. 10), caule scabro, ramis 1-cephalis, foliis oppositis petiolatis elliptico-lanceolatis, invol. squamis adpressis radio dimidiò brevioribus. — ad rupes circa Monte-Video legit Commerson. Ligulæ numerosæ latæ flavæ. Discus fusco-purpureus. Invol. squamæ scabræ fimbriatæ.

20. *R.?* *DECUMBENS* (Smith in Rees cycl. 30. n. 11), caule decumbente repente, foliis oppositis petiolatis lineari-spathulatis fortè lævibus, pedunculis erectis solitariis angulatis longissimis gracilibus leviter pilosis, invol. squamis ovatis acutis adpressis radio dimidiò brevioribus. — ad rupes circa Monte-Video legit Commerson. Genus ex auct. ipso valde dubium.

21. *R.?* *ELLIPSOIDES* (Smith in Rees cycl. 30. n. 12), caule decumbente, foliis oppositis petiolatis ovatis scabris, pedunculis erectis solitariis longissimis apice inflatis et velutinis, invol. squamis ovatis adpressis radio dimidiò brevioribus. — ad Monte-Video legit Commerson.

++ *Species exclusæ.*

*R. ALATA* Jacq. = *Helenium quadridentatum*.

*R. AMPLEXICAULIS* Vahl. = *Dracopis amplexicaulis*.

*R. AMPLEXIFOLIA* Jacq. = *Dracopis amplexicaulis*.

*R. ANGUSTIFOLIA* Linn. = *Helianthus angustifolius*.

*R. CINEREA* Hort. = *Obeliscaria pinnata*.

*R. CITRIODORA* Hort. = *Obeliscaria pinnata*.

*R. COLUMNARIS* Sims. = *Obeliscaria columnaris*.

*R. COLUMNIFERA* Fras. = *Obeliscaria columnaris*.

*R. DIGITATA* Willd. = *Obeliscaria pinnata*.

*R. HISPIDA* Hoffmans. = *Echinacea serotina*.

*R. OPURATA* Hort. = *Obeliscaria pinnata*.

- R. OPPOSITIFOLIA Linn. = *Heliopsis lævis*.  
 R. PAUCIFLORA. ex Steud. = *Zinnia pauciflora*.  
 R. FRUFOLIATA Cav. = *Dracopis amplexicaulis*.  
 R. PINNATA Vent. = *Obeliscaria pinnata*.  
 R. PURPUREA Linn. = *Echinacea purpurea*.  
 R. PURPUREA  $\beta$  Nutt. = *Echinacea serotina*.  
 R. SEROTINA Sweet. = *Echinacea serotina*.  
 R. TAGETES James. = *Obeliscaria Tagetes*.

CCCXXII. DRACOPIS Cass. dict. 35. p. 273 et 46. p. 400. Less. syn. 226 (excl. syn. Raf. et Neck.?). — Rudbeckia sp. auct.

Capitulum multiflorum heterogamum, fl. radii neutris ligulatis, disci hermaphr. 5-dentatis. Invol. biseriale, squamis ext. 6-8 linearibus patulis, int. parvis adpressis. Recept. conicum, paleis squamis int. interiores referentibus. Cor. disci subcylindraceæ, dentibus reflexis. Anth. inclusæ. Styli rami appendiculo lineari terminati. Achænia teres basi attenuatum; areolâ laterali, apice primâ fronte calvum, reversâ pappo coroniformi brevissimo integerrimo coronatum. — Herba annua Bor.-Americana glabra ramosa. Caulis ramique striato-sulcati. Folia caul. alterna cordato-amplexicaulia lævia integra acuta. Capitula terminalia solitaria, ligulis flavis, disco fusco.

1. D. AMPLEXICAULIS (Cass. l. c.). ① in Louisiana infer. ad ripas fl. Mississippi (Pursh) et in Mexico (Cav.). Rudbeckia amplexicaulis Vahl. act. soc. sci. nat. Hafn. 2. p. 29. t. 4. Schkuhr handb. 3. t. 259. Rudb. amplæxicaula Jacq. ic. rar. 3. t. 592. Rudb. perfoliata Cav. ic. 3. p. 27. t. 252. (v. v. c.)

CCCXXIII. OBELISCARIA Cass. dict. 46. p. 401. — Lepachys et Ratibida Less. journ. phys. 1829. p. 100. — Lepachys Less. syn. 225. — Rudbeckia sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis 1-serialibus neutris, disci hermaphr. tubulosi 5-dentatis. Invol. 1-aut 2-seriale. squamis ext. paucis linearibus patulis, inter. parvis obtusis à paleis fortè non distinctis. Recept. elongatum spiciforme, paleis complicatis apice hirsutis achænia disci semiamplexiculis. Styli rami breves et appendicem semilanceolatam parçè barbellatam producti. Achænia radii trigona hispida abortiva, disci bilateraliter compressa ovalia calva anticè margine subalata aut apice unidentata. — Herbae Bor.-Americane erectæ speciosæ glabriusculæ. Caulis striato-sulcatus. Folia alterna aut irregulariter subverticillata, pinnatisecta, petiolo striato, lobis 8-9 nunc indivisis, nunc pinnatipartitis. Rami apice longè aphylli 1-cephali. Ligulæ flavæ aut crocæ. Discus cylindricus, floribus parvis fuscis innumeris.

#### Sect. 1. LEPACHYS Raf. l. c.

Invol. biseriale, serie int. brevissimâ. Ligulæ angustæ. Achænia anticè subalata apice calva.

1. O. PINNATA (Cass. dict. 46. p. 402), tenuissimè pubescens, foliis pinnatisectis segmentis oblongo-lanceolatis utrinquè acuminatis parçè serratis, terminâ majore trinervi, superioribus indivisis. 2. in partibus occidentalibus Carolinæ et Georgiæ (Pursh), in regione Illinoensi. Rudbeckia pinnata Vent. l. h. cels. t. et t. 71. Michx. fl. bor. am. 2. p. 144. Smith exot. bot. 1. t. 38. Schrad. bot. journ. 2. p. 61. Sims bot. mag. t. 2310. Rudb. digitata Willd. sp. 3. p. 111. excl. syn. Lepachys pinnatifida Raf. et Less. ll. cc. Rudbeckia odorata et l.

cinerea hortul. Flores subodorati. Recept. dissectum odorem aromaticum spirant. Ligulæ flavæ pollicem et ultra longæ dependentes. Achænia anticæ alata. (v. v. c.)

## Sect. II. RATIBIDA Raf. l. c.

<sup>1</sup> Invol. uniserial. Ligulæ elongatæ angustæ. Achænia anticæ subalata apice calva.

2. *O. COLUMNARIA*, puberulo-hirtella, foliis pinnatisectis, segmentis lineari-lanceolatis acutis integerrimis, terminali cæteris subæquali, infimis gradatim subminoribus.  $\frac{1}{2}$  in campis Louisianæ superioris (Pursh), ad flum. Missouri (Sims), Saskatchewan et Red-River (Hook.). *Rudbeckia columnaris* Pursh fl. bor. am. 2. p. 575. Sims bot. mag. t. 1601. *Rudb. columnifera* Fraas cat. 1813 ex Pursh. *Ratibida sulcata* Raf. l. c. Ligulæ elongatæ flavæ dependentes apice bidentatæ, interdum ex Nutt. croceo-rubræ ut in *Tagetes patulâ*. (v. v. c.)

## Sect. III. MONODONTA.

Invol. biserial. serie int. brevissimâ. Ligulæ obovatæ. Achænia exalata ad marginem int. apice unidentata.

3. *O. PULCHRARIMA*, glaberrima, foliis pinnatisectis, segmentis pinnatipartitis, lobis linearibus distantibus integerrimis.  $\frac{1}{2}$  ? in Mexici provinc. Texas ad San-Fernando de Bejar, et in sinu Spiritûs-Sancti ad lacum Sancti-Nicolai legit cl. Berlandier (pl. exs. n. 1545 et 1960). Ligulæ fusco-croceæ (ferè *Tagetis* seu *Calliopsisidis* flor. æmulæ) margine et apice luteæ, patulæ demum deflexæ, 8-9 lin. longæ. (v. s. sp.)

+ *Non satis nota.*

4. *O. TAGETES*, hirsuta ramosissima, foliis radicalibus subintegris spatulatis (?) linearibus seu pinnatifidis, caulinis interruptè pinnatifidis lobis sæpius linearibus, pedunculis brevibus oligocephalis, ligulis recurvis. — In Amer. bor. ad montes Scopulosos (*Rocky-Mountain*). *Rudbeckia Tagetes* James exp. *Rocky-Mount.* 2. p. 353. *Rudb. hirsuta* Lais. 1824. p. 236. Herba pedalis multicaulis habitu *Marutæ* cotulæ. Ligulæ rubro-fuscæ. Discus fuscus minùs elongatus quàm in affini *O. columnari* dicitur.

## CCCXXIV. ANDRIEUXIA.

Capitulum multiflorum heterogamum, fl. radii circ. 20 uniserialis ligulatis fœmineis sterilibus, disci hermaphr. tubulosis 5-dentatis. Invol. biserial. squamis oblongis foliaceis vix acutis apice patentibus. Recept. convexum, paleis membranaceis acutis subconvolutis et flosculos semiinvol. Ligulæ subcoriaceæ cum ovario continuæ. Cor. disci cylindricæ glabræ. Stylus radii profundè bifidus glaber, disci rami subrevoluti apice summo barbati aut cono brevissimo superati. Achænia trigona, radii abortiva pubescentia, disci angustiora glabra omnino calva. — Herba Mexicana erecta parçè ramosa. Caulis angulatus, hinc indè subseriatum piloso-hirsutus, basi glaber. Folia opposita petiolata ovata acuta serrata pubescentia. Caulis apice pedunculiformis 1-cephalus subcapitulo obconicus velutinus. Flores lutei. Capitulum ferè *Rudbeckiæ*. Genus affine *Zinnia* et *Phylactidi* ex ligulis cum ovario continuis sed differt achæniis cortice tenui donatis et epapposis: *Rudbeckiæ* et *Gymnolomati* ex styli cano brevissimo, recept. convexo, etc., sed diversum ligulis distinctè styliferis persistentibus, achæniis trigonis omnino calvis. — Dicavi cl. Andrieux qui Mexicum peragravit et plantam ipsam cum pluribus aliis hinc indè citatis detexit.

1. *A. MEXICANA*. (1) ? in Mexico ad montem San-Felipe prope Oaxacam julio

*Spicula legit cl. Andrieux. Gymnolepis? Andr. l. pl. exs. n. 303. Herba squipedalis. Ligula 9 lin. longa. Anthera subnigrescentes. (v. s.)*

#### COCCOEY. ANOMOSTEPHIUM.

Capitulum multiflorum heterogamum, fl. radii & corollalibus ligatis neutris, disci hermaphrod. 5-dentatis. Invol. campanulatum pluri- (aut fortè uni-) seriale, squamis ext. ovatis acutis foliaceis late erectis, exterioribus oblongis membranaceis obtusis prioribus angustioribus ad longitudinem aequalibus. Recept. plenum, paleis membranaceis apice rufis subtendendis ceterum convolutis et tubosis involventibus. Ligula tubo gracili longo, limbo ovali. Cor. disci tubo brevi, fauce longè ampl. Antheræ exsertæ. Styli disci rami puberi, solum in flor. marginalibus obtusi, in centralibus cono apertissimo superati. Achenia lineari-subtragona gracilia, radii glabri apice breviter bicornia, disci adpressi villosa pappo irregulari coroniformi superata. — Herba Brasiliensis aut Caribæa basi suffruticosa erecta. Caulis subsimplex hispidus teris. Folia opposita sessilia utrinquè sparsè et adpressè piloso-scabra. Capitula ad apices caulis et ramorum pauca subsessilia pedicellata. Cor. luteæ antheris nigris. Genus habitu Wedelia, sed ligula neutra.

#### SECT. I. ANOMOSTEPHIA VERA.

Involucrum plariseriali disco brevius.

1. *A. OVATIFOLIUM*, caule confertè hirsuto, foliis sessilibus late ovatis trinervi basi obtusis apice vix acutis, capitulis breviter pedicellatis subconformis. ? in Brasiliæ prov. Minas-Geralium prope Sabara legit cl. Venthier (pl. exs. n. 301). Paleis recept. et invol. squamis post anthesis patentibus stultis. Antheræ exsertæ. (v. s.)
2. *A. BUPHTHALMOIDES*, caule folisque pube brevi rigidè asperissimis, foliis lanceolatis basi in petiolum attenuatis acuminatis vix suberratis penninervi pedicellis elongatis folium superantibus. ? in Guadalupe (Bertero! Krauss! loco dicto Pointe-à-Pitre (Perrottet!)), Buphthalmum asperissimum Spreng. n. entdeck. 2. p. 140 falsò demùm ab ipso auct. cum Wedelia calycina confusum. Antheræ inclusæ. Styli rami tennes. (v. s.)

#### SECT. II. ANOMOSTEPHIA DUBIA.

Involucrum subuniseriali disco longius aut æquale. — An genus diversum?

3. *A. ? OBLONGIFOLIUM*, caule sparsè hirsuto, foliis oblongo-lanceolatis atrinervi acuminatis subtriplinnerviis serrulatis, infimis ellipticis, pedicello terminis foliis longiore. — in Brasiliæ prov. Minas-Geralium ad Serra do Frio legit cl. Venthier (pl. exs. n. 317!). Herba hispida simplex. Invol. squamis circ. oblongo-lanceolatis disco longiores foliaceæ hirtæ. Paleis recept. membranaceis acutis glabris modicè convolutis. Ligula elongata apice dentata. Antheræ in classis fortè subeffortæ. Stylus longè exsertus, ramis crassiusculis. Achenia pinnoria, radii linearia pappo bidentato, disci villosa pappo substriato oblique-formi dentes 2 proferente, matura ignota et ideo de genere dubius noster. (v. s.)
4. *A. ELLIPTICUM*, caule sparsè subhirtello, foliis ellipticis et superioribus oblongis vix subserrulatis trinerviis subscabridis ciliolatis, pedicello foliis longior invol. squamis oblongis disco circ. aequalibus. ? in Brasiliæ prov. Sancti-Pauli. Valdè affinis *A. oblongifolio* sed multò minus hirtum, folia trinervia nec tripinnaria, nervis lateralibus nequè à basi nec longè supra basin ortis. (v. s. in Mus. reg. Par. à Mus. imp. Brasiliensi sub n. 409 missa.)



CCCXXVI. *ASPILIA* Pet.-Th. gen. mad. p. 12. n. 40. Cass. dict. 3. suppl. p. 57.

Capitulum multiflorum radiatum, ligulis 5-10 neutris 1-serialibus apice dentatis, fl. disci tubulosis 5-dentatis hermaphr. Invol. biseriale, squamis oblongis disco vix æqualibus. Recept. planiusculum paleis oblongis complicatis achænia foventibus apice acuminatis onustum. Styli rami cono brevi superati. Achænia linearia, adpressè villosa, superata pappo coroniformi dentato-ciliato. — Herbæ Madagascarienses procumbentes. Caules graciles puberuli. Folia opposita subsessilia lanceolata integra aut subdentata, suprâ sparè, subtus ad nervos setoso-sabra. Capitula è dichotomiis longè pedicellata solitaria. Flores flavi. — Car. ex A. Bojeri, nec ab A. Thouarsii mihi ignotâ.

1. A. BOJERI, invol. squamis erectis, achæniis (immaturis) lævibus. 2. in siccis inter dumeta per omnem insulæ Madagascar prov. Kmrnensem legit cl. Bojer. Ximenesia procumbens Bojer! in litt. 1835. Ligulæ 8-10. (v. a. comm. à cl. Bojer.)

2. A. THOUARSII, invol. cylindrici squamis 5 exterioribus apice reflexis, achæniis oblongis apice latioribus. — in ins. Madagascar, verisimiliter prope Foulpointe. *Aspilina* Pet.-Th. l. c. Ligulæ 5-6. An certè prioris congener?

CCCXXVII. *GYMNOPEPSIS*. — *Gymnolomia* H. B. et Kunth nov. gen. am. 4. p. 217 non. Ker. — *Aldama* Lallav. et Lex. nov. gen. 1. p. 14? Less. syn. 227.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis neutris, disci tubulosis hermaphrod. Invol. biserialia; squamis ext. subfoliaceis. Recept. planum convexum aut fortè conicum paleaceum. Styli disci rami appendiculati. Achænia pappo coroniformi dentato brevissimo coronata. — Herbæ aut suffrutices Americæ. Folia opposita petiolata tri- aut triplinervia dentata. Capitula pedicellata. — Genus malè definitum, differt à *Wulffia* achæniis non verè calvis nunquam carnosis, à *Montaguzia* paleis apice non spinosis, ligulis homom. nec heterochromis, stylique ramis longè appendiculatis. Species omnes fortè non sunt strictè congeneres nec satis notæ. — Nomen Kunthii falsum, Lallavei dubium, non admisi ne confusio oriatur, sed quàm minimè prius mutavi. Caract. pappi in 4 Kunthianis spec. observavit Ad. Brongniart et eorum icones humanissimè communicavit.

1. G. DENTATA, caule procumbente dichotomo tereti, foliis petiolatis sublanco-latis remotè serratis utrinquè asperis, capitulis è dichotomiis ortis longè pedunculatis, invol. squamis circ. 12 biserialis ciliatis, ligulis 8-10 linearibus, recept. conico, paleis achænia foventibus corrogatis apice encullatis. — in inundatis rivuli Huehneyapa ad Sonetum-Josephum del Corval Mexic. *Aldama dentata* Lallav. et Lex. nov. veg. 1. p. 14. Flores aurei. De pappo silent cl. auct.

2. G. SCHNEIDERANA, caule erecto hirsuto subdichotomo, foliis breviter petiolatis scabro-hirsutis triplinerviis oblongo-ovatis acuminatis remotè serrulatis, invol. squamis biserialis adpressis ciliatis, ext. ovatis, int. obovatis, recept. convexo, paleis achænia foventibus carinatis, achænio subtrigono-compresso. — inter segetes ad margines dumetorum prope Jalappam. *Aldama dentata* Less. in linnæ 1830. p. 154. 1834. p. 271. Flores aurei. Achænia pappo brevissimo coroniformi superata, intima sæpè abortiva.

3. G. VERAKSINORDRA, caule erecto subtetragono pubescente, foliis petiolatis ovato-lanceolatis basi cuneatis apice acuminatis vix subserratis triplinerviis utrinquè puberulis subcabridis, capitulis in dichotomiis apicibusque breviter pedicellatis, invol. biserialis squamis paucis, ext. lanceolatis, inter. ovalibus, ligulis 5-7, paleis complicatis rectis, achæniis glabris pappo coroniformi dentato

- superatis. — ad Trinitatem legit cl. Lockart (sed loci plures sunt hujus nominis?). Petioli 4 lin. longi. (v. s. comm. à cl. Hooker.)
4. *G. HELIANTHOIDES*, caule teretiusculo hirsuto et aspero, foliis breviter petiolatis lanceolatis aut subpandureformibus acuminatis triplinerviis hinc inde subdentatis utrinque strigilloso-scabris, petiolis ciliatis, capitulis ternis pedicellatis terminalibus, invol. biserialis squamis lanceolatis hirtellis, ligulis 10-12, recept. planiusculo, paleis membranaceis complicatis rectis, achæniis oblongis compressis, pappo coroniformi dentato. — ad Bonariam legit cl. Bacle. Capitula lutea. (v. s.)
5. *G. WEDELIOIDES*, caule erectiusculo tereti apice hispidulo, foliis subpetiolatis oblongo-lanceolatis subacuminatis basi trinerviis et obtusis serratis supra strigillis subtilis sparse puberis, capitulis pedicellatis terminalibus, invol. squamis biserialis subæqualibus oblongis foliaceis strigillosis, paleis complicatis cuspidatis rectis glabris, ligulis 8-10, recept. vix convexo, achæniis pappo coroniformi ciliato superato. — in ins. Caribæis, fortè in Tabago (h. L'Her). Affine *G. tenellæ* sed formâ foliorum potius oblongâ quàm ovatâ jam primâ fronte differt. (v. s.)
6. *G. TENELLA*, caule procumbente subvolubili tereti hispidulo, foliis petiolatis ovatis acutis basi subcordatis utrinque hispido-scabris dentatis, capitulis pedunculatis ramos terminantibus, invol. squamis circ. 20 lineari-lanceolatis acutis laxè imbricatis, ligulis circ. 10, paleis linearibus acutis subcarinatis, recept. convexo, achæniis obovatis subtetragonis puberulis, junioribus pappo coroniformi brevissimè dentato superatis  $\frac{2}{3}$  in Novâ-Granatâ prope Itagm et Contreras. Flores flavi. *Gymnolomia tenella* H. B. et Kunth nov. gen. am. 4. p. 218. t. 373.
7. *G. RUDBECKIOIDES*, volubilis, ramis sulcato-angulatis verrucosis, junioribus pilosis, foliis petiolatis ovatis acuminatis basi rotundatis crenato-serratis trinerviis utrinque hispido-pilosis, pedunculis axillaribus et terminalibus pilosis, invol. squamis 9-10 subbiserialis lanceolato-linearibus pilosis, paleis carinatis ovato-lanceolatis, recept. convexo, ligulis circ. 7, achæniis compresso-tetragonis, junioribus adultisque pappo vix dentato brevissimo coronatis.  $\frac{2}{3}$  in temperatis prope Ayavaca Peruvianorum. *Gymnolomia rudbeckioides* H. B. et Kunth nov. gen. am. 4. p. 218.
8. *G. HONDENSIS*, caule tereti piloso, foliis petiolatis ovatis subacuminatis basi rotundatis trinerviis crenato-dentatis utrinque hispido-pilosis, capitulis longe pedunculatis axillaribus, invol. squamis 15-20 lanceolatis laxè imbricatis hispidis, recept. convexo, ligulis 2-3-fidis, achæniis junioribus apice pappo inæquali acutè et longiusculè dentato coronatis.  $\frac{2}{3}$  in calidissimis prope Hondâ Novo-Granatensium. *Gymnolomia hondensis* H. B. et Kunth nov. gen. am. 4. p. 218. Cl. auct. dubitat an à *G. tenellâ* satis differt, sed, ex icone specim. authenticæ ab amic. Ad. Brongniart missæ, satis distincta pappo videtur.
9. *G. TRIPLINERVIA*, ramis teretibus glabriusculis junioribus adpresso-pilosis, foliis petiolatis ovatis angustato-acuminatis crenato-dentatis basi acutis triplinerviis utrinque adpresso-hispidulis, capitulis axillaribus longè pedunculatis, invol. squamis oblongo-lanceolatis laxè imbricatis, achæniis obovatis brevibus pappo coroniformi inæqualiter acutè et longiusculè dentato coronatis.  $\frac{2}{3}$  in altâ planitie Bogotensi. *Gymnolomia triplinervia* H. B. et Kunth nov. gen. am. 4. p. 219.
10. *G. ? EUXENIOIDES*, erecta setulis adpressis scabra, foliis ovato-lanceolatis basi cuneatis acuminatis serratis, capitulis 2-3 ad apices ramorum, invol. squamis polyphylo. — in Brasiliâ ad viam Felisbertiam. *Euxenia radiata* Nees et Mart. act. soc. nat. cur. 19. p. 7. Ab *Euxeniâ* differt capitulis radiatis et ipsis docent. auct. accedit ad *Gymnolomiam*.

† *Species exclusa.*

*GYMNOLOMA MACULATUM* Kef. = *Wulffia maculata*.

CCCXXXVIII. WULFFIA *Neck. elem.* 1. p. 35. *Cass. dict.* 46. p. 398 et 403. — *Chakistella Cass. dict.* 43. p. 37 et 46. p. 402. — *Chylodia Rich. ex Cass. journ. phys.* 1823. mai. p. 214. non *Brown*. — *Gymnoloma Ker bot. reg. t.* 662. *Less. syn.* 227 non *Kunth*. — *Coreopeidis, Helianthi, Verbesinæ et Cassulæ* sp. *auct.*

Capitulum multiflorum sæpius heterogamum, fl. radii 1-serialibus ligulatis neutris, rarius homogamum discoideum fl. tubulosis hermaphr. Invol. subhemisphæricum, squamis laxè imbricatis sublanceolatis. Recept. subconvexum, paleis lanceolatis subcarinatis planisve persistentibus. Styli rami cono superati. Achænia radii abortiva, disci crassa obtusè tetragona plùs minùs carnosà obtusà, apice umbilicatà calvā glabrā aut apice summo velutino-villosa. — *Herbæ Austr.-Americane.* Caules erecti aut subsarmentosi sæpius scabro-asperī sæpè (fortè semper) maculati. Folia opposita petiolatà latè ovata acuminatà serratà, nervis ad basin folii utrinquè 3-pinnatis tertio majore et indè folia pseudo-triplinervia. Capitula pedicellatà, terminalia aut axillaria, sæpè ternà, post anthesin subglobosa. Flores flavi. — Genus admodum naturale!

\* *Capitulis radiatis.*

1. *W. MACULATA*, caule erecto tetragono vix subscabro maculato, foliis oblongo-lanceolatis basi cuneatis apice acuminatis serratis supernè scabris subtus glabrisculis, pedunculis ternis terminalibus, ligulis circ. 8 involuero duplò longioribus.  $\frac{1}{2}$  in Brasilià (Ker), circa Rio de Janeiro (Lund!). *Gymnoloma maculatum* Ker bot. reg. t. 662. Petioli 3-4 lin. longi. Flores aurantiaco-flavi. (v. s.)
2. *W. STENOGLOSSA*, caule erecto subsarmentoso tetragono scabrido submaculato, foliis latè ovatis acuminatis basi vix cuneatis serratis utrinquè scabris asperimis membranaceis, pedunculis ternis ad apices ramorum et ramulorum axillariorum, ligulis 8-10 invol. duplò longioribus, invol. squamis ext. ovalibus adpressè hirtellis.  $\frac{1}{2}$  in Cayennâ. *Verbesina membranifolia* Poir. suppl. 3. p. 18. *Helianthus sarmentosus* Rich. act. soc. h. nat. Par. p. 105. Pers. ench. 2. p. 475. n. 9. *Verbesina oppositiflora* Poir. l. dict. 8. p. 460. *Chakiatella stenoglossa* Cass. dict. 29. p. 490 et 46. p. 406. *Chylodia sarmentosa* Rich. ined. ex *Cass.* *Coreopsis baccata* Linn. f. suppl. 380? *Pascalina baccata* Spreng. syst. 3. p. 602? Petioli 7-8 lin. longi. Ligulæ angustæ pallidè flavæ. (v. s.)
3. *W. PLATYGLOSSA*, caule erecto striato-sulcato adpressè hirtello, foliis petiolatis latè ovali-lanceolatis serratis utrinquè scabridis, capitulis ternis terminalibus pedicellatis, pedicello medio breviorè, invol. squamis ext. ovatis foliaceis, int. paleiformibus, paleis oblongo-lanceolatis achænia amplect., ligulis paucis invol. vix longioribus.  $\frac{1}{2}$  in Guianâ gallicâ legit cl. Martin. *Crodisperma aspera* Poit. in h. Gay. ex Cass. *Chakiatella platyglossa* Cass. dict. 46. p. 403. Cor. flavæ. Ligulæ luteæ plurinerviæ. Folia petiolo excepto 5 poll. longa. 3 poll. lata. Stirps à cl. Poiteau lecta et à cl. Cassinio descripta tantum à stirpe Martini differt foliis paulò angustioribus, obs. cl. Ad. Brongniart. (v. s. comm. à Mus. reg. Par.)
4. *W. HAVANENSIS*, caule hexagono erecto scabro, foliis petiolatis latè ovatis acuminatis basi obtusis subcordatis callosè et remotè dentatis utrinquè asperis, nervis 3 inferioribus approximatis, capitulis ad apices ramorum et ramulorum axillariorum ternis, invol. squamis ext. lanceolatis paleisque apice subuncinatis, ligulis 9-12.  $\frac{1}{2}$  in insulâ Cnbâ juxta Havanam legit cl. Ant. de La Ossa. Petioli 5 lin. longi. Limb. fol. 5-6 poll. longus, 2  $\frac{1}{2}$ -3 poll. latus. Ligulæ pallidè flavæ. (v. s.)
5. *W. SALZMANNI*, caule tereti striato subscabro, foliis breviter petiolatis ovatis acuminatis basi obtusis serratis utrinquè scabris, nervis 3 infer. approximatis, capitulis ad apices ramorum ternis medio brevius pedicellato, invol. squamis ob-

longis obtusiusculis, paleis oblongis glabris subcoloratis, ligulis 7-10 disco paulo longioribus,  $\frac{1}{2}$  in umbrosis circa Bahiam Brasiliæ legit. cl. Salzmann qui fructum verè baccatum in vivo vidit. Caulis 4-pedalis. Petioli vix 3 lin. longi. (v. s.)

6. *W. OBLONGIFOLIA*, caule hexagono erecto subpubero, foliis petiolatis oblongo-lanceolatis acuminatis serratis basi acutis supra sparse puberis subtus molliter villosis, capitulis terminalibus ternis, paleis rigidis apice uncinato-recurvis, ligula parvis paucis albidis.  $\frac{1}{2}$  circa Rio-Janeiro legit fructiferam cl. Gaudichaud et floridam ni fallor vidi in h. Mus. imp. Bras. sub. n. 120. ex prov. Mato-Gross. Petioli 3 lin. longi. Limb. fol. 2 poll. longus, 7 lin. latus. (v. s.)

\*\* *Capitulis discoideis.*

7. *W. BLANCHETII*, caule teretiusculo scabro, ramis subangulatis, foliis brevissimè petiolatis ovato-lanceolatis acuminatis serratis utrinquè scabris basi obtusis aut subcordatis, capitulis terminalibus ternis breviter pedicellatis, invol. squamis lanceolatis subscabris, ligulis nullis.  $\frac{1}{2}$  in Brasiliâ circa Bahiam legit cl. Blanchet. Petioli 1-2 lin. longi. Limb. fol. 2  $\frac{1}{2}$  poll. longus, pollicem latus. Cor. intensè lateæ. (v. s.)

8. *W. SCANDENS*, caule scandente subhexagono subaspero, foliis petiolatis ovatis valdè acuminatis subserrulatis utrinquè scabris basi obtusis, capitulis ad apicem ramorum pedicellatis, invol. squamis ovalibus subacutis, paleis acutis rectis, ligulis nullis.  $\frac{1}{2}$  in Peruvâ legit cl. Poeppig. Cæsulia scandens Poepp. ! pl. exs. n. 24. diar. 1663. Petioli 9 lin. longi. Limb. fol. 4 poll. longus, 20 lin. latus. Cor. intensè flava. Fructum non vidi. (v. s.)

CCCXXIX. MONTAGNÆA. — *Eriocoma* H. B. et Kunth nov. gen. am. (1820) 4. p. 267. t. 396. Cass. dict. 15. p. 193. Less. syn. 226. non Nutt. (1818). — *Montanoa* Lallav. et Lex. nov. veg. mex. 2. (1825) p. 11. Less. syn. 490. — *Eriocarpha* Cass. dict. 59. (1829) p. 236. — *Priestleya* fl. mex. ic. ined. non DC.

Capitulum multiflorum heterogamum, fl. radii 5-10 uniseriatis ligulatis neutris, disci tubulosi 5-dentati hermaphrod. Invol. biserialè, squamis ext. 5 oblongis patulis, inter. sæpius 10 paleas referentibus. Recept. convexum, paleis basi latis apice spinoso-mucronatis achæna plus minus involventibus. Styli disci rami cono nunc brevè nunc longiusculo superati. Achænia radii abortiva, disci subcompressa obovata cuneata calva, nectario interdum prominulo. — Suffrutesc frutescive Americani plerique Mexicani. Rami teretes apice sæpius hirsuti. Folia opposita petiolata, ovata aut cordata, serrata aut lobata, sæpius subtus subtomentosa. Capitula corymbosa. Ligulæ albæ (aut fortè roseæ). Discus albidus. — Nomen paulò emendavi ut melius cum nomine cl. doct. Lud. Montanæ (pronunt. Montagna) nec Montano congruat. Genus affine *Sclerocarpo*.

§. 1. *Eriocarphæ, ligulis 5, paleis hirsutissimis.*

1. *M. FLORIBUNDA*, caule herbaceo? oppositè ramoso villosopubescente, foliis oppositis petiolatis deltoideo-ovatis propè basin triplinerviis subserratis supra scabrisculis subtus hirsutopubescentibus et canescentibus, corymbis fastigiatis polycephalis capitulis longè pedicellatis, invol. squamis biserialibus oblongo-lanceolatis obtusiusculis, paleis ovatis acuminatis albo-lanatis, ligulis 5.  $\frac{1}{2}$  in Mexico inter urbem et sanctuarium Guadalupense. *Eriocoma floribunda* H. B. et Kunth nov. gen. am. 4. p. 267. t. 396.
2. *M. TOMENTOSA* (Lallav. et Lex. nov. veg. mex. 2. p. 11), caule fruticosopubescente villosotomentoso, foliis oppositis petiolatis latè ovato-lanceolatis subserratis apice acuminatis sinuatis incisive prope basin triplinerviis supra subpubescentibus subtus petiolisque villosotomentosis, corymbo composito fastigiato, invol. squamis, 5 ext. oblongis subobtusis dorso villosis disco sub-

æqualibus, inter. paleisque dorso hirsutissimis in acumen rectum spinulosum productis, ligulis 5.  $\frac{3}{2}$  in Mexico (Alam.), circa urbem Mexici ad ripas fossarum (Lallav.), ad Real del Monte (Nées), ad Tlapujahu (Keerl. in h. Mart.). Vulgò *Cihnapatl* Mexic. dicitur. *Priestleya corymbosa* fl. mex. ic. ined. *Eriocoma fragrans* Don in Sweet br. fl. gard. ser. 2. t. 44. *Eriocoma heterophylla* Schrad. sem. h. Goett. 1833. p. 3. (v. s.)

ℓ. *cordifolia*, foliis basi latè et profundè cordatis, petiolis hinc iudè alato-appendiculatis  $\frac{3}{2}$  in horto Mexicano legit cl. Berlandier (pl. exs. n. 574!). Folia supernè glabriuscula, subtùs minùs tomentosa, magn. ferè dupla. (v. s.)

γ. *ternifolia*, foliis terno-verticillatis oblongis basi obtusis apice acuminatis, parèe dentatis.  $\frac{5}{2}$  in Mexico ad Chapolopa legit cl. Berlandier (pl. exs. n. 574!). Unicum specim. habeo et floribus junioribus donatum, undè dubius hæreo an sit spec. aut var. (v. s.)

§. 2. *Acanthocarpæ*, *ligulis circ. 10. paleis parèe villosis evidentiùs mucronatis demùm spinescentibus.*

3. *M. serotina*, caule herbaceo striato subangulato glabro ad apices subvillosò, foliis oppositis sessilibus basi in petiolum alatum crispum subsinuatum attenuatis longè supra basin triplinerviis irregulariter pinnatilobatis (lobis acuminatis subintegris) suprà villosis subtùs molliter et densè villosò-tomentosis, ramis oppositis paniculatis nudis 1-cephalis, ligulis 10 elongatis, paleis recept. convexi basi latis apice spinescenti-subulatis  $\frac{2}{2}$ ? in Mexico ad Cuernavaca octob. flor. legit cl. Berlandier (pl. exs. n. 1057). Folia 8-9 poll. longa, 3 poll. lata. Capitula diam. sesquipollic. (v. s.)

4. *M. grandiflora*, suffruticosa oppositè ramosa, ramis apice pubescentibus, foliis in petiolum alatum dentatum attenuatis longè suprà basin triplinerviis suprà strigilloso-scabris subtùs puberulis, infer. irregulariter pinnatifidis, lobis utrinquè 2-3 lato-lanceolatis subobtusis, summis ovato-lanceolatis indivisis, capitulis ad apices ramorum ternis pedicellatis, invol. squamis lineari-lanceolatis reflexis, paleis parèe villosis basi latis in spinulam productis, lignis invol. longioribus circ. 10.  $\frac{5}{2}$  in Mexico. *Eriocoma grandiflora* Alam.! in litt. 1831. *Priestleya longifolia* fl. mex. ic. ined.? *Ligulæ* pollicem longæ apice grossè 2-3 dentatæ. (v. s. comm. à cl. Alam.)

5. *M. Karwinskii*, caule herbaceo? tereti glabriusculo, foliis oppositis petiolatis ovatis obtusè sinuatis basi 3-5-nerviis aut subtriplinerviis parciatimè pubescentibus, paniculis laxis polyccephalis, invol. squamis lineari-lanceolatis acuminatis inæqualibus, ligulis 10 obovato-oblongis  $\frac{2}{2}$ ? in Mexico legit cl. de Karvinski. A duobus prioribus differt petiolis evidentiùs omninò immarginatis, à sequentibus foliis sinuatis seu obtusè lobatis. *Ligulæ* 5-6 lin. longæ, 2-3 lin. latæ. (v. s.)

6. *M. frutescens* (Maire! in litt. 1833), caule suffruticoso glabro, foliis oppositis et super. alternis breviter petiolatis lanceolatis acutis basi in petiolum non productis triplinerviis subserratis suprà asperis subtùs glabris, capitulis laxè corymbosis, invol. uniseriatis squamis 5 obtusis disco brevioribus, paleis recept. glabris demùm convolutis apice recurvo rigido achænium involv. et multò superantibus, ligulis 8-10 invol. longioribus.  $\frac{3}{2}$  in Mexico (Alam.! Maire!), circa urbem (Berl.! pl. exs. n. 908), et in valle Toluacanà (Berl.! n. 1154). *Priestleya squarrosa* fl. mex. ic. ined. *Eriocoma frutescens* Alam.! in litt. *Ligulæ* 6 lin. longæ. Paleæ per anthesin breves, dein longè accretæ et capitulum fructiferum squarrosam formantes. (v. s.)

7. *M. arborescens*, caule fruticoso oppositè ramoso, ramis pubescenti-villosis, foliis oppositis petiolatis lanceolatis basi acutis apice acuminatis subintegerrimis subtriplinerviis suprà asperis subtùs pubero-hirtellis, corymbis laxis trifidis, capitulis longè pedicellatis, invol. squamis biseriatis oblongis disco ferè æqualibus, paleis parèe villosis mucronatis demùm apice induratis inflexis, ligulis circ. 8.  $\frac{3}{2}$  in Mexico (Alam.! Maire!), ad Cordilleram Cuchilaqua (Berl.!

pl. exs. n. 1006). *Eriocoma arborescens* Alam. in litt. 1831. *Montanoa floribunda* Cerv. in h. acad. Monac. Ligulæ 7-8 lin. longæ, 3 lin. latæ obtusæ. (v. s.)

8. *M. OVALIFOLIA*, caule fruticoso subsicante ad apices ramorum molliter hirsuto, foliis oppositis petiolatis ovalibus ovatisve acutis propè basin triplinerviæ parçè serratis suprâ petiolisque sparsè villosis subtus villosa-tomentosis cæscenscentibus, capitulis corymbosis longè pedicellatis, invol. squamis ext. oblongis hirsutis subreflexis, inter. paleisque parçè villosis in acumen spinulosum productis, ligulis circ. 10.  $\frac{3}{4}$  ad Santa-Fé de Bogota. Ligulæ ex sicco fortè roseæ, 5-6 lin. longæ. (v. s. comm. ab amic. Benj. Delessert.)

CCCCXX. *SCLEROCARPUS* Jacq. act. helv. v. 9. (1786) p. 34. t. 2. f. 1. Cass. dict. 48. p. 148. Thonn. in Schum. pl. guin. 394.

Capitulum multiflorum heterogamum, fl. radii 3 neutris minimis irregulariter ligulatis ante squamas inv. interiores sitis, disci 8-12 tubulos hermaphroditis, centralibus sæpè abortivis. Invol. duplex, squamis ext. foliaceis subpetiolatis majoribus (fortè bracteis), int. 5 minoribus distantibus. Recept. parvum convexum, paleis disci flores involventibus et iis brevioribus. Cor. disci pubescens 5-dentata persistens. Anth. vix concretæ apice in appendicem subulatam productæ. Styli rami subulati hispidi. Achænia radii abortiva tenuissima, disci obovata levia calva intrâ paleam curvato-subcomplicatam induratum intus hiantem apice in collum contractam inclusa. — Herba Africana erecta. Caulis scaber teres apice præsertim ramosus. Folia alterna petiolata ovata acuminata serrata triplinervia suprâ villosa scabra subtus puberula. Capitula ad apices ramulorum solitaria subsessilia, Cor. flavæ.

1. *S. AFRICANUS* (Jacq. ic. rar. 1. t. 176), foliis latè ovatis acutis serratis. (In Guineâ (Thonn.), et in cultis inundatis regionis Walo ad flum. Senegal (Perrotet!); etiâ sed verisim. introductum in Indiâ orient. legit cl. Royle et cl. Baranquillas Americæ austr. cl. Bertero. Hic ultimus est *Melampodium Banzquillæ* Spreng. syst. 3. p. 619! (v. s. sp. et v. c.)

† *Species valdè dubiæ.*

2. *S. ? GRACILIS* (Smith in Rees cycl. v. 31), foliis lineari-lanceolatis integris, pedunculis 4-5 ad apices caulis foliosis. — ad oram boreali-occid. Americæ septentrionalis. Paleæ fructiferæ glandulis prominentibus onustæ. Cor. ign. et plerique car. valdè incerti. †.
3. *S. ? EXIGUUS* (Smith l. c.), foliis lineari-lanceolatis integris, pedunculis natis folio triplò longioribus. († cum priore. Herba 3-4 pollic. Folia ferè *Myosotis*. Flores et fructus vix noti. †.

CCCCXXI. *ENCELIA* Adans. fam. 2. p. 128. Cass. dict. 14. p. 447. 59. p. 119. Less. syn. 227. — *Pallasia* L'Hér. diss. 1784. Ait. kew. ed. 1. v. 3. p. 253 non Linn. — *Coreopisid* sp. Jacq.

Capitulum multiflorum radiatum, ligulis 1-serialibus neutris, fl. disci hermaphr. 5-dentatis. Invol. biseriatum disco æquale subimbricatum. Recept. planum (nec conicum ut vult Less.), paleis membranaceis navicularibus. Styli rami cono superati. Achænium plano-compressum margine albo-lanatum emarginatum calvum totum villorum. — Suffrutescentia Americani ramosi plus minus, saltè juniores, cæcescentes. Folia alterna petiolata ovata aut oblonga tri- seu triplinervia integra aut subrepanda. Capitula pauca pedicellata in paniculam disposita. Cor. flavæ.

1. *E. CANESCENS* (Cav. ic. 1. p. 45. t. 61), erecta, foliis latè ovatis integris obtusis mollior canescentibus, invol. aquamis villosis ciliatis.  $\S$  in Peruvîâ (Adans. Gaud.!), circa Limam (Domb.!), nec verisim. in Mexico (ut dixit Cavanilles ex hort. Madr.) Lindl. bot. reg. t. 909. Kunth nov. gen. am. 4. p. 206. *Coreopsis Limensis* Jacq. ic. rar. 3. t. 594. *Pallasia halimifolia* L'Hér. ! herb. et mus. Willd. sp. 3. p. 2260. *E. alternifolia* et fortè *Eucalia canescens* Ræusch. nom. 251? (v. v. c. et a. sp.)
2. *E. PARVIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 206), procumbens, ramulis sericeo-pubescentibus, foliis rhombeo-ovatis scutis basi-cuneatis canescenti-pubescentibus.  $\S$  in Peruvîâ ad littora Oceani pacifici. *Pallasia procumbens* Spreng. syst. 3. p. 610. Frutex humilis ramosissimus proximus *E. canescenti*.
3. *E. HALIMIFOLIA* (Cav. ic. 3. p. 6. t. 210), erecta, foliis ovatis subrepandis dentatisve glabriusculis margine valdè ciliatis.  $\S$  in Mexico ex Cav., sed verisim. potius in Peruvîâ. Cass. dict. 14. p. 448. *Pallasia grandiflora* Willd. sp. 3. p. 2261. (v. v. c. in h. Par.)
4. *E. OBLONGIFOLIA*, erecta, foliis oblongis utrinquè subattenuatis integerrimis canescentibus, invol. basi præsertim hirsutissimis.  $\S$  in Chili (Hænk.!), circa Coquimbo (Gaudichaud! Macrae.!). Ligulæ elongatæ lineares. Folia var. 1-2 pollic. longa, 4-8 lin. lata. (v. s.)

† *Species exclusæ.*

*E. DENTATA* Poir. = *Verbesina dentata* ?

*E. MEXICANA* Mart. = *Simaia auriculata*.

CCCXXXII. PHILOGLOSSÆ.

**Capitulum** multiflorum heterogamum, fl. radii ligulatis innumeris 1-2-? seriatis neutris, disci tubulosi hermaphroditis ad medium 5-fidis. Invol. 4-5-seriale, squamis ext. inter se alternis biseriatis lanceolatis foliaceis ciliatis, inter. 3-ser. membranaceis albidis glabris lanceolatis. Recept. planum, paleis lineari-subulatis scabridis caducissimis. Ligulæ basi hirtæ angustissimæ. Cor. disci tubo brevi, lobis 5 acuminatis profundis. Anth. nigricantes. Stylus apice summo subbifidus longè hispidus. Achænia obovata lævia calva, areolâ terminali minimâ. — *Herba Limensis* villosa-hispida. Caulis adscendens. Folia caulina opposita, ramorum flor. quaterno-verticillata, ovalia aut obovata basi cuneata sessilia tenuissimè triplinervia integra aut subdentata. Pedicelli filiformes hispidi axillares terminalesque 1-cephali, anthesi ineunte folio breviores, demùm longiores. Flores lutei ferè calendulacei.

1. *P. PERUVIANA*. (L.)? in Peruvîâ circa Limam legit cl. Dombey et recentius cl. Abadia. (v. s.)

CCCXXXIII. CHRYSOSTEMMA *Less. syn.* 227. — *Coreopsis* sp. *Linn.* —  
*Anacis* sp. *Schrank.*

**Capitulum** multiflorum heterogamum, fl. radii neutris 1-serialibus ligulatis, disci hermaphr. tubulosi 5-dentatis. Invol. biseriale, squamis foliaceis basi coadunatis, ext. paucis parvis linearibus patentibus, inter. erectis ovali-oblongis margine subscariosis. Recept. planum, paleis linearibus angustissimis ferè filiformibus. Styli rami cono superati pubescente descendente instructi. Achænium plano-obcompressum obovato-ellipticum angulis alatum glaberrimum apice emarginatum, pappo coroniformi lacero coronatum, disco epigyno minuto, alis albidis. — *Herba Boreali-Americana* nitida erecta. Folia opposita petiolata, ima

subpedato-pinnatifida, super. trisecta, segmentis lanceolatis integerrimis. Pedunculi subgemini 1-cephali. Capitula radio luteo, disco brunneo.

1. *C. TRIPTERIS* (Less. l. c.),  $\frac{1}{2}$  in America bor. montibus altis à Virginià ad Carolinam (Pursh), et in Georgià occid. (Ell.). *Coreopsis tripteris* Linn. sp. 1282. Michx. fl. bor. am. 2. p. 138. Ell. sketch 2. p. 447. *Anacis triptero* Schrank. in akad. Münch math. et nat. 5. p. 7. — *Moris. oxon.* 3. s. 6. l. 3. f. 44. (v. v. c.)

CCCXXXIV. *CALLIOPSIS* Reichenb. mag. n. 70. Cass. dict. 59. p. 321. Lem. syn. 228. — *Diplosastera Tausch hort. Can. fasc. 1.* — *Calliopsisidis* sp. Spreng. *Coreopsisidis* sp. Nutt.

Capitulum multiflorum heterogamum, fl. radii neutris 1-serialibus ligulatis grosse 3-5-dentatis, fl. disci hermaphr. tubulosis 5-dentatis. Invol. biseriale, squamis ext. brevibus squarrosis, int. majoribus erectis coadunatis. Recept. planum paleis linearibus deciduis floribus brevioribus onustum. Stylus disci ramis truncatis apiceque solo penicillatis. Achænia obcompressa calva incurva truncata, disco epigyno minuto. — Herba Boreali-Americana elegans glabræ. Folia opposita uni- aut bipinnatisecta, lobis integris. Capitula ramos terminantia, radio luteo basi fusco-purpureo maculato, disco luteo aut subpurpureo. — Genus vix à *Coreopside* differt.

1. *C. TINCTORIA*, glabra, foliis radicalibus pinnatipartitis seu bipinnatipartitis lobis integerrimis, superioribus tripartito-multifidis lobis linearibus, sup. ext. brevissimo acuto, ligulis trifidis, invol. int. duplò longioribus, achæniis oblongis exalatis utrinquè minutè tuberculatis. (1) in territorio Arkansas ad ripas flum. Rubri in pratis inundatis detexit cl. Nuttall. *Coreopsis tinctoria* Nutt. journ. acad. Philad. 1821 oct. p. 114. Radius schrif. ges. Leips. 181. p. 159. t. 3. Bart. fl. am. bor. 2. t. 45. ex bot. mag. t. 2512. bot. reg. t. 84. Link et Otto abb. t. 33. Sweet fl. gard. t. 72. *Diplosastera tinctoria* Tausch hort. canal. fasc. 1. icon et descr. *Calliopsis bicolor* Reichenb. mag. t. 70. ligulæ basi fuscae cæterum aureæ. Herba elegans in hortis ubiquè culta et jam hinc inde in Europâ seminibus hortensibus sata. (v. v. c.)
2. *C. CARDAMINEFOLIA*, foliis infer. uni- aut bipinnatisectis lobis obovatis oblongis sæpè trifidis, super. bipinnatisectis lobis lineari-spathulatis, invol. ext. brevissimo obtusiusculo, ligulis trifidis invol. interiore duplò longioribus, achæniis alatis utrinquè lævibus aut minutissimè tuberculatis. (1) in Meco prov. Texas inter Bejar et flum. Trinitatis, ad Matamoros et ad lacum Sancti-Nicolai in sina Sancti-Spiritus legit cl. Berlandier (pl. exs. n. 1668, 1936, 1478, et 2312). Ligulæ basi fuscae apice aureæ ut in *C. tinctoriâ* ad quam accedit, sed achæniis diversa. (v. s. sp.)
3. *C. ATKINSONIANA* (Hook. fl. bor. am. 1. p. 311), foliis radicalibus bipinnatisectis lobis integerrimis, caulinis pinnatisectis lobis linearibus subspathulatis, invol. ext. squamis lineari-oblongis margine submembranaceis, ligulis invol. int. triplò longioribus obtusè 3 dentatis, achæniis alâ brevi marginatis utrinquè lævissimis.  $\frac{1}{2}$  ad ripas fluminis Colombiæ Amer. bor. et insul. Merries s. Menzies, legit cl. Douglas. *Coreopsis Atkinsoniana* Dougl. in Lindl. bot. reg. t. 1376. Similis *C. tinctoriæ* sed elatior, perennis, et achæniis distinctissima. (v. v. c.)

CCCXXXV. *PERAMIBUS* Raf. ann. nat. 1 (1820), p. 14. Cass. dict. 38. p. 416. — *Coreopsisidis* sp. Nutt. — *Calliopsisidis* sp. Spreng.

Omnia *Coreopsisidis* sed achænia triangularia lævia nuda. — Genus valdè incertum, soli Rafinesquio notum et botanicis Americanis commendatum. †.



1. *P. hirtus* (Raf. l. c.), hirtus, caule angulato flexuoso ferè dichotomo, foliis æmiamplexicaulibus lanceolatis acuminatis medio serratis, inferioribus ovato-lanceolatis, floribus ferè corymbois subsessilibus, invol. squamis lineari-lanceolatis acutis, ligulis ellipticis. 2/ in collibus ad flum. Liking in Kentucky. *Coreopsis hirta* Raf. ex ipso. †.
2. *P. scabra* (Raf. l. c.), ramis remotis elongatis subulatis foliisque imis oppositis, summis alternis, foliis integris (lanceolatis?) scaberrimis, achæniis oblongis angulatis. 2/ in Louisianâ. *Coreopsis scabra* Raf. fl. lud. 72. Herba bipedalis. An eadem ac sequens ex Raf.? †.
3. *P. acutus* (Raf. l. c.), foliis ovato-lanceolatis acutis denticulatis subhirtis, capitulis corymboso-paniculatis. ② in Georgiâ. *Coreopsis acuta* Pursh fl. bor. am. 2. p. 569. Cæet. ign. †.

Div. III. COREOPSIDÆ *Less. syn. p. 228.* — *Coreopsis* et *Heliantheæ* prototypæ *Cass. opusc. 3. p. 58 et 59.*

Capitula heterogama, fl. radii neutris. Achænia crostris pappo nunc aristis 2-4 tantum, nunc aristis et squamellis constante.

CCCXXXVI. AGARISTA.

Capitulum multiflorum radiatum, ligulis 1-seriatis neutris obovato-truncatis multinerviis, fl. disci tubulosis 5-dentatis hermaphrod. Invol. campanulatum duplex, squamis acutis, ext. 4 latè ovatis basi subconcretis, inter. 8 ovali-oblongis duplò ferè longioribus submembranaceis. Recept. paleæ cum fructu deciduæ, ad latus hujus ext. applicatæ, lineari-lanceolatæ subhyalinæ. Styli rami cono superati. Achænia obcompressa, radii ovalia inania glabra calva, disci oblonga sub paleâ glabra, margine utroque et facie interiore villosissima. Pappus bisquamellatus, squamellis elongatis membranaceis corollâ parùm brevioribus ex angulis achæniis ortis. — Herba Californica glaberrima annua erecta *Calliopsidis* seu *Coreopsidis* facie. Caulis teres ramosus. Folia alterna multifida rachi lobisque linearibus elongatis. Rami apice aphylli 1-cephali. Capitula erecta aurea. Genus differt ab *Heliantheis* ferè omnibus pappo ut in *Helenieis* squamellato nec aristato, sed tamen affinitate naturali ad hunc locum revocatum. — *Agarista* mythologicè erat *nympha* venustissima.

1. *A. calliopsidea*. ① in Californiâ legit cl. Douglas. Ligulæ circ. 8 basi cuneatæ, 8 lin. longæ, 4 lin. latæ, lineis 15-20 striatulæ. (v. s. comm. ab hon. soc. hortie. Lond.)

CCCXXXVII. COREOPSIS *Linn. gen. n. 981. (excl. spec.) Less. syn. 228.* — *Coreopsis* et *Coreopoides* *Mench meth. p. 593 et 594.* — *Coreopsis* et *Leachea* *Cass. dict. 10. p. 419. 25. p. 388 et 59. p. 320 et 321.* — *Coreopsis* et *Anacis* sp. *Schrank.* — *Coreopsis* et *Chrysomelea* *Tausch.* — *Coreopsis* et *Aspersum* *Neck. elem. n. 63 et 64.*

Capitulum multiflorum heterogamum, fl. radii circ. 8 neutris ligulatis, disci hermaphr. 5-dentatis. Invol. duplex utrumque circ. 8-phyllum imâ basi subcoadunatum, squamis ext. foliaceis angustioribus subpatulis, int. latioribus submembranaceis. Recept. planum paleaceum, paleis persistentibus. Styli rami apice hispidi cono superati. Achænia obcompressa-plana, rarò incurva, margine plùs minùs alata, apice nunc bidentata, nunc biaristata, aristis lævibus aut sursùm ciliato-serratis (nec retrorsum pilosis). — Herbæ Americanæ rarò suffrutescentes ple-

rumquē glabræ. Folia opposita, rariùs alterna. Pedunculi 1-cephali solitarii aut corymbosi. Capituli discus luteus, radius concolor rari albus.

\* *Foliis alternis.*

1. *C. ANGUSTIFOLIA* (Ait. kew. 3. p. 253), glabra, foliis alternis lineari-lanceolatis integerrimis lævibus, ligulis oblongis trifidis, laciniâ mediâ majore.  $\frac{1}{2}$  in Carolinâ et Floridâ. Ait. (v. s. ?) †.
2. *C. GLADIATA* (Walt. car. p. 215), caule glabro supernè dichotomo, foliis alternis in petiolum semiamplexicaulem attenuatis angustis aut ovali-lanceolatis puberulis glabrisve integerrimis, invol. squamis exter. lineari-lanceolatis inter. brevioribus, ligulis 3-dentatis, achæniis obovatis alatis, alis serrulatis, pappo bisetoso.  $\frac{1}{2}$  in pinetis humidis et sphagnosis umbrosis à Novâ-Cæsariâ ad Carolinam (Pursh). *C. gladiata* Nutt. gen. am. 2. p. 180. Ell. sketch 1. p. 444. *C. dichotoma* Michx. fl. bor. am. 2. p. 137. Variat foliis angustioribus pilosis et latioribus glabris. (v. s. sp.)
3. *C. ASPERA* (Pursh fl. bor. am. 2. p. 570), foliis lanceolato-linearibus asperis, superioribus alternis, inferioribus oppositis, caule monocephalo. — in Maryland. Cæc. ign.

\*\* *Foliis oppositis indivisis.*

4. *C. LINEARIFOLIA*, caule suffruticoso adscendente glabro simplici basi folioso in pedunculum nudum 1-cephalum longè productum, foliis oppositis linearibus integerrimis basi parçè ciliatis, invol. ext. squamis linearibus glabris, int. lanceolatis hirsutis, ligulis ellipticis multiradiatis nervulis alternis brevioribus, styli ramis cono acuto superatis.  $\frac{1}{2}$   $\frac{3}{4}$  in Mexico ad Tlapujahu legiti cl. Kent ex h. Mart! Species distinctissima, vix pedalis. Ligulæ intense flavæ nigri striatæ. Discus fuscus. Achænia ex statu immaturo omninò calva ut in *Caliopside* sed styli rami cono ut in *Coreopside* superati. (v. s. in h. Mart.)
5. *C. VENUSTA* (H. B. et Kunth nov. gen. am. 4. p. 229), fruticosa glabra, foliis oppositis linearibus acutiusculis integerrimis uninerviis subcoriaceis, capitula terminalibus solitariis longè pedunculatis, invol. squamis exter. oblongis obtusis patentibus, ligulis oblongis subtridentatis, achæniis (junioribus) marginè albo-lanatis biaristatis, aristis adpresso-pilosis.  $\frac{3}{4}$  prope Loxam et in scopulis Parami de Saragura. Ligulæ aurantiacæ, discus flavus.
6. *C. LANCEOLATA* (Linn. sp. 1283), caule tereti striato, foliis oppositis connatis seu oblongo-lanceolatis integerrimis basi ciliatis, infer. petiolatis, pedunculis longissimis simplicibus nudis, invol. squamis exter. lanceolatis interioribus subæquantibus, ligulis apice 5-dentatis, achæniis orbiculatis alatis scaberrimis biaristulatis.  $\frac{1}{2}$  in Carolinâ (Linn.) et Virginiâ (Pursh), ad lacum Superiorem et Huronensem (Hook.). *Leachea lanceolata* et verisim. *L. crassifolia* Cass. dict. 25. p. 388. *Chrysomelea lanceolata* Tausch hort. Canal. f. 1.  
*a. succisæfolia*, caule erectiusculo ramoso foliisque glabris, ligulis amplioribus. — Hæc varietas hortensis videtur typus speciei et generis. Dill. elub. 55. t. 48. f. 56. Linn. hort. cliff. 420. Ligulæ amplæ 10-12 lin. longæ. (v. s. c.)
- ß. *glabella* (Michx. fl. bor. am. 2. p. 137), caule abbreviato glabro, foliis margine ciliatis utrinquē glabriusculis angustioribus, ligulis mediocribus.  $\frac{1}{2}$  in humidis Carolinæ. Foliorum limbus circ. pollicem longus, petiolo ferè æqualis. Ligulæ 6-7 lin. longæ. (v. s. sp.)
- γ. *villosa* (Michx. l. c.), caule abbreviato hirsutissimo, foliis utrinquē canescenti-villosis, pedunculis glabris.  $\frac{1}{2}$  in pinetis et locis siccioribus et Ell. *Coreopsis crassifolia* Ait. kew. 3. p. 253. Ell. sketch 2. p. 434. (v. s. comm. à cl. B. Greene.)
7. *C. INTEGRIFOLIA* (Poir. suppl. 2 (1811), p. 352), caule glabro tereti simpliciter, foliis oppositis ovatis integris utrinquē acutis glabriusculis, infer. petiolatis imâ basi connatis, summis sessilibus, capitalis panicis trichotomocorymbosis, invol. ext. brevi subpatente, ligulis trifidis, achæniis breviter

bistaratis. — in Carolinā legit cl. Bosc. Coreopsis bidens Boeck herb. non Walt. Capitula lutea. Habitus Bidentis sed achæniis aristæ non retrorsum scaræ. Ovaria juniora tantum vidi. (v. s.)

8. *C. ORMLERI* (Ell. sketch 2. p. 435), glabra, caule angulato, foliis oppositis sessilibus lato-lanceolatis utrinque acutis integerrimis, pedunculis axillaribus terminalibusque subdichotomè corymbosis, invol. ext. brevi, ligulis integris, achæniis cuneatis marginatis leviter dentatis. — in Carolinā ad junctionem flum. Broad et Saluda. Affinis dicitur *C. latifolia*.
9. *C. LATIFOLIA* (Michx. fl. bor. am. 2. p. 137), foliis oppositis ovatis acumina-  
tis breviter petiolatis dentatis, dentibus mucronatis, ligulis integris, achæniis cuneato-oblongis apteris apice nudis.  $\frac{1}{2}$  in montibus altis Carolinæ. Habitus Silphii sed capitula minora. — An Silphium glaucum Weinm. hort. dorp. 143? sed à Coreopside et à Silphio alienum esset si recept. verè nudum ut vult Weinmann.
10. *C. ARGUTA* (Pursh fl. bor. am. 2. p. 567), glabra, foliis oppositis petiolatis lanceolato-ovatis sensim attenuatis argutè serratis, pedunculis axillaribus terminalibusque dichotomè corymbosis.  $\frac{1}{2}$  in Carolinā ex Pursh, sed ab Elliott. non reperta. An ex Nutt. var. *C. latifolia*? Flores aurei. *C. serrulata* Poir. suppl. 2. p. 352 huc à Spreng. admissa, videtur potius *Bidens grandiflora*.

\*\*\* *Foliis oppositis partitis sectivis.*

11. *C. MUTICA*, suffruticulosa glabra, foliis oppositis serratis, aliis ovato-lanceolatis acuminatis serratis, aliis trisectis, segmento medio folia indivisa referente, lateralibus minoribus, capitulis ad apices ramorum pedicellatis ternis, invol. squamis ext. oblongis, ligulis multistriatis, achæniis elliptico-oblongis submarginatis apice vix ac ne vix obtusè bidentatis.  $\frac{3}{4}$  in Mexico ad Tlapujahua legit F. W. Keerl ex h. Mart. Flores lutei. Paleæ membranaceæ striatæ. (v. s. in h. Mart.)

$\beta$ . *subvillosa*, ramulis foliis subtus involucrique squamis exterioribus villosis. — in Mexico legit cl. de Karvinski. (v. s. in h. Acad. reg. Monac.)

12. *C. AURICULATA* (Linn. sp. 1281), caule erecto ramoso breviter puberulo, foliis oppositis aliis indivisis aliis trisectis, lobis lanceolatis acutis integerrimis utrinque brevissimè puberulis, terminali triplo quadruplè majore, petiolo ciliato, capitulis subsolitariis pedicellatis, invol. squamis ext. lineari-oblongis subacutis glabris demum reflexis, ligulis 4-5-dentatis, achæniis suborbiculatis alatis brevissimè bistaratis.  $\frac{1}{2}$  in Amer. bor. montibus Virginie et Carolinæ. Schkuhr handb. 3. t. 260 opt. *C. auriculata* var.  $\alpha$ . Ell. sketch 2. p. 436. *Lea-ebos trifoliata* Cass. dict. 25. p. 389. Coreopsoides lanceolata Mœnch meth. 594. *Anacis auriculata* Schrank akad. Münch. nat. 5. p. 7. *Chysomeles auriculata* Tausch h. Can. f. 1. — Pluk. alm. t. 83. f. 5. (v. v. c.)

$\beta$ . *glabrata*, caule glabro, foliis glabriusculis, petiolis vix basi ciliatis.  $\frac{1}{2}$  cum var.  $\alpha$  in hortis sæpius culta, fortè var.  $\beta$  è culturâ aut loco orta. Utraque var. variat achæniis plûs minùs ut in *Calliopside incurvis*, intus extrinseque muricato-tuberculatis aut lævibus, aristis minimis mox deciduis aut fortè nullis. (v. v. c.)

13. *C. DIVERSIFOLIA* (DC. non Jacq.), caule cespitoso brevissimo pilis retrorsis patulis hispidis, foliis radicalibus oppositis glabriusculis, aliis ovatis indivisis, aliis trisectis lobo medio ovato lateralibus multò minoribus, petiolis hispidulis limbo multò longioribus, pedunculis longissimis glabris, invol. squamis ext. lanceolatis, ligulis 4-5-dentatis, achæniis orbiculatis alatis, alâ subdentatâ. (Ell.)  $\frac{1}{2}$  in Carolinā mediâ (Whitl. in Ell.), et boreali (B. Greene!). Coreopsis auriculata  $\beta$  diversifolia Ell. sketch 2. p. 437. — Pluk. alm. t. 242. f. 4. Habitu et car. indicatis satis à *C. auriculatâ* differe videtur. In specim. meo ut in icone Pluken. folia radicalia longè petiolata indivisa, caulina trisecta, sed Elliott dicit folia inf. trisecta, sup. indivisa. *C. diversifolia* Hook. bot. mag. t. 3474 in agro Texano à cl. Drummond lecta videtur eadem. (v. s. comm. à cl. Greene.)

14. *C. SENIFOLIA* (Michx. fl. bor. am. 2. p. 138), caule erecto trichotomo, foliis

# DICOTYL. SEU EXOGENÆ.

COROEPSIS.

sitis sessilibus trisectis, segmentis ovato-lanceolatis integerrimis pube brevitate utrinque velutinis membranaceis, capitulis subcorymbosis, involucri ext. lineari-oblongis obtusis, ligulis oblongis integris, acheniis oblongis angustè alatis glabris integerrimis, aristis minimis deciduis. 2 in Virginie (Pursh), Carolinæ et Georgiæ (Lyon.), Kentucky (Raf.), in (Fras.) sabulosis (Michx.), et pinetis (Ell.). Pursh fl. bor. am. sketch 2. p. 438. Coreopsis major Walt. car. 214. C. stellata Bask. ex Pursh. (v. s.)

2. *rigida* (Nutt. gen. am. 2. p. 180), glabrescens rigida, foliis indivisis, super. tripartitis aut bipartitis imò subindivisis, segmentis angustè lanceolatis supra scabridis. — in Georgia (Nutt.), in Carol. merid. ex specim. à cl. Fraser et Mitchell missis. An forte species? (v. s. sp.)

3. *DELPHINIFOLIA* (Lam. dict. 2. p. 108), glabra, foliis oppositis sessilibus si trisectis, segmentis indivisis seu pinnatipartitis, lobis lineari-lanceolatis integerrimis, capitulis pinnatis, squamis ext. lineari-oblongis, ligulis integris acutis, acheniis obovatis subulatis breviter alatis, discis siccis Amer. bor. in Virginia (Duby!). C. verticillata var. Linn. 1281 (excl. syn. Pluk.). C. verticillata Willd. sp. 3. p. 225t. bot. maj. 156. Schkuhr handb. 3. t. 260 b. f. s. Delphinium foliis Vaill. ac. Par. 1720. Ehret. pict. t. 61. f. 1. Delphinium non sunt verticillata sed opposita. Variat caule folisque glabris aut minutè subpuberulis. Discus fuscus, ligulæ flavæ. (v. s. et s. sp.)

16. *C. TENUIFOLIA* (Willd. sp. 3. p. 225t), glabra, foliis oppositis sessilibus bipartitis, lobis linearibus integerrimis, capitulis pinnatis, squamis ext. lineari-oblongis obtusis, ligulis integris acutis, acheniis obovatis subulatis minutè alatis, discis siccis Virginie (Pursh) et Carolinæ (Rose!). Schkuhr handb. 3. t. 260 a. fig. dextr. C. verticillata Linn. sp. 1281 (excl. syn. Vaill.). C. verticillata Lam. dict. 2. p. 108. C. verticillata & tenuifolia Michx. fl. bor. am. 2. p. 139. — Pluk. mant. t. 344. p. 4. C. delphinifolia affinis, sed lobi folii angustiores semper glabri, discus luteus, etc. (v. s. et s. sp.)

17. *C. GRANDIFLORA* (Hogg. mss. ex Sweet brit. fl. gard. 2. t. 175), caule erecto ramoso sulcato glabro, foliis oppositis sessilibus glabris basi ciliatis, infer. bipartitis, super. trisectis tripartitiave, lobis linearibus elongatis integerrimis supra canaliculatis, pedunculis elongatis 1-cephalis, invol. squamis ext. lanceolatis acuminatis interiores æquantibus, ligulis acutè 5-dentatis, acheniis elliptici vix alati aristis subpaleaceis cristatis. 2 in Americâ bor. sine loc. propr. design. à Noveboraco missa. Capitulum poll. 2 ferè diam. disco radiato luteo concoloribus. (v. s. in h. Barclay ubi etiâ Sweet suam vidit, à Fries coll.)

18. *C. MITIS* (Michx. fl. bor. am. 2. p. 140), glaberrima, caule obtusè tetragono brachiato, foliis oppositis bipinnatipartitis, lobis linearibus serrulatis, capitulis paniculatis, invol. ext. squamis linearibus serrulatis, ligulis obovatis apertè obscure dentatis, acheniis oblongis serrulatis biaristatis. (2) ex Pursh. In humidis Carolinæ (Michx.) et in agro Missouriensi (Lew. ex Pursh.). C. coronata Walt. car. 2. p. 15 ex Ell. sketch 2. p. 441.

19. *C. TRICHOSPERMA* (Michx. fl. bor. am. 2. p. 139), glabra, caule obtusè tetragono, foliis oppositis, lobis lanceolato-linearibus subdentatis, superiorum in limbum lineari-lanceolatum elongatum dentatum confluentibus, capitulis subcorymbosis, invol. squamis ext. oblongo-linearibus obtusis subciliatis, ligulis integris obtusis, acheniis planis puberulis, aristis cor. disci longitudinem subæquantibus. 2 Ell. (2) Pursh. In humidis Carolinæ super. (Michx.). Nova-Cæsareæ (Greene!), Bostoniæ (Big.), Noveboraci et Nova-Cæsareæ (Torrey). Habitus omnino Bidentis sed aristæ non retrorsum scabræ. (v. s.)

20. *C. AUREA* (Ait. kew. ed. 1. v. 3. p. 252), caule glabro striato, foliis oppositis tripartitis trisective pubescenti-pilosis, lobis linearibus grossè serratis alio-

divisis aut terminali tripartito, invol. squamis ext. linearibus patentibus, ligulis amplis obovatis integris, achæniis cuneatis glabriusculis apice bicornibus. ② in Amer. bor. fortè circa Noveboracum. Lindl. bot. reg. 1228. *C. trichosperma* β aurea Nutt. gen. am. 2. p. 180. Capitula ampla aurea.

21. *C. ARISTOSA* (Michx. fl. bor. am. 2. p. 140), pubescens, foliis oppositis quinato-pinnatisectis, lobis serratis, ligulis integris lato-ovalibus, achæniis cuneato-obovatis biaristatis, aristis longissimis divaricatis. ♀ in Carolinâ infer. (Pursh) et agro Illinoensi (Michx.). *C. aristata* Willd. sp. 3. p. 2253. Capitula ampla. Ligulæ latissimæ.

22. *C. PUBESCENS* (Ell. sketch 2. p. 441), pubescens, ferè subtomentosa, foliis quinato-pinnatisectis, lobis lanceolatis obtusis integerrimis, lateralibus parvis, invol. ext. squamis ovato-lanceolatis glabris, ligulis undulatis? achæniis orbiculatis angustè alatis emarginatis, pappo diphylo brevi subulato pubescente. ♀ in Georgiâ occidentali.

23. *C. MULTIFIDA*, herbacea glabra erecta, foliis oppositis petiolatis circumser. suborbiculatis pinnatipartitis, lobis pinnatifidis, lobulis lanceolatis brevibus hinc inde incisus, capitulis pedunculatis solitariis paucisve corymbosis, invol. squamis lanceolatis subæqualibus, ligulis ellipticis integris, achæniis (immaturis) linearibus apice attenuatis subciliatis in aristas 2 rigidulas glaberrimas abeuntibus. — in Peruvîâ legit cl. Pavon ex h. Thibaud. Huc refero specimina duo: unum spontaneum etate proevectius ligulis nullis deciduis, alt. hortense junius ligulis albidis cæterum simillimum. (v. s.)

β. *mutica*, caule magis angulato, foliorum lobulis paulò acutioribus, achæniis exaristatis. — in Peruvîâ undè semina ad hort. Par. olim misit cl. Dombey. Var. α simillima præter notas indicatas. Ligulæ ex sicco vid. roseæ. (v. s. ex h. L'her.)

24. *C. ANTHEMOIDES*, caule herbaceo demisso ramoso radicante subpuberulo, foliis oppositis glabris multifidis, lobis linearibus acutis, pedunculis terminalibus 1-cephalis erectis, invol. squamis ext. linearibus interiores subæquantibus, ligulis ellipticis integris, achæniis (immaturis) linearibus glabris brevissimè bidentatis, ♀? in Mexico ad Guachilapam in Cordillera legit cl. Berlandier (pl. exs. n. 1030), fl. octob. Herba demissa habitu ferè Anthemidis arvensis. Flores latei. Ligulæ 3-4 lin. longæ. (v. s.)

25. *C. NÆSI*, fruticulosa? ramis teretibus glabris trichotomis, foliis oppositis basi hirsuto-ciliatis lineari-filiformibus acutis, aliis indivisis, aliis tripartitis, capitulis pedunculatis solitariis, invol. extus hirsuto, squamis subæqualibus, ligulis elliptico-oblongis integris, achæniis glaberrimis brevissimè bidentatis, ♂? in Amer. calid. legit cl. Née, sed locus propr. ign. Fl. flavi. Ligulæ 6-7 lin. longæ. Achænia matura ign. (v. s. ex h. Thibaud. specim. manc.)

26. *CAPILLACRA* (H. B. et Kunth nov. gen. am. 4. p. 230), fruticosa glabra, ramis tetragonis, foliis oppositis petiolatis tripartitis, lobis filiformibus integerrimis, capitulis solitariis pedicellatis, invol. squamis exter. paulò brevioribus lanceolato-linearibus ciliatis, ligulis lanceolato-oblongis subtridentatis, achæniis margine lanato-pilosis biaristatis, aristis corollas disci æquantibus. ♀ in Andibus Peruviae. Flores disci flavi. Ligulæ aurantiacæ.

27. *C. TRIPIDA* (Lam. ill. t. 704. f. 2. Poir. suppl. 2. p. 353), glabra, caule subflexuoso, foliis trifidis subulatis in axillâ fasciculatis, pedunculis terminalibus 1-cephalis, ligulis angustis obtusè et grossè 3-dentatis. — patr. et cat. ign. Folia ex icone opposita, ex descr. alterna. †

† *Species non satis notæ.*

28. *C. PALMATA* (Nutt. gen. am. 2. p. 180), caule simplici compresso sub-1-cephalo, foliis oppositis sessilibus subcoriaceis palmato-3-lobis margine scabris, lobis lineari-oblongis obtusis, invol. utroque 8-partito, achæniis oblongo-ellipticis nudis. — in planitiis terit. Michiganensis, Illinoensis et in Louisiana infer. Calliopis palmata Spreng. syst. 3. p. 611. Flores lutei. An Perambibi sp. ex Raf.

29. *C. NUDATA* (Nutt. gen. am. 2. p. 180), caule humili simplici apice dichotomo foliis alternis glabris subulato-linearibus remotissimis, super. minimis, radiis obsoletis 3-dentatis, acheniis nudis. — in Florida occid. circa urbem Sanctæ Mariæ. *Calliopsis nudata* Spreng. syst. 3. p. 611. Ligulæ rubræ. Invol. est minutum. An *Peramibi* sp. ex Raf.† †.
30. *C. ROSEA* (Nutt. gen. am. 2. p. 179), parva glaberrima, caule sæpius simplici apice dichotomo oligocephalo, foliis oppositis linearibus integris, capitulis longè pedunculatis terminalibus, ligulis inæqualiter 3-dentatis, acheniis integerrimis nudis. ¼ in paludosis à Novâ-Cesareâ ad Georgiam. *Calliopsis rosea* Spreng. syst. 3. p. 611. Ligulæ roseæ. An *Peramibi* sp. ex Raf.† †.
31. *C. PAUCIFLORA* (hort. Hamb. ex linnæâ 1835. litt. p. 76), caule herbaceo simplici, foliis oppositis sessilibus tripartitis glabris, lobis lanceolatis obtusiusculis margine scabris, medio longiore, capitulis terminalibus subsolitariis. 1 in Americâ boreali.
32. *C. BITERNATA* (Lour. fl. coch. 622), caule herbaceo erecto tetragono, foliis biternatis ovato-lanceolatis serratis, paniculâ diffusâ, ligulis 6 neutris, pappi bicorni ramoso, recept. nudo. — prope Cantonem Sinarum. *Actines biternata* Spreng. syst. 3. p. 474. Caulis integrè luteus dicitur. Genus omninò ambiguum ex recept. nudo dicto et fortè ad *Actineam* rectè relatum. †.

†† *Species exclusæ.*

- C. ACUTA* Pursh = *Peramibus acutus*.  
*C. ALATA* Cav. = *Actinomeris tetraptera*.  
*C. ALATA* Pursh = *Actinomeris alata*.  
*C. ALBA* Linn. = *Bidens alba*.  
*C. ALTERNIFOLIA* Linn. = *Actinomeris alternifolia*.  
*C. AMPLEXICAULIS* Cav. = *Simsia amplexicaulis*.  
*C. ANGUSTIFOLIA* Pav. = *Bidens procera*.  
*C. ARTEMISIAEFOLIA* Jacq. = *Cosmos sulphureus*.  
*C. ATKINSONIANA* Dougl. = *Calliopsis Atkinsoniana*.  
*C. BACCATA* Linn. = *Wulffia stenoglossa*.  
*C. BIDENS* Walt. = *Bidens chrysanthemoides*.  
*C. CHRYSANTHA* Spreng. = *Bidens Coreopsidis*.  
*C. CORONATA* Linn. = *Bidens leucantha*.  
*C. CORONATA* Spreng. = *Bidens oxyodonta*.  
*C. DIVERSIFOLIA* Jacq. = *Bidens grandiflora*.  
*C. FERULÆFOLIA* Jacq. = *Bidens Coreopsidis*.  
*C. FLAMMULA* Banks. = *Bidens chrysanthemoides*.  
*C. FOETIDA* Cav. = *Simsia ficifolia*.  
*C. FORNICULACEA* fl. mex. = *Bidens fœniculifolia*.  
*C. FORMOSA* Bosc. = *Cosmos bipinnatus*.  
*C. FRUTICOSA* Vest. = *Bidens fruticosa*.  
*C. FRUTICOSA* Forst. = *Campylothea micrantha* Æ.  
*C. GEORGINA NUDA* Cass. = *Dahlia variabilis*.  
*C. GEORGINA PRUINOSA* Cass. = *Dahlia coccinea*.  
*C. HETEROPHYLLA* Cav. = *Echinacea heterophylla*.  
*C. HIRTA* Raf. = *Peramibus hirtus*.  
*C. INCISA* Ker. = *Bidens Coreopsidis*.  
*C. LEUCANTHA* hort. = *Bidens leucantha*.  
*C. LEUCORHIZA* Lour. = *Bidens leucorhiza*.  
*C. LIMENSIS* Jacq. = *Encelia Limensis*.

- C. ODORATA Lam. = *Bidens odorata*.  
 C. OVATA Cav. = *Actinomeris ovata*.  
 C. PARVIFLORA Jacq. = *Cosmos parviflorus*.  
 C. PREPOLIATA Bosc. = *Bidens chrysanthemoides*.  
 C. PROCERA Ait. = *Actinomeris alternifolia*.  
 C. PURPUREA fl. mex. = *Bidens purpurea*.  
 C. REPTANS Linn. = *Bidens Coreopsis*.  
 C. SCABRA Raf. = *Perambius scaber*.  
 C. SERRULATA Poir. = *Bidens grandiflora*.  
 C. TETRAGONA Lellav. = *Bidens tetragona*.  
 C. TINCTORIA Nutt. = *Calliopsis tinctoria*.  
 C. TRIPTERIS Linn. = *Chrysostemma tripteris*.

CCCXXXVIII. *ACTINOMERIS* Nutt. gen. am. 2. (1818) p. 131. Less. syn. 228. — Ridan Adans. fam. 2. (1763) p. 130. — Pterophyton Cass. bull. philom. 1818. p. 76. dict. 44. (1826) p. 48. 59. p. 139. — Actimeris Raf. 1820. — Coreopsis sp. Linn.

Capitulum multiflorum radiatum, ligulis neutris paucis 1-serialibus elongatis, fl. disci herm. tubulosi 5-dentatis. Inv. 1-3-seriale, squamis subinæqualibus foliaceis acuminatis. Recept. convexum paleaceum, paleis achæniorum marginem amplexantibus. Styli rami appendiculati acuti. Achænia bilateraliter plano-compressa angustè alata, aristis binis triquetris sublaevibus coronata. — Herbæ Boreali-Americanae scabrae biennes perennesve. Caules erecti ramosi teretes foliis longè decurrentibus alati. Folia ovata aut lanceolata serrata alterna rariùs subopposita. Capitula corymbosa. Corollæ flavæ (aut var. fortè ligulæ albæ). — Differt à Coreopside achæniis lateraliter compressis nec obcompressis, à Verbesinâ ligulis neutris.

1. *A. ALTERNIFOLIA*, caule supernè pubescente, foliis alternis lato-lanceolatis serratis scabris, paniculâ corymbosâ laxâ foliosâ, invol. patente uniseriali, recept. subgloboso. ♀ in Virginiâ (Pluk. Pursh), Carolinâ (Fras.), Georgiâ (Ell.), in territorio Arkansano è semin. à cl. Pourtales lectis. Coreopsis alternifolia Linn. sp. 1283. Jacq. hort. vind. t. 110. Ridan Adans. fam. 2. p. 130. Pterophyton alternifolium Cass. dict. 44. p. 49. Actinomeris squarrosa Nutt. gen. am. 2. p. 131. Ell. sketch 2. p. 413. Verbesina Coreopsis Michx. fl. bor. am. 2. p. 134. — Pluk. alm. t. 159. f. 3. Ligulæ 7-8 flavæ. Variat etiâ ligulis nullis ex Michx. Folia infima opposita aut subverticillata, undè verisim. Coreopsis procera Ait. kew. 3. p. 258 ab hac non differt. (v. v. c. et s. sp.)

♀? *alba* (Ell. sketch 2. p. 413), ligulis albis, foliis angustius lanceolatis magis scabris, corollis disci glaberrimis. ♀ in Carolinâ inferiore. An species propria?

2. *A. HELIANTHOIDES* (Nutt. gen. am. 2. p. 181), caule subscabro, foliis alternis lanceolatis acutis serratis scabris subtus villosis, paniculâ corymbosâ subsimplici oligocephalâ coarctatâ, invol. biseriali, recept. convexo. ♀ in Georgiâ circa Louisville (Ell. sketch 2. p. 412), in Ohio, Kentucky, Tennessee et Louisiana infer. (Nutt.). Verbesina helianthoides Michx. fl. bor. am. 2. p. 135 non Kunth. Ligulæ 10-12 aureæ elongatæ.

3. *A. TETRAPTERA*, caule subscabro, foliis oppositis et superioribus alternis ovato sen rhombeo-lanceolatis scaberrimis basi cuneatis à medio serratis, pedunculis ad capitulum usquè alatis subcorymbo, ligulis 14-15. ♀ in Mexico (Cav.). Coreopsis alata Cav. ic. 3. p. 3. t. 260 non Pursh. Helianthus tetrapteris Ort. dec. 6. p. 74. Moc. fl. mex. ic. ined. Pterophyton alatum Cass. dict. 44. p. 49. — Hern. mex. p. 189. ic. Ligulæ flavæ. (v. v. c.)

4. *A. TETRAGONA*, foliis etiā superioribus oppositis ovatis serratis basi angustatis, pedunculis alatis 1-cephalis.  $\frac{1}{2}$  in Mexico circa urbem (Alam. Berl. f. pl. exs. n. 647), ad Temisci prædium saccharinum ubi *Contrayerva* audit (fl. mex. mss.). *Helianthus tetragonus* Moq. fl. mex. ic. et descr. ined. Mamantlaeod. Heru. mex. p. 383. f. 2. Affinis *A. tetrapteris* et fortè ferè varietas sed ex Hernandez et Moqino videtur diversa. Antiseptica et febrifuga ulcera putrida cito et mirificè sanat ex Moq. (v. s.)
5. *A. OVATA* (Nutt. gen. am. 2. p. 181 in adnot.), caule glabrinsculo, foliis alternis ovato-oblongis crenulatis, capitulis corymbosis, pedunculis apteris, invol. squamis linearibus obtusis, ligulis 8.  $\frac{1}{2}$  in Mexico. *Coreopsis ovata* Cav. ic. 3. p. 41. t. 280. Ligulae flavae.
6. *A. PEDUNCULOSA*, foliis oppositis ovato-rhombeis dentatis triplinerviis, pedunculis terminalibus longissimis 1-cephalis omnino exalatis.  $\frac{1}{2}$  in Mexico. Folia utrinquè scabro-aspera, longè secus caulem decurrentia, apice ferè obtusa. Pedunculi 7-8 poll. longi. (v. s. comm. à cl. Alaman.)
7. *A. ALATA* (Nutt. gen. am. 2. p. 181), foliis oppositis indivisis ovatis acuminatis serratis petiolatis triplinerviis. (2) in Virginia (ex Bartr. in h. Banks). *Coreopsis alata* Pursh fl. bor. am. 2. p. 567 non Cav.
8. *A. PAUCIFLORA* (Nutt. in Sillim. journ. 5. (1822) p. 301), foliis oppositis oblongis serrulatis, pedunculis nudis oligocephalis. — in Florida orientali. Caulis videtur simplex. Invol. simplex squamis lanceolatis. Achenia breviter bicornia. Flor. ign.

CCCXXXIX. *ARMANIA* Berter. herb. — *Hopkirkia* sp. Spreng.

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris 1-serialis, disci hermaphroditis cor. tubo brevi, fauce latiore cylindrica, limbo 5-dentato. Invol. 3-seriale, squamis imbricatis adpressis. Recept. planum paleaceum. Styli fl. hermaphrod. rami breves truncati apice barbati exappendiculati. Achæmium obcompressum ellipticum corollagineum adpressè pubescens, setis 2 rigidis anticè (nec retrorsum scabris) coronatum. — Frutex Austr.-Americanus, ramis teretibus foliisque adpressè villosis. Folia petiolata ovata acuminata subcrenata penninervia. Capitula pauca (2 in specim.) pedicellata terminalia. Flores flavi. — Genus à Salmeà (quæ *Hopkirkia* Spr.) diversissimum capitulis radiatis, stigmate truncato apice barbato, recept. plano etc.; à *Bidentate* differt involucre, stigmate, formâ achæнии et habitu toto.

1. *A. FRUTICULOSA* (Berter. herb.).  $\frac{3}{4}$  ad Sanctam-Martham Amer. merid. aë legit amic. Bertero. *Hopkirkia fruticulosa* Spreng. syst. 3. p. 444. (v. s.)

CCCXL. *OYEDÆA*.

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris 1-seriatis, disci tubulosi 5-dent. hermaphr. Invol. squamæ 3-seriales subæquales foliaceæ. Recept. planum, paleis complicatis mucronatis terminatis coriaceo-subscariosis. Cor. disci tubus gracilis, faux cylindrica, dentes 5 erecti extus puberuli. Anth. nigricantes incluse. Styli rami graciles appendice hispidâ superati. Achænia radii abortiva linearia pappo 2-3-aristato aristellisque minimis superata; disci compressa alata subcuneata, apice hinc sæpè gibba, pappo vario, aristis nempe 2 caducis elongatis et pluribus aliis brevibus intermediis constante. — Frutex Austr.-Americani *Verbesinæ* facie. Rami teretes adpressè villosi subhirti. Folia opposita in petiolum attenuata ovata acuminata dentata supra sparse villosa subtus villosa-subcanescentia aut tomentosa, inter-grossè triplinervia, summa subpenninervia. Capitula pedunculata



*rymbosa*. Flores lutei.—Genus à *Verbesinâ* ligulis neutris, ab *Heliantho* et *Viguierâ* achæniis alatis diversum, ab *Actinomeride* squamellis inter duas aristas pappi intermediis. Dicavi cl. Hispano navigatori Alphonso Oyeda, qui, primus post Colombium, Americam adiit et imò in nave suâ vehebat Americum Vespucium nimio honore postea decoratum cum Oyeda ferè oblitus. (Conf. bibl. voy. 6. p. 212.)

1. *O. VERBERINOIDES*, foliis subtus villosis, invol. squamis lineari-lanceolatis acuminatis pubescentibus.  $\frac{3}{4}$  circa Caracas legit cl. Vargas. Folia inf. pedem longa, 7-8 poll. lata. Ligulæ pollicem longæ, apice 2-3-dentatæ. Achænia glabra sparsè maculata. (v. s.)
2. *O. BUPHTHALMOIDES*, foliis subtus secus nervos villosos-strigosis inter nervos tomentosis, invol. squamis ovatis mucronatis, ext. densè villosis, inter. submembranaceis longè ciliatis.  $\frac{3}{4}$  in Peruvâ legit cl. Poeppig. *Bupthalmum trifolium* Poepp. pl. exa. n. 22 et 1624. Caules ramique densè hirsuti. Folia opposita, nec in meo specim. ternâ, 9 lin. longa. Achænia radii 3- disci 2-aristata, aristellis minimis paucis hinc inde interjectis. (v. s. comm. à cl. Poepp.)

CCCXLI. *SIMSIA* Pers. ench. 2 (1807). p. 478. Cass. dict. 59. p. 136 non Brown. (1810). — *Ximenesia* sp. Kunth Less. — *Simsia* et *Ximenesia* sp. Loud. h. brit.

Capitulum multiflorum heterogamum, fl. radii neutris ligulatis 1-serialibus paucis (5-12), disci hermaphr. tubulosis 5-dentatis. Invol. squamæ 2-3-seriales erectæ lineares subæquales. Recept. planum, paleis membranaceis complicatis. Styli rami appendiculati. Achænia plano-compressa glabriuscula exalata apice biaristata. — Herbe Mexicanæ, *Viguieræ* facie, pleræque hispide. Folia inf. opposita, sup. alterna trinervia petiolata aut sessilia, sæpè basi auriculato-amplexicaulia. Capitula pedicellata laxè corymbosa. Flores omnes lutei. — Differt à *Viguierâ* squamellis inter aristas nullis, ab *Heliantho* invol. non imbricato, à *Ximenesiâ* ligulis neutris et achæniis exalatis.

1. *S. LAGASCÆFORMIS*, caule parçè subhispido, foliis inf. oppositis, sup. alternis petiolo exaurito longè piloso donatis ovatis acuminatis serratis utrinquè mollioribus subvillosis, corymbo laxo subpauciculato, inv. squamis lanceolatis, exter. paucis dorso hispidis dimidiò brevioribus, inter. glabris subcoloratis disco æqualibus, ligulis circ. 5 breviter bidentatis, achæniis glabris. — in Mexico. Species ab omnibus aliis faciliè differt invol. serie externâ dimidiò internâ breviorè. (v. s. comm. à cl. Mairet.)
2. *S. FICIFOLIA* (Pers. ench. 2. p. 478), ramis hispidis et pube glandulosâ pulverulento-velutinis, foliis utrinquè puberulis petiolatis exauriculatis cordatis profunde trilobis dentatis, infer. oppositis, sup. alternis, capitulis paucis subpaniculatis, invol. squamis ext. hispidis, int. margine membranaceis, ligulis 8 vix denticulatis, achæniis glabris apice vix ciliatis. (X) in Mexico. *Cœreopsis fetida* Cav. ic. t. p. 55. t. 77. *Ximenesia fetida* Spreng. syst. 3. p. 606. Herba graveolens. Folia exactè Fici aut Mori. (v. s.)
3. *S. HETEROPHYLLA* (DC. non Cass.), ramulis hispidis, foliis inf. oppositis, sup. alternis, omnibus ovali-oblongis membranaceis utrinquè pilosisculis denticulatis, aliis petiolatis petiolo nudo exauriculato, aliis sessilibus obtusè auriculatis, nunc variè trilobis nunc indivisis, invol. hispidi squamis lineari-subulatis, ligulis 7-8 tridentatis, achæniis glabris margine ciliatis.  $\frac{2}{3}$  in Mexico prope Zinepequaro (H. et B.) inter Tampico et Real del Monte (Berl. pl. exa. n. 442) *Ximenesia heterophylla* H. B. et Kunth nov. gen. am. 4. p. 227. t. 380. *Simsia Kunthiana* Cass. dict. 59. p. 137, sed *Simsia heterophylla* Pers. è genere excludenda et ad *Echinaceam* referenda. (v. s.)
4. *S. AURICULATA*, ramis hispidis, foliis utrinquè villosis latè ovatis irregulariter

trilobatis acuminatis dentatis basi breviter cuneatis, petiolo hispidulo immarginato basi latè auriculato, corymbo composito laxo polycephalo, invol. squamis lineari-lanceolatis acuminatis, ext. hispidis, int. margine membranaceis, ligulis 6-8 minutè 2-3-dentatis, achæniis glabrisculis. — in Mexico circa artem legit cl. Berlandier (pl. exs. n. 850 et 927). *Encelia Mexicana* et *Ximenesia hirta* Mart. ! herb. A. S. heterophyllâ differt foliis multò latioribus, petiolisque basi amplè auriculatis. Variat foliis oppositis alternis, limbo basi truncato aut ovato. (v. s.)

5. *S. HISPIDA* (Cass. dict. 59. p. 137), caule apice hirtulo-hispidulo, foliis alternis sessilibus ovato-oblongis obsolete serratis suprâ piloso-subtùs sericeo-hispidis, invol. squamis hispidis lanceolatis, ligulis circ. 12, achæniis .... 2½ in montanis Mexici prope S. Rosa de la Sierra et Valladolid. *Ximenesia hispida* H. B. et Kunth nov. gen. am. 4. p. 227. *S. heterophylla* similis dicitur.
6. *S. CORDATA* (Cass. dict. 59. p. 137), ramis pilosis, foliis inf. oppositis, sup. alternis deltoideo-ovatis acuminatis cordatis basi cuneatis grosse crenatis utrinque adpressè hispidulis, petiolis basi amplè dilatatis, invol. squamis lanceolatis-linearibus piloso-hispidis, ligulis 6-8 emarginato-bidentatis, achæniis sericeo-pubescentibus. 2½ in Mexico prope Capula et Valladolid de Mechoacan. *Ximenesia cordata* H. B. et Kunth nov. gen. am. 4. p. 228. *Ximenesia cuneatis* proxima dicitur ! An fortè ideò *Ximenesia* spec?, sed ligulas dicuntur neutre et achæniis ala marginalis ex descr. videtur nulla.
7. *S. DOMBEYANA*, caule tereti sparsè hispido et inter setas minutè puberulo-glanduloso, foliis superioribus alternis petiolatis latè ovatis irregulariter repando-dentatis hinc indè sublobatis acutis utrinquè setis hispidulis et puberulo-glandulosis, capitalis paucis breviter pedicellatis, invol. squamis linearilanceolatis acuminatis disco longioribus, ligulis paucis minimis, achæniis obcordatis biaristatis margine ciliolatis. — in Amer. austr. verisim. in Peruvia legit Dombey. Petioli valdè hispidi, setis at in totâ plantâ longis patulis mollioribus. (v. s. comm. à Mus. reg. Par.)
8. *S. AMPLEXICAULIS* (Pers. ench. 2. p. 478), canescens, foliis alternis subpinnato-trilobis, petiolo basi auriculato-semiamplexicauli. — in Mexico. *Coreopsis amplexicaulis* Cav. cors. descr. p. 226. Lag. et Rod. in annal. scienc. nat. 1801 v. 5. p. 75. Ait. kew. ed. 2. p. 134. Folia sæpius sub-5-loba, infima apertè obliqua indivisa. Flores crocei. (Pers.)

† *Species exclusa.*

3. *HETEROPHYLLA* Pers. = *Echinacea heterophylla*.

CCCXLII. *VIGUIERA* H. B. et Kunth nov. gen. am. 4. p. 224. — *Viguiera* Lessert. 229.

Capitulum multiflorum heterogamum, fl. radiis neutris ligulatis, f. disci hermaphr. 5-dentatis. Invol. semi-globosum, squamis 1-2-seriatim subæqualibus apice in appendicem foliaceam productis. Recept. paleis persistentibus achænia amplectentibus obsitum. Styli rami exerti subulati hispidi. Achænia compressa obovato-cuneata pubescentia squamellis 4 ovatis subdenticulatis aristisque 2 ex angulis ortis coronata. — Herbar. Americanæ ramosæ. Caules glabri, rarius villosi. Folia opposita aut alterna petiolata triplinervia ovato-lanceolata sæpè scabra. Capitula pedicellata laxè paniculato-corymbosa. Cor. luteæ. — Genus *Heliantho* et *Leighiæ* affine, sed caractere Kunthianorum dilatato, apprimè naturale.

§. 1. *Receptaculo acutè aut obtusè conico.*

1. *V. HELIANTHOIDES* (H. B. et Kunth nov. gen. am. 4. p. 226. t. 379), caule glabro, foliis alternis ovatis acuminatis integerrimis trinerviis suprâ subtè adpressè pilosis, petiolis villosis-ciliatis, invol. squamis uni?seriatis

linearibus hispidulis apice dilatato-subspathulatis, recept. longo conico cavo. ① in insulâ Cubâ prope Havanam frequens.

2. *V. SAGRAMANA*, caule glabro, foliis oppositis et alternis ovato-lanceolatis subintegerrimis triplinerviis suprà subcabriusculis subtis petiolisque parè pubescentibus, invol. squamis biserialibus linearibus in appendicem foliaceam acutam productis, recept. conico demùm cavo. ①? in insulâ Cubâ juxtâ Havanam legit cl. Ramon de La Sagra. Valdè affinis *V. helianthoidi*, sed folia frequentius opposita subintegra, petioli non villosi-ciliati, invol. diversa, etc. (v. a. comm. cum seq. à cl. de La Sagra).
3. *V. MICROCLINA*, caule glabro, foliis oppositis ovato-lanceolatis integerrimis trinerviis suprà pube minutissimâ subcabris subtis petiolisque villosi-canescantibus, invol. squamis biserialibus in appendicem foliaceam acutam apice productis, recept. conico pleno. ①? in insulâ Cubâ prope Havanam legit cl. Ramon de La Sagra. Habitus *V. helianthoidis*, sed juxta hujus descr. et iconem facili diversa. Ligulæ 5-7. Achænia adpressè villosa. (v. a.)
4. *V. CANESCENS*, caule villosi, foliis alternis ovalibus acuminatis in petiolum longè attenuatis integerrimis suprà pilosulo-scabris subtis petiolisque villosi-canescantibus, invol. squamis biserialibus dorso canescentibus in appendicem linearem acutiusculam productis, recept. obtusè conico. ① in Mexico prov. Leoninâ ad occid. nrbs Guanajuato legit cl. Mendez. Habitus ferè *Ximenensis*. Cor. croceo-flavæ. (v. a.)
5. *V. TRIQUETRA*, caule glabro, foliis serratis suprà scabris subtis pubescentibus, infer. oppositis, summis subalternis, petiolis glabriusculis, invol. squamis linearibus acutis biserialis, recept. obtusè conico demùm cavo. ①? in valle Mexicanâ prope S. Angelo legit cl. Alaman et etiam Berlandier (pl. exs. n. 827 et 937). Panícula trichotoma, pedunculis angulatis subtriquetris. *Helianthus triquetra* Ort. dec. p. 76. (v. a.)
6. *V. DENTATA* (Spreng. syst. 3. p. 615), caule glabro, foliis serratis oppositis, petiolis subciliatis, paniculâ oligocephalâ, invol. squamis 2-seriatis acutis, recept. convexo. — in Mexico. *Helianthus dentatus* Cav. ic. 3. p. 10. t. 220. Folia basè latiora apice magis acuminata quàm priorum. Pedicelli graciles vix angulati. (v. a.)

### §. 2. *Receptaculo planiusculo.*

7. *V. PROSTRATA*, caule diffuso debili pubescente, foliis oppositis breviter petiolatis ellipticis utrinquè acutis apice subserratis triplinerviis utrinquè pubescenti-scabridis, ramis elongatis floridis apice nudiusculis 1-cephalis, invol. squamis lineari-oblongis biserialis canescenti-villosis, paleis indivisis, achæniis glabris angulatis. ½ in Amer. bor. ex Willd., sed ab hodiernis in confus. Anglo-Americ. non reperta! fortè Mexicana? *Helianthus prostratus* Willd. sp. 3. p. 2242 non Poir. Pappi aristæ tenuissimæ, squamellæ minimæ. Caulis ex Link non verè prostratus. (v. a.)
8. *V. SESSELIFOLIA*, caule suffruticoso adscendente laxè ramoso sparsè hispidi demùm glabrato, foliis inf. oppositis, rameis alternis, sessilibus ovatis acutis subintegerrimis utrinquè villosis, paniculâ laxâ oligocephalâ subnudâ, invol. squamis 2-3-seriatis oblongo-linearibus villosi-hispidis, ligulis 5-6, paleis lanceolatis acuminatis, achænio adpressè sericeo. ½ in Mexico circa Villalpando legit cl. Mendez et ad Tlapujahua cl. F. W. Keerl (h. Mart.). Paleæ recept. glabræ, apice virescentes, cor. disci æquales. Pappus squamellis 2 aristatis, 4 brevibus, omnibus ciliato-pubescentibus. Rec. planiusculum. Ligulæ pallidè flavæ nervis nigricantibus lineatæ. (v. a.)
9. *V. BREVIFLORA*, caule herbaceo puberulo laxè ramoso, foliis breviter petiolatis ovato-lanceolatis acuminatis subintegerrimis triplinerviis utrinquè puberulo-scabris, petiolis villosi-ciliatis, paniculâ laxâ, invol. squamis 2-3-seriatis lanceolatis acutis dorso subtrinerviis, paleis lanceolatis acutis membranaceis, achæniis adpressè subpubescentibus. — in Mexico circa Bejar legit cl. Berlandier (pl. exs. n. 1905). Habitus omninò *V. helianthoidis*. (v. a.)

\*\* *Folii oppositis.*

9. *L. ANGUSTIFOLIA*, caule suffruticuloso ramosissimo tereti basi glabro, ramis puberulo-hirtellis, foliis oppositis sessilibus linearibus acutis integerrimis utrinque piloso-hirtis, pedicellis terminalibus 1-cephalis, invol. squamis 2-3-seriatis, ext. foliaceis dorso hirtellis disco longioribus, intim. membranaceis, paleis apice calloso-mucronatis, ligulis invol. duplò longioribus, acheniis villosis 2-aristatis et 4-squamellatis.  $\frac{1}{2}$   $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 884.)

*ß. breviaristata*, foliis squamisque inv. exterioribus magis acuminatis paulò rigidius strigoso-hirtis, paleis in acumen subrotundum productis, acheniis villosis, pappo breviter biaristato et minutissimè subsquamellato.  $\frac{1}{2}$   $\frac{1}{2}$  in prov. Rio-Grande. An species propria? (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. n. 1037.)

10. *L. ARILLATA*, caule basi suffruticuloso adscendenti-erecto ramosissimo puberulo scabrido tereti, foliis oppositis sessilibus lineari-lanceolatis hinc inde grossè dentatis aut minutè serratis penninerviis, suprà sparè, subtùs secus nervos pubero-hirtellis, pedunculis terminalibus 1-cephalis, invol. squamis 3-seriatis, ext. lanceolatis dorso hirtellis, mediis latioribus submembranaceis, acheniis villosis radii abortivis 3-aristatis, disci obovato-oblongis 2-aristatis et paucisquamellatis basi utrinque membranè adpressà subrotundà ex areoli ortà quasi arillatis!  $\frac{1}{2}$   $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Species *L. calendulacæ* valdè affinis. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. n. 883.)

11. *L. CALENDULACRA*, caule suffruticuloso? gracili adscendente subsimplici puberulo, foliis oppositis sessilibus lineari-lanceolatis acutis utrinque villosis subscabridis integerrimis, pedunculis terminalibus solitariis 1-cephalis, invol. squamis oblongo-lanceolatis dorso puberis apice squarrosis ciliatis, ligulis 9-10, paleis lanceolatis, acheniis sericeo-villosis.  $\frac{1}{2}$   $\frac{1}{2}$  in Brasiliæ littoralis parte dictà Banda orientale legit cl. Bacle. Cor. ex sicco videntur aurantiacæ. Pappi aristæ breves. (v. s.)

12. *L. BUPHTHALMIFLORA*, caulibus suffruticulosis adscendentibus setis crassis inæqualibus præsertim ad apices hirtis, foliis oppositis sessilibus elongato-lanceolatis acuminatis serratis triplinerviis suprà sparè strigillosis secus nervulos hirtellis, ramis apice pedunculiformibus nudis 1-cephalis, invol. squamis 3-seriatis, ext. lanceolatis scabris laxis disco longioribus inæqualiter foliaceis, paleis membranaceis acuminatis, acheniis villosis biaristatis et paucisquamellatis.  $\frac{1}{2}$   $\frac{1}{2}$  in Brasiliæ prov. Rio-Grande. Capit. magnitudine colorè habitu simile cap. *Bupthalmi salicifolii* (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 879.)

13. *L. URTICIFORMIS*, caule suffruticuloso tereti hispidissimo apice scabro, foliis oppositis sessilibus lanceolatis subserratis basi obtusis trinerviis apice acuminatis suprà scabridis subtùs villosis, capitulis terminalibus solitariis, invol. squamis appendice patulà squarrosis, exter. acuminatis, inter. obtusis, acheniis 2-aristatis, squamellis intermediis minimis acutis.  $\frac{1}{2}$  in Mexico circa Villapaz propè urbem Guanaxnato legit cl. Mendez. Capit. flava, diam. sesquipedalia. Folia 2 poll. longa, 7 lin. lata. (v. s.)

14. *L. SPHEROCEPHALA*, caule herbaceo oppositè ramoso striato pubescenti-losso, foliis oppositis petiolatis ovato-lanceolatis acuminatis triplinerviis et dentatis, suprà scabris, subtùs secus nervos villosis, cæterum pubescentibus, capitulis ternato-corymbosis, invol. squamis villosis squarrosis acutis, paleis complicatis mucrone rigido recurvo, acheniis villosis ciliatis.  $\frac{1}{2}$  in Mex legit cl. Alaman. Pedicelli striati. Capitula diam. pollic. ferè globosa. Cor. ex invol. reflexo quasi exserta. (v. s.)

† *Species subdubie.*

15. *L. ? LEPTOCEPHALA*, fruticosa, ramis petiolisque pube brevi confertis, foliis oppositis petiolatis ovatis acuminatis basi breviter cuneatis etc.

utrinque sparso et minuto puberulis, paniculâ parvâ corymboâ, invol. oblongi squamis linearî-lanceolatis apice patentî-squarrosis parçè hirtis, lignis 5-6 apice dentatis, squamellis setiformibus plurimis inter aristâ setis achænium coronantibus.  $\frac{3}{4}$  in Mexico? Habui ex herb. Thib. sub nomine evid. falso Coreopidâs fœtidæ. Habitus à cæteris Leighiis differt, sed ex caract. ad nullum aptius genus adjungere quivi. (v. s.)

16. L. ? *ECLIPSAFOLIA*, caule herbaceo erecto glabro, ramulis erectis parvis versus apices adpressè puberulis, foliis oppositis sessilibus elongato-lanceolatis linearibusve adpressè et minutissimè puberulis, aliis integerrimis, aliis utrinquè undentatis et triplinerviis, capitulis terminalibus solitariis pedicellatis, invol. squamis linearî-lanceolatis apice squarrosis foliaceis, achæniis villosis nunc breviter 1-aristatis nunc exaristatis et squamellis coroniformibus constante.

(?) in Brasiliæ provinciâ Rio-Grande. Folia ferè Ecliptæ erectæ, 4 poll. longa, 4 lin. lata. Flores flavi. Pappus ex ovario judicatus à genere subdiversus. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 882.)

17. L. ? *OVATIFOLIA*, ramis teretibus glabrisculis, foliis suboppositis alterniave sessilibus distantibus ovatis acutiusculis integerrimis suprà sparso substrigillosis subtus velutino-hirantibus, pedicellis terminalibus et axillaribus apice puberulis, invol. squamis pluriseriâlibus lanceolatis acutis ciliatis adpressiusculis, lignis circ. 10 ellipticis invol. paulò longioribus, achæniis radii glabris abortivis, disci villosis 1-2-aristatis et squamellis plurimas parvas gerentibus. — in Brasiliâ prov. Sancti-Pauli. Ex invol. coriaceo adpresso paulò magis ad Harpalium fortè accedit. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 410.)

18. L. ? *SPECIOSA*, caule erecto tereti, foliis petiolatis cordatis indivisis trilobisque crenatis, pedunculo 1-cephalo supernè incrassato, invol. squamis apice foliosis acutis squarrosis, paleis acuminatissimis cuspidatis longitudine florum disci, cor. disci basi inflatis ad faucem cylindricis, achæniis obtusè 4-gonis biaristatis et brevissimè 4-squamellis.  $\frac{2}{3}$  in Mexico ad Jorullo. Helianthus speciosus Hook. bot. mag. t. 3295. Herba habitu Tithoniæ, caractere Leighiæ. Ligulæ circ. 12-13 intensè crocæ.

### † *Species exclusæ.*

L. *NICOLOR* Cass. = *Helianthus angustifolius*.

CCCKLIV. *HARPALIMUM* Cass. dict. 20. p. 299. 25. p. 437. — *Helianthi* sect. Less. — *Helianthi* sp. auct.

Capitulum multiflorum heterogamum,  $\beta$ . radii ligulatis neutris 1-seriâlibus. Invol. squamæ imbricatæ ovatæ inappendiculatæ obtusæ. Recept. convexum, paleis oblongis achænia amplectentibus. Cor. et genit. Helianthi. Achænium compressum parçè villisum. Pappus disci squamellis 2 aristæformibus ex angulis achæniî ortis et 4 brevibus obtusis intermediis. — Herbæ Americanæ alterni- aut oppositifoliæ, habitu Vigueræ et Helianthi. Genus à priori invol. pluriseriâli imbricato inappendiculato, à posteriore pappi squamellis intermediis diversum.

### Sect. I. EUHARPALIMUM.

Ligularum pappus nullus.

1. H. *RIGIDUM* (Cass. dict. 20. p. 300), caule erecto rigidè piloso, foliis infer. oppositis triplinerviis basi attenuatis, super. remotis subsessilibus lanceolatis vix dentatis rigidis pube brevi confertâ utrinquè cinereis, ramis apice nudis 1-cephalis, invol. hemisphærici squamis ovalibus obtusis imbricatis, rec. convexo, paleis obtusis, achæniis hispidis.  $\frac{2}{3}$  in Americâ boreali. Helianthus rigidus Desf. cat. h. Par. ed. 3. p. 184. Folia subpallida seu cinerascientia. Capit. æquale cap. Helianthi multiflori. Cor. disci fusca. Styli ligulæque lutei. Pappi squamellæ 2 longæ aristæformes, 4 breves oblongæ. (v. v. in hort. Girard. Genève.)
2. H. *TRUXILLENSE* (Cass. dict. 25. p. 438), caule herbaceo erecto pilosiusculo, foliis

alternis sessilibus lanceolato-oblongis acutis basi rotundatis integerrimis triplicinerviis rigidis scaberrimis, capitulis solitariis pedicellatis, invol. squamis adpressis lanceolatis extus hispidis, ligulis 12-15, paleis lanceolatis scariosis, achæniis glabro biaristato squamellis pluribus interjectis. ♀ in maritimis Peruvia juxta Truxillo. *Helianthus Truxillensis* H. B. et Kunth nov. gen. am. 4. p. 223.

3. *H. aureum* (Cass. dict. 25. p. 438), caule herbaceo, ramis hispidis-villosis, foliis alternis? petiolatis oblongis utrinque acutis subintegerrimis subtriplicinerviis rigidis supra pilosis subtus incanis et villosulo-lanatis, capitulis solariis longè pedunculatis, invol. squamis lanceolato-oblongis acutis adpressis hispidopilosis, ligulis 16-20, paleis lineari-carinatis obtusis, achæniis adpresso-pubescentibus biaristatis et plurisquamellatis. ♀ in apricis prope Alausi Quilmesan Ligulæ aurantiacæ. *Helianthus aureus* H. B. et Kunth nov. gen. am. 4. p. 224

### Sect. II. HARPALIZIA.

Achænia disci sericeo-villosissima, pappo aristis 2 et squamellis latis ovatis fimbriatis aristas ferè æquantibus constante; radii abortiva gracilia glaberrima pappo amplo coronæformi dentato superata.

4. *H. ? sericeum*, frutescens, ramis jun. sericeo-canescens, foliis alternis petiolatis oblongo-lanceolatis basi acutis apice acuminate integerrimis subtriplicinerviis adpresso sericeo-pubescentibus, pedicellis solitariis folio æqualibus 1-cephalis, invol. squamis 3-seriatis adpressis, ext. ovatis obtusis, int. obovato-mucronatis, omnibus dorso canescenti-villosis, ligulis 9-10 multistriatis. ♀ in Americâ calid. legit. cl. Née, sed locus propr. ign. Habitus ferè *Helianthi argentei*, sed achænia diversissima. Car. fruct. *H. aurei*, sed fruticoseum. (v. c.)

CCCXLV. *TITHONIA* Desf. in Juss. gen. p. 189. et in ann. Mus. h. n. Par. : p. 46. t. 4. Cass. dict. 35 (1825), p. 277. 54. p. 454 et 59, p. 147. *B. in megarth. Wein.* 1828. p. 156. Less. syn. 229. non *Rausch.* (1)

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris disci hermaphr. 5-fidis. Invol. 2-3-seriale, squamis ovatis basi multistriatis subcallosis apice foliaceo appendiculatis obtusis. Recept. convexum, paleis lanceolatis membranaceis achænia plus minus amplexantibus. Cor. disci tubo brevissimo, fauce inflatâ multinerviâ. Stylus subulati hispidi exserti revoluti. Achænia radii obcompressa seu tetragetra, pappo brevissimo parco, disci subtetragona compressa leviter pappo 1-seriali squamellis pluribus denticulatis, et in omnibus æt saltè in fl. centralibus setis 1-2 ex angulis fructus majoribus ortis coronata. — Herbaræ annuæ ex Americâ æquin. Folia alterna serrata triplicinerviâ. Rami apice pedunculiformes 1-cephali subnudi. Cor. erecta seu flavæ. — Genus *Heliantho* propinquum.

\* *Pedunculis apice obconicis incrassatis.*

1. *T. TAGETIFLORA* (Desf. ! ann. Mus. 1. p. 46. t. 4. cat. h. Par. ed. 3. p. 44) foliis plerisque trilobis, pedunculis apice incrassatis, invol. biseriali breviter velutino erecto, ligulis ovali-oblongis. (†) ad Vera-Cruz Mexic. legit olim Thierry de Menonville. Specimen habeo à cl. Bertero in Sancto-Dominico, sed forte cultum. Lam. ill. t. 708. mal. Poir. ill. 3. p. 284 et 519. n. 1c. fl. Mex. ined. Cass. dict. 54. p. 456. Don in bot. reg. t. 591 ? Ligulæ sericeo-aurantiacæ. Achænia pleraque squamellis tantum et centralia tantum insuper coronata. Folia triloba ferè Fici folia referunt. (v. v. c.)
2. *T. TUBÆFORMIS* (Cass. dict. 35. p. 277 et 54. p. 454), foliis omnibus indiv. pedunculis apice incrassatis, invol. hirsuto subpatente, ligulis oblongis. (†) Mexico (Ort.), ad Tlapujahua (Keerl in h. Mart.). *Helianthus tubæformis*

(†) Genus sine descr. videtur *Rudbeckiæ* sp.

Ort. dec. p. 101. Jacq. h. Schombr. 3. p. 65. t. 375, Lindl. bot. mag. t. 1519. Tith. helianthoides B. in mag. gart. Weim. 1818. p. 156. ex bull. For. sc. nat. 16. p. 432. *Ligula flavæ*. Achænia pleraque biaristata. (v. v. c.)

\*\* *Pedunculis cylindricis.*

3. *T. pachycephala*, caule juniore hinc indè hirsutissimo demùm scabro, foliis ovato-lanceolatis serratis atrinquè scabris trinerviis reticulatis, pedunculis teretibus, invol. squamis pluriserialibus, exter. ciliatis squarrosis, achæniis exterioribus uni-, interioribus biaristatis. ½? in Mexici prov. Leoninà ad occid. urbem Guanajuato legit cl. Mendez. Capitulum amplum crassum. Recept. hemisphæricum. Cor. omnes croceo-flavæ. Pili caulis è basi tuberculosa fasciculati decidui. (v. s. sp.)

4. *T. excelsa*, caule scabro et hinc indè bifariam villosa, foliis ellipticis acutis apice serratis triplinerviis utrinquè scabris, pedunculis teretibus, invol. squamis pluriserialibus ciliatis adpressis, achæniis plerisque biaristatis. ½ in Mexico (Cav.), ad S. Angelum (Berl.) pl. exs. n. 940. *Helianthus giganteus* Cav. ic. 3. p. 10. t. 219 non Lin. nec Lour. *Helianthus excelsus* Willd. sp. 3. p. 2243. *Helianthus trinervius* hort. Madr. et Monsp. 1813. *Herba orgyalis*. Capitula flavæ. (v. v. c. et sp.)

CCCXLVI. *HELIANTHUS* Linn. gen. n. 979. Cass. dict. 20. p. 351. 59. p. 140. — *Chrysis* Ren. 84. t. 83. — *Corona-Solis* Tourn. inst. p. 489. t. 279. — *Vosacæ* Adams. fam. 2. p. 130. — *Helianthi* sect. 1<sup>a</sup> Less. — *Helianthi* sp. auct.

Capitulum multiflorum heterogamum, fl. radii ligulatis neutris 1-ser., disci tubulosis hermaphroditis. Invol. irregulariter imbricatum, squamis ext. foliaceis acutis appendiculatis inappressis, intimis minoribus paleiformibus. Recept. planum aut convexum, paleis semimimiplectentibus oblongis acutis. Cor. disci tubus brevis, faux amplior cylindracea, dentes 5. Styli rami appendiculati. Achænia lateraliter compressa aut subtetragona, parè villosa aut glabra. Pappus squamellis 2 ex angulis achæniis ortis aristæformibus constans, et rarissimè 2 alias conformes ex angulis lateralibus ortas exhibens. — *Herbæ* (rarius suffrutices) Boreali-Americanae pleraque scabræ aut villosæ. Folia opposita, super. interdum alterna, integra aut dentata, sæpius triplinervia. Capitula ampla solitaria rarè corymbosa. Cor. disci lutæ aut fuscae, radii luteæ aut subaurantiacæ.

\* *Annui.*

1. *H. annuus* (Linn. sp. 1276), radice fibrosa, caule subimplici erecto, foliis alternis petiolatis cordatis seu latè ovatis trinerviis grossè serratis, capitulis maximis cernuis, invol. squamis latè ovatis abruptè acuminatis ciliatis, paleis subindivisis, achæniis disci biaristatis. (1) in Peru (Lob. Dod. Cam.), Mexico? (Linn.), nunc in hortis totius ferè orbis frequens et hinc indè per campos aceminibus vulgatus. Mill. illustr. ic. capituli. *H. platycephalus* Cass. dict. 20. p. 352. — Lobel. io. t. 592. f. 2. Pappi paleola dum subconcave lanceolatae denticulatae. Herba 5-6-petalis. (v. v. c.)

*H. indicus*, involucri squamis saltem exterioribus in folia subpetiolata expansis. — *H. indicus* Linn. mant. 117. Moustrum potius quàm varietas. (v. s.)

γ. *pumilus*, pumilus multò hirsutior. — *H. pumilus* Pers. ench. 2. p. 475. (v. v. c.)

2. *H. macrocarpus*, caule erecto vix ramoso strigoso, foliis alternis petiolatis ovatis acuminatis dentatis subscabris grossè triplinerviis, summis juxta capitulum subconfertis etiàm petiolatis, invol. squamis scabris ciliatis acuminatis, ligulis basi pubentibus oblongo-lanceolatis, achæniis corollas disci per anthesin duplò superantibus. (1) patr. ign. Vagat in hortis sub nom. hybrido Hel.

longicarpi. Ligulæ citrino-flavæ, apice acutè dentatæ,  $2\frac{1}{2}$  poll. longæ, 8-9 lin. latæ. Cor. disci basi villosæ, flavæ. Pappi et recept. palcæ pallidæ. (v. v. c.)

3. *H. ovatus* (Lehm. cat. h. Hamb. 1828. p. 16 et in Linnaea 1830. p. 376. pl. nov. n. 7), caule erecto subsimplici strigoso, foliis longè petiolatis ovatis acuminatis remotè subdentulatis triplinerviis strigoso-asperis, inf. oppositis, super. alternis, pedunculis terminalibus incrassatis, invol. squamis oblongo-lanceolatis setoso-ciliatis, ligulis numerosis integris. (1) in Mexicanis terris. Vagat in hortis sub nomine evidentissimè falso *Hel. petiolaris*. Capitula 3-5 poll. diam. Corollæ disci basi aurantiacæ, lobis intus subfasciæ, tubo ad partem dilatatam annis piloso cineto. (v. v. c.)

4. *H. LENTICULARIS* (Dougl. in bot. reg. t. 1265), caule erecto hispido, foliis alternis petiolatis ovatis attenuatis apice acuminatis triplinerviis serratis, capitulis amplis subcernuis, pedunculo tereti, invol. squamis expansis ovatis acuminatis dorso scabris, paleis acutè trifidis, acheniis biaristatis. (1) in Amer. bor. temperatis ad oram occident. et reg. interioribus legit Douglas, in rep. Arkansanæ ex sem. Poir. Hook. fl. bor. am. 1. p. 313. Valdè accedit ad *H. annuum*, sed capitula minora. Corollæ disci ad apicem purpureo-fusce, basi villosæ. Pappi et receptaculi palcæ apice fusco-rubentes. Achenia obovata fere per anthesin longitudine corollæ. Ligulæ intensè flavæ 2 poll. longæ, fere pilicem latæ. (v. v. c.)

5. *H. PATENS* (Lehm. ind. sem. h. Hamb. 1821. p. 8. ind. schol. 1828. p. 19), caule erecto asperrimo patentim ramoso, foliis alternis longè petiolatis orato-subcordatis serratis undulatis triplinerviis scabris subtus canescentibus, pedunculis longissimis sub capitulo incrassatis, invol. squamis linearis-lanceolatis infer. reflexis, acheniis pilosis. (1) in Americâ calid. sed loc. propr. ign. Discus fusco-purpureus. Ligulæ saturatè flavæ.

6. *H. PETIOLARIS* (Nutt. in Journ. ac. nat. sc. Philad. 1821. oct. p. 115), caule erecto ramoso-subpubescente foliis alternis longissimè petiolatis ovatis acuminatis trinerviis integrisculis scabris, invol. squamis lanceolatis acuminatis breviter scabris, ligulis circ. 20 integris. (1) in arenosis ad ripas fl. Arkansas Amer. bor. detexit cl. Nuttall. Sweet! brit. fl. gard. ser. 2. v. 1. t. 5. Desf. cat. h. Par. ed. 3. p. 403. Capitula longè pedunculata. Ligulæ flavæ magn. circ. H. multiflori. Discus fuscus. Achenia villosa. Pappus biaristatus. (v. v. c. et a. sp.)

\*\* *Perennes*, disco atropurpureo. — *Discomela Raf. neogen.* 1825. p. 3.

7. *H. ATRO-RUBENS* (Linn. sp. 1279), caule erecto pilis longis sparsis hispido, foliis oppositis oblongo-spathulatis seu ovatis crenatis triplinerviis scabris, invol. squamis ovatis acutis vix discerni æquantibus glabriusculis, ligulis circ. 15 acheniis jun. villosis biaristatis. (1) in Amer. bor. à Pennsylvania ad Carolinam in glaucosis et siccis necnon ad ripas fl. Saskatchewan (Hook.). Ker bot. reg. t. 59. H. Missuricus Spr. pug. 21. n. 44. H. diffusus Sims bot. mag. t. 2020. — Benth. t. 111. t. 94. f. 110. Discus atropurpureus. Capitula longè pedicellata subultra ex icon. recentiorum, aut brevius pedicellata ternato-corymbosa ex ic. B. Recept. dicitur convexum. Eadem fortè species ex Desf. ac H. sans rigidus, s. pappus ex Ell. biaristatus nec squamellatus.

8. *H. LATIFLORUS* (Pers. ench. 2. p. 476. n. 26), caule erecto aspero, foliis oppositis, sup. alternis ovali-lanceolatis basi cuneato-attenuatis apice longè acuminatis serratis utrinquè asperis, pedunculis elongatis 1-cephalis invol. squamis lanceolatis acuminatis ciliatis disco fere longioribus, 12-15-16, acheniis glabris biaristatis. (1) in Americâ bor. (hortul.), in Ger. occid. ? (Ell.). *H. atro-rubens* Lam. dict. 3. p. 86 non L. H. sparsiflorus sketch 2. p. 415. Discus subatro-rubens. Palcæ integræ apice callosæ resvirentes. (v. v. c.)

9. *H. ORGYALIS*, caule elato lævi, foliis alternis sessilibus linearibus planis subdentulatis vix subscabridis 1-nerviis, capitulis 5-7-corymbosis longè pedunculatis, invol. squamis linearibus acuminatis ciliatis, recept. palcæ lineæ-



achænensis integris apice subciliatis, achænio glabro 3-4- aristato.  $\frac{1}{2}$  (Linn.)  
in Amer. bor. Virginâ (Linn.), agro Arkanziano (Fourcade?). Helianthus angustifolius Linn. sp. 1279 non Michx. Coreopsis Mill. in t. 224. f. 2. Sæpè cum *H. angustifolio* Michx. confusum et stricto jure angustifolium dicendum, sed nomen special Michx. servavi nè angustar confusio. Herba 6-re-pedalis. Discus atro-purpureus. (v. v. c. in h. Genes. ex sem. Fourcadeana.)

*H. angustifolius* (Michx.) f. bor. am. 2. p. 141. caule erecto parè villosa, foliis sessilibus linearibus margine revolutis subscabridis 2-nerviis, inf. oppositis alternis alternis, capitula penda longè pedicellatis, invol. squamis lanceolatis disco aequalibus, recept. paleis ovatis apice serratis trifidis, achænio glabro biaristato.  $\frac{1}{2}$  in Amer. bor. à Virginâ ad Floridam (Michx.), Carolinæ hamidis pinetis (Ell. Boal), Nova-Cæsarâ (Torr. Green.). Ell. sketch 2. p. 415. Sims bot. mag. t. 2051. Rudbeckia angustifolia Linn. sp. 1281. excl. syn. Mill. *Leighia bicolor* Cass. dict. 25. p. 436. Discus atro-rubens. Rec. anabæntum. *Légume* 12-15. (v. v. c.)

*H. glaberrimus*, caulis pluribus stricte glabris, foliis oppositis linearibus lanceolatis basi attenuatis apice acuminatis undulatis integerrimis glande 2-nerviis utrinquè glabris margine præsertim ad basin rigidè ciliatis, capitulis solitariis terminalibus, invol. squamis imbricatis adpressis latè ovatis subscabris ciliatis. *Légume* circ. 8-10, achæniis glabris biaristatis.  $\frac{1}{2}$  in Mexico prope Reynosa de Tamaulipas apr. flor. legit cl. Berlandier. *Rhinoma ligustrina*. Caules pedales. *Légume* 5 lin. longæ, flavæ, striæ circ. 12 fasciæ notatæ. Discus subfuscus involuero longior. (v. v. c.)

\*\*\* *Perennes, disco luto, foliis omnibus oppositis.*

*H. mollis* (Lam. dict. 3. (1789) p. 81 non Willd.), caule erecto piloso, foliis oppositis sessilibus semisimplexlinearibus cordato-ovatis acuminatis adpressè serratis suprâ pubero-asperis subtus molliter villosis, invol. squamis lanceolatis linearibus hirsuto-canescens, recept. demum convexo, paleis acuminatis integris, ligulis circ. 20, achæniis glabris biaristatis.  $\frac{1}{2}$  in America bor. pratis humidis, in Alabama, Georgiâ (Ell.). Boches 12. ex hort. Paris. *H. pubescens* Willd. sp. 3. p. 2240. Ker bot. reg. t. 524 (excl. syn. Vahl. ex descr. et specim. h. Hafn.). Tausch in flor. 1828. p. 503. *H. canescens* Michx. f. bor. am. 2. p. 140. (v. v. c.)

*H. donosissimus* (Lam. dict. 2. (1789) p. 84), caule erecto basi levi æterum pubescenti-scabrido, foliis sessilibus oppositis et sup. alternis ovato-lanceolatis serratis adpressè serratis suprâ subscabridis subtus pubescentibus, invol. squamis linearibus lanceolatis puberulis margine longè ciliatis, recept. paleis trifidis lobo medio longiore acuminato, ligulis circ. 12, achæniis glabris biaristatis.  $\frac{1}{2}$  in Amer. bor. siccis subfertilibus à Pennsylvania ad Carolinam. *H. pubescens* Vahl symb. 2. (1791) p. 92 (ex spec. hort. Hafn. et conf. descr.). Hook. bot. mag. t. 2778 non Willd. *H. mollis* Willd. sp. 3. p. 2240 (non Lam. et excl. syn. Michx.). Ell. sketch 2. p. 418. *H. Hookeri* Don in Lond. h. brit. p. 358. *H. pilosus* Tausch in flor. 1828. p. 502? (v. v. c.)

*H. macrophyllus* (Willd. hort. berol. t. 70), caule erecto piloso-scabro, foliis oppositis breviter petiolatis ovato-aut oblongo-lanceolatis serratis utrinquè scabris triplinerviis, petiolo ciliato, capitulis pendis subcorymbosis, invol. squamis linearibus lanceolatis basi longè ciliatis et dorso glabriusculis apice scabridis patulo-squarrosis, paleis 3-dentatis, achæniis glabris biaristatis.  $\frac{1}{2}$  in Amer. bor. An *H. strumosus* Linn. sp. 1277. Ell. sketch 2. p. 420? — Bocc. sic. t. 27. f. 4. pensim. Zan. hist. t. 56. med. *H. serotinus* Tausch in flor. 1828. p. 504? (v. v. c.)

*H. divaricatus* (Linn. sp. 1279 excl. syn. Mor.), caule erecto glabro levi, foliis oppositis subsessilibus oblongis aut ovali-lanceolatis acuminatis serratis suprâ scabris subtus pubescentibus imâ basi ciliatis, corymbo trichotomo, invol. squamis lanceolatis brevissimè ciliatis patulo-squarrosis, paleis acute 3-dentatis, achæniis glabris biaristatis.  $\frac{1}{2}$  in Amer. bor. à Canadâ (Hook.) et

- Noveboraco ad Carolinam. Lam. diot. 3. p. 86. Tausch in florâ 1828. p. 504. *H. hispidulus* Ell. sketch 2. p. 419? *H. leiostemon* Raf. in litt. — *Moss.* ex s. 6. t. 3. f. 62 (nec t. 7. f. 66 quâ citatione falsâ decepti plurimi auct. Amer. ab hoc nom. alteram spec. foliis longè petiolatis habuerunt; tales sunt Elliott, Beck., etc.). *Radix repens.* (v. v. c. et a. sp.)
16. *H. DECAPETALUS* (Linn. sp. 1277), caule erecto lavi apice scabro, foliis oppositis ovatis petiolatis acuminatis grossè serratis triplinerviis utrinque scabris, invol. squamis lineari-lanceolatis squarrosis hispidulis ciliatis, ligulis 10. nec. paleis 3-dentatis. 2/ in Canadâ (Kalm ex L.), in siocis lapideis sylvaticis Canadâ ad Virginiam (Pursh). Affinis *H. multifloro*, sed caulis glaber, folia angustiora, ligulæ pauciores angustiores breviores. Rob. ic. t. 235 ex Pursh. (v. v. c.)
17. *H. RUDDLEIAFORMIS*, caule suffruticuloso tereti pubescenti-scabrido, foliis oppositis breviter petiolatis oblongo-lanceolatis subciliatis trinerviis acutis supra scaberrimis subtus secus nervos hirtis, capitulis pedicellatis corymbis plerisque ternis, invol. squamis appendiculatis, appendice patulo-reflexo squarrosâ, ext. acuminatis, cæteris obtusissimis, acheniis biseriatis. 3 in Mexico circa Villalpando legit cl. Mendez et Alaman. *Verbesina mollis*? Allen. in l. Accedit ad *Leighiam urticiformem.* (v. s.)
18. *H. TRUNCATUS* (Schwein. in Ell. sketch 2. p. 416), caule gracili glabro, foliis oppositis arcuè sessilibus ovatis superne attenuatis serratis pilosis scabris, capitulis paucis, invol. squamis lanceolatis, paleis indivisis apice fimbriatis. 2 in Carolinâ boreali (Schw.) et Georgiâ occid. (Ell.)
19. *H. LONGIFOLIUS* (Pursh fl. bor. am. 2. p. 571), glaberrimus, caule paucis ramis summitate oligocephalis, foliis oppositis subsessilibus lanceolatis triplinerviis integerrimis, infer. serratis, invol. squamis acutis, ext. linearibus divaricatis, ligulis circ. 10, paleis lanceolatis 3-dentatis. 2/ in humidis fertile Georgiâ occid. Ell. sketch 2. p. 417. *Habitus Coriopsisidum aquaticarum.* Pappus subulatus caducus.
20. *H. TENUIFOLIUS* (Ell. sketch 2. p. 420), caule levigato, foliis oppositis lanceolatis ovato-lanceolatis acuminatis grossè serratis utrinque scaberrimis membranaceis, capitulis parvis paucis, invol. squamis lanceolatis ciliatis compressis, ligulis circ. 10?, paleis 3-dentatis fimbriatis. 2/ in Georgiâ districtu occidentalibus. Affinis *H. spatulato*. Pappus subulatus pubescens.
21. *H. SPATULATUS* (Ell. sketch 2. p. 421), caule superne scabro, foliis oppositis in petiolum attenuatis spatulato-ovatis paulò acuminatis serratis supra scabridis subtus pubescentibus, ramulis oppositis, invol. squamis lanceolatis bipidis apice subulatis, ligulis 10-12?, paleis acuminatis integris. 2/ in Georgiâ occidentali. Pappus subulatus pubescens. Ligulæ pollicem longæ.
22. *H. TRICUSPIS* (Ell. sketch 2. p. 422), scaberrimus, caule oppositè ramoso, foliis oppositis oblongis ovato-lanceolatis subintegris margine subrevolutis, capitulis paucis, invol. squamis lato-subulatis ciliatis, ligulis 14-16, paleis pedalis tricuspidatis. 2/ in Georgiâ occidentali. Pappus subulatus lacerus.
23. *H. DIVERSIFOLIUS* (Ell. sketch 2. p. 423), caule scabro oppositè ramoso, foliis oppositis breviter petiolatis supra scaberrimis subtus pubescentibus, infer. ovato-lanceolatis acuminatis, superioribus cordato-ovatis mucronatis, invol. squamis lanceolatis ciliatis, ligulis 10-12, paleis subtridentatis apice villosis. 2/ in Georgiâ occidentali. Affinis *H. tricuspidi*. Pappus subulatus pubescens achenio longior.
24. *H. SCABERRIMUS* (Ell. sketch 2. p. 423), caule scabro parvè ramoso, foliis oppositis basi attenuatis lanceolatis utrinque scaberrimis subintegerrimis, invol. squamis ovatis adpressis, ligulis 16-20, paleis integris dorso ciliatis. 2/ in Georgiâ occid. Pappus subulatus pubescens.
25. *H. PAUCIFLORUS* (Nutt. gen. am. 2. p. 177), caule nudo trichotomo oligocephalo, foliis oppositis linear-lanceolatis acuminatis serratis fere levibus invol. arcuè imbricatis ovatis. 2/ in Louisianâ inferiore. Folia interdum ternato-verticillata, subtus pallidiora.

\*\*\*\* *Perennes, disco luteo, foliis superioribus alternis.*

3. *H. PROCUMBENS* (Pers. ench. 2. p. 475), caule diffuso-procumbente vix apice pubescente, foliis alternis subsessilibus ovalibus utrinquè obtusis basi triplinerviis apice suberratis, capitulis solitariis pedicellatis, inv. squamis biserialis pubescentibus subciliatis lineari-oblongis, paleis indivisis, acheniis tenuissimè 1-2-aristatis.  $\mathcal{L}$  in Peruvia ex hort. Par. H. prostratus Desf. cat. ed. 2 nec ed. 3. Ligulæ 6 lin. longæ. — *Sanvitalia helianthoides* Willd. sp. 3. p. 2190 hæc à Persoonio citata videtur diversa caule erecto, capitulis duplò majoribus, etc. H. procumbens Raf. fl. Lud. p. 70, omninò diversus à genere removendus sed vix notus. (v. v. c.)

4. *H. MICRANTHUS* (Spreng. syst. 3. p. 618), herbaceus, ramis retrorsum scabro-hirtellis, foliis inf. oppositis sessilibus lineari-oblongis obtusis subintegerrimis hispido-scaberrimis rigidis, capitulis solitariis longissimè pedicellatis, invol. squamis 10 lanceolatis adpressis hispidis, paleis acutis, acheniis glabris biaristatis.  $\mathcal{L}$  prope Santa-Rosa Mexicanorum. H. parviflorus H. B. et Kunth nov. gen. am. 4. p. 222. t. 378 non Bernh.

5. *H. AMPLEXICAULIS*, caule herbaceo ramoso subscabrido, foliis semiamplexicaulibus dentatis, inf. oppositis basi obtusè auritis longè supra basin triplinerviis supra auriculas angustatis dein ovato-lanceolatis ferè panduræformibus, mediis oppositis indivisis, summis alternis, paniculà laxà, pedicellis hirtellis, invol. squamis 2-3-seriatis lineari-lanceolatis acutis basi hirtociliatis adpressis, recept. paleis acuminatis integris, acheniis aristatis. — in Mexici prov. Leoninà ad occid. urbis Guanajuato legit cl. Mendez. Habitus Vignieræ sed recept. vix convexum et pappus purè biaristatus nec squamellas ullas exhibens. (v. s.)

6. *H. TRACHELIPOLIUS* (Willd. sp. 3. p. 2241), caule scabro supernè ramoso, foliis ovato-lanceolatis acuminatis serratis triplinerviis utrinquè scaberrimis, inf. oppositis, sup. alternis, paniculà laxà, invol. squamis lineari-lanceolatis ciliatis, ligulis 10-12, paleis apice 3-dentatis villosis.  $\mathcal{L}$  in Amer. bor. sylvis et palud. exsiccatis in Novâ-Cæsariâ (Torr. !), à Canadâ ad Carolinam (Pursh), in montibus (Ell.). Hook. fl. bor. am. 1. p. 312. H. gigas Michx. fl. bor. am. 2. p. 141. (v. s. et v. c.)

7. *H. CORNIFOLIUS* (H. B. et Kunth nov. gen. am. 4. p. 223), herbaceus, ramis angulatis hispidis, foliis oppositis alternisque petiolatis oblongo-ellipticis acuminatis basi rotundatis remotè denticulatis triplinerviis rigidis suprâ asperimis subtus hispidis, invol. squarrosis hispidis, paleis acuminato-pungentibus, acheniis adpressè pubescentibus biaristatis.  $\mathcal{L}$  in asperis Mexici prope Santa-Rosa de la Sierra.

8. *H. TOMENTOSUS* (Michx. fl. 2. p. 141 ? Ell. sketch 2. p. 424), caule aspero parè ramoso, foliis ovato-lanceolatis supernè attenuatis acutis serrulatis suprâ scabris subtus tomentosis plerumquè alternis, capitulis grossè pedunculatis, invol. squamis foliaceis squarrosis lanceolatis ciliatis, ligulis 12-14, paleis trifidis  $\mathcal{L}$  in Georgia occid. et inter flum. Chatahouchia et Alabama (Ell.), in pratis Illinoensium ? (Michx.)

9. *H. CALIFORNICUS*, caule elato lævi apice laxè paniculato, foliis inf. ...., sup. alternis remotis elongato-lanceolatis integerrimis basi attenuatis subciliatis apice acuminatis utrinquè scabris triplinerviis, pedunculis scabris, invol. squamis lineari-sublanceolatis disco paulò longioribus squarroso-patulis puberulo-subhirtellis, acheniis glabris biaristatis.  $\mathcal{L}$  ? in Californiâ legit cl. Douglas. Affinis *H. giganteo* sed caulis lævis, folia angustiora integerrima, invol. squamæ vix ciliatæ, etc. (v. s. comm. ab hon. Soc. hortie. Lond.)

10. *H. GIGANTEUS* (Linn. sp. 1278 non Cav.), caule elato scabro, foliis sparsis lanceolatis serratis obsolete triplinerviis scaberrimis basi attenuatis et ciliatis, corymbo stricto, invol. squamis lanceolatis linearibus disco longioribus squarroso-patulis basi villosociliatis apice scabris, acheniis glabris, squamellis 2 acuminatis subfimbriatis.  $\mathcal{L}$  in Amer. bor. à Canadâ ad Carolinam. Mill. dict. n. 5. Tausch in florâ 1818. p. 501. H. virgatus Lam. dict. 3. p. 85. H. altis-

simus Jacq. h. vind. 2. t. 163 non Linn. *H. giganteus* et *altissimus* Willd. sp. 3. p. 2242. — *Moria*. ox. 3. a. 6. t. 7. f. 66 et 67. Herba elata in hortis frequens variat caule plus minus scabro, purpurascens aut viridi; foliis angustioribus latioribusve plus minus serratis; corymbo oligo-aut polyccephalo; invol. squamis in appendicem foliaceam sæpe longissime productis, semper disco longioribus; ligulis 12-15, etc. Inter has variationes limites non satis certos vult ut varietates sint totò distinguendæ. (v. v. c. et s. sp.)

34. *H. ALTISSIMUS* (Linn. sp. 1278. excl. syn. Mor.), caule erecto glaberrimo, foliis alternis oblongo-lanceolatis in petiolum ciliatam attenuatis acuminatis obtusè serratis triplinerviis supernè scabris. invol. squamis lanceolatis d. æqualibus dorso glabris margine longè ciliatis, achæniis glabris, squamulis acuminatis fimbriatis. ♀ in Amer. bor. sed locus propr. incertus ob confusiones hujus sp. cum *H. giganteo*, *H. altissimus* Tausch in flor. 1828. p. 54. *H. giganteus* Hill. veg. syst. 2. t. 38. f. 6? Ligulæ 10-12. Paleæ apicè villosæ ciliatæ acutæ integræ. Pedicelli adpressè pilosi scabrinaculi. Discus latus. (v. v. c.)

β. foliis exsertè et argutè serratis ♀. (v. v. c.)

35. *H. MULTIFLORUS* (Linn. sp. 1277), rhizomate tereti inflexo, caule erecto ramoso scabrido, foliis petiolatis dentatis triplinerviis scabris, inf. cordatis, super. ovatis. invol. squamis ext. linear-lanceolatis ciliatis, inter. lanceolatis, ligulis oblongis numerosis. ♀ in montanis siccis sylvaticis Amer. bor. à Canadâ (Hook.), à Pensylvaniâ ad Carolinam (Pursh). Cort. bot. mag. t. 227. Hel. polyccephalus Cass. dict. 20. p. 35a. — *Fluk. phyt.* t. 159. f. 2. Tab. ic. t. 764. Invol. potentia imbricatâ nec squarrosa. Capitula erecta 2 poll. diam. Folia alterna, æternum variantia opposita, imò ternato-verticillata. (v. v. c.)

36. *H. TUBEROSUS* (Linn. sp. 1277), radice repente tubercula oblonga gerens. caule erecto ramoso scabro, foliis alternis petiolatis triplinerviis scabris serratis. inf. cordato-ovatis, super. ovato-acuminatis, petiolis basi ciliatis, invol. squamis linear-lanceolatis ciliatis. ♀ in Brasiliâ (C. Rauh.), nunc in Europâ ob æberes edules cultus sub nom. *Topinambour*. Jacq. h. vind. t. 161. Cass. dict. x. p. 353. *Geis. ic. f. 1. t. 19. Schkhar handb.* 3. t. 258. — *Col. cephr.* 2. p. 11. t. 13. Pappus 1-4-aristatus aristis ciliatis. Ligulæ rudimenta organ. farinorum retinent. Variat tuberibus subrotundis et oblongis subulcibus ex Tausch in fl. 1828. p. 506. (v. v. c.)

β. *oblongifolius*, foliis oblongo- seu ovali-lanceolatis. — Specim. sub nom. *H. tuberosi* ex horto Calcuttensi leger. d. Heyne et Wallich, cat. a. 3. Sat similis *H. decapetalo* sed folia alterna. Radices ign. (v. a. comm. a. hon. Cur. merc. ind. or. Lond.)

37. *H. HOOKERIANUS*, caule simplici striato hispido, foliis alternis oblongo-lanceolatis obtusis longissimè in petiolum attenuatis, pedunculo elongato terminali 1-cephalo, invol. squamis patentibus lato-lanceolatis ciliatis, pappo solo solitaria binisve inæqualibus validis et margine lacero constante. — in hujus planitiebus Columbiæ ad Oceanum pacificum, in vallibus montium scopulorum et in Californiâ ad San-Francisco. Hel. longifolius Hook. fl. bor. am. p. 313 non Pursh. An fortè *Leighizæ* seu *Harpalii* spec.?

\*\*\*\*\* Fruticosi.

38. *H. QUINQUERADIATUS* (Cav. ic. 3. p. 38. t. 272), caule fruticoso, juniore subtomentoso, foliis alternis petiolatis ovato-lanceolatis basi cuneatis serris crenulatis utrinquè subtomentosis, corymbo subconferto, invol. squamis biseriatis acuminatis, ext. apice reflexis, ligulis 6 ovatis, paleis rec. tridentatis obtusis. ♂ in Mexico. An hujus generis?

39. *H. ARGENTUS* (H. B. et Kunth nov. gen. am. 4. p. 221. t. 376), fruticose ramosissimus, ramulis foliisque subtus sericeo-lanatis, foliis alternis breviter petiolatis lanceolatis integerrimis trinerviis suprâ sericeo-pilosis, capsulis pedicellatis solitariis erectis, invol. squamis oblongis acutis subquarternis ligulis circ. 15, paleis oblongo-lanceolatis membranaceis, achæniis glabris

3 in Andibus Asuayensium prov. Quitensis. Achenia dicuntur biaristata. An rectè? Habitus Leighiæ.

40. *H. VRABEINOIDES* (H. B. et Kunth nov. gen. am. 4. p. 221), fruticosus ramosissimus, ramulis lanatis, foliis alternis petiolatis ovato-lanceolatis acuminatis integerrimis trinerviis supra adpresso-pilosis subtus incanis et sericeo-tomentosis, capitulis subcorymbosis, invol. squamis oblongis obtusis ext. subseparatis, ligulis circ. 10, paleis linearibus obtusis, achenio glabro. 3 in temperatis Peruviae juxta Guacabamba. An Leighiæ sp.?

† *Species non satis notæ.*

41. *H. ? ARISTATUS* (Ell. sketch 2. p. 428), caule erecto scabro, foliis infer. oppositis acutè sessilibus ovali-lanceolatis acutis dentatis scabris subtus pilosis, capitulis paucis corymbosis, invol. squamis ovato-lanceolatis pubescentibus, paleis ovato-oblongis, aristis 2 persistentibus. 2 in siccis sterilibus Georgiæ occid. Car. Actinomeridis ex Ell. sed folia non decurrentia.
42. *H. ? COCHINCHINENSIS* (Pers. ench. 2. p. 476. n. 21), caule recto scabro parè ramoso, foliis alternis lanceolatis integerrimis enerviis scabris, capitulis solitariis terminalibus, invol. squamis oblongis imbricatis patentibus squarrosis, ligulis multis longissimis, pappo biaristato. — in Cochinchinâ et Chinâ cultus. *H. giganteus* Lour. fl. coch. ed. w. 2. p. 623 non Linn. nec Cav.
43. *H. FRONDOSUS* (Linn. amœn. 4. p. 290), caule solitario infernè glabro, foliis ovatis serratis, pedunculis scabris, invol. squamis squarrosis undulatis frondosis ciliatis hispidis. 2 in Canadâ. Species dubia, hodiernis ignota et fortè varietas. *H. decapetali* invol. in folium mutato. Confer Hooker fl. bor. am. 1. p. 312.
44. *H. HIRSUTUS* (Raf. ann. nat. 1820. p. 14), hirsutus, caule simplici, foliis remotis breviter petiolatis lanceolatis acutissimis serratis subtus pallidis subtrinerviis, capitulis terminalibus solitariis subsessilibus, invol. squamis adpressis ovatis acutis ciliatis striatis. 2 in collibus Kentuckensibus.
45. *H. INCANUS* (Pers. ench. 2. p. 475), incano-tomentosus, foliis ovatis integerrimis trinerviis, invol. viloso. — in Peruvia. *H. canescens* Juss. herb. Affinis dicitur *H. molli*. Cœt. ign.
46. *H. PARVIFLORUS* (Bernh. ex Spreng. syst. 3. p. 617 non Kunth), caule herbaceo ramoso glabro, foliis oppositis alternisque petiolatis ovato-oblongis acuminate subserratis triplinerviis scabrescentibus, invol. squamis linearibus laxissimis subreflexis integerrimis. — in Amer. bor. Cœt. mihi ign. Bernh. descr. in libris mihi obvis non reperire quivi.
47. *H. TRILOBATUS* (Link enum. h. ber. 2. p. 352), caule herbaceo hirsuto superne ramosissimo, foliis longè petiolatis trilobatis scaberrimis, lobis serratis acutis, invol. squamis pilosis, ext. acutis reflexis, achenio breviter biaristato. 2 in Mexico. Capitulum in genere minimum. An certè generis?

†† *Species exclusæ.*

- H. AUREUS* H. B. K. = *Harpalum aureum*.  
*H. SUPRATHALMOIDES* Hort. = *Heliopsis canescens*.  
*H. CORDATUS* Lamb. = *Heliopsis canescens*.  
*H. CORYMBOSUS* Poepp. = *Flourensia thurifera*.  
*H. DENTATUS* Cav. = *Viguiera dentata*.  
*H. EXCELSUS* Willd. = *Tithonia excelsa*.  
*H. GIGANTUS* Cav. = *Tithonia excelsa*.  
*H. GLUTINOSUS* Hook. et Arn. = *Flourensia thurifera*.  
*H. LAEVIS* Linn. f. = *Heliopsis laevis*.  
*H. LINEARIS* Cav. = *Leighia linearis*.  
*H. MICROPHYLLUS* H. B. K. = *Leighia microphylla*.

- H. OLEIFER Wall. = *Gnizotia oleifera*.  
 H. PROSTRATUS Willd. = *Viguiera prostrata*.  
 H. RIGIDUS Desf. = *Harpalium rigidum*.  
 H. REVULARIS Poepp. = *Leptocarpa rivularis*.  
 H. SARMENTOSUS Rich. = *Wulfia membranifolia*.  
 H. SPECIOSUS Hook. = *Leighia speciosa*.  
 H. SQUARROSUS H. B. K. = *Leighia linearis*.  
 H. TETRAPTERUS Ort. = *Actinomeris tetraptera*.  
 H. THURIFER Mol. = *Flourensia thurifera*.  
 H. TRINERVIS Hort. = *Tithonia excelsa*.  
 H. TRIQUETER Ort. = *Viguiera triquetra*.  
 H. TRUXILLENSIS H. B. K. = *Harpalium Truxillense*.  
 H. TUBÆFORMIS Ort. = *Tithonia tubæformis*.

## CCCXLVII. FLOURENSIA.

Capitulum multiflorum homogamum aut heterogamum, fl. radi-  
 ligulatis sterilibus neutris seu organa fœminea rudim. gerentibus, &  
 disci hermaphr. 5-dentatis. Invol. 2-3-seriale, squamis oblongis sub-  
 æqualibus. Recept. subconvexum, paleis complicatis deciduis flore bre-  
 vioribus. Cor. disci faux vix tubo latior. Styli rami exserti elongati  
 ferè linguiformes obtusi extus parè hispidi. Achænia compressa basi  
 attenuata villosissima 1-4-aristata, aristis membranaceis lineari-subu-  
 latis ciliato-fimbriatis, 1-2 longioribus, 2 lateralibus minoribus sæpe  
 nullis. — Suffrutices aut frutices Americani glaberrimi sæpius glutinosi.  
 Folia alterna sessilia oblongo-lanceolata coriacea sæpius penninervi  
 reticulata. Pedunculi axillares terminales subcorymbosi. Corollæ om-  
 nes luteæ. — Genus spectabile dicatum clar. et amicissimo Flourensi  
 physiologo acutissimo, Acad. scient. Paris. secretario perpetuo.

\* *Capitulis radiatis*, Chilenses.

1. F. THURIFERA, glaberrima, foliis breviter petiolatis oblongo-lanceolatis re-  
 pando-serratis penninerviis, capitulis radiatis, invol. squamis oblongo-lan-  
 ceolatis. ♂ in Chili circa Valparaiso (Gay! Poepp. ! pl. exs. n. 235) et Coguibo  
 (Gaudich.!) in fruticetis sterilibus calidisque collinis (Bert.!). *Helianthus* thi-  
 rifer Mol. chil. ed. gall. p. 336 ex Gay, Bertero et Poepp. et ex usu populi. Hel.  
 glutinosus Hook. et Arn. pl. Beech. t. p. 32. *Diomedea thurifera* Bert. ! herb.  
 Colla mem. acad. Turin. 38. p. 35, n. 76. t. 31. Vulgò Chilensibus *Maravilla*  
 seu *Maravilla del Campo*. Resinam thuris succedaneam fundit. (v. s.)

♀. *angustifolia*, foliis lineari-oblongis integerrimis aut vix subserratis,  
 achæniis disci exaristatis aut 1-2-aristatis. — in Chili legerunt cl. Humb.  
 Dombey et Née. An forè species distincta? (v. s.)

2. F. CORYMBOSA, glabriuscula, foliis breviter petiolatis ovali- seu oblongo-lan-  
 ceolatis integerrimis triplinerviis minutissimè puberulis, capitulis radiatis corym-  
 bosis, invol. squamis lanceolatis. ♂ & ♀ in Amer. australi legit cl. Poeppig. *He-*  
*lianthus corymbosus* Poepp. ! pl. exs. n. 791. Caulis striatus. Capitula longè  
 pedicellata. Invol. squamæ ext. 2-3-seriales rigide adpressæ lanceolatae, intus  
 breviores. Recept. convexum. Paleæ subacutæ. Achænia villosissima longè  
 aristata, squamellis 1-2 hinc inde intermediis. (v. s. comm. à cl. Poeppig.)

\*\* *Capitulis discoideis*, Mexicanæ.

3. F. LAURIFOLIA, glaberrima, foliis petiolatis elliptico-lanceolatis integerrimis  
 acuminatis, capitulis discoideis erectis. ♂ in Mexico inter Vittoria et Tula legit

cl. Berlandier (pl. exs. n. 22051). Arbor 5-10-pedalis menae nov. florens. Folia juniora et involucria glutinosa. Invol. squamæ adpressæ subobtusæ disco breviores. Achænia villosissima, compressa aut subtriquetra, aristis 1-2 superata. (v. s. sp.)

4. *F. CRANUA*, caule ramisque subpuberis incanis, foliis subæqualibus ellipticis utrinquæ acutis integerrimis suprà glabris subtùs pubescentibus, capitulis discoideis cernuis. 5 in Mexici provinciâ Novâ-Leoninâ ad Monterey legit cl. Berlandier (pl. exs. n. 1401). Frutex ramosissimus cinerascens-viridis. Folia 7-9 lin. longa, 3-5 lata. Capitula ad apices ramorum subracemosa adgregata. Pedicelli foliosi apice recurvi. Achænia villosissima biaristata. (v. s.)

**DIV. IV. BIDENTIDÆ** Less. syn. 229. — *Coreopsidearum* gen. *Cass.* opusc. 3. p. 58.

Capitula heterogama fl. radii neutris, rarissimè homogama discoidea. Achænia apice rostrata, rarius calva, sæpius producta in aristas 1-4 ferè semper retrorsum aculeatas.

**CCCXLVIII. CAMPYLOTHECA** Cass. dict. 51 (1827). p. 475. Less. in *linnæâ* 1831. p. 508. syn. p. 230. — *Bidentis* sp. *auct.*

Capitulum multiflorum heterogamum, fl. radii ligulatis multinerviis neutris, disci hermaphr. tubulosis 5-dentatis, tubo brevi, fauce longissimâ 10-nerviâ. Recept. planum paleis inter flores instructum. Invol. pauciseriale, foliolis oblongo-ellipticis demùm reflexis. Achænia conformia elongata subcontorta aut curva subtetragono-obcompressa angustissimè alata, alis apice in mucrones minimos sæpè deciduos aut subnullos abeuntibus. — Frutices (aut fortè herbæ) ex insulis Polyneis orti, erecti trichotomi. Folia opposita ternatim biternatimve secta, segmentis ovato-lanceolatis dentatis aut integris. Capitula lutea pedicellata.

1. *C. MICRANTHA* (Cass. l. c.), fruticosa glabra, foliis ternatim pinnatimve sectis, segmento impari sæpè trilobo, fructibus incurvis apice biaristulatis. 5 in ins. O-Wahu (Cham.), et insulis Sandwicensibus (Gaudich.). *Bidens micrantha* Gaudich. bot. voy. Freyc. p. 464. t. 85. *Camp. australis* Less. in *linnæâ* 1831. p. 509. Capitula 2 lin. longa. Cor. lineam longæ.

2. *C. AUSTRALIS*, *Coreopsis* fruticosa Forst. prod. 91. *Bidens australis* Spreng. syst. 3. p. 458. An stirps australis eadem ac Sandwicensis? *Bidens pinnatifida* Forst. in h. Mus. Par.? Specim. valdè imperfectum. Differt toto cælo à pl. Gaudichandi. Caulis fruticosus gracilis bi- nec trichotomus. Folia superiora pinnatifida, lobis linearibus integerrimis. Capitula videntur minora sed fl. fract. et invol. destituta. Ex Forst. prod. crescit inter tropicos. (v. s. in h. Mus. Par.)

3. *C. GRANDIFLORA*, herbacea? puberulo-hirtella, foliis ternatim sectis, segmentis lanceolatis dentato-serratis, fructibus subspiraliter tortilibus nodis aut junioribus vix bisetosia. — in ins. O-Wyhee Sandwicensium ad montem Kaah legit cl. Macrae. Capitula 7 lin. longa. Cor. 4-5 lin. longæ. (v. s. comm. ab hon. soc. hort. Londin.)

**CCCXLIX. BIDENS** Linn. gen. n. 932. *Gertn. fr.* 2. p. 412. t. 167. Less. syn. 230. — *Bidens* et *Kerneria* *Manch. meth.* 569 et 595. *Cass. dict.* 24. p. 397 et 402. 51. p. 473 et 59. p. 321, 328 et 329. — *Pluridens* et *Edwardsia* *Neck. elem.* n. 156 et 157. — *Ceratocephalus* *Vaill. Rich. non Manch.*

Capitulum multiflorum, nunc homogamum discoideum, nunc in iisdem aut diversis speciebus sæpius radiatum, fl. radii ligulatis neutris. Invol. squamæ biseriales, seriebus similibus dissimilibusve. Recept.

planiusculum paleaceum. Styli rami cono brevi superati. Achenium plus minus obcompressum aculeatum rostro ab achenio vix diversum superatum, desinens in aristas 2-5 rigidas retrorsum piloso-scabras. — Herbae sæpius annuæ, pleraque Americanæ, paucæ Gerontogæ. Caulis tetragoni teretesve sæpius oppositè ramosi. Folia opposita, summa interdum alterna, indivisa, partita aut secta, lobis incisus serratisve. Capitula sæpius flava, rarius radio albo, in unicâ specie disco radioque purpureis.

### SECT. I. PLATYCARPÆA.

Achenia ovalia seu obovato-cuneiformia obcompressa — plana setulis omnia aut superioribus retrorsis ad margines ciliata, imâ basi vix in areolam dilatata. Capitula discoidea lutea aut radiata, ligulis flavis rarius albis. Folia sæpius indivisa rarissimè in segmenta divisa.

1. *B. TRIPARTITA* (Linn. sp. 1165), glabra, foliis tripartitis, lobis lanceolatis, dentatis, capitulis discoideis basi bracteatis, bracteis basi ciliatis capitulum superantibus, invol. squamis marg. vix membranaceis, acheniis sæpius biaristatis. (1) in Europæ fossis et inundatis, etiâ ad Caucasum (Stev.), et in Sibiriâ et Dahuriâ (Turcz.). Huds. fl. angl. 2. p. 355. opt. Sow. engl. bot. 1. 116. *B. frondosa* a Lam. dict. 1. p. 413. ill. t. 668. f. 4. *B. radiata* Thuill. fl. Par. ed. 2. p. 422. quæ discoidea in ipsissimo auct. specim. — Blakw. t. 519. Achenia, ut in *B. cernua*, obovato-cuneata obcompressa erostris. (v. v.)

2. *B. pinnatifida* (Turcz. ! in litt.), foliis tripartitis, lobis angustis pinnatis — in Dahuriâ ad flum. Argun. (v. s. sp.)

3. *B. hybrida*, foliis sub-5-partitis, lobis longo-linearibus. — circa Parisiâ *Bidens hybrida* Thuill. fl. Par. ed. 2. p. 422.

4. *B. tenuis*, capitulis paucifloris, foliis indivisis aut vix basi grossius truncatis. (1) cum var. *a.* — *Verbesina minima* Dill. ex Ray syn. angl. 1. t. 7. f. 2. *B. minima* Linn. sp. 1165. ex syn. Ray? *B. cernua* tenuis ! in litt. Similis var. *γ* *B. cernua* mihi ignotæ, sed achenia bi- nec 4-angula. Ab Hudsonio tantum rite distincta. (v. v. et s.)

5. *B. connata* (Muhlenb. in Willd. sp. 3. p. 1718), caule glabro raris foliis glabris, infimis ternatisectis lobis lateralibus in petiolum decurrentibus super. indivisis oblongo-lanceolatis, utrinque attenuatis serratis, capitulis discoideis breviter pedicellatis, invol. squamis foliaceis disco longioribus, achenio ad margines retrorsum ciliato apice 2-3-aristato. (1) in sylvis et agris Americae borealis frequens à Canadâ ad Carolinam. Ell. sketch 2. p. 430. Affinis *B. frondosæ* sed foliis et achenio facile differt. Varietatem foliis ternato-vernatis notat cl. Hooker fl. bor. am. 1. p. 314. (v. s. comm. à cl. Jauin.)

6. *B. frondosa* (Linn. sp. 1166), caule lævi sulcato, foliis glabris, infer. ternatisectis bijugis cum impari, mediis trisectis, lobis omnium lanceolatis serratis, capitulis pedicellatis discoideis erectis, invol. exter. squamis foliaceis basi ad disco longioribus, acheniis biaristatis marginibus subciliatis, pilis pro supernè retrorsis infernè erectis. (1) in humidis Americæ borealis in Can. (Hook.), et à Pennsylvania ad Carolinam. Ell. sketch 2. p. 431. Willd. sp. 1718. — Moris. ox. 16. t. 5. f. 21. mal. Invol. nunc paulò nunc multo longius. Achenia subrugosa. (v. s. sp. et v. c.)

7. *B. cernua* (Willd. sp. 3. p. 1716), glabra, foliis indivisis longe lanceolatis, capitulis cernuis basi bracteatis, bracteis subintegris capitulum superantibus, invol. squamis margine membranaceis coloratis, acheniis ad margines retrorsum ciliatis. (1) in Europæ et nunc in Americæ borealis fossis et aquis frequens. Sm. fl. brit. 858. DC. fl. fr. n. 1388

*a. discoidea*, capitulis discoideis cernuis caule ramoso. *B. cernua* Linn. sp. 1165. Engl. bot. 1. 1114. (v. v.)

*a. radiata*, capitulis radiatis. *Coreopsis Bidens* Linn. sp. 1165. (v. v.)



nna Schkuhr handb. t. 234. Barr. ic. t. 1209. — Interdum in iisdem speciminibus discoidea et radiata. Fl. dan. t. 841. (v. v.)

7. *minima*, capitulis discoideis erectis caule subsimplici. Bid. minima Linn. sp. 1165? excl. Ray. syn. — Fl. dan. t. 312. j.

5. *B. HELIANTHOIDES* (H. B. et Kunth nov. gen. am. 4. p. 230), erecta ramosa glabra, caule teretiusculo, foliis sessilibus lanceolatis serratis, pedunculis terminalibus et axillaribus 1-cephalis, capitulis radiatis, invol. squamis ovalibus obtusis, exter. foliaceis subpatentibus, int. subcoloratis, achæniis obcompressis oblongo-cuneatis margine retrorsum ciliato-serratis apice biaristatis. 2/ in paludosis et fossis Mexici ad montem Chapultepec (H. et B.), ad Yiotla (Andr.) pl. exs. n. 312, in montibus circa urbem (Berl.) pl. exs. n. 605, et in Amer. austr. circa urbem Angustura (Bert.), et circa Buenos-Ayres (Bacle!). *Kerneria helianthoides* Cass. dict. 24. p. 399. *Coreopsis serrata* ic. fl. mex. ined. Flores lutei. Ab omnibus differt achæni marginibus retrorsum serrato-scabris. (v. s.)

6. *B. CHRYSANTHEMOIDES* (Michx. fl. bor. am. 2. p. 136. non Pursh), glabra erecta, foliis oblongis utrinque attenuatis dentato-subserratis basi connatis, capitulis cernuis radiatis, invol. squamis ext. oblongis rigidè ciliatis patulis, inter. ovalibus margine membranaceis, ligulis ellipticis invol. duplò longioribus, achæniis retrorsum scabris biaristatis. (1) in oryzetis et sphagnosis Carolinæ (Ell. sketch 2. p. 430), et ad oram occid. Amer. bor. (Hook. fl. bor. am. 1. p. 314). *Coreopsis Bidens* Walt. car. 215 C. perfoliata Bosc! herb. Affinis *B. cernue* et *B. quadraristatæ*, sed achænia bi- nec 4-aristata, *B. helianthoidi* sed invol. squamæ ext. rigidè ciliatæ. (v. s. ex h. Bosc et Fraser.)

7. *B. QUADRARIATA*, glabra, foliis oblongis utrinque attenuatis dentatis basi connatis capitulis erectis, invol. ext. longo patente squamis foliaceis linguiformibus basi rigidè ciliatis, ligulis invol. interiore quadruplò longioribus ovalibus, achæniis 4-aristatis. (1) in Amer. bor. humidis, circa Bostoniam. *B. chrysanthemoides* Pursh fl. bor. am. 566. Big. fl. bost. ed 2. p. 294. non Michx. Capitula amplissima aurea. Ligulæ 12-15 lin. longæ, 6-8 latæ. Paleæ rec. apice coloratæ. (v. s.)

8. *B. BACKII* (Torr. in Spreng. neu. entd. 2. p. 135), foliis infer. capillaceo-multifidis, super. lanceolatis connatis acutè serratis, pedunculis terminalibus 1-cephalis, ligulis invol. longioribus. 2/ ? in Amer. bor. stagnosis aquis ad Schenectady Noveboracensium (Torr.), in Noveboraco occid. (Beck), et ad flum. St.-Laurent prope Mont-Real. Gold. pl. canad. in Edimb. phil. journ. 1822. apr. Beck bot. p. 207. Folia inf. omnino Ranunculi aquatilis sed opposita, sup. sæpè variè incisa. Ligulæ oblongæ, 6 lin. longæ, 1 1/2 lin. latæ. (v. s. sine fr.)

9. *B. BULLATA* (Linn. sp. 1167), caule hirtò, foliis aliis indivisis latè ovatis grossè dentatis scabris, aliis tripartitis lobis lateralibus ovali-lanceolatis parvis, terminali folia indivisa referente, capitulis terminalibus discoideis, invol. ext. squamis ovatis ciliatis disco paulò longioribus, achæniis basi sursum, apice retrorsum ciliatis, biaristatis. (1) in Italià ad Patavum, Ticinum, etc., ex Americà dicitur orta, tamen in floribus Americ. nullibi indicatur. Willd. sp. 3. p. 1723. Ard. specim. 2. p. 37. f. 18. optim. Cor. latez. Folia interdum omnia indivisa, crassa, et quasi bullata. (v. v. c.)

10. *B. MONIFLORA* (Linn. sp. 1165), caule hirtò dichotomo, foliis ovatis oblongave utrinque 1-2-dentatis, capitulis ad alas seu apices ramorum inter foliorum par ultimum sessilibus solitariis discoideis, invol. cylindrico, achæniis aristatis 4 (2 longiores et 2 breviores) gerentibus. (1) in Benghalo Willd. sp. 3. p. 1716. — Dill. elth. 5a. t. 44. f. 5a.

11. *B. REPENS* (Don prod. fl. nep. 180?), glabra, radice repente, foliis indivisis tri- aut rariùs pinnatipartitis, lobis oblongo-lanceolatis dentibus acutiusculis, terminali longiore, capitulis discoideis erectis sparsè bracteatis, invol. squamis acutis inter se æqualibus, achæniis apice biaristatis. 2/ in Nepalià. *B.*

tripartita Roxb. h. calc. p. 60. Wall. cat. et herb. n. 297. A. excl. litt. sq. Valde accedit ad B. tripartitam, sed radice repente perenni, foliorumque lobis angustioribus differt. (v. s. comm. ab ampl. aqlâ mere. angl. Ind. or.)

## Sect. II. PSILOCARPÆA.

Achenia lineari-tetragona, glabra aut margine sursum ciliata, basi vix in aulam subdilatata. Capitula raro discoides flava, sæpius radiata, ligulis flavis in paucis albis.

\* *Folii indivisi integris dentatisve sæpius lanceolatis.*

12. B. FRUTICOSA, fruticosa glabra, ramis angulatis, foliis oppositis oblongo-lanceolatis acutè dentatis in petiolum attenuatis, pedunculis axill. et termin. capitulis subcorymbosis longè pedunculatis radiatis, acheniis subtetragonis 4-arithatis. ♀ patr. ign. Coreopsis fruticosa Vest in florâ 1820. p. 7. non Font. sed auctor ipse dixit aristas retrò aculeatas undè ad Bidentem pertinet. Lig. aurantiacæ.
13. B. DECOLORATA (H. B. et Kunth nov. gen. am. 4. p. 233), caule erecto-ramoso glabro tetragono, foliis petiolatis lanceolatis serratis suprâ pilosiusculis, capitulis subcorymbosis longè pedunculatis radiatis, radio heterochromo, pedunculis pubescentibus, invol. squamis linearibus ciliatis, exter. patulis, acheniis biaristatis. ♀ in Mexico inter Capula et Valladolid de Mechoaca. Ligulæ albæ obovato-ellipticæ multinerviæ 5 lin. longæ.
14. B. TETRAGONA, glabra, caule tetragono, foliis oppositis lanceolatis, inter serratis, super. integris, breviter petiolatis subconnatis, pedunculis longiusculis solitariis, capitulis radiatis, invol. squamis exter. linearibus ciliatis, inter oblongis subcoloratis, ligulis 8 ovatis crenatis bidentatis, acheniis oblongis biaristatis. — in pratis humidis et fossis circa urbem Mexici. Coreopsis tetragona Lallav. et Lex. nov. veg. t. p. 31. Cl. auct. ipsi dicunt aristas setis instructas et ideo species Bidentis. Ligulæ flavæ.
15. B. TRIPLINERVIA (H. B. et Kunth nov. gen. am. 4. p. 231), caule erecto sulcato ramisque pubescenti-hirtis, foliis petiolatis oblongo-lanceolatis grossè serratis triplinerviis utrinquè glabris, capitulis longè pedunculatis solitariis radiatis, invol. squamis lanceolatis subhispidulis, acheniis. . . . glabris biaristatis. ♀ in Mexico circa urbem et prope St.-Augustin de Las Cuevas (H. et B. Ligulæ flavæ ellipticæ multinerviæ subsemipollicares.
16. B. HIRTELLA (H. B. et Kunth nov. gen. am. 4. p. 232), tota pubescente hirta, caule erecto sulcato subsimplici, foliis petiolatis lanceolatis serratis infernè subincisis, capitulis longè pedunculatis radiatis, invol. squamis lanceolatis adpressis, acheniis compressis biaristatis. ♀ in Mexico? cum B. triplinervia? cui valde affinis: differt foliis rigidioribus non triplinerviis et acheniis.
17. B. PROCUMBENS (H. B. et Kunth nov. gen. 4. p. 232), caule procumbente simplici 1-cephalo bifariam piloso, foliis petiolatis lanceolatis serratis subtripplinerviis hispidulis, pedunculo glabro elongato, capitulo radiato, invol. achenio biaristato. ♀ prope Xalapam Mexicanorum?
18. B. ARGUTA (H. B. et Kunth nov. gen. am. 4. p. 231), erecta glabra, caule tetragono oppositè ramoso, foliis lanceolatis petiolatis argutè serrato-dentatis, capitulis terminalibus subcorymbosis longè pedunculatis radiatis, invol. æqualis squamis lanceolatis, ext. ciliatis patulis, acheniis scabris linearibus obcompressis biaristatis. ♀ in Mexico juxta Valladolid (H. et B.), ad Iloacam et in Cordillera Guichilaquâ (Berl. pl. exs. n. 1091 et 1001), ad Tlapacote (Keerl ex h. Mart.). Cor. radii flavæ ovales obtusissimæ multistriatæ. Aristæ apice tantum glochidiatæ. (v. s. comm. à cl. Alaman et Berlandier.

♀. luxurians (Willd. enum. 847), invol. exteriore intetius separato. — in Mexico (Willd.) et forte in insulis Sandwich ex Hook. et Arn. Beech. p. 86.

19. *B. LONGIFOLIA*, herbacea glabra aut subglabriuscula, caule tereti, foliis inf. oppositis, super. alternis elongato-lineari-lanceolatis utrinque acuminatis integerrimis aut vix lobulo incisus, pedunculis longis 1-cephalis subcorymbosis, capitulis radiatis, invol. squamis ext. linearibus subpatulis scabro-ciliatis, acheniis lineari-subtetragonis glabris biaristatis. — in valle Mexicanâ legit cl. Berlandier (pl. exs. n. 717) et etiam à cl. Alaman recepti. Flores flavi. Folia distantia 3 poll. longa, 3 lin. lata. (v. s.)

\*\*\* *Foliis aliis indivisis ovatis lanceolatisve, aliis basi incisus aut trisectis, segmentis serratis.*

20. *B. HETEROPHYLLA* (Ort. dec. p. 99. t. 12), caule simplici glabro vix apice pubescente, foliis glabris, inf. pinnatipartitis, mediis tripartitis, summis indivisis, lobis lineari-lanceolatis serratis, terminali longissimo, capitulis subcorymbosis erectis radiatis, invol. squamis subæqualibus ciliatis aliis linearibus aliis lanceolatis, acheniis bi- (rarius 3- ex Ort.) aristatis. ½ in Mexico. Petioli vix subconnati basi subciliati. Capitula flava. Ligulæ oblongo-cuneatæ obtusæ disco duplò longiores. (v. v. c.)

21. *B. SERRATA* (Pavon in h. L'hér.), caule tetragono ramosissimo villosa, foliis petiolatis lanceolatis utrinque acutis subpuberulis serratis aliis indivisis aliis basi incisus imò subtrilobis, capitulis pedunculatis subsolitariis radiatis, invol. xophyllo basi imprimis piloso, squamis adpressis ext. lanceolatis int. linearibus, achenio lineari-teretiusculo bicorni. — in Peruvîâ. Coreopsis L'hér.! uss. Capit. floridum 18-20 lin. diam. Ligularum achenium calvum (L'hér.! mss.). Folia cum petiolo 2 poll. longa, 6 lin. lata. (v. s. sine fl. in h. L'hér.! et flor. descr. ex L'hér. mss.)

22. *B. DICOTOMA* (Desf.! cat. h. par. 1804. p. 104), caule tetragono glabro tri-ebotomè ramoso, foliis petiolatis glabris, aliis indivisis ovatis acuminatis serratis, aliis basi 1- seu 2-lobis accessoriis terminali minoribus acutis et bi-trisectis, capitulis subcorymbosis pedicellatis discoideis, invol. squamis inter se et disco æqualibus, ext. patulis puberis, int. glabris erectis margine scariosis, acheniis lineari-tetragonis glabris 3-4-aristatis. ½ in Brasiliâ ex specim. Commers.! Lund! Pers. ench. 2. p. 393. Poir. appl. 1. p. 630. An *B. reflexa* Linken. h. berol. 2. p. 306 ex Mexico orta? Habitus ferè *B. rubifolia*, sed discoideis. Pili pauci in ramulis. (v. v. c. et s. sp.)

23. *B. STRIATA* (Sweet brit. fl. gard. t. 237), caule erecto subtetragono striato puberulo oppositè ramoso, foliis glabris petiolatis trisectis, segmentis ovato-lanceolatis acuminatis serratis, medio paulò majore et petiolulato, capitulis corymbosis pedicellatis radiatis, invol. squamis lanceolatis puberis ext. sub-reflexis, acheniis lineari-subtetragonis margine scabris biaristatis. (1) ? in Mexico. Hook. bot. mag. t. 155. Ligulæ amplæ albæ multiradiatæ.

\*\*\* *Foliis 3-5-sectis, segmentis ovato-lanceolatis serratis non verò partitis.*

24. *B. DONDÆVOLIA* (Less. in linnæâ 1830. p. 155), glabra, ramis subtetragonis, foliis trisectis, segmentis sessilibus obovatis acutis obtusive apice serratis basi cuneatis, summis indivisis, capitulis radiatis geminis longissimè pedunculatis, invol. squamis ext. squarrosis, acheniis biaristatis. — prope Vera-Cruz.

25. *B. PILOSA* (Linn. sp. 1166), caule erecto subtetragono glabriusculo, foliis infer. impari-pinnatisectis, super. trisectis, omnium lobis glabriusculis ovatis acutis serratis vix decurrentibus, petioliis basi ciliatis, capitulis pedicellatis subcorymbosis discoideis rarius radiatis, invol. squamis inter se et disco subæqualibus, acheniis lineari-angulatis glabriusculis 2-3-aristatis, ext. brevioribus aristis divergentibus, centralibus longioribus aristis rectis. (1) in agris et cultis Americæ bor. à Pensylvaniâ ad Carolinam (Pursh), unde fortè migravit in Teneriffam (Bucht Brouss.), ins. Mauritiî (Bout.), Novam-Zeelandiam (Forst. Lesson). — Dill. elth. 1. t. 43, sed caulis nimis hispidus. Ceratocephalus pilosus Rich. cat. h. med. p. 91. *Kerneria dubia* Cass. diet. 24. p. 393. *Kerneria tetragona* Moench meth. 595. (v. v. c. et s. sp.)

26. *B. LEUCANTHA* (Willd. sp. 3. p. 1282), caule glabro subtetragono erecto, foliis glabriusculis, infer. impari-pinnatisectis 2-3-jugis, sup. unijugis, lobis omnium ovatis scutis serratis basi subæqualibus, capitulis subcorymbosis pedicellatis radiatis, radio heterochromo, invol. squamis lanceolatis inter se et disco subæqualibus, achæniis lineari-subtetragonis, aliis glabris, aliis puberulis, 2-3-jugatis. (I. in Amer. merid. (Linn.), in Chili et insulis Caribæis (Bert.), ad Bahiam in ruderatis (Salzm.), in Mexico circa Jalapam (Schied.). Ex Amer. migravit ad Maderam (Lowe!), insulas Mauritanas ubi dicitur *Herbe à Sorne* (Bory!) et ad Cap. Bonæ-Spei (Boiss. herb.). Ex hac regione habui speciem (Eckl. n. 872. Drege! u. 5084) huic speciei simillimum sed aut discoideum aut ligulis casu quodam orbatum. *Coreopsis leucantha* Linn. sp. 1282. amss. ac. 4. p. 297. *B. leucantha* H. B. et Kunth fortè diveras? *B. Chinensis* Willd. sp. 3. p. 1719 huc forsân revocanda? — Icon Plum. ed. Burm. t. 53. f. 2 huc certe pertinet et ideò *Coreopsis coronata* Linn. sp. 1281 eadem, sed species obscuræ remanet et *C. coronata* Spreng. Steud. et hortul. omninò diveras. (v. r. c. c. s. sp.)
27. *B. SUNDAICA* (Blum.! bijdr. p. 913), glabriuscula aut subpubera, foliis infer. super. ternatisectis, mediis subpinnatisectis, segmentis ovatis acuminatis serratis, lateralibus basi obliquè attenuatis, terminali ovato-lanceolato reflexo longiore, capitulis longè pedunculatis laxè corymbosis subradiatis, invol. biserialis squamis marginæ membranaceis, achæniis glabris apice triaristatis — in graminosis humidis Javæ ad Buitenzorg (Bl.), Zeylonæ (Lesch.) et fere Indiæ contin. B. tripartita Wall. cat. et herb. n. 3187. A Javanis sub *Harraja* fortè non immeritò cum *B. Chinensi* confusa et hujus forsân varietas? Achænia 4-5 lin. longa. (v. s.)
28. *B. WALLICHII*, glabriuscula aut subpubera, foliis infer. pinnatis, super. ternatisectis, segmentis ovatis acuminatis incisò-serratis dentatisve, lateralibus basi obliquè attenuatis, capitulis longè pedunculatis laxè corymbosis subradiatis, invol. biserialis squamis subæqualibus, inter. ovali-oblongis marginæ membranaceis, achæniis linearibus angulatis striatis glabris aristas 3 apice gerentibus. — in locis incultis prope Bataviam (Blume) et in Indiâ contra montes Dendigul (Wight!), Promé, Tavoy, Segain, etc. (Wall.), eandem? fallor in Casorum regione legit cl. Drege, n. 5083? *B. Chinensis* Wall. cat. n. 3189. comp. 499. DC. in Wight contr. p. 19 sed species homonyma Willd. sp. 3. p. 1719 ex Chinâ et Japoniâ orta videtur diversa. — *Agrimonia Molucca* Rumph. amb. 6. t. 15. f. 2? sed ligulæ dicuntur flavæ. Ligulæ in pl. spec. paniceæ albæ neutræ. Achænia 5-6 lin. longa. (v. s. comm. ab hon. aulâ am. angl. Ind. or.)
29. *B. PEDUNCULARIS* (Gaudich. voy. (1826) p. 464), glabra, foliis ternatis pinnatisve sectis, segmentis ovatis acuminatis argutè serratis, panicula 1-2-chotomâ, involuero exter. interiori æquante, achæniis aristas 3 apice tantum hamosas gerentibus, radio brevi. — in insulâ Rawak (Gaudich.), in O-Wallis archip. Sandwicensium (h. Acad. imp. Petrop.). *B. Sandwicensis* Less. in linn. 1831. p. 508. Simillima dicitur *B. leucantha*, sed radius luteus, achænia tantum longa, aristæ brevissimæ achænio nempè sextuplò breviores interioribus nullæ. (v. s.)
30. *B. RUBIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 237. t. 381), glabra, caule tetragono, foliis trisectis, super. indivisis, segmentis serratis supra minus lateralibus ovatis acutis sessilibus, medio maximo ovato-oblongo apice angustato-acuminato petiolato, capitulis terminalibus corymbosis radiatis, invol. squamis ext. linearibus patentibus, int. ovato-oblongis, achæniis linearibus trigonis ad angulos hispidulis biaristatis. 2? 3? in provinciâ Quitensi? *Arra-rubifolia* Cass. dict. 24. p. 399. Ligulæ flavæ. Achænia 5 lin. longa.
31. *B. TRÆTICAULIS*, caule subsicante tereti glabro, ramulis vix stratis, foliis petiolatis trisectis, segmentis ovato-lanceolatis acuminatis serratis sparse pilosulis, medio petiolulato, capitulis pedicellatis subpaniculatis radiatis, invol. squamis subæqualibus acutis, ext. reflexo-squarosis. achæniis apice radi-

biaristatis. — In Mexico prope Tantoyuca legit Berlandier (pl. exs. n. 2150 et 2148); eadem ni fallor adest ex Brasiliæ prov. Sancti-Pauli in herb. Mus. Par. (h. imp. Bras. n. 412). *B. squarrosa* Less. in Linnæa 1830. p. 156 (à sylvia Jalapensis orta) excl. utroq. *Kunthii* syn. Differt à *B. rubifolia*, caule tereti nec tetragono, à *B. squarrosa* foliis glabrisculis, ab utraq. capitulo minore, foliis etiã superis trisectis et patriâ. (v. s.)

32. *B. Conzorsinis*, glabra, caule tereti scandente, foliis oppositis petiolatis pinnatim trisectis, segmentis ovato-lanceolatis grossè dentatis acuminatis, capitula subcorymbosis breviter pedicellatis radiatis, invol. squamis ext. linearibus patulis, ligulis oblongis integris, acheniis biaristatis.  $\mathcal{L}$  in Jamaicâ (Bert.) inter sepes et dumeta (Sloan.). *Coreopsis chrysantha* Spreng.! syst. 3. p. 614 non Linn. — Sloan. hist. 1. p. 261. t. 154. f. 2 et 3. Caulis certè scandens ut jam dixit Sloane, fortè non repens. Flores lutei. Achenia immatura vidi. Aristas jam in flore evidenter retrorsum aculeato-pilose. (v. s. comm. à cl. Bert.)

? *reptans*, glabra, foliis aliis pinnatim sectis, aliis indivisis, ligulis apice denticulatis.  $\mathcal{L}$  in Jamaicâ. *Coreopsis reptans* Linn. amæn. ac. 5. p. 407 (excl. syn. *Brownei* ex Smith et Sloane! ex Ker.). Smith spicil. 2. n. et t. 22. — An hæc aut var.  $\alpha$ , in Barbados à cl. Maycock lecta?

? *incisa*, fruticosa, petiolis foliisque pilosiusculis, foliis pinnatim sectis, aliis uni-, aliis bijugis cum impari, ligulis integerrimis.  $\mathcal{L}$  in Jamaicâ. *Coreopsis incisa* Ker bot. reg. t. 7. An species propria?

33. *B. squarrosa* (H. B. et Kunth nov. gen. am. 4. p. 238), erecta, ramis teretibus glabris, foliis trisectis, super. indivisis, segmentis ovato-oblongis acuminatis serratis super. pilosiusculis subtus pubescenti-hirtis, later. subsessilibus, medio longè petiolato majore, capitulis racemoso-corymbosis radiatis, invol. squamis ext. squarroso-reflexis linearibus, int. lanceolatis, acheniis lineari-subtetragonis ciliatis biaristatis.  $\mathcal{L}$  inter. Caracas et montem Buenavista. Achenia 4 lin. longa.

34. *B. floribunda* (H. B. et Kunth nov. gen. am. 4. p. 238), glabra erecta, ramis tetragonis, foliis infer. tripartitis?..., sup. indivisis ovatis acuminatis basi angustato-cuneatis serratis, corymbis terminalibus paniculatis foliatis, capit. ...., invol. squamis ext. spatulato-linearibus patulo-reflexis, int. lanceolatis, acheniis lineari-tetragonis margine piloso-hispidis biaristatis.  $\mathcal{L}$  prope Quito ad radices montis Javirac. Affinis *B. squarrosæ* dicitur.

35. *B. andicola* (H. B. et Kunth nov. gen. am. 4. p. 237), caule erecto quadrangulo glabrisculo, foliis trisectis pubescenti-hispidis, segmentis oblongis acutis serratis basi cuneatis, lateralibus subsessilibus, medio majore petiolato, capitulis pedunculatis radiatis, invol. squamis lanceolatis, exter. hispidulis, acheniis lineari-subtetragonis scabrisculis biaristatis. ( $\mathcal{L}$ ) in declivitate montis Chimborazo. Ligulæ albæ (Kunth), flavæ (Willd.). Achenia 3 lin. longa.

36. *B. samuelpoza* (Cav. ic. 3. p. 15. t. 219), glabra, foliis decursivè pinnatipartitis, lobis oblongo-lanceolatis serratis, ultimo paulò majore, capitulis radiatis, invol. squamis æqualibus disco brevioribus, ligulis disco duplò longioribus, acheniis 4-aristatis.  $\mathcal{L}$  5 in Peruvia (Cav. L'hér. ms.). et fortè in Mexico? Cor. coccineæ. Ligulæ interdum bilabiatae. Species lobis fol. secus petiolum decurrentibus faciliè distincta. (v. v. c.)

37. *B. hispida* (H. B. et Kunth nov. gen. am. 4. p. 237), caule erecto tetragono sulcato piloso-hispido, foliis infer. impari-pinnatisectis, super. trisectis, lobis ovatis acuminatis crenato-serratis secus petiolum plerisque decurrentibus in petiolo nervisque subtus hirsutis, capitulis pedicellatis subcorymbosis discoideis, invol. squamis inter se et disco subæqualibus, acheniis lineari-tetragonis biaristatis glabris. ( $\mathcal{L}$ ) in siccis prov. Venezuela (H. et B.) et olim in hortis cnita sub nom. *B. decussatæ* Pav. ex Peruvia. Icon Dill. elth. 1. t. 43 huc fortè pertinet excl. fig. 6-10. (v. s. cum descr. in h. L'hér. nunc meo.)

38. *B. californica*, caule tetragono oppositè ramoso basi glabro apice scabrisculo, foliis parè pilosiusculis impari-pinnatisectis infer. bi-super. 1-jugis,

segmentis ovatis acuminatis serratis, capitulis pedicellatis subpaniculatis discoideis aut ligulas paucas parvas gerentibus, invol. squamis lanceolatis subaequalibus, ext. patulis, omnibus demum reflexis, acheniis lineari-tetragonis apice subscabris 3-4-aristatis. (1) in California legit cl. Douglas. Achenia 4 lin. longa. Caulis circ. pedalis à basi ramosus. Capitula parva. (v. s. comm. ab hoc soc. hortie. London.)

39. *B. EXARISTATA*, caule erecto tetragono glabriusculo, ramis oppositis tetragonis, foliis glabriusculis pinnatisectis, segmentis oppositis alternisve ovatis aut ovali-lanceolatis dentato-serratis, terminali elongato, petiolis ad has segmentisque subciliatis, capitulis pedicellatis laxè corymbosis radiatis, invol. squamis aequalibus acutis glabris, acheniis glabris lineari-subtetragonis apice attenuatis truncatis exaristatis. — in Mexico inter Vittoriam et Tulam legit Berlandier (pl. exs. n. 2220). Herba circ. bipedalis ferè omnino glabra tridactoma laxè corymbosa. Ligulæ pluriaristatæ pallidæ flavæ. Achenia longiora in iisdem capitulis varia, et etiã per anthesin calva. (v. s.)

40. *B. SUBALTERNA*, glabra erecta, caule striato, foliis inf. ..., sup. altera petiolatis pinnatisectis, segmentis sessilibus circ. bijugis cum impari, subaenariis ovali-lanceolatis basi cuneatis apice acuminatis serratis, infer. sæpè tripartitis, capitulis pedicellatis subcorymbosis radiatis, invol. squamis et brevioribus linearibus patulis pedicellisque puberis, acheniis lineari-tetragonis glabris 3-4-aristatis. (1) in locis cultis circa Bahiam legit cl. Salzmann. Ligulæ flavæ. Achenia 4 lin. longa abortiva et interdum subhirta. (v. s. comm. à Salzmann.)

41. *B. QUADRANGULARIS*, caule erecto tetragono glabriusculo oppositè ramoso, foliis petiolatis infer. pinnatisectis, segmentis 3 ovato-lanceolatis serratis petiolatis medio longius stipitato, super. pinnatipartitis lobis bijugis cum impari oblongis acutis sessilibus subserratis, capitulis pedunculatis subcorymbosis radiatis, invol. squamis ext. linearibus ciliatis subpatulis, int. lanceolatis, acheniis lineari-tetragonis glabris 4-aristatis. (1) ? circa Rio-de-Janeiro legit cl. Ponson. Ligulæ paucae ex sicco videntur albæ. Achenia 4 lin. longa, riora hinc inde 2-3-aristata. (v. s. comm. à cl. inv.)

42. *B. BRACHYCARPA*, glabra, caule erecto tetragono trichotomo, foliis petiolatis pinnatisectis, segmentis 3-4-jugis ovalibus utriusque acuminatis dentato-serratis pari paulò majore, infer. basi inciso-partitis, capitulis radiatis pedunculatis laxè corymbosis-paniculatis, invol. squamis subaequalibus lanceolatis margine scariosis, acheniis linearibus scabridis 2-aristatis invol. paulò longioribus — in Mexico ad Tampico de Tamaulipas legit cl. Berlandier (pl. exs. n. 5611). Ligulæ paucae in specim. meo disco paulò longiores, flavæ? Achenia 4 lin. longa. (v. s.)

43. *B. CARACASANA*, glabra erecta ramosissima, caule sulcato-angulato, petiolatis pinnato-trisectis, segmentis ovali-lanceolatis basi cuneato-trisectis apice acuminatis serratis, impari majore, capitulis longè pedunculatis subpaniculatis radiatis, invol. squamis int. lanceolatis margine subscariosis, ext. linearibus subreflexis, acheniis lineari-tetragonis setulis rigidis valde attenuatis apice biaristatis. (1) circa Caracas legit cl. Vargas. Ligulæ obovatæ, acutæ sicco pallidæ, disco elevato paulò longiores, patentes. Achenia 5-6 lin. longa. (v. s. comm. à cl. Vargas.)

44. *B. OXODONTA*, glabra erecta, caule tetragono subramoso, foliis petiolatis pinnatisectis, segmentis petiolatis lanceolatis utriusque acuminatis argenteo-ovatis 2-3-jugis, inferioribus sæpè trisectis, terminali indiviso aut tripartito, capitulis pedicellatis in paniculam dispositis radiatis, invol. squamis linearibus subaequalibus, ext. subpatulis, acheniis lineari-tetragonis apice subscabridis biaristatis. — in Sancto-Domingo legit Bertero. Coreopsis et Bidens Spreng. syst. 3. p. 614 sed ab icone Plum. à Linnaeo citatâ certe distat. Bidentis nec Coreopsidi adgreganda. Ligulæ pallidæ in sicco. (v. s.)

45. *B. ANTHRISCOIDES*, tota pubescenti-hirta, caule suberecto tetragono, petiolatis trisectis, inferiorum segmentis ovatis acutis subaequalibus ad

dissimili, superiorum segmentis lateralibus ovatis medio duplò longiore petiolato lanceolato utrinque attenuato, omnibus dentatis, ramis floridis nudis elongatis 1-3-cephalis discoideis, invol. squamis disco et inter se subæqualibus dorso pubescenti-hirtellis, achæniis lineari-tetragonis spice scabris biaristatis. (X) in Mexici Cordillera de Guachilagua legit cl. Berlandier (pl. exs. n. 1070 et 1152) octob. flor. Capitula parva, in sicco flavo-subfusca. Radix obliqua seu caulis stricto sensu adscendens sed ferè erecta. Achæn. 6 lin. longæ. (v. s.)

*B. angustiloba*, caule foliisque glabrisculis, foliorum lobis elongato-lanceolatis in apice caulis ferè linearibus. (X) in Mexico circa urbem legit etiam Berlandier (pl. exs. n. 875). (v. s.)

46. *B. PORTORICENSIS* (Spreng. in litt. ad Balb.), caule glabro erecto tetragono, foliis glabris petiolatis, aliis biternato-sectis, aliis impari-pinnatisectis, omnium segmentis ovatis acutis basi cuneatis dentato-serratis, capitulis pedicellatis discoideis, invol. squamis disco brevioribus acutis inter se æqualibus, achæniis lineari-tetragonis glabris 4-aristatis. (X) in Porto-Ricco legit cl. Bertero. Valdè affinis *B. bipinnatæ*, sed satis distincta notis indicatis videtur. (v. s. comm. ab amic. Bertero.)

47. *B. DAUCIFOLIA*, caule suberecto tetragono sulcato pubescenti-hirtello, foliis villosa-hirsutis pinnatipartitis, lobis circ. 3-jugis cum impari lineari-lanceolatis inciso-serratis acutis, pedunculis paucis 1-cephalis, capitulis radiatis, invol. squamis disco brevioribus inter se æqualibus ciliatis, achæniis lineari-tetragonis striatis scabriusculis biaristatis. — in Mexico legit cl. Berlandier fortè in valle Toluacanà (commixt. sub n. 1138). Ligulæ disco longiores ovales obtusissimè multistriatæ flavæ 3-4 lin. longæ. (v. s.)

48. *B. RIPARIA* (H. B. et Kunth nov. gen. am. 4. p. 236), erecta, caule ramisque glabris, foliis duplicato-trisectis supra pilosiusculis margine ciliato-scabris, lobis ovatis inciso-serratis, sup. trifidis, lobulis terminalibus angustato-acuminatis, capitulis pedunculatis radiatis, invol. hispidi squamis exter. linearibus duplò longioribus, achæniis lineari-tetragonis 5-aristatis, aliis glabris, aliis brevioribus scabris. (X) in calidissimis ad ripam flum. Magdalenæ prope Badillas.

49. *B. ABADIA*, caule tetragono sulcato hispido erecto ramoso, foliis utrinque villosa-hirsutis bipinnatisectis segmentis 2-3-jugis ovatis grossè inciso-dentatis cum terminali trilobo basi cuneato, dentibus obtusis mucronatis, capitulis pedunculatis laxè et rigidè paniculatis radiatis, invol. squamis glabris lauceolatis subæqualibus disco brevioribus, achæniis lineari-subtetragonis vix involucri longioribus apice subciliatis biaristatis. — in Peruvia circa Limam legit cl. Abadia. Ligulæ ex sicco albæ 5-striatæ. Caulis apice subglabratus. (v. s. comm. à cl. Abadia.)

\*\*\* *Foliis variè sectis, segmentis angustis integris linearibus aut multifidis, lobis linearibus subintegris.*

50. *B. ANGUSTISSIMA* (H. B. et Kunth nov. gen. et. sp. 4. p. 233), erecta glabra, ramis tetragonis, foliis tripartitis, lobis angustissimè linearibus acuto-mucronatis integerrimis subcarneis, intermedio longiore, capitulis terminalibus longè pedunculatis radiatis, invol. squamis exter. patentibus brevioribus, int. oblongis extus hispidulis, achæniis compressis biaristatis. (X) in frigidis Mexici prope Los Joares et Santa-Rosa de la Sierra (H. et B.). Cor. flavæ. Achænia matura ign.

51. *B. CRITHMIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 234), caulibus diffusis procumbentibus hirtò-pubescentibus, foliis pinnatis quinque-sectis glabris, lobis linearibus integerrimis, capitulis longè pedunculatis solitariis radiatis, invol. squamis exter. linearibus obtusis, inter. lanceolatis, achæniis 4-gonis ? biaristatis ? (X) prope Quito. Ligulæ flavæ.

52. *B. GLABERRIMA*, herbacea erecta ex omni parte glabra, ramis (caulibusve) teretibus, foliis oppositis petiolatis pinnato-multifidis internodio multò longioribus, lobis oblongo-linearibus crassiusculis, capitulis solitariis pedicellatis terminalibus radiatis, invol. squamis ext. linearibus obtusis subpetalis, int.

oblongis, ligulis invol. triplò longioribus multinerviis, acheniis glabris biaristatis. — in Amer. calidiore legit el. Née ex h. Thibaud, sed locus propr. ign. Ligulæ flavæ ellipticæ obtusæ, 6-7 lin. longæ, nervulis 15-16 nigrellis striatæ. (v. s.)

53. *B. NUMILIS* (H. B. et Kunth nov. gen. am. 4. p. 234), diffusa subprocumbens glabra, foliis irregulariter bipinnato-quinque-partitis, lobis linearilanceolatis, petiolis ciliatis, capitulis longè pedunculatis solitariis radiatis, invol. squamis ext. linearibus obtusis, int. lanceolatis, acheniis lineari-quadrangularibus scabriusculis biaristatis.  $\mathcal{L}$  in radicibus montis Cotopaxi prope Malab. Quitensium. Ligulæ flavæ oblongo-ellipticæ multinerviæ 3 lin. longæ. Achenia 2; lin. longa.

54. *B. SCANDICINA* (H. B. et Kunth nov. gen. am. 4. p. 235), caule erecto angulato glabro, ramis pilosis, foliis pinnato-quinque-partitis subtus hispidis, lobis incisus, inf. tripartitis, petiolis ciliatis, pedunculis pilosis, capitulis radiatis, invol. hispiduli squamis ext. linearibus, int. lanceolatis, acheniis linearibus compresso-quadrangularibus scabriusculis 2-3-aristatis. (I) prope Llacungu et Hambato Quitensium. Ligulæ flavæ ex Kunth, albæ ex Bonpl. Achenia 5 lin. longæ.

55. *B. ALAUSENSIS* (H. B. et Kunth nov. gen. am. 4. p. 235), caule erecto glabro, ramis tetragonis, foliis pinnato-quinque-partitis pilosis, lobis oblongis inciso-serratis, petiolis ciliatis, capitulis longè pedunculatis radiatis, invol. squamis ext. linearibus, int. lanceolatis, acheniis linearibus compresso-quadrangularibus scabriusculis biaristatis.  $\mathcal{L}$  prope Alausi Quitensium (H. B.), et in summo Serro-Colorado Mexici (Schied.). *B. diversifolia* Willd. brò Ligulæ flavæ. Achenia 4 lin. longæ. Tota plūs minūs hirsuta ex Less. in Lemée, p. 155.

56. *B. DELPHINIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 234), erecta glabrescens, caule sulcato supernè ramoso, foliis irregulariter bipinnatisectis, lobis linearibus, capitulis longissimè pedunculatis... invol. ..., acheniis linearibus tetragonis scabriusculis triaristatis (I) ? ad urbem Quito ? Ligulæ flavæ.

57. *B. PARVIFLORA* (Willd. enum. 840), glabra, foliis multipartitis, lobis linearibus oblongo-lanceolatis incisus, summorum linearibus, capitulis discoideis longè pedunculatis laxè corymboso-paniculatis, invol. biseriali, interdum exterior longè superante, acheniis sparse pubescentibus longè aristatis 2 rectas apice gerentibus. (I) in Sibiria ad lacum Baikal (Fisch.) in pibus ad Selengam (Turcz.!), in China (Staunt.!). *B. macrosperma* Kuhn. hort. Gorenk. *B. multifida* Desf. ! cat. h. Par. ed. 3. p. 186. *B. pauciflora* Petrop. suppl. 1. p. 630. *B. Messerschmidii* Turcz. ! in litt. *B. chinensis* herb. Petrop. sed vix Willd. Flores 3-6 quadridentati. Achenia inæqualia centum majora, 8-9 lin. (aristis excl.) longæ. (v. v. c. et s. sp.)

58. *B. DECOMPOSITA* (Wall. ! cat. et herb. n. 298), caule glabro tetragono ramoso, foliis bipinnatisectis, lacinis lanceolatis basi cuneatis mucronatis utrinque scabridis, capitulis longè pedunculatis laxè subpaniculatis discoideis, invol. biseriali demum reflexo, acheniis elongatis biaristatis, aliis brevibus stricte scaberrimis, aliis basi glabris apice subtrigosis in eodem capitulo. — in orient. ad Monghir prov. Bahar et in Nepalia Coreopsis corymbifolia Smilt. ! ex Wall. (v. s. comm. ab hon. aulæ merc. angl. Ind. or.)

59. *B. GRANDIFLORA* (Balb. cat. h. taur. 1812. p. 19), caule suberecto teretibus, foliis glabris impari-pinnatisectis, segmentis 3-5-jugis, aliis oblongo-serratis aut subincisis, aliis in lobulos lineares pinnatisectis, capitulis pedunculatis radiatis, invol. squamis ciliatis disco subbrevioribus, ext. patentibus, acheniis lineari-subtetragonis glabris 2-4-aristatis. (I) in Mexico circa urbem Mexici exs. n. 1168). DC. cat. h. monsp. p. 83. Horn. h. batn. 2. p. 788. *B. colorata* Dum. Cours. bot. cult. 4. p. 241 non Cav. Coreopsis diversifolia Schlegel. 1. p. 80. t. 54. opt. *Bideus diversifolia* hortul. *B. serrulata* Cass. cat. h. par. 1829. p. 186. *Kerneria serrulata* Cass. dict. 51. p. 4-4. *Cuscuta* Sims bot. mag. t. 1689. — Ab hac ex desc. videtur diversa Coreopsis



- rulata Poir. suppl. 3. p. 35a etiam à cl. Desf. admissa. Canlis sæpè basi radicatis. Ligulæ 9-10 lin. longæ, 3-3½ latæ, 9-10 nervulis striatæ. Variat invol. glabro et extus plùs minùs hirsuto. (v. v. c. et s. sp.)
60. *B. ODOBRATA* (Cav. ic. 1. p. 9. t. 13), glabra erecta, caule tetragono ramosissimo, foliis bipinnatisectis lobis cuneato-trifidis, capitulis pedunculatis solitariis radiatis, invol. squamis lanceolatis apice ciliatis, exter. subreflexis, achæniis lineari-subtetragonis ad angulos scabris biaristatis, lateralibus brevioribus. (1) in Mexico (Cav.) fortè etiàm in insulis Societatis ex Hook. et Arn. bot. Beech. p. 66? *Coreopsis odorata* Lam. ill. t. 704. f. 1. Accedit ad *B. grandifloram*, sed radius albus, flores odorati, achænia scabra, et fol. lobi cuneati.
61. *B. FRUTULIFOLIA*, glabra, caule herbaceo teretiusculo, foliis oppositis bipinnatisectis, lobis linearibus vix sublancoelatis integerrimis costam segmentorum latitudine non superantibus, capitulis pedicellatis laxè corymbosis radiatis, invol. squamis linearibus demùm patulis, int. ovali-lanceolatis margine membranaceis, ligulis obovatis obtusis multinerviis invol. quadruplò longioribus, achæniis linearibus glabriusculis biaristatis. (2) in Mexico (Willd.), *Coreopsis frutulaefolia* Jacq. h. Schænb. 3. p. 65. t. 375. bot. mag. t. 2059. Quid sit var. odoratissima à Persoonio citata ignoro. *Kerneria frutulaefolia* Cass. dict. 51. p. 43. Aristæ certè retrorsum scabræ in ipso specim. à Jacquinio filio comm. et apprimè iconi conveniente. An idèò aliam plantam sub oculis habuit cl. Lessing cnm inter *Coreopsides* veras admisit? (v. s. et v. c.)
62. *B. PROCERA* (Don in bot. reg. t. 684), glabra striata ramosa, foliis decursivè 2-3-pinnatisectis, lobis linearibus acutis canaliculatis integris flaccidis, capitulis subcorymbosis pedicellatis amplè radiatis, invol. squamis ext. linearibus obtusis pubescentibus, achæniis glabris compresso-cuneiformibus biaristatis. (2) in Mexico. *Coreopsis angustifolia* Pavon in h. Lamb. ex Don. Ligulæ flavæ 10 lin. longæ. Ex habitu et lobis foliorum integris ad *Cosmum* pertinere videtur, sed achænium (junius equidèm descriptum) potiùs *Bidentis*.
63. *B. FOMICULIFOLIA*, caule erecto acutè 4-6-angulato glabriusculo, ramis acutè tetragonis subpuberulis, foliis oppositis bi- aut tripinnatisectis glabriusculis, lobis linearibus acutis integerrimis aut hinc indè lobulatis, capitulis sæpiùs ternis pedicellatis subcorymbosis radiatis, invol. squamis linearibus ciliatis, inter. ovato-lanceolatis margine scariosis longit. æquantibus, ligulis obovatis obtusis multinerviis inv. triplò longioribus, achæniis lineari-tetragonis glabris vix apice scabridis biaristatis. — in Mexico ad occid. urbis Guanaxuato in prov. Leoninà leger. et mecum comm. cl. Mendez et Alaman. *Coreopsis fomiculacea* ic. fl. mex. ined. Achænia 3-3½ lin. longæ. Flores intensè lutei. (v. s.)
64. *B. CYNAPIIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 235), glabra, caule erecto subtetragono, ramis floridis divaricatis, foliis irregulariter tripinnatispartitis, lobis oblongis basi cuneatis, terminali longè acuminato, capitulis longè pedunculatis breviter radiatis, invol. squamis linearibus ciliatis disco subæqualibus, exter. reflexis, achæniis lineari-tetragonis 4-aristatis, aliis elongatis glabris, aliis brevioribus pilosulis. (1) in Insulâ Cubâ prope Guanaxacoa (H. et B.), circa Havanam (La Sagr.!). Ligulæ parvæ angustæ, in sicco pallidæ. (v. s.)
65. *B. MIPINNATA* (Linn. sp. 1166), glabra, caule erecto obtusè tetragono, foliis bipinnatisectis, lobis lanceolatis, capitulis pedicellatis radiatis, ligulis paucis (2-4) vix disco longioribus, invol. squamis subinæqualibus disci circ. longitudine, ext. reflexis, achæniis lineari-tetragonis glabris 3-4-aristatis. (1) in sicca agris Amer. bor. frequens; nunc in Senegaliâ (Pers.!), et in agrum Tyrolense quasi spontè introducta (Koch!). Ell. sketch 2. p. 432. — Herm. parad. t. 123. Moris. oxon. s. 6. t. 7. f. 23. Ligulæ flavæ, primâ fronte vix perspicuæ. (v. s. sp. et v. c.)
66. *B. CHILENSIS*, glabra, caule erecto tetragono sulcato vix ramoso, foliis subciliatis petiolatis biternatisectis seu pinnatisectis, segmentis bijugis cum impari tripartitis, lobis obovatis basi cuneatis acutis grossè serratis dentibus mucrona-

B. SCANDENS Br. = *Salmon scandens*.

B. SCANDENS Sw. = *Salmon eupatoriæ*.

B. TENUIFOLIA Labill. = *Glossogyne tenuifolia*.

CCCL. COSMOS Cav. ic. 1. (1791). p. 9. t. 14 et 79. *Cass. dict.* 11. p. 4. — *Cosmos Willd. sp.* 3. (1803). p. 2250. *Hoffmans. verz.* 1824. p. 206 *Less. sp.* 230. — *Cosmos Pers. ench. a. n.* 1917. — *Georgia Spreng.*

Capitulum multiflorum radiatum, ligulis neutris. Invol. duplex, utrumque squamis 8-10 basi plūs minus concretis apice acuminatis. Recept. planum paleaceum, paleis membranaceis in filum elongatum productis. Stylī rami apice incrassati hispidi in conum subulatum producti. Antheræ apice appendice scariosa cordatâ superati. Achænium tetragenum exalatum rostratum et interdum stipitatum, aristis 2-4 retrorsum piloso-scabris deciduis coronatum. — Herbariæ Americanæ annuæ glabræ aut vix pilosulæ elatæ ramosæ. Folia bipinnatisecta, lobis linearibus lanceolatis sæpius margine integerrimis. Capitula ad apices ramorum longè nudorum solitaria. Discus intensè luteus. Radius versicolor.

1. C. REPENSATUS (Cav. ic. 1. p. 9. t. 14), glaberrimus, foliis bipinnatisectis, lobis linearibus remotis integerrimis, invol. ext. squamis ovato-lanceolatis acuminatis, achæniis glabris breviter rostratis triaristatis. (I) in Mexico (Cav.), Georgia bipinnata Spreng. syst. 3. p. 612. *Coreopsis formosa Bonate in diss. Pisaur.* p. 21. t. 2. Ligulæ 7-9, amplæ, 7-8 lin. longæ, purpureæ roseæ seu albide nunquàm flavæ.

f. *exaristatus*, aristis ad apicem achæniū nullis. (I) in Mexico (Alam.) ad Pezenaro (H. et B.), Oaxaca (Andr.! pl. ex. n. 307). S. Angel. (Berl. pl. ex. n. 938). Hæc var. sola in hortis mihi notis nunc colitur et sola invenit in herb. Mexic. (v. s. sp. et v. c.)

2. C. PARVIFLORUS (H. B. et Kunth nov. gen. am. 4. p. 241), glaberrimus, foliis bipinnatipartitis, lobis capillaceo-linearibus integerrimis, invol. squamis ext. 8 lanceolatis acutis brevioribus, achæniis glabris ad angulos scabridis breviter rostratis 3-aristatis. (I) in Mexico (Jacq.) inter Puerto-de-Variantos et S. Juan de la Sierra (H. et B.). *Coreopsis parviflora Willd. sp.* 3. p. 2251. Ligulæ roseæ aut albæ, 3-dentatæ, 3 lin. longæ. (v. s. et v. c.)

3. C. TENELLUS (H. B. et Kunth nov. gen. am. 4. p. 240), foliis bipinnatipartitis glabris, lobis lanceolatis acutis, inferioribus basi subserratis, terminalibus superioribus, invol. squamis exter. lineari-spathulatis brevioribus, achæniis scabris.... aristis.... (I) in frigidis Mexici inter urbem et Huehuetoca. Ligulæ roseæ?

4. C. CAUDATUS (H. B. et Kunth nov. gen. am. 4. p. 240), caule glabro, foliis bipinnatipartitis, lobis lanceolatis cuspidatis margine scabris, petiolis ciliatis, invol. squamis exter. lineari-lanceolatis acutis ciliatis patulis vix basi concretis interiores ferè æquantibus, achæniis glabrescentibus apice scabris longissimè rostratis, aristis 3 patulis. 2/ in insulâ Cubâ juxta Havanam (H. et B.), Porto-Rico (Berl. Wydl.), Sancti-Thomasii agris saccharo cultis (Less. in *linnæa* 1801. p. 72<sup>1</sup>). Martinicâ (herb. Duby!), etiâ in Indiâ or. (ex h. Wight) et insulâ Mauritiî (Bont!), sed introduct. *Bidens Berteriana Spr.*! syst. 3. p. 454. DC. in Wight contrib. p. 19. Ligulæ roseæ apice trifidæ. Achænia ferè pollicem longæ. Flores disci 12 ex Kunth, circ. 20 in sp. meis. (v. s. sp.)

5. C. SELPURUS (Cav. ic. 1. p. 56. t. 79), caule piloso, foliis bipinnatipartitis, lobis lanceolatis mucronatis margine subsabridis, petiolis ciliatis, invol. squamis ext. linearibus adpressis quàm interiores adhuc brevioribus et multi angustioribus, achæniis glabrescentibus apice scabris longè rostratis, aristis 3 patulis. 2/ in Mexico (Cav.) circa Oaxacam (Andr.! pl. ex. n. 306) ad Villalpando (Mendel!), *Cosmos selpurus Willd. sp.* 3. p. 2250. *Coreopsis selpurus Jacq. ic.* 3. t. 59. f. 1. Achænia ferè pollicem longæ, apice puberis. Achæniū g.

petiolatis lato-lanceolatis acuminatis glabris serratis acumine integerrimo, paniculâ patente terminali, capitulis parvis parçè radiatis, involacro brevi, acheniis dentibusque 2 brevibus hispidis. 3 in insulâ Otahiti. — Mixtum eam succo nucis Cocœ et ebullitione coctum est catharticum, motu dictum.

72. *B. TENELLA* (Linn. amœn. ac. afr. 47), foliis linearibus, pedunculis capillaribus, invol. sub-4-phyllis, acheniis 5 erectis. ① ad Cap. Bonæ spei. †.
73. *B. ? CHRYSANTHA*, foliis ternatisectis segmentis ovato-oblongis serratis, capitulis longè pedunculatis subcorymbois, radio concolore. — in insulis Caribæis. *Coreopsis chrysantha* Linn. sp. 1282. — Plom. ed Burm. t. 53. f. 1. Affinis videtur *B. scandenti*.
74. *B. ? LEUCORHIZA*, caule herbaceo erecto tetragono, foliis quinquatis lanceolatis serrato-incisis, pedunculis oligocephalis, ligulis 6 ovatis integerrimis acheniis 3-cornutis. — prope Cantonem Sinarum. *Coreopsis leucorhiza* Lour. fl. coch. 622. Radix perpendicularis crassa alba. Capitula integrè crocea. Ligulæ neutre. Aristæ setis inversis ramosæ, undè ad *Bidentem* potiùs pertinere videtur.
75. *B. ALBA*, caule debili flexuoso, foliis oppositis petiolatis trisectis rariùs 5-sectis, segmentis obovatis serratis, capitulis pedicellatis radiatis, ligulis 8 obovatis, acheniis linearibus biaristatis. — in insulâ Sanctæ-Crucis. *Coreopsis alba* Linn. sp. 1282. — Herm. parad. 124. t. 124. Pluk. alm. t. 160. f. 3. Ligulæ albæ. Aristæ more *Bidentium* ex descr. Merm. scabræ. Affinis videtur *B. Portoricensis* et fortè hujus varietas radiatæ. — Syn. Brownei à Linnæo citatum videtur alienum ex recept. nudo dicto.
76. *B. ? VERTICILLATA* (Linn. sp. 1167), caulibus procumbentibus subsimplicibus, foliis inf. alternis, longè sup. oppositis breviter petiolatis lanceolato-ovatis subintegris suprà glabris subtùs tomentoso-albidis, capitulis in axillis sup. binis discoidis, achenio tenni setis a patulis coronato. — in Verâ-Cruce. *Bidens* Linn. hort. Cliff. 399. n. 4.

### †† *Species exclusæ.*

- B. ACMELLA* Lam. = *Spilanthes acmella*.  
*B. ACMELLOIDES* Big. = *Spilanthes oleracens*.  
*B. AGUSTIFOLIA* Lam. = *Spilanthes cereus*.  
*B. APIIFOLIA* Linn. = *Chrysanthellum procumbens*.  
*B. ARBORNA* Roxb. = *Petrobium arboreum*.  
*B. ATRIPLEXIFOLIA* Linn. = *Isocarpha atriplexifolia*.  
*B. AUSTRALIS* Spreng. = *Campylothea micrantha* f.  
*B. BARDANA* Wall. = *Chrysanthellum Indicum*.  
*B. BERTERIANA* Spreng. = *Cosmos caudatus*.  
*B. CROCATA* Cav. = *Verbesina crocata*.  
*B. FRUVIDA* Lam. = *Spilanthes oleracea*.  
*B. FRUTESCENS* Mill. = *Salmea scandens*.  
*B. FUSCA* Lam. = *Spilanthes oleracea fusca*.  
*B. HIRSUTA* Sw. = *Salmea hirsuta*.  
*B. INSIPIDA* Stend. = *Spilanthes insipida*.  
*B. MICRANTHA* Gandich. = *Campylothea micrantha*.  
*B. NIVEA* Linn. = *Melanthera hastata*.  
*B. NIVEA* Sw. = *Melanthera deltoidea*.  
*B. OCYMIFOLIA* Lam. = *Spilanthes alba*.  
*B. PINNATIFIDA* Wall. = *Glossogyne pinnatifida*.  
*B. PINNATIFIDA* Forst. = *Campylothea micrantha* f.  
*B. RAMTILLA* Wall. = *Guizotia oleifera*.

B. SCANDENS Br. = *Salmea scandens*.

B. SCANDENS Sw. = *Salmea eupatoria*.

B. TENUIFOLIA Labill. = *Glossogyne tenuifolia*.

CCCL. COSMOS Cav. ic. 1. (1791). p. 9. t. 14 et 79. Cass. dict. 11. p. 4. — *Cosmea Willd. sp. 3.* (1803). p. 2250. *Hoffmans. verz. 1824. p. 206* Less. *fl. 230.* — *Cosmos Pers. ench. 2. n. 1917.* — *Georgia Spreng.*

Capitulum multiflorum radiatum, ligulis neutris. Invol. duplex, utrumque squamis 8-10 basi plus minus concretis apice acuminatis. Recept. planum paleaceum, paleis membranaceis in filum elongatum productis. Styli rami apice incrassati hispidi in conum subulatum producti. Antheræ apice appendice scariosâ cordatâ superati. Achænium tetragonum exalatum rostratum et interdum stipitatum, aristis 2-4 retrorsum piloso-scabris deciduis coronatum. — *Herbæ Americanæ annuæ glabræ aut vix pilosulæ elatæ ramosæ. Folia bipinnatisecta, lobis linearibus lanceolatis sæpius margine integerrimis. Capitula ad apices ramorum longè nudorum solitaria. Discus intensè luteus. Radius versicolor.*

1. C. BIPINNATUS (Cav. ic. 1. p. 9. t. 14), glaberrimus, foliis bipinnatisectis, lobis linearibus remotis integerrimis, invol. ext. squamis ovato-lanceolatis acuminatis, achæniis glabris breviter rostratis triaristatis. (I) in Mexico (Cav. *Georgia bipinnata* Spreng. syst. 3. p. 611. *Coreopsis formosa Bonato in d. Pisaur. p. 21. t. 2. Ligulæ 7-9, amplæ, 7-8 lin. longæ, purpureæ roseæ subbidæ nunquam flavæ.*

β. *exaristatus*, aristis ad apicem achæniî nullis. (I) in Mexico (Alam. d. Pezcuaro (H. et B.), Oaxaca (Andr.! pl. ex. n. 307). S. Angel. (Berl. p. ex. n. 938). Hæc var. sola in hortis mihi notis nunc colitur et sola venit in herb. Mexic. (v. s. sp. et v. c.)

2. C. PARVIFLORUS (H. B. et Kunth nov. gen. am. 4. p. 241), glaberrimus, foliis bipinnatipartitis, lobis capillaceo-linearibus integerrimis, invol. squamis 8 lanceolatis acutis brevioribus, achæniis glabris ad angulos scabridis breviter rostratis 3-aristatis. (I) in Mexico (Jacq.) inter Puerto-de-Varientos et Sierra de la Sierra (H. et B.). *Coreopsis parviflora Willd. sp. 3. p. 2251. Ligulæ roseæ aut albæ, 3-dentatæ, 3 lin. longæ.* (v. s. et v. c.)
3. C. TENELLUS (H. B. et Kunth nov. gen. am. 4. p. 240), foliis bipinnatipartitis glabris, lobis lanceolatis acutis, inferioribus basi subserratis, terminalibus serratis, invol. squamis exter. lineari-spathulatis brevioribus, achæniis subscabris... aristis.... (I) in frigidis Mexici inter urbem et Huehuetoca. *Ligulæ roseæ?*
4. C. CAUDATUS (H. B. et Kunth nov. gen. am. 4. p. 240), caule glabro, foliis bipinnatipartitis, lobis lanceolatis cuspidatis margine scabris, petiolis ciliatis invol. squamis exter. lineari-lanceolatis acutis ciliatis patulis vix basi ciliatis interiores ferè æquantibus, achæniis glabrisculis apice scabris longissimè rostratis, aristis 2 patulis. (I) in insulâ Cubâ juxta Havanam (H. et B.), Porto-hæc (Berl. Wydl!), Sancti-Thomasii agris saccharo cultis (Less. in *linneæ. p. 727*), Martinicâ (herb. Duby!), etiâ in Indiâ or. (ex h. Wight et Bont. Mauriti) (Bont.), sed introduct. *Bidens Berteriana Spr.! syst. 3. p. 454. D. Wight contrib. p. 19. Ligulæ roseæ apice trifidæ. Achænia ferè pollicem longæ. Flores disci 12 ex Kunth, circ. 20 in sp. meis.* (v. s. sp.)
5. C. SULFUREUS (Cav. ic. 1. p. 56. t. 79), caule piloso, foliis bipinnatipartitis lobis lanceolatis mucronatis margine subscabridis, petiolis ciliatis, invol. squamis ext. linearibus adpressis quàm interiores adhuc brevioribus et multo gustioribus, achæniis glabriusculis apice scabris longè rostratis, aristis 2 patulissimis. (I) in Mexico (Cav.) circa Oaxacam (Andr.! pl. ex. n. 305. Willalpando (Mendez!), *Cosmea sulfurea Willd. sp. 3. p. 2250. Cosmea attenuatifolia Jacq. ic. 3. t. 595. Ligulæ sulfuræ. Cor. disci subnudi superne pubescentes. Achæniis 9-10 lin. longum.* (v. s. sp.)

6. *C. CHRYSANTHEMOIDES* (H. B. et Kunth nov. gen. am. 4. p. 239. t. 382), glaber, foliis pinnato-quinque-partitis, lobis obovato-cuneatis bi- aut trilobulatis, petiolis ciliatis, invol. squamis exter. lanceolato-linearibus paulò brevioribus, achæniis glabris biaristatis.  $\mathcal{L}$  in Mexico. *Cosmea chrysanthemifolia* Spreng. syst. 3. p. 615. Ligulæ violacæ 8 lin. longæ. Achænium, ex ovario, videtur compressum subcuneatum, potius Bidentis quàm Cosmi.
7. *C. PILOSUS* (H. B. et Kunth nov. gen. am. 4. p. 241), caule folisque hispidopilosus pinnatipartitis, lobis bijugis oblongo-ovatis serratis, terminali paulò majore, invol. squamis exter. linearibus obtusis brevioribus, achæniis glabris biaristatis.  $\mathcal{L}$  in aridis Mexici prope Santa-Rosa de la Sierra. Ligulæ rosæ ellipticæ 3-dentatæ 5-6 lin. longæ. Achænium junius obcompressum ut in Bidente.
8. *C. CRITHEMIFOLIUS* (H. B. et Kunth nov. gen. am. 4. p. 242), glaber, foliis pinnatipartitis, lobis angustè linearibus integerrimis rigidis, infer. rariùs bifidis, invol. squamis exter. lanceolatis acutis brevioribus, achæniis... (I) ? prope Valladolid Mexicanorum (H. et B.). Ligulæ rosæ? — Huc fortè pertinet n. 1667 Berl. pl. exs. circa Toluccam lectus. Flores in specim. desiderantur. Achænia tetragona rostrata apice subscabra 6 lin. longæ, aristis 2-3-erectis superata. (v. s.)
9. *C. SCABIOSOIDES* (H. B. et Kunth nov. gen. am. 4. p. 242), caule pubescente, foliis pinnatipartitis ambus hispidolis, lobis bijugis lanceolato-oblongis acutis basi cuneatis apice subserratis, terminali majore, invol. squamis exter. lanceolatis acutis brevioribus, achæniis glabris 3-aristatis. (I) ? in Mexici montanis Mechuacansibus prope Peruaro (H. et B.) et ad Tlapujhua (Keerl. in h. Mart.). *Cosmea scabiosa* Spreng. syst. 3. p. 615. Ligulæ violaceo-purpureæ latæ obtusæ semipollicares. (v. s. in h. Mart.)

CCCLI. ADENOLEPIS Less. in *linnæa* 6. 1831. p. 510. t. 6. f. 6-10. syn. 230.

Capitulum pluriflorum heterogamum, fl. radii circ. 4 ligulatis neutris ligulâ subellipticâ emarginatâ, disci tubulosis hermaphr. 5-dentatis, limbo basi inflato dentibus revolutis. Recept. planum bracteolatum. Invol. cylindraceum disco æquale, squamis biserialibus linearibus, inter. foliaceis glandulâ magnâ terminatis. Anth. exsertæ ecaudatæ. Achænia conformia subtriangulato-obcompressa exalata, breviter rostrata seu apice attenuata omninò calva, in disco sæpè linearia sterilia. — Herba annua Sandwicensis gracilis parva apice puberula. Folia opposita petiolata ternatisecta aut superiora sessilia triloba. Capitula parva lutea terminalia solitaria.

1. *A. FULCHELLA* (Less. ll. cc.). (I) in O-Wahu Sandwicensium detexit cl. de Chamisso.

Div. v. VERBESINÆ Less. syn. 230. — Helianthæ Spilanthæ et Verbesinæ pleræque *Cass. opusc.* 3. p. 58.

Capitula heterogama, fl. radii ligulatis fœmineis, rariùs homogama discoidea. Achænia cortice tenni donata, ferè semper compressa aut obcompressa, rariùs 3-sulcata seu trigona, nunc calva, nunc sæpiùs in aristas rigidas tot quot achæniis anguli producta, rarissimè aut pappo vaginæformi aut setis fragilibus rigidis superata,

CCCLII. LASIANTHÆA. — Lasianthus Zucc. in *h. Acad. Monac. non DC. nec Jack.*

Capitulum multiflorum radiatum, fl. radii 1-serialibus ligulatis fœmineis, disci longè tubulosis hermaphr., lobis 5 supernè pube crassâ hirsutissimis. Invol. campanulatum, squamis pluriserialibus imbricatis

ovatis obtusis, exter. ciliatis. Recept. planum, paleis membranaceis complicatis. Styli fl. herm. ramis apice summo barbatis exappendiculatis. Antheræ appendice lineari-lanceolata superatæ. Achænia glabra, radi triquetra triala 3-aristata, disci compressa bialata et triaristata, aris radii et disci basi alato-dilatatis. — Frutex Mexicanus di- seu trichotomus. Rami teretes pubero-asperi. Folia opposita breviter petiolata ovata sublancoolata, vix subdentata, supra sparsè strigilloso-pubes, subtus subvelutino-villosa reticulato-nervosa. Capitula circ. 5 ad apic ramorum breviter pedicellata umbellata crassa. Flores lutei. — Nomen paululum mutavi ne cum Lasiantho (sect. Gordonis) confundatur.

1. *L. HELIANTHOIDES* (Zucc. l. c.). ♂ in temperatioribus intra Saltepos et Cals Mexicanorum legit septembr. flor. cl. de Karvinski. (v. s. in h. Acad. reg. Monac.)

CCCLIII. *PERYMENIUM* Schrad. ind. sem. hort. Gœtt. 1830. et in Ex. dur. et io. — *Schistocarpa* Less. in Linnæa 1831. p. 387. syn. 387.

Capitulum multiflorum radiatum, ligulis 1-serialibus femininis, 1 disci hermaphr. 5-dentatis. Invol. ovati aut campanulati squamis imbricatis ovatis obtusis aut subacutis. Recept. planiusculum, paleis membranaceis complicatis. Stylus disci ramis apice breviter appendiculatis seu cono terminatis. Achænia subconformia subcompressa aut marginalia obcompressa mutica aut apice submarginata exaristata. Pappus setis rigidulis fragilibus 8-10 uniserialibus inæqualibus constans. — Suffrutescens frutescens Mexicani setulis adpressis scabridi. Folia opposita plus minus petiolata seu triplinervia ovata aut oblonga densa. Capitula pedicellata solitaria aut corymbosa. Flores croceo-flavi. — Genus *Verbesinæ* et *Lipochætæ* affine.

1. *P. DISCOLOR* (Schrad. l. hort. Gœtt. ind. sem. 1830 et in litt. ic. et descr.) fruticosum, ramis cinereo-puberis, foliis oppositis subsessilibus oblongis vix acutis triplinerviis integerrimis margine subrevolutis supra et subtus in nervis adpressè villosis subtus inter nervos canescenti-velutinis, corymbis compositis trichotomis polycephalis, invol. ovati squamis ovalibus obtusis ciliatis, ligulis ovalibus disco paulò longioribus, ♂ in Mexico (Schrad.), in ditione Oaxacana legit cl. Andrieux (pl. exs. n. 3051). *Schistocarpa bicolor* Less. in Linnæa 1831. p. 409. Folia ferè Rosmarini. Flores croceo-flavi. Ligule apice appendice lineari auctæ subbilabiata. (v. s.)
2. *P. VERBESINOIDES*, fruticosum, ramis striatis scabridis, foliis oppositis petiolatis ovatis basi vix acutis apice acuminatis serratis triplinerviis supra acutis subtus molliter pubescenti-velutinis, capitulis 7-9 pedicellatis subcorymbosis, invol. ovati squamis ovatis acuminatis dorso villosis, ligulis 9-10 oblongo-linearibus, ♂ in Mexico. Valde simile *Verbesinæ* serratae sed pappo et involucre certè distinctum (v. s. ex hort. Mexic. comm. à cl. Berlandier sub n. 565.)
3. *P. BERLANDIERII*, fruticosum, ramis striatis vix subasperis, foliis oppositis brevè petiolatis ovato-lanceolatis triplinerviis remotè et parvè serratis supra sparsè subtusque in nervis minutissimè strigillois, capitulis 7-9 apice in axillis supremis corymbosis breviter pedicellatis, invol. ovati squamis ovatis subobtusis extus tenuè pubescentibus, ligulis 8-10 oblongo-linearibus, ♂ in Mexici montibus Guichilapensis legit cl. Berlandier (pl. exs. n. 1040). Medium inter *P. verbesinoides* et *P. Cervantesii*, sed ab utroque distinctum. (v. s.)
4. *P. MENDEZII*, fruticosum, ramis teretibus, junioribus pedicellisque adpressè pubescentibus, foliis oppositis brevè petiolatis ovato-lanceolatis triplinerviis subintegerrimis utrinquè puberulo-scabris, capitulis pedicellatis ad apic ramorum subumbellatis, invol. squamis ovatis obtusis infer. extus pubescentibus.

- tibus, ligulis 3-4 oblongis.  $\zeta$  in Mexico circa Villalpando legit cl. Mendez. Medium inter P. Berlandieri et Cervantesii. Pedicelli vix semipollicem longi. Styli rami breves cono crasso hirsutissimo terminati. Achænia matura regulariter punctis rufis maculata. Pappus 8-10-setus 1-2 subaristatiformibus. (v. s.)
5. P. CERVANTESII, suffruticosum, ramis adpressè piloso-scabris, foliis oppositis ovatis acutis basi obtusiusculis brevissimè petiolatis hinc inde serratis triplicinerviis suprâ piloso-scabris subtus pubescenti-hirtellis, capitalis 3-5 pedicellatis corymbosis, invol. campanulati squamis ovatis paucis dorso substrigillosis obtusis, intimis membranaceis oblongis, ligulis 5-10 ellipticis.  $\zeta$  in Mexico (Alam.), circa urbem in montanis (Berl. pl. exs. n. 634). Verbesina Mexicana Cerv. ined. ex Alam. Variat pedicellis 3 ferè bipollicaribus, aut 5 dimidiò et ultrâ brevioribus. (v. s.)
6. P. BUPHTHALMOIDES, suffruticulosum, ramis adpressè puberulis, foliis oppositis subsessilibus ellipticis lanceolatis acuminatis basi 3-5-nerviis grossè serratis suprâ sparè subtus præcipuè in nervis puberis, pedunculis terminalibus elongatis subsolitariis, invol. campanulati squamis oblongo-lanceolatis subacutis dorso adpressè puberis, ligulis 7-10 oblongis.  $\zeta$  in Mexico legit cl. Alaman. Habitu refert Verbesinam buphthalmoidem Link (abb. t. 49), sed faciliè differt. Pappus brevior quàm priorum. (v. s.)
7. P. BARCLAYANUM, suffruticosum subscandens, ramis adpressè villosis, foliis petiolatis ovatis acutis subdentatis trinerviis suprâ sparè subtus secus nervos strigilloso-pilosis, pedicellis solitariis elongatis 1-cephalis, invol. campanulati squamis ext. latè ovatis obtusis strigillosis, ligulis 5-7 oblongis.  $\zeta$  in Mexico ex hort. Barclay, ubi pro genere novo à cl. Don descriptum anno 1830 habebatur. Ligulæ croceo-flavæ 6-8 lin. longæ. Pappus 12-13-setus longior quàm priorum. (v. s. à Filio coll.)
8. P. GYMNOLOMOIDES, fruticosum scandens? ramis.... foliis...., invol. campanulati squamis ext. obtusissimis extus hirsutis, int. elliptico-obovatis, ligulis 5 disco paulò longioribus, tubo hispidiusculo, ovarii subulatis ciliatis.  $\zeta$  in dumetis Misanthe Mexicanorum. Lipotriche gymnomoides Less. in Linzæ 1831. p. 408. Simillimum dicitur Gymnolome microcephalæ.

CCCLIV. PSATHUROCHÆTA.

Capitulum multiflorum radiatum, ligulis 1-serialibus fœmineis, fl. disci hermaphrod. tubnlosis 5-dentatis. Invol. 1-2-seriale, squamis lanceolatis disco brevioribus. Recept. convexum, paleis lanceolato-linearibus acuminatis. Cor. lobi puberuli. Stylus disci ramis cono elongato acuminato hispido terminatis. Achænia acuta tetragona subcompressa brevia, setis disci plurimis 8-10 fragilibus inæqualibus deciduis coronata, setis radii paucioribus brevioribus. — Herba Australi-Africana, Verbesinæ facie. Caulis teretiusculus sex-sulcatus, oppositè ramosus et ideò trichotomus, pube brevi scabrido-canescens. Folia opposita petiolata utrinquè scabra trinervia sæpius triloba, lobo medio longiore et magis acuminato. Pedunculi ad apices ramorum alterni subterni 1-cephali nudi. Cor. intensè flavæ. — Genus affine Perymenio et Lipochætæ; à priore involucro 1-2-seriali, achæniis tetragonis, styli ramis longius acuminatis; à posteriore angulis fructus in aristas non productis; ab utroque recept. convexo diversum. Nomen ex  $\psi\alpha\theta\upsilon\rho\varsigma$  fragilis, et  $\chi\alpha\iota\tau\alpha$  seta.

- x. P. DREGEI.  $\zeta$ ? in Africâ australi ad Omsamconbo, ferè ad maris altitudinem, detexit cl. Drege. An varietates, an fortè species duæ distinguendæ? nempè:

a. *latifolia*, foliis submembranaceis basi latioribus trilobis, lobis dentatis

lateralibus sæpè sinuato-bilobis, nervulis subtus amplè areolatis vix prominulis. (v. s.)

β. *reticulata*, foliis rigidioribus, lobis serratis, lateralibus brevibus indivisis, nervulis subtus crebrè reticulatis elevatis. (v. s.)

CCCLV. LIPOCHÆTA. — *Lipotriche* Less. *syn.* 231. et in *linnæa* 1831. p. 3. non Brown. — *Texmenia* Lallav. et *Lex. nov. veg.* 1. p. 13? — *Verbesina* Gayd.

Capitulum multiflorum radiatum, fl. radii 1-serialibus feminis: disci hermaphr. 5-dentatis. Invol. ovati aut campanulati squamis 2-serialibus ovalibus adpressæ. Recept. planiusculum, paleis membranaceis complicatis. Stylus disci ramis apice appendiculatis. Achænia triquetra vix subalata, angulis sæpius ternis in aristam persistentem productis, dentibus setiformibus paucis interjectis; disci compressi 1-2-aristata ad latus interius subalata. — Suffrutices aut herbæ. Foliis opposita breviter petiolata aut sessilia ovato-lanceolata subserrata tripplinervia. Capitula pedicellata solitaria aut corymbosa. Flores lutei. — Habitus *Verbesinarum* oppositifoliarum.

\* *Species Americana.*

1. *L. MONOCEPHALA*, caule fruticoso tereti glabro, foliis petiolatis ovato-lanceolatis serratis tripplinerviis utrinquè scabris, petiolis intus villosis, pedicellis ex axillis ramorum solitariis 1-cephalis, invol. squamis biariatis parvis ovalibus submembranaceis, ligulis 4-5, achæniis cuneatis alatis, pappo equaliter pluriaristato. β patr. ign. olim in horto Paris. culta sub nom. *Verbesinae* fruticosæ et inde fortè ex ins. Caribæis orta. Flores lutei. (v. s.)

2. *L. STRIGOSA*, suffruticosa, ramis teretibus glabrisculis, foliis petiolatis ovatis acuminatissimis serrulatis suprâ rigide et sparse strigosius subtus adpressè pubescenti-villosis, pedicellis terminalibus 1-cephalis, invol. squamis biariatis, ext. foliaceis strigosius, achæniis elongatis angulatis 2-3-aristatis et paleolas totidem inter aristas gerentibus. β in Mexico prope Oaxacâ australi circa Tehuantepec legit cl. Andrieux (pl. exs. n. 313!). Fol. petiol. super. pili basi tuberculosi. Flores non vidi et ex fructu tantum hæc n. (v. s.)

3. *L. UMBELLATA*, fruticulosa, ramis adpressè scabridis, foliis brevissime petiolatis ovato-lanceolatis utrinquè subacutis minutè serrulatis suprâ brevè sparse strigillosis subtus secus ramulos hirtellis, nervis utrinquè 3 prope in medio ortis, petiolis subciliatis, capitulis plurimis umbellatis, pedicellis triplo longioribus, invol. ovatis squamis ext. obtusissimis, int. acutis et dimidiò ferè brevioribus. β in Mexico prope Cuernavaca legit cl. Berlandier (pl. exs. n. 1065), et ad Acapulco (h. Hænk!). Achænia radii 3-gona imbricè 3-aristata, disci compressa longè biaristata. (v. s.)

β. *conferta*, capitulis numerosioribus confertis, pedicellis capitulo brevioribus, invol. disco paulò breviorè. β ad Cuernavaca Mexicanorum legit cl. Berlandier pl. exs. n. 1053! (v. s.)

4. *L. FASCICULATA*, caule fruticoso glabro, ramulis vix apice puberulis, petiolatis oblongo-lanceolatis utrinquè acuminatis subserratis suprâ subtus puberulis, pedicellis axillaribus terminalibusque, infer. solitariis fasciculatis capitulo diacoideo sexuplo longioribus, invol. squamis ovalibus minimis, ext. pubescentibus, int. longioribus glabris. β in Mexico legit cl. Berlandier (pl. exs. n. 2134!). Frutex ergalis. Pedicelli inter axillares sæpius 2-3-serialiter dispositi! Ligula nulla. Achænia ext. triquetra, int. compressa biaristata, squamellis intermediis omnium minimis subaffinis *L. umbellatæ*. (v. s.)



" *Species Sandwicensis.*

5. *L. LOBATA*, fruticulosa, ramis pubero-scabris, foliis breviter petiolatis ovato-lanceolatis grossè dentatis sæpè tri- aut rariùs 5-lobatis 3-5-plinerviis suprà aspero-scabris subtùs secus nervos strigillosis inter eos adpressè puberis, capitulis 3-5 pedicellatis ad apices ramorum subumbellatis, invol. ext. squamis latè ovatis, ligulis 5-10 disco duplò longioribus. ♂ in insul. Sandwich (Gaud. Hook.), nempè Woahoo sen O-Wahu (Macr. Cham.). *Verbesina lobata* Gaud. voy. Freyc. bot. p. 464. Hook. et Arn. bot. Beech. 87. *Lipotriche australis* Less. in linnæa 183. p. 310 (v. s. coll. à cl. Macrae comm. ab hon. hort. soc. Lond.)
6. *L. NASTULATA* (Hook. et Arn. in Beech. bot. 87. sub *Verbesinâ*), suffruticosa, foliis oppositis brevè petiolatis ovato-lanceolatis 3-nerviis verans basin utrinquè sursum lobulatis grossè serratis suprà scabris subtùs hispidis. ♂ in insulis Sandwich. Affinis *L. connatæ* dicitur, sed ex phrasi vix à *L. lobatâ* differe videtur. †.
7. *L. CONNATA* (Gaudich. in bot. Freyc. voy. 464. sub *Verbesinâ*), suffruticosa, foliis sessilibus connatis rhombeo-ovatis argutè et grossè duplicato-serratis suprà scabris subtùs densè hispidis. ♂ in insulis Sandwich. Cæt. ign. (v. s. in h. Mus. Par. sed non car. carp. vidi.)
8. *L. LAVARUM* (Gaudich. in bot. Freyc. voy. 464. sub *Verbesinâ*), fruticulosa, foliis lanceolatis argutè serratis utrinquè strigoso-sericeis argenteis. ♂ in montanis insularum Sandwich, ubi dicitur *Kokolao* (v. s. in h. Mus. Par. sed car. carp. non vidi.)
9. *L. SUCCULENTA* (Hook. et Arn. in Beech. bot. 87. sub *Verbesinâ*), herbacea glabra nitida succulenta, foliis oblongo-ovatis apice obtusis mucronulatis basi in petiolum attenuatis suprà medium crenulato-serratis, pedunculis solitariis, invol. squamis orbiculatis. — in maritimis Sandwichianis inter rupes volcanicas ins. Ooneheow ubi dicitur *Neke* et *Neneke*. Achænia compressa sen trigona, angulo uno acutissimo interruptè subalato, aristis 2-3 brevibus coronata.

† *Species vix nota.*

10. *L. ? SERRATA*, caule sarmentoso 12-pedali et ultrà, ramis oppositis teretibus pubescentibus, foliis oppositis sublanceolatis petiolatis serratis asperis, junioribus subtùs pubescentibus, inflorescentiâ terminali, squamis inv. 10-12 biserialibus ovatis apice revolutis, paleis apice dilatatis. — in San-José del Corral Mexici, ad vias. *Zexmenia serrata* Lallav. et obs. nov. veg. 1. p. 13.

CCCLVI. DIPLOTHRIX.

**Capitulum** multiflorum radiatum, ligulis circ. 5 ovatis fœmineis cum ovario continuis persistentibus, fl. disci tubulosos 5-dent. hermaphr. **Invol.** ovatum imbricatum, squamis ovatis. Recept. paleis complicatis **flores** involv. onustum. Styli rami cono superati. Achænia radii obcompressa (seu triquetra) calva seu denticulis minimis superata, disci compressa, exter. subcalva, centralia in aristas 2 producta. — Suffrutices **Mexicani** ramosissimi. Folia opposita conferta acerosa acuta integerrima vix subpuberula 1-nervia, marginibus subcallosis. Capitula ad apices ramorum solitaria breviter pedicellata. Ligulæ ex sicco albidæ. — **Nomen** (olim ab Oppiano Pino consecratum) alludit et ad habitum foliorum ad Coniferas accedentem et ad aristas binas florum centralium.

11. *D. ACEROSA*, pedicellis involucri et folio vix longioribus, achæniis radii obcompressis. ♂ in Mexici ditione Sancti-Ludovici de Potosi decemb. flor. legit cl. Berlandier (pl. exs. n. 1343!). Folia tenuia 3 lin. longa. Capitula vix 3 lin. (v. s.)

2. D. JUNIPERIFOLIA, pedicellis capitulo et folio longioribus, achæniis radii triquetris. 3 cum priore legit cl. Berlandier (pl. exs. n. 1359!), sed ab eâ differt. foliis 6 lin. longis, ad margines callosio-nervosis, pedicello 6-12 lin. longo, capitulo duplò majore. In meo specim. nimis ætate provecto caract. florum non ritè vidi. (v. a.)

CCCLVII. SELLOA H. B. et Kunth nov. gen. am. (1820) 4. p. 265. t. 395. Cass. dict. 55. p. 273. Less. syn. 231 non Spreng. — Fœna Spreng. syst. 3. (1826) p. 581 non Bory (1824.)

Capitulum multiflorum radiatum, ligulis 1-serialibus 10-15 fœmineis et filam. antheris destituta gerentibus, fl. disci hermaphr. tubulosis 5-dentatis. Invol. 2-3-seriale, squamis circ. 10-15, ext. ovato-ellipticis, int. oblongis acutis. Recept. convexum, paleis lineari-setiformibus. Achænia cuneata pentagona lævia coronata setulis 3 (5?) hispidulis caducis tubo corollæ adpressis. — Herbæ Mexicanæ subscapiformes. caule nempe basi folioso apice longè denudato 1- rariùs 3-cephalo. Folia opposita petiolata ovato-elliptica subintegerrima triplinervia glabra. Capitulum ferè Senecionis elegantis. Ligulæ albæ extus purpurascens 9-nerviæ apice trilobæ.

1. S. PLANTAGINEA (H. B. et Kunth nov. gen. 4. p. 265. t. 395), caule apice nudo. foliis ovali-oblongis subdentatis triplinerviis. 2/ in montibus Mexicanis inter Aguasarco et Ario. Fœna plantaginea Spreng. syst. 3. p. 581.  
2? S. LINEARIS, caule etiàm apice foliato, foliis spatulato-linearibus cæcis integerrimis. 2/ in Monte-Video. Fœna linearis Spreng. syst. 3. p. 581. Spec. ex Less. syn. p. 231 omninò obscura. †.

#### † Species exclusæ.

- S. CAPENSIS Spreng. = Denekia Capensis.  
S. GLUTINOSA Spreng. = Gymnosperma glutinosum.  
S. NUDATA Nutt. = Gymnosperma nudatum.

CCCLVIII. VERBESINA Less. syn. 231. — Verbesinæ et Siegesbeckia sp. Lx: — Verbesina et Platypteris Kunth. — Verbesina, Platypteris et Hamulus Cass. — Verbesina et Locheria Neck.

Capitulum multiflorum, sæpiùs radiatum ligulis fœmineis, rariùs homogamum discoideum. Invol. bi- aut pluriseriale, squamis subæqualibus aut imbricatis. Recept. planum aut subconvexum, paleis minus complicatis et flores amplectentibus. Stylus disci ramis appropinquatis. Achænia erostris plano-compressa ad angulos alarum in aristas binas rigidas producta. — Herbæ aut suffrutices. Spec. genuinæ omnes Americanæ. Folia opposita aut alterna, petiolata sessilia aut decurrentia dentata aut pinnatilobata. Capitula pedicell. solitaria aut corymbosa. Flores nunc omnes lutei, rariùs omnes albi. interdum in disco flavi in radio albi. — Genus idè multiflorum sed a caract. ægrè in plurima divellendum.

Sect. 1. VERBESINARIA. — Verbesina Cass. dict. 59. p. 142.

Capitula radiata, ligulis uniseriatis. Achæniis aristæ rectæ sæpiùs æquales, unâ interdum breviorè aut deficiente.

§. 1. *Apteræ, nempe foliis non decurrentibus et ideo caule non alato.*\* *Oppositifoliæ.*

1. *V. SERRATA* (Cav. ic. 3. p. 7. t. 214), caule herbaceo jun. tomentoso demum subscabro, foliis oppositis breviter petiolatis ovato-lanceolatis grossè serratis suprà scabris subtus tomentosis, petiolis villosis subconnatis, capitulis compositè corymbosis, ligulis paucis aut (ex Kunth) nullis. 2/ in Mexico (Cav.), in temperatis inter Temascatio et Guanajuato (H. et B.), in insul. Cubà (h. Lem.). Cass. dict. 59. p. 142. Folia summa subalterna interdum integerrima. Invol. squamæ biseriales oblongæ foliaceæ disco breviores. Ligulæ breves sæpè difformes. Achenium maturum glabrum nigricans alà albà cinctum. Variat foliis super. alternis et capitulis discoideis. (v. v. c. et s. sp.)
2. *V. CANTHOPIFOLIA* (Willd. sp. 3. p. 2225), caule erecto, foliis oppositis brevè petiolatis ovatis serratis quintuplinerviis utrinquè pilis brevibus obsitis subtus scabris, pedunculis elongatis racemoso-paniculatis. — in Mexico prope Acapulco. Cœt. ign. †.
3. *V. ATRIPICIFOLIA* (Juss. et Desf. hort. Par. Poir. dict. 8. p. 460), caule fruticoso tereti apice villosulo-velutino, foliis oppositis petiolatis latè ovatis triplinerviis acuminatis dentatis indivisis aut trilobatis suprà scabridis subtus petiolisque molliter villosulo-velutinis, capitulis subternis pedicellatis, invol. squamis lanceolatis acutis disco brevioribus, ligulis circ. 12 apice dentatis. 5 patr. ign. Spreng. 3. p. 577. L'Hér. descr. ined. Colla hort. rip. p. 144. t. 31. Galinsoga discolor Spreng. nov. prov. 19. Clerodendron phlomisoides hortul. ital. Ligulæ fortè steriles nunc stylo donatæ nunc destitutæ, albæ. Discus flavus. Achenia ex statu immatura subvillosa biaristata? Hinc fomè species duæ confusæ, sed certè ambæ à *V. Phaetusa* diversissimæ. (v. s.)
4. *V. BUPHTHALMOIDES* (Link et Otto abb. p. 105. t. 49), caule decumbente suffrutescente supernè pubescente, foliis suboppositis lanceolatis acutis serrulatis in petiolom attenuatis pubescentibus, pedunculis elongatis solitariis 1-cephalis, invol. 2-3-serialis squamis lanceolatis subæqualibus, ligulis 10-12 tridentatis, acheniis pubescentibus exalatis biaristatis. 5 2/ in Monte-Video. Cor. flavæ.
5. *V. AUREA*, fruticosa, ramis teretibus adpressè canescenti-villosis, foliis oppositis sessilibus ovato-lanceolatis basi obtusis apice subacuminatis serratis utrinquè pubero-scabris, capitulis ad axillas super. solitariis et terminalibus, ideo ultimis ternis, invol. squamis ext. ovalibus obtusis dorso pubero-scabris, ligulis 7-9 involucri duplò longioribus, acheniis radii obcompressis biaristatis, disci triquetris inæqualiter aristatis. 5 in Mexico. Wedelia aurea Hook. Bot. mag. t. 3384. Flor. radii feminei sed fortè tamen steriles. (v. s. c. in hort. Barcl.)

\*\* *Alternifoliæ.*

6. *V. HELIANTHOIDES* (H. B. et Kunth nov. gen. am. 4. p. 204 non Michx.), caule herbaceo sulcato hirsuto-piloso, foliis alternis subsessilibus lanceolato-oblongis acuminatis remotè denticulatis basi cuneate-angustatis suprà scabris subtus pubescenti-pilosis, corymbis ramosis, invol. semiglobosi squamis imbricatis oblongis obtusis subscariosis, ligulis circ. 15, paleis oblongo-lanceolatis, acheniis glabris angustissimè alatis 2-3-aristatis. 2/ in regno Quitensi? Cor. flavæ? Verbesigia Humboldtii Spreng. syst. 3. p. 577.
7. *V. SORDESCENS*, rhizomate lignoso, caulibus suffruticosis teretibus pubero-scabris apice subhirsutis, foliis alternis sessilibus oblongo-lanceolatis basi obtusis apice acutis serratis utrinquè puberulis, capitulis ramulos terminantibus in corymbum dispositis, invol. squamis oblongis obtusis disco brevioribus, paleis vix acutis, acheniis glabris angustissimè alatis aut subexalatis longe biaristatis. 2/ 5 in Brasiliæ prov. Rio-Grande verisim. in campis. Caulis circ. pedalis obscurè viridis. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 876.)

8. *V. ACUMINATA*, caule subherbaceo glabro striato, foliis alternis subpetiolatis oblongo-ellipticis utrinque longè acuminatis integerrimis suprâ glabris subtus pallidioribus subpuberulis, corymbo composito ramosissimo polycephalo, capitulis oblongis pedicellatis, invol. squamis paucis obtusis, ligulis 1-2, achæniis puberis angustissimè alatis.  $\frac{1}{2}$   $\frac{3}{4}$  circa Caracas legit cl. Vargas. Folia 6 poll. longa, 15 lin. lata, in sicco marginibus subrevoluta. Ligula alba aut pallida. (v. s.)
9. *V. ELEGANS* (H. B. et Kunth nov. gen. am. 4. p. 204), caule herbaceo ramoso hirtio-villoso, foliis alternis subsessilibus oblongis acutis argutè denticulatis hirtio-pilosis corymbis subsimplicibus, invol. squamis paleisque lanceolatis acutis, ligulis circ. 15, achæniis apice ciliato-hispidis.  $\frac{1}{2}$  ? in regno Quitensi? Color flor. dubius.
10. *V. DENTATA* (H. B. et Kunth nov. gen. am. 4. p. 205), caule herbaceo ramoso pubescenti-canescente, foliis alternis subsessilibus approximatis oblongis grosse dentatis basi cuneatis subcoriaceis suprâ hirtio-pilosis subtus cano-lanatis, corymbis subsimplicibus, invol. semiglobosi imbricati squamis ext. obtusis, inter. acutis, ligulis circ. 20, paleis acutis, achæniis glabris alatis ciliolisque subbiaristatis, aristâ alterâ corollam æquante, alterâ brevissimâ.  $\frac{1}{2}$  in cæcis regni Quitensis. Cor. flavæ. *Pallasia dentata* H. et Bonpl. pl. equin. 2. p. 11. t. 111.
11. *V. SALICIFOLIA* (H. B. et Kunth nov. gen. am. 4. p. 205), caule herbaceo glabro, ramis virgatis striatis apice pedunculisque pubescentibus, foliis alternis subsessilibus lineari-sublanceolatis utrinque angustatis, corymbo ramoso ramis foliaceis superato, pedicello centrali brevi crasso, invol. squamis lineari-oblongis, ligulis 7-8, paleis vix apice subuncinatis.  $\frac{1}{2}$  in Mexico (Maire), cum Xalapam (H. B.). Valde accedit ad *V. virgatam*. Flores pallidius flavi. Folia 3 poll. longa, 4-5 lin. lata. (v. s.)
12. *V. PERSICIFOLIA*, suffruticosa glabra, foliis elongato-lineari-lanceolatis ètante calloso-serratis utrinque acuminatis breviter petiolatis penninerviis corymbo terminali ramoso polycephalo, invol. squamis lanceolatis, latis paucis? parvis? recept. paleis lineari-cuneatis, achænio glabro latè alato cordato ad latus interius 1-aristato, ad exterius mutico.  $\frac{3}{4}$   $\frac{1}{2}$  in Mexico (Santander et Victoria legit cl. Berlandier (pl. exs. n. 2209). Capitula nunc ferè globosa. Affinis *V. virgatæ*, sed folia nec auriculata nec decurrentia. Folia 5 poll. longa, 10-18 lin. lata. (v. s.)
13. *V. TOMENTOSA*, suffruticosa, ramis teretibus alternis apice cano-tomentosis foliis alternis petiolatis ovato-lanceolatis sublobatis inæqualiter dentato-serratis suprâ araneoso-pubescentibus subtus petiolisque cano-tomentosis, corymbo composito confertè polycephalo ramoso, invol. squamis lineari-oblongis, ligulis 5-8, achæniis glabris latè alatis emarginatis biaristatis.  $\frac{3}{4}$  in America? legit cl. Née, sed locus propr. ign. Cor. disci cylindricæ basi tomentosæ. fructifer.)
14. *V. OAXACANA*, caule fruticoso apice summo pedicellis subpubescentibus foliis alternis in petiolum attenuatis ellipticis utrinque acuminatis serratis penninerviis suprâ glabris subtus molliter cano-subtomentosis, corymbo simplici 5-8-cephalo, capitulis subsphæricis, invol. squamis biserialibus foliaceis subobtusis, ligulis plurimis brevibus obtusis, paleis submembranaceis acutiusculis, achæniis glabris.  $\frac{3}{4}$  in Mexici ditione Oaxacanâ prope Tepec tepec angust. flor. legit cl. Andrieux (pl. exs. n. 301) et fortè ad Acapulco. h. Hænke! Flores lutei. (v. s.)
15. *V. SUBCORDATA*, fruticosa, ramis pube brevi confertâ velutinis, foliis alternis sessilibus oblongo-lanceolatis utrinque attenuatis basi obtusè subcordatis èt dentatis subrepandis suprâ scabris subtus velutinis, pedicellis velutino-tomentosis 1-cephalis subcorymbosis, invol. squarrosis, ligulis 10-12, achæniis pubescentibus angustatis vix marginatis biaristatis.  $\frac{3}{4}$  circa Buenos-Ayres legit Baile. Flores flavi. (v. s. comm. à cl. inv.)

§. 2. *Pterocaulæ*; *nempè foliis in alas foliaceas secus caulem decurrentibus.*

\* *Foliis pinnatilobatis alternis.*

16. *V. GIGANTEA* (Jacq. ic. rar. 1. t. 175. coll. 1. p. 53), caule fruticoso simplici glabro tereti, foliis alternis infer. decurrentibus, sup. sessilibus auritis amplexicaulibus basi cuneatis suprà pinnatipartitis lobis utrinquè 3-4 lanceolatis subintegerrimis suprà pubescentibus subtùs molliter villosis canescentibus, corymbo composito, invol. squamis lanceolatis, ligulis paucis aut nullis, achenio glabro.  $\S$  in insulis Caribæis (Jacq.), fruticetis montosis Jamaicae (Swartz fl. Ind. occ. 3. p. 1368). Capitula ovato-oblonga circ. 12-flora. Cor. albæ. (v. s.)
17. *V. PINNATA* (Clark. ex herb. et mss. L'Hér.), caule suffruticoso tereti molliter villosa alato, foliis alternis confertis breviter decurrentibus pinnatipartitis basi cuneatis, junioribus subvillosis, adultis utrinquè pubescentibus, lobis utrinquè 4-5 lanceolatis serratis.  $\S$  ? patr. ign. ex L'Hér. mss. Corymbus compositus villosus confertus. Flores albi, ligulis 2-3 sæpè nullis. Achenia jun. subvillosa. Invol. subimbricatum. Confusa cum *V. gigantea*, sed differt foliis etiã summis evidenter decurrentibus multò minùs villosis, caule villosa, etc. (v. s. sine flor. ex h. L'Hér.)
18. *V. TURBACENSIS* (H. B. et Kunth nov. gen. am. 4. p. 203), caule suffruticoso ramosissimo apice cano-tomentoso, foliis (superioribus) alternis decurrentibus sinuato-laciniosis basi valdè angustatis suprà piloso-hirtis subtùs tenuiter canescenti-tomentosis, corymbis ramosissimis, invol. campanulatis squamis spatulato-lanceolatis, ligulis 6, acheniis glabris.  $\S$  ad Turbaco Novo-Granatenisum. Herba procera affinis *V. giganteæ*. Cor. flavæ.
19. *V. SINUATA* (Ell. sketch 2. p. 411), caule herbaceo striato pubescente infernè alato, foliis alternis, sup. sessilibus inf. decurrentibus, sinuatis pinnatifidisve basi attenuatis apice acutis suprà scabris subtùs pubescentibus, corymbo composito, invol. squamis subimbricatis, ligulis 3-5, acheniis subalatis.  $\mathcal{L}$  in Carolinà merid. (Nutt.), in arenosis insularum maritimarum Edding et Hilton-Head (Ell.). *V. laciniosa* Nutt. gen. am. 2. p. 170 excl. syn. Lam. Flores albi. — Fortè ex Ell. hinc referenda *Siegesbeckia laciniosa* Poir. dict. 7. p. 158 à cl. Boec in Carolinà lecta ?
20. *V. DIVERSIFOLIA*, caule herbaceo glabresculo, foliis alternis decurrentibus ovatis acuminatis basi longè cuneatis suprà sparsè puberulis subtùs molliter pubescenti-velutinis, aliis grossè hinc indè incisis, aliis trilobis lobis lanceolatis, summis sessilibus indivisis, corymbo composito polycephalo, invol. oblongi squamis lineari-lanceolatis, ligulis paucis, acheniis glabris.  $\mathcal{L}$  circa Bahiam Brazilæ leg. cl. Blanchet et Salzmann. Affinis *V. pinnatifida* sed folia alterna, *V. giganteæ* sed suprema etiã decurrentia. Flores albi. Caulis 5-6-pedalis. Var. habeo foliis indivisis. (v. s.)

\*\* *Foliis pinnatilobatis oppositis.*

21. *V. PINNATIFIDA* (Cav. ic. 1. p. 67. t. 100 non Swartz), caule herbaceo tomentoso alis 4 aucto, foliis oppositis longè decurrentibus pinnatifidis basi longè cuneatis utrinquè præsertim in nervis puberulo-hirtis, lobis utrinquè 2-4 lanceolatis hinc indè subincisis, paniculâ oppositè ramosâ apice corymbosâ, invol. squamis lineari-lanceolatis acutis, ligulis circ. 12 oblongis, acheniis glabris alatis.  $\mathcal{L}$  in Mexico (Cav.). Flores pallidè flavi. Invol. nigricantia. Achenia radii interdum 3-alata ex Cav. (v. s.)

$\beta$ . *undulata*, alis caulinis valdè undulato-crispatis, caule minùs tomentoso basi suffruticoso, foliorum lobis regulariter dentatis. — Jacq. h. Schœnbr. 3. t. 305.

\*\*\* *Foliis indivisis oppositis.*

22. *V. PHANTUSA* (Cass. dict. 51. p. 476 et 59. p. 149), caule herbaceo alis 4 foliaceis integerrimis aucto, foliis oppositis decurrentibus ovato et ferè rhombeo-lan-

ceolatis triplinerviis glabrisculis serratis, paniculâ trichotomâ polycephalâ apice corymbosâ, invol. squamis paucis obtusis, ligulis 1-5 homochromis, achæniis villosis nunc exaristatis nunc 1-2-aristatis pilis nonnullis intermediis. 2/ in Amer. bor. sed locus propr. ign. Frequenter in hortis bot. Eur. culta ut abortu subditioca evadit. *Phaetusa Americana* Gærtn. fr. 2. p. 465. t. 169. Lm. syn. 216 excl. syn. Poir. *Siegesbeckia occidentalis* Linn. spec. 1269. *Phaetusa borealis* Spreng. syst. 3. p. 591. — Pluk. mant. t. 342 fig. penult. Certè specificè differt à *V. Siegesbeckiâ* Michx. Foliis grossè triplinerviis et à *Verbesina* caract. paulò recedit ob achænia inter aristas seriem pilorum brevium præsentia, sed non admittere queo sententiam cl. Lessingii, qui *Phaetuse* genus in tribu *Verbesinæ* omninò alienâ collocavit. (v. s. c.)

23. *V. Siegesbeckia* (Michx. fl. bor. am. 2. p. 134), caule herbaceo glabro aliâ foliaceis integerrimis aucto, foliis oppositis decurrentibus ovali-lanceolatis utrinquè acuminatis penninerviis glabrisculis grossè serratis, paniculâ trichotomâ polycephalâ apice corymbosâ, invol. squamis paucis obtusis, ligulis 1-5 homochromis, achæniis exalatis subpuberis, aristis rigidis, pilis nullis intermediis. 2/ in umbrosis Amer. bor. sylvia à Virginiâ ad Carolinam (Pursh, Bor.) et in Georgiâ (h. Duby!). Willd. sp. 3. p. 2224. *Verbesina occidentalis* Walt. fl. car. p. 213. *Siegesbeckia* Gron. virg. 128 sed *Siegesbeckia occidentalis* Lænei qui expressis verbis dixit folia triplinervia ad *V. Phaetusam* potius referenda. (v. s. sp. nec unquam in hort. Eur. cult. vidi.)

\*\*\* *Foliis indivisis alternis.*

24. *V. microptera*, caule suffruticoso apice pubescenti-velutino, foliis alternis decurrentibus ovatis basi longè petioliformibus cuneato-angustatis obtusè obtusè sinuato-dentatis suprâ scabridis subtus vix pubescentibus, ramulis subnudis, capitulis 3-10 paniculatis, invol. squamis linearibus, ligulis 6-8 achæniis cuneato-linearibus vix alatis. 2/ 5 in Mexico inter Laredo et Bq. febr. florentem legit cl. Berlandier (pl. exs. n. 1442). Flores ex sicco vid. care. Capitula subrotunda. Ligulæ obovatæ disco paulò longiores. (v. s.)

25. *V. virgatâ* (Cav. ic. 3. p. 38. t. 275), caule suffruticoso glabro, ramulis pedicellisque pubescentibus, foliis alternis lineari-lanceolatis utrinquè attenuatis serratis brevissimè petiolulatis, aliis basi nudis, aliis auriculâ subrotunda subtus què auctis, aliis breviter decurrentibus, corymbo terminali, pedicellis corymbis brevissimis, invol. squamis lanceolatis acutis, ligulis 10-12, paleis lanceolatis, jun. apice rigidè uncinatis. 5 in Mexico undè habui lectam à cl. Alaman, Mairet. Flores lutei. Achænia pubescenti-hispida margine alata alata biaristata. (v. s.)

2/ *conyzoides*, foliis suboppositis basi auriculatis. *V. conyzoides* fl. icon. ined. (v. ic.)

26. *V. polycephala*, caule herbaceo? brevissimè pubescenti-velutino, foliis alternis decurrentibus ellipticis acuminatis basi longè attenuatis integerrimis suprâ sparsè scabridis subtus molliter pubescenti-velutinis subincanis, corymbo composito polycephalo, invol. oblongi squamis lineari-lanceolatis paleis ligulis 3-4, achæniis striatis hispidulis latè alatis. 2/ 5 in Mexico prope Bejar angusto flor. legit cl. Berlandier (pl. exs. n. 1813). Caulis 4-6-ped. Flores albi. Pedunculi albido-tomentosi. (v. s.)

27. *V. pterocaulis* (fl. mex. icon. ined.), caule herbaceo glabrisculo, foliis alternis longè decurrentibus lineari-lanceolatis utrinquè acuminatis acutè serratis glabris, corymbo 10-12-cephalo, invol. squamis linearibus obtusis, paleis 10-12, achæniis angustè alatis et ciliatis. 2/ in Mexico. Cor. 2/ Habitùs *V. virgatæ* sed folia omnia decurrentia. (v. ic.)

28. *V. virginica* (Linn. sp. 1270), caule herbaceo pubescente inferne serrato apice aptero, foliis alternis penninerviis lanceolatis seu obovato-lanceolatis serratis glabrisculis, infimis decurrentibus, corymbo composito, invol. squamis subobtusis pubescentibus ligulis 3-4 heterochromis, achæniis angustè alatis. 2/ in sylvis umbrosis à Pennsylvania ad Carolinam (Pursh, Fl.

in Kentucky orient. (Raf.), Georgia (Ell. sketch 2. p. 410). *Anomantia* Raf. I in litt. 1831. *V. paniculata* Poir. I dict. 8. p. 456. Ligulæ albæ ovales. Achæniæ aristæ a pilis nullis intermediis. (v. s. sp.)

29. *V. MOZZIS* (H. B. et Kunth nov. gen. am. 4. p. 203), caule herbaceo simpliciter densè hirsuto, foliis alternis decurrentibus lanceolatis obtusè serratis suprâ densè piloso-hirtis subtus albo-lanatis, capitulis ternis subcorymbois, invol. semiglobosis squamis ligulisque lineari-lanceolatis, achænio glabro margine ciliato, aristis valdè inæqualibus. (1) in Mexico inter Guanajuato et Villalpando. Flores flavi. *V. alata* ex radice annuâ et aristis inæqualibus affinis.

Sect. II. *HAMULIUM*. — *Hamulium* Cass. dict. 20. p. 260.

Capitulum radiatum, ligulis numerosis subbiseriatis. Achæniæ aristæ 2 inæquales, minore rectâ majore apice uncinatâ. — Herba annua, foliis decurrentibus.

30. *V. ALATA* (Linn. sp. 1270), caule herbaceo glabresculo alato, foliis alternis longè decurrentibus oblongis ovatisve obtusè undulatis et sinuato-dentatis glabresculis, capitulis subglobosis solitariis terminalibus, invol. squamis oblongis, achæniis hispiduli alati ciliolati aristis inæqualibus, majore apice uncinatâ. (2) in Curaco (Volk. nor. t. 106), Surinamo (Comm. h. amst. 1. t. 3), Jamaica (Sloan. Br.), in Mexico (fl. mex. ined. ic.), ins. Cubâ (H. B.). Swartz obs. p. 313. Sims bot. mag. t. 1716. *Hamulium alatum* Cass. dict. 20. p. 261. — Magn. hort. t. 40. Cor. crocæ, ligulæ numerosæ subbiseriales breves. (v. v. c. et s. sp.)

Sect. III. *PLATYPTERIS*. — *Platypteris* H. B. et Kunth nov. gen. am. 4. p. 200. Cass. dict. 50. p. 259.

Capitula discoides, ligulis planè nullis.

§. 1. *Involucro imbricato, foliis decurrentibus oppositis, caule herbaceo.*

31. *V. CROCATA* (Less. syn. 232), herbacea, ramis tetrapteris apice hispido-scabris, foliis oppositis decurrentibus irregulariter pinnatilobatis, lobis ovalibus erosè-dentatis, terminali subdeltoideo, capitulis pedunculatis solitariis, invol. squamis imbricatis paleisque lineari-subulatis, ligulis nullis, achæniis glabris latè alatis biaristatis. (2) in Mexico propè Tusco (H. et B.) et alibi. *Bidens crocata* Cav. ic. 1. p. 66. t. 99. *Spilanthus crocata* Sims bot. mag. t. 1627. *Platypteris crocata* H. B. et Kunth nov. gen. am. 4. p. 201. Cass. dict. 50. p. 259. Capitula globosa crocæ. (v. s.)

§. 2. *Invol. circ. bi- seu triseriali non sensim imbricato, foliis alternis non decurrentibus, caule fruticoso.*

32. *V. AURICULATA*, caule suffruticoso tereti brevissimè et confertè velutino, foliis alternis obtusè auriculatis sessilibus ellipticis acuminatis basi longè attenuatis penninerviis suprâ scabris subtus molliter et sordidè tomentosis subdentatis, capitulis plurimis breviter pedicellatis ad apices ramorum racemoso-corymbois, invol. squamis ovali-oblongis subacutis 2-3-serialibus, rec. paleis elongatis rigidis apice subulatis, ligulis nullis. (5) in Mexici ditione Oaxacanâ ad Tehuantepec august. flor. legit cl. Andrieux (pl. exs. n. 302). Folia 7 poll. longa, 2 poll. lata. (v. s. florib. non satis evol.)
33. *V. ARBOREA* (H. B. et Kunth nov. gen. am. 4. p. 202), arborea, ramis teretibus tenuiter fuscescenti-villosis, foliis alternis petiolatis oblongis acuminatis integerrimis suprâ hirtis subtus lanato-tomentosis et fuscescenti-panicatis, corymbis ramosissimis foliatis, invol. imbricati campanulati squamis obtusis oblongo-linearibus, ligulis nullis, flosculis 10-12, paleis obtusis. (5) in frigidis montosis agri Quitensis præes. ad montem Pichincha.

† *Verbesinæ non satis notæ.*

\* *Americanae.*

34. *V. æSTUANS* (Lam. dict. 3. p. 262), caule simplici, foliis oppositis petiolatis subspathulatis acutis penninerviis dentatis subtus tomentosis, capitulis pedunculatis, invol. ovato adpressè imbricato, ligulis linearibus numerosissimis. — in Amer. calid. Plum. ed. Burm. t. 41. f. 2. Asteris sp. Tourn. Helenii sp. Vall. Inula æstuaus Linn. sp. 1236. Propius ad Verbesinam quàm ad alia generis citata accedere videtur, sed car. flor. et fruct. vix noti. †.
35. *V. FOLIACEA* (Spreng. syst. 3. p. 578), herbacea, foliis oppositis petiolatis ovatis crenatis pubescentibus, pedunculis adgregatis axillaribus abbreviatis hirsutis, invol. foliaceo. — in Guadalupâ (Bert.), in Brasiliâ (Sell.). An plantæ duæ diversæ? An generis?
36. *V. ILICIFOLIA* (Poir. dict. 8. p. 460), caule..., ramis teretibus glabris, foliis oppositis petiolatis coriaceis elliptico-lanceolatis utrinquè acutis grossè indentatis triplinerviis suprâ glabris subtus petiolisque sparsè et minutè puberulis, capitulis pedicellatis subsolitariis, invol. squamis lanceolatis.... — in Sancto-Domingo. *V. fruticosa* Linn. sp. 1271? non Willd. *V. frutescens* Plum. ed. Burm. t. 52? Cor. lateæ. Cæ. car. ign. An Flourensiae species? (v. s. folia.)
37. *V. MEGAPOTAMICA* (Spreng. syst. 3. p. 578), caule herbaceo simplici oligocephalo, foliis oppositis breviter petiolatis lanceolatis integerrimis subtrigilosis, pedunculis axillaribus elongatis, radio angusto. — in Brasiliæ provinciâ Rio-Grande.
38. *V. MONTEVIDENSIS* (Spreng. syst. 3. p. 578), caule herbaceo simplici raphalo hispido, foliis oppositis subfasciculatis lanceolatis integerrimis hispidissimis, capitulo terminali solitario. — in Monte-Video. Affinis ex phras. vid. *V. buphthalmoides* sed descr. incompl.
39. *V. TRIDENTATA* (Spreng. syst. 3. p. 577), fruticosa, foliis oppositis cuneatis in petiolum decurrentibus 3-dentatis subserratis scabris triplinerviis, pedunculis 1-cephalis, invol. simpliusculo. ♀ in Amer. merid. (Willd.). in Martiniâ (Spr.). *V. fruticosa* Willd. sp. 3. p. 577 excl. syn. ex Spreng. †.

\*\* *Gerontogææ.*

40. *V. CHINENSIS* (Linn. sp. 1270), subtomentosa fruticosa, caule tereti, foliis alternis petiolatis ovato-lanceolatis serratis, capitulis terminalibus solitariis pedunculatis, invol. hemisphærico, achænio 4-gono setis 4 lævibus cœmatis. ♀ in Chinâ. Fl. lutei.
41. *V. SPICATA* (Lour. fl. coch. 2. p. 620), herbacea ramosa, foliis alternis longipetiolatis serratis subtus tomentosis, inf. palmato-incisis, sup. lanceolatis, capitulis spiciformibus congestis terminalibus, invol. squamis subulatis linearibus, ligulis 5 unilateralibus invol. æqualibus, achænio calvo. — Culinæ in hortis Cochinchinæ et Chinæ. Cum acetariis mensis apponitur. Flores albi radiati. A Verbesinâ videtur omnino aliena! sed locum aptiorem ignoro.

†† *Species exclusæ.*

- V. ACMELLA* Linn. = *Spilanthes Acmeella*.  
*V. ALBA* Linn. = *Eclipta erecta*.  
*V. ASTEROIDES* Linn. = *Amellus Lychnitis*.  
*V. BIFLORA* Linn. = *Wollastonia biflora*.  
*V. BIFLORA* Blume. = *Wollastonia scabriuscula*.  
*V. BIFLORA* var. Wall. = *Wollastonia urticæfolia* E.  
*V. BOSWALLIA* Linn. f. = *Glossocardia Boswallia*.  
*V. CALENDULACEA* Linn. = *Wedelia calendulacea*.



- V. CANESCENS Gaudich. = *Wollastonia canescens*.  
 V. CONNATA Gaudich. = *Lipochæta connata*.  
 V. CORROPSIS Michx. = *Actinomeris alternifolia*.  
 V. DICHOTOMA Murr. = *Blainvillea rhomboides*.  
 V. DICHOTOMA Wall. = *Blainvillea latifolia*.  
 V. HASTULATA Hook. = *Lipochæta hastulata*.  
 V. HELIANTHOIDES Michx. = *Actinomeris helianthoides*.  
 V. LANCEOLATA Poir. = *Blainvillea rhomboides* β.  
 V. LAVARUM Gaudich. = *Lipochæta lavarum*.  
 V. LAVENIA Linn. = *Adenostemma Lavenia*.  
 V. LOBATA Gaudich. = *Lipochæta lobata*.  
 V. MEMBRANIFOLIA Poir. = *Wulffia membranifolia*.  
 V. MOLUCCANA Blume. = *Wollastonia Moluccana*.  
 V. MONTANA Blume. = *Wollastonia montana*.  
 V. MUTICA Linn. = *Chrysanthellum procumbens*.  
 V. MUTICA Sw. = *Chrysanthellum Swartzii*.  
 V. NODIFLORA Linn. = *Synedrella nodiflora*.  
 V. OPPOSITIFLORA Poir. = *Wulffia membranifolia*.  
 V. OVATA hort. Par. = *Heliopsis bupththalmoides*.  
 V. PROSTRATA Linn. = *Eclipta prostrata*.  
 V. PSEUDO-ACMELLA Linn. = *Spilanthes Pseudo-Acmella*.  
 V. PUSILLA Poir. = *Eclipta pusilla*.  
 V. REPANDA Pers. = *Lepidophorum repandum*.  
 V. SATIVA Roxb. = *Guizotia oleifera*.  
 V. SCABERRIMA Wall. = *Wedelia urticæfolia* γ.  
 V. SUCCULENTA Hook. = *Lipochæta succulenta*.  
 V. STRIGULOSA Gaudich. = *Wollastonia aristata*.  
 V. URTICÆFOLIA Blume. = *Wedelia urticæfolia* γ.

CCCLIX. DITHRICHUM *Cass. bull. philom. 1817. febr. dict. 13. p. 272. — Spilanthes sp. Less.*

Capitulum pluriflorum homogamum, fl. hermaphr. tubulosis, cor. extus pilosis. Invol. floribus longius cylindraceum biseriale, squamis ext. brevissimis laxis, int. longis erectis coriaceis acuminatis. Recept. planum, paleis in appendicem subulatam productis. Achænia compressa obovata glabra. Pappus biaristatus, aristis ex angulis ortis crassis subbarbellulatis. — Herba elata. Caulis teres pubescens. Folia alterna sessilia subdecurrentia oblongo-lanceolata irregulariter sinuato-lobata scabra ferè pedalia. Capitula paniculato-corymbosa. Flores flavi.

1. D. MACROPHYLLUM (Cass. ll. cc.). — Patr. ign. verisim. hab. in America; confusa enim latebat in herb. Juss. sub nomine *Conyzæ lobatæ*.

## CCCLX. MICRACTIS.

Capitulum multiflorum, fl. radii minutissimè ligulatis fœmineis, disci tubulosis 4-dentatis hermaphroditis, centralibus fortè sterilibus. Invol. 1-2-seriale, squamis oblongis subconcavis dorso subpuberulis. Recept. palæ etiam complicato-concavæ sed glabræ, ut squamæ inv. achænia semi-involventes. Stylus radii bifidus, ligulâ integrâ concavâ

paulò longior, disci inclusus. Anth. 4 ecaudatæ. Achænia tetragono-compressa ferè obovata glabra omninò calva. — Herba Madagascariensis annua erecta minutissimè puberula. Folia opposita oblonga acuta vix subcrenata scabra, super. integra. Capitula ad apices ramorum terna congesta breviter stipitata. Achænia nigra inter paleas persistentes seu tardè deciduas semioccultata. Flores minutissimi flavi. — Ambiguit inter capitula monoica (ex Bojer) et heterogama, sed potius ad hæc meo sensu referenda. An potius ad Millericas seu Silphicas rejicienda?

1. M. BOJER. (1) in ins. Madagascar legit cl. Bojer. Habitus plebeius sed structura singularis præsertim ob ligularum minutiam, et florum disci stylum exiguum fortè sæpius abortivum. (v. s. comm. à cl. Bojer.)

CCCLXI. SPILANTHES Jacq. Amer. 212. Schreb. gen. 1266. — Spilanthes Linn. mant. 475. Gærtn. fr. 2. t. 167. Less. syn. 232. — Pyrethrum Medic. act. pal. 3. p. 237 non Willd. — Ceruchis Gærtn. mss. ex Schreb. — Spilanthes et Acmella Rich. in Pers. ench. 2. p. 293 et 472. Cass. dict. 59. p. 137 et 138. 50. p. 257 et 258. p. 24. 328. — Spilanthes et Athronia Neck. elem. 1. p. 17 et 32.

Capitulum multiflorum, nunc heterogamum fl. radii ligulatis femininis sæpius parvis, nunc homogamum fl. omnibus hermaphrod. tubulosis 4-5-dentatis. Invol. biserialè adpressum disco brevius, squamis ext. subfoliaceis, inter. submembranaceis complicatis. Styli fl. hermarum apice truncati et penicillati. Anth. nigricantes. Achænia disci compressa erostria sæpius ad latera ciliata seu exaristulata, radii (ubi adest) triquetra seu obcompressa. — Herba sæpius annua interdum perennes aut suffruticulosa dichotoma aut vagè ramosa, Geronto et sæpius Neogæ. Folia opposita subintegra. Pedunculi terminales aut è dichotomiis orti erecti 1-cephali. Capitula ovata aut conica. Cor. flava rarius albæ, ligulis disco concoloribus, rarius discoloribus. — Herba pleræque gustu fervida et salivariæ.

Sect. 1. Acmella. — Acmella Rich. in Pers. ench. 2. p. 472. Cass. dict. 24. p. 531. — Athronia Neck. elem. 1. p. 32.

Capitula radiata, ligulis basi hispidis. Achænia radii sæpius trigona triaristata seu rarius calva, aut angulo interiore nudo biaristata, aut angulo inter. oblitterato ferè obcompressa.

1. S. ARNICOIDES, caule repente adscendente hirtello parè ramoso, foliis ad basin subconfertis oppositis in petiolum longè attenuato-cuneatis ellipticis ovato-ve sublanceolatis subdentatis glabriusculis, pedunculo longissimo 1-cephalo erecto sparsè puberulo, capitulo subgloboso, invol. squamis circ. 13 ovali-oblongis obtusis, ligulis 12-15 invol. triplò longioribus, achæniis radii abortivis, disci ciliatis biaristatis. 2. in Brasiliæ prov. Rio-Grande. Cor. intense luteæ ferè crocæ, in specimine majori pallidius flavæ, in altero (verisimè conspecifico) capitulum majus et magis depressum quàm in cæteris sp. (v. s. in h. Mus. reg. Par. à Mus imp. Bras. sub n. 1023 miss.)
2. S. DORONICOIDES, caule basi subrepente adscendente basi glabro inter folia confertè hispido, foliis lanceolatis obtusis basi longè cuneato-attenuatis subpetiolatis integerrimis hinc indè ad nervum et margines subcætosia, pedunculo elongato rigido 1-cephalo subangulato adpressè puberulo, capitulo ovato-obtusato, invol. squamis ovali-oblongis obtusis, ligulis triplò longioribus obtusis 3-dentatis, achæniis subciliatis, radii 3-, disci 2-angulatis aristaticque. 1. in Brasiliæ prov. Rio-Grande. Affinis S. arnicoidi. Cor. flavæ. Herba flores ra-

pedalis, pedunc. 4-poll. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1038.)

3. *S. LOWEFOZIA*, caule erecto simplici hispido ad medium foliato supernè in pedunculum longissimum 1-cephalum sub capitulo hirsutum producto, foliis oblongo-lanceolatis basi attenuatis integerrimis medio subtriplinerviis, infer. brevioribus glaberrimis, super. longioribus hispidis, invol. squamis linearibus circ. 15-16, ligulis invol. duplò longioribus.... 2½ in Brasiliæ prov. Rio-Grande. In specim. malè servato car. fruct. non vidi, sed ex analogiâ cum *Sp. arnicoide* de genere non dubito. Folia 3 poll. longa, 5 lin. lata. Ligulæ crocæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1029 miss.)
4. *S. BLEPHARICARPA*, caule basi repente adscendente mox erecto infra foliorum juga hirtè cæterùm glabriusculo, foliis lineari-lanceolatis obtusis basi longè cuneato-attenuatis integerrimis glabris, pedicellis elongatis striatis 1-cephalis, capitulis ovatis obtusissimis, invol. squamis ovalibus substriatis, ligulis invol. duplò longioribus, acheniis radii 3- disci 2-angulatis aristatisque. angulis densè et longè ciliatis. 2½ in Brasiliæ prov. Rio-Grande (h. Mus. imp. Bras. n. 1030!), et fortè eandem in prov. Sancti-Pauli legit cl. Lund. Herba pedalis ad *S. macropoda* accedens. Flores flavi. (v. s.)
5. *S. MACROPODA*, caule herbaceo striato subsimplici glabriusculo, foliis oppositis petiolatis oblongo-lanceolatis subacutis basi longè attenuatis integerrimis glabris margine scabris, pedicellis longissimis paucis 1-cephalis apice subpuberis, capitulis ovato-globosis, invol. squamis ovatis circ. 10 dorso trinerviis, ligulis circ. 10 elongatis 3-dentatis, acheniis glabris radii 3- disci 2-aristatis. — in Brasiliæ prov. Rio-Grande. Caulis circ. pedalis. Pedicelli 10-12 poll. longi. Cor. flavæ. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1049 miss.)
6. *S. LEPTOPHYLLA*, glaberrima, caule basi repente mox adscendente basi folioso parè ramoso, foliis linearibus integerrimis rigidulis, pedunculis nudis 1-cephalis caule longioribus, invol. squamis 10-12 oblongis, capitulis ovato-globosis, ligulis invol. triplò ferè longioribus, acheniis.... 2½ in Brasiliæ prov. Rio-Grande inter muscos. Car. fruct. in specim. caris subdestructo non vidi. Affinitas summa cum *S. macropoda*. Flores flavi. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1026.)
7. *S. STOLONIFERA*, caule à basi ramoso stolonifero, ramis floridis adscendentibus hinc indè hispidulis, foliis lanceolatis basi attenuatis subobtusis integerrimis triplinerviis glabris basi vix subsetoso-ciliatis, pedicellis terminalibus elongatis apice adpressè pubescentibus, capitulis globosis, invol. squamis elliptico-oblongis subacutiusculis, ligulis invol. duplò triplòve longioribus 3-dentatis, acheniis ciliatis aristulatis. 2½ in Brasiliæ prov. Rio-Grande. Habitus ferè *Hieracii Auriculæ*. Flores flavi. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1077.)
8. *S. SPHÆROCEPHALA*, glabra debilis, caule tereti gracili repente, foliis oppositis in petiolum attenuatis oblongis basi trinerviis vix subserratatis, capitulis ramos apice nudos terminantibus globosis aut vix ovatis, invol. squamis lanceolatis, ligulis 8-10 invol. duplò longioribus apice dentatis, acheniis ciliatis, radii triquetris 3-aristatis, disci compressis biaristatis. — in Brasiliæ prov. Rio-Grande. Flores flavi. Caules 8-12 poll. longi. Foliorum juga remota. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 1075 miss.)
9. *S. FIMBRIATA* (H. B. et Kunth nov. gen. am. 4. p. 208), caulibus erectis vel procumbentibus subimpticibus glabris, foliis oppositis petiolatis ovatis acutis basi subrotundatis dentatis glabris, pedicellis terminalibus demùm alaribus elongatis, capitulis breviter radiatis, invol. squamis circ. 7 oblongis adpresso-pilosis, acheniis radii triangularibus, disci oblongis, omnibus ciliatis 2-3-? aristulatis. (I) in Andibus Novo-Granatensium prope Ybague et Passo del Machin. *Acmella fimbriata* Cass. l. c. Flores latei.
10. *S. CILIATA* (H. B. et Kunth nov. gen. am. 4. p. 208), caule erecto? tetragono glabro, ramis junioribus hispidulis, foliis oppositis petiolatis ovatis acu-

minatis basi paulò angustatis grossè dentatis glabris petiolisque ciliatis, pedicellis terminalibus demùm alaribus folio longioribus, capitulis breviter radiatis, invol. squamis 6-7 ovato-oblongis acutis glabris, achæniis lævibus? ciliatis aristulatis. (1) in Amer. merid. prope Chipò et Santa-Fé de Bogota. *Acmella ciliata* Cass. l. c. Flores flavi. Affin. *Sp. Mutisii* et *fimbriatæ* dicitur.

11. *S. MUTISII* (H. B. et Kunth nov. gen. am. 4. p. 209), caulibus erectis vel procumbentibus subdichotomis glabriusculis, foliis oppositis petiolatis ovatis acutis basi subrotundatis grossè dentatis glabris, pedicellis subhispidulis terminalibus folio longioribus, capitulis subrotundo-ovatis radiatis, invol. squamis circ. 10 ovato-oblongis pubescentibus, ligulis 10 invol. longioribus acutè 3-dentatis, achæniis exaristatis, radii subverrucosis ciliolatis, disci lævibus glabris (1)? prope Santa-Fé de Bogota. *Anthemis Americana* Mutis in Linn. f. suppl. 378. *Anth. occidentalis* Willd. spec. 3. p. 2183. *Anth. oppositifolia* Lam. Dec. 1. p. 576. *Acmella occidentalis* Pers. ench. 2. p. 473. *Heliopsis*? dubia Dunal mem. Mus. Par. 5 p. 58? *Acmella Mutisii* Cass. dict. 24. p. 331. Ligulæ mixtilo piloso majores quàm in cat. spec., corollæque disci flavæ. (v. s.)
12. *S. GRAYII*, caule erecto ramoso hispidulo, foliis oppositis ellipticis in petiolum cuneato-attenuatis apice obtusis triplinerviis subintegerrimis glabris, pedunculis folio triplò longioribus 1-cephalis striatis rigidulis, capitulis se-globosis, invol. squamis ellipticis obtusis, ligulis invol. duplò triplòve longioribus apice dentatis, achæniis subciliatis aristatis. — in Brasiliæ prov. Rio-Grande. Herba sesquipedalis. Folia 3 poll. longa, 8-10 lin. lata. Flores flavi ligulis ex sicco subtus striato-purpureascentibus. (v. s. in h. Mus. reg. Par. et h. Mus. imp. Bras. sub. n. 1005.)
13. *S. BECCABUNGA*, caule basi repente glabro dein ascendenti-erecto subhispido, foliis oppositis petiolatis latè ovatis subacutis repandulo-subdentatis triplinerviis glabriusculis, pedicellis terminalibus glabriusculis folio 2-3 longioribus, capitulis rotundo-ovatis radiatis, invol. squamis circ. 12 ovato-oblongis, ligulis circ. 10 invol. longioribus sub-3-dentatis, achæniis glabris aristatis. 2l in Mexico in valle Mexici et Tolucae legit cl. Berlandier (pl. exs. n. 720 et 1160!). Flores lutei. Capitula ferè magnit. *S. oleracei*. Folia 3 poll. longa, pollicem lata, petiolo 3-4 lin. (v. s.)
14. *S. PÖPPIGII*, caule erecto subtetragono glabriusculo, foliis oppositis petiolatis ovatis subacutis dentatis trinerviis supra glabris subtus secus nervos strigilloso-pilosis, petiolis ciliatis, pedicellis terminalibus demùm alaribus folio longitudine, capitulis ovatis radiatis, invol. squamis ext. circ. 8 ovato-ciliatis, ligulis parvis, achæniis submarginatis ciliatis exaristatis aut aristatis minimam gerentibus. — in Peruvia legit cl. Pöppig. (pl. exs. n. 32. sub n. 1205!). Cor. disci 4-dentatæ. Ligulæ flavæ. (v. s.)
15. *S. LUNDII*, caule à nodis radicante demisso, ramis erectis subpuberulis foliis oppositis petiolatis ovato-lanceolatis dentatis glabriusculis, petiolis ciliatis, pedicellis terminalibus folio longioribus, capitulis ovato-conicis radiatis, invol. squamis ext. 7-9 ovato-lanceolatis ciliatis, ligulis totidem paulò longioribus, achæniis ciliatis tenuè aristulatis. 2l? ad rivulos circa Rio Janeiro legit cl. Lund. Ligulæ et cor. disci flavæ. (v. s.)
16. *S. SUBHIRSUTA*, caulibus repentibus ascendentibus vix ramosis petiolis pedicellisque hispidò-subhirsutis, foliis oppositis petiolatis ovatis aut summis lanceolatis subacutis acutè serratis sparsè hispidulis, pedicellis terminalibus folio longioribus, capitulis subrotundo-ovatis parè radiatis, invol. squamis circ. 10 lanceolatis, ligulis obovatis 3-dentatis invol. paulò longioribus, achæniis glabris aristatis. 2l in Mexico circa Tampico de Tamaulipas legit cl. Berlandier (pl. exs. n. 149!). Ligulæ corollæque disci flavæ. (v. s.)
17. *S. CESPITOSA*, caule à basi ramoso subradicante procumbente caespitosa glabro, foliis oppositis confertis petiolatis lanceolatis basi acutis apice subacutis vix subdentatis glabris, pedicellis terminalibus folio multo et caulis longioribus, capitulis subrotundis ovatis parè et minutè ligulatis, invol. squamis

mis 8-10 lanceolatis, achæniis ciliatis exaristatis. ♀ in Brasiliæ prov. Minarum-Generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 3081). Affinis S. Salzmanni, sed habitus diversissimus et achænia aut ad latas superius aut utrinquæ ciliata. Basis caulis ferè suffrutescit. (v. s.)

18. S. KARVINSKIANA, caule imâ basi suffruticuloso ramoso erecto diffuso glabro sub nodis hirsutulo, foliis oblongo-lanceolatis sessilibus acutis subintegerrimis glabriusculis setis longis hinc inde ciliatis, pedicellis axillaribus 1-cephalis folii longitudine et ex dichotomiis folio longioribus, invol. squamis oblongo-linearibus extûs hirtellis, ligulis parvis oblongis indivisis, achæniis glabris calvis angulatis. ♀ inter plantas Mexicanas Karvinskii sine loci indicatione hanc video. Ligulæ parvæ subtûs virescentes. Valdè affinis S. Mariannæ. (v. s. in h. acad. reg. Monac.)

19. S. MARIANNÆ, glabra herbacea erecta debilis, foliis oppositis in petiolatû attenuatis ovali-lanceolatis utrinquæ acuminatis triplinerviis ferè quintuplinerviis subtûs glabris suprà et margine junioribus subpuberis, pedicellis axillaribus 1-cephalis folio longioribus, invol. squamis oblongo-linearibus hirtellis, ligulis minimis linearibus indivisis bipartitisve, achænio glabro calve angulato. ① ? in Brasiliæ prov. Minarum-Generalium ad Mariannam legit cl. Vauthier (pl. exs. n. 322). Capitula parva. Flores flavi. Achænia nigra. (v. s.)

20. S. CAULIRHIZA, caule prostrato sub petiolis radicante, foliis breviter petiolatis ovali-rhombeis subcrenatis ramisque glabriusculis, pedunculis è dichotomiis aut apice ramorum ortis scabridis folio triplò duplòve longioribus, capitulis subglobosis, ligulis obovatis 3-dentatis involucrio longioribus, achæniis glabris calvis. — in Sennar Africæ borealis. *Acmella caulirhiza* Caill. et Delil. pl. afr. p. 45. t. 64. f. 7.

♀. *Madagascariensis*, glabriuscula foliis ovatis paulò longius petiolatis, capitulo paulò majore, recept. breviori. — in insulâ Madagascar legit cl. Rojer. Fortè species propria ? (v. s. comm. à cl. Rojer.)

21. S. ACMELLA (Linn. syst. veg. 610), caule basi subradicante erecto aut adscendente, foliis petiolatis ovato-lanceolatis subintegerrimis dentatisve glabriusculis, pedunculis folio triplò longioribus, capitulis ovatis, ligulis 5-6 minimis, achæniis ciliatis 1-2-aristatis. ① in Indiâ orient. *Verbesina Acmella* Linn. mant. 475. *Acmella* Linnæi Cass. dict. 24. p. 330. et fortè *Acm. brachyglossa* v. 50. p. 258 ex ipso in Desf. cat. ed. 3. p. 186. Omnino refert S. pseudacmellam et fortè hujus mera varietas. Synonyma omnia confusa videntur. Icones omnes ad alteram citatæ ob ligularum minutiam etiâ hic citari possent. (v. s.)

22. S. REPENS (Michx. fl. bor. am. 2 p. 131), caule repente ramoso sparso subhirtello, foliis oblongo-lanceolatis brevissimè petiolatis acutis subintegerrimis, pedunculis terminalibus demùm alaribus folio longioribus, capitulis ovatis radiatis, invol. squamis circ. 12 lanceolatis, ligulis totidem sub-3-dentatis invol. duplò longioribus, achæniis scabridis exaristatis ad angulos eciliatis. ♀ in Carolinæ humidis. *Anthemis repens* Walt. car. 211. Pursh fl. 2. p. 562. *Acmella repens* Pers. ench. 2. p. 473. Ell. sketch 2. p. 406. Discus radinsque flavus. Folia pollicem longa, 2-3 lin. lata. (v. v. c.)

23. S. SALZMANNI, caule imâ basi breviter adscendente radicante cæterum erecto glabriusculo, foliis oppositis petiolatis ovatis aut ovato-lanceolatis subdentatis, pedicellis terminalibus demùm alaribus folio longioribus, capitulis ovato-conicis parçè radiatis, invol. squamis 8-10 lanceolatis, ligulis invol. æquantibus, achæniis glaberrimis eciliatis exaristatis. ① ? in paludosis circa Bahiam Brasiliæ legit cl. Salzmann. Habitu accedit à S. uliginosam, sed major, cor. omnes flavæ et præ. achænia diversa. (v. s.)

24. S. AFRICANA, caule adscendente hinc inde setulis aspero, foliis oppositis petiolatis ovato-lanceolatis acutè et inæqualiter dentatis margine rigidè et brevè ciliatis, pedicellis folio duplò longioribus, capitulis ovato-conicis, ligulis involucrio paulò longioribus, invol. squamis oblongis vix acutis, achæniis glabris

- eciliatis exaristatis. — in Africâ austr.-orient. ad Zw. Omtata et Omsamwelo legit cl. Drege n. 5086! Valdè affinis S. Salzmanni, sed ciliis foliorum præcipuè diversa. Folia petiolo incluso var. 1  $\frac{1}{2}$ -2  $\frac{1}{2}$  poll. longa. (v. s.)
25. S. ULIGINOSA (Swartz prod. 110. flor. Ind. occ. 1291), caule erectissimo dichotomo glabro aut apice subpuberulo, foliis petiolatis ovato- aut oblongo-lanceolatis subdentatis glabriusculis, pedicellis terminalibus demùm alaribus folio longioribus, capitulis jnn. subrotundis demùm ovato-conicis, invol. squamis circ. 10-12 exter. ellipticis, ligulis paucis minimis obtusis, achæniis ciliatis breviter biaristatis. (1) in uliginosis argillosis Jamaicæ australis (Swartz). Porto-Ricci (Ledru!), Sanctæ-Crucis? (Vest.l.). *Acmella uliginosa* Cass. dict. 1. p. 531. Cor. disci 4-dentatæ flavæ, ligulæ albæ. (v. s.)
26. S. TENELLA (H. B. et Kunth nov. gen. am. 4. p. 211), caule adscendente dichotomo glabro, foliis oppositis petiolatis ovato-oblongis acutis basi pauli angustatis subintegerrimis glabris margine scabriusculis, pedicellis folio longioribus angulatis pilosiusculis, capitulis parvis breviter radiatis, invol. squamis circ. 5 oblongis acutis, achæniis ciliatis 2-aristulatis, radii triangulari-compressis, disci compressis. (2) circa Caracas. *Acmella tenella* Cass. l. c. Cor. flavidæ. An var. Sp. debilis?
27. S. DEBILIS (H. B. et Kunth nov. gen. am. 4. p. 210), caule adscendente dichotomo glabro, foliis oppositis petiolatis ovatis acutis basi rotundatis subintegerrimis glabris margine scabriusculis, pedicellis terminalibus et alaribus longissimis, capitulis radiatis, invol. squamis 5-6 elliptico-oblongis ciliatis, ligula obovatis brevissimis obtusè 3-dentatis, achæniis ciliatis biaristulatis. (1) ad ripam fluminis Orinoci Amer. merid. *Acmella debilis* Cass. l. c. Cor. flavæ. Affinis dicitur Sp. exasperatæ.
28. S. INTERMEDIA, caule procumbente glabro, foliis ovato-lanceolatis dentatis, pedunculis terminalibus. — in Americâ. *Acmella intermedia* Rich. in Pers. ench. 2. p. 472. Cor. truncus omnibus speciebus radiatis ferè conveniens.

Sect. II. SALIVARIA — *Spilantes* Cass. dict. 50. p. 257. — *Spilantes* Pers. ench. 2. p. 493.

Capitula discoidea. Achænia omnia conformia compressa.

29. S. RADICANS (Schrader! cat. h. Goett. 1831. p. 5. non Jacq.), caulibus ramis procumbentibus basi radicanibus, ramis adscendentibus magis minusve pilosis, foliis oppositis petiolatis ovatis basi cuneatis à medio crenato-serratis margine scabriusculis, pedicellis terminalibus folio longioribus, capitulis ovato-oblongis, invol. squamis 12-15 elliptico-oblongis subciliatis, corollis radii elongatis tubulosis. (2) patr. incert. Flores lutei. Capitula subradiata dixit cl. Schrader et reverà fl. marginales elongatos non verò ligulatos, sed verisim. steriles in specim. video. (v. s. comm. à cl. Schrader.)
30. S. OLERACEA (Jacq. hort. Vind. 2. t. 135), caule ramoso diffuso, foliis oppositis petiolatis latè ovatis basi obtusis truncatis aut subcordatis repando-dentatis, pedicellis terminalibus 1-cephalis folio longioribus, capitulis crenatis ovatis discoideis, invol. squamis 15-16 ovali-oblongis, achæniis ad angulum ciliatis biaristatis mucicæve. (1) in Indiâ orientali? (Willd.), sed potius in Americâ merid. (Lam. Desf. Plam. sp. 10). Vulgò *Cresson de Para* hort. Pyrethrum *Spilanthus* Medic. act. pal. 3. phys. p. 242 et 273. *Bidens acmelloides* Berg. act. holm. 1768. *Bidens fervida* Lam. dict. 1. p. 415. Herba densè virens, corollis etiã in medio disci flavis. Folia, imò herba tota, gustu ferrugine et intensè salivaria! (v. v. c.)
- β. ? *fusca*, corollis in medio disci fuscis, foliis obscure viridibus. (1) patr. fortè Brasilia ex nom. sp. Brasiliensis ab hortul. olim admissio. *Spilantes fusca* Hort. Par. *Bidens fusca* Lam. dict. 1. p. 416. An suad. cl. Lamarck et Desf. species distincta? Constans enim à dimidio sæculo in horti culta servatur, sed caract. excepto colore nulli. (v. v. c.)

31. *S. ALBA* (Willd. sp. 3. p. 1714), caule ramoso adscendente, foliis oppositis et alternis petiolatis ovatis subrepandis, pedicellis terminalibus folio longioribus, capitulis ovatis discoideis, invol. squamis 15-20 oblongo-linearibus, achæniis ad latera ciliatis biaristatis. (1) in Peruvia legit cl. Dombey! *Spilanthus albus* L'Hér.! stirp. p. 7. t. 4. *Spil. Salivaria* Morr. comm. Gœtt. 1783. p. 3. t. 1. *Bidens ocyimifolia* Lam. dict. 1. p. 416. ill. t. 668. f. 3. Corollæ albæ. Sapor et virtus *Sp. oleraceæ*. Variat in eisdem aut diversis individuis foliis longiusculis aut brevissimè petiolatis. (v. s.)
32. *S. ZEUCANTHA* (H. B. et Kunth nov. gen. am. 4. p. 210. t. 370), caule volubili? sexangulato glabro, foliis oppositis petiolatis ovato-oblongis acuminatis basi rotundatis dentatis rariùs integerrimis quintuplinerviis suprâ margineque scabriusculis, pedicellis terminalibus aut è dichotomiis ortis folio longioribus, capitulis ovatis discoideis, invol. squamis circ. 15 ovato-lanceolatis, achæniis glaberrimis ciliatis exaristatis. (1) in collibus siccis prope Alausi Quitensium. Cor. albidæ 5-dentatæ.
33. *S. URENS* (Jacq. Amer. 212. t. 126. f. 1), caule prostrato aut adscendente basi radicante, foliis oppositis lanceolatis integerrimis glabris basi attenuatis, pedunculis terminalibus folio longioribus glabriusculis, capitulis ovato-globosis discoideis, invol. squamis ovatis sublanceolatis ciliatis, achæniis ciliatis biaristatis. (2) in arenosis maritimis Carthagenæ (Jacq.), insulis Caribæis (Plum.), Sancto-Domingo (Bert.!), in Peruvia prope Truxillo (H. B. et Kunth nov. gen. am. 4. p. 211). *Bidens angustifolia* Lam. dict. 1. p. 416. et ill. t. 668. f. 2. (var. angustifolia). *Sp. repens* Spr. in h. Balb. *Spil. urens* et verisim. *S. Portoricensis* Spreng. syst. 3. p. 444. *Cotula Spilanthus* Linn. mant. 1. p. 116. *Spilanthus urens* et *Cotula Spilanthus* Willd. Cor. albidæ. Anthæræ nigricantes. (v. s.)  
*β. hispidula*, foliis utrinquè pedunculisque hirtellis. — circa Limam legit cl. D'Urville. (v. s.)
34. *S. MAURITIANA*, caule repente adscendente, foliis petiolatis ovali-lanceolatis subdentatis utrinquè ramisque subpuberulis, pedunculis folio duplò longioribus, capitulis ovato-globosis discoideis, achæniis glabris calvis. — in insula Mauritii (Belang.!), in graminosis montium excelsorum (Bout.!), speciatim ad montem Pollicis (Bojer!). Foliis petiolus 3 lin., limbus 6 lin. long., 3 lin. latus. Capitulum parvum. *Acmella Mauritiana* Rich. in Pers. ench. 2. p. 472. (v. s.)
35. *S. PSEUDO-ACMELLA* (Linn. syst. veg. 610), caule erecto subpubero, foliis petiolatis ovalibus lanceolatisve dentato-serratis asperiusculis acuminatis pedunculatis folio triplò longioribus, capitulis ovatis discoideis, achæniis lævibus ciliatis biaristatis. (3) in Indiâ orient. frequens. *Verbesina pseudo-Acmella* Linn. sp. ed. 2. p. 1270. mant. 475. *Pyrethrum Acmella* Medic. act. pal. phys. 3. p. 243. t. 19. *Sp. Acmella* Blum. bijdr. 912. Wight herb. 2. n. 30. — *Sebastes*. 1. p. 19. t. 10 et 11. Rumph. amb. 6. t. 65. Pluk. alm. t. 159. f. 4. (v. s. sp. et v. c.)
36. *S. CALVA* (DC. in Wight contrib. p. 19), caule adscendente basi repente spice hirsutulo, foliis petiolatis ovatis obtusiusculis serratis crenatisve glabriusculis basi ciliatis, pedunculis folio triplò longioribus, capitulis ovatis discoideis, achæniis glabris calvis. — in Indiâ orientalis agris Mysorensi et Travancorensi legerunt cl. Heyne et Wight (fl. ind. 2. n. 29!). *Sp. Acmella* Wall. cat. et herb. n. 295. F. H. An à *S. pseudo-Acmella* satis differt? (v. s.)
37. *S. PANICULATA* (Wall. cat. et herb. n. 296), caule erecto ramosissimo glabriusculo, foliis breviter petiolatis ovatis acutis dentatis, pedunculis innumeris folio triplò longioribus, capitulis ovatis discoideis, achæniis ciliatis biaristatis ad utrumque latus setoso-scabris. — in Indiâ orient. ad Martaban legit cl. Wallich. (v. s. comm. à cl. Car. merc. angl. Ind. or.)
38. *S. RUGOSA* (Blum. in litt. 1831), caule adscendente subpubero, foliis breviter petiolatis lanceolatis parçè et grossè dentatis rugosis suprâ sparsè setoso-scabris subtus secus nervos subscabris, pedunculis folio quadruplò longioribus, capitulis discoideis ovatis, achæniis glabris calvis. — in Javæ orientalis prov. Rou-

juerangie legit cl. Blume et mecum comm. Hirsutie et foliis rigidioribus obscurioribus grossius serratis à *S. pseudo-Acmella* differt. (v. s.)

39. *S. LEOGARTIA*, caulibus diffuso-procumbentibus, foliis oppositis sessilibus obovato-oblongis integris glabris tenuissimè triplinerviis, pedicellis terminalibus solitariis vix folio longioribus, capitulis globosis subdepressis, invol. squamis ovatis obtusis, achæniis glabris ciliatis inæqualiter biaristatis. *T. in Peruvia* (Dombey!), ad sinum Chorillo (Macrae!). Recept. paleæ obtusæ. (r. c. comm. à Mus. reg. Par. et Soc. hortie. Lond.)

40. *S. EXASPERATA* (Jacq. ic. rar. 3. t. 584), caule dichotomo ascendente basi radicante exasperato glabrinsculo, foliis oppositis petiolatis ovatis repandis pedicellis terminalibus folio longioribus, capitulis ovatis discoideis, invol. squamis ext. 5 lanceolatis puberis, inter. obovatis subconvolutis apice truncatis denticulatis, achæniis ciliatis biaristatis. *(1) in Venezuela*. Spil. radicem Jacq. coll. 3. p. 229 non Schrad. Cor. albæ sæpius 4-fidæ. Achæniis superius utraque ex icone glabra.

β. *Cayennensis*, petiolis sparsè pilosiusculis, foliis ovato-lanceolatis amplexatis, achæniis utrinque sparsè puberulis. *(1) in Cayennâ* legit cl. Pers. Pars inf. plantæ mihi ignota. (v. s.)

41. *S. ? INSIPIDA* (Jacq. amer. 216. t. 126. f. 2. folium), caule frutescente procumbente ramoso, foliis oppositis obovatis subdentatis sessilibus glaberrimis. *5 in Havannâ* rupestribus maritimis. Pedunculi et capitula omnino refert *S. exasperatam* ex Jacq. Cæc. ign. †.

42. *S. BRASILIENSIS* (Spreng. syst. 3. p. 444), caule erecto, foliis ovato basi attenuatis mucronato-dentatis glabris, pedunculis solitariis, capitulis discoideis. — *in Brasiliâ*. Car. mancus fere omnibus speciebus discoideis conveniens. †.

#### † *Species dubia.*

43. *S. NITIDA* (Lallav. et Lex. nov. veg. mex. 1. p. 28), caule volubili suffulculoso, foliis oppositis brevè petiolatis ovato-lanceolatis subintegerrimis glabris, corymbis terminalibus axillaribusve, invol. imbricati calyculati squamis ovatis ciliatis, capitulis discoideis, achæniis ciliatis et biaristatis. *5 in Mexico* dicit José del Corral ad vias in montosis. In specim. siccis folia juniora alba et cuprino nitore splendent. — An *Salmæ* potius species?

#### †† *Species exclusæ.*

*S. ARBOREA* Forst. = *Petrobium arboreum*.

*S. CROCATA* Sims. = *Verbesina crocata*.

*S. PSEUDO-GUMMIFERA* Forst. mss. = *Petrobium arboreum*.

*S. TETRANDBA* Roxb. = *Petrobium arboreum*.

*S. TINCTORIA* Lour. = *Adenostemma tinctorium*.

#### CCCLXII. DUNANTIA.

Capitulum multiflorum homogamum, fl. omnibus hermaphr. tubulosis 5-dentatis. Invol. squamæ 1-seriales adpressæ lanceolatae aut oblongæ. Recept. elongatum. Paleæ exteriores squamis invol. similium, interiores sensim tenuiores submembranaceæ. Styli rami exserti appendicem filiformem hispidam longè producti. Achænia tetragona gracilia erostris glabra calva. — Herbae Mexicanæ. Caulis erectus teres pubescens. Rami sæpius oppositi subdivaricati apice nudi. Folia opposita brevissimè petiolata lanceolata basi acuta triplinervia pubescentia integerrima. Capitula ovata ad apices ramorum solitaria bina aut ternata, lateraliter ubi adsunt subsessilia. Cor. ex sicco albidæ. — Genus affiné *Spilanthi* sed distinctissimum, involucri stigmatibus et achæniis



structurâ, præ cæteris analogiam inter squamas invol. et paleas recept. summam exhibet. — Genus dixi in honorem amic. Phil. Dunant itineris Berlandierani fautoris, qui herbarium ditissimum sollicitè curat et botanicis humaniter communicat!

1. *D. ACHYRANTHES*, caule pube basi confertâ velutinâ, foliis acuminatis, invol. squamis et recept. paleis acuminatis indurato-subspinosepimentibus. 2/7 in Mexico ad Tantoyuca legit cl. Berlandier (pl. exs. n. 2155!) et in Panamæide cl. Hænke! Caulis 1-2-pedalis, habitu fortè *Achyranthis porrigentis*. (v. a.)

2. *D. MICROCEPHALA*, caule vix pubescente, foliis oblongis subsæctis, invol. squamis et recept. paleis obtusis. 2/ in Mexico legit cl. Hænke. Folia pollicem longa, 3 lin. lata. Capitula ovata. (v. a. in h. Hænke à cl. de Sternberg comm.)

CCCLXIII. *XIMENESIA* Cav. ic. 2. p. 60. t. 178. — *Ximenesia* sp. Less. syn. 232. — *Ximenesia* sp. Kunth.

Capitulum multiflorum heterogamum, fl. radii foemineis ligulatis 1-serialibus, disci hermaphr. tubulosi 5-dentatis. Invol. circ. biseriale disco ferè longius, squamis patulis angustis acutis foliaceis subsqualibus. Recept. plano-subconicum, paleis amplectentibus membranaceis lanceolatis. Cor. omnium tubus hispidus. Styli disci ramis apice appendiculatis. Achænia radii plano-compressa alata apice profundè emarginata subpilosa biaristata, aristis cum lateribus alarum subconcretis. — Herbe annue pleræque Mexicanæ plûs minûs canescentes. Folia opposita aut alterna in petiolum sæpius subulatum et basi auriculatum attenuata, cordata ovata aut oblonga, dentata. Capitula laxè et irregulariter corymbosa. Corollæ luteæ. — Genus affinë *Verbesinæ*, semper enim vidi stylos in ligulis et embryonem in achæniis radii, sed ignoro an radius sit fertilis?

1. *X. KNECHTIOIDES* (Cav. ic. 2. p. 60. t. 178), achæniis disci villosiusculis undiquè alâ cinetis, apice emarginatis, radii apteris rugosis, ligulis profundè tridentatis. ① *Pallasia serratifolia* Smith in Rees cycl. v. 26. n. 3. Hæc quatuor stirpes fortè in posterum distinguendas adhuc servo, nempe:

a. *hortensis*, foliis infer. cordatis, super. latè ovatis, alâ achæniæ pellucidâ, lobis acutis. — Frequens in hortis botanicis sed nunquam specimen certè spontaneum vidi. (v. v. a.)

β. *pachyptera*, foliis infer. cordatis, super. latè ovatis, alâ achæniæ crassiusculâ opacâ, lobis acutiusculis. — in Mexico legit cl. Berlandier (pl. exs. n. 2068 et 2286!). (v. a. sp.)

γ. *oblongifolia*, foliis infer. ovatis, super. oblongis, alâ achæniæ subopacâ parè ciliatâ, lobis subobtusis. — in Senegaliâ ad ripas flum. regionis Walo legit Perrottet, in insulis Mauritanis Sieber (fl. maur. 2. n. 118!) et Belanger, et in hort. Calcuttensi Wallich. (v. a.)

δ? *cana*, foliis omnibus ovali-oblongis valdè cano-villosis, achæniis villosissimis alis opacis obtusis. — in Mexico circa Laredo legit cl. Berlandier (pl. exs. n. 2068 et 2074). Facile ut species propria admittenda videatur. (v. a.)

2. *X. MICROPTERA*, achæniis disci glabriusculis apice summo tantum parè alatis et brevissimè aristatis. — circa Buenos-Ayres legit cl. Bacle. Folia latè ovata subtus pubescentia suprâ scabriuscula. Petioli alii nudi, alii ad basin brevè et acutè auriculati. Ligulæ minutè tridentatæ. Achænium sæpè hinc tantum ad apicem subulatum. (v. a.)

† *Species exclusæ.*

*X. CAVANILLESII* Spr. = *Echinacea heterophylla*.

*X. CORDATA* H. B. K. = *Simsia cordata*.

*X. PORTIDA* Spr. = *Simsia sicifolia*.

X. HETEROPHYLLA H. B. K. = *Simsia heterophylla*.

X. HIRTA Mart. = *Simsia auriculata*.

X. HISPIDA H. B. K. = *Simsia hispida*.

CCCLXIV. SANVITALIA Gualt. in Lam. journ. h. nat. 2. (1792) p. 176. t. 31 illustr. (1797) t. 686. Cav. ic. 4. (1797) t. 351. Cass. dict. 47. p. 294. Less. gn. 232. — Lorentea Ort. dec. 4. (1797) p. 42. t. 5 non Less.

Capitulum multiflorum radiatum, ligulis foemineis persistentibus 1-serialibus, fl. disci hermaphr. tubulosis 5-dentatis. Invol. 2-3-seriale, squamis adpressis subimbricatis, inter. paulò longioribus disco aequalibus. Recept. conicum convexumve, paleis oblongis semiamplexentibus. Cor. disci super ovarium articulata pilifera. Styli rami cono superat. Achenia radii grossa triquetra lævia aristis 3 crassis lævibus conico divergentibus coronata, disci compressa biformia, exter. muricata fere calva, interiora alata ciliata aristellis 2 minimis superata. — Herba Americanæ trichotomæ annuæ sæpius villosa. Folia opposita ovata integra triplinervia in petiolum ciliatum attenuata. Capitula solitaria terminalia inter ultima folia bracteantia sessilia. Ligulæ luteæ. Disci atro-purpureus, convexusque ut in Rudbeckiâ.

1. S. PROCUMBENS (Lam. journ. 2. p. 176. t. 33. ill. t. 686), caule procumbente aut diffuso, foliis ovatis, ligulis aristas achenii superantibus, acheniis disci exterioribus alâ continuâ ciliatâ cinctis et aristellis binis superatis. (1) in Mexico (Cav. Ort. Née), circa urbem (Berl. pl. exs. n. 872). Sanvitalia villosa Cav. ic. 4. t. 351. Lorentea atro-purpurea Ort. dec. 4. p. 42. t. 5. — Pluk. phyt. t. 34. fig. antepenult. (v. v. c. et s. sp.)

♀. oblongifolia, foliis oblongis, petiolis vix subciliatis, capitulis dimidiis minoribus, ligulis 6-7. — in Mexico ad Chapultepec legit Berlandier (pl. exs. n. 372). An species propria? sed fructus non satis vidi. (v. s.)

2. S. ACINIFOLIA, caule procumbente aut diffuso, foliis oblongis, ligulis aristas achenii superantibus, acheniis disci exter. muricatis, inter. lævibus alâ dentatis et ciliatâ cinctis vix aristellatis. (1) in Mexico legit cl. Alaman. Habitus priorum. Ligulæ magis ovales. (v. s.)

3. S. OCYMOIDES, caule diffuso erectiusculo, foliis ovatis, aristis radii ligulas superantibus, acheniis disci compresso-tetragonis, exter. muricatis exalatis brevissimè biaristatis, inter. lævioribus subalatis paulò longius aristatis. (1) in Mexico prov. Texas legit cl. Berlandier (pl. exs. n. 2102!). Herba et folia loc. Oeymi. Petioli ciliati. Ligulæ minimæ. Achenia longiora quàm priorum. (v. s.)

4. S. TRAGIÆFOLIA, caule procumbente aut diffuso, foliis ellipticis, aristis radii ligulas superantibus, acheniis disci fere omnibus compresso-tetragonis coloribus exalatis aut vix subalatis. (1) in Mexico ad San-Fernando de las Presas legit cl. Berlandier (pl. exs. n. 2233!) et in ditione Oaxacanâ Andrieux (pl. ex. n. 3001). Folia supernè magis scabra quàm priorum.

5. S. HELIANTHOIDES (Rich. in Willd. sp. 3. p. 2190), caule erecto, foliis ovato-oblongis apice serratis. — in Peru. Spreng. syst. 3. p. 580 excl. syn. Capitula magnitudine Helianthi gigantei, pedunculata. An certè congener? Cat. fig. f.

#### CCCLXV. ANAITIS.

Capitulum multiflorum radiatum, ligulis foemineis multistriatis persistentibus, fl. radii herm. tubulosis.... Invol. campanulatum 2-3-seriale, squamis imbricatis obtusis. Recept. convexum, paleis deciduis complicatis calloso-acuminatis. Styli radii brevissimè exserti, disci. Achenia glabra minutissimè et parcè muriculata, radii trigona rostrata calva, disci plano-compressa subalata emarginata aut bidentata. —

*Suffruticulus Acapulcensis* ramis teretibus divaricatis, junioribus pubescentibus. Folia opposita oblonga basi attenuata integerrima pubescentia. Rami apice longè aphylli 1-cephali. Flores radii lutei, disci subfusci. — Anatisdis deæ pudicitiz nomen selegi ob stylos florum fœmineorum vix exsertos.

1. *A. ACAPULCENSIS*. ♂ in Mexico occidentali circa Acapulco legit cl. Hænke. Folia 5-6 lin. longa, 1-1½ lata. Pedunculi circ. 5 pollic. (v. s. in h. Hænke à cl. de Sternberg comm.)

## CCCLXVI. OLIGOGYNE.

Capitulum pluriflorum heterogamum, fl. radii 1-5 fœmineis ligulatis, disci 6-8hermaphr. tubulosi. Invol. 1-seriale erectum, squamis 5 ovali-oblongis foliaceis. Recept. planum, paleis lineari-lanceolatis planis. Styli fl. herm. rami exserti hispidi. Achænia obcompressa seu tertiuscula submuriculata glabra aristis 2-4 conicis rigidis superata. — *Herba Americana* habitu *Sanvitaliæ*, diffusæ, puberulæ. Folia opposita petiolata ovata serrata subtripplinervia. Pedicelli axillares 1-cephali. Capitula parva. Flores flavi. — Genus affine *Sanvitaliæ*.

1. *O. BAHIENSIS*, pedicellis capitulo ferè brevioribus, achæniis teretiusculis. — circa Bahiam Brasiliæ legit cl. Blanchet (pl. exs. n. 1706!). Caules debiles pedales glabriusculi. Folia sparse puberula 2 poll. longa, pollicem lata, petiolo 8-10 lin. (v. s.)
2. *O. MZOAPOTAMICA*, pedicellis capitulo brevioribus, achæniis obcompressis. — in Brasiliæ provinciâ Rio-Grande (h. Mus. imp. Bras. n. 874). Valdè affinis *O. Bahiensis*. Caules basi subrepentes adscendentes ad apicem adpressè pubescentes. Involucrum adpressè villosum. Ligulæ 3-4 pallidæ. (v. s. in h. Mus. reg. Par.)
3. *O. TAMPICANA*, pedicellis capitulo multò longioribus, achæniis exter. obcompressis. — circa Tampico de Tamaulipas Mexicanorum legit cl. Berlandier (pl. exs. n. 61!). Folia 9-12 lin. longa, 6-9 lata, petiolo 4 lin. longo. Caules 6-9 poll. longi, ramosissimi, adpressè pubescentes. (v. s.)

CCCLXVII. *SYNEDRELLA* Gærtn. fr. 2. p. 456. t. 171. Rich. in Pers. ench. 2. n. 1901. Cass. dict. 51. p. 470. Less. syn. 233. — *Ucacon* sp. Adans. — *Verbesinæ* sp. Linn.

Capitulum pluriflorum heterogamum, fl. radii circ. 8 subbiseriatis ligulatis fœmineis, fl. disci circ. 8 hermaphr. tubulosi. Invol. oblongum oligophyllum erectum. Recept. parvum planum paleaceum. Ligulæ disco circ. æquales. Achænia radii obcompressa ovalia biaristata aut mutica margine alata, alis laciniatis, disci linearia subtriquetra integra 2-3-aristata. — *Herba Americana* dichotoma. Caulis teres glabriusculus. Folia opposita petiolata ovalia utrinquè attenuata integra parcè villosa. Capitula in axillis ramorum aut foliorum superiorum sessilia subglomerata. Flores lutei.

1. *S. NODIFLORA* (Gærtn. l. c.). ① in littore Mexicano prope Campêche (H. et B.), ins. Cubâ (La Sagr.), Jamaicâ (Sl.), Guianâ (Perr.) et verisim. introducta in Senegaliâ (Perr.). *Verbesina nodiflora* Linn. amœn. 4. p. 290. Willd. sp. 3. p. 2226. — Dill. elth. 53. 1. 45. f. 53. Sloan. hist. 1. t. 154. (v. s.)

CCCLXVIII. *CALYPTOCARPUS* Less. syn. 221 et in Linnaâ 1834. p. 269.

Capitulum multiflorum heterogamum, fl. radii ligulatis 1-seriatis fœmineis, disci hermaphrod. tubulosi 5-dentatis. Invol. 5-phyllum.

Recept. paleaceum. Achænia conformia erostris plano-obcompressa biaristata interruptè et angustè alata. — Herba Mexicana gracilis dichotoma annua procumbens. Folia decussata petiolata scabra dentata. Capitula solitaria lutea brevissimè radiata pedunculata terminalia demum alaria. — Genus proximum *Heterospermo*, *Synedrella*, etc., sed achæniis cortice crasso donatum.

1. *C. VIALIS* (Less. ll. cc.). (1) ad vias prope Jalapam Mexic. Caulis basi rãcens. Folia ovata rhombea albo-hirsuta scabra. Invol. squamæ acuminatæ æquales. Paleæ scariosæ floribus breviores.

CCCLXIX. ELECTRA DC. non Panz. (1).

Capitulum pluriflorum heterogamum, fl. radii circ. 5 ligulatis fœmineis, disci circ. 8-10 tubulosis hermaphr. 5-dentatis. Invol. dupler, squamis exter. 5 linearibus distantibus, 5 inter. ovali-oblongis submembranaceis basi inter se et ferè cum ext. concretis. Recept. angustum planum, paleis oblongo-linearibus coloratis persistentibus flores ferè superantibus onustum. Cor. disci glabræ abruptè in faucem dilatata. Styli rami cono hispido terminati, fl. radii glabri. Achænia obcompressa plana elliptico-oblonga utrinquè subangustata apice obtusè truncata glaberrima calva. — Fruticulus Mexicanus glaber erectus. Rami juniores tetragoni, adulti teretes. Folia opposita in petiolum brevem angustata oblongo-lanceolata acuminata grossè serrata penninervia. Capitula pedicellata sæpius terna corymbosa ovali-oblonga. Cor. et recept. paleæ luteæ, undè nomen ad electrum seu succinum alludens. *Electra* uxor *Atlantis* alludit etiam ad plantam transatlanticam.

1. *E. MEXICANA*. 5 in Mexico circa Villalpando ultra Guanajuato legit. a mecum comm. cl. Mendèz et Alaman. Etiam habui sine loci design. ex ill. Neesii, (v. s.)

CCCLXX. CHRYSANTHELLUM Rich. in Pers. ench. (1810) 2. p. 471. Cat. dict. 9. p. 150. Less. syn. 234. — *Chrysanthellina* Cass. dict. 25. (1822) p. 394. — *Sebastiania* Bertol. opusc. 1822. p. 37 non Spreng. — *Collæa* Spreng. n. 3. (1826) p. 622 non DC. 1825. — *Anthemidis Bidentis* et *Verbesinæ* sp. Linn.

Capitulum pluriflorum radiatum, ligulis fœmineis 1-serialibus linearibus bidentatis, disci hermaphr. 5-dentatis. Invol. biseriale, ext. squamis 5 paulò brevioribus erectis, int. squamis 10 subcoloratis. Recept. planum, paleis linearibus. Styli fl. herm. ramis in appendicem productis. Achænia calva, exter. oblonga truncata emarginata aut truncata alâ minimâ crassâ cartilagineâ utrinquè aucta. — Herba annua glabræ parvulæ, ramulis subnudis apice 1-cephalis. Folia inter se diversa, radicalia petiolata ovalia basi cuneata, summa oblongo-linearâ apice tridentata. Capitula parva. Flores lutei. — Genus ex habitu naturale, ex caract. facilè dividendum, sed tunc genera tot quot sunt species admittenda, meo sensu invitâ naturâ.

1. *C. PROCUMBENS* (Rich. l. c.), achæniis omnibus glabris lævibus, exter. retinensculis collo brevissimo superatis, inter. obcompressis truncatis. (1) in paucis subhumidis insularum Caribæarum nempe in Jamaica? (Sloan?), Santo-Domingo (Bert.). *Anthemis Americana* Linn. sp. ed. 1. p. 895. *Bidens apiculata* Linn. amœn. 5. p. 403. *Euphthalmum* Linn. hort. cliff. 414. *Verbesina natica* Linn. sp. ed. 2. p. 1273. Lam. ill. t. 686. f. 2 bona. *Chrysanthellum*

(1) *Electra* Panzer = *Schismus* ex Steud.

fasciculata et fortè *C. gracilis* Cass. dict. 25. p. 391? *Sebastiania heterophylla* Bectol. opusc. scient. 1822. p. 37. Collæ procumbens Spreng. syst. 3. p. 632. — Sloan. jam. 1. t. 155. f. 1? Plum. ed. Barm. t. 86. f. 2. Variet caulibus erectis procumbentibusve et formâ foliorum. (v. s. comm. à cl. Bertero.)

2. *C. SWARTZII* (Cass. dict. 25. p. 392. sub *Chrysanthellinâ*), achæniis omnibus obcompressis, exter. glabris, inter. ad angulos ciliatis. ① in insulis Caribæis nempe in pascuis subhumidis Jamaicæ (Sw.) et ins. Cubæ circa Havanæ (R. de La Sagra!). *Verbesina mutica* Swartz obs. 314. t. 8. f. 1. Herba variat etiam procumbens diffusa et erectiuscula et foliis polymorphis. (v. s.)

3. *C. SENEGALENSIS*, achæniis omnibus obcompressis glabris, exter. tuberculosis immarginatis, inter. lævibus alâ marginatis. ① in demissis humidis Senegambiæ circa Casam Senegalensem dictam, april. fl. legit cl. Perrottet. Bidens marginata Perrottet in litt. 1833. Folia hinc indè opposita multipartita. (v. s. sine flor.)

4. *C. INDICUM*, achæniis omnibus obcompressis apice brevissimè emarginatis, margine callosis glabris, hinc lævibus planis, illinc convexis striatis. ① in Indiâ orient. ad Gajpur et Sukanaghur. Bidens? *Bardana* Wall. cat. et herb. n. 3231. comp. n. 401. *Achillea*? *Bardana* Hamilt. herb. ex Wall. Ligulæ paucæ parvæ. (v. s. comm. ab hon. Cur. Ind. or. et à cl. Royle.)

CCCLXXI. *NEURACTIS* Cass. dict. 34. p. 496. Less. syn. 232.

Capitulum pluriflorum heterogamum, fl. radii 3 ligulatis foemineis, disci hermaphr. tubulosi 5-dentatis. Invol. subbiseriale disco multò brevius, squamis oblongis, exter. brevioribus. Recept. planum medio nudum, paleis ad margines oblongo-linearibus planis. Ligulæ amplæ ultrâ medium trifidæ. Styli fl. herm. rami exerti revoluti in appendicem gracilem longissimam producti. Achænia obcompressa? oblonga glabra apice breviter in collum producta. Pappus nullus aut margo incompletus dentatus. — Herba Javanica glabra diffusa ramosa multifloris. Folia alterna petiolata bipinnatifolata, lobis oblongis integris mucronatis. Capitula pedicellata solitaria. Cor. flavidæ. Ligulæ nervis fuscis lineatæ. Genus *Chrysanthello* valdè (fortè nimis) affine, sed mihi ignotum.

1. *N. LESCHERNAULTII* (Cass. l. c.). ① in Javâ.

CCCLXXII. *GLOSSOCARDIA* Cass. dict. 19. p. 62. Less. syn. 233. — *Verbesinæ* sp. Linn. f. — *Zinnia* sp. Retz.

Capitulum pluriflorum, fl. radii 1-2 (imò interdum nullis) ligulatis foemineis, disci circ. 5 hermaphr. tubulosi 4-dentatis. Invol. oblongum, squamis paucis subimbricatis, externis brevissimis. Recept. paleis linearibus deciduis membranaceis inter flores onustum angustissimum. Stylus fl. herm. ex Less. apice appendiculatus. Achænium disci plano-obcompressum in angulis 4 villosissimum, subbialatum, aristis binis rigidis lævibus ex angulis lateralibus ortis coronatum. — Herba Indica annua multicaulis diffusa. Folia alterna basi linearia. Capitula pedunculis nudis brevibus insidentia solitaria flava.

1. *G. BOSVALLÆ* (DC. in Wight contrib. p. 19). ① in Indiâ orient. (Koen. ex Linn. f.), in dumetis aridis circa Madras (Wight!), in Carnatic (Belang.). *Verbesina Bosvallæ* Linn. suppl. (1781) p. 379. *Verbesina Boswallia* Willd. sp. 3. p. 2223. Linn. herb. et Roxb. fl. ind. 3. p. 443 ex Wight! *Glossocardia linearifolia* Cass. dict. 19. p. 62. *Pectis meifolia* Wall. cat. et herb. n. 308. Folia non linearia sed lineariloba! Herba esculenta odore et sapore Feniculi (Koen. in Linn. f.). (v. s. comm. ab ampl. Anlâ merc. angl. Ind. or.)

CCCLXXIII. *HETEROSPERMUM* Willd. sp. 3. (1803) p. 2129. Cass. dict. 12. p. 128. H. B. et Kunth nov. gen. am. 4. p. 245. Less. syn. 233. — *Heterosperma* Cav. ic. 3. (1794) p. 34. t. 267.

Capitulum pluriflorum heterogamum, fl. radii 1-3 ligulatis foemineis, disci 5-15 tubulosis hermaphr. Invol. duplex utrumque oligophyllum, squamis erectis oblongis, exter. ciliatis. Recept. angustum, paleis oblongo-linearibus planis obtusis. Styli rami apice appendiculati. Achænia obcompressa glabra alis incrassatis cincta difformia, radii ovalia extrinseca calva, disci lineari-oblonga rostrata biaristata, aristis retrorsum aculeolatis. — Herbarum Americanarum annuæ oppositè ramosæ. Caulis glaberrimus aut bifariam pilosus. Folia opposita dentata lobata aut pinnatipartita glabra. Pedunculi ex dichotomiis aut apicibus ramorum graciles 1-cephali. Cor. flavæ.

1. *H. PINNATUM* (Willd. sp. 3. p. 2129), caule glabro, foliis pinnatipartitis, lobis linearibus acutis integris aut apice dentatis. (1) in Mexico (Cav.) circa urbem et in valle Tolucanâ (Berl. pl. exs. n. 576 et 1081!), in prov. Leoninâ ad occidentem urbis Guanajuato (Mend.!). *Heterosperma pinnata* Cav. ic. 3. t. 267. (v. s. sp. et v. c.)
2. *H. DIVERSIFOLIUM* (H. B. et Kunth nov. gen. am. 4. p. 246. t. 384), caule bifariam piloso, foliis aliis obovato-cuneatis indivisis apice crenato-serratis, aliis trifidis. (1) in aridis prope urbem Quito et in convalle Guallagambenâ (v. c.)
3. *H. MARITIMUM* (H. B. et Kunth nov. gen. am. 4. p. 245. t. 383), caule bifariam piloso, foliis obovato-cuneatis indivisis apice crenato-serratis. (1) in maritimis Peruviae prope Truxillo. *Heterosperma ovatifolia* Cav. dem. bot. p. 204? *H. ovatum* Willd. l. c.?

CCCLXXIV. *GLOSSOGYNE* Cass. dict. 51. p. 475 non Less. *Bidentis* sp. am.

Capitulum multiflorum heterogamum, fl. radii ligulatis foemineis, disci hermaphrod. tubulosis 5-dentatis. Recept. planum demum convexum paleolatum. Invol. 2-seriale breve adpressum. Stylus fl. herm. brevis in stigmata 2 hirta longissima productus. Achænia linearia angulata aristis 2 retrorsum setosas gerentia. — Herbarum glabrae erectae, basi suffruticosae dichotomae, nunc frutescentes pedunculos plurimos nudos simplices gerentes. Folia alterna ad collum conferta breviter pinnatipartita, lobis linearibus acutis integerrimis, summa pauca parva linearia indivisa. Capitula erecta ebracteata lutea. — Cassinius *Bidentis*, Lessingius *Salmea* affinem judicant.

1. *G. PINNATIFIDA* (DC. in Wight contrib. p. 19), caulibus dichotomis, achæniis aristis valde divaricatis. 2. in India orient. ad Goruckpore, Kamisphur, etc. legerunt cl. Heyne et Wight. *Bidens pinnatifida* Wall. cat. et herb. n. 30. *Coreopsis*? *Bacana Heyne*! herb. Zinnia *Bidens* Retz obs. 5. p. 28 ex Roxb. l. ind. 3. p. 435 teste Wight. (v. s. comm. ab hon. Cur. merc. angl. Ind. or. et à cl. Wight.)
2. *G. TENUIFOLIA* (Cass. et Less. ll. cc.), caulibus dichotomis, achæniis aristis erectis. 2. in Novâ-Caledoniâ (Labill.!), Novâ-Hollandiâ ad Port-Jackson et insulâ Guam Marianarum (Gandich.), *Bidens tenuifolia* Labill. ! sert. calif. p. 44. t. 45. (v. s. comm. à cl. Labill.)
3. *G. PEDUNCULOSA*, caulibus fruticosis basi simplicibus apice pedunculos plurimos longissimos nudos monocephalos gerentibus, achæniis aristis erectis; in Novâ-Hollandiâ parte tropicâ in collibus rupestribus ad Caput Cleveland legit cl. Cunningham. Caulis crassitie pennae auserinae. Folia elongata linearia pinnatipartita, lobis paucis alternis integerrimis. Pedunculi pedales cauli

Capitulum fructigerum et cor. disci omninò generis sed ligulas in specim. deflorato non vidi. (v. s. comm. à cl. Cunningham.)

*N. B. Corsoopals Tannensis* Forst. ex Spreng. syst. 614 ad hoc genus pertinet, ex patriâ ad secundam speciem, ex testimonio Sprengelii (in litt. ad Wall.) ad primam referenda. *Glossogyne Chinensis* Less. syn. 212 mihi omninò ignota. *Bidens Chinensis* mihi et ut antumo Botaniciis cognita, est certè *Bidentis species*!

## CCCLXXV. DELUCIA.

Capitulum multiflorum radiatum, fl. radii 1-seriatis fœmineis ligulatis, fl. disci hermaphr. tubulosis 5-dentatis. Invol. duplex subæquale, squamis lineari-lanceolatis erectis, exter. foliaceis, inter. coloratis. Recept. planum paleaceum, paleis linearibus. Anth. nigricantes. Styli fl. herm. rami cono superati. Achænia disci et radii obcompressa linearia aristis 3 retrorsum hispidis coronata. — Herba Mexicana subsimplex vix pedalis glabra. Folia opposita petiolata trisecta, segmentis ovatis utrinquè acutis grossè dentato-serratis. Capitula solitaria longè pedicellata lutea. — Genus affinè *Bidenti* sed ligulæ fœmineæ fertiles, Narvalinæ sed ligulæ plures, aristæ achæniî 3. persistentes et achænia linearia immarginata. Genus montanum dixi in honorem cl. geologi et physici Th. Andr. Deluc, qui laboribus circa barometri et thermometri usum geographiæ botanicæ progressus egregiè paravit.

1. *D. oenanthoides*. 2f? in Mexici montanis circa urbem legerunt cl. Berlandier (pl. exs. n. 920), Alaman, Mairet, etc. Folia ferè omninò Imperatoriæ Ostruthii sed opposita. Capitulum è longinquo ferè Trollii Asiatici. (v. s.)

CCCLXXVI. *NARVALINA* Cass. dict. 38. p. 17. 59. p. 320. Less. syn. 234. — *Needhamia* Cass. dict. 34. p. 335 non Scop. nec Brown.

Capitulum pluriflorum radiatum, ligulis solitariis (aut 2-3) fœmineis, fl. disci circ. 8 hermaphr. 5-dentatis. Invol. biseriale, exter. breve squamis circ. 3 foliaceis acutis, inter. longius disco brevius squamis circ. 5 oblongis erectis apice coloratis. Recept. planum, paleis squamas inter. referentibus. Ovaria obcompressa ovalia amplà latè marginata ciliata. Pappus caducus biaristatus, aristis crassis fragilibus coloratis triquetrosulcatis secus angulos spinellas strigosas gerentibus. — Frutex Domingeensis glaberrimus. Folia opposita petiolulata obovata serrata coriacea. Capitula corymbosa. Flores lutei.

1. *N. DOMINGENSIS* (Cass. dict. 38. p. 17). 5 in Sancto-Domingo. *Needhamia Domingensis* Cass. dict. 34. p. 335. — Ramum huius speciei pertinentem in horto Paris. lectum recepi à cl. Adolph. Brongniart qui eum specimine Cassinii comparavit. Differt à descr. Cass. tantùm ligulis 2-3 nec solitariis (v. s.)

CCCLXXVII. *THELESPERMA* Less. in *linnaea* 1831. p. 511. syn. p. 234.

Capitulum multiflorum homogamum. Invol. biseriale utrumque circ. 8-phyllo, exter. breve squamis lanceolatis vix inter se concretis, inter. campanulatum squamis longè concretis striatis. Recept. planum, paleis oblongis medio nervigeris marginibus latè albo-membranaceis. Styli rami apice appendiculati. Achænia obcompressa-tetretiuscula erostria. Pappus biaristatus, aristis oppositis retrorsum fimbriatis trigono-conduplicatis sulcatis persistentibus. — Herba Brasiliensis glabra perennis. Caulis teres oppositè et ideò trichotomè ramosus, rigidus. Folia

opposita biternatim seu bipinnatim secta, lobis linearibus elongatis acutis integerrimis. Capitula ramos apice nudos terminantia. Cor. obscure luteæ ex sicco videntur. — Ex invol. gamophyllo subglanduloso et stigmatum structurâ ad Tagetines accedit, sed pappus Heliantheorum.

1. *T. SCABIOSOIDES* (Less. l. ll. cc.). 2l. in Brasiliâ (Sellow. l.), in provinciâ Rio-Grande (h. imp. Bras. n. 910). Herba bipedalis habitu ferè Scabiosæ columbariæ Gramuntinæ sed glabra et tenuius secta. (v. s. comm. à Mus. reg. Berol.)

CCCLXXVIII. ISOSTIGMA Less. in *linnæa* 1831. p. 513. *syn.* 235.

Capitulum multiflorum radiatum, ligulis fœmineis oblongis 1-seriis, fl. disci hermaphr. tubulosis 5-dentatis. Invol. campanulatum 2-3-seriale, squamis liberis, exter. brevioribus angustioribus. Recept. planum paleaceum. Styli rami in fl. ♀ et etiam ♀ in appendicem lineari-filiformem longissimam et hispidam producti. Achænium lineare striatum plano-obcompressum subalaturn rostro achænio simili terminatum, alis in mucrones aristasve 2 divergentes læves productis. — Herbæ Brasilienses perennes glaberrimæ. Rhizoma liguosum subhorizontale. Caules erecti nudi 1-cephali simplicissimi. Folia ad collum orta alterna, nunc indivisa, sæpius biternatim seu bipinnatim secta, lobulis integerrimis. Capitula magna speciosa, in sicco fusca, cuncta aquam colore luteo tingentia. — Ligulæ in herbar. sæpè nullæ, verisim. deciduæ.

1. *I. SIMPLICIFOLIUM* (Less. in *linnæa* 1831. p. 513), caule apice equamato tereti, foliis integris linearibus basi angustatis apice subdentatis raris trifidis, achæniis aristis latitudinem ovarii æquantibus. 2l. in Brasiliâ. (Less.)
2. *I. PRUCEDANIFOLIUM* (Less. l. c. p. 514), caule tereti equamato, foliis multipli-pinnatisectis, segmentis trigonis supra canaliculatis, achæniis aristis divergentibus latitudinem ovarii non æquantibus, interdum subnullis. 2l. in Brasiliâ (Sellow. l.), in prov. Rio-Grande (h. Mus. imp. Bras. n. 1060). *Tragoceras pencedanifolium* Spreng. *syst.* 3. p. 576. Caulis 2-3-pedalis. Capitula 15-16 lin. diam. (v. s. comm. à Mus. reg. Berol.)
3. *I. SPACIOSUM* (Less. l. c. p. 515), caule apice incrassato squamigero, foliis ternatim seu pinnatim sectis bisectisve, segmentis subtrigonis crassiusculis, achæniis aristis vix dimidiam ovarii latitudinem æquantibus. 2l. in Brasiliâ cum præcedente. (Less.), in prov. Rio-Grande ex h. Mus. imp. Bras. n. 1039 (v. s. in h. Mus. reg. Par.)
4. *I. CRITHMIFOLIUM* (Less. l. c. p. 515), caule tereti squamato, foliis ternatim biternatimve sectis, segmentis planis striatis, achæniis aristis latitudinem ovarii superantibus. 2l. in Brasiliâ cum præcedentibus (Less.), in prov. Rio-Grande ex h. Mus. imp. Bras. n. 1058! (v. s. in h. Mus. reg. Par.)

Subtrib. III. *FLAVERIÆ* Less. *syn.* 233. — Gen. *Heliantheorum* Milliarum Cass.

Capitula nunc uni- vel pauciflora, densè aggregata, alia heterogama è uno fœmineo et reliquis hermaphroditis, altera intermixta homogama, nunc multiflora fl. fœmineis pluriserialibus reliquis hermaphroditis. Achænia calva exalata teretiuscula basi longè attenuata. Styli fl. hermaphr. rari sæpius exappendiculati. Recept. in capitulis paucifloris epaleaceum, in multifloris paleaceum. — Herbæ glabriusculæ Austr.-Americanae. Folia opposita tri- seu multiplinervia. Flores flavi.



CCCLXXIX. FLAVERIA Juss. gen. (1789), p. 186. Cass. dict. 17. p. 127. Less. syn. 233. — Vermifuga Ruiz et Pav. syst. p. 210. prod. fl. per. t. 24. — *Milleria* sp. Cav.

Capitula pauciflora glomerato-fasciculata, nunc omnia homogama, nunc heterogama, fl. unico ligulato fœmineo, cæteris hermaphrod. 5-dentatis. Invol. 3-4-phyllum connivens oblongum, squamis longitudine æqualibus, unâ exter. majore subconcavâ. Recept. parvum epaleaceum. Styli fl. herm. rami apice exappendiculati. Achænia oblonga calva striata glabra. — Herbæ annuæ Americanæ glabræ aut vix secus ramos puberulæ. Folia opposita sessilia interdum basi attenuata dentata aut integra sæpius trinervia. Capitula secus ramulos unilateralia plus minus conferta et ideò cymas glomerulosve referentia. Cor. luteæ.

1. *F. CONTRAYERBA* (Pers. ench. 2. p. 489), erecta, foliis lanceolatis serratis trinerviis basi attenuatis, capitulis confertissimis, omnibus flosculosis aut uniligulatis, ligulâ discum non superante. (1) in Amer. austr. regionibus occidentalis nempe in Chili (Fenill.), Peruvia (Ruiz et Pav. Domb. | Gaudich. etc.), nec non in Mexico inter Fernando et Santander (Berl. | pl. exs. n. 2259), Tampico et Real del Monte (Berl. | n. 38), et ad occid. urbis Guanaxuato (Mend. |), demum in insulâ Cubâ (Rio de La Sagra). Sims bot. mag. t. 2400. *Milleria Contrayerba* Cav. ic. 1. p. 2. t. 4. *M. contrahierba* Lam. dict. 4. p. 183. illustr. t. 710 f. 4. *Vermifuga corymbosa* R. et Pav. | syst. 210. *Flaveria capitata* Juss. ex Smith in Rees cycl. v. 14. n. 1. *Ethulia Bidentis* Linn. mant. p. 110 ex Smith. — Fenill. obs. 3. t. 14. Herba alexipharmaca et colore luteo pannos inficiens, vulgò *Contrayerba* et *Matagusanos* dicta. (v. v. c. et s. sp.)
2. *F. ANGUSTIFOLIA* (Pers. ench. 2. p. 489), erecta, foliis lineari-lanceolatis sessilibus trinerviis, capitulis corymbosis uniligulatis, ligulâ discum superante. (1) in Mexico (Cav. Berl. | pl. exs. n. 640). *Milleria angustifolia* Cav. ic. 3. p. 12. t. 223. Variat foliis apice serrulatis et integerrimis, ligulis majoribus et dimidiis minoribus, corymbis fastigiatis terminalibus et etiam ad apices lateraliû ramorum. An fortè duæ species hic confusæ? *Flav. integrifolia* et *Fl. radicans* icon. fl. Mex. ined. An icon Cav. 1. t. 4 hæc etiam, saltè quoad florum analyses, pertinet? Sæpè vidi in hortis sub nom. *Fl. Contrayerba* cultam. (v. v. c. et s. sp.)
3. *F. LINEARIS* (Lag. nov. gen. (1815), p. 33. n. 407), procumbens, foliis sessilibus connato-vaginantibus linearibus integerrimis carnosis enerviis, capitulis confertè corymbosis uniligulatis. (1) in insule Cebæ maritimis prope Havanam. *Fl. maritima* H. B. et Kunth nov. gen. am. 4 (1820), p. 285.
4. *F. BONARIENSIS*, erecta, foliis sessilibus subconnatis linearibus acuminatis membranaceis tenuissimè trinerviis, capitulis cymoso-corymbosis. (1) circa Bonariam legit cl. Bacle. Flores ign. (v. s.)

† *Species exclusæ.*

*F. PERUVIANA* Juss. = *Piqueria artemisioides*.

*F. REPANDA* Lag. = *Broteroa trinervata*.

CCCLXXX. BROTEROA. — *Brotera Spreng. in Schrad. journ.* 1800. p. 2. p. 186. t. 5. Pers. ench. n. 1959. Cass. dict. 34. p. 304. 59. p. 236 non alior. (1). — *Naueemburgia Willd. sp.* 3 (1803). p. 2393. pessimè! *Hoffmans. verz.* 1824. p. 237. Less. syn. 232 non *Mænch* (1802). — *Flaveria* sp. *Spreng.* (1826). Lag.

Capitula in glomerulum ovatum densè spicato-conferta sessilia inter se varia, alia biflora heterogama seu homogama, alia 1-flora femi-

(1) *Brotera* Cav. (1799). = *Pentapteris* sp. — *Brotera* Willd. (1803). = *Cardapatum*. — *Brotera* Spreng. (1802). = *Hyptis* sp.

nea aut hermaphrodita. Invol. squamæ nunc solitariae concavæ amplæ, nunc 2-3, alteræ amplæ concavæ, alteræ parvæ. Recept. punctiforme nudum. Cor. tubulosæ extus pilis articulatis confertis hispidæ 5-fido, fœmineæ filiformes subligulatæ? Styli fl. herm. ramis apice exappendiculatis. Achænia obcompressa obovato-oblonga glabra calva, fl. herm. minora quàm fl. fœm. ex Cass. — Herba Austr.-Americana glabra, ramis paucis divaricatis. Folia opposita sessilia ovali-lanceolata acuminata trinervia integerrima serrata aut subrepanda. Glomeruli ad apices ramorum aut axillas super. pauci subumbellati. Cor. flavæ. — Nomen paulò mutavi ut melius cum nomine ill. viri congruat et à diversis homonymis distinguatur.

1. B. TRINERVIA (Pers. ench. 2. p. 498). ①? ②? in America austr. circa Caracas legit cl. Vargas, in Mexico ubi dicitur *Azumiatas* (Lag.). Brotera Contrayerva Spreng. in Schrad. journ. 1800. p. 2. p. 186. 2. 5. excl. syn. Cav. et Feuill. Brotera Sprengelii Cass. dict. 34. p. 304. Nanemburgia trinervia Willd. sp. 3. p. 2393. Flaveria repanda Lag. nov. gen. p. 33 ex Spreng. syst. 3. p. 500. neu. entd. 3. p. 42. Herba basi indurata ferè lignosa et verisim. perennis, etiam ex hortis Europæis annua dicitur. Variat foliis termo-verticillatis (v. s. sp. comm. à cl. Vargas.)

#### CCCLXXXI. CLAIRVILLEA.

Capitula fasciculato-capitata pluriflora heterogama, fl. radii ligulatis circ. 5 uniserialibus fœmineis, fl. disci 4-5 tubulosis hermaphr. 5-dentatis. Invol. ovatum, squamis ovatis arcè imbricatis. Recept. angustum paleis oblongo-linearibus in margine onustum, centro epilaceum. Styli fl. herm. rami apice appendiculati hispiduli. Achænia ovata glabra calva. — Frutex Austr.-Americanus. Rami teretes, juniores densè albo-tomentosi, adulti glabri. Folia opposita connato-vaginantia ovato-lanceolata acuminata serraturis distantibus callosospinellos serrata, suprâ 5-nervia, juniora araneosa, adulta rugosa glabriuscula, subtus densè albo-tomentosa. Capitula subsessilia in glomerulum ovatum terminalem densè aggregata, centralia præcociora. — Genus Broteræ affine sed ligulis circ. 5, styli ramis appendiculatis et dispositio capitulorum faciliè diversum. Dixi in memoriam cl. de Clairville Gail botanici et entomologici de historiâ naturali Helveticâ benè meriti.

1. C. QUINQUENERVIA. ③ in Amer. australi legit ill. Dombey, sed locus ibi non design. verisim. in Peruvia (v. s. comm. à Mus. reg. Par.)

CCCLXXXII. ENHYDRA. — Enydra Lour. fl. coch. (1790), ed. Willd. 2. p. 614 — Meyera Schreb. gen. (1791), n. 1318. Less. syn. 236. — Sobrya R. et P. fl. per. prod. (1794) p. 109. t. 24. — Enydra, Meyera et Sobrya Pers. cat. n. 1964, 1896 et 1966. — Hingcha Roxb. (1814) sine descr. — Tetraotis Reuss. (1824), in Blum. bijdr. 892. — Phyllimena Blum. in litt. — Meyeræ sp. Br. — Enydræ sp. Cass.

Capitulum multiflorum heterogamum, fl. omnibus paleâ complicatè involutis, ext. pluriserialibus fœmineis, int. hermaphr. sterilibus. Recept. parvum planiusculum. Paleæ quasi bivalves, subcoriæ arcè conniventes apice hirtellæ. Invol. 4-phyllum, squamis foliaceis 2 oppositis majoribus. Cor. tubo radii filiformi, disci subconico, limbo radii subligulato 5-4-fido, disci infundibuliformi 5-fido. Stigm. 2 et 3-4-serta revoluta versùs apicem hispidula. Achæmium obcompressum et v. crostre paleâ omninò involutum. — Herbæ Asiaticæ et Amer.

canæ fortè etiam Australasicæ, aquaticæ, radicales, oppositifoliæ, Cryphiospermum referentes. Folia oblonga interdum biauriculata. Capitula ad axillas superiores solitaria. Flores albi (an in omnibus?)

\* Asiaticæ, foliis brevissimè petiolatis.

1. *E. FLUCTUANS* (Lour. fl. coch. 2. p. 625), tota glabra, foliis subhastatis serratis brevissimè petiolatis, capitulis pedunculatis. — in aquosis paludibus Cochinchinæ. Meyera fluctuans Spreng. syst. 3. p. 602. excl. syn. R. et Pav. (v. s.?)
2. *E. LONGIFOLIA*, caule glabro apice summo puberulo, foliis basi subhastatis vix petiolatis distanter et minutè serrulatis glabris margine scabris, capitulis sessilibus. ½ in paludosis ins. Javæ prope Bataviam. Tetraotis paludosa Reinw. in Blum. ! bijdr. 892. Phyllimena Blum. ! in litt. Folia lineari-lanceolata, 2 poll. longa, 3 lin. lata. (v. s. in h. Blum.)
3. *E. PALUDOSA*, caule hispido, foliis basi subhastatis vix petiolatis grossè serratis asperis, capitulis sessilibus. ½ in paludosis ins. Javæ prope Bataviam. Tetraotis paludosa Reinw. in Blum. ! bijdr. 892 ! Phyllimena paludosa Blum. ! in litt. Capitula interdum terminalia. Folia prioris sed pilis scabrida et grossius serrata. (v. s. in h. Blum.)
4. *E. HELONCHA*, caule basi glabro, apice hirtello, foliis basi subhastatis vix petiolatis serratis glabris, capitulis sessilibus. ½ in aquosis Indiæ or. in prov. Silhet (Wall.), ad Dumdumma et Golpara (Ham.). Meyera Heloncha Wall. ! cat. n. 3195. comp. 305. Hingcha repens Roxb. cat. calc. 62. Duabus prioribus valdè affinis. Vernac. dicitur *Heluncha* et *Helumochika*. (v. s.)

\*\* Americanæ, foliis sessilibus.

5. *E. SOBRYEA*, glabra, foliis sessilibus basi sagittatis lanceolatis serratis, capitulis.... ½ in Peruvæ stagnosis. Sobryea sessilifolia fl. per. syst. 1. p. 197. Sobrya sessilifolia Pers. ench. 2. p. 473.
6. *E. OBLONGA*, erecta glabra, foliis sessilibus oblongis basi rotundatis obscure dentatis, capitulis breviter pedunculatis. ½ in Peruvæ stagnosis. Sobryea oblonga fl. per. syst. 1. p. 197. Sobrya oblonga Pers. l. c. (v. s. comm. ex herb. Dombey à Mus. reg. Par.)
7. *E. SÆSILIS*, caule erecto glabro sub foliorum jugis puberulo, foliis sessilibus oblongis basi rotundatis apice subacutis parcè serratis glabris, capitulis sessilibus. ½ in aquosis Jamicæ australis (Sw.), Sancti-Domingi (Bert.). Eclipta sessilis Swartz prod. 114. Meyera sessilis Swartz fl. Ind. occ. 1364. Spreng. syst. 3. p. 602. Variat foliis subhirtis ex Sw. (v. s.)
8. *E. MARITIMA*, glabra, foliis oblongis acutis basi rotundatis obsolete serratis, subcarnosis, capitulis sessilibus. ½ in littore Oceani pacifici prope El Callao de Lima (H. et B.). Meyera maritima H. B. et Kunth nov. gen. am. 4. p. 269. Willd. herb. ex Less.
9. *E. SUBCUNEATA*, demissa glabra, foliis sessilibus oblongis basi cuneato-attenuatis apice vix acutis obtusè et parcè dentatis, capitulis ad axillas solitariis sessilibus. ½ in fossis circa Rio-Janeiro speciatim ad viam Brocco leger. cl. Lund et Gandichand. A quatuor prioribus facile differt folia basi nec sagittatis nec auriculatis nec obtusis. An fortè *E. maritima* Less. in linnæa 1831. p. 159 à cl. Chamisso ad fretum Sanctæ-Catharinæ lecta, huc ex loco natali et foliis oblongo-obovatis referenda? (v. s.)

† *Species exclusæ.*

ENTYDRA CÉSULIOIDES Cass. = Cryphiospermum repens.

MEYERA GUINEENSIS Spr. = Cryphiospermum repens.

MEYERA CAPITATA Spreng. quæ Tilesia capitata Mey. esseq. 251. omninò aliena.

TETRAGONOTHECA L'Hér. etiam à Spreng. admissa est omninò diversa.

Subtrib. IV. *TAGETINÆ*. — *Tagetineæ* Cass. dict. 20. p. 367. 59. p. 61. ann. sc. nat. 1829. p. 12. opusc. 3. p. 54. — *Tagetineæ*, *Pectideæ* et pars *Chrysocomearum* Less.

Capitula multiflora, nunc heterogama fl. radii femineis ferè semper ligatis, nunc homogama discoidea. Invol. nunc uniseriale squamis plùs minus concretis, nunc pluriseriale paleis exterioribus bracteiformibus liberis interioribus concretis. Recept. planum epaleaceum. Achæniû erostre striatum elongatum basi attenuatum. Pappus aristis squamellis setisve constans. — *Herbæ* interdum suffruticosæ sæpius annuæ Americanæ glandulas pellucidas grossas in contextu foliorum et involucrorum gerentes et ideò sæpè odoriferæ. Folia opposita aut alterna. — *Pectideæ* à cl. Cassini hûc fortè non inmerito adnexæ, ad *Vernoniaceas* à cl. Lessing rejectæ, nexum inter duas tribus indicant.

Div. I. *TAGETÆ*. — *Tagetineæ*, *Dyssodieæ* et *Archetypæ* Cass. dict. 59. p. 61. — *Tagetineæ* et pars *Senecionearum* Less. 79. p. 236 et 388.

Capitula sæpius heterogama radiata. Invol. squamæ saltè interiores inter concretæ. Cor. disci regulariter 5-dentatæ. Styli rami cono superati. Pappus inæqualis 1-2-serialis aristis squamellisve constans.

CCCLXXXIII. *ADENOPHYLLUM* Pers. ench. 2. (1807) n. 1870. — *Willdenowia* Cav. ic. 1. p. 61. t. 89 non Thunb. — *Schlechtendalia* Willd. sp. 3. (1806) p. 2125 non Less. 1830. nec Spreng. 1829 (1). — *Dyssoda* sp. Lag. — *Tagetis* sp. Less.

Capitulum radiatum. Invol. squamæ lineares concretæ apice productæ in aristam subulatam basi 1-glandulosam, extûs cinctum bracteis sub-1-serialibus aristato-subulatis basi uniglandulosis. Recept. alveolatum seu fimbrilliferum. Achænia elongata. Pappus duplex, ext. paleæ breves truncatæ, inter. elongatæ apice acutè subtrifidæ. — *Herbæ* Mexicanæ *Tagetis* facie. Folia opposita et superiora alterna pinnatisecta, segmentis ovatis aut ferè linearibus aristato-dentatis, rachi inter segmenta aristas hinc indè gerente. Pedunculi 1-cephali sub capitulo incrassati. Cor. crocæ seu coccineæ, in disco dilutiones.

1. *A. COCCINEUM* (Pers. l. c.), foliorum segmentis aliis ovatis dentatis, aliis intermediis capillaceo-subulatis. — in Mexico juxta Doscaminos oppidum legit Ne (Lag.). *Willdenowia* glandulosa Cav. ic. 1. p. 61. t. 89. Lam. ill. t. 685. ic. Cav. *Schlechtendalia* glandulosa Willd. sp. 3. p. 2125. *Tagetes* punctata L. Mexic. ic. ined. *Boebera* Cavanillesii Spreng. syst. 3. p. 544 ex hac et *Dyssodia* porophyllâ mixtis facta videtur. *Dyssoda* Cavanillesii Lag. elench. h. Madr. p. 29. n. 371. Variat statu abnormali nempe floribus disci 6-8-fidis et stylo trifido, sed potius monstrum quàm varietas et perperam Cavanillesius hanc in carnem gener. recepit. Confer Lag. et Cass. ll. cc.

2. *A. CAPILLACEUM*, foliorum lobis omnibus capillaceo subulatis, aut in lobulis subulatos ad basin usque fissis. — in Mexico. *Schlechtendalia* capillacea fl. Mex. ic. ined. An prioris var.? (v. ic. pict.)

(1) Stricto jure nomen *Willdenovianum* servandum esset, sed *Personianum* admodum augeatur synonymia et ne botanici peritissimi nomen generi subdubio sit consecratum.

CCCLXXXIV. *LEBETINA* Cass. dict. 25 (1822) p. 394. 59. p. 68. — *Bæbera* Less. syn. 237 non Willd. — *Dyssodia* Willd. enum. 900 non Cav.

Capitulum radiatum. Invol. cylindraceo-campanulatum 1-seriale, squamis circ. 20 infernè concretis apice variè liberis, bracteis 1-2-seriatis pinnatipartitis lobis apice subulato-piliformibus cinctum. Recept. convexum fimbriatiferum. Achæmium cylindraceum minutissimè pubescens. Pappus duplex, ext. paleis 10 oblongis integris, int. 10 setis basi cuneato-paleatis apice ramoso-pinnatis filiformibus. — Herba Mexicana glabra, Tagetis seu *Dyssodia* facie, erecta ramosa. Folia alterna pinnatipartita, lobis serratis, serraturis in pilum longum desinentibus. Rami apice subnudi, foliis supremis paucis in bracteas desinentibus, 1-cephali. Cor. flavo-aurantiacæ.

1. *L. CANCELLATA* (Cass. dict. 25. p. 394). (1) in Mexico ad occid. urbis Guanaxuato legit cl. Mendez, circa Tamanlipas cl. Berlandier (pl. exs. n. 2140!), et ad Tlapajhua F. W. Keerl (h. Mart.). *Dyssodia* Porophyllum Willd. enum. p. 900 non Cav. *Bæbera* Porophyllum Less. syn. 237. excl. syn. Habitus *Dyssodia* porophyllæ, sed ligulæ multò majores invol. superantes, recept. magis convexum longius fimbriatiferum et pappus duplex! (v. s.)

CCCLXXXV. *DYSODIA* DC. non Lour. (1). — *Dyssodia* Cav. anal. scienc. nat. 6. (1802) p. 334. Cass. dict. 25. p. 396. 59. p. 61. Less. syn. 388. — *Bæbera* Willd. sp. 3. (1803) p. 2125. Cass. dict. 5. suppl. p. 2. non Less. — *Dyssodia* sp. Lag.

Capitula radiata aut rarissimè discoidea. Invol. proprium 1-seriale, squamis plùs minus inter se concretis, rariùs nudum, sæpè extùs bracteis 1-2-seriatis calyculatum. Recept. planum brevissimè hirsuto-fimbriatiferum aut subalveolatum. Styli rami cono pubero superati. Achæmium elongatum tetragono-compressum. Pappus 1-serialis, paleis imà basi indivisis et in pilos setaceos scabros aut barbellato-palmatis fissis, et idèò quasi polyadelphus dicendus. — Herbæ Mexicanæ Tagetis facie. Folia opposita aut rariùs alterna, indivisa aut sæpiùs pinnatipartita, serraturis acutis aut in pilos abeuntibus.

#### Sect. 1. *EUDYSODIA*.

Involucrum bracteatum, bracteis pinnatipartitis, lobis in pilos abeuntibus. Recept. densè hirsuto-fimbriatiferum. — Folia alterna pinnatisecta.

1. *D. POROPHYLLA*, erecta glabra, foliis alternis pinnatipartitis, lobis extrorsum latioribus dentatis, dentibus pilo longo apiculatis, invol. bracteis pinnatipartitis apice setigeris obvallato, ligulis invol. vix longioribus aut nullis. (1) in Mexico. *Dyssodia* porophylla Lag. elench. h. madr. 29. *Dyssodia* glandulosa Less. syn. 388. excl. syn. *Bæbera* Porophyllum Kunth nov. gen. am. 4. p. 198. (excl. Willd. syn.) Spreng. neu. entd. 3. p. 44. Habitus *Lebetina* sed pappus omninò diversus et ligulæ multò minores oblongæ aut nullæ.

a. *radiata*, ligulis oblongis involucrum paulò superantibus. *Dyssodia* porophylla Cav. anal. 6. p. 334. *Bæbera* alternifolia fl. mex. ic. 1. 636. (v. v. c.)

β. *discoidea*, ligulis nullis. *Pteronia* Porophyllum Cav. ic. 3. p. 13. t. 225. Willd. sp. 3. p. 1782. (v. s. sp. in pl. exs. Berl. n. 957.)

(1) *DYSODIA* Lour. = *Serissa*.

Sect. II. BOEBERA. — *Bœbera Willd. Pers.*

Invol. bracteatum, bracteis oblongis linearibusve indivisis. Recept. brevissimi fimbriiferum aut subnudum. — Folia opposita pinuatisecta.

2. *D. CHRYSANTHEMOIDES* (Lag. elench. h. madr. p. 29. n. 364), adscendens aut diffusa subpuberula, foliis oppositis subpinnatisectis, lobis linearibus apice dentatis, capitulis solitariis, pedunculis foliosis, invol. 8-phyllo squamis basi concretis constante bracteis circ. 8 linearibus basi ciliatis cincto, ligulis parvis involucri paulò superantibus. (I) in America frequens ad ripas flum. Missouri et Mississippi (Pursh), in Mexico (Alam.! Mair.!), ad Santa-Fe Bogotensium ubi ex odore dicitur *Ruda* (Lag.). *Tagetes papposa* Vent.! h. cels. t. 36. Michx. fl. 2. p. 132. *Tagetes pumila* hortul. *Dyssodia glandulosa* Cav. dem. bot. 202. *Bœbera chrysanthemoides* Willd. sp. 3. 2125. *Bœbera glandulosa* Pers. ench. 2. p. 459. Variat caule adscendente aut diffuso, glabrescente aut puberulo, foliis variè incisis angustissimis aut sublatioribus, bracteis plus minus ciliatis. An fortè species plures hic confusæ? in omnibus video recept. nudum aut vix alveolatum, achenia compressa tetragona glabriuscula pappo longiora. (v. v. c. et s. sp.)
3. *D. FASTIGIATA*, erecta puberula, foliis oppositis pinnatipartitis, lobis linearibus apice serratis, capitulis corymbosis, pedunculis 1-cephalis subaphyllis, invol. 8-phyllo bracteis 6-8 indivisis cincto, ligulis circ. 8 invol. paulò longioribus. (I) in Mexici montanis (Alam.!) prope Ario Pacuaru et lacum Cuicero (H. et B.). *Dyssodia tagetiflora* Lag. elench. h. madr. p. 29? *Bœbera tagetiflora* Spreng. l. c.? Valde accedit ad *D. chrysanthemoidem*. Fœniculam redolens ex Kunth. (v. s. comm. à cl. Alaman et Berl.)
4. *D. PUBESCENS* (Lag. elench. h. madr. p. 29. n. 365), caule pubescente, foliis pinnatisectis, lobis linearibus apice acutè 3-5-inciso-serratis cæteris integerrimis, pedunculis 1-cephalis foliosis, invol. squamis 8 basi concretis bracteis 6-8 indivisis ciliolatis cincto. — in Mexico (Lag. Hænke!), in Less prope Guanaxuato (Mend.!). *Aster pinnatus* Cav. ic. 3. p. 6. t. 212. *Bœbera pubescens* Spreng. syst. 3. p. 544. Variat caule patulo et erecto. Species *D. chrysanthemoidi* proxima, sed lobis latioribus subcuneatis faciliè distincta. (v. s.)
5. *D. INCANA*, caule erecto hirsuto, foliis pinnatisectis infernè subhirsutis, lobis linearibus acutis canaliculatis, aliis integerrimis, aliis subcuneatis trifidis, terminali à cæteris non distincto, pedunculis 1-cephalis. — in Mexico (Lag. Alam.!), circa urbem (Berl.! pl. exs. n. 631). *Bœbera incana* Lindl. bot. reg. t. 1602. opt. *Dyssodia subintegerrima* Lag. elench. p. 29. n. 366. *Bœb. subintegerrima* Spreng. l. c. Nomen Lag. vetustius sed decipiens et sublobatum. Pappus fusco-nigricans (v. s.)
6. *D. APPENDICULATA* (Lag. elench. h. madr. p. 29. n. 368), glabra, foliis oppositis pinnatisectis, lobis 5-7 ovato-sublanceolatis inæqualiter argutè serratis glandulâ oblongâ apice appendiculatis, invol. bracteato. (I) in Mexico prope Acapulco legit. cl. Née et ad occid. urbis Guanaxuato cl. Mendès. Pappus *D. chrysanthemoidis*. Invol. *D. porophyllæ* si cilia exterioris exceperis. (v. s. comm. à cl. Mendès et Alaman.)

## Sect. III. BOEBEROIDES.

Invol. bracteatum, bracteis ovatis acutis subserratis. — Folia opposita indivisa.

7. *D. GRANDIFLORA*, erecta glabra, foliis oppositis ovali-lanceolatis argutè serratis acuminatis, capitulis ad apices ramorum solitariis, involucri bracteis ovatis subdentatis acuminatis cincto, ligulis invol. multò longioribus, achenio pubescenti-villoso. — in Mexico. *Bœbera grandiflora* ic. fl. mex. ned. t. 635. Paulò refert *Zinniam coccineam*. (v. s. in herb. Hænke. à cl. de Sternberg comm.)

## Sect. IV. GYMNOLENA.

Invol. ebracteatum omninò nudum. Recept. subalveolatum. — Folia opposita indivisa.

8. *D. serratifolia*, erecta glaberrima, foliis oppositis indivisis oblongo-lanceolatis utrinquè attenuatis argutè serratis minutè punctatis, capitulis longè pedicellatis corymbosis, invol. nudo, ligulis oblongis tubulisque involucri ovali-oblongo longioribus, achænio adpressè villosò. — in Mexici prov. Oaxaca ad Atitla et locos vicinos. Habitus ferè Tagetis lucidæ, sed folia ciliata et pappus diversissimus. Recept. vix subalveolatum. (v. s. comm. à cl. Alaman.)

## Sect. ? v. ACIPHYLLÆA.

Invol. nudum inter folia suprema acerosa subbracteiformia sessile. Recept. minutè alveolatum. — Folia opposita integerrima. — Habitu à genere valdè recedit sed cum caract. generico quadrat.

9. *D. acerosa*, suffruticosa ramosissima erecta aut subadscendens glabra, foliis oppositis distinctis subulatis acutis, capitulis inter folia suprema subbracteantia sessilibus solitariis, involucri disco breviorè obtusè 8-dentato, ligulis 5-6 oblongis, achænio glabro. 5 in Mexici prov. Sancti-Ludovici de Potosi decemb. fl. legit cl. Berlandier (pl. exs. n. 1340!). Fruticulus vix semipedalis. Folia 3-4 lin. longa erecta rigidula integerrima. Invol. oblongo-cylindricum subcostatum. Pappus sordidè subgriseus. (v. s.)

† *Species exclusa.*

*BORRERA POROPHYLLUM* Less. = *Lebetina cancellata*.

CCCLXXXVI. *CLOMENOCOMA* Cass. dict. 9. p. 416. v. 59. p. 61 et 66. — Bartolina *Adans. fam.* 2. p. 24 (ex car. et syn. *Houst.*) non Brown.

Capit. multiflorum radiatum, ligulis 1-serialibus fœmineis, fl. disci tubulosi 5-dent. hermaphr. Invol. imbricatum, squamis linearibus acutis dorso versus apicem glandulam grossam gerentibus. Recept. tenuiter fimbriiferum. Achæmium elongatum multistriatum glabriusculum. Pappus squameis 10 uniseriatis basi polyadelphis supernè lineari-setiformibus. — Herbæ Austr.-Americanae Tagetis facie. Folia opposita pinnatisecta. Capitula terminalia pedunculata solitaria. Flores aurantiaci. — Genus recognoscendum. Differt à *Dysodiâ* involucri imbricato nec uniseriali.

1. *C. aurantia* (Cass. dict. 9. p. 416. 59. p. 66), foliis ad basin petioli lobulos setaceos utrinquè gerentibus, segmentis bijugis cum impari ovalibus acutis serratis. — in Verâ-Cruce (Houst.). *Aster aurantius* Linn. sp. 1232. Inula aurantia Spreng. syst. 3. p. 524. — *Aster*, etc. Houst. reliq. t. 18. Caulis dicitur basi radicans.

† *Species dubia.*

2. *C. ? pinnata*, foliorum segmentis multi- (5-7) jugis, omnibus aequalibus linearibus basi integris apice latioribus inciso-dentatis. — in Novâ-Hispaniâ, prope Hapam. *Aster pinnatus* Cav. ic. 3. p. 6. t. 222. Habitus prioris sed pappus videtur setis paucis simplicibus constans. Fortè genericè separanda, sed *Clomenocomæ* certè multò magis affinis quàm *Asteri*. †.

CCCLXXXVII. *HYMENATHERUM* Cass. bull. philom. 1817 jan. et 1818 decemb. dict. 22. p. 313. Less. syn. 236.

Capitulum multiflorum radiatum, ligulis 10 fœmineis ovalibus suprâ velutinis. Invol. turbinatum campanulatum disco brevius squamis

10-14 uniseriatis concretis obtusis grossè glandulosus constans. Recept. planum omnino nudum. Cor. subinæqualiter 5-dentatæ. Styli rami elongati. Achænia gracilia striata. Pappus paleis 10 rarius 5 subuniseriatis infernè indivisis membranaceis supernè trifidis, lobis nunc omnibus rigidulis aristatis scabris, nunc lateralibus membranaceis medio aristato, nunc alternè trifidis aristatis, alternè oblongis membranaceis. — Herbæ annuæ, omnes fortè Chilenses, diffusæ. Caules angulati. Folia saltè inferiora opposita. Capitula ramos terminantia solitaria verisim. lutea.

§. 1. *Pappo 1-seriali, foliis pinnatisectis.*

1. H. ? *NEMER*, foliis infer. oppositis, super. alternis pinnatipartitis, lobis filiformibus integerrimis, pappi paleis 10 basi scariosis trifidis, lobis lateralibus membranaceis, medio elongato aristæformi scabro. (I) in Chili? legit cl. Nees. Herba vix 5-6-pollic. apice ramosa, ramis foliosis 1-cephalis. Invol. campanulatum ebracteatum 12-14-dentatum. Flosculi 70 quorum 10 ligulati. An potius *Dysodia* species? (v. s.)
2. H. *TENUILOBUM*, caule herbaceo duriusculo erecto aut adscendente ramoso, foliis pinnatipartitis subpuberulis, lobis subulatis acutis integerrimis, pedunculis elongatis 1-cephalis subnudis, pappi 1-serialis squamellis 10, omnibus basi submembranaceis apice trifidis, lobo medio setiformi scabro, lateralibus brevibus submembranaceis. (I) in Mexico circa Bejar legit cl. Berlandier (pl. exs. n. 2063!). Etià habeo sine loci indic. ex itin. Nees. H. tenuifolium Cass. dict. 22. p. 313 dictum cum dubio è Chili ortum, huc fortè pertinet. (v. s.)
3. H. *BERLANDIERI*, caule suffruticuloso erecto ramosissimo, foliis confertis puberulis pinnatipartitis, lobis subulatis acutis integerrimis, pedunculis elongatis 1-cephalis squamulas paucas gerentibus, pappo 1-seriali, squamellis alternè 5 ovali-oblongis obtusis membranaceis, 5 duplò longioribus in aristæ scabram productis basi margine scariosis. § in Mexico inter Santander et Vittoria novemb. fl. legit cl. Berlandier (pl. exs. n. 2253!). Similiter H. tenuilobo, sed pappo facillè distinctum. (v. s.)
4. H. *PENTACÆTUM*, caule suffruticuloso erecto ramoso, foliis pinnatipartitis puberulis demùm glabratis, lobis subulatis acutis, infer. brevioribus approximatis, pedunculis elongatis 1-cephalis subnudis, pappi squamellis 5 basi submembranaceis subtrifidis, lobis lateralibus brevibus aut nullis, medio setiformi elongato scabro. § (I) in Mexico circa Monterey legit cl. Berlandier jan. flo. (pl. exs. n. 1382!). Facillè cum H. Berlandieri et tenuilobo primi fructus confundendum, sed pappo 5- nec 10-seto distinctum. (v. s.)

§. 2. *Pappo biseriali, foliis indivisis.*

5. H. ? *KUNTHII* (Less. syn. 236), foliis in vaginam connatis linearibus integerrimis pilosiusculis. (I) in Chili. Herba gracilis nigrinosa pauci-pollicaris simplex aut parùm ramosa. Folia decussata, 6-9 lin. longa, vix semilineam lata. Invol. 6-8-dentata, dentibus ciliatis subæqualibus. Confer cum *Coilopodio* Chilense.

CCCLXXXVIII. *TAGETES* Tourn. inst. 488. t. 278. Linn. gen. n. 964. *Græc. fr. 2. p. 437. t. 172. Less. syn. 236.* — *Tagetes*, *Diglossus* et *Eualcidea* Cass. c. 59. p. 62 et 69-70.

Capitulum radiatum aut rarissimè subdiscoideum. Invol. squamæ 1-seriales concretæ in cupulam campanulatam cylindricamve apice 5-dentatam. Recept. subalveolatum. Achænia elongata basi attenuata compresso-tetragona. Semen in achænio pedicellatum. Pappus simplex paleis inæqualibus, aliis obtusis subconcretis, aliis aristatis elongatis liberis. — Herbæ annuæ Americanæ sæpius fœtidæ rarò suaveolentes.



glandulis in foliorum laminis subrotundis pellucidis et in invol. lineari-  
bus plus minus obsitæ. Folia opposita aut alterna, indivisa serrata aut  
pinnatisecta, segmentis serratis integerrimisve. Capitula solitaria aut  
fasciculata plus minus pedicellata. Cor. crocæ seu flavæ. Ligulæ sæpè  
(præsertim in capitulis plenis) bilabiatae.

§. 1. Pectidiformes, foliis indivisis oppositis serratis, involucrio  
cylindrico.

1. T. LUCIDA (Cav. ic. 3. p. 33. t. 264), caule erecto parè ramoso, foliis lan-  
ceolatis argutè serratis basi aristato-serratis, invol. cylindraceo, ligulis 2,  
pappo 5-mero, paleis 3 obtusis, setis 2 scabris cæteris superantibus. (1) in  
Mexico (Cav.), ad montes circa urbem (Berl. pl. exs. n. 22, 709 et 2132), ad  
Tlapachua (Keerl. in h. Mart.), ad Tamaulipas (Berl. n. 2132). Willd. sp. 3.  
p. 2126. Variat foliis inf. obtusis, super. acutis, angustioribus aut latioribus.  
Nonquàm folia vidi connata ut vult Cavan. (v. v. c. et s. sp.)
2. T. FLORIDA (Sweet brit. fl. gard. ser. 2. t. 35), caule erecto ramoso, foliis lan-  
ceolatis argutè serratis basi aristato-serratis, ligulis sæpius 3, achænio paleis  
membranaceis 2-3 æqualibus aristisque nullis coronato. (1) in Mexico (Sw.).  
Similis T. lucidæ sed ligulæ majores et aristæ pappi nullæ. (v. s. ex hort. Barcl.  
in quo etiàm Sweet descr.)

§. 2. Macrocephalæ, foliis pinnatisectis, capitulis multi- (30 et ultra)  
floris, ligulis subrotundis amplis, invol. ovatis campanulatisve.

3. T. ELLIPTICA (Smith in Rees cycl. v. 35. n. 7), erecta ramosa, foliis pinnati-  
sectis, segmentis 3-4-jugis cum impari ellipticis subintegerrimis mucronatis.  
infer. minoribus, summis gradatim majoribus, eorymbo subpaniculato, capi-  
tulis pedicellatis bracteolis subulatis 1-2 stipatis, invol. levibus obovato-oblu-  
gis, ligulis amplis. — in Peruvia (Domb. 1). Tagetes n. 27. Poepp. pl. exs.  
Peruv. Folia Rutam pinnatam subreferunt. Glandulæ foliorum rotundæ, in-  
volucri lineares. Spec. distinctissima. (v. s. ex coll. Domb. et Poepp.)
4. T. ZYPAQUIRENSIS (Humb. et Bonpl. pl. eq. 2. p. 18. t. 73), caule erecto pani-  
culato, ramis oppositis subfastigiatis, foliis oppositis pinnatisectis, segmentis  
4-6-jugis cum impari oppositis oblongis obtusis serratis subtus pubescentibus,  
rachis puberulâ hinc inde setigerâ, capitulis longè pedunculatis solitariis sub-  
corymbosis, involucrio oblongo-campanulato, ligulis oblongis 10-11 elongatis.  
(1) in Novâ-Granatâ propè urbem Zypaquira. H. B. et Kunth nov. gen. am. 4.  
p. 195. Ligulæ flavæ. Pappi squamæ circ. 5 lineari-subulatae subinæquales.
5. T. PATULA (Linn. sp. 1249), caule erecto, ramis patulis, foliis pinnatisectis,  
segmentis lineari-lanceolatis serratis, serraturis præsertim superiorum aristatis,  
pedunculis 1-cephalis elongatis subcylindricis, involucri levibus. (1) in Mexico  
et nunc in hortis totius fere orbis culta. Eandem aut valdè affinem circa Quito  
leger. cl. Humb. et Bonpl. Sims bot. mag. t. 150. — Dill. elth. 273. t. 279.  
f. 361. Dod. pempt. 255. Hern. mex. p. 156. f. 1 et fortè 2 et 3? Knorr del. 1.  
t. S. 8. Cor. aureo-fuscæ seu fulvæ. Capitula duplò minora et longius peduncu-  
lata quàm T. erectæ, eodem modo variantia. (v. v. c.)
6. T. BARBATA (Linn. sp. 1249), caule ramisque erectis, foliis pinnatisectis, seg-  
mentis lanceolatis serrulatis, serraturis præsertim superiorum aristatis, pedun-  
culis 1-cephalis apice ventricosissimo-increasatis, invol. subangulatis. (1) in Mexico  
sponte et nunc in hortis orbis fere totius culta. Lam. ill. t. 684. Cempoaxochilit  
Hern. mex. 154. T. major Gärtn. fr. 2. p. 434. t. 172. — Knorr del. 1. t. S. 6.  
Cor. unicolores citrino-flavæ. Capit. duplò majus quàm T. patulæ, brevius  
pedunculatum. Variat capitulis proliferis, corollis exter. tantum ligulato-subbi-  
labiatis, aut omnibus ligulato-subbilabiatis, etc. Odor herbæ foetidus. (v. v. c.  
et s. sp. ex Berl. pl. exs. n. 737.)
7. T. CORYMBOSA (Sweet brit. fl. gard. ser. 1. t. 151), caule ramisque erectis sul-

cato-angulatis, foliis oppositis alternisque pinnatisectis, segmentis 6-8-jugis cum impari lineari-lanceolatis acuminatis serratis punctatis, fol. sup. serratis aristatis, rachi inter segmenta hinc inde setigerâ, pedunculis 1-cephalis subnatis corymbosis, invol. oblongo-subturbinato 5-angulato sub-5o-floro, ligulis 20 amplis subrotundis. (1) in Mexico (Sweet), ad Tlapujahua (Kerl. in h. Mart.). Tag. tenuifolia H. B. et Kunth nov. gen. 4. p. 196 non Cav. Ligula supernè fusco-erocæe. Achænia glabra nigra. Pappi paleæ subconcretæ obtusæ, 1 sæpius libera longior acuminata. In hortis occurrit sæpè sub nom. falso T. bonariensis Pers. (v. s. c.)

8. T. PEDUNCULARIS (Lag. et Rod. in annal. scienc. nat. 1802. v. 5. p. 74. Cav. conc. p. 201), caule erecto subdiffuso, ramis foliisque inf. oppositis, sup. alternis, foliis pinnatisectis 5-6-jugis, segmentis lineari-oblongis grossè et acutè serratis, serratis hinc inde setigeris, ramis longè denudatis 1-cephalis, invol. oblongo 5-dentato circ. 3o-floro, ligulis 5 amplis subrotundis. (1) in Mexico circa urbem leger. cl. Alaman, Mairet et Berl. ! pl. exs. n. 1183. T. elongata Willd. sp. 3. p. 2127? Ligulæ flavæ nec fusco maculatæ. Pappi paleæ membranacæe obtusæ, 1 duplò longior libera lineari-subulata. (v. s. sp.)

9. T. PAUCILOBA, glabra erecta, caule basi suffruticoso, foliis oppositis 1-2-jugis, rachi lobisque linearibus acutis integris grossè glandulosis, capitulis nudis solitariis terminalibus pedicellatis, invol. oblongo 5-dentato, ligulis circ. 5 subrotundis. 3 inter Chilenses herb. Hænkeani servatur. Folia  $1\frac{1}{2}$ -2 poll. longa, lobis 6-8 lin. longis, lineam latis, terminali pollicem longo. Invol. 7 lin. longum, 3 lin. latum. (v. s. in h. Hænke.)

10. T. TENUIFOLIA (Cav. ic. 2. p. 54. t. 169), erecta vix ramosa, foliis oppositis alternisve pinnatisectis, segmentis 6-8-jugis cum impari linearibus serratis, fol. summorum serratis aristatis, pedunculis 1-cephalis subnatis paucis subcorymbosis, invol. obovato lævi disco breviorè, ligulis amplis subrotundis. (1) in Peru, ubi vulgò *Chinchimali* dicitur. T. tenuifolia Pers. ench. n. 8 (excl. q. Ort.). Corollæ flavæ concolores immaculatæ. Achænia glabra nigra. Pappi paleis 5, quarum 4 concretæ obtusæ et quinta sublibera longior acuminata. (v. v. c. in h. Madr. et Par.)

§. 3. *Leptocephalæ*, foliis pinnatisectis, rariùs bipinnatisectis, capitulis paucis (5-20) floris, invol. cylindraceis, ligulis parvis aut nullis.

\* *Radiatæ*.

11. T. GRAVEOLENS (L'Hér. ! herb. et mss.), caule erecto subtetragono oppositis ramoso, foliis oppositis pinnatisectis, segmentis 10-11-jugis cum impari elongato-lanceolatis serratis, super. decurrentibus subconfluentibus, infimis parvis in lobos aristatos incisus ad fol. conjunctionem confertis, paniculis laxis subsæcis, capitulis pedicellatis, invol. subtetragono 4-5-dentato, ligulis 4-5 parvis invol. paulò longioribus. (1)? in Peruvia olimplegit Pavon. Herba 3-organis graveolens. Folia ferè Ebuli, segmenta 2 poll. longa, 4 lin. lata. Pappi paleæ inæquales, una subulata longissima. Species ex foliis distinctissima. Flores in herb. desiderantur, sed in mss. L'Hér. descr. (v. s. sine fl.)

12. T. GLANDULIFERA (Schrank pl. rar. h. Mon. 2. t. 54), caule erecto, ramis erectis brevibus, foliis alternis pinnatisectis, segmentis lineari-ambulantibus utrinquè acuminatis serratis 6-8-jugis cum impari, super. subdecurrentibus. inf. cujusque folii parvis pinnatifidis lobis aristatis, capitulis corymboso-fasciculatis ad apicem ramorum plerisque subsessilibus circ. 7-floris, invol. cylindrico, ligulis 3 parvis. (1) in rudatis et fossis ad San-Jago (Bert.) et Valparaíso Chilensium (Gaud.), Bonariâ (Dill.), Brasiliæ merid. Banda orient. (Baclet) et prov. Rio-Grande (h. Mus. imp. Bras. n. 987). T. minuta Jacq. Feuille. obs. 3. p. 64. Dill. elth. 2. p. 374. t. 280. f. 362. T. minuta Linn. f. 1250. T. Bonariensis, Pers. 2. p. 459. n. 5. T. glandulosa Link. ensm. 1 p. 339. Colla mem. ac. Tur. 33. p. 165. app. 3. p. 53. T. montana hort.

Lagd. 1821. Ligulæ non albicantes ut vult Dill. sed pallidæ flavæ. Caulis nunc semipedalis et nomini Linnæano sat conveniens, nunc sæpius 1-2-pedalis. Odor gravis. Segmenta foliorum sesquipollicem fere longa, a lin. lata. Glandulæ in foliis sparsæ grossæ. Ligulæ nunc integræ subrotundæ, nunc apice 2-3-dentatæ. Achænium subpubescens. Pappi palææ membranacæ breves obtusæ, una aristata longa corollâ tamen brevior. (v. s. sp. et v. c.)

13. *T. DAUCIFORMIS* (Schrad.! ex sem. hort. Gœtt. 1833. p. 5. et in Linnæa 1835. litt. p. 71), erecta alternè ramosa, foliis alternis pinnatisectis 10-11-jugis elongato-linearibus lanceolatis grossè et argutè inciso-serratis, lacinialis hinc inde serratis, segm. inf. parvis in dentes longè setigeros fissis, pedunculis axill. terminalibusque sæpius trifidis, pedicellis involucri oblongo 5-dentato brevioribus, flosculis circ. 15-18, ligulis 5 parvis subrotundis. (I) patr. ign. Affinis *T. foetidissima* et glandulifera. Folia parçè glandulosa 3-4 poll. longa; segm. 15 lin. longa, 2-2  $\frac{1}{2}$  lin. lata, rarisimè pinnati lobata. Pappus ligul. palæis a elongatis et cæt. obtusis, disci palæa 1-aristatâ, cæteris obtusis. (v. v. c.)
14. *T. FORTISSIMA* (hortul.), caule erecto paniculato-ramoso, foliis alternis pinnatisectis, segmentis oblongo-lanceolatis serratis 10-14-jugis, summis decurrenti-subconfluentibus, infer. parvis in lobos aristatos incisissimis, capitulis pedicellatis solitariis subcorymbosis circ. 12-flores, invol. ovato cylindraceo, ligulis 5 parvis oblongis. (I) in Mexico (Alam.), ad Guchilacæ Cordilleram (Berl.! pl. exs. n. 1043). Affinis *T. glandulifera* et cum eâ sæpè sub nov. *T. glandulosa* confusa. Herba 1-3-pedalis foetidissima. Segm. fol. 6 lin. longa, a fere lata. Pappi palææ 3 membranacæ breves, 2 aristatæ corollam fere superantes. (v. s. sp. et v. c.)
15. *T. TERNIFLORA* (H. B. et Kunth nov. gen. am. 4. p. 196), caule erecto ramoso sulcato apice corymboso polycephalo, ramis patulis oppositis, foliis oppositis pinnatisectis, segmentis subsexjugis cum impari oppositis lanceolatis argutè serratis, capitulis subternis breviter pedicellatis, invol. clavato tubuloso 9-10-flores 5-dentato. (I) prope Quito ad radices montis Pichinchæ. Pappi palææ 3 minimæ membranacæ, a linearibus subulatis. Ligulæ indescr.
16. *T. MULTIFLORA* (H. B. et Kunth nov. gen. am. 4. p. 197), caule erecto ramoso tereti apice corymboso polycephalo, foliis alternis pinnatisectis, segmentis subquadrijugis subalternis lanceolatis serratis, capitulis fasciculatis longè pedunculatis, invol. tubuloso-subventricosum 14-15-flores, ligulis 4-5 subrotundis. (I) prope Quito in radicibus montis Pichinchæ. Pappi duplicis palææ 5 exteriores lineares subæquales, 5 interiores minimæ obtusæ.
17. *T. MULTISETA*, caule erecto alternè ramoso, foliis alternis bipinnatisectis, lobis linearibus subulatis fere omnibus in setam elongatam productis, pedunculis solitariis longis nudis 1-cephalis, invol. oblongo 5-dentato, flosculis 15-20, ligulis 5 parvis retusis obcordatisve. (I) in Mexico (Alam.! Hanke!), ad Coenavacam (Berl.! pl. exs. n. 969). An *T. subulata* Lallav. et Lex. nov. veg. mex. 1. p. 31 sed descr. non suff. Herba 6-12 poll. Folia hinc glandulosas grossas gerentia, inf. setis destituta. Pedunc. 2-4 pollic. longi. Pappi palææ fere omnes concretæ, a longiores subliberæ. (v. s.)
18. *T. GRACILIS*, erecta apice tantum parçè ramosa, foliis oppositis pinnatisectis, segmentis oblongo-linearibus grossè pauci serratis subalternis, inf. parvis in dentes setigeros incisissimis, pedunculis trifidis corymbosis, pedicellis involucri cylindrico lævi longioribus, flosculis 10-12, ligulis paucis vix invol. longioribus. (I) in Peruvia? Radix tenera. Caulis pedalis teres. Folia distantia 1-1  $\frac{1}{2}$  poll. longa, segmentis glandulosis 4-5-serratis 4 lin. longis, lineam sesquilineamve latis. Pappi palææ 3 breves membranacæ, 2 aristatæ elongatæ. Affinis vid. *T. ternifloræ* et multifloræ. (v. s. comm. à Mus. reg. Par. ex coll. Dombey sine loci propr. design.)
19. *T. OLIGOCEPHALA*, caule erecto simplici aut infernè oppositè, supernè alternè, ramoso, foliis oppositis, pinnatisectis 5-7-jugis cum impari, segmentis linearibus lanceolatis acuminatis serratis, infer. foliis et omnibus fere ad apicem caulis serrato-longè-aristatis, pedunculis paucis elongatis 1-cephalis,

invol. oblongo circ. 20-floro, ligulis nullis aut fortè hinc indè minimis. (I) in Mexici Cordilleris Guachilapensis legit cl. Berlandier (pl. exs. n. 1033 et 1197) Pappi paleæ 3 membranaceæ et 2 elongatæ aristatæ. Folia vix pollicem longa, segm. 4-6 lin. longis, 1-1  $\frac{1}{2}$  latis. (v. s.)

20. *T. pusilla* (H. B. et Kunth nov. gen. am. 4. p. 194). procumbens ramosa, foliis oppositis alternisve bipinnatisectis, lobis linearibus integerrimis, pedunculis solitariis 1-cephalis, invol. tubuloso-clavato, flosc. 12-13, ligulis 1-3 parvis. (I) prope Chillo Quitensium ubi dicitur *Amicillo*, herba enim animæ redolet. Ligulæ albidæ. Pappi paleæ 3 obtusæ membranaceæ breves, 3 aristæformes alternantes. (v. s. sine flor.?)

21. *T. angustifolia* (H. B. et Kunth nov. gen. am. 4. p. 194), caulibus diffusis subdichotomis, foliis alternis pinnatisectis, segmentis linearibus integerrimis, terminali subdentata, capitulis terminalibus solitariis longè pedunculatis circ. 15-floris, invol. clavato-tubuloso. (I) in frigidis montium Mexicanorum prope Toluca, Pappi paleæ 2 longiores lanceolatæ, 3<sup>a</sup> truncata. Ligulæ indescrite, an nullæ?

22. *T. caracasana* (Humb. in Willd. sp. 3. p. 2128), caule erecto sulcato apice corymboso, foliis pinnatisectis, segmentis lanceolatis apice serratis, pedunculis elongatis erectis 1-cephalis alternis, invol. cylindraceis. (I) ad Caracas. Catign. †.

23. *T. dianthifolia* (H. B. et Kunth nov. gen. am. 4. p. 197), caule erecto ramoso striato-sulcato apice corymboso polycephalo, foliis alternis impari-pinnatisectis, segmentis trifugis oppositis lanceolatis subintegerrimis, pedunculis 1-cephalis, invol. ventricoso-tubuloso 7-9-dentato, flosc. circ. 20, ligulis 7 lanceolato-oblongis. (I) in altis Peruviz inter flum. Catamayo et villam Locarque Ligulæ flavæ. Pappi squamæ 3 exter. subulatæ, 3 inter. alternantes minime

24. *T. coronopifolia* (Willd. enum. suppl. p. 60), caule ramisque erectis terribus, foliis oppositis alternisve pinnatisectis, segmentis 6-8-jugis cum impari linearibus integerrimis rarius dentatis, pedunculis solitariis 1-cephalis nudis, invol. oblongo-cylindraceo, ligulis brevissimis subrotundis. (I) in Mexico Jacq. eclog. 1. p. 118. t. 80. *T. clandestina* Lag. eleuch. h. Madr. p. 28. n. 362? Lea in linnæa 1830. p. 160. *Enalcida pilifera* Cass. dict. 14. p. 443. Affinis *T. urticifida*, sed capitula solitaria nec sensim corymbosa. Capit. 5-6 lin. longa, 1-2 lata. Pappi paleæ membranaceæ apice deniatæ concretæ sæpius inæquales, sed nullæ aristatæ. Achænia ad angulos puberula.

25. *T. forniculacea* (Poepp. pl. exs. peruv. n. 26), erecta stricta parcè ramosa, foliis oppositis à basi pinnatisectis, segmentis 6-8-jugis cum impari linearibus subulatæ integerrimis, pedunculis solitariis 1-cephalis nudis, invol. cylindricis, ligulis subsolitariis parvis. (I) in Peruvia legit cl. Poeppig. Herba ferè sesquipedalis glabra omninò media inter *T. micrantham* et *T. multifidam*. Diglossæ variabilis Cass. dict. 13. p. 241 ad hanc (fortè cum aliâ specie mixtam?) pertinere videtur. (v. s.)

26. *T. micrantha* (Cav. ic. 4. p. 31. t. 352), erecta aut diffusa oppositè ramosa, foliis oppositis pinnatisectis, segmentis 1-2-jugis cum impari lineari-subulatæ integerrimis, pedunculis solitariis 1-cephalis apice nudis, involuero cylindrico, ligulis 1-3 parvis obtusissimis. (I) in Mexico circa urbem (Berl. pl. exs. n. 471) et ad Queretaro (Née). Herba semipedalis tenuis glaberrima. Cor. flavæ parvæ. Pappus paleis 2-3 membranaceis truncatis inæqualibus et aristis 2-3 scabro rigidulis longis alternantibus constans. (v. s.)

\* \* *Flosculosa.*

27. *T. multifida*, erecta ramosa, foliis alternis uni- aut bipinnatisectis, segmentis linearibus acutis in setas non productis, capitulis corymbosis breviter pedunculatis, invol. oblongo subtruncato levi, flosculis 20-25 omnibus tubulosis. (I) in Mexico ad Lerma (Berl. pl. exs. n. 1143). Variat segmentis 4-6-jugis integerrimis dentatis aut pinnatifidis. Pappi paleæ 3 aut 4 membranaceæ nuncate dentatæ. 2 seu 1 liberæ lineares elongatæ non verè aristatæ. An I 3:

*folia* Lag. *elench.* 25, sed *ligulæ* dicuntur minutæ *lonulatæ* dùm in *meâ nullæ*? (v. s.)

28. *T. FLOSCULOSA* (Spreng. *synt.* 3. p. 571), caule...., foliis *pinnatifidis filiformibus elongatis integerrimis summis indivisis*, pedunculis *terminalibus solitariis*, invol. *turbinato*, radio *obsoleto*. — in *Monte-Video*. Cst. ign. †.

† *Species non satis notæ.*

29. *T. LUNULATA* (Ort. *dec.* p. 44), caule *erecto subangulato*, ramis *patulis*, inf. *oppositis*, sup. *subcorymbosis*, foliis *pinnatisectis*, segmentis *alternis oppositisque lineari-lanceolatis obtusè serratis*; intimis *minoribus eciliatis*, pedunculis *longissimis 1-cephalis*, ligulis 5 *obcordatis*. (†) in *Mexico*. *Ligulæ* *maculâ sanguineâ lunatâ maculatæ*. Pappi *paleæ* 2 *lineari-subulatæ*, reliquæ *breves truncatæ*. Numerus *florum*, eorum *proportio*, et *forma invol.* *ignorantur*.

30. *T. SUBVILLOSA* (Lag. *elench.* h. *Madr.* p. 29 n. 363), caule...., foliis *pinnatisectis villosulis*, segmentis *lanceolatis serratis*, capitulis *laxè corymbosis*. (†) in *Mexico*. Cst. ign. †.

31. *T. VERTICILLATA* (Lag. et Rod. in *annal. scienc. nat.* 1802. v. 5. p. 73), caule *erecto ramosissimo*, foliis *ramisque verticillatis ternis*, capitulis *corymbosis*, ligulis *brevissimis*. — ad *Cumanam*.

Quid *T. ozolis* Spreng. à *Steud.* citata?

†† *Species exclusæ.*

*T. FRUILLÆ* Bert. = *Lasthenia Chilensis*.

*T. PAPPOSA* Vent. = *Dysodia chrysanthemifolia*.

*T. FUMILA* hortul. = *Dysodia chrysanthemifolia*.

*T. PUNCTATA* fl. *mex.* = *Adenophyllum coccineum*.

CCCLXXXIX. THYMOPHYLLA Lag. *nov. gen. in elench. h. Madr.* p. 25.

Capitulum *multiflorum homogamum discoideum*. Invol. *campanulati apice dentati squamæ concretæ*. Recept. *nudum*. Pappus *paleis 5 truncatis brevibus 1-serialibus*. — Suffrutex *Mexicanus*, ramis *oppositis*. Folia *opposita conferta subsetacea parva (vix sesquilineam longa) tomentosa*. Pedunculi *terminales 1-cephali 1-2-phylli villosuli*. Invol. *extus pulverulento-tomentosum achæniis æquale*. Cor. *ex sicco atropurpureæ*. — Genus mihi *ignotum ad Tagetineas à cl. Cassini et Lessing relatum*, sed *tomentum fol.* obstat et de *glandulis silet cl. Lagasca qui solus plantam vidit*.

1. *T. SETIFOLIA* (Lag. l. c. n. 325). — in *Mexico*.

Div. II. POROPHYLLÆ. — Tagetinearum et Pectidearum pars. Cass. *opusc.* 3. p. 55.

Capitulum *semper homogamum discoideum*. Invol. *squamæ liberæ aut vix basi concretæ*. Cor. *disci inæqualiter fissæ subbilabiatae*. Styli *rami in appendicem subulatam producti*. Pappus *setosus pluriserialis*.

CCCXC. POROPHYLLUM Vaill. *act. acad. Par.* 1719. p. 407. *Linn. hort. cliff.* 494. *Adans. fam.* 2. p. 122. *Cass. dict.* 43. p. 56. — *Kleinia Jacq. amer.* 215. *Willd. spec.* 3. p. 1738. *Less. syn.* 196 non *Juss. nec Haw.* (1). — *Calam sp. Linn.*

Capitulum *multiflorum homogamum*. Recept. *nudum punctatum*. Invol. *1-seriale*, squamis 5- (in spuriis 9-12) *oblongo-linearibus*, ju-

(1) Ex juribus collectis *prioritatis*, *facilitatis*, *etymologiæ*, nomen *Vaillantii* htc

nioribus basi subconcretis. Cor. tubulosæ regulariter 5-dentatæ sub  
dente unico profundius diviso sublabiatæ. Styli rami exserti, basi intus  
series stigmaticas 2 gerentibus, apice undiquè (ferè ut in Vernonia-  
ceis) hispidi acuti. Achænia elongata angulata sæpius scabra, apice  
plùs minùs in rostrum attenuata. Pappus pluriserialis pilosus scaber.  
— Herbæ aut suffrutices. Species genuinæ omnes glabræ Americanæ.  
Folia alterna aut rarius opposita, sæpius glauca, ovata aut oblonga,  
integra aut dentata, sæpius glandulis pellucidis amplis oblongis no-  
tata. Pedicelli elongati x-cephali. Invol. sæpè cæruleo-purpureo col-  
orata et glaucescentia. Cor. sordidè purpureæ. — Affinitas subdubia;  
an Senecioneis, an Vernoniaceis propinqua?

## Sect. I. EUPOROPHYLLUM.

Invol. squamæ 5. — Folia sæpè poris glandulosis notata. — Species Ame-  
ricanæ.

§. 1. *Capitulis cylindraceis elongatis, invol. squamis mucrone cal-  
loso apiculatis, cor. regulariter 5-dentatis, achænio evidentiùs  
rostrato.*

1. P. MACROCEPHALUM, herbaceum erectum, foliis longè petiolatis ellipticis ob-  
tusis crenatis ad sinus pellucido-glandulosis, limbis petiolo brevioribus, pe-  
dunculis inflato-turbinatis, invol. squamis mucrone calloso apiculatis. (†) in  
Mexico circa Villalpando legit cl. Mendez. *Tzinoma* Hern. mex. 434. f. 3.  
Kleinia glandulosa flor. mex. icon. ined. nimis fortè P. elliptico affine, differt  
petiolis longioribus, foliis parvis 6 lin. longis, 4 lin. latis, pedunculis cranio-  
ribus, invol. pollicem longis distinctius glandulosis. Vulgò in Mexico dicitur  
*Tzinoma* ex Hern., *Popoloquolite* ex Moq. (v. s.)

2. P. ELLIPTICUM (Cass. dict. 43. p. 46), herbaceum erectum, foliis petiolatis  
ellipticis oblongisve obtusis mucronatisve crenatis ad sinus aut in disco pel-  
lucido-glandulosis, invol. squamis mucrone calloso acuminatis. (†) in Peruvia  
(Cav.!), ad Turbaco Novo-Granatensium (H. et B.), in Brasiliâ (Cham.!), ad  
Bahiam (Salzm.!), in Santo-Domingo (Menz. in h. Lindl.!), Martinicâ (Sw.),  
in aridis et argillosis. *Cacalia Porophyllum* Linn. sp. 1169. (excl. syn.) Cav.  
ic. 3. p. 11. t. 222. Kleinia *Porophyllum* Willd. sp. 3. p. 1738. — Pluk. sin.  
t. 161. f. 1. mal. Variat achæniis in diversis aut iisdem individuis hirtella,  
puberis aut glabris! (v. v. et s.)

Æ. *intermedium*, foliis ellipticis oblongisve crenatis opacè glandulosis aut  
subglandulosis. (†) in Brasiliâ (Gaudich.!), ad Bahiam (Blanch.!), in Serri-  
dos Orgaos prov. Rio-de-Janeiro (Vanth.!) pl. exs. n. 305). Valde accessit  
ad P. ruderale, sed huc potius referendum videtur. Variat ut var. a achæ-  
niis hirtis et lævibus. An species propria? (v. s.)

3. P. RUDERALE (Cass. dict. 43. p. 56), herbaceum erectum, foliis petiolatis  
elliptico- aut lanceolato-oblongis acuminatis subintegerrimis impunctatis. (†)  
in ruderatis et cultis insularum Caribæarum, Jamicæ (Sw.!), Sancti-Domingi  
(Jacq.), Martinicæ (Jacq.), Guadalupæ (Perr.!), Porto-Ricci (Wyd.!), Cuba  
(La Sagr.!). *Cacalia ruderalis* Sw. prod. 110. Kleinia *ruderalis* Jacq. am. 315  
t. 127. (v. s.)

4. P. VIRIDIFLORUM, fruticosum erectum, foliis breviter petiolatis obovatis sub-  
undulato-repandis aut sinuatis integerrimis, ad margines glandulas pellucidas  
paucas gerentibus. § locis temperatis Mexici juxta Valladolid (H. B. K.)  
Cuernavaca (Berl.!) pl. exs. n. 936). Kleinia *viridiflora* H. B. et Kunth non

admissi, nam Kleinia nomen generi in quo *Cacalia* Kleinia Linn. militabat, evidentè  
præcæntibus Haworthio et ipso Linnæo servandum erat.

KLEINIA Juss. = *Jaumea*.

- gen. am. 4. p. 157. Frutex orgyalis affinis P. elliptico. Cor. virescentes. Invol. squamæ obtusiusculæ glandulis biseriatis notatæ. (v. s.)
5. *P. NUMMULARIUM*, herbaceum erectum, ramis foliisque oppositis, foliis petiolatis elliptico-orbiculatis obtusissimis subcrenatis ad sinus opacè glandulosis, invol. squamis mucrone callosos apiculatis dorso glandulosis. — in Mexici umbrosis ad Tehuantepec et loc. vicin. Herba foetida angusto florens (Alam.). Rami primà fronte videntur folia pinnata! Invol. 4-5 lin. longa. Limbus folii 8 lin. longus, 6 lin. latus. Petioli 4-lin. (v. s. comm. à cl. Alaman.)
- β. foliis obovatis. — (v. s. in h. Hænke.)
6. *P. JORULLENSIS* (Cass. dict. 43. p. 57), fruticosum ramossissimum, foliis longè petiolatis oblongis obtusis subrepandis eglandulosis, invol. squamis lanceolato-linearibus acutis. § in Mexicani montis ignivomi Jorullo declivitate orientali. Kleinia Jorullensis H. B. et Kunth nov. gen. am. 4. p. 156. t. 356.
7. *P. LANCEOLATUM*, glabrum strictum, caule imà basi suffruticuloso, foliis alternis lanceolatis basi attenuatis subacutis subintegerrimis eglandulosis, ramis erectis apice nudis 1-cephalis, achænio scabro. ¶ § in Brasiliæ prov. Rio-Grande: Affine P. prenanthoidi, sed folia alterna, pedunculi strictè erecti, achænia scabra. (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 936. miss.)
8. *P. PRENANTHOIDES*, glabrum erectum, caule virgato basi suffruticuloso, foliis oblongo-lanceolatis utrinquè attenuatis subintegerrimis impunctatis, infer. oppositis, super. alternis, paniculâ laxâ, capitulis nutantibus oblongis. § in prov. Brasiliæ Sancti-Pauli. Habitus ferè Prenanthis. Achænia scabriuscula. An capitula exsiccatione aut naturâ propriâ nutant? (v. s. in h. Mus. reg. Par. à Mus. imp. Bras. sub n. 402. miss.)
9. *P. HÆNKEI*, fruticosum erectum, ramis foliisque oppositis, foliis lineari-lanceolatis utrinquè attenuatis eglandulosis, capitulis breviter pedicellatis, achæniis scabris. § in Cordilleris Chilensibus ex herb. Hænk. Affine P. lineari, sed folia latiora omnia opposita, pedunculi breviores, capitula paulò minorâ. An P. oppositifolium Spreng., sed patria diversa et folia non breviora dicenda cum 15 lin. longa, 2-3-lata? (v. s. in Hænk. à cl. Sternb. miss.)
10. *P. LINEARE*, fruticosum erectum glabrum, foliis alternis oppositisque linearibus integerrimis acutis eglandulosis, invol. oblongi squamis apice breviter mucronatis, achæniis scabris. § in Brasiliæ prov. Rio-Grande. Folia sesquipedalica. longa, lineam lata. Pedunculi post anthesin incrassati striati subtruncati. Capit. 6 lin. longum. (v. s. in h. Mus. Par. à Mus. imp. Bras. sub n. 933 et fortè 934.)
11. *P. LINIFOLIUM*, suffruticosum glabrum multicaule subcaespitosum, foliis alternis linearibus integerrimis subacutis eglandulosis, invol. ovali. § in Brasiliâ (Ard.), ad Monte-Video (Comm.), in prov. Rio-Grande (h. Mus. Bras.), fortè in Peruvîâ et Mexico (h. Hænk.). Cacalia linifolia Linn. mant. 109. Kleinia suffruticosa Willd. sp. 3. p. 1739 non Lodd. Cacalia Linaria var. β pumila Poir. suppl. 2. p. 6. — Lam. ill. t. 673. f. 4. Valdè accedit ad P. Linariam, sed multò minus et magis caespitosum. Folia 4-6 lin. longa. Achænia glabriuscula. (v. s. in h. Mus. reg. Par. ex. Mus. imp. Bras. sub n. 924, 935 et fortè 940 miss.)
- §. 2. *Capitulis ovali-oblongis brevioribus, invol. squamis obtusis margine scariosis, corollis lobo uno profundiore sublabiatis (an semper?), achænio apice non sensim attenuato.* — *Hunteria fl. mex. icon. non Roxb.*
12. *P. LINARIA*, suffruticosum glabrum erectum, ramis virgatis, foliis angustè linearibus acutis integerrimis glandularum pellucidarum serie duplici notatis, invol. squamis obtusis coloratis. § in Mexico (Cav.) Cacalia Linaria Cav. ic. 3. p. 29. t. 257 non Poir.? Hunteria columbina icon fl. mex. Invol. ovali-

oblongum 5 lin. longum, 4 lin. latum, colore fusco-cærulescente subumbili decorum. Achaenium suberosum. (v. s. ex hort. Mex. commun. à cl. Bert. pl. exs. n. 881.)

13. *P. TAGETOIDES*, herbaceum? erectum, foliis breviter petiolatis lanceolatis linearibus acutis subintegerrimis margine pellucido-punctatis subtus glauciscentibus, invol. squamis ovato-oblongis obtusis externè pruinoso-glaucescentibus. (1) in hortis Mexicanis leger. cl. Humb. et Bonpl. *Kleinia tagetoides* E. B. et Kunth. nov. gen. 4. p. 158. Valdè affine dicitur *P. linifolia*.
14. *P. DECUMBENS*, suffruticosum multicaule decumbens subcæspitosum, folia lineari-oblongis basi attenuatis obtusiusculis integerrimis glaucissimis apice glanduloso-callois cæterum eglandulosi, invol. squamis vix acutis. 2/5 in Mexico (Lodd.). *Kleinia suffruticosa* Lodd. bot. cab. t. 1561 non Willd. *Kleinia linearis* hort. Par. 1817! Capitula ferè *P. Linariae*, folia *P. colorati* vel obtusiora, caules *P. linifolii*. (v. v. c. et s.)
15. *P. COLORATUM*, herbaceum erectum, ramis virgatis, foliis lineari-lanceolatis subintegerrimis margine hinc inde pellucido-punctatis, invol. squamis obtusiusculis coloratis. (1) in Mexico circa Guanajuato et Temascatio (H. et B.) ad Tlapujahu (Keerl). *Kleinia colorata* H. B. et Kunth nov. gen. an. 4. p. 158. Mart. amoen. monac. p. 23. t. 15. *Hunteria fuscata* icon. flor. am. Invol. fusco-seu purpureo-cærulescens. Folia pollicem longa, sequilinem lata.
16. *P. OBTUSIFOLIUM*, suffruticosum multicaule, caulis erectis striatis, folia oblongis obtusis basi attenuatis subintegerrimis glandulas pellucidas parvas ad margines gerentibus, nervo medio albo prominulo, invol. squamis obtusiuscoloratis. 3/5 in Mexico propè Villalpando legit cl. Mendez. Stirps circ. pedalis. Folia sup. ferè linearia subacuta. Invol. ovalia pulchrè cæruleo-purpureascentia. (v. s.)

#### Sect. II. CUSIMBUA.

Invol. squamæ 9-12. — Folia poris glandulosis destituta (saltè in desomissa). — Species Indicae. — An genus proprium?

17. *P. CUSIMBUA*, basi fruticosum erectum, caulis sulcatis apice panicatis, foliis semiamplexicaulibus oblongo-lanceolatis longè acuminatis argutè serratis basi integerrimis, pedicellis squamulosis capituli longitudine, invol. cylindrici squamis plurimis (ultra 9) linearibus. 3/5 in Nepaliâ ubi Nawaris dicitur *Cusimba-Swa*. *Cacalia Cusimbua* Don prod. fl. nep. 179. *Kleinia Cusimba* Less. in Linnæ 1831. p. 133. Folia ex Don subtus pubescentia, ex Less. pilidiora. Ovaria glaberrima.
18. *P. CACALIOIDES*, herbaceum erectum, foliis petiolulatis carnosulis oblongo-ovatis acuminatis retrorsum mucronulato-denticulatis, capitulis longè pedunculatis racemoso-subcorymbosis, invol. 9-phyllo cylindraceo, squamis linearibus. — in insulâ Luzoniâ. *Kleinia cacalioides* Less. in Linnæ 1831. p. 133. Cor. apice purpurea? limbo à tubo non distincto. Stylus basi bulbosus. Affine dicitur *P. hieracioidi*.
19. *P. HIERACIOIDES*, herbaceum erectum simplex, foliis petiolatis ovatis dentatis, capitulis subcorymbosis breviter pedunculatis, invol. squamis 12 lanceolatis. (1) in Indiâ orientali. *Cacalia hieracioides* Willd. sp. 3. p. 1731? *Kleinia hieracioides* Less. in Linn. 1831. p. 133. Stirps Lessingiana videtur diversa Willdenowianâ. Folia ex Willd. obtusa, ex Less. longè acuminata. †

#### Sect. III. KUGAIA.

Invol. basi calyculatum. — Folia pinnatifida. — Species Japonica. — An genus proprium?

20. *P. JAPONICUM*, herbaceum, caule debili carnosio villosio, foliis petiolatis incom-pinnatifidis glabris, lobis oblongis acutis dentatis, capitulis paniculatis, pedicellis subsquamosis, involucri calyculato. 2/5 in Japoniâ circa Nagasaki ubi



vulgò *Kwugai Sai* dicitur. *Senecio Japonicus* Thunb. fl. jap 315. Willd. sp. 3. p. 1978. *Kleinia Japonica* Less. in linnæa 1831. p. 134. syn. 133.

† *Species vix notæ et verisimiliter à genere excludendæ.*

21. P. ? *OPPOSITIFOLIUM* (Spreng. syst. 3. p. 438, sub Kleinia), suffrutescens, caule ramisque filiformibus, foliis oppositis sessilibus linearibus brevibus involucrique pubescenti-pulverulentis, capitulis solitariis pedicellatis. ♀ ad Buenos-Ayres. *Cacalia oppositifolia* Poir. ! suppl. 2. p. 7. (v. s. sine fl.)
22. P. ? *SALLOI* (Spreng. l. c. sub Kleinia), fruticosum simplicissimum, foliis decussato-oppositis sessilibus lanceolatis obtusiusculis crenato-serratis eglandulosis hirsutulis, pedunculis racemosis terminalibus. ♀ ad Rio-Grande Brasiliæ. Cæt. ign. †.
23. P. ? *FILIFOLIUM*, (Spreng. l. c. sub Kleinia), herbaceum ramosum, foliis fasciculatis lineari-filiformibus glabris, pedunculis 1-cephalis terminalibus. — in Monte Video. Cæt. ign. An eadem ac P. linifolium ? †.
24. P. *OSCURUM* (Spreng. l. c. sub Kleinia), herbaceum elatum supernè divinum glaucum, foliis alternis remotis linearibus glabris, pedunculis axillaribus terminalibusque 1-cephalis. — ad Rio-Grande Brasiliæ. Cæt. ign. †.

Subtrib. V. *HELENIÆ* Cass. dict. sc. nat. 1829. p. 13. opusc. 3. p. 36. Less. syn. 237

Capitula sæpius heterogama, fl. radii ligulatis (raro tubulosis ampliatis) fœmineis neutrisve 1-serialibus, reliquis hermaphr., rarius homogama, nunquam dioica. Antherae sæpè nigricantes, basi subproductæ nec caudatæ, articulo antherifero tereti. Styli rami nunc apice truncati, nunc cono imò appendice supra pilos collectores primarios gerente aucti. Achæninum erostre. Pappus multipaleaceus, squamellis planis distinctis 1-serialibus scariosis obtusis seu acuminatis, rarissimè aut abortu nullis, aut in vaginulam concretis. Invol. squamæ plurimæ liberæ. — Folia sæpius alterna interdum opposita. Species pleræque Americanæ.

Div. 1. *GAILLARDIÆ*. — *Galardieæ* Nutt. gen. am. 2. p. 177. — *Galardieæ* et *Rosilleæ* Less. syn. 237 et 245.

Pappi paleæ membranaceæ sæpiissimè integræ. Recept. epaleaceum nudum aut alveolatum fimbriiferumve. — Species notæ omnes Americanæ.

Subdiv. 1. *EUGAILLARDIÆ*.

Ligulæ neutrae.

CCCXCI. *GAILLARDIA* Foug. mem. acad. sc. Par. 1786. p. 1 et 6 (sub *Gail-lardâ*). Cass. dict. 18. p. 17. 55. p. 264. — *Galardia* Lam. dict. 2. p. 285. illustr. t. 708. Less. syn. 237. — *Calonnea* Buchoz icon. 1786 t. 126. — *Virgilia* L'Hér. diss. cum ic. non Lam. — *Galordia* Rœusch. nom. 251.

Capitulum multiflorum radiatum, ligulis neutris multinerviis glandulosus apice palmato-trifidis, fl. disci hermaphroditi, tubo brevi, limbo pilis articulatis hispido 5-fido. Recept. timbrilliferum convexum, invol. squamæ 2-3-seriales, appendice foliaceâ acuminatâ disco longiore. Styli rami appendice longâ subulatâ hispidâ terminati. Achænia oblonga villosa. Pappi paleæ uninerviæ in aristam longè acuminatæ. — *Herbæ Boreali-Americanæ* erectæ subpilosæ. Folia alterna, super. integra sessilia aut semiamplexicaulia. Rami longè denuati 1-cephali. Capituli discus fuscus seu flavus, ligulæ flavæ seu basi crocæ.

§. 1. *Annua, invol. squamis interioribus abruptè in appendicem lineari-subulatam angustatis.*

1. *G. DRUMMONDII*, annua subpubera, foliis angustè lanceolatis indivisis ac incis, invol. squamis basi brevissimè glabrè appendice subulatà scabro-ciliat, corollis hirsutis pappo longioribus, pappi paleis basi latè dilatatis, ligula basi cuneatis involucri paulò longioribus. (1) in campis Mexicanæ prov. Tera ad San-Fernando de Bexar (Berl. pl. exs. n. 2027 et 2067), et ad Rio-Braz (Drum.). *Gaillardia bicolor* var.  $\beta$  *Drummondii* Hook. bot. mag. t. 266. Ligulæ 5 lin. longæ, purpureæ, vix apice flavæ. Recept. fimbrillæ densè rigidæ. Folia alia indivisa, alia pinnatilobata, parçè pubera. (v. s. sp.)
2. *G. PULCHELLA* (Foug. l. c. Cass. dict. 18. p. 17), annua subhirsuta, invol. squamis basi glabriusculis appendice abruptè angustatà rigidè ciliat, corollis disci parçè villosis pappi longitudine, pappi paleis basi angustis rigidulis, ligulis basi valdè angustatis maculatis. (1) in Louisianà olim legit com. de sales et in hortos Gallicos introduxit anno 1786 sed anno (ni fallor) 1791 semina omnia ad maturitatem non pervenerunt et species in hortis amissa fuit. *Gaillardia bicolor* Lam. dict. 2. p. 585. ill. t. 708. *Calounea pulcherrima* Lechoz icon. t. 126. *Virgilia helioides* L'Hér. diss. cum. icon. 2. Garten mag. 1810 p. 42. t. 6. *G. amara* Raf. fl. lud. p. 69? *G. alternifolia* Rausch. l. c. Ligulæ 8-9 lin. longæ, circ. 3 lin. latæ. (v. s. ex h. L'Hér.)

§. 2. *Perennes, invol. squamis lineari-lanceolatis appendice nuda basi distinctà.*

3. *G. LANCEOLATA* (Michx. fl. bor. am. 2. p. 142), perennis, invol. squamis extus villosis-subhirsutis lineari-lanceolatis, corollis disci hirsutis pappi circ. longitudine, pappi paleis basi subdilatatis, ligulis obovato-cuneatis basi maculatis. 2/ à Carolinà ad Floridam in aridis (Michx.), in pinetis siccis Georgiæ mediæ (Ell.), ad fl. Saskatchewan et in montibus scopulosis (Hook. fl. bor. am. t. p. 315). *Gaillardia rustica* Cass. dict. 18. p. 20. *Gaillardia bicolor* Ell. sketch 2. p. 449. Sims bot. mag. t. 1602. (excl. syn. ferè omn.) Ligula 7 lin. longæ, 5 lin. latæ, basi crocæ, cæterum flavæ. Cor. disci crocæ. (v. v. c.)
4. *G. ARISTATA* (Pursh fl. bor. am. 2. p. 573), perennis, invol. basi hirsutis squamis lineari-lanceolatis, cor. disci hirsutissimis pappo longioribus, pappi paleis basi latè dilatato-membranaceis, ligulis basi longè cuneatis unicoloribus. 2/ in collibus siccis ad Rocky-Mountains (Pursh), in collibus herbosis ad fl. Missouri (Nutt.). Lindl. bot. reg. t. 1186. Hook. bot. mag. t. 2940. fl. bor. t. p. 315. *G. bicolor* var.  $\beta$ . Nutt. gen. am. p. 175. Magis quàm priores hirsuta. Ligulæ 12-14 lin. longæ, 5 lin. latæ, flavo-subaurantiæ non maculatæ. Cor. disci fusco-purpureæ, lobis elongatis. Achæmium rufo-villosissimum. Pappus albus, hanc et 2 antecedentes sub nom. *G. bicoloris* confudit Spreng. syst. 1. p. 618. excl. syn. Walt. (v. v. c.)

† *Species exclusæ.*

*G. ACAULIS* Pursh = *Actinella acaulis*.

*G. FIMBRIATA* Michx. = *Leptopoda puberula*.

CCCXCH. *BALDUINA* Nutt. gen. am. 2. p. 175. Ell. sketch 2. p. 447. lam. syn. 238.

Capitulum multiflorum radiatum, ligulis neutris apice tridentatis aut trifidis, fl. disci hermaphrod., cor. tubulosis 4-5-dentatis extus glanduliferis basi corneis. Invol. imbricatum campanulatum. Recept. convexum alveolato-fimbrilliferum. Antheræ basi breviter bisectæ. Styli rami inclusi vix appendiculati. Achænia in foveolis immersa se-

ricea obconica. Pappi paleæ membranaceæ breves. Herbæ Boreali-Americanae perennes glabriusculæ. Caules teretes simplices aut ramosi, ramis apice nudis 1-cephalis. Folia alterna sessilia integra. Cor. flavæ. — Affinitas ex antheris bisetosis dubia!

Sect. I. BALDUINA *Ell. in obs. 2. p. 447.*

Pappus achænium æquans biserialis paleaceus, paleis inter. acutis, exter. sub-fimbriatis. Invol. pluriserialis.

1. *B. UNIFLORA* (Nutt. et Ell. ! ll. cc.), caule simplici 1-cephalo pubescente, foliis angustè obovatis.  $\mathcal{U}$  in humidis et paludum marginibus Carolinae et Georgiae. (v. s. comm. à cl. Elliott.)

Sect. II. ACTINOSPERMUM *Ell. sketch p. 448. in observ.*

Pappus brevis 12-14-paleaceus, paleis in copulam patentem dispositis obtusis. Invol. biserialis.

2. *B. MULTIFLORA* (Nutt. et Ell. ll. cc.), caule ramoso glaberrimo polycephalo, foliis linearibus ferè setaceis.  $\mathcal{U}$  in collibus arenosis secus Altamaha ad acidulam Barrington.

CCCXCIII. LEPTOPODA *Nutt. gen. am. 2. p. 174. Ell. sketch 2. p. 445. Cass. dict. 26. p. 79. 55. p. 264 et 272. Less. syn. 238.*

Capitulum multiflorum, fl. radii circ. 20 ligulatis trifidis neutris, fl. disci hermaphr. tubulosi 3-5-dentatis. Invol. biserialis, squamis exter. linearibus subulatis, int. sublancoatis. Recept. convexum nudum. Styli rami breves exappendiculati. Achænia subcylindracea. Pappus paleis membranaceis circ. 8 brevibus fimbriatis in aristam non productis. — Herbæ Boreali-Americanae erectæ simplices 1-cephalæ. Folia alterna linearibus lanceolata subdentata. Capitula terminalia solitaria. Flores flavi. — Habitus Helenii sed ligulæ neutrae.

1. *L. PUBERULA* (Macbr. in Ell. sketch 2. p. 445), caule viscido-pubescente fistuloso stricto, foliis semiamplexicaulibus glabris punctatis, caulinis inciso-dentatis, achæniis villosis subtruncatis.  $\mathcal{U}$  in paludosis apricis Carolinae et Floridae. Galaridia fimbriata Michx. fl. bor. am. 2. p. 142. Helenium vernale Walt. fl. car. p. 210.

2. *L. HELENIUM* (Nutt. gen. am. 2. p. 174), caule glaberrimo solido, foliis decurrentibus glabris denticulatis, achæniis glabris cylindricis.  $\mathcal{U}$  in humidis pinetis Carolinae et Georgiae. Lept. helenioides Cass. dict. 26. p. 80. Lept. decurrens Macbr. in Ell. sketch 2. p. 446.

Subdiv. 2. EUHELENIÆ.

Ligula foemineæ seu nulla.

CCCXCIV. GUTIERREZIA *Lag. olench. h. madr. p. 30.*

Capitulum pluri- (8) florum radiatum, ligulis 3 foemineis. Invol. imbricatum, squamis apice reflexis. Recept. favosum dentato-paleaceum. Pappus polyphyllus paleaceus. — Planta Mexicana suffruticosa glabra resinosa. Folia sparsa sessilia linearia acuta integerrima. Corymbus terminalis oligocephalus. Cæt. ign. — Genus mihi ignot. et dubium. A Galinsoga, quæcum junxit Sprengel, differt recept. favoso nec ut videtur ex descr. Lag. verè paleaceo, invol. imbricato nec n-seriali, etc. †.

1. *G. LINEARIFOLIA* (Lag. l. c.) § 2 in Mexico. *Galinsogea linearifolia* Spreng. syst. 3. p. 579. Folia pollicem et amplius longa, vix lineam lata, sup. breviora.

CCCXCV. *ACHYROPAPPUS* H. B. et Kunth nov. gen. am. 4. p. 257. t. 39 nec Bieb. — *Achyropappi* sect. Less. syn. 239. — *Chamæstephanum* Willd. in Berl. mag. 1807. p. 140. ex Kunth.

Capitulum multiflorum radiatum, fl. radii 3-5 fœmineis. Invol. squamæ quinque 1-seriales ovals obtusissimæ. Recept. parvum nudum alveolatum. Cor. disci 5-fidæ vix pappo longiores. Styli rami breviter appendiculati. Achænia lineari-cuneata 3-4-gona. Pappus paleis 6-membranaceis obovato-muticis. — Herbæ Mexicanæ annuæ vix subpuberulæ erectæ. Folia opposita et super. alterna, multipartita, lobipinnatipartitis oblongis seu linearibus obtusis. Capitula longè pedicellata laxissimè corymbosa. Cor. flavæ (nec radio albo ut sphalmate legitur in opere Kunth). Genus vix à Schkuhriâ, rectè monente cl. Nees del. sem. Bonn. 1831, satis diversum.

1. *A. ANTHEMOIDES* (H. B. et Kunth 4. p. 259. t. 390), caule folisque pubescentibus, fol. lacinii linearisublancoatis, ligulis 5. (1) in Mexici agris frigidis circa Toluccam ad pagum Islahuaca. (v. s. in h. acad. reg. Monac.)  
2. *A. SCHKUHRIOIDES* (Link et Otto abb. neu. gew. gart. Berl. n. et t. 30), caule folisque subsæbris, fol. lacinii linearibus, ligulis 1-3. (1) in Mexico (Link ad occid. urbis Guanaxuato (Mendes l.). *Schkuhria senecioides* Nees del. ex h. Bonn. 1831. An à priore satis differt? (v. v. c. et s. sp.)

† *Species exclusa.*

*A. PEDATUS* Less. = *Florestina pedata*.

CCCXCVI. *SCHKUHRIA* Roth cat. bot. 1. (1797), p. 116. Cass. dic. 1. p. 87. 55. p. 263. Less. syn. 239. non Manch. — *Tetracarpum* Manch. sp. (1802), p. 241. — *Pectidis* sp. Lam. Ort. — *Mieria* Lallier. et Lex. nov. mex. 2. p. 9. nec Link.

Capitulum pauciflorum, ligulis 1-2 fœmineis, fl. disci 5-8 tubulosi 4-5-dentatis brevibus. Invol. squamæ circ. 5 uniserialis obovate submembranaceo coloratæ. Recept. parvum nudum. Styli rami breviter appendiculati. Achænia tetragona basi attenuata. Pappi paleis 8, in angulis achæntii ortis acuminato-aristatis, 4 alternis nunc priores revertentibus nunc iis brevioribus et obtusioribus margine subfimbriatis. — Herbæ Americanæ annuæ ramosissimæ glabræ. Folia alterna tripinnatisecta, segmentis linearibus tenuissimis integerrimis. Capitula ovata solitaria longè pedicellata. Cor. luteæ.

1. *S. OCTOARISTATA*, pappi paleis 8 æqualibus acuminato-aristatis corollis brevioribus, ligulâ involucri longiore. (1) in Peruvîa legit cl. Dombey quænti simillima et cum eâ in herb. interdum confusa, sed pappo (asus diversus. (v. s.)  
2. *S. ABROTANOIDES* (Roth cat. 1. p. 116), pappi paleis 4 acuminato-aristatis 4 alternis brevioribus obtusatis omnibus corollis disci brevioribus, ligulâ involucri longiore. (1) in Peruvîa (Dombey!), ad Huanacum urbem (Ort. in Mexico. — Schkuhr. handb. 3. t. 250. b. *Pectis pinnata* Lam. juss. nat. 2. p. 150. t. 31. *Pectis multifida* Ort. dec. p. 45. *Tetracarpum pinnatisecta* Mørch. l. c. In Peruvîa dicitur *Asha-Pichana* id est scopulæ amara (Ort. in specim. Dombeyano basi induratus suffrutescens. (v. v. c. et s. sp.)  
3. *S. VIRGATA*, pappi paleis 4 acuminato-aristatis, 4 alternis brevioribus

satis, ligulâ involucri breviorē, interdum nullâ. ① in Mexico circa urbem (Berl.! pl. exs. n. 708), ad colles Taenbaya et aliis locis Mexico vicinis (Lallav.), ad occid. urbis Guanajuato (Mend.!). *Mieria virgata* Lallav. et Lex. nov. veg. mex. 2. p. 9. Invol. basi squamulis accessoriis duabus instructum, nec ut in aliis nudum. (v. s. sp.)

† *Species exclusa.*

S. *SENECIONIDES* Nees = *Achyropappus* Schkurnioides.

CCCXCVII. FLORESTINA *Cass. bull. philom.* 1815. p. 175. *journal. phys.* 1816. p. 144. 1817. p. 11. *dict.* 17. p. 155 et 55. p. 266. *atl.* 1816. f. 3. t. 8. *opuscul. phyt.* 1. p. 197. t. 8. — *Lepidopappus fl. mex. ic. ined.* — *Stevia* sp. Cav. — *Agerati* sp. *Ort.* — *Hymenopappi* sp. *Lag.* — *Achyropappi* sect. *Less.*

Capitulum homogamum discoideum. Invol. floribus brevius, squamis 8 uniserialibus oblongis obtusis. Recept. parvum planum subulveolatum. Cor. tubo brevi, lobis 5 inæqualibus papilloso-hirtellis. Styli rami in appendicem subulatam hirtam producti. Achænia subtetragona oblonga hispidula. Pappus paleis 8-12, uniseriatis scariosis suborbiculatis nervo saltē basi opaco. — *Herbæ Mexicanæ, Steviæ facie, pube adpressâ brevissimâ subcanescentes, affines Hymenopappo. Folia alterna pedatipartita seu indivisa. Capitula latē corymbosa seu subpaniculata. Cor. albidæ aut purpurascens. — Si ut sectio habenda, potiùs meo sensu Hymenopappo quàm Achyropappo adjungenda.*

1. F. *PEDATA* (Cass. ll. ec.), foliis petiolatis pedatipartitis, lobis 5 oblongis integris, summis indivisis. ① in Mexico (Cav.), circa urbem in montanis (Berl.! pl. exs. n. 707 et 743). *Stevia pedata* Cav. ic. 4. p. 33. *Ageratum pedatum* Ort. dec. p. 38. *Hymenopappus pedatus* Lag. nov. gen. p. 28. n. 356. *Achyropappus pedatus* Less. syn. 239. Invol. sæpè purpurascens. (v. s. sp. et v. c.)
2. F. *TRIPTERIS*, foliis petiolatis trisectis, segmentis ovatis aut ovali-oblongis aut in apice caulis linearibus, omnibus integerrimis. ① in Mexico ad Larri- (Berl.! pl. exs. n. 2077). *Lepidopappus tripteris* fl. mex. ic. ined. Pappi paleæ acutæ, margine hyalinæ, nervo medio crassiusculo. (v. s. sp.)
3. F. *CALLOSA*, foliis linearibus integerrimis, ramis subdichotomis glanduloso-muricatis. ①? in territorio Arkansano Amer. bor. *Stevia callosa* Nutt.! *journal. ac. Philad.* 1821. p. 121. et in linnæa litt. 1829. p. 40. *Annua à Nuttallio* dicitur, sed in specim. ab eo misso caulis suffruticulosus videtur. Pappi paleæ membranaceæ obtusæ nervo non perspicuo. Cor. purpurascens. Styli rami longè appendiculati, appendice hispidâ stylum *Vernoniæ* ferè simulantē. (v. s. comm. à cl. Nutt.)

#### CCCXCVIII. ACTINOLEPIS.

Capitulum pluriflorum, fl. radii 5-5 ligulatis fœmineis, disci tubulatis 5-dentatis bisexualibus, stylo abortivo masculis? Invol. ovato-oblongum bracteis foliaceis paucis cinctum, squamis paucis obtusis adpressis dorso molliter tomentosis. Recept. angustum nudum. Ligulæ latæ breves 2-3-dentatæ, stylo bifido exserto. Tubuli tubo tereti, fauce dilatata, antheris albis, stylo simplici apice capitellato aut subnullo? Achænia oblonga subangulata, radii pubentia pappo squamellis circ. 5 scariosis acuminatis constante superata, disci glabra calva. — *Herba annua tenella multicaulis Californica. Caules tenuēs apice subaraneosi. Folia opposita parva latē obovata, basi cuneata et integrā, apice obtusissima 5-dentata. Capitula in axillis supremis approximatis solitaria*

subcongesta parva. Flores flavi. — An *Heleniea* ex pappo radii squamellato? An *Partheniea* si discus verè masculus?

1. A. *MULTICAULIS* (1) in Californiâ legit cl. Douglas. Radix exilis simplex. Caules circ. 2-pollic. Folia 2-3 lin. longa et lata, superiora araneosa. Capitula cano-tomentosa. Achænia nigra. (v. s. comm. ab hon. soc. hortie. Londæ)

CCCCXCIX. BAHIA. — Bahia et *Eriophyllum* Lag. nov. gen. in elench. h. nat. 1815. p. 28. et 30. Less. syn. p. 238. — *Trichophyllum* Nutt. gen. am. (1818) 2. p. 166. nec Ren. — *Phialis* Spreng. gen. 2 (1831), n. 631. — *Helenia* Spreng. syst.

Capitulum multiflorum radiatum, ligulis fœmineis 5-11, fl. hermaphr. 5-dentatis extus glanduloso-subsetosis. Invol. ovatis campanulatis squamæ adpressæ 1-2-seriales. Recept. epalaceum nudum aut parcè alveolato-fimbriiferum. Styli disci rami obtusi appendiculati. Achænia turbinato- aut lineari-tetragona, ferè semper glabra. Pappi paleæ 4-8 ovales aut oblongæ obtusiusculæ membranaceæ. — Herbæ perennes aut suffrutices Americani. Folia opposita aut alternà integra aut partita, suprâ glabra aut araneosa, subtus tomentosa. Pedunculi nunc elongati 1-cephali, nunc apice corymbosi. Flores lutei, saltè in genuinis, purpurei in specie dubiâ. — Genus fortè dividendum, sed car. generum nunc propositorum non sufficiunt.

### §. 1. Foliis alternis indivisis.

1. B. *ANGUSTIFOLIA* (Less. in linnæâ 6. 1831. p. 519 sub *Trichophyllo*), cæcis ramis hirsutis, foliis alternis subsessilibus lanceolato-linearibus subentasisabris, pedunculis elongatis glabris. 2? in Monte-Video. Ageratum asperifolium Spreng. syst. 3. p. 446. Capitula radiata lutea. An sequentium? monente Less., omninò congener?
2. B. *STÆCHADIFOLIA*, suffrutescens?, foliis alternis linearibus obtusis marg. subrevolutis suprâ araneosis glabratissve subtus cano-tomentosis, apice corymbosis, invol. squamis et pappi paleis 8-10. 2? in Mexico (Lag., in Real del Monte (Hænk!)). *Eriophyllum stæchadifolium* Lag. elench. h. nat. p. 28. n. 361. *Helenium stæchadifolium* Spreng. syst. 3. p. 574. Folia ex l. interdum opposita, in specim. obvio alternà, sed partes inferiores desidentium elongatum subvillosum. Pappi paleæ oblongæ vix obtusæ. (v. s. à Hænk. à cl. de Sternb. miss.)
3. *Californica*, capitulis paucioribus, invol. pedicellisque cano-tomentosis — in Novâ-Californiâ legit Douglas. An fortè distinguenda? (v. s. comm. ab hon. soc. hort. Londæ.)
3. B. *INTEGRIFOLIA* (Hook. fl. bor. am. 1. p. 316. sub *Trichophyllo*), inferioribus obovato-spathulatis, superioribus lineari-spathulatis integerrimis vel rariissimè apice subtrifidis utrinquè albo-lanatis. — in Amer. bor. occid. cum B. lanatâ.

### §. 2. Foliis oppositis trilobis tripartitisve.

4. B. *SINUATA* (Less. in linnæâ 1831. p. 160), fruticosa, foliis oppositis ternis nerviis sapius trilobis basi truncatis suprâ pubescentibus subtus incanamentosis. 3 in Mexici rupibus prope Hacienda de La Laguna.
5. B. *OPPOSITIFOLIA* (Nutt. gen. am. 2. p. 167 sub *Trichophyllo*), decumbens ramosissima pubescens, foliis oppositis palmato-trifidis, lobis ligulatis implexibus vel divaricato-subdivisis, pedunculis filiformibus plurimum ædotomis vix folio longioribus. 2? in collibus sterilibus Amer. bor. circumf. Mandan. *Helenium oppositifolium* Spreng. syst. 3. p. 573. Caulis salsus. Herba amara (nec aromatica), 6-12-pollic. Ligulæ brevissimæ.

6. *B. AMBROSIOIDES* (Lag. l. c.), fruticosa, foliis oppositis tripartito-multifidis pubescentibus, lobulis subobtusis.  $\S$  in Chili fruticetis rupestribus maritimis (Bert. l.), circa Conceptionem (Lag.), Valparaiso (Bert. l. Gay! Macr. l. Gand. l.). Vulgò *Manzanilla-Simarona* dicitur. *Achyropappus maritimus* Poepp. l. pl. exs. n. 238. Ligulæ albæ elliptico-oblongæ discum longè excedentes. (v. s.)

$\S$ . 3. *Foliis alternis pinnatilobatis.*

7. *B. ARTEMISIAEFOLIA* (Less. l. in linnæa 1829. p. 160. 1831. p. 258), suffruticosa, foliis alternis pinnatipartitis suprà subglabris subtus cano-tomentosis, lobis 2-3-jugis integris, capitulis corymbosis, invol. ovati glabriusculi squamis oblongis.  $\S$  in Californiâ septentr. (Menz.), ad Sanctum-Franciscum (Chamisso l.). Ligulæ ovales parvæ. Invol. cylindraceum 1-2-seriale, squamis subæqualibus oblongis subconvolutis 2 lin. longis. Recept. planum alveolato-simbrilliferum. Pappi paleæ lineares. (v. s.)
- $\beta$ . *Douglasii*, pedicellis involucriisque magis tomentosis, capitulis duplò ferè majoribus, ligulis etiàm majoribus.  $\S$  in Californiâ legit Douglas. (v. s. comm. ab hon. Soc. hort. Lond.)
8. *B. CONFERTIFLORA*, caule herbaceo erecto araneoso aut densè brevissimèque tomentoso, foliis alternis pinnatipartitis, lobis linearibus integris subtus sub-tomentosis, capitulis confertè corymbosis, invol. ovati squamis obovatis retusis arachnoideo-tomentosis.  $\mathcal{L}$  in Novâ-Californiâ legit cl. Douglas. Folia ferè *B. lanatæ*, sed angustiora. Capitula confertè corymbosa nec solitaria, duplò minorâ, magis ovata. Ligulæ paucæ obovatæ. (v. s. comm. ab hon. Soc. hort. Londin.)
9. *B. TENUIFOLIA*, herbacea erecta tota pube adpressâ aut subaraneosâ canescens, foliis alternis apice pinnatipartitis margine revolutis, rachi lineari integrâ, lobis 1-3-jugis linearibus dentatis integrisve, corymbo laxo, invol. tomentoso, ligulis obovatis inv. duplò longioribus.  $\mathcal{L}$  in Californiâ legit cl. Douglas. Habitus lanugine exceptâ Senecionis artemisiæfoliæ. Capitula parva aurea. Pappi squamellæ oblongæ vix acutæ. (v. s. comm. ab hon. Soc. hort. Lond.)
10. *B. LANATA* (Nutt. gen. am. 2. p. 168 sub *Trichophyllo*), decumbens sæpè erecta parçè ramosa, caule tereti foliisque subtus lanuginoso-tomentosis, foliis alternis pinnatifidis, summis linearibus integris, pedunculis elongatis 1-cephalis tomentosis.  $\mathcal{L}$  ad fontes flum. Columbia (Pursh l. Dougl. l.) et Kooskoosky (Hook.). *Actinella lanata* Pursh fl. bor. am. 2. p. 560. *Eriophyllum cæspitosum* Dougl. l. in Lindl. bot. reg. t. 1167. *Helenium lanatum* Spreng. syst. 3. p. 574. *Phalis* Spreng. gen. 2. p. 634 excl. car. Involucrum enim non verè gamophyllum sed squamis liberis 1-seriatis. Herba pedalis *Hymenopappum artemisiæfolium* aut *Bahiam artemisiæfoliam* foliis referens. Ligulæ magnæ. (v. s. comm. ab hon. Soc. hort. Lond.)
11. *B. LEUCOPHYLLA*, tota cano-lanuginosa, caule herbaceo erecto, foliis oppositis et alternis obtusè sinnato- aut pinnatilobatis, lobis foliisque summis dentatis, pedunculis elongatis 1-cephalis, capitulis multi-(25-30) floris, acheniis glaberrimis tetragonis, pappi paleis octo, 4 lanceolatis acutis, 4 minimis alternantibus.  $\textcircled{1}$   $\S$  ad Nootka et Mullgrave Amer. occid.-bor. leg. cl. Néel et Hanke! Ex specim. malè servato priori videtur affine. Herba 8-10-pollicaris Senecionis incanos referens, tota, etiàm in invol., lanugine densè vestita. (v. s.)
12. *B. ACHILLEOIDES*, basi fruticosa ramosa, ramis virgatis teretibus tomentosis apice nudis parçè ramosis, foliis alternis basi cuneato-attenuatis apice pinnatipartitis, lobis 2-4-jugis approximatis dentatis suprà araneosis subtus tomentosis, capitulis ramulos terminantibus solitariis, invol. subgloboso lanato.  $\S$  in Californiâ legit cl. Douglas. Folia 7-9 lin. longa apice cristato-partita, lobulis apice callosis. Ligulæ oblongæ flavæ. Pappus squamellatus parvulus. (v. s. comm. ab hon. Soc. hort. Lond.)
13. *B. TROLLIFOLIA*, tota subincana, caule herbaceo, foliis alternis, infer. tripartitis multifidis dentatis, super. indivisis linearibus, pedunculis elongatis

1-cephalis, invol. 9-11-phylo, pappi paleis 4 lanceolatis acutis canaliculatis 2/4 in Mexico legit cl. Née. *Eriophyllum troiliifolium* Lag. elench. h. Mad. p. 28. n. 360. *Helenium troiliifolium* Spreng. syst. 3. p. 573. Cor. 9-11 purpureæ (Lag.). An verè congener?

CD. HYMENOPAPPUS *L'Hér. diss. cum icon. sine anno. Cass. dict. 55. p. 166 et 279. Less. syn. 238. — Rothia Lam. journ. h. nat. 1792. v. 1. p. 16. t. 1 non Schreb. — Hymenopappi sp. Juss.*

Capitulum homogamum. Recept. parvum nudum foveolatum. Invol. squamæ 2-3-seriales elliptico-oblongæ obtusissimæ, interiores interdum majores petaloides. Cor. tubo basi dilatato extius glanduloso-setigero, limbo campanulato 5-fido. Anth. exsertæ. Styli lobi apice breviter appendiculati. Achænia turbinate striata basi attenuata subpedicellata, matura subtetragona. Pappus paleis circ. 12-16 membranaceis brevibus elliptico-oblongis obtusis 1-serialibus. — Herbae Carlinianæ perennes. Caules sulcato-angulati ramosi infernè glabri, supernè tomentosi aut velutini. Folia alterna pinnatipartita, lobis sæpè incisis aut multifidis. Capitula corymbosa pedicellata. Corolla et inv. squamæ majores albidæ.

1. *H. tenuifolius* (Pursh fl. bor. am. suppl. p. 742), foliis glabriusculis subbipinnatipartitis, lobis linearibus integris, corymbo composito subconfero, invol. squamis 5-6, interioribus vix flosculos superantibus, achæniis villosissimis. (2) (Nutt.) in Louisianâ superiore (Pursh), ad Red-River (Nutt. ! herb.) et ad ripas fl. Missouri in collibus glareosis (Nutt. gen. am. 2. p. 139). A sequente certè videtur distinctus. (v. s.)
2. *H. scabiosæus* (L'Hér. ! l. c.), foliis subtus caudicanti-lanuginosis pinnatipartitis, lobis lineari-oblongis vix subdentatis, corymbo simpliciusculo laxo, invol. squamis interioribus 10 majoribus membranaceo-petaloides flosculos superantibus, achæniis parè villosis. 2/4 (Ell.) in Carolinâ (Michx. ! Frax.) et Georgiâ (Ell. sketch 2. p. 313). Michx. ! fl. bor. am. 2. p. 104. Pursh fl. 2. p. 519 et 742. *Rothia Carolinensis* Lam. l. c. et ill. t. 667. *Hym. laxiflorum* L'Hér. ! mss. (v. s.)
3. *H. artemisiæfolius*, foliis subtus cano-tomentosis, infer. in petiolum longè attenuatis oblongis sinuato-dentatis, cæteris sessilibus pinnatipartitis, lobis 3-4-jugis integris acutis cum terminali majore, corymbo composito laxo, invol. squamis ovalibus disco vix longioribus, achæniis villosissimis. (3) in Mexicanâ prov. Texas legit cl. Berlandier (pl. exs. n. 1532). Bractee semipedicellos obovate membranaceæ obtusæ. Cor. albidæ. Folia *Artemisiæ campestris* aut *Bahiæ artemisiæfoliæ*, (v. s.)
4. *H. douglasii* (Hook. fl. bor. am. 1. p. 316), lanâ albidâ deciduâ tomentosa, ramis elongatis subcorymbosis, foliis bipinnatifidis, lobis linearibus sinuatis, capitalis solitarii, invol. squamis sub-20 linearibus pubescentibus disco brevioribus, corollis filiformibus apice pilosis pappi paleas lineares obtusas nec vix superantibus. — in siccis arenosis Columbiæ à loco dicto Great-Falls ad Rocky-Mountains. *H. scabiosæus* Dougl. mss. non L'Hér. (v. s. comm. ab nom. ab hon. Soc. hort. Lond.)
5. *H. filifolius* (Hook. fl. bor. am. 1. p. 317), lanâ albâ deciduâ tomentosa, ramis paniculatis sparsis, foliis bipinnatifidis, lobis filiformibus elongatis integerrimis, capitulis solitariis, invol. squamis sub-12-ellipticis pubescentibus disco subbrevioribus, corollæ tubo supernè campanulato, pappo brevissimo. — in aridis flum. Columbiæ ad Wella-Wella et ad ripas flum. Spoken et Flat-Head Amer. bor.-occid. *H. tenuifolius* Dougl. mss. non Pursh.



† *Species exclusæ.*

H. ANTHEMOIDES Juss. = *Hymenoxys anthemoides*.

H. GLAUCUS Spreng. = *Cephalophora glauca*.

H. PEDATUS Lag. = *Florestina pedata*.

CDI. CHÆNACTIS.

Capitulum multiflorum homogamum, fl. radii amplis basi tubulosis sursum subbilabiatis aut subpalmatis, disci tubulosis regulariter et obtusè 5-dentatis. Invol. campanulatum circ. 20-phyllum, squamis erectis 1-2-seriatis linearibus acutiusculis. Recept. epaleaceum alveolatum. Cor. tubo glabro, lobis hirsutulis. Styli disci rami hispidi cono obtuso brevi superati. Achænia lineari-tetragona basi attenuata subpedicellata adpressè pubescenti-villosa. Pappus paleis 5-6 membranaceis, radii brevibus obtusis, disci longioribus subacutis corollæ tubi dimidiam partem æquantibus. — Herbæ Californicæ erectæ habitu *Hymenopappum tenuifolium* referentes. Folia alterna pinnatipartita lobis linearibus integris. Rami apice nudi monocephali. Flores lutei. — Genus inter *Hymenopappum* et *Bahiam* medium; à priori fl. radii majoribus hiantibus, pappo squamellis 5-6 nec 15-16; à posteriore fl. radii non ligulatis nec fœmineis; ab utroque, pappo radii et disci diverso.

1. C. GLABRIUSCULA, caule ramoso, foliis involucriisque glabriusculis seu hinc inde subaraneosis, fl. radii palmatifidia expansis disco evidenter longioribus.  $\mathcal{U}$  5 in Novâ-Californiâ legit cl. Douglas. Herba pedalis, junior araneoso-villosa, adulta glabra. Foliorum lobi utrinquè 5-6 obtusiusculi. Folia summa linearia integerrima. Rami ad apicem per intervallum 2  $\frac{1}{2}$  poll. aphylli. (v. s. comm. ab hon. Soc. horticult. Lond.)

2. C. LANOSA, caulibus è collo plurimis foliis involucriisque cano-lanatis, floribus radii tubuloso-infundibuliformibus vix disco majoribus. ( $\mathcal{U}$ ) in Novâ-Californiâ etiam legit cl. Douglas. Radix exilis simplex. Folia rarissimè glabrata cæterum cano-lanata, summa linearia parva. Ramorum pars nuda 6 pollic. circ. longa. (v. s. comm. cum prior.)

CDII. POLYPTERIS Nutt. gen. 2. p. 139? Ell. sketch 2. p. 313? non Less.

Capitulum homogamum. Invol. squamæ biseriales foliaceæ ovali-lanceolatæ breviter acuminatæ, jun. erectæ demum reflexæ. Recept. hemisphæricum omninò nudum. Cor. tubo brevi, fauce amplâ inflatâ, lobis 5 subulatis hirtellis. Styli rami in appendicem linearem hirtam longè producti. Achænia turbinata sericeo-villosa. Pappi paleæ 9 achænio ferè longiores basi angustatæ membranacæ grossè uninerviæ, nervo in aristam longam rigidam producto. — Herba Boreali-Americana erecta ramosa pubescenti-scabra. Folia alterna integra oblongo-linearia sessilia. Capitula ramos terminantia solitaria. Cor. fuscæ aut pallidè flavæ, lobis fuscis. Achæniæ setæ et pappi aristæ fuscæ. — Genus forsan à *Cercostylo* non satis diversum.

1. P. INTEGRIFOLIA.  $\mathcal{U}$  ? in Carolinâ meridionali olim legit Fraser. P. integrifolia Nutt. à nostrâ differt invol. squamis oblongo-ellipticis membranaceis albidis, achænio subtetragono scabro nec villosa, pappi paleis 12-14. An verò ut vult Less. ad *Palafoxiam* referenda? P. integrifolia Ell. differt tantum achænio leviter scabro. Nostra certè ad *Palafoxiam* non referenda. Ex notolâ herb. L'Hér. ad *Hymenopappum* referebat cl. Michaux sed in Michx. fl. desideratur. (v. s.)

† *Species exclusa.*P. BRASILIENSIS Less. = *Cercostylos Brasiliensis*.

## CDIII. ESPEJOA.

Capitulum 7-florum homogamum, fl. tubulosi 5-fidis glabris hermaphrod. Invol. 4-seriale, squamis 8 oppositis latè ovatis imbricatis apice mucronatis. Recept. alveolatum. Styli rami obtusi parè puberuli. Anth. breves inclusæ. Achænia obcompressa-cuneata villis sericeis rufis confertissimis onusta, disco epigynò lato orbiculari. Pappus palès circ. 16 uniserialibus membranaceis ovali-oblongis 1-nerviis mucronulatis corollæ ferè longitudinem adæquantibus, villis sericeis achæniæ basi cinctis. — Herba glabriuscula erecta dichotoma Mexicana. Caulis subangulatus versus apicem pilosiusculus. Folia opposita ovali-oblonga sessilia integerrima. Pedunculi axillares 1-cephali. Flores ex sicco flavo-subpurpurascentes. — Genus Mexicanum dicavi cl. navigatori Hispano Antonio de Espejo qui anno 1533 plagas Mexicanas primus adiit.

1. E. MEXICANA. (1) ? in Mexico ad Divum-Dionysium septemb. flor. legit cl. amic. Alamau. Stobæa glabrata var. Alam. in litt. 1831. (v. s.)

CDIV. CERCOSTYLOS Less. syn. 239. — *Polypteris* Less. in *linnæa* 1831. p. 518 nec Nutt.

Capitulum homogamum. Invol. campanulatum biseriale, squamis longè acuminatis. Recept. convexum epaleaceum densè paleaceo-fimbriiferum. Cor. 5-dentatæ. Styli rami appendice lineari angustissimâ. Achæmium turbinatum villosum. Pappi palææ decem 1-nerviæ apice laceræ in aristulam acuminatæ. — Herba Brasiliensis basi lignescens, foliis sessilibus vel decurrentibus integerrimis alternis pilosiusculis canescentibus, capitulis solitariis caulem ramosque elongatos et superæ aphyllis superantibus. Cor. luteæ. — Unica species adhucdum desc.

1. C. BRASILIENSIS (Less. syn. 239). 5 in Brasiliâ. *Polypteris Brasiliensis* Less. in *linn.* 1831. p. 518. Capit. magn. Gaillardia lanceolatæ. Folia oblongo-obovata punctata, 1½-2 poll. lata, vix semilineam lata.

## CDV. HOPKIRKIA. DC. non Spreng (1).

Capitulum 3-florum homogamum. Invol. oblongum 4-5-phyllum. squamis obovato-oblongis subscariosis, bracteolis minimis 2-3 las stipatum. Recept. nudum angustum. Cor. tubulosæ pappo breviores inæqualiter 4-5-dentatæ. Styli rami breves apice capitellati. Achænia tetragona basi attenuata substipitata, angulis densè villosa-ciliatis. Pappi squamellæ 8 scariosæ medio 1-nerviæ apice subtrifidæ nervis aristulam producta. — Herba Mexicana ramosa adscendens glabra Folia alterna pinnatipartita, lobis 1-2-jugis cum impari angustè linearibus elongatis integerrimis. Capitula solitaria pedicellata terminalia parva. Flores lutei. — Cum *Hopkirkia* Spreng. sit omninò delatæ. dicavi hoc novum genus cl. Thom. Hopkirk auctori Floræ anomor. nè honore debito fraudetur.

2. H. ANTHEMOIDRA. — in Mexici agro Regiomontano legit cl. Hænke. (v. s. h. Hænke à cl. de Sternberg comm.)

(1) *Hopkirkia* Spreng. (nov. prov. 1818) est *Salmea* DC. (cat. h. monsp. 1818)

CDVI. HYMENOXYS *Cass. dict. 55. p. 278.*

Capitulum multiflorum discoideum, homogamum seu ligulis 8-10 uniserialibus fœmineis, fl. disci hermaphr. tubulosis 5-dentatis. Invol. biserialia, squamis rigidulis adpressis, interioribus longioribus. Recept. conicum epaleaceum alveolatum, glandulis parvis subonustum. Stylus disci ramis apice truncatis barbellatis. Achænia conformia turbinata villosa sericeis erectis villosissima. Pappus paleaceus, paleis 5-8 inæqualibus membranaceis ovali-lanceolatis aristato-acuminatis erectis. — Herbae Americanæ erectæ ramosæ glabriusculæ, trita odorem Anthemidearum spirans. Folia alterna aut in eodem caule rarius opposita, in lobos lineares multipartita et Schkuhræ aut Achyropappi æmula. Rami apice nudi pedunculiformes. Cor. luteæ.

Sect. I. EUHYMENOXYS. — *Hymenoxys Cass. l. c.*

Capitula discoidea homogama.

1. *H. ANTHEMOIDES* (*Cass. dict. 55. p. 278*), invol. squamis obovatis subspathulatis. — in Bonariâ (*Comm.*). *Hymenopappus anthemoides* *Juss. ann. mus. Par.* 2. p. 426. Incautè, ex accuratâ *Cass. descr.*, cl. *Jussieu* pappum duplicem et *Persoon* folia decurrentia dixerunt. Capitula egregiè nota, sed herbae descr. non sufficiens. — *Cephalophora anthemoides* *Less. syn. 240* et in *linnæa* 1831. p. 518. videtur ex capitulo radiato diversa.
2. *H. HÆNKEANA*, invol. squamis ellipticis obtusiusculis. — in Peruvia legit cl. *Hænke*. An ab *H. anthemoides* mihi ignotâ satis distincta? Caulis teres divaricato-ramosus. Folia pinnatipartita, rachi lobisque linearibus integerrimis. Rami apice nudi et pedicelli ex axillis supremis orti 1-cephali apice striati. Capitula globosa. Recept. in centro conicum. Achænia tereti-turbinata longè sericea villosa. Pappi paleæ basi ovales in aristam cuspidatæ. Corollæ viz pappo longiores tubo breviteret. Styli inclusi. (v. s. in *h. Hænke*. à cl. de *Sternberg* comm.)

Sect. II. OXYAPPUS. — *Cephalophoræ* sect. 3. *Less. syn. 240.*

Capitula radiata.

3. *H. ODORATA*, invol. squamis lanceolatis acuminatis. ? in Mexico legit cl. *Berlandier*, sed sched. deperditâ, locus proprius ign. *Cephalophora anthemoides* *Less. in Linn. 1831. p. 518. syn. 240 excl. syn. Juss.* Folia multò tenuius pinnatipartita quàm prioris. Pedunculi foliis duplè triplè longiores. Ligulæ 5-6 flavæ apice grossè tridentatæ. Pappi squamellæ lanceolatæ acuminatæ. Capitulum tritum valdè odoratum. (v. s.)
4. *H. CHRYSANTHEMOIDES*, caule herbaceo? ramoso polycephalo glabro apice villosopubescente, foliis petiolatis subpubescentibus subbipinnatilibus multipartitis, lobis linearibus, ramis apice nudis 1-cephalis, invol. squamis biserialibus, pappi paleis 5 acuminato-subulatis. ? propè *Moran Mexicanorum* (*H. B.*) et in dititione *Chalco* (*Andr.!* pl. exs. n. 316). Actines chrysanthemoides *H. B.* et *Kunth* nov. gen. am. 4. p. 298. t. 411. Capitula flava magnitudine *Leucanthemi*. Achænia juniora lineari-cuneata sericeo-hirsuta. Ligulæ 12-15 apice 3-lobæ (v. s.)

CDVII. CEPHALOPHORA *Cav. ic. 6. p. 79. t. 599. — Græmia Hook. exot. fl. t. 189. — Cephalophoræ* sect. 1a. et 2a. *Less. syn. 240.*

Capitulum sphæricum homogamum aut radiatum, ligulis fœmineis tridentatis. Invol. squamæ biserialia reflexa lineari-lanceolatæ flosculis æquales aut iis breviores. Recept. nudum foveolare. Cor. extus hispidula, tubo brevissimo, fauce obovoideâ tubulosâ, limbo 5-dentato

connivente subcalloso, Anth. inclusæ. Stylus filiformis. Styli rami exserti patulo-revoluti, apice hispiduli exappendiculati. Ovarium turbinato-4-gonum pilis squamosis adpressis tectum. Pappi paleæ 5-8 membranaceæ margine erosæ apice apiculatæ. — Herbariæ aut suffrutices Chilenses erecti glauci, caule striato ramoso brevissimè puberulo, foliis alternis oblongo-linearibus acutis. Capitula ramos nudiusculos terminantia. Cor. flavæ, juniores apice fuscæ. Involucri squamæ cum flosc. externis interdum mixtæ.

## Sect. I. EUCEPHALOPHORA.

Capit. homogamum.

1. *C. GLAUCA* (Cav. l. c.), caule suffruticoso ramoso, foliis oblongo-lanceolatis, infer. inciso-serratis, summis integerrimis. ③ et ex Bert. variat ② in Chili propè Cucha-Cucha (Née!), in pascuis et fruticetis collium propè Valparaíso (Bert.!). *Ethulia repanda* Née! herb. *Hymenopappus glaucus* Spreng. syst. 3. p. 449. *Santolina tinctoria* Mol. chil. ed. gall. p. 336. *Santolinoides* vulgò *Poquill.* Feniell. obs. 3. t. 45. f. 2. Pedunculi teretes in omnibus æcim. sylvestr. et etiã in Neaxanis cultis, nec ut in ic. Cav. incrassati. Folia exsiccatæ ad colorem luteum utuntur Chilenses. (v. v. et s. sp.)
2. *C. PLANTAGINEA*, caule herbaceo sparsè ramoso, foliis radicalibus in petiolum attenuatis oblongo-lanceolatis basi cuneatis apice obtusis repando-dentatis, caulinis distantibus linearibus acutis integerrimis. ② in Chili? Radix subsimplex elongata. Rami circ. 3 valdè elongati. Pappus rufo-fuscus. (t. s. comm. à Mus. reg. Par. ex coll. Dombey sine loci design.)
3. *C. AROMATICA* (Schrader. ind. sem. h. Goett. 1830. Less. syn. 240), caule herbaceo ramosissimo, foliis lineari-lanceolatis subundulatis semiamplexicaulis, infer. pinnatifidis, sup. vix subdentatis. ① in Chili (Hook.) in pascuis et fruticetis collium propè Valparaíso (Bert.!) ubi (ex odore aromatico ad Chamomillam accedente) dicitur *Manzanilla-del-Campo*. *Graminia aromatica* Hook. exot. bot. 2. t. 189. opt. (non Poepp. quæ *Euxenia Mitiqui* var. ②). *Cephaloph. tenera* Cass. opusc. 3. p. 93. Capitula dimidio quàm *C. glauca* minora cæterum similia. (v. s. sp. et v. c.)

Sect. II. ACTINELLA Less. syn. 240. — *Actinella Pers.* 2. p. 460. n. 1888. — *Actinea Juss. ann. mus. Par.* 2. p. 426. — *Actinea* et *Dugaldea Cass. dict.* 55. p. 270. — *Actineæ* sp. *Spreng.*

Capitula radiata, ligulis fœmineis apice 3-dentatis.

§. I. *Invol.* 1-3-seriali adpresso disco breviori aut subæquali *Acheniis turbinatis.* — *Folia non decurrentia.*

\* *Caule fruticoso folioso.*

4. *C. HETEROPHYLLA* (Less. syn. 240), caule fruticoso ramoso rufescente, foliis sessilibus lineari-lanceolatis, infer. grossè serrato-incisis, super. integerrimis, ramis apice nudis 1-cephalis, invol. squamis 1-2-serialis lanceolatis disco brevioribus, pappi paleis 8 oblongis aristatis. ③ in Bonariâ. *Actinea heterophylla* Juss. ann. Mus. 2. p. 425. t. 61. f. 1. *Actinella heterophylla Pers. ench.* 3. p. 46. *Actineæ alternifolia Spreng. syst.* 3. p. 574. *Achænia oblonga villosa, pappi biapiculatis. Ligulæ (albæ?) apice trilobæ* Cass. dict. 55. p. 270.
5. *C. RADIATA* (Less. in *linnæâ* 1831. p. 516), caule basi fruticuloso ramosissimo glabro, foliis sessilibus linearibus integerrimis punctatis, ramis oppositis 1-cephalis, invol. biserialis squamis ext. ovato-linearibus acumatis, inter. oblongis vix mucronatis, pappi paleis oblongo-ellipticis aristatis. ④ ⑤ in campis ad Rio-Negro et Rio-Grande Brasilæ. Folia pollicem long.

lineam lata. Capit. magnitudine Anthemidis arvensis. Recept. hemisphaericum. Cor. disci limbo apice purpureo, ligulæ pallide lutes. Achænia turbinata fusco-villosa. Anth. obsol. caudatæ. (v. s. in h. Mus. reg. Par. ex. h. Mus. imp. Bras. sub n. 1013. 1042. 1051 et 1053.)

\* \* *Caulis herbaceo aphylo scaposo, foliis radicalibus.*

6. C. ACAULIS, scapo nudo elongato 1-cephalo, foliis radicalibus lineari-spathulatis integris punctatis villosis, invol. squamis circ. biseriatis oblongo-lanceolatis, pappi paleis 5-6 apice aristatis.  $\frac{1}{2}$  in collibus Missouriensibus ad Fort-Mandan Amer. bor. Galaridia acaulis Pursh. fl. bor. am. 2. p. 743. Actinella acaulis Nutt. gen. am. 2. p. 173. Capitula magnit. Chamomillæ, latè flava. Ligulæ 8-10 apice 3-dentatæ.
7. C. SCAPOSA, villosa, caule aphylo stricto simplici 1-cephalo, foliis radicalibus lineari-lanceolatis basi attenuatis, aliis integerrimis, aliis pinnatifidis lobis paucis acutis, invol. ext. squamis disco brevioribus obtusis adpressis.  $\frac{1}{2}$  in Mexicanæ prov. Texas district. orientalibus legit cl. Berlandier (pl. exs. n. 1867 et 1882). Scapus 8-12 poll. longus. Folia 2-3 poll. longa, 2-3 lin. lata, acuta. Ligulæ 4-nerviæ 3-dentatæ. Pappi paleæ uninerviæ, margine scariosæ, ovales, abruptè aristatæ, cor. disci paulò breviores. (v. s.)

§. 2. *Invol. biseriale, squamis exter. linearibus reflexis. Achænia tetragona. — Folia ut in Helenio decurrentia.*

8. C. DECURRENS (Less. in Linnæa 1831. p. 517), caule herbaceo dichotomo pubero, foliis decurrentibus oblongis subrepandis puberis, infer. obtusis, super. acuminatis, ramis apice aphyllis 1-cephalis, invol. squamis linearibus reflexis, pappi paleis 6 ellipticis pinnatifidis aristatis.  $\frac{1}{2}$  in Californiâ. An ex Less. Helenium pubescens Ait. kew. 3. p. 227? Ligulæ disci diametro breviores. Achænia tetragona. Anthere obsolete caudatæ.

† *Species exclusæ.*

- C. ANTHEMOIDES Less. = Hymenoxis odorata.  
 ACTINELLA COMMUTATA Spreng. = Calea pinnatifida.  
 A. LANATA Pursh. = Bahia lanata.

CDVIII. JAUMEA Pers. ench. 2. p. 397. n. 1816. — Kleinia Juss. ann. Mus. 2. p. 424. t. 61. f. 1. non Linn. nec Willd.

Capitulum multiflorum discoideum homogamum. Invol. latum patens imbricatum, squamis subrotundis triplici ordine dispositis. Recept. nudum. Anth. ecaudatæ appendice ovato-lanceolatâ terminatæ. Styl. rami vix ac ne vix subpapilloso cono brevi terminati. Achænia teretiuscula costis minimis longitudinalibus notata. Pappi squamellæ 8-10 breves inæquales denticulatæ acutæ. — Suffrutex Bonariensis glaber. Folia opposita connata linearia carnosa integra. Capitula terminalia solitaria nutantia. Car. ex Juss. descr. et ic. nec non ex notis et icone ineditis à cl. Adr. de Jussieu et Decaisne juxta specim. ipsum Commers. paratis et mecum benevolenter communicatis.

1. J. LINEARIFOLIA (Pers. l. c.).  $\frac{3}{4}$  in Bonariâ ad ostium flum. Plata. — Flores ex sched. Commers. vix comparebant lutei.

#### CDIX. BURRIELIA.

Capitulum multiflorum heterogamum, fl. radii ligulatis fœmineis obovatis, disci plurimis tubulosis 5-fidis hermaphroditis aut interdum abortu sterilibus. Involucri patenti-campanulati squamæ ovales acumi-

natæ disco paulò longiores uni- aut biseriales subalternæ æquales. Recept. epaleaceum. Cor. disci lobis extus tenuè barbellatis. Styli disci ramis cono brevi terminatis. Achænia gracilia lineari-subtetragona, radii ferè obcompressa pappo longè bi- seu triaristato, disci pappo 3-4-phylo, paleis lanceolato-aristatis corollam longitudine subæquantibus. — Herbæ Californicæ annuæ graciles parcè puberulæ aut glabræ subsimplices erectæ. Folia opposita linearia longa integerrima. Capitula ramos caulesve apice nudos terminantia. Flores aurei — Dixi in honorem cl. Job. Marc. Burriel qui anno 1758 iter Venegasii in Californiam edidit et auxit.

1. *B. GRACILIS*, ligulis et invol. lobis 10-12. (1) in Novâ-Californiâ legit cl. Douglas. Herba 5-6 pollic. Folia 2 poll. longa, lineam lata. Capitulum lei Coreopsidis Bidentisve. (v. s. comm. cum seqq. ab hon. Soc. hort. Lond.)
2. *B. TENERRIMA*, ligulia et invol. lobis circ. 5. (1) in Novâ-Californiâ legit cl. Douglas. Herba 3-4-pollic. Magis adhuc glabra et tenera quàm prior. Folia 6 lin. longa, vix  $\frac{1}{2}$  lin. lata. Achænia disci 3-aristata. (v. s.)
3. *B. MICROGLOSSA*, ligulis et invol. squamis circ. 3. (1) in Novâ-Californiâ legit cl. Douglas. Herba habitu *B. tenerrimæ* sed villosiuscula et primâ fronde discoidea. Folia  $\frac{1}{2}$ -1 lin. lata. Ligulæ 2-3 brevissimæ integræ stylo suo superatæ. Achænia omnia 2-3-aristata. (v. s.)

CDX. *LASTHENIA* Cass. opusc. 3. p. 88 (1834). DC. in Lindl. bot. reg. t. 1. p. 8. — Rancagua Poepp. et Endlich. nov. gen. ac spec. pl. amer. t. p. 15. t. 24. p. 25 (1835).

Capitulum multiflorum subhomogamum, fl. radii nempè feminis paucis parvis tubulosis obliquè truncatis (ex Cass.), Recept. conicum epaleaceum papillosum. Invol. squamæ uniseriales concretæ in cupulam 5-8-15-dentatam, dentibus acutis ciliatis. Corollæ breves, fauce inflatâ, limbo 5-dentato. Anth. inclusæ, Styli in radio exserti, in disco subinclusi, rami breves extus arcuati cono brevi superati. Achænia compressa adpressè pubescentia, areolâ basilari corneâ cavâ. Pappus paleis 5-10 apice dentato-laceris acutis corollam subæquantibus est nullis. — Herbæ Occid.-Americanae, glabræ aut subpilosiusculæ, erectæ, ramosæ, in uliginosis nascentes, Enhydræ ferè habitu. Folia opposita linearia subintegerrima ad apicem limbi et lobulorum callosa-obtusâ. Rami nudi apice 1-cephali sub capitulo in conum inversum cavum dilatati. — Genus affine Gamolepidi ex involuero gamophyllo, Coinogyni ex foliis oppositis, ab utroque distinctum capitulo subhomogamo et pappo paleaceo; à Tagetinceis differre videtur glandalis nullis.

### §. 1. Pappo paleaceo.

1. *L. GLABERRIMA*, tota glaberrima, involuero 15-dentato, pappo 5-paleaceo. (1) in uliginosis Californiæ legit cl. Douglas. Lasthenia Californica (DC. à Lindl. bot. reg. n. 1780 excl. car.) Cum error quidam per litteras in textu Lindleyano irreperit et nunc altera species in Californiâ detecta fuerit, nomen deleo nec confusio persistat. (v. s. comm. ab hon. Soc. hortie. Londin.)
2. *L. ORTUSIPOLIA* (Cass. opusc. 3. p. 89), glaberrima, involuero 5-8-dentato, pappo 10-paleaceo. (1) in Chili ad fossarum margines locis nidis propè Rancagua legit cl. Bertero. Tagetes Fenillei Bert. ! pl. exs. n. 680. Colla mem. à Turin 38. p. 34. n. 15. t. 30. sed à Tagete Chilensi Feuill. omnino differre videtur. Rancagua Feuillei Poepp. et Endlich. nov. gen. ac sp. amer. t. p. 15. t. pl. Folia 2 poll. longa, lineam lata, integerrima vix obtusa (undè nomen speciei Cassinianum minus aptum), summa 2 lin. longa dente grosso utrinque medi-

aucta. — Herba procumbens. Invol. 5-dentatum. Pappi paleæ lineares, corollæ breviores, apice lacero bi- vel trifidæ. Poepp. et Endlich. l. c. (v. s.)

β. *Bridgesii*, erecta, pappi corollam æquantis paleis ovato-lanceolatis duplicato-serratis. Rancagua Bridgesii Poepp. et Endlich. l. c. t. 25. An species distincta?

### §. 2. *Pappo nullo.*

3. L. GLABRATA (Lindl. bot. reg. t. 1780), ramulis pedicellisque tenuissimè pubescentibus, invol. 15-dentato, pappo nullo, achæniis apice mucronatis. ① in Californiâ legit cl. Douglas. Similis L. glaberrimæ sed non omnino glabra, capitula paulò majora, invol. basi vix attenuatâ et potiùs campanulâtâ. (v. s. comm. ab hon. soc. hortie. Lond.)

### CDXI. PICRADENIA Hook. fl. bor. am. t. p. 317. t. 108.

Capitulum multiflorum radiatum, ligulis 8-10 fœmineis obovato-oblongis grossè tridentatis, fl. disci hermaphr. 5-dentatis extùs glandulosus. Invol. subhemisphærici squamæ imbricatæ, externæ basi coadunatæ. Recept. conicum nudum. Styli disci rami inclusi breviusculi apice incrassati dorso hirsuti. Achænia oblonga villosa. Pappi paleæ 6-7 erectæ ovatæ acuminatissimæ nudæ membranaceæ. — Herba Boreali-Americana. Radix crassa. Folia alterna linearifiliformia rigida pinnatifida glanduloso- et pellucido-punctata. Capitula terminalia flava. Glandulæ cor. disci amarissimæ undè nomen. — Cl. auctor ad Helenieas suum refert genus et hîc ex ejus sententiâ admisi sed ex stigmate ab ipso Hookero delineatq potiùs ad Astereas Achyrideas pertinere videtur.

1. P. RICHARDSONI (Hook. l. c. p. 317. t. 108). 2. in Amer. bor. circa Carlton-House ad Saskatchawan.

### CDXII. HECUBÆA.

Capitulum multiflorum; fl. radii ligulatis fœmineis 3-5-fidis, disci planiusculi fl. hermaphroditis 5-fidis, lobis brevibus fusco-hirsutis. Invol. biseriale, squamis exterioribus foliaceis patulis radium æquantibus, int. brevioribus linearibus discum vix paulò superantibus. Recept. nudum planiusculum. Styli rami apice truncati subcapitati hispiduli. Achænia angulata glabra glandulosa calva. — Herba Mexicana glabra pedalis simplex, caule tereti ad medium folioso apice nudo pedunculiformi 1-cephalo, pedunculo sub capitulo obconico fusco-subvillosa. Folia alterna decurrentia oblongo-lanceolata acuminata integerrima. Capitula majora et plauiora quam Helenii, ligulis flavis, disco fusco-purpurascente. — Genus Helenio valdè affine Hecubæam, ex Hecubâ Heleni matre, ob analogiam summam dixi.

1. H. SCORZONERÆFOLIA. 2. in Mexico legit cl. Alaman qui cum plur. aliis communicavit et genus novum in schedulâ indicavit. (v. s.)

CDXIII. HELENIIUM Linn. gen. n. 961. Lam. ill. t. 688. — Helenia Lin. hort. cliff. 418. Manch meth. 589. Gærtn. fr. 2. p. 438. t. 169. — Helenium et Tetradus Cass. Less.

Capitulum multiflorum radiatum, fl. radii fœmineis 1-seriatis ligulatis aut rarò tubulosis subirregulariter 3-5-fidis, fl. disci hermaphrod. tubulosis brevissimè 4-5-dentatis. Recept. convexum globosum aut oblongum, nudum. Invol. squamæ biseriales, exter. plurimæ foliaceæ elongatæ sublineares reflexæ aut patentissimæ, inter. subpauciores

breviores acuminatæ recept. paleas simulantes, sed extra flores sitæ. Styli rami obtusi exappendiculati. Achænia turbinato-obovata multifariam villosa. Pappi paleæ 5-6 membranaceæ plus minus apiculatæ. — Herbæ Boreali-Americanae glabræ aut pubescentes. Folia alterna decurrentia subtus punctata integra dentata aut subpinnatifida. Capitula ad apices caulis et ramorum solitaria. Cor. luteæ, extus puberulæ.

Sect. 1. *HELENIA* Linn. *Gærtn. Mæsch II. cc.* — *Helenium Cass. dict. 55. p. 262. Less. syn. 240.*

Cor. disci 5-dentatæ. Pappi paleæ apice aristatæ aut distinctè apiculatæ. Recept. convexum vel globosum.

1. *H. AUTUMNALE* (Linn. spec. 1120), glabrum, caule erecto ramoso polycephalo, foliis lanceolatis serratis longè decurrentibus, ligulis planis 3-5-fidis, pappi paleis ciliato-subfimbriatis in aristam desinentibus cor. disci triplo brevioribus.  $\frac{1}{2}$  in humidis secus rivulos frequens à Canadâ ad Carolinam. Schleich. handb. 3. t. 250. Lam. ill. t. 688. Hook. bot. mag. t. 2994. *Helenia decurva* Mæsch meth. 589. *Helenia autumnalis* Gærtn. — Corn. canad. p. 63. in (v. c. et s. sp.)

*ß. pubescens* (Pers. ench. 2. p. 475), totum brevissimè pubescens, caule erecto ramoso polycephalo, foliis lanceolatis grossè serratis decurrentibus, ligulis planis 3-fidis, pappi paleis basi ciliatis in aristam desinentibus cor. disci triplo brevioribus.  $\frac{1}{2}$  in Amer. bor. *H. pubescens* Ait. bot. ed. 1. v. 3. p. 287? An, suadente cl. Persoon, mera var. hortensâ *H. autumnalis*? An spec. propria? (v. s. c.)

2. *H. MEXICANUM* (H. B. et Kunth nov. gen. am. 4. p. 299?), glabrescens, caule erecto ramoso polycephalo, foliis lineari-lanceolatis elongatis integerrimis decurrentibus, ligulis planis trifidis subtus pubescentibus, pappi paleis scariosis basi ovatis apiculatis cor. disci quintuplo brevioribus.  $\frac{1}{2}$  in Mèxici ditione Oaxaca legit cl. Andrieux (pl. exs. n. 317!). *Achænia nervoso-angulata*, secus angulos pilis albis adpressis vestita. Ligulæ disco breviores. (v. s.)
3. *H. TUBULIFLORUM*, minutè et sparsè puberulum, caule erecto apice ramoso, ramis subnudis apice 1-cephalis, foliis membranaceis elliptico-lanceolatis subdentatis, ligulis tubulosis inæqualiter 5-fidis, pappi squamis lineari-lanceolatis acuminatis scariosis basi ciliatis cor. disci dimidiò brevioribus. — Ex Amer. bor. occid. semina in hortos Anglicos introduxit cl. Douglas. (v. s. in hort. Soc. hort. lect. 1830.)
4. *H. CANALICULATUM* (Lam. journ. h. nat. 2. p. 213. t. 35), glabrescens, caule erecto simplici 1-cephalo, foliis lanceolatis dentatis decurrentibus, ligulis concavo-canaliculatis apice tridentatis, pappi paleis acuminatis cor. disci triplo brevioribus. — in Amer. bor.?

Sect. II. *TETRODUS*. — *Tetrodus Cass. dict. 55. p. 262. Less. syn. 240.*

Cor. disci 4-dentatæ. Pappi paleæ breves. Recept. ovatum aut oblongum.

5. *H. QUADRIDENTATUM* (Labill. act. soc. h. n. Par. 1. p. 22. t. 4), glabrescens, caule erecto apice ramoso polycephalo, foliis decurrentibus oblongo-linearibus, infer. pinnatifidis, super. integerrimis, ligulis glabris, invol. brevissimo, achæniis squamellis vix apiculatis. (I) in Amer. bor. nempe in Carolina (Fras.), Louisiana (Labill.), Mexico (Berl. ! pl. exs. 436). Lam. ill. t. 688 f. 2. Don in Lindl. bot. reg. t. 598. *Rudbeckia alata* Jacq. ic. rar. 3. t. 591. coll. 5. p. 154. *Hel. Milleri* Schult. ex rec. Salz. *Mesodetra alata* Raf. fl. Ind. suppl. p. 141. (v. v. c. et s. sp.)
6. *H. VARIUM* (Schrad. ! hort. Gætt. sem. 1831. et in litt. 1832), glabrum, caule erecto subsimplici sublaxè ramoso, foliis decurrentibus subtrinerviis, caulibus



- integerrimis, infer. . . . , invol. squamis radio brevioribus, ligulis trifidis sub-  
tus vix subpuberulis, pappi squamellis aliis obtusis aliis brevissimè apiculatis.  
① in Mexico (Alam.!), etiā in horto Mexicano (Berlandier! pl. exs. n. 436),  
et Gœttingensi (Schrad.!), cultum. Folia 2-2  $\frac{1}{2}$  poll. longa, basi 4-8 lin. lata.  
Capitula rotunda 5-6 lin. diam. (v. s.)
7. *H. PUBERULUM*, tenuissimè puberulum, caule erecto ramoso polycephalo,  
foliis decurrentibus oblongo-lanceolatis integerrimis, infimis incisis, ligulis  
trifidis pabero-velatinis, invol. brevissimo, pappi squamellis acuminatis. ①?  
in Californiā legit cl. Douglas. Affine *H. quadridentato*, sed majus et ex omni  
parte puberulum. Capitulum globosum 6 lin. diam. Ligulæ parvæ. Pedicelli  
elongati striati. (v. s. comm. ab hon. Soc. hort. London.)
8. *H. MATAPOXYLLUM*, glabrum, caule erecto apice corymboso-ramoso polyce-  
phalo, foliis decurrentibus, inf. obtusè pinnatifidis, mediis sinuatis serratave,  
summis integerrimis acutis, invol. squamis apice subulatis radio ferè longiori-  
bus, ligulis planis 3-fidis, pappi paleis ovatis breviter aristatis obtusiusculis  
cor. disci triplò duplòve brevioribus. ① in Mexico prov. Tamaulipas circa  
Tampico legit cl. Berlandier (pl. exs. n. 107, 190 et 2113). Herba pedalis. Ca-  
pitulum ovato-globosum diam. 3-4 lin. (v. s.)
9. *H. ELZEANS*, glabrum, caule erecto stricto apice corymboso-ramoso oligoce-  
phalo, foliis decurrentibus lanceolato-linearibus integerrimis, utrinquè glan-  
duloso-punctatis, infer. . . . , invol. squamis apice subulatis radio subbrevio-  
ribus, ligulis basi villosis apice obtusè 3-4-dentatis, pappi squamellis brevissimis  
obtusis. ①? in Mexico circa Bejar legit cl. Berlandier (pl. exs. n. 1651!) Li-  
gula ex sicco basi purpureo-croceæ, cæterum luteæ. (v. s.)
10. *H. MICROCEPHALUM*, glabrum, caule erecto ramosissimo polycephalo, foliis  
decurrentibus lanceolatis integerrimis subpunctatis, infer. . . . , invol. brevis-  
simo, ligulis tridentatis glabris, pappo brevissimo obtuso. — in Mexico inter  
Lando et Bejar legit cl. Berlandier (pl. exs. n. 1429!). Ligula flavæ. Capitulum  
subrotundum 3 lin. diam. (v. s.)

† *Species non satis notæ.*

11. *H. LONGIFOLIUM* (Smith. in Rees cycl. n. 2), foliis lineari-lanceolatis inte-  
gris lævissimis, pedunculis nudis elongatis. — in Amer. boreali. *H. autumnale*  
Mill. dict. ed. 8. ex Sm. †.
12. *H. FUMILUM* (Willd. enum. h. Ber. suppl. p. 60), foliis oblongis subinteger-  
rimis, invol. squamis lanceolatis patentibus. ♀ patr. et cæt. car. ign. in h.  
Willd. non reperitur (Link). †.
13. *H. QUADRIPARTITUM* (Link enum. h. Ber. 2. p. 338), foliis lanceolatis de-  
currentibus, cor. radii quadripartitis. ② patr. et cæt. car. ign. An var. *T. qua-*  
*dridentati* seu *Hel. autumnalis* ligulas sæpè quadrifidas gerentis? †.

†† *Species exclusæ.*

- H. LANATUM* Spreng. = *Bahia lanata*.  
*H. OPPOSITIFOLIUM* Spreng. = *Bahia oppositifolia*.  
*H. PUBESCENS* Ait. = *Actinella decurrens*?  
*H. STOCHADIFOLIUM* Spreng. = *Bahia stoechadifolia*.  
*H. TROLLIIFOLIUM* Spreng. = *Bahia trolliifolia*.

CDXIV. AMBLYOLEPIS.

Capitulum multiflorum radiatum, fl. radii fœmineis 1-serialibus  
ligulatis apice 3-fidis, fl. disci hermaphrod. tubo brevi, fauce amplā,  
lobis 5 lanceolatis callosio-apiculatis nervo marginatis. Invol. biseriale,  
squamis ser. ext. circ. 6-7 ovatis acutis foliaceis adpressis disco æqua-  
libus, ser. int. obovato-rotundis obtusissimis hyalinis enerviis. Recept.

planiusculum alveolatum. Antheræ longè et acutè appendiculatæ. Styli rami exappendiculati. Achænium turbinaum adpressè villosissimum. Pappi paleæ 5 uniseriales obtusissimæ enerviæ corollæ tubo æquales. — Herba Mexicana annua erecta simplex, hinc indè secus caulem margines foliorum et squamarum invol. exteriorum pilis longis ciliatopilosa. Folia alterna semiamplexicaulia nec decurrentia ovali-lanceolata subintegerrima. Capitula terminalia solitaria flava, ferè Pyrethri Myrenis magnit. et facie. — Nomen (ex *αμβλυνε* obtusa et *λεπτις* squama) alludit ad squamas invol. exteriores et ad paleas pappi obtusas.

1. *A. SETIGERA*. (J) in Mexicanà prov. Texas inter. Bejar et Austin legit cl. Berlandier (pl. exs. n. 1613!). Herba pedalis paucifolia. Folia inf. obtusa, sup. acuminata. (v. s.)

## CDXV. ARGYROXIPHIMUM.

Capitulum multiflorum heterogamum, fl. radii 1-serialibus ligulatis fœmineis, disci hermaphrod. tubulosis 3-dent. Recept. nudum planum. Invol. campanulatum, squamis 2-3-seriatis lanceolato-linearibus subæqualibus disco paulò brevioribus. Ligulæ obovato-cuneatæ apice late incisæ. Styli stigm. 2 divergenti-revoluta subfiliformia, in disco apice pilosiuscula. Antheræ ecaudatæ apice appendiculatæ. Achænium elongatum glabrum ad angulos 2 vix subciliatum compressum, latere uno alterove nervo seu angulo subexserto instructum (in specim. mihi obl. cassum). Pappus persistens 1-serialis paleaceus radii auricularis ad latus exterius situs acuminatus integer, disci paleis 2-5 irregularibus serratis subfoliaceis rigidis. — Herba Sandwicensis crassicaulis Tussilagines, Cinerarias seu Ligularias Europæas texturâ referens. Folia ab e basi subrigida elongata subensiformia integerrima villis sericeis adpressis copiosissimis argentea, alia basi teneriora virescentia pube glandulosâ instructa. Involucra etiâ hirta et fol. supremis texturâ similia. Pedunculi ferè ex axillis foliorum orti solitarii glanduloso-hirti subaphylli 1-cephali. Capitula subglobosa nutantia.

1. *A. SANDWICENSE*. ♀ in insulis Sandwicensibus ad Owwhyhee floridam mense junio legit cl. Macrae. Herba mirabilis et distinctissima circ. bipedalis. Folia argyræa pedem longa, 3-4 lin. lata. Capitula ferè diam. pollic., ex siccis fortè purpurascentia. (v. s. in herb. cl. et am. Lindley.)

## CDXVI. ROSILLA Less. syn. 245.

Capitulum radiatum, ligulis fœmineis. Invol. cylindræci squamis scariosæ, exter. sublineares. Recept. epaleaceum. Styli rami appendiculati. Achænium obcompressum basin versus angustatum. Pappi paleis plumoso-fimbriatis. — Herba annua Mexicana, floribus lutes.

1. *R. LUTEA* (Less. l. c.). (J) in Mexico. Cœt. ign.

Div. II. GALINSOGÆ. — Galinsogæ et Caleinæ *Cass. dict.* 55 p. 264 et 265. *Less. syn.* 241 et 245.

Pappus 1-serialis, paleis integris aut plumosis constans, rarissimè nullis. Recept. paleaceum. Invol. squamæ planiusculæ, achænia radii non involvens

## Subdiv. I. EUGALINSOGÆ.

Ligulæ fœmineæ seu nullæ. Pappi squamellæ planiusculæ nec activate contortæ.

KVII. *LEMMATTUM*. — *Caleacte Less. in linnæa* 1830. p. 158. *syn.* 248. *non Brown.*

Capitulum 6-7 florum heterogamum, fl. radii circ. 3 ligulatis foemina, disci etiā 3-4 tubulosis hermaphrod. 5-fidis, lobis longiusculis glabris. Invol. cylindricum, squamis imbricatis oblongis, exter. foliaceis, inter. membranaceis. Recept. angustum, paleis convolutis tri-oblongis flores involventibus et eos æquantibus. Styli rami exendiculati. Achænia trigono-prismatica ad angulos villosa. Pappus membranaceus vaginans apice dentatus, squamellis concretis verisim. stans, cor. faucem adæquans. — Frutex Brasiliensis. Caulis densè et viter hirsutus teres seu obtusè angulatus. Folia opposita sessilia subunda basi cordata crenato-dentata coriacea, suprā puberula, subtus aut quintuplinervia reticulata villosa-scabra. Capitula subsessilia in umbellum amplum valdè compositum fastigiatum digesta, bracteis ovatis foliaceis instructa. Flores lutei. — Genus habitu et stigmatibus ianthæ affine sed achæniis et corollæ lobis diversum, et potius Cærmo et Caleæ affine, ab omnibus Helenieis pappi squamellis in vānam concretis distinctum, sed tamen meo sensu iis potius quāmransanthemeis adjungendum. Nomen ex λεμματιον vaginula.

*L. ROTUNDIFOLIUM*.  $\frac{5}{2}$  in Brasiliæ prov. Minarum-Generalium legit cl. Vauhier (pl. exs. n. 284). *Caleacte rotundifolia Less. ll. cc.* Folia 2  $\frac{1}{2}$  poll. diam. (v. s.)

KVIII. *CALYDERMOS* *Lag. nov. gen. p. 24.* — *Calydermos* et *Calebrachys Cass. dict. 55. p. 265 et 277. Less. syn. p. 225 et 241.* — *Caleæ* sp. *Kunth.*

Capitulum multiflorum homogamum. Invol. oblongum imbricatum, squamis subscariosis obtusissimis. Recept. angustum conicum, paleis caribus acutis. Cor. tubulosæ 5-fidæ. Styli rami cono terminati. hænium 4-5-angulatum pilosum. Pappi paleæ 1-seriales lineares oblongæ 1-nerviæ aut nullæ. — Herba Mexicanæ erectæ scabræ aut pilosæ vix ramosæ. Folia opposita sessilia ovata aut elliptico-acuminata serrata trinervia. Rami longè aphylli, apice breviter trifidi, caula fasciculato-corymbosa gerentes. Invol. sublutescentia. Flores si.

St. I. *EUCALYDERMOS*. — *Calydermos Lag. Cass. et Less. ll. cc.* Achænia calva.

*C. SCABER* (Lag. nov. gen. et sp. n. 323), caule hispidulo, foliis ovato-lanceolatis utrinquè scaberrimis subtus præcipuè villosis, achæniis calvis.  $\frac{2}{2}$  in flexici montanis circa Guanaxuato, Acapulco et Real del Monte (Lag.), Vilalpando (Mend.), et urbem Mexici (Berl.) pl. exs. n. 646). Herba pedalis et esquipedalis. An fortè Caleæ peduncularis var. epapposa H. B. et Kunth 4. 1. 296. t. 408. f. 5 huc referenda? (v. s.)

*C. LONGIFOLIUS* (Lag. nov. gen. et sp. n. 324), caule subpiloso, foliis oblongo-lanceolatis lævisculis.  $\frac{2}{2}$  in Mexico. Variat foliis oppositis et ternato-verticillatis.

Sect. II. *CALEBRACHYS*. — *Calebrachys Cass. Less. ll. cc.*

Achænia pappo 1-seriali paleis 10-12 oblongo-linearibus obtusis coronato.

*C. PEDUNCULARIS*, caule hirsuto, foliis inf. brevissimè petiolatis, sup. sessilibus ovato aut oblongo-lanceolatis utrinquè pubescenti-hirsutis, achænio paleis 8-10

hermaphr. rami exappendiculati. Achænium teres compressum angulatumve. Pappus multipaleaceus, paleis 5-20 sæpius circ. 10 linearilanceolatis scariosis acuminatis uninerviis subæqualibus. — Frutices (rarius herbæ) ex Americâ æquinoctiali indigeni. Folia opposita, rarissimè terna, sæpius 3-nervia, scabra integra aut dentata. Capitula terminalia aut axillaria. Flores lutei.

## Sect. I. AMPHICALÆA.

Capitula discoidea 4-flora. Receptaculum epaleaceum. — An genus proprium? sed habitus valdè ad *C. cymosam* accedit. Fortè flores omninò marginales et idè paleæ ut flores centrales ipsi deficientes?

1. *C. GENTIANOIDES*, erecta herbacea glaberrima, caulibus teretibus striatis simplicibus, foliis elliptico-oblongis sessilibus basi attenuatis apice obtusissimis integerrimis prope basin triplinerviis, pedunculis oppositè corymboso-pauculatis capitula 3 pedicellata discoidea 4-flora gerentibus, invol. squamis oblongis adpressis, recept. nudo! pappi paleis angustis acuminatis. ? in Brasiliæ prov. Sancti-Pauli. Herba gentianoidea circ. pedalis. Inv. squamæ ext. parvæ, int. submembranaceæ multò majores. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 406.)

Sect. II. DISCOCALEA. — *Calea Brown l. c. Cass. dict. 55. p. 265.*

Capitulum homogamum discoideum multiflorum.

2. *C. CORNIFOLIA* (Swartz fl. ind. occ. 3. p. 1326), fruticosa, ramis oppositis divergentibus teretibus glabris, foliis petiolatis cordatis acuminatis serratis trinerviis scabris, corymbis terminalibus et axill. 3-8-cephalis, capitulis ovatis discoideis 16-18-floris, invol. squamis membranaceis obtusis, bracteis lanceolatis acutis, achæniis compressis, pappi paleis circ. 20 striato-penninerviis lanceolatis cor. disci æqualibus. ? in frutetis montium Jamaicæ australiore. Flores luteo-purpurascens.
3. *C. JAMAICENSIS* (Linn. sp. 1179), fruticosa, ramis teretibus laxis pubescentibus, foliis petiolatis ovatis acuminatis subserratis trinerviis suprâ scabris subis hirtis, capitulis pedicellatis ad apices ramulorum ternis ovatis discoideis, invol. squamis ovatis extus hirtis, paleis lanceolatis, pappi paleis serratis. ? in Jamaicâ. Swartz fl. ind. occid. 1327. obs. 302. Santolina Jamaicensis Linn. amœn. 5. p. 404. — Sloan. jam. 1. p. 257. t. 151. f. 3.
4. *C. BERBERIANA*, suffruticosa? ramis teretibus hirtellis demùm glabrescens, foliis petiolatis ovatis acutis grossè dentatis subtrinerviis suprâ sparè pubes subitis densè villosis-hirtis, capitulis pedicellatis ad apices ramorum umbellatis discoideis ovatis, invol. squamis ext. acutis subfoliaceis, int. obtusis submembranaceis, achænio puberulo squamellis circ. 20 linearibus acuminatis, receptaculo planiusculo. ? in Amer. austr. ad Sanctam-Martham legit Bertero. Marchallia australis Spreng. ! in h. Balb. 1822 sed in ejus oper. deacid. Valdè affinis *C. pinnifolia*. (v. s.)
5. *C. ZACATECHICHI* (Schlect. in linnæâ 1834. p. 589), canle fruticoso tereti multirichotomo, foliis petiolatis ovatis acutis subtripplinerviis grossè crenatis supra petiolisque hispidulis, ramulisque pubescentibus, capitul. 12-floris in cymis irregulares dispositis, invol. cylindracei squamis concavis obtusis margine scarioso-undulatis, achænio turbinato, pappo brevi. ? in collibus apricis Mex. prope Hacienda de La Laguna et Jalapam. Vulg. Mexic. dicitur *Zacatechicli*.
6. *C. PRUXIPOLIA* (H. B. et Kunth nov. gen. am. 4. p. 294. t. 406), fruticosa, ramis junioribus subangulatis hirtis-scabris, foliis petiolatis ovatis remote crenato-serratis utrinquè scabris, capitulis longè pedicellatis ad apices ramorum umbellatis ovatis discoideis, invol. squamis ext. acutis, interioribus bracteisque obtusis, achænio compresso-tetragono hispido, pappi paleis linearibus

acuminatis achænio paulò longioribus.  $\frac{3}{4}$  in ripâ fluminis Magdalena prope urbem Honda. Fl. centrales masculi aut steriles. Recept. oblongum.

7. *C. TRIPHYLLA* (H. B. et Kunth nov. gen. am. 4. p. 294), fruticosa, ramis subteretibus hirtis-pubescentibus, foliis ternis verticillatis petiolatis ovatis acutis basi angustatis serratis suprâ scabris subtus hirtis, corymbis in apice ramulorum sessilibus trichotomis, capitulis discoideis circ. 7-floris, invol. campanulati squamis 10 ovatis, extimis minutis hirtellis, bracteolis oblongis, achænio hispidio, pappi paleis lanceolatis margine fimbriatis.  $\frac{3}{4}$  in ripâ flum. Magdalena prope Honda.
8. *C. SOLIDAGINEA* (H. B. et Kunth nov. gen. am. 4. p. 295. t. 407), suffruticosa, ramulis pubescentibus demum glabris, foliis petiolatis ovato-oblongis acuminatis basi angustatis obsolete serratis suprâ scabris subtus pubescentibus, capitulis breviter pedicellatis ovatis discoideis ad apices ramorum et ramulorum axillarum umbellatis, invol. squamis ovatis subacutis, bracteolis ovato-oblongis acutis, achænio compresso-tetragono apice subhirsuto, pappi paleis lineari-subulatis corollas disci æquantibus.  $\frac{3}{4}$  in declivitate montis Silla de Caracas.
9. *C. PEDUNCULOSA*, caule basi suffruticoso demum glabro, ramis dense hirtis basi foliosis apice in pedicellum elongatum 1-cephalum puberulum productis, foliis sessilibus latè ovatis acutis basi obtusis grosse serratis utrinquè hirtellis et impresso-punctatis, invol. campanulati squamis obtusissimis, exter. foliaceis dorso subpuberulis, inter. membranaceis lineato-striatis, achænio angulato villosio, pappi squamellis lineari-lanceolatis fimbriatis acuminatis achænio longioribus.  $\frac{2}{4}$   $\frac{3}{4}$  in Brasiliæ prov. Rio-Grande. Habitus *C. oligocephalæ*. Recept. paleæ paucæ lineares complicatæ acuminatæ. (v. s. in h. Mus. reg. Par. ex h. Mus. imp. Bras. sub n. 1050 miss.)
10. *C. SENSILIFLORA* (Less. in Linnæâ 1830. p. 158), fruticosa..., foliis basi obtusissimis et subcordatis, capitulis discoideis cylindræcis circ. 10-floris..., invol..., bracteolis muticis.  $\frac{3}{4}$  in Mexico. *Chrysosphaerium gnaphalioides* Willd. herb. Cæc. ign. †.
11. *C. SERRATA* (Less. in Linnæâ 1830. p. 158), caule herbaceo..., foliis subtus hirsuto-pubescentibus, capitulis discoideis plurimifloris..., bracteolis muticis.  $\frac{2}{4}$  in Brasiliâ. Cæc. ign. †.

Sect. II. CALZACTE. — *Calezacte Brown l. c. Cass. dict. 55. p. 265 non Less.*

Capitulum radiatum. Fl. radii 5-10. — Invol. squamæ siccae nec exteriores foliaceæ bractæiformes.

12. *C. AXILLARIS*, ramulis teretibus petiolisque subhirsutis, foliis petiolatis ovato-lanceolatis basi breviter attenuatis apice longè acuminatis grosse serratis suprâ vix scabris subtus villosio-hirsutis, capitulis pedicellatis radiatis ex axillis apicibusque plurimis ortis subumbellatis, invol. squamis siccis, exter. brevioribus acutis, inter. obtusis, bracteolis scariosis oblongis aristatis disco brevioribus, achænio subangulato apice hispidulo, pappi paleis numerosis linearibus acuminatis cor. disci paulò brevioribus.  $\frac{3}{4}$  in Amer. calid. (Née!), in Mexico (Hänke). Spec. *C. solidagineæ* affinis. *Mocinna serrata* Lag. nov. gen. p. 31. n. 384 quæ *Galinsogea serrata* Spreng. syst. 3. p. 579, à Nemo in Mexico circa Salmanticam urbem lecta, videtur nostra. (v. s.)
13. *C. VERBENÆFOLIA*, ramis teretibus pubescenti-hirtis, foliis petiolatis ovatis acutis serratis, suprâ sparè puberulis, subtus trinerviis reticulatis sparè et præcipuè in nervis puberulo-scabris, capitulis breviter pedicellatis ad apices ramulorum dispositis in thyrsum terminalem digestis ovatis 5-radiatis, invol. squamis siccis obtusis, exter. subacutis, bracteolis linearibus obtusis, pappi paleis 3-4 deciduis, achænio lineari glabro circ. æqualibus.  $\frac{3}{4}$  in Amer. calid. legit cl. Née, sed locus propr. in herb. Thibaud desideratur. (v. s.)
14. *C. BRACHIATA*, fruticosa, ramis brachiatis, foliis ovatis obtusè mucronatis, capitulis fasciculatis plerisque terminalibus.  $\frac{3}{4}$  in Panamæide. *Mocinna bra-*

C. PINIFOLIA Forst. = *Oenothermus pinifolius*.

C. SCOPARIA Linn. = *Baccharis scoparia*.

C. SPECTABILIS Labill. = *Apalochlamys spectabilis*.

CDXXII. ALLOCARPUS H. B. et Kunth nov. gen. am. 4. p. 291. t. 405. Cal. dict. 55. p. 265. Less. syn. 242. — *Alloispermum Willd. mag. naturf. Ber.* 1807. p. 139. ex Cass.

Capitulum multiflorum radiatum, fl. radii circ. 5 ligulatis fœmineis. disci hermaphr. tubulosis 5-dentatis. Invol. hemisphærici squamæ lute imbricatæ, circ. 10 scarioso-membranacem. Recept. subplanum paleis lanceolatis persistentibus scariosis onustum. Styl. fl. hermaphrod. rami exappendiculati. Achænia radii cuneata compressa calva, disci cuneato-cylindracea pappo 1-seriali multipaleaceo coronata. Paleæ pappi acuminatæ, ex icon. pectinatæ ex descr. integræ. — Herba Americana ramosa. Folia opposita breviter petiolata, tripli- seu 5-7-nerviâ, ovalia utrinquè acuminata integra villosa. Capitula pedicellu flava aut fortè albida.

1. A. CARACASANA (H. B. et Kunth l. c. t. 405), caule erecto hirsuto, ramis spinoso-corymbiferis, foliis ovali-lanceolatis acuminatis hinc inde dentatis triplicinerviis villosis. 2? in temperatis propè urbem Caracas. *Alloispermum divaricatum* Willd. l. c. *Galinsoga allocarpa* Spreng. syst. 3. p. 579. Pappi paleæ margine fimbriatæ.
2. A. INTERFOLIUS, caule herbaceo tereti glabro virgato, foliis breviter petiolatis ovali-lanceolatis acuminatissimis integerrimis 5-7-plinerviis utrinquè strigillis brevibus apertis scabridis, corymbo composito polycephalo. — inter plantas Mexicanas à cl. Karvinski lectas video in h. aed. reg. Monac. sed locus non indicatur. Herba fortè subscandens? Folia 5 pollic. longa, pollic. lata, membranacea subnitida. Capitula fortè albida, dimidiò minorâ quàm prioris, circ. 15-flora. Pappi paleæ vix margine scabræ. (v. s. in h. cit.)
3. A. SABAZIOLDES (Less. syn. 242 et in linnæâ 1834. p. 690), herbacea procumbens monocephala, foliis petiolatis ovatis acutis crenatis triplicinerviis hinc bracteolis scariosis acuminatissimis bidentatis. — in Mexico prope San-Miguel del Soldado et La Joya.

#### CDXXIII. VARGASIA.

Capitulum heterogamum, fl. radii ligulatis paucis 1-serialibus inter invol. squamas et rec. paleas ext. et majores occultatis, disci tubulosi 5-dentatis. Recept. convexum paleaceum, paleis membranaceis angustissimis, ext. majoribus. Invol. hemisphæricum, squamis subbiseriatis ovalibus mucronulatis. Ligulæ breves obovatæ apice tridentatæ. Stylus brevis inclusus, ramis cono brevi terminatis? Achænia rotunda obcompressa inter squamam int. et paleam ext. inclusa, disci subtrigona, omnia basi attenuata subpuberula. Pappi paleæ variegatæ, setiformes parvæ, disci 5 ovatæ erectæ serrato-ciliatæ acutæ coram triplo longiores. — Herba Austr.-Americana æquinoctialis annua erecta gracilis, parvè dichotomè ramosa, sparsè hispida. Folia opposita breviter petiolata ovata acuminata dentato-serrata. Pedunculi è dichotomiis orti, 1-cephali ebracteati. Capitula subglobosa, invol. glabro. Flores pallidè purpurascentes, à *Galinsoga* differt ferè ut *Allocarpus* à *Calea*. — Genus affine *Galinsoga* dixi in honorem cl. doct. Vargas Caracasanum rerum naturalium sagacis observatoris.

1. V. CARACASANA. (1) ad Caracas legit cl. Vargas. (v. s. comm. à cl. interve-

CDXXXIV. GALINSOGA *R. et Pav. fl. per. prod.* (1794) p. 110. t. 24. *Cass. dict.* 18. p. 96. — *Galinsoga* Zucc. in *florâ* 1821. p. 612. — *Wiborgia* Roth cat. 2. (1800) p. 112. *Less. syn.* 245 non *Spreng. Thunb. Mönch* (1). — *Galinsoga* spec. *Cav. Willd.*

Capitulum pluriflorum radiatum, ligulis 4-5 fœmineis suborbiculatis, fl. disci hermaphr. tubulosi 5-dent. Invol. 1-seriale, squamis 5 ovatis margine hyalinis. Recept. conicum paleis lineari-lanceolatis acutis integris trifidisve onustum. Styli rami acuti vix appendiculati. Achænium angulatum adpressè hispidulum. Pappus 1-serialis, paleis oblongis brevibus plumoso-fimbriatis. — *Herbæ* Americanæ annuæ trichotomoramosissimæ. Folia opposita ovata triplinervia serrata. Pedunculi axillares et terminales 1-3-cephali. Capitula parva, disco luteo, radio albo.

1. *G. PARVIFLORA* (Cav. ic. 3. p. 41. t. 281), glabrescens, receptaculo conico, pappi paleis 8-16. ① in Peruvia uliginosis et ruderatis (*R. et Pav. Cav.*), in Chili ad Coquimbo (*Hook. et Arn.*), in Novâ-Granatâ circa Ybague (*H. B.*), Mexico (*Keerl.*), nunc circa Erlang, etc., seminibus ex hort. bot. egressis quasi spontanea (*Zucc.*). *Gal. quinqueradiata* *R. et Pav. syst.* 198. *Wiborgia Acmella* Roth cat. 2. p. 112. — *Veill. per.* 2. p. 744. t. 32. Vulgò dicitur Peruvianis *Paica-Jullo* (*Feuill.*) seu *Pacoyuyu-Fino* (*R. et Pav.*). Vulneraria et antiscorbutica habetur. (v. v. c.)

fl. ligulis 3-4. — *Galinsoga quadriradiata* *R. et Pav. syst.* 198. — in Peruvia ad Limam et Chanoy. Vulgò *Pacoyuyu-Cimarron* ex *R. et P.* Vidi in hortis specim. 4-ligulata certè à var. *a* non diversa. (v. v. c.)

γ. *hispida*, caule foliisque sparse hispidis, ligulis 5. ① in Mexico circa urbem (*Berl.* pl. exs. n. 615), in Chilensibus montibus (*H. Hænke!*). Caulis crassior. Folia majora. Pedunculi abbreviati. (v. s.)

2. *G. ? OBLONGIFOLIA* (*Hook. misc. bot.* 2. p. 226 sub *Wiborgia*), tota pilis adpressis albis aspera, foliis oblongo-lanceolatis triplinerviis grossè serratis basi attenuatis subsessilibus, invol. polyphylo, rec. paleis linearibus hispidis, pappo piloso brevissimo. — ad Luriam prope Limam legit *cl. Cruikshanks*. Pedicelli binuclei. Ligulis bidentatis. Videtur præsertim ex pappo à genere exclud. et novam varietatem constituendum.

+ *Species viz notæ.*

3. *G. ? ANGUSTIFOLIA* (*Spreng. neu. entd.* 2. p. 138. *syst.* 3. p. 590), caule basi suffruticoso hirsuto, ramis glabriusculis angulatis, foliis oppositis lineari-lanceolatis dentatis margine revolutis, infer. hirsutis, super. glabris integris, pedicellis terminalibus 1-cephalis, invol. squamis subrotundis coloratis, ligulis crenatis, pappo squamellato. 3 in Brasiliâ. Flores lutei.

4. *G. ? MEXICOTAMICA* (*Spreng. syst.* 3. p. 580), herbacea glabra, foliis linearibus tripartitis integerrimis, ramis ternis 1-floris rigidis. — in Brasiliâ ad Rio-Grande. Cæt. ign.

5. *G. ? UNIFLORA* (*Spreng. syst.* 3. p. 580), caule herbaceo simplicissimo 1-cephalo supernè hirsutulo, foliis sparsis linearibus mucronatis margine scabris. — ad Monte-Video. Cæt. ign.

6. *G. ? RESINOSA* (*Hook. et Arn. bot. Beech.* 1. p. 32), caule fruticoso angulato ramoso, ramulis foliisque angustissimè linearibus resinoso-glandulosis, capitulis paniculato-corymbosis, invol. imbricato inæquali viasoso, recept. nudo. 5 in Chili ad Coquimbo et Valparaiso. Species ex recept. nudo potius Bahiæ quàm *Galinsogæ* referenda videtur. Valdè dubia nec satis descr. †.

(1) *Wiborgia* Thunb. seu *Viborgia* Spreng. et DC. est genus in prod. jam inter Leguminosas admissum (2. p. 135). *Viborgia* Mönch (1794) est sect. Cytisi mihi *Tubocytisus*. *Viborgia* Ort. (1797) est *Varennea* DC. prod. 2. p. 522.

++ *Species exclusæ.*

- G. ALLOCARPA Spreng. = *Allocarpus* Caracasensis.  
 G. BALBISIOIDES H. B. K. = *Sogalgina* balbisioides.  
 G. BRACHIATA Spreng. = *Calea* brachiata.  
 G. RESINOSA Hook. et Arn. = *Bahia* resinosa.  
 G. SERRATA Spreng. = *Calea* axillaris.  
 G. TRIFIDA Pers. = *Sogalgina* trilobata.  
 G. TRILOBATA Cav. = *Sogalgina* trilobata.

CDXXV. SOGALGINA Cass. bull. philom. 1818. p. 31. dict. 49. p. 397 et 55. p. 265. — *Galinsogæ* sp. Cav. — *Galinsogæ* et *Sogalgina* Less. syn. 245. — *Galinsogæ* Kunth nov. gen. am. 4. p. 253 non R. et Pav. — *Sogalgina* Send.

Capitulum multiflorum radiatum; fl. radii fœmineis 1-serialibus bilabiatis, labio ext. amplo apice dentato, int. lobulis 2 linearibus nunc discretis nunc concretis, fl. disci hermaphrod. tubo brevi, fauce cylindricâ, limbo 5-dentato. Recept. hemisphæricum pauciseriale, squamis submembranaceis ovalibus obtusissimis. Recept. convexum paleis ovali-acuminatis flores amplectentibus onustum. Styli rami appendiculo filiformi longo terminati. Achænium teretiusculum. Pappi paleæ 1-seriales conformes pinnatipartitæ seu plumoso-fimbriatæ. — Herbe Mexicanæ glabræ aut pilosiusculæ trichotomæ. Folia opposita lateraliter incisa. Capitula pedunculata terminalia. Flores lutei. — Nomen anagramma *Galinsogæ*.

1. S. TRILOBATA (Cass. dict. 49. p. 397), ligulis cuneatis labio inter. bipartito donatis, pappi paleis achænio brevioribus. (1) in Mexico. *Galinsoga* trilobus Cav. ic. 3. p. 42. t. 282. Spreng. nov. prov. p. 19. Less. syn. 245. Variat foliis vel trilobatis aut pinnato-incis, caule foliisque glabriusculis aut pilosis, pedunculis apice pubescentibus aut glanduloso-puberis, achænio glaberrimo (*Galinsoga* trifida Pers. ench. 2. p. 472) puberulo aut hirtulo, pappi paleis brevissimis vix pinnatilobatis aut longioribus plumoso-fimbriatis, aequantibus sed semper brevissimis. An plures forsan species hic confusæ? (r. v. c. et s. sp.)
2. S. BALBISIOIDES (Cass. dict. 49. p. 398), ligulis latissimis subreniformibus labio inter. indiviso donatis, pappi paleis achænio longioribus evidenter plumosis. (1) in temperatis Mexici inter Valle de Santiago et lacum Palango. *Galinsoga* balbisioides H. B. et Kunth nov. gen. am. 4. p. 253. t. 386. *Ptilostephium* balbisioides Spreng. syst. 3. p. 570. *Sogalgina* Less. syn. 245. Herba pilosa subhirta. Folia ovato-lanceolata inciso-serrata aut integra. Achænium juncatum pubescenti-sericeum.

CDXXVI. PTILOSTEPHIUM H. B. et Kunth nov. gen. 4. p. 255. — *Ptilostephium* et *Carphostephium* Cass. dict. 44. p. 62. Less. syn. 246.

Capitulum homogamum. Cor. fl. interiorum regulares, tubo brevissimo, fauce tumidiore elongatâ, dentibus 5 regularibus, fl. ext. majores inæqualiter 5-lobæ, lobis ext. majoribus. Invol. paucifolium imbricatum campanulatum, squamis margine scariosis. Recept. planum paleis oblongo-linearibus scariosis onustum. Styli rami breviter appendiculati. Achænium villosum basi attenuatum. Pappus 1-serialis, paleis 15-20 ciliato-plumosis. — Herbe Mexicanæ annuæ trichotomæ pilosiusculæ. Folia opposita irregulariter palmati- aut pinnatilobata, lobis



linearibus acutis. Capitula longè pedunculata. Cor. luteæ. Species duæ habitu similimæ nec meo sensu ritè separandæ.

1. *P. TRIFIDUM* (H. B. et Kunth l. c. t. 388), invol. squamis exter. plurimis emarginatis, pappi squamis membranaceis ciliato-fimbriatis. (I) in Mexico inter Guanajuato et Valladolid ad lacum Cuiseo (H. et R.). *Carphostephium trifidum* Cass. dict. 44. p. 63. — Specimina habeo circa Mexico à cl. Berlandier (pl. exs. n. 768) lecta, iconi Kunthianæ involucri et pappi similia, sed folia alia trifida alia subpinnatifida ut in sequente. An tertia species? (v. s.)
2. *P. coronopifolium* (H. B. et Kunth l. c. t. 387), invol. squamis exter. obtusis mucronatisve, pappi paleis medio nervigeris margine plumosis. (I) in convalli Mexicanâ. Folia variant pinnatifida et trifida. Invol. squamas nunquàm verè acuminatas sed hiuc inde mucronatas et æpius obtusas video. (v. s.)

CDXXVII. *TRIDAX* Linn. hort. Cliff. 418. gen. n. 972. Brown trans. Linn. soc. 12. p. 103. Less. syn. 246. — *Balsisia Willd. sp. 3. p. 2214. Cass. dict. 3. suppl. 169 et 55. p. 262 et 265 non DC. — Bartolina Adans. fam. 2. p. 124 (ex syn. Linn.) non Brown.*

Capitulum multiflorum radiatum, fl. radii fœmineis ligulatis, ligulâ ovatâ trilobatâ, disci tubulosis 5-dent. hermaphr. Invol. ovati squamæ laxè imbricatæ, exter. 6-8 ovales foliaceæ dorso villosæ, inter. membranaceæ glaberrimæ paleas referentes sed latiores. Recept. paleaceum, paleis lauceolato-linearibus. Styli fl. hermaphr. rami appendiculo filiformi. Achænia villosa, disci obpyramidata, radii obovata obcompressa. Pappus 1-serialis, paleis 15-16 pinnatisectis (seu plumosis axi rigidulâ), radii minoribus. — Herba Americana procumbens hirsuta ramosa. Folia opposita petiolata ovata inciso-dentata. Rami apice nudi valdè elongati 1-cephali. Capit. disco luteo, radio pallidè flavido.

1. *T. PROCUMBENS* (Linn. sp. ed. 1. p. 900). (I) in Mexico (Ort.), Verâ-Cruce (Linn.), Sanctâ-Marthâ (Bert.), Cubâ (La Sagra!). Etiam reperitur in insulâ Martinii (ex specim. à cl. Bojer et Bouton missis) et in Indiâ or. (Wall. cat. n. 307) sed ex hortis migrata! *Balsisia elongata* Willd. spec. 3. p. 2214. *Balb. pedunculata* Hoffmans. verz. 1824. p. 228. *Amellus pedunculatus* Ort. ex Willd. *Balsisia divaricata* Cass. ann. sc. nat. 1829. dict. sc. nat. v. 60 et opusc. 3. p. 91 certè ex specim. Bojer. et Bout. qui plantam ad Cassin. olim miserunt. Caulis plùs minus hirtus. (v. v. c. et s. sp.)

*B. canescens*, caule hirsuto subtomentoso ferè incano. — ad Sanctam-Martham confin. Amer. austr. *Balsisia canescens* Pers. ench. 2. p. 469. (v. s. comm. à cl. Balbis.)

CDXXVIII. *BLEPHARIPAPPUS* Hook. fl. bor. am. 1. p. 316.

Capitulum multiflorum radiatum, ligulis paucis fœmineis lato-cuneatis profundè trifidis, fl. disci hermaphr. 5-dent. Invol. hemisphærici squamæ 1-seriales æquales oblongæ seu ellipticæ. Recept. paleaceum. Corollæ apice longè pilosæ. Styli rami in fl. hermaphr. lineares longè recurvi glanduloso-pubescentes, in radio brevissimi. Achænia clavata. Pappi paleæ circ. 8 lineari-acuminatæ fimbriatæ ciliatæ. — Herba annuæ ramosæ Boreali-Americanae. Folia alterna linearia integra. Capitula terminalia solitaria, disco flavo, radio albo.

1. *B. GLANDULOSUS* (Hook. l. c.), hispidus, foliis lato-linearibus, superioribus involucrique glandulosis, glandulis nigris pedicellatis, pappi paleis elongatis basi longè pilosis, radii achæniis nudis calvis. (I) in arenosis planis Columbiæ Amer. bor.-occid.

2. *B. SCABER* (Hook. l. c.), foliis angusto-linearibus scabris, involocris subbescentibus, glandulis pedicellatis nullis, achæniis omnibus villosis, paleis pappi breviusculis supernè præcipuè fimbriato-ciliatis. ① hab. cum. priore.

CDXXXIX. *MARSHALLIA* Schreb. *gen.* (1791), n. 1762. *Cass. dict.* 55. p. 261. *Less. syn.* 241. — *Persoonia* Michx. fl. bor. am. 2. p. 104 non Sm. — *Phytomopals* Juss. *ined. Michx. l. c. icon.* — *Trattenickia* Pers. *ench.* n. 1825. n. Willd. — *Athanasia* Walt. fl. car. 201. non Linn.

Capitulum homogamum. Invol. 1-2-serialis squamæ linearilanceolatæ disco æquales. Recept. convexum paleis linearibus acutis onustum. Cor. tubulosæ 5-dentatæ extus hirsutæ. Styli rami exappendiculati. Achænium turbinato-subquingangulum pubescens seu villosum. Pappi paleæ 5-ennerviæ ellipticæ acuminatæ integræ. — Herba Brasiliæ Americanæ perennes. Folia alterna integra. Capitula terminalia scitaria purpurascentia non malè Scabiosarum capit. referentia. — A satis ab hoc genere differt *Therolepta* Raf. nov. *gen.* 1825. p. 5?

1. *M. CÆSPITOSA* (Nutt. ! in litt. 1825), glabra cæspitosa, caule simplici apice 1-cephalo, foliis elongato- et lato-linearibus subobtusis integerrimis, inter squamis oblongo-linearibus obtusis, paleis recept. linearibus. ½ in America ad Red-River legit cl. auctor. Habitus ferè *Armeriæ*. Caulis 8-10-poll. erectus striatus. Cor. ex sicco albæ, fortè subrosæ. Achænium villosum. (v. s.)
2. *M. LANCEOLATA* (Pursh fl. bor. Am. 2. p. 519), caule simplici inferè hirsuto supernè nudato, foliis glabris, radicalibus petiolatis obovatis triplinerviis, caulinis sessilibus oblongo-lanceolatis, invol. squamis ovali-oblongis obtusis. Recept. paleis lineari-spathulatis. ½ in Carolina montanis. Ell. sketch 2. p. 33. *Persoonia lanceolata* Michx. fl. bor. am. 2. p. 105. *Trattenickia lanceolata* Pers. *ench.* 2. p. 403. *Athanasia obovata* Walt. car. p. 201. (v. s.)
3. *M. LATIFOLIA* (Pursh l. c.), caule simplici, foliis glabris lanceolato-oblongis acuminatis trinerviis sessilibus, infimis vaginantibus, invol. squamis recept. paleis angusto-linearibus. ½ in Carolina montanis. *Persoonia latifolia* Michx. fl. bor. amer. 2. p. 105. t. 43. *Athanasia trinervia* Walt. car. 2. *Trattenickia latifolia* Pers. l. c. *Marshallia Schreberi* Tratt. arch. gew. 1. p. 123. cum icon. Michx. mut.
4. *M. ANGUSTIFOLIA* (Pursh l. c. p. 520), caule ramoso vix ad apices pubescenti foliis infer. angusto-lanceolatis, super. linearibus obscure 3-nerviis glabris, invol. squamis acuminatis rigidulis, recept. paleis linearibus. ½ in Tennessee, Missouri in pinetis humidis Carolinae et Georgiae (Ell. Fras.!). *Persoonia angustifolia* Michx. l. c. *Trattenickia angustifolia* Pers. *ench.* 2. p. 403. *Athanasia angustifolia* Walt. car. p. 200. (v. s.)

5. *cyrananthera* (Ell. sketch 2. p. 317), caule simplici apice magis pubescente, foliis evidentius 3-nerviis, invol. pubescente. ½ in Georgiae St.-Mary. (v. s.?)

† *Species exclusæ.*

*M. ALIENA* Spreng. = *Heterothalamus brunnioides*.

*M. AUSTRALIS* Spreng. = *Calea Berteriana*.

*M. BULBOSA* Raf. fl. lud. 61 et ideò Spreng. *syst.* = *Liatridia spec.*?

CDXXX. *DUBAUTIA* Gaudich. in voy. Freyc. bot. 468. t. 84. Hook. bot. Beech. 88. *Less. syn.* p. 247.

Capitulum multi- (8-10) florum homogamum. Recept. ... campanulatum, squamis circ. 8 liberis 1-serialibus sibi mutuò incumbentibus. Cor. tubulosæ regulares 5-dentatæ. Antherae ecaudatæ breviter appendiculatæ. Filam. apice vix incrassata. Stigmata dilatata

ciliata. Achænium hirsutum oblongum seu obpyramidatum, areolâ terminali. Pappus 1-serialis paleaceus, paleis angustis ciliato-plumosis. — Suffrutices Sandwicenses, ramis strigilloso-pilosis teretibus apice foliosis, infernè cicatricato-annulatis. Folia opposita sessilia rigida subparallelè venosa dentata. Capitula in racemos paniculatos aggregata. — Genus affine Tridaci ex Less. et Hook. aut Railliardæ.

1. D. PLANTAGINEA (Gaudich. bot. Freyc. voy. 468. t. 84); foliis amplexicaulis lineari-lanceolatis acuminatis crenatis glabris, paniculâ foliatâ, capitalis fasciculatis. ♂ in insul. Sandwich. montanis ad 550-600 hexap. altit., in ins. O-Wahu ex Less. in linnæâ 1831. p. 164. Facies Echli fruticosi! (v. s. comm. à cl. de Chamisso.)
2. D. LAXA (Hook. et Arn. bot. Beech. 87.) foliis oblongo-lanceolatis basi attenuatis argutè serratis subtus strigoso-hispidis, paniculâ corymboseâ laxâ nudiusculâ. ♂ in insulis Sandwich.

#### Subdiv. 2. SPHENOGYNEÆ.

Ligulæ neutræ. Pappi paleæ obovatæ obtusissimæ more corollarum contortarum convolutæ.

CDXXXI. SPHENOGYNE R. Brown h. Kew. ed. 2. v. 5. p. 142. Cass. dict. 50. p. 204. Less. syn. 242. — Oligerion Cass. dict. 2. p. 75. 29. p. 187. — Thelythamnus Spreng. f. suppl. 25. — Spermothylla Neck. elem. 1. p. 2. excl. sp. — Arctotidis sp. Linn. Thunb. — Ursinæ sp. Lam. ill. t. 716. f. 1-2. Spreng. syst. 3. p. 409.

Capitulum multiflorum radiatum, fl. radii ligulatis 1-serialibus neutris, disci tubulosis 5-dentatis hermaphroditis. Recept. paleaceum, paleis scariosis flores amplexantibus sæpius apicè truncatis. Invol. campanulatum multiseriale, squamis imbricatis, inter. majoribus apice amplè scariosis. Stylus in ramos apice truncatos ferè cuneatos desinens. Antheræ appendice terminali subcordiformi superatæ. Achænium pilis mollibus è basi ortis cinctum, cylindraceum vix basi angustatum. Pappus 1-serialis paleaceus, laminis obovatis cuneatisve obtusis ante anthesin contortarum more spiraliter dispositis, junioribus pellucidis, adultis lacteis opacis. — Herbæ aut suffrutices Capenses Anthemidis facie. Folia alterna sæpius pinnatiloba, rarius trifida, imò integerrima. Rami apice pedunculiformes nudi 1-cephali. Cor. flavæ aut apice fuscæ, ligulis interdum subtus rubris.

#### Sect. 1. THELYTHAMNOS.

Involucri squamæ exteriores absque margine scarioso manifesto.

##### §. 1. Species annuæ. — Anthemoides Less. syn. 242.

1. S. ANTHEMOIDES (R. Brown h. Kew. ed. 2. v. 5. p. 142), herbacea glabra, foliis irregulariter bipinnatipartitis lacinii linearifiliformibus, pedicellis elongatis nudis, recept. paleis conformibus truncatis, invol. squamis externis acutiusculis, ligulis discoloribus, paleis truncatis. (†) ad Cap. Bonæ-Spei. Less. syn. 242. Arctotis anthemoides Linn. sp. 1307? Berg. cap. 324. Sims bot. mag. t. 544 bona. Arctotis pinnata et A. cernua Thunb. fl. cap. 711 et 712. Arctotis paradoxa Ait. h. Kew. ed. 1. v. 3. p. 373. Ursinia anthemoides Poir. dict. 8. p. 257. (v. s. c.)
2. S. VERSICOLOR, herbacea multicaulis diffusa glabrescula, junior anabpnbescens, foliis bipinnatipartitis lobis linearibus, omnibus integris linearibus vix acutis, infer. brevioribus, pedicellis elongatis undis, invol. squamis ext. ovatis albo

submarginatis, ligulis subdiscoloribus, paleis truncatis. (1) in Africa australi ad Camisbergen alt. 4-500 ped. legit cl. Drege pl. exs. n. 2808! Affinis *S. antheroidi* sed folia simpliciter nec duplò pinnatisecta et capitula paulò minor. Ligulæ colore variabiles, intus basi atro-purpureæ cæterum croceæ, extus janiore modicè rubentes, demùm omninò purpurascentes. (v. s.)

3. *S. CHRYSANTHEMOIDES* (Less. syn. 242), herbacea....., foliis....., recept. paleis extimis amplis truncatis appendiculo parvo et rotundo superatis, interioribus spathulatis multòque angustioribus, invol. squamis extimis acuminatis, radii ligulis discoloribus. (1) ad Cap. Bonæ-Spei. Cæter. ign.
4. *S. ADONIDIFOLIA*, herbacea erecta ramosissima glaberrima, foliis sessilibus pinnatipartitis, lobis infer. brevibus integris, super. incisis pinnatipartitis, lobulis mucronatis, ramis apice nudis 1-cephalis, invol. squamis exter. orbisub lanceolatis subobtusis nigro marginatis, ligulis discoloribus, paleis truncatis pappo maturo longioribus, acheniis muriculatis. (1) in Cap. Bonæ-Spei ad Paarl. alt. 50-100 ped. legit cl. Drege pl. exs. n. 6014! Folia ferè *Senecionia* seu *Adonidis*. Paleæ elongatæ citrinæ et pappus lacteus amplus faciliè, maturitate inenante, speciem distinguunt. (v. s.)
5. *S. FENICULACEA* (Less. syn. 242), herbacea glabra erecta, foliis bipinnatisectis, lobis linearibus acutis, pedicellis nudis elongatis, invol. squamis ovatis subacutis, recept. paleis conformibus truncatis, ligulis concoloribus. (1) ad Cap. Bonæ-Spei. Barch. l. n. 6166 et fortè Eckl. n. 1232. 1226. et 106. Plura hic fortè confusæ remanent species? *Arctotis fœniculacea* Jacq. h. Schœnbr. t. 156? et idèò *Urauria fœniculacea* Poir. dict. 5. p. 258? Accedit ad *S. antheroidem*, sed capitulum minus, fol. lobi acutiores, ligulæ concolores pallide flavæ. (v. s.)
6. *S. CALENPULEFLORA*, herbacea erecta ramosa glabra, foliis bipinnatisectis in petiolum attenuatis, lobis acutis rachibusque linearibus, ramis apice longe nudatis pedicelliformibus, invol. squamis exter. apice subrufo-scariosis, ligulis concoloribus invol. quadruplò longioribus, paleis truncatis. (1) ad Cap. Bonæ-Spei loco dicto Zilverfontain alt. 3-400 ped. legit cl. Drege pl. exs. n. 284! Herba pedalis. Pedicelli 4-6-pollic. Ligulæ croceæ pollicem longæ. (v. s.)
7. *S. RUSILLA*, herbacea ramosa glabra, foliis crassiusculis pinnatipartitis, lobis linearibus obtusiusculis nunc integerrimis nunc in lobos 2-3 partitis, pedicellis plurimis 1-cephalis subcorymbosis caulibus ferè longitudine, invol. squamis exter. lanceolatis subacutis, ligulis concoloribus, paleis truncatis, acheniis muricatis. (1) in Cap. Bonæ-Spei ad Olifant-Rivier legit cl. Drege pl. exs. n. 6011! Herba 3-6-poll. Invol. pallidum, squamis int. apice subscariosis. Cor. flavæ Pappus amplus lacteus. (v. s.)
8. *S. PALLIDA*, herbacea erecta tenella glabra, foliis pinnatipartitis 7-9-jugis lobis inf. parvis acutis integris, sup. in lobulos lineares acutiusculos apice 3-partitis, pedicellis elongatis nudis 1-cephalis, invol. squamis pallide stramineis, exter. fusco marginatis acutis, ligulis concoloribus invol. paulò longioribus. (1) in Africa australi ad Clam-William legit cl. Ecklon. Herba 5-6-pollic. (v. s.)  
*ℳ. immarginata*, invol. squamis stramineis immarginatis. — ad Camisbergen legit cl. Drege n. 5026! Fortè species propria, sed ex specim. carie excoriat vix admittenda. (v. s.)
9. *S. MICROCEPHALA*, herbacea erecta subsimplex sparsè hispida, foliis pinnatipartitis lobis paucis linearibus integris acutiusculis, caule ramisque apice nudis pedunculiformibus 1-cephalis, invol. squamis exter. margine nigris lanceolatis acutis, ligulis discoloribus, paleis recept. truncatis. (1) in Africa austr. ad Berg rivier ferè maris altit. legit cl. Drege (pl. exs. n. 6014!). Capitulum ligula exceptis circ. 3. lin. diam. Ligulæ angustæ 2 lin. longæ. Pappus mat. non vidi. (v. s.)

§. 2. *Species perennes aut frutescentes.* — *Thelythamnus Less.*  
syn. p. 243.

\* *Foliis pinnatim incis, seu uni- aut bipinnatisectis.*

a. *Foliis hispidis sericeis tomentosive.*

10. *S. CORONOPHOLIA* (Less. syn. 244), herbacea tota tomentoso-lanata canescens erecta, foliis confertis oblongo-ligulatis per totam longitudinem et apice acutè inciso-serratis, summis linearibus acutissimis integerrimis, ramis apice nudis 1-cephalis involucrique squamis exter. linearibus lanatis, intimis apice acuminatis scariosis, ligulis concoloribus.  $\mathcal{L}$  in Africæ Capensis distr. Tulbagh et Worcester legit cl. Ecklon! Species distinctissima! (v. s.)
11. *S. HISPIDA*, suffruticulosa erecta molliter hispida, foliis pinnatisectis laciniis linearibus mucronato-acutis integerrimis, capitulo supra suprema folia subsessili, invol. squamis exter. acutis, recept. paleis apice 3-dentatis, radii ligulis concoloribus brevibus.  $\mathcal{Z}$  ad Cap. Bonæ-Spei legit cl. Burchell. Species distinctissima? An hujus, an sequentis sectio? Duratio et pappus maturus ign. (v. s. comm. à cl. Burch. sub n. 6998 et 7035.)
12. *S. DREGIANA*, caulibus è collo plurimis albo-tomentosis, foliis molliter hispidis subcanescentibus pinnatipartitis, lobis linearibus mucronatis integerrimis utrinque 6-7, pedunculis longissimis nudis glabriusculis 1-cephalis, invol. squamis exter. lineari-lanceolatis puberulis, ligulis concoloribus, paleis obtusè truncatis.  $\mathcal{L}$   $\mathcal{Z}$  ad Cap. Bonæ-Spei in Giftberg cum plur. rariss. et novis stirp. legit cl. Drege pl. exs. [n. 6005! Caules ferè cæspitosi, pedunculo triplò breviores. Invol. squamæ int. in amplam scariosam appendicem productæ. Capit. florens diam. pollicari. (v. s.)
13. *S. SUBHIRSAUTA*, fruticosa ramosa adscendens, ramis pedunculis folisque præsertim ad costam molliter subhirsutis, foliis pinnatisectis, lobis infer. 4-5-jugis brevibus integris, super. 3-4-jugis longis pinnatipartitis, lobis lobulisque acutis, pedicellis nudis folio triplò ferè longioribus, invol. squamis exter. rufo-marginatis apice brevissimè scariosis, ligulis concoloribus, recept. paleis truncatis.  $\mathcal{Z}$  in Africæ Capensis distr. Swellendam legit cl. Ecklon. Accedit ad *S. abrotanifoliam*. Fruct. ign. (v. s.)
14. *S. ABROTANIFOLIA* (Brown h. kew. ed. 2. v. 5. p. 143. Less. syn. 243), fruticosa ramosa pube brevi patulâ hirsutâ subcanescens, foliis bi-tri-pinnatisectis, segmentis divergentibus acutis, infer. brevibus subdivisis, pedunculis folio quadruplò longioribus, invol. tomentosi squamis ext. lanceolatis, intimis amplè scariosis, radii ligulis concoloribus, paleis recept. apice denticulatis.  $\mathcal{Z}$  ad Cap. Bonæ-Spei ad Paarlberg alt. 2-300 ped. legit cl. Drege pl. exs. n. 1760. Ursinia abrotanifolia Spreng l. c. (v. s.)
15. *S. SERICEA* (Less. l. c.), fruticulosa sericeo-villosa cana, foliis bipinnatisectis laciniis brevibus elongatisve lineari-filiformibus integris, pedunculis folio multò longioribus, invol. squamis extimis subtomentosis ovatis acutis, radii ligulis concoloribus, paleis recept. apice denticulatis.  $\mathcal{Z}$  ad Cap. Bonæ-Spei (Th.), ad Zeederbergen alt. 3-400 ped. (Drege! pl. exs. n. 1978), Tulbagh (Ecklon!). *Arctotis sericea* Thunb. fl. cap. 713. *Sph. peduncularis* E. Mey! in Dreg. herb. (v. s.)
16. *S. MACROPODA*, caule suffruticoso adpressè sericeo-villoso, foliis junioribus sericeo-villosis, adultis glabrat, pinnatipartitis, lobis paucis racheque linearibus elongatis integerrimis, pedicellis longissimis glabris, invol. squamis ext. lineari-lanceolatis glabris, ligulis concoloribus, recept. paleis apice subdenticulatis, pappi laminis 3-5 albis obovatis cor. dimidiam longit. æquantibus.  $\mathcal{L}$   $\mathcal{Z}$  ad Cap. Bonæ-Spei loco dicto Giftberg alt. 2-300 ped. legit cl. Drege n. 6008! Ligulæ aureo-flavæ. Foliorum lobi 9-12 lin. longi. Pedicelli ferè pedales. (v. s.)

$\mathcal{L}$ . *ambigua*, foliorum lobis brevioribus obtusis, invol. squamis exter. sub-

villosa, ligulis paulò magis pallidis, pappo dimidiò breviora.  $\frac{1}{2}$   $\frac{1}{2}$  ad Zanderbergen alt. 3-400 ped. legit cl. Drege pl. exs. n. 6008! An forte species distinguenda? (v. s.)

$\beta$ . *Foliis glabriusculis, lobis brevibus dentatis, dentibus apice setigeris.*

17. *S. DENTATA* (R. Brown h. Kew. ed. 2. v. 5. p. 143), fruticosa glabra, foliis pinnatipartitis, lobis brevibus integris aut trifidis, dentibus setigeris, pedunculis valdè elongatis, invol. squamis exter. ovatis acutis, radili ligulis discoloribus.  $\frac{1}{2}$  ad Cap. Bonæ-Spei circa urbem. Burch! cat. Afr. austr. n. 438. Drege! n. 13. Eckl. n. 1920. 460. 858. *Arctotis dentata* Linn. sp. 1307. N. L. Burm.! prod. 44. non Thunb. nec Willd. *Arctotis piligera* Berg. cap. 325. *Arctotis pectinata* Thunb. fl. cap. 710. *Ursinia dentata* et pilifera Polr. dict. 8. p. 256. Speng. syst. 1. c. — Burm. Afr. t. 64. Pluk. alm. t. 276. f. 2. et fortè 382. f. 5. (v. s.)

$\beta$ . *marginata*, invol. exter. nigro marginatis.  $\frac{1}{2}$  ad Cap. Bonæ-Spei Burch! cat. Afr. austr. n. 6698. Drege! pl. exs. Eckl. n. 1257? 1547. 1622. (v. s.)

18. *S. ASTRIGATA*, suffruticosa, caule inter folia puberulo, foliis pinnatisectis glabris, segmentis utrinque 4-5-apice 2-4-fidis, dentibus longè setigeris, pedunculo elongato folia sparsa linearia integra apice setigera gerente, invol. squamis exter. lanceolatis acutis nigro-marginatis, ligulis discoloribus invol. triplò longioribus, recept. paleis truncatis.  $\frac{1}{2}$  in Africa austr. ad Cap. Aphie legit cl. Drege pl. exs. n. 6009! Valdè accedit ad *S. dentatam*, sed capitula duplò majus et setæ fol. frequentiores et longiores. (v. s.)

$\gamma$ . *Foliis glabris seu glabriusculis, lobis apice non setigeris.*

19. *S. NUDICAULIS* (Less. 1. c.), glaberrima, basi suffruticosa, foliis subradicalibus pinnatisectis, lobis 6-7-jugis, inferioribus brevibus indivisis, superioribus longioribus variisimè partitis, lobis lobulisque linearibus vix acutis, apice lobis multò longiore nudo stricto, invol. squamis extimis ovatis acutis, ligulis discoloribus invol. duplò longioribus.  $\frac{1}{2}$  ad Cap. Bonæ-Spei (Burch! n. 632. a. montem Tabularem alt. 4-500 ped. (Drege! n. 6007. Eckl. n. 1918. 659. 1000). *Arctotis nudicaulis* Thunb. fl. cap. 711. Variat parte caulis fruticosa nec emissa aut rhizoidea ascendente, nunc erecta et verè caulescente. (v. s.)

20. *S. SCAPIFORMIS*, glaberrima, rhizomate suffruticoso, foliis subradicalibus pinnatipartitis, lobis 10-12-jugis, inf. parvis indivisis acuminatis, sup. longioribus in lobos lineares 2-3 partitis, pedicello longissimo stricto nudo nullo, invol. squamis exter. ovato-lanceolatis acutis, ligulis concoloribus invol. circ. longitudine.  $\frac{1}{2}$   $\frac{1}{2}$  ad Cap. Bonæ-Spei loco dicto Langekloof. alt. 200 ped. legit cl. Drege pl. exs. n. 6004 et ad Talbagh cl. Ecklon n. 1361. et Worcester n. 1446! An mera *S. nudicaulis* var.? Differt præter notas indic. titulo dimidiò minore. Stirps Eckl. n. 1743, 1047, 213 ad Worcester ut utrasque ferè conjungit. (v. s.)

21. *S. CHAMOMILLEFOLIA*, glaberrima, caule imò basi suffruticulosos radice subadscendente aut erecto, foliis pinnatipartitis, lobis linearibus acutis 4-jugis, lob. inf. indivisis, sup. 2-3-lobatis longioribus, pedicellis valdè obsoletis nudis 1-cephalis, invol. squamis ext. minimis acutis, mediis ovali-oblongis obtusis, summis scarioso-appendiculatis, ligulis discoloribus, paleis truncatis subdenticulatis.  $\frac{1}{2}$   $\frac{1}{2}$  in Cap. Bonæ-Spei ad Draakensteenberg alt. 3-400 ped. legit cl. Drege pl. exs. n. 6006! ad Worcester (Ecklon! n. 923). Assidua *S. scapiformem*. (v. s.)

22. *S. ANETHIFOLIA* (Less. syn. 243), fruticosa erecta glabra, foliis pinnatisectis punctatis pinnatisectis, lobis lineari-filiformibus acutis, inf. brevibus, sup. apice 2-3-fidis, pedicellis elongatis 1-cephalis, invol. squamis ligulis discoloribus invol. paulò longioribus.  $\frac{1}{2}$  ad Cap. Bonæ-Spei. Lobi fol. utrinque 7-8. Ligulæ 3 lin. longæ. An certè notè giana ex descriptionis brevitate dubia? (v. s. comm. à Verreaux.)

23. *S. ANETHOIDES*, fruticosa erecta glabra aut subpuberula, foliis subcoriaceis pinnatisectis, lobis lineari-filiformibus acutis subpunctulatis, infer. integris, super. apice 3-fidia, pedicellis elongatis 1-cephalis, invol. squamis exter. lanceolatis acutis, ligulis concoloribus invol. duplò longioribus, paleis truncatis.  $\zeta$  ad Cap. Bonæ-Spei prope Breederivier et Draakensteensberg legit cl. Drege alt. 1-200 ped. pl. exs. n. 6015, 6019, 6020! et ad Worcester cl. Ecklon n. 1058 592. 212. 1860! Proximè accedit ad *S. anethifolium*. (v. s.)  
 $\beta$ . *brachyglossa*, ligulis vix invol. longioribus.  $\zeta$  ad Piquetberg alt. 2-300 ped. legit cl. Drege pl. exs. n. 6012! (v. s.)  
 $\gamma$ . *ramosissima*, caule laxè ramosissimo, foliis brevioribus brevius lobatis.  $\zeta$  ad Giftberg alt. 2-300 ped. legit cl. Drege pl. exs. n. 6011! (v. s.)
24. *S. ORACILIS*, fruticulosa erecta virgata glabrescens, foliis pinnatipartitis brevibus, lobis inf. indivisis, sup. subternis 3-dentatis vix 3-lobatis, omnibus lineari-filiformibus mucronulatis brevibus, ramis apice longè denudatis, invol. squamis exter. paucis minimis acutis, cæteris obtusis, ligulis concoloribus vix invol. longioribus, paleis truncatis.  $\zeta$  in Africâ australi ad Zeederbergen alt. 4-500 ped. legit cl. Drege pl. exs. n. 6010! Affinis *S. anthemoidi*, sed capit. dimidiò minora et ligulæ minimæ. Villi baseos fructûs longissimi. (v. s.)
25. *S. CHAMÆMELOIDES*, suffruticosa glabra, foliis pinnatisectis, lacinii linearibus acutis, superiorum integris, inferiorum sæpè trifidis, invol. squamis exter. triangularibus acuminato-aristatis, radii ligulis discoloribus.  $\zeta$  ad sinum Algoæ Africæ australis (Forbes!). Rami floriferi parçè foliosi 1-2 poll. longi. Invol. squamæ plurimæ scarioso-membranaceæ. Ligulæ 5-7-nerviæ. (v. s.)
26. *S.?* *RIGIDULA*, suffruticosa erecta aut ascendens glabra, foliis pinnatipartitis subepinescentibus, inf. brevibus integris, sup. longioribus trilobatis, pedicellis elongatis 1-cephalis, invol. squamis ext. ovali-oblongis margine scariosis subobtusis, ligulis discoloribus, paleis obtusis.  $\zeta$  in Africâ austr. ad Piquetberg alt. 2-300 ped. legit cl. Drege pl. exs. n. 6013! Aff. *S. chamæmelifolæ* et cum eâ fortè ad Uriniam repellenda? Fruct. ign. Pedicelli 3-6 poll. longi. Folia 10-12 lin. longæ. (v. s.)
27. *S. BREVIFOLIA*, fruticosa erecta ramosissima puberula aut glabrescens, foliis rigidulis glaucescentibus pinnatipartitis paucijugis, lobis mucrone subuncinato-acutis, infer. 2-3-jugis minimis interdum nullis, super. 3-5 approximatis longioribus, pedicellis elongatis folia linearia integerrima erecta basi gerentibus, invol. squamis ext. ovali-oblongis apice subscariosis obtusis rufè marginatis, ligulis discoloribus, paleis obtusis.  $\zeta$  in Africâ Capensis distr. Uitenhage et Caledon legit cl. Ecklon n. 275. 258. 963. 1431 et 1257. Fruct. ign. Folia 5 lin. longæ. (v. s.)
28. *S. PALMATA* (Less. syn. 244), fruticosa glabrescens, foliis pinnatisectis, segmentis margine revolutis, invol. squamis ext. numerosis angustè linearibus acuminatis conduplicatis subpatulis, radii ligulis discoloribus.  $\zeta$  ad Cap. Bonæ-Spei. Arctotis paleacea Thunb. fl. cap. 712.
29. *S. GRANDIFLORA*, fruticosa glabra, foliis punctatis pinnatisectis, lacinii linearibus acutis, inf. brevioribus, super. longioribus fastigiatis, pedunculo foliis duplò longiore, invol. squamis linearibus vix acutis, radii ligulis concoloribus squamis inv. inter. duplò longioribus.  $\zeta$  ad Cap. Bonæ-Spei in Hortentots-Holland (ex h. Deless.) ad Draakensteensberg alt. 3-400 ped. legit cl. Drege l. n. 6018. Eckl. l. n. 257. Valdè affinis *S. punctatæ* et forsàn cum eâ à plurimis conjuncta, sed ligulæ 8-9 lin. longæ, 2 lin. ferè latæ, 5-7-nerviæ, evidentiùs quàm prioris glandulosæ. (v. s.)
30. *S. MACULON* (Less. syn. 243), fruticosa tomentosa (Th.), foliis coriaceis pinnatisectis segmentis integerrimis, invol. squamis extimis minutis ovatis acutis, radii ligulis discoloribus.  $\zeta$  ad Cap. Bonæ-Spei. Arctotis dentata Thunb. fl. cap. 712. excl. syn. ex Less. Capitula solitaria (Less.). Fol. segmenta subulata brevia. (Thunb.)
31. *S. CRITHMIFOLIA* (R. Brown h. Kew. ed. 2. v. 5. p. 142), fruticosa glabra,

45. *S. serrata*, suffruticosa puberula, foliis semimplexicaulibus oblongis lanceolatisve grossè serratis, pedunculis pluribus 1-cephalis subcorymbois foliis triplò longioribus, radii ligulis concoloribus.  $\S$  ad Cap. Bonæ-Spei Burch! n. 5965. Eckl. n. 1351. *Arctotis serrata* Thunb. fl. cap. 705. Linn. f. suppl. 385. *Ursinia serrata* Poir. et Spreng. ll. cc. *Sphen. Sommeratii* Cass. dict. 5. p. 205. (v. s.)
46. *S. incisa*, suffruticosa puberula, foliis elongato-oblongis apice subulatis inciso-serratis, serraturis acuminatis, inferioribus parvis distantibus, super. approximatis longioribus, ramis in pedunculos 1-cephalos in parte medià folio quadruplò longioribus, radii ligulis concoloribus.  $\S$  ad Cap. Bonæ-Spei in distr. Zwellendam legit cl. Burchell et mecum sub n. 7008 cat. geogr. com. Folia sesquipoll. longa, basi 2-3 apice 4-5 lin. lata. (v. s.)
47. *S. pilifera* (Ker bot. reg. t. 60), fruticosa ramoso-diffusa, foliis pinnatisectis subcarnosulis patentim et brevè hispidulis, lobis linearibus acutè terminatis, pedicellis elongatis subhispidulis, invol. squamis inf. margine scarioso angustò cinctis obtusis, ligulis discoloribus, paleis truncatis, pappi lamina obovatis achænio paulò brevioribus.  $\S$  ad Cap. Bonæ-Spei. Less. syn. 244. *Arctotis pilifera* Thunb. fl. cap. 712 non Berg.
- $\beta$ . *crithmoides*, foliis glabriusculis, lobis brevioribus.  $\S$  *Arctotis crithmoides* Jacq. h. Schönbr. 2. t. 155 non Berg. *Ursinia crithmoides* Poir. dict. 8. p. 258.
- $\gamma$ . *subcanescens*, ramulis foliisque novellis pube adpressâ plus minus cunctibus, foliorum lobis abbreviatis.  $\S$  ad Zwellendam legit cl. Drege, pl. exs. n. 6367. (v. s.)
48. *S. scariosa* (R. Brown et Less. ll. cc.), fruticosa glabra, foliis pinnatisectis segmentis lineari-filiformibus integerrimis mucronulatis, pedunculis folio duplò triplòve longioribus, invol. squamis extimis omninò scariosis subrotundis, radii ligulis subdiscoloribus, rec. paleis truncatis, achæniis molliter et parte villosis.  $\S$  ad Cap. Bonæ-Spei. *Arctotis scariosa* Willd. sp. 3. p. 236 excl. Thunb. syn. *Ursinia scariosa* Poir. dict. 8. p. 257. — Burch. l. cat. afr. austr. n. 5451. Sæpè in hortis sub nom. *Arctotidis paleaceæ* vagat. (v. s. sp. et c.)
- $\beta$ . *brevipes*, capitulis supra ultima folia brevissimè pedicellatis.  $\S$  ad Cap. Bonæ-Spei Drege! pl. exs. n. 6016. (v. s.)
49. *S. odorata* (R. Brown l. c.), fruticosa glabra, foliis planis apice inciso-pinnatifidis, pappo obsoleto, radii ligulis discoloribus.  $\S$  ad Cap. Bonæ-Spei. *Anthemis odorata* Ait. kew. 3. p. 238. Willd. sp. 3. p. 2186. *Achænia glabra* ex Br.
- CDXXXII. *URSINIA* Gærtn. fruct. 2. p. 462. t. 174. Less. syn. 244. — *Ursina* sp. Poir. Spreng. — *Arctotidis* sp. auct.

Omnia *Sphenogynes*, sed 1°. achænium obovatum turbinatum basi distinctè attenuatum glaberrimum nervulis 5 minimis subcostatis. 2°. Pappus duplex, ext. laminis 5 amplis obovatis lacteis spiraliter revolvato-contortis ut in *Sphenogyne*, int. filis 5 concoloribus angustissimis imâ basi subconcretis. — Herbæ aut frutes Capenses. Folia univ. et bipinnatisecta. Invol. squamæ pleræque apice in appendicem scariosam productæ.

### §. 1. *Species perennes aut frutescentes.*

1. *U. MONTANA*, fruticosa ascendens humilis parçè ramosa, caule ramisve hinc inde subpuberis, foliis confertis pinnatipartitis crassinerviis glabris puberis. lobis inf. minimis acutissimis, sup. linearibus mucronulatis, pedicellis ant. longitudine caulis, invol. ovatis squamis ext. ovato-lanceolatis obtusis margine nigrello et scarioso albo cinctis, ligulis parvis discoloribus, paleis truncatis.  $\S$  ad Cap. Bonæ-Spei ad Willbergen alt. 8-900 ped. legit cl. Drege, pl. exs. n. 6027. Fruct. matur. non vidi. (v. s.)



2. *U. APICULATA*, fruticosa caespitosa demissa glabra, foliis pinnatipartitis, lobis omnibus apiculo albo mucronatis seu albo-sphaeclatis, sup. linearibus indivisis longis trilobisve, pedicellis caule ferè longioribus, invol. squamis ext. ovatis nigro marginatis, ligulis discoloribus, paleis obtusis subdentatis.  $\mathcal{U}$   $\mathfrak{z}$  in Cap. Bonæ-Spei ad Sneeuwbergen alt. 5-600 ped. legit cl. Drege, pl. exs. n. 2132. Habitus *U. montanae*, sed ab hac et ab omnibus facilè differt fol. lobis apice mucrone albisimo terminatis. (v. s.)
3. *U. TENULOSA*, glaberrima, caule suffruticuloso subdemisso brevi parèe ramoso, foliis pinnatipartitis, lobis linearibus aristato-acuminatis, infer. brevioribus integris, super. 2-3-lobis, pedicellis caule longioribus, paleis apice obtusè dentatis, pappo int. laminis exterioribus paulò brevioribus.  $\mathcal{U}$   $\mathfrak{z}$  in Cap. Bonæ-Spei ad Kat-Riviersberg alt. 5-600 ped. legit cl. Drege, pl. exs. n. 3698 et in distr. Ultenhage prope Zwartkop-Rivier cl. Eckl. n. 107. Urnsia forniculacea Eckl. l. c. non Spreng. Folia ferè Santolinæ viridis. Habitus Achillæ moschatæ. Pappus ext. amplius. (v. s.)
4. *U. PARADOXA* (Gærtn. et Less. ll. cc.), suffruticulosa ascendens glabriuscula, foliis pinnatisectis, lobis inf. brevibus integris, sup. longioribus 2-3-fidis, lobulis oblongo-linearibus apice piliferis, ramis apice nudis pedicelliformibus, invol. squamis omnibus margine scariosis obtusis, ligulis discoloribus, pappo int. exteriore breviori.  $\mathcal{U}$   $\mathfrak{z}$  ad Cap. Bonæ-Spei legit cl. Drege, pl. exs. n. 531. *Arctotis paradoxa* Linn. amœn. 4. p. 330 (excl. syn.). Thunb. fl. cap. 712. *Sphenogyne anthemoides* E. Mey. in Drege coll. non Brown. Habitus omnino *Sphaen. anthemoidis*, sed differt caract. gen. (achæn. glabro et pappo duplice) nec non fol. apice piliferis, et invol. squamis omnibus margine scarioso albo dactis. Capitulum non malè refert *Anacyclum Pyrethrum* seu icon. bot. mag. t. 544. (v. s.)
5. *U. ALBICAULIS*, suffruticulosa erecta, ramis sublanatis, foliis pinnatipartitis glabriusculis, lobis linearibus mucronatis integerrimis aut apice 2-3-fidis, ramis apice nudis pedicelliformibus glabriusculis, invol. squamis ext. in appendicem scariosam lanceolato-acuminatam productis, pappo inter. externum longitudine subsequente. — ad Cap. Bonæ-Spei ad Zwellendam t'Koussé et Gariep, alt. 2-300 ped. legit cl. Drege. Pedicelli circ. 2-3 pollic. longi. Ligulæ intus flavæ subdiscolores 9-10 lin. longæ. Pappus lacteus, exterior amplius. (v. s.)
6. *U. PUNXULOSA*, subherbacea, ramis laxè sublanatis, foliis glabris bipinnatisectis, lobis linearibus mucronatis, ramis longè denudatis glabris, invol. squamis ext. in appendicem scariosam lanceolato-acuminatam productis. — ad Cap. Bonæ-Spei loco dicto Klein-Namaqualand alt. 2-300 ped. legit cl. Drege, pl. exs. n. 2807. An ab *U. albicaulis* certè diversa? Fruct. matur. ign. tamen in specim. quodam (Eckl. n. 322) huc ex invol. et foliis relato vidi achænia turbinata lævia albida pappo (verisimiliter deciduo) omnino destituta. (v. s.)
7. *U. RADICANS*, caule subherbaceo decumbente basi repente glabriusculo, foliis pinnatipartitis, lobis inf. simplicibus, super. 2-3-lobatis acutè subdentatis, omnibus mucrone apiculatis, pedicellis modicè elongatis 1-cephalis, invol. squamis ext. lanceolatis appendice brevi scariosa subapiculatis, ligulis discoloribus, paleis obtusis apice subdentatis, pappo int. exteriore paulò breviori.  $\mathcal{U}$  ad Cap. Bonæ-Spei in Algoa-bay. verisim. in arenosis maritimis legit cl. Drege, pl. exs. n. 2131. Habitus *Pyrethri maritimi*. Accedit ad *U. speciosam*, sed habitu decumbente et repente facilè diversa. (v. s.)
8. *U. FILICAULIS*, herbacea glabra, caule filiformi prostrato hinc indè radicante, foliis pinnatipartitis, lobis acutis rachique linearibus, infer. indivisis, super. trifidis, pedicellis ascendentibus 1-cephalis, invol. campanulatis, squamis ext. in appendicem scariosam acuminatam productis, ligulis discoloribus, paleis obtusis.  $\mathcal{U}$  ad Cap. Bonæ-Spei loco dicto Kaaspeche-Vlakte legit cl. Drege, pl. exs. n. 6021 in distr. Caledon? (Eckl. n. 454?). Car. gener. non vidi sed huc refero ex involuero, achænio juniore, et analogiam maximam cum *U. radicante*. Caules circ. 2 ped. longi graciles. Folia remota. Ligul. crocæ, subtus subrubræ, 7-8 lin. longæ. (v. s.)

§. 2. *Species annuæ.*

9. *U. SPECIOSA*, herbacea erecta ramosa glabra, foliis pinnatipartitis, lobis inf. brevibus integris, sup. longioribus incis. pinnatifidisve, lobulis omnibus un. apiculatis, pedicellis nudis elongatis, invol. squamis ext. obtusiusculis scarioso-marginatis, ligulis concoloribus, paleis truncatis, pappo intern. lamina ext. ternis paulo brevior. (1) in Cap. Bonæ-Spei loc. dict. Zwellendam (Kome et Gariep, alt. 2-300 ped. legit cl. Drege. Herba et infl. ferè *Chrysanthemi* carinati.

*α. croceo-flava*, ligulis croceo-flavis. Drege! pl. exs. n. 2745. (v. s.)

*β. albida*, ligulis albidis. Drege! pl. exs. n. 6368. (v. s.)

10. *U. CAKILEFOLIA*, herbacea erecta ramosissima glabra, foliis crassiusculis pinnatipartitis, lobis linearibus acutis, inf. integris, sup. partitis, pedicellis elongatis nudis, invol. squamis ext. in appendicem subscariosam acuminatam productis, ligulis concoloribus, paleis ext. 3-dentatis, pappo inter. externi longitudinem æquante. (1) in Cap. Bonæ-Spei ad Giftberg, alt. 2-300 ped. legit cl. Drege, pl. exs. n. 6023! Capitula parva. Flores flavi. (v. s.)

*β. crocea*, ligulis croceis paulo majoribus, invol. squamis appendicem subnigricantibus, paleis maturis rufidulis. (1) ad Cap. Bonæ-Spei legit cl. Drege, pl. exs. n. 6025! (v. s.)

11. *U. INDECORA*, herbacea ramosissima glabrescens, foliis crassiusculis pinnatipartitis, lobis linearibus, inf. integris, sup. tri- aut pluripartitis, pedicellis ovatis 1-cephalis elongatis, invol. squamis ext. ovatis subacutis, ligulis concoloribus invol. paulo longioribus, paleis truncatis, pappo int. externum longitudine æquante. (1) ad Cap. Bonæ-Spei in Carro, alt. 3-400 ped. legit cl. Drege, pl. exs. n. 7721! Habitus Senecionis. Affinis *U. cakilefoliæ*. Ligulæ 3-4 lin. longæ. (v. s.)

12. *U. NANA*, herbacea glabra pluricaulis subdiffusa, foliis crassiusculis pinnatipartitis, lobis linearibus obtusiusculis, aliis integerrimis, aliis 2-3-lobatis, rami apice pedicelliformibus 1-cephalis, invol. squamis ext. nigro marginatis ex subacutis int. apice albo-scariosis rotundatis, ligulis concoloribus invol. paulo longioribus, pappi int. setis lamina ext. brevioribus. (1) in Cap. Bonæ-Spei ad Silver-Fontain, alt. 2-300 ped. legit. cl. Drege, pl. exs. n. 2782! Habitus ferè *Sphenogynæ* pusillæ, sed magis diffusa et ligulæ minores. Arhenia *U. glabra* basi attenuata nervulis 5 costulata. An hæc aut ad sequentem referenda Urs. annua Less. in linnæa 1831. p. 165, nomine tantum nota? (v. s.)

13. *U. PYGMÆA*, herbacea glabra pluricaulis suberecta, radice exili, foliis pinnatipartitis, lobis paucis lineari-filiformibus, capitulis pedicellatis solitariis, invol. squamis ext. acuminatis. (1) in Africæ Capensis distr. Namaqualand legit cl. Ecklon, n. 27. Specimen manum tantum vidi sed ex fracto ad Uraliam referendum: ex invol. squamis acuminatis ab *U. nanâ* distinguendum. Herba pollicaris. (v. s.)

† *Species exclusæ.*

*U. ABROTANIFOLIA* Spreng. = *Sphenogyne abrotanifolia*.

*U. ANTHEMOIDES* Poir. = *Sphenogyne anthemoides*.

*U. CRITHMOIDES* Poir. = *Sphenogyne pilifera β.*

*U. DENTATA* Poir. = *Sphenogyne dentata*.

*U. FILIFORMIS* Griess. = *Sphenogyne pinnata*.

*U. FOENICULACEA* Poir. = *Sphenogyne ulacea*.

*U. FILIFERA* Poir. = *Sphenogyne*

*U. SCARIOSA* Poir. = *Sphenogyne*

*U. SERRATA* Poir. = *Sphenogyne*

*U. TENUIFOLIA* Poir. = *Sphenogyne*

## DIV. III. MADIÆ (1).

Receptaculum margine aut ferè totum paleaceum. Invol. squamæ 1-2-seriales convolutæ aut complicatæ achænia radii semper calva includentes. Achænia disci pappo paleaceo donata aut calva, fertilia aut sterilia. — Folia alterna. Flores flavi.

CDXXXIII. MADIA *Mol. chil. ed. germ.* 113. *ed. gall.* 336. *Cav. ic.* 3. p. 50. — Madia et Biotia *Cass. dict.* 34. p. 308. — Madia sp. *Don, Less.*

Capitulum multiflorum heterogamum, fl. radii 10-12 uniserialibus foemineis ligulatis aut monstrosè irregulariter et amplè tubulosus, fl. disci hermaphr. tubulosus regulariter 5-dentatis. Invol. subglobosum 1-seriale, squamis (tot quot ligulæ) dorso carinatis complicatis achænia radii involventibus. Recept. planum paleas inter radium et discum 1-2-seriales gerens, in centro omnino nudum. Ligulæ basi glabræ invol. vix longiores. Styli fl. disci rami infernè ad latus internum series duas stigmatico-glandulosas gerentes, apice pilis rigidis erectis onusti. Achænia compressa recta basi attenuata glabra calva utrinquè in medio nervo longitudinali aucta et ideò 4-5-gona. — Herba Chilensis annua erecta villosa et apice præsertim glanduloso-pilosa. Folia infima opposita, sup. alterna, semiamplexicaulia oblonga integra. Capitula breviter pedicellata in racemum digesta foliis parvis sæpius bracteata. Cor. luteæ. Semina oleosa.

1. M. SATIVA (Mol. et Don II. cc.). (1) in Chili spontanea et ad oleum è seminibus eliciendum culta sub nom. vernaculis *Madi* et *Melosa* ex Bertero! Eandem fortè cultam in Californiâ legit cl. Douglas. M. sativa, M. viscosa et M. melloso Willd. sp. 3. p. 195 vix varietates ex Don. M. viscosa Cav. ic. 3. p. 50. t. 298. M. melloso Jacq. hort. Schœnbt. 3. p. 29. t. 302. M. sativa DC. not. jard. gen. VII. n. 5. mem. soc. Gen. 7. p. 277. (v. v. c. et s. sp.)

\* monstrosa, ligulis in tubulos ampliatos irregulariter imutatis. — Biotia Cass. l. c. (v. v. c.)

+ Species exclusa.

M. ELEGANS Don. = Madaria elegans.

CDXXXIV. MADARIA DC. mem. soc. Gen. v. 7. p. 280. not. pl. rar. jard. Gen. VII. n. 6. — Madia sp. Don.

Capitulum multiflorum radiatum, ligulis 10-12 uniserialibus foemineis, fl. disci tubulosus 5-dentatis masculis seu sterilibus. Invol. subglobosum 1-seriale torulosum, squamis complicatis achænia radii omnino involventibus. Recept. subconvexum, margine inter ligulas et discum squamas 1-2-seriales gerens, medio epaleaceum densè et breviter fimbriatissimo-hirsutum. Ligulæ involuero duplò longiores et corollæ tubulosæ basi puberulo-hirtellæ. Styli rami apice hispiduli. Achænia radii compressa nervis in latere utroque aucta et ideò sub-4-5-gona, glabra, calva basi rostello incurvulo donata, disci omnino abortiva. — Herba Californicæ annuæ erectæ villosæ et sæpè glandu-

(1) Genera sex hic habitu toto consociata inter se characteribus normalibus valdè differunt, et strictissimo sensu Hartmannia et Madaroglossa fortè ad Galinsogaea, Hemizonia et fortè Madaria ex disci achæniis calvis et abortivis ad Parthenias, Madia et Oxyura ex disci achæniis calvis et fertilibus ad Heliospidea essent referenda? sed in ordine naturali, ut in civili, summum jus sæpè summa injuria.

loso-pilosæ. Folia sessilia lanceolata aut linearia integra. Capitula in corymbum laxum disposita, longius pedicellata chracteata. Flores flavi. — Nomen ex *μαδαρος* calvus.

1. *M. ELIOGANS*, caule foliisque pilis apice glandulosis et setis eglandulosis mixtis hispido. ① in Californiâ detexit cl. Douglas. *Madia elegans* Don in Lindl. bot. reg. t. 1458. Ligulæ basi purpuro-maculatæ, lineari-cuneatæ, apice acute infide. Achænia radii intrâ inv. squamas inclusa subbiserialia nigra compresso-trigona, in quoque latere uninervia, nudè ferè 5-gona apice truncata omni calva. (v. s. sp. et v. c.)
2. *M. COMYMOSA*, caule involucriisque pilis apice glandulosis et setis eglandulosis hispido, foliis linearibus villosis subhispidis eglandulosis. ① in Amer. bot. oecd. ad Columbia-Rivier legit cl. Douglas. Variat pilis glandulosis plurimis aut paucissimis. Achænia compresso-trigona, in specim. fortè non commutata, rufa submaculata. (v. s.)

β. *hispida*, caule foliis involucriisque pilis omnibus eglandulosis paucis hispida. ① in Californiâ legit cl. Douglas. An fortè species propria? (v. s. comm. cum var. α ab hon. Soc. hort. Lond.)

#### CDXXXV. HEMIZONIA.

Capitulum multiflorum radiatum, ligulis 1-serialibus femineis invol. paulò longioribus, fl. disci tubulosis 5-dentatis masculis seu strobilibus. Invol. squamæ 1-seriales concavæ et achænia radii semi-involventes. Recept. planum paleaceum aut saltè inter radium et discum paleas 1-seriales gerens et medio paleaceum punctatum. Cor. bas. glabræ. Styli rami obtusiusculi. Achænia radii obovata ferè obcompressa extus convexa intus plana glabra calva, disci abortiva. — Herba Californicæ erectæ ramosæ basi suffruticosæ pilosiusculæ interdum glanduliferæ. Folia alterna conferta lanceolata aut linearia, integra aut serrata. Capitula terminalia solitaria aut pauca, bracteis foliaceis plurimis cincta. Flores flavi. — Genus omninò medium inter *Mazarian* et *Hartmanniam*. Nomen ex *ἡμιζωνος* (semicinctus) alludat achænia invol. squamis semicincta.

#### Sect. I. MADARIOIDES.

Recept. margine paleas 1-seriales gerens, medio nudum planum punctatum.

1. *H. CONGESTA*, caule herbaceo erecto ramoso, ramis pilis glandulosis et eglandulosis molliter hispidulis, foliis lineari-lanceolatis parè villosis-glandulosis capitulis bracteatis congestis, achæniis obovatis apice obtusis. — in Calif. bot. reg. legit cl. Douglas. Achænia nigra, potiùs obcompressa quàm lateribus compressa, matura basi producta in stipitem rostelliformem insigniter recurvam areolam albam gerentem. Recept. paleæ subcoriaceæ foliaceæ. (v. s. comm. ab hon. Soc. hort. Lond.)
2. *H. ANGUSTIFOLIA*, caule ramosissimo imò basi suffruticuloso, ramulis ferè involucriisque minutè puberulis, foliis bracteisque angustè linearibus integerrimis, capitulis bracteatis subsolitariis, achæniis obovatis apice cono luto acuto mucronatis. ③ in Californiâ legit cl. Douglas. Recept. paleæ angustè membranaceæ. Achænia estipitata areolâ obliquâ, supernè cunulo breviter coronata. Eandem nō fallor video in h. Hænke, sed *Regiomontana* dicitur. (v. s. comm. cum priorib.)

#### Sect. II. OLOCARPHA.

Paleæ recept. discum totum occupantes.

3. *H. LUZULIFOLIA*, caule molliter hirsutulo, foliis inferioribus oppositis, superioribus alternis, elongato-linearibus plurinerviis subdenticulatis molliter villosis eglandulosis.

delosis, corymbo laxo, invol. squamis hispidulis, achæniis radii muticis. — in Novâ-Californiâ legit cl. Douglas. Ligulæ 15-20. Capitula parvisimâ bractea. Achænia cætipitata areolâ laterali. (v. s. comm. ab hon. Soc. hort. Lond.)

4. *H. MACRADENIA*, caule suffruticoso ramosissimo, ramulis foliisque pilosiusculis, foliis lineari-lanceolatis, infer. parce serratis, cæteris integris, ramis super. bracteisque glandulâ grossâ truncatâ terminatis, invol. squamis grossè et callosè inciso-serratis, paleis apice subulatis ciliatis, achæniis obovato-turbinatis apice incurvis rostro subadscendenti. ♂ in Californiâ legit cl. Douglas. Species priori primâ fronte (simillima sed verè distinctissima et structurâ in ordine anomalâ spectabilis. Affinis *Calycadeniæ* sed ligulæ planæ, etc. (v. s. comm. ab hon. Soc. hort. Lond.)

CDXXXVI. *OXYURA DC.* in *Lindl. nat. syst. bot. ed. 2. p. 256.*

Capitulum multiflorum radiatum, ligulis 10-12 uniserialibus foemineis, fl. disci tubulosis 5-dentatis bisexualibus, centralibus plurimis abortivis. Invol. squamæ 1-seriales basi rigidè ciliatæ convolutæ et achænia radii omninò involventes, supernè in appendicem linearem foliaceam productæ. Recept. totum paleis membranaceis inter flores onustum. Ligulæ basi pubentes apice grossè dentatæ. Tubuli obcompressi ad lobulos pubentes. Anth. nigricantes. Styli disci rami in appendicem hispidam acutissimam producti. Achænia omnia obcompressa obovato-oblonga glabra calva, centralia sæpè abortiva. — Herba Californica annua erecta *Chrysanthemum* segetum habitu referens. Caulis teres purpurascens glaber. Folia alterna ciliata pinnatifida, lobis 5-8-jugis integris. Capitula ramulos apice nudiusculos hirsutos terminantia. Flores flavi. Bractæ paucæ ciliatæ. — Nomen ex *ὄξυς*, acutus, et *ὤρα*, cauda.

1. *O. CHRYSANTHEMOIDES*. (I) in Novâ-Californiâ legit cl. Douglas. Radix simplex perpendicularis. Caulis circ. pedalis. (v. s. comm. ab hon. Soc. hort. Lond.)

CDXXXVII. *HARTMANNIA*.

Capitulum pluri- aut multiflorum heterogamum, fl. radii uniseriatis foemineis ligulatis grossè apice dentatis, disci hermaphr. tubulosis 5-dentatis. Invol. squamis oblongo-lanceolatis acutis convolutis flores radii semiamplectentibus, bracteis nonnullis linearibus extus cinctis. Recept. paleaceum, paleis squamas referentibus sed planiusculis. Cor. disci vix apice puberulæ. Antheræ nigricantes. Stylus, disci ramis cono acuto apiculatis. Achænia radii tri- seu tetragona obovata subpedicellata, dorso subconvexa, apice calva, areolâ terminali obliquâ interiùs sitâ interdum papilloso-exsertâ, disci ovali-oblonga obcompressa rarò calva sæpiùs paleis 5 membranaceis enerviis obtusis corollâ dimidiò brevioribus coronata. — Herba Californicæ erectæ. Caulis pilis patulis hispidus simplex apice corymboso-ramosus. Folia alterna, infima interdum opposita, sessilia lineari-oblonga, pinnatifida aut serrata pubera. Capitula ad apices ramorum corymbosa aut fasciculata. Cor. flavæ. — Genus meo sensu ex habitu naturale, sed in sectiones tot quot species faciliè distinguendum, Helenieis ex pappo paleaceo et antheris nigris pertinet, sed transitum ad Anthemideas exhibet! — Genus distinctissimum dixi in honorem cl. peregrinatoris Jos. Adolph. Hartmann qui, anno 1739, Californiam adiit et ejus (descript. edidit.

1. *H. FASCICULATA*, foliis inciso-dentatis inâ basi obtusè et brevè auriculatis,

- P. 66, lin. 47, sub D. reticulato, adde syn.: *Rolinkia centas*  
ind. 2. p. 14. t. 14.
- P. 109, lin. 7, t. 3?, lege t. 5?
- P. 110, lin. 17, MIKANOIDES, lege MIKANIOIDES.
- P. 125, lin. 11, dele syn. Nuttallii.
- P. 255, lin. 27, adde syn.: *Galatella punctata* Lindl. bot. reg.
- P. 275, lin. 25, dele syn. Lessingii.
- P. 283, lin. 8, adde: DC. non Bess. — *Heterochaeta* Bess. est A
- P. 295, lin. 9, adde: *Erigeron filiforme* Spreng. est ex Schleg  
pappus pinifolius Less. ined.
- P. 298, lin. 14, disci, lege radii.
- P. 313, lin. 38, adde syn.: DC. pl. rar. jard. Gen. not. VII, p. 1
- P. 374, lin. 20, LESTADIA, lege LÆSTADIA.
- P. 376, lin. 27, LÆNNECIA, lege LAENNECIA.
- P. 398, lin. 8 à fine, Tursonia, lege Tursenia.
- P. 415, lin. 11, dele syn. Sprengelii ex Schlect. in litt., ad Lucil  
referendum.
- P. 428, lin. 8 à fine, t. 27, lege t. 47, et dele *Chrysocoma punct*
- P. 449, lin. 41, dele et *Chlænobolus*.
- P. 485, lin. 47, adde syn.: Hook. bot. mag. t. 3466.
- P. 503, lin. 45, *Petrantha*, lege *Tetrantha*.
- P. 511, lin. 2, *Guardiola*, lege *Guandiola*.
- P. 521, lin. 35, Poir., lege Poiteau.
- P. 524, lin. 37, ACANTHOLENA, lege CENTROLENA.
- P. 531, lin. 11, xovia, pulveris, lege xovis, pulvis.
- P. 543, lin. ult., *Jægera* Raeusch. et Gisek. est genus inter Scita
- P. 551, lin. 38, adde syn.: DC. pl. rar. jard. Gen. not. VII, p. 5
- P. 595, lin. 10, adde syn.: DC. pl. rar. jard. Gen. not. VII, p. 5

# INDEX NOMINUM ET SYNONYMORUM

## PARTIS QUINTÆ.

*Obs.* Nomina admissa litteris romanis, Synonyma italicis traduntur, nempe in utraq[ue] classe:

Ordinalia majoribus,  
Tribus alia majusculis,  
Generica vulgaribus,  
Sectionalia minoribus.

|                                |              |                               |              |
|--------------------------------|--------------|-------------------------------|--------------|
| <i>Acantholoma DC.</i>         | 460          | <i>Albertinia Spreng.</i>     | 80           |
| <i>Acanthosperma Arrab.</i>    | 3            | ALBERTINIEÆ DC.               | 78           |
| <i>Acanthospermum Schrank.</i> | 521          | <i>Aleina Cav.</i>            | 520          |
| <i>Acanthoxanthium DC.</i>     | 523          | <i>Aleina DC.</i>             | <i>ibid.</i> |
| ACHYRIDEÆ DC.                  | 212, 312     | <i>Alcinia Steud.</i>         | <i>ibid.</i> |
| <i>Achyrocoma Cass.</i>        | 15           | <i>Alciope DC.</i>            | 209          |
| <i>Achyropappus H. B. K.</i>   | 654          | <i>Aldama Lallav. et Lex.</i> | 561          |
| <i>Acicarpha Juss.</i>         | 3            | <i>Alibum Less.</i>           | 97           |
| <i>Acilepis Don.</i>           | 15           | <i>Alix Comm.</i>             | 319          |
| <i>Acilepis DC.</i>            | <i>ibid.</i> | <i>Allagopappus Cass.</i>     | 477          |
| <i>Aciphylla DC.</i>           | 641          | <i>Allagopappus DC.</i>       | <i>ibid.</i> |
| <i>Acispermum Neck.</i>        | 569          | <i>Allocarpus H. B. K.</i>    | 676          |
| <i>Acmella Rich.</i>           | 620          | <i>Alloispermum Willd.</i>    | <i>ibid.</i> |
| <i>Acmella DC.</i>             | <i>ibid.</i> | <i>Alomia H. B. K.</i>        | 105          |
| <i>Acosta DC.</i>              | 90           | ALOMIÆ Less.                  | 103          |
| <i>Actinea Juss.</i>           | 662          | <i>Alymania Neck.</i>         | 515          |
| <i>Actinella Pers.</i>         | <i>ibid.</i> | <i>Alymania DC.</i>           | <i>ibid.</i> |
| <i>Actinella Less.</i>         | <i>ibid.</i> | <i>Amblyolepis DC.</i>        | 667          |
| <i>Actinolepis DC.</i>         | 655          | <i>Ambrosia Linn.</i>         | 525          |
| <i>Actinomeris Raf.</i>        | 575          | AMBROSIACÆ Less.              | 500          |
| <i>Actinomeris Nutt.</i>       | <i>ibid.</i> | AMBROSIAE Cass. Less.         | 498, 522     |
| ACTINOPHYTUM Neck.             | 4            | AMELLEÆ DC.                   | 211, 213     |
| <i>Actinoptera DC.</i>         | 543          | AMELLOIDÆ Less.               | 213          |
| <i>Actinospermum Ell.</i>      | 653          | <i>Amellus Adans.</i>         | 226          |
| <i>Adenocaulon Hook.</i>       | 207          | <i>Amellus Cass.</i>          | 213          |
| <i>Adenocyclus Less.</i>       | 10           | <i>Amellus P. Br.</i>         | 544          |
| <i>Adenolepis Less.</i>        | 607          | <i>Amellus Linn.</i>          | 98, 213      |
| <i>Adenophyllum Pers.</i>      | 630          | <i>Ampherephus H. B. K.</i>   | 70           |
| <i>Adenostemma Forst.</i>      | 110          | <i>Amphiaschysis DC.</i>      | 313          |
| ADENOSTYLEÆ Cass. DC.          | 126          | <i>Amphicalca DC.</i>         | 672          |
| <i>Adenostyles Cass.</i>       | 203          | <i>Amphirhapis DC.</i>        | 343          |
| <i>Agarista DC.</i>            | 569          | <i>Amphistephium DC.</i>      | 273          |
| <i>Agathæa Cass.</i>           | 223          | <i>Anacis Schrank.</i>        | 567          |
| AGERATEÆ Less.                 | 107          | <i>Anaglypha DC.</i>          | 311          |
| <i>Ageratoides DC.</i>         | 108          | <i>Anaitis DC.</i>            | 626          |
| <i>Ageratum Linn.</i>          | <i>ibid.</i> | <i>Andrieuxia DC.</i>         | 559          |
| <i>Ageratum Auct.</i>          | 107, 108     | <i>Andromachia Cass.</i>      | 96           |
| <i>Aglossa DC.</i>             | 543          | <i>Andromachia H. B. K.</i>   | 95           |
| <i>Agrianthus Mart.</i>        | 125          | <i>Andryaloides DC.</i>       | 377          |
| <i>Aiolothea DC.</i>           | 508          | ANGIANTHÆÆ DC.                | 500          |
| <i>Alarçonia DC.</i>           | 537          | <i>Anhebecarpa DC.</i>        | 221          |

|                             |               |                                   |          |
|-----------------------------|---------------|-----------------------------------|----------|
| <i>Anisochæta</i> DC.       | 109           | BELLIDÆ DC.                       | 212, 322 |
| <i>Anisotrichia</i> DC.     | 80            | <i>Bellis</i> Linn.               | 324      |
| <i>Anoplostephium</i> DC.   | 560           | <i>Bellium</i> Linn.              | 325      |
| <i>Anonymus</i> Walt.       | 128           | <i>Berlandiera</i> DC.            | 517      |
| ANTENNARIÆ.                 | 500           | <i>Berthelotia</i> DC.            | 375      |
| <i>Anthalmum</i> Neck.      | 487           | <i>Bidens</i> Linn.               | 523      |
| ANTHEMIDÆ.                  | 499           | BIDENTIDÆ Less.                   | 498, 523 |
| <i>Anthemopsis</i> Boj.     | 547           | <i>Bigelovia</i> DC.              | 329      |
| <i>Anthemoides</i> Less.    | 681           | <i>Biotia</i> DC.                 | 224      |
| <i>Anvillea</i> DC.         | 487           | <i>Biotia</i> Cass.               | 69       |
| <i>Apalus</i> DC.           | 507           | <i>Blainvillea</i> Cass.          | 492      |
| <i>Aphanostephus</i> DC.    | 310           | <i>Blanchetia</i> DC.             | 5        |
| <i>Apodiscus</i> DC.        | 350           | <i>Blepharipappus</i> Hook.       | 679      |
| <i>Aplopappus</i> Cass.     | 345           | <i>Blepharisperмум</i> Wight, DC. | 328      |
| <i>Arctogeron</i> DC.       | 260           | <i>Blepharodoa</i> DC.            | 248      |
| <i>Arctotis</i> Auct.       | 681, 688      | <i>Blumea</i> DC.                 | 424      |
| <i>Argophylla</i> DC.       | 266           | <i>Bobartia</i> Petiv.            | 554      |
| <i>Argyrochæta</i> Cav.     | 532           | <i>Boebera</i> Less.              | 639      |
| <i>Argyrochæta</i> DC.      | ibid.         | <i>Boebera</i> Willd.             | 224      |
| <i>Argyroxiphium</i> DC.    | 668           | <i>Boebera</i> DC.                | 649      |
| <i>Armania</i> Bart.        | 576           | <i>Boeberoides</i> DC.            | 224      |
| <i>Arthenachne</i> Cass.    | 328           | <i>Bojeria</i> DC.                | 94       |
| ARTEMISIÆ Less.             | 499           | BOJERIÆ DC.                       | 224      |
| <i>Artemisioides</i> DC.    | 105           | <i>Boltonia</i> L'Hér.            | 322      |
| <i>Ascaricida</i> Cass.     | 15, 60        | <i>Bombycilana</i> DC.            | 420      |
| <i>Ascaricida</i> DC.       | 60            | BOOPIDÆ Cass.                     | 1        |
| <i>Aspilula</i> Pet.-Th.    | 561           | <i>Boopis</i> Juss.               | 2        |
| <i>Astemma</i> Less.        | 502           | <i>Borrchia</i> Adans.            | 428      |
| <i>Aster</i> Adans.         | 253           | <i>Botryadenia</i> F. et M.       | 329      |
| <i>Aster</i> Nees.          | 226           | <i>Brachycome</i> Cass.           | 325      |
| ASTERFÆ.                    | 211           | <i>Brachycome</i> Gaud.           | 280      |
| ASTERINEÆ DC.               | 211, 213      | <i>Brachycome</i> Cun.            | 325      |
| ASTEROIDÆ Less.             | 8, 211        | <i>Brachyglossa</i> DC.           | 265      |
| <i>Asteriscus</i> Moench.   | 486           | <i>Brachyglottis</i> Forst.       | 219      |
| <i>Asteroides</i> Tourn.    | 483, 485, 486 | <i>Brachylæna</i> Brown.          | 429      |
| <i>Asteromea</i> Blum.      | 302           | <i>Brachyris</i> Nutt.            | 312      |
| <i>Asteropsis</i> Less.     | 260           | <i>Brachyscoma</i> Cass.          | 325      |
| <i>Athanasia</i> Walt.      | 680           | <i>Brauneria</i> Neck.            | 553      |
| ATHANASIÆ.                  | 499           | <i>Brickellia</i> Ell.            | 182      |
| <i>Athenæa</i> Adans.       | 11            | <i>Brotera</i> Spreng.            | 635      |
| <i>Athroisma</i> DC.        | 368           | <i>Broteroa</i> DC.               | 222      |
| <i>Athronia</i> Neck.       | 620           | <i>Babonium</i> DC.               | 463      |
| <i>Aurelia</i> Cass.        | 314           | <i>Bulbostylis</i> DC.            | 138      |
| BACCHARIDÆ Less.            | 212, 368      | BUPHTHALMÆ Less.                  | 213, 483 |
| BACCHAREÆ DC.               | 328           | <i>Buphthalmum</i> Cass.          | 423      |
| <i>Baccharis</i> Linn.      | ibid.         | <i>Burrielia</i> DC.              | 663      |
| <i>Baccharoides</i> Linn.   | 60            | <i>Bustamenta</i> bot. mex.       | 166-168  |
| <i>Bahia</i> Lag.           | 656           | <i>Bustia</i> Adans.              | 425      |
| <i>Bailliera</i> Cass.      | 505           | <i>Cacalia</i> Tourn.             | 203      |
| <i>Balsisia</i> Willd.      | 679           | <i>Cacosmia</i> H. B. K.          | 9        |
| BALBISIÆ DC.                | 500           | <i>Cæsulia</i> Roxb.              | 222      |
| <i>Balduina</i> Nutt.       | 652           | CAESULIÆ DC.                      | 222      |
| <i>Balsamorhiza</i> Hook.   | 551           | <i>Calea</i> R. Br.               | 671      |
| <i>Baltimora</i> Linn.      | 509           | <i>Calea</i> Linn.                | 222      |
| BALTIMOREÆ Cass.            | 500           | <i>Caleacte</i> R. Br.            | 673      |
| <i>Bartolina</i> Adans.     | 641, 679      | <i>Caleacte</i> Less.             | 679      |
| <i>Bechium</i> DC.          | 70            | <i>Caleacte</i> DC.               | 673      |
| <i>Bellidiastrum</i> Michx. | 226           | <i>Calebrachys</i> Cass.          | 679      |
| BELLIDÆ DC.                 | 212, 304      | <i>Calebrachys</i> DC.            | 679      |



|                               |          |                                           |          |
|-------------------------------|----------|-------------------------------------------|----------|
| <i>CALEVEE</i> Cass.          | 668      | <i>Chromolena</i> DC.                     | 133      |
| <i>Caliceras</i> Cav.         | 2        | <i>Chronolena</i> DC.                     | 397      |
| <i>Calimeris</i> Cass. Nees.  | 258      | <i>Chronopappus</i> DC.                   | 84       |
| <i>Callilepis</i> DC.         | 671      | <i>Cryptopetalum</i> Cass.                | 101      |
| <i>Calliopsis</i> Reichenb.   | 568      | <i>Cryptopetalum</i> DC.                  | ibid.    |
| <i>Callistemma</i> Cass.      | 274      | <i>Chrysactinium</i> H. B. K.             | 97       |
| <i>Callistephus</i> Cass.     | ibid.    | <i>Chrysanthellina</i> Cass.              | 630      |
| <i>Callophora</i> DC.         | 542      | <i>Chrysanthellum</i> Rich.               | 630      |
| <i>Calonnea</i> Buchoz.       | 651      | CHRYSANTHEMEAE.                           | 499      |
| <i>Calotis</i> B. Br.         | 302      | <i>Chrysis</i> Ren.                       | 585      |
| <i>Calycadenia</i> DC.        | 696      | <i>Chrysocoma</i> Cass.                   | 353      |
| <i>Calycera</i> Rich.         | 2        | <i>Chrysocoma</i> Linn.                   | 351, 353 |
| CALYCEREAE R. Br.             | 1        | CHRYSOCOMEAE DC.                          | 212, 310 |
| <i>Calycium</i> Ell.          | 316      | <i>Chrysogonum</i> Linn.                  | 510      |
| <i>Calydermos</i> Lag.        | 669      | <i>Chrysomelea</i> Tausch.                | 569      |
| <i>Calydermos</i> Cass.       | ibid.    | <i>Chrysophania</i> Kunth.                | 553      |
| <i>Calydermos</i> Spreng.     | 11       | CHRYSOPSIDAE DC.                          | 212, 326 |
| <i>Calypocarpus</i> Less.     | 629      | <i>Chrysopsis</i> Nutt.                   | 326      |
| <i>Campuloclinium</i> DC.     | 136      | <i>Chrysospharium</i> Willd.              | 673      |
| <i>Campylothea</i> Cass.      | 593      | <i>Chrysostemma</i> Less.                 | 567      |
| <i>Camutia</i> Bonat.         | 520      | <i>Chthonia</i> Cass.                     | 98       |
| <i>Cappa</i> DC.              | 469      | <i>Chylodia</i> Rich.                     | 563      |
| <i>Carelia</i> Less.          | 125      | CICHORACEAE Paill.                        | 4, 9     |
| <i>Curelia</i> Ponted.        | 108      | <i>Clairvillea</i> DC.                    | 636      |
| <i>Carphephorus</i> Cass.     | 132      | <i>Clavigera</i> DC.                      | 127      |
| <i>Carpophostephium</i> Cass. | 678      | <i>Clibadium</i> Linn.                    | 565      |
| <i>Curtesia</i> Cass.         | 71       | <i>Clomenocoma</i> Cass.                  | 641      |
| <i>Carthamus</i> Hill.        | ibid.    | <i>Celestina</i> Cass.                    | 107      |
| CASSINIEAE.                   | 500      | <i>Coleosanthus</i> Cass.                 | 136, 138 |
| <i>Celmisia</i> DC.           | 209      | <i>Colloea</i> Spreng                     | 630      |
| <i>Celmisia</i> Cass.         | ibid.    | <i>Commidendrum</i> DC.                   | 344      |
| <i>Centauropsis</i> Bojer.    | 93       | COMPOSITAE Paill.                         | 4        |
| <i>Centipeda</i> Less.        | 371      | COMPOSITIFLORE Gært.                      | ibid.    |
| <i>Centropalus</i> Cass.      | 15, 60   | <i>Goniothele</i> DC.                     | 531      |
| <i>Centratherum</i> Cass.     | 70       | <i>Conoclinium</i> DC.                    | 135      |
| <i>Centratherum</i> Less.     | 69       | <i>Contarena</i> Adans.                   | 88       |
| <i>Centrocarpha</i> Don.      | 555      | <i>Conyza</i> Less.                       | 377      |
| <i>Centrolæna</i> DC.         | 524      | <i>Conyza</i> Ait. Lam.                   | 69       |
| <i>Centrospermum</i> Kunth.   | 521      | <i>Conyza</i> Linn.                       | 377, 396 |
| <i>Cephalophora</i> Cav.      | 661      | <i>Conyza</i> Wall.                       | 77       |
| <i>Ceratocephalus</i> Vaill.  | 593      | CONYZEAE Less.                            | 212, 368 |
| <i>Ceratochilæna</i> DC.      | 522      | COREOPSIDAE Less.                         | 498, 569 |
| <i>Cercostylus</i> Less.      | 660      | <i>Coreopsis</i> Linn.                    | 569      |
| <i>Ceruana</i> Forsk.         | 488      | <i>Coreopsis</i> Auct. 562, 568, 569, 575 | 575      |
| <i>Ceruchis</i> Gært.         | 620      | <i>Coreopsoides</i> Moench.               | 569      |
| <i>Chanaetis</i> DC.          | 659      | <i>Corethrogyne</i> DC.                   | 216      |
| <i>Chataetis</i> DC.          | 317      | <i>Corona-solis</i> Tourn.                | 585      |
| <i>Chætopappa</i> DC.         | 301      | <i>Corvisartia</i> Merat.                 | 463      |
| <i>Chætophora</i> Nutt.       | ibid.    | <i>Corvisartia</i> DC.                    | ibid.    |
| <i>Chakintella</i> Cass.      | 563      | CORYMBIFERE Paill.                        | 4        |
| <i>Chamaestephanum</i> Willd. | 654      | <i>Corymbium</i> Linn.                    | 88       |
| <i>Charies</i> Cass.          | 350      | <i>Cosmea</i> Willd.                      | 606      |
| <i>Chiliadenus</i> Cass.      | 476      | <i>Cosmos</i> Cuv.                        | ibid.    |
| <i>Chiliadenus</i> DC.        | ibid.    | <i>Cosmos</i> Pers.                       | ibid.    |
| <i>Chiliophyllum</i> DC.      | 554      | COTULEAE Less.                            | 499      |
| <i>Chiliotrichum</i> Cass.    | 216      | <i>Crassicola</i> DC.                     | 373      |
| <i>Chlenobolus</i> Cass.      | 449, 453 | <i>Crassina</i> Scop.                     | 534      |
| <i>Chlenobolus</i> DC.        | 453      | <i>Crinita</i> Moench.                    | 351      |
| <i>Chresta</i> Arrab.         | 85       | <i>Crinitaria</i> Cass. Less.             | ibid.    |
| <i>Chromocheta</i> DC.        | 351      | <i>Critonia</i> P. Br.                    | 140      |



|                           |          |     |
|---------------------------|----------|-----|
| Eucalypta DC.             |          | 67  |
| Euerigeron DC.            |          | 619 |
| Euvax DC.                 |          | 484 |
| EUGALINOGEEAE DC.         | 499, 6A  | 320 |
| Eugamelia fl. mex.        | 502      | 298 |
| Euharpalum DC.            | 583      | 459 |
| EUHELENIEAE DC.           | 653      | 460 |
| Eubeliopsis DC.           | 550      | 368 |
| Eubymenoxys DC.           | 661      | 654 |
| EUINULEAE DC.             | 463      | 187 |
| Eujasonia DC.             | 476      | 503 |
| Euliatrix DC.             | 128      | 502 |
| Eulychnophora DC.         | 79       | 298 |
| Eumelampodium DC.         | 518      | 671 |
| Eumonosis DC.             | 77       | 308 |
| EUPATORIACEÆ Less.        | 8, 103   | 485 |
| EUPATORIÆAE DC.           | 103      | 646 |
| EUPATORIÆAE Less.         | 106      | 457 |
| Eupatorium Linn.          | 141      | 455 |
| Eupatorium Auct.          | 141, 187 | 550 |
| Eupatoriophalacron Vaill. | 489      | 325 |
| EUPECTIDEAE DC.           | 98       | 77  |
| Euphanja DC.              | 114      | 65  |
| Euphrosyne DC.            | 530      | 1   |
| Eupiqueria DC.            | 104      |     |
| Euporophyllum DC.         | 648      |     |
| Eupytrocoma DC.           | 350      |     |
| Eurybia Cass.             | 265      |     |
| Eurybia Nees.             | 264      |     |
| Eurybiopsis DC.           | 260      |     |
| EUSENECIONEAE DC.         | 500      |     |
| Eusphæranthus DC.         | 369      |     |
| Euthamia Cass.            | 330      |     |
| Euthamia DC.              | 341      |     |
| Euthamia Nutt.            | ibid.    |     |
| Eutarchonanthus DC.       | 431      |     |
| Eutrageroceras DC.        | 533      |     |
| EUTUSSILAGINEAE DC.       | 207      |     |
| EUVERNONIEAE DC.          | 10       |     |
| Euxanthioides DC.         | 524      |     |
| Euxanthium DC.            | 523      |     |
| Euxenia Cham.             | 501      |     |
| EUXENIÆAE DC.             | 498, 500 |     |
| Evax Gærtn.               | 458      |     |
| Fæea Spreng.              | 612      |     |
| Felicia Cass.             | 218      |     |
| Felicia DC.               | ibid.    |     |
| Ferdinanda Lag.           | 552      |     |
| Filago Willd.             | 458      |     |
| Fimbrillaria Cass.        | 387      |     |
| Fimbrillaria DC.          | ibid.    |     |
| Flaveria Juss.            | 635      |     |
| Flaveria Juss.            | 104      |     |
| FLAVERIEAE Less.          | 498, 605 |     |
| FLOSCULOSÆ Tourr.         | 4        |     |
| Florestina Cass.          | 655      |     |
| Flourensia DC.            | 590      |     |
| Fougeria Moench.          | 509      |     |
| Fougerouxia DC.           | ibid.    |     |
| Galea                     |          | 619 |
| Galea                     |          | 484 |
| Galea                     |          | 320 |
| Galea                     |          | 298 |
| Galea                     |          | 459 |
| Galea                     |          | 460 |
| Galea                     |          | 368 |
| Galea                     |          | 654 |
| Galea                     |          | 187 |
| Galea                     |          | 503 |
| Galea                     |          | 502 |
| Galea                     |          | 298 |
| Galea                     |          | 671 |
| GALINSOGIA L.             |          | 308 |
| GALINSOGIA L.             |          | 485 |
| Galordia Tourr.           |          | 646 |
| Gamocarpha L.             |          | 457 |
| Garuleum Cass.            |          | 455 |
| Geigeria Less.            |          | 550 |
| Georgia Spreng.           |          | 325 |
| GEORGIA Willd.            |          | 77  |
| GLOSSARIPHYTUM            |          | 65  |
| Glossocardia Cass.        |          | 1   |
| Glossogyne Cass.          |          |     |
| Glaphiphyllaea DC.        |          |     |
| Glutinaria Comm.          |          |     |
| Glutinaria DC.            |          |     |
| GNAPHALIEAE.              |          |     |
| Gnaphalium Vaill.         |          |     |
| Gnaphalodes Tourr.        |          |     |
| Graemia Hook.             |          |     |
| Grangea Adans.            |          |     |
| GRANGEINEAE DC.           |          |     |
| Grindelia Willd.          |          |     |
| Guardiola Steud.          |          |     |
| Guardiola H. B.           |          |     |
| Guizotia Cass.            |          |     |
| Gundelia Tourr.           |          |     |
| Gundelsheimera Cass.      |          |     |
| Guttierrezia Lag.         |          |     |
| Gymnactis DC.             |          |     |
| Gymnanthemum Cass.        |          |     |
| Gymnanthemum DC.          |          |     |
| Gymnarhena Desf.          |          |     |
| Gymnocoronis DC.          |          |     |
| Gymnolœna DC.             |          |     |
| Gymnoloma Ker.            |          |     |
| Gymnolomia H. B. K.       |          |     |
| Gymnopsis DC.             |          |     |
| Gymnosperma Less.         |          |     |
| GYMNOSPERMEAE DC.         |          |     |
| Gymnostephium Less.       |          |     |
| Gynema Raf.               |          |     |
| Gyneheteria Willd.        |          |     |
| Gyneteria Spreng.         |          |     |
| Hacub Vaill.              |          |     |
| Hamulium Cass.            |          |     |
| Hamulium DC.              |          |     |

- Haplostephium Mart.* 583  
*Harpalium Cass.* 583  
*Harpalizia DC.* 584  
*Hartmannia DC.* 693  
*Haxtonia Hortul.* 271  
*Haynea Spreng.* 16  
*Haynea Willd.* 14  
*Hebecarpæa DC.* 219  
*Hebeclinium DC.* 136  
*Hebephora DC.* 453  
*Hecubæa DC.* 665  
*Hectorea DC.* 95  
*Heleastrum DC.* 263  
*HELENIEAE Cass.* 499, 651  
*Helenia Linn.* 666  
*Helenia DC.* *ibid.*  
*Helenium Cass.* *ibid.*  
*Helenium Linn.* 665  
*HELIANTHEAE Less.* 498, 534  
*Helianthus Linn.* 585  
*Helichroa Raf.* 554  
*HELICHRYSÆAE.* 500  
*HELIOPSIDEAE DC.* 458, 534  
*Heliopsis Pers.* 550  
*Heliophthalmum Raf.* 555  
*Hemiachyris DC.* 313  
*Hemizonia DC.* 692  
*Henanthus Less.* 356  
*Henotogyna DC.* 432  
*Henricia Cass.* 362  
*Herderia Cass.* 13  
*Heterochaeta DC.* 282  
*Heterocoma DC.* 14  
*HETEROCOMEAE DC.* *ibid.*  
*HETEROPAPPEAE DC.* 212, 297  
*Heteropappus Less.* 297  
*Heterosperma Cav.* 632  
*Heterospermum Willd.* *ibid.*  
*HETEROTHALAMEAE DC.* 216  
*Heterothalamus Less.* *ibid.*  
*Heterotheca Cass.* 316  
*HETEROTHECEAE DC.* 212, 316  
*Hidalgoa Less.* 511  
*Hidalgoa Lallav.* 521  
*Hingcha Roxb.* 636  
*HIPPIEAE.* 499  
*Hololepis DC.* 16  
*Holophyllæa DC.* 670  
*Homochroma DC.* 324  
*Homogyne Cass.* 204  
*Honrebeckia Corn.* 346  
*Hoplophyllum DC.* 73  
*Hopkirkia DC.* 660  
*Hopkirkia Spreng.* 493  
*Hunteria fl. mex.* 649  
*Hunteria DC.* *ibid.*  
*Hybridella Cass.* 554  
*Hymenatherum Cass.* 641  
*Hymenopappus L'Hér.* 658  
*Hymenoxys Cass.* 661  
*Hysterophorus Vaill.*  
*Ichthyothere Mart.*  
*Inula Gärtn.*  
*Inula Linn.* 463, 469  
*INULEAE Cass.*  
*Inulopsis DC.*  
*Iphiona Cass.*  
*Isocarpha R. Br.*  
*Isocarphoides DC.*  
*Isomeria Wight.*  
*Isonema Cass.*  
*Isostigma Less.*  
*Isotrichia DC.*  
*Iva Linn.*  
*IVEAE Cass.*  
*Ixauchenus Cass.*  
*Jageria H. B. K.*  
*Jasonia Cass.*  
*Jaumea Pers.*  
*Kahiria Forsk.*  
*Kalimeris Cass.*  
*Kallias Cass.*  
*Kallias DC.*  
*Karelinia Less.*  
*Kaulfussia Nees.*  
*Keerlia DC.*  
*Kerneria Mönch.*  
*Kleinia Jacq.*  
*Kleinia Juss.*  
*Kugaia DC.*  
*Kuhnia Linn.*  
*Kyberia Neck.*  
*Laennecia Cass.*  
*Laestadia Kunth.*  
*Lagasca DC.*  
*Lagasca Cav.*  
*Lagascea H. B. K.*  
*Lagenifera Cass.*  
*Lagenophora Cass.*  
*Lasianthæa DC.*  
*Lasianthus Zucc.*  
*Lasthenia Cass.*  
*Latreillea DC.*  
*Lavenia Swartz.*  
*Lavenia Wall.*  
*Laxmannia Forst.*  
*Leachea Cass.*  
*Lebetina Cass.*  
*Leichanum DC.*  
*Leighia Cass.*  
*Leighia Scop.*  
*Lejica Hill.*  
*Leiogonia DC.*  
*Lemmatium DC.*  
*Leontophthalmum Less.*  
*Leontophthalmum Willd.*

NOMINUM ET SYNONYMORUM.

703

|                              |          |                             |          |
|------------------------------|----------|-----------------------------|----------|
| <i>Lepachys</i> Raf.         | 558      | <i>Meyeria</i> DC.          | 670      |
| <i>Lepachys</i> DC.          | 558      | <i>Micractis</i> DC.        | 619      |
| <i>Lepidaploa</i> Cass.      | 15, 26   | <i>Microelium</i> Forsk.    | 489      |
| <i>Lepidaploa</i> DC.        | 26       | <i>Microglossa</i> DC.      | 320      |
| <i>Lepidopappus</i> fl. mex. | 655      | <i>Microgyne</i> Less.      | 296      |
| <i>Lepidophyllum</i> Cass.   | 314      | <i>Micropsis</i> DC.        | 459      |
| <i>Lepidopogon</i> Tausch.   | 458      | <i>Micropus</i> Linn.       | 460      |
| <i>Leptocarpa</i> DC.        | 495      | <i>Microtrichia</i> DC.     | 366      |
| <i>Leptocoma</i> Less.       | 280      | <i>Nieria</i> Lallav.       | 654      |
| <i>Leptoderis</i> DC.        | 3-3      | <i>Mikania</i> Willd.       | 187      |
| <i>Leptogyne</i> Ell.        | 450      | <i>Milleria</i> Cass.       | 503      |
| <i>Leptopoda</i> Nutt.       | 653      | MILLERIEÆ DC.               | 498, 502 |
| <i>Leptospermoides</i> DC.   | 17       | <i>Minuria</i> DC.          | 298      |
| <i>Leptosyne</i> DC.         | 531      | <i>Mocinna</i> Lag.         | 671      |
| <i>Leptothamnus</i> DC.      | 367      | <i>Molina</i> R. et Pav.    | 398      |
| <i>Lessingia</i> Cham.       | 351      | <i>Molpadia</i> Cass.       | 485      |
| <i>Leucopsis</i> DC.         | 348      | <i>Monactis</i> H. B. K.    | 646      |
| LEYSSERIEÆ.                  | 500      | <i>Monarrhenus</i> Cass.    | 457      |
| LIABEÆ Cass.                 | 94       | <i>Monoteles</i> Labill.    | 455      |
| <i>Liabum</i> Adans.         | 96       | <i>Monodonta</i> DC.        | 559      |
| <i>Liatris</i> Schreb.       | 128      | <i>Monogyria</i> DC.        | 325      |
| <i>Limbarda</i> Adans.       | 470      | <i>Monosis</i> DC.          | 77       |
| <i>Limbarda</i> DC.          | ibid.    | <i>Montagnæa</i> DC.        | 664      |
| <i>Linosyris</i> DC.         | 351      | <i>Montana</i> Lallav.      | ibid.    |
| <i>Lipochæta</i> DC.         | 610      | <i>Munychia</i> Cass.       | 222      |
| <i>Lipotriche</i> R. Br.     | 544      | MUTISIAEÆ.                  | 9        |
| <i>Lipotriche</i> Less.      | 610      | <i>Myriactis</i> Less.      | 308      |
| <i>Locheria</i> Neck.        | 612      | <i>Myriadenus</i> Cass.     | 476      |
| <i>Lorentea</i> Less.        | 101, 102 |                             |          |
| <i>Lorentea</i> Lag.         | 98       | <i>Nardosmia</i> Cass.      | 205      |
| <i>Lorentea</i> Ort.         | 628      | <i>Narvalina</i> Cass.      | 633      |
| <i>Lychnocephalus</i> Mart.  | 83       | NASSAUUVIACEÆ.              | 9        |
| <i>Lychnophora</i> Mart.     | 78       | <i>Naumburgia</i> Willd.    | 635      |
| <i>Lychnophoroides</i> DC.   | ibid.    | <i>Nauplius</i> Cass.       | 486      |
|                              | ibid.    | <i>Needhamia</i> Cass.      | 633      |
| <i>Machæranthera</i> Nées.   | 261      | <i>Neja</i> Don.            | 325      |
| <i>Madaria</i> DC.           | 691      | <i>Neuractis</i> Cass.      | 632      |
| <i>Madarioides</i> DC.       | 692      | NEUROLAENEÆ.                | 500      |
| <i>Madaroglossa</i> DC.      | 694      | <i>Nidorella</i> Cass.      | 321      |
| <i>Madia</i> Sol.            | 301      | <i>Niebuhria</i> Scop.      | 509      |
| <i>Madia</i> Mol.            | 691      | <i>Nocca</i> Cass.          | 92       |
| <i>Madia</i> auct.           | 691-692  | <i>Noccea</i> DC.           | ibid.    |
| MADIEÆ DC.                   | 499, 691 | <i>Noccea</i> Jacq.         | 91       |
| <i>Mahometa</i> DC.          | 457      | <i>Nolletia</i> Cass.       | 366      |
| <i>Mairia</i> DC.            | 217      | <i>Noticastrum</i> DC.      | 279      |
| <i>Mairia</i> Nées.          | ibid.    | <i>Nothites</i> Cass.       | 186      |
| <i>Margarita</i> Gaud.       | 226      |                             |          |
| <i>Marshallia</i> Schreb.    | 680      | <i>Obeliscaria</i> Cass.    | 658      |
| <i>Matamoria</i> Lallav.     | 87       | <i>Obeliscotheca</i> Vaill. | 555      |
| <i>Meisteria</i> Scop.       | 14       | <i>Odontocarpa</i> DC.      | 71       |
| MELAMPODIEÆ Less.            | 498, 517 | <i>Odontoloma</i> H. B. K.  | 11       |
| MELAMPODINEÆ Less. DC.       | 497, 500 | <i>Odontospermum</i> Neck.  | 486, 488 |
| <i>Melampodium</i> Linn.     | 517      | <i>Ogiera</i> Cass.         | 545      |
| <i>Melanthera</i> Rich.      | 544      | <i>Ogiera</i> Spreng.       | 501      |
| <i>Melanthera</i> Rohr.      | ibid.    | <i>Oiospermum</i> Less.     | 11       |
| <i>Melanodendron</i> DC.     | 279      | <i>Olearia</i> Mœnch.       | 271      |
| <i>Mendezia</i> DC.          | 532      | <i>Oligactis</i> Cass.      | 95       |
| <i>Meratia</i> Cass.         | 502      | <i>Oligactis</i> DC.        | ibid.    |
| <i>Mesoretra</i> Raf.        | 666      | <i>Oligarion</i> Cass.      | 681      |
| <i>Meyera</i> Schreb.        | 636      | <i>Oliganthes</i> Cass.     | 15       |

- Oligogyne* DC. 629  
*Oligolepis* Cass. 379  
*Olecarpha* DC. 692  
*Oolinium* DC. 133  
*Oreina Bertol.* 104  
*Oreina Bertol.* 498  
*Oswalda* Cass. 505  
*Oswalda* DC. *ibid.*  
*Oxylobus* Moq. 115  
*Oxylobus* DC. *ibid.*  
*Oxyrappus* DC. 651  
*Oxyura* DC. 693  
*Oyedma* DC. 578  
  
*Pachyderis* Cass. 360  
*Pachyderis* DC. *ibid.*  
*Pachourina* Ait. 14  
*Pachourinopsis* Cass. *ibid.*  
*Palafoxia Lag.* 124  
*Palafoxia* Cass. *ibid.*  
*Pallaria* L'Hér. 588  
*Pollenia* Cass. 487  
*Poquerina* Cass. 307  
**PARTHENIEAE** Less. 498, 530  
*Partheniastrum* Niss. 531  
*Partheniastrum* DC. 531  
*Partheniata* DC. *ibid.*  
*Parthenium* Linn. 531  
*Pascalina* Ort. 549  
**PECTIDEAE** DC. 10, 94  
**PECTIDEAE** Less. 98  
*Pectidium* Less. *ibid.*  
*Pectidopsis* DC. *ibid.*  
*Pectis* Less. *ibid.*  
*Pectis* Linn. *ibid.*  
*Pectis* Cass. *ibid.*  
*Pectinellum* DC. 109  
*Pegoletia* Cass. 481  
*Pentameranthes* DC. 495  
*Pentanema* Cass. 474  
*Peramibus* Raf. 568  
*Perconia* Michx. 680  
*Perymenium* Schrad. 608  
*Petrobium* R. Br. 501  
**PETASITEAE** DC. 204  
*Petasites* Toura. 206  
*Phagnalon* Cass. 396  
*Phalacraea* DC. 105  
*Phalacroloma* Cass. 297  
*Phalacromesus* Cass. 457  
*Phania* DC. 114  
*Phialis* Spreng. 656  
*Philactis* Schrad. 534  
*Philoglossa* DC. 567  
*Phyllinema* Blum. 630  
*Phyllocephalum* Blum. 66  
*Phyllocephalum* DC. *ibid.*  
*Phylloneja* DC. 325  
*Phytumopsis* Juss. 680  
*Pieradenia* Hook. 665  
  
*Pingra* Cass. 398  
*Pinillosia* Ossa. 398  
*Piptocoma* Cass. 692  
*Piqueria* Cav. 133  
*Piquerioides* DC. 104  
*Pirarda* Adans. 498  
*Pithecocterus* Mart. 505  
*Platoclypta* DC. *ibid.*  
*Platycarpa* DC. 115  
*Platycarpa* Less. *ibid.*  
*Platypterus* H. B. K. 651  
*Platypterus* DC. 693  
*Pleionactis* DC. 578  
*Pluchea* Cass. 360  
**PLUCHINEAE** Cass. 379  
*Pluridens* Neck. 14  
*Podanthus* Lag. 104  
*Podocoma* Cass. *ibid.*  
*Podoneja* DC. 124  
*Pollalesta* H. B. K. *ibid.*  
*Polylopsis* Less. 588  
*Poloa* DC. 487  
*Polyactidium* D. C. 307  
*Polyactis* Less. 498, 530  
*Polyarrhena* Cass. 531  
*Polycephalus* Forst. 531  
*Polycephalus* DC. *ibid.*  
*Polytmia* Linn. 531  
*Polytmiastrum* Lam. 549  
*Polypappus* Less. 10, 94  
*Polypterus* Nutt. 98  
*Polypterus* Less. *ibid.*  
**POROPHYLLEAE** DC. *ibid.*  
*Porophyllum* Vaill. 499  
*Praxelis* Cass. 114  
*Priestleya* fl. mex. 56  
*Pronacron* Cass. 56  
*Proteopsis* Mart. 56  
*Proustia* Spreng. 56  
*Psathurocheta* DC. 56  
*Pseudo-elephantopus* Robr. 56  
*Pseudovax* DC. 56  
*Psidia* Jacq. 56  
**PSIADIEAE** DC. 312, 317  
*Psilocarpa* DC. 204  
*Pterigeron* DC. 206  
*Pterocaulon* Ell. 396  
*Pterocaulopsis* DC. 105  
*Pteronia* Linn. 297  
*Pteronia* Linn. f. 457  
*Pteropappus* Less. 114  
*Pteropappus* DC. 656  
*Pterophora* Neck. 534  
*Pterophorus* Cass. 567  
*Pterophorus* DC. 630  
*Pterophyton* Cass. 66  
*Ptiostephium* H. B. K. *ibid.*  
*Pycnocephalum* DC. 325  
*Pyrarda* Cass. 680  
*Pyrarda* DC. 665

# NOMINUM ET SYNONYMORUM.

|                    |          |                                |          |
|--------------------|----------|--------------------------------|----------|
|                    |          |                                | 705      |
| um Med.            | 620      | Skirroglyphis DC.              | 533      |
| ta DC.             | 349      | Sobreyra R. et P.              | 638      |
| ma Hook.           | 350      | Sobrya Pers.                   | ibid.    |
| ia DC.             | 477      | Sogalgina Cass.                | 678      |
| ia Gärtn.          | 478      | Sogaligna Steud.               | ibid.    |
|                    |          | Solenogyne Cass.               | 367      |
| TA Toura.          | 4        | SOLENOGYNEAE DC.               | 212, 365 |
| a DC.              | 551      | SOLIDAGINEAE DC.               | 212, 328 |
| ua Poepp. et Endl. | 664      | Solidago Linn.                 | 330      |
| a Raf.             | 558      | Sommea Bory.                   | 3        |
| DC.                | 559      | Sommerfeltia Less.             | 302      |
| NIEAE Less.        | 500      | Sparganophorus Vaill.          | 11       |
| rium Desf.         | 463      | Sparganophorus Michx.          | 114      |
| lina Less.         | 275      | Spermophylla Neck.             | 681      |
| ospermum Reinw.    | 296      | SPHAERANTHEAE DC.              | 212, 368 |
| Adans.             | 575      | Sphaeranthoides Cunn.          | 456      |
| ertia Cass.        | 503      | Sphaeranthus Linn.             | 369      |
| ia DC.             | 345      | Sphenogyne R. Br.              | 681      |
| ra Rothb.          | 90       | SPHENOGYNEAE DC.               | 499, 681 |
| ia Zenk.           | 646      | Spilanthes Jacq.               | 620      |
| ia.                | 183      | Spilanthes Cass.               | 624      |
| Less.              | 668      | Spilanthus Linn.               | 620      |
| Lam.               | 658      | Spilanthus Pers.               | 624      |
| ia Cass.           | 555      | Spiracantha H. B. K.           | 90       |
| ia Auct.           | 554, 558 | Spixia Schrank.                | 7        |
| CKIEAE Cass.       | 498, 554 | Spongotrichum Nées.            | 265      |
| dia DC.            | 549      | Spongotrichum DC.              | 269      |
|                    |          | Stachyanthus DC.               | 84       |
|                    |          | Stammarium Willd.              | 102      |
| Cass.              | 496      | Stammarium DC.                 | ibid.    |
| DC.                | 624      | Starkia Juss.                  | 96       |
| DC.                | 493      | Starkea Willd.                 | ibid.    |
| lia Gualt.         | 628      | Steiractis DC.                 | 345      |
| hemum Cass.        | 367      | Stemmodontia Cass.             | 538      |
| Neck.              | 288      | Stenactis Less.                | 298      |
| DC.                | 492      | Stenactis Cass.                | 281      |
| yne Cass.          | 475      | Stephananthus Lebm.            | 398      |
| endalia Willd.     | 638      | Stevia Cav.                    | 115      |
| arpha Less.        | 608      | Stilpnopappus Mart.            | 76       |
| ria Roth.          | 654      | Stobaea Spreng.                | 71       |
| ia Mönch.          | 496      | Stokesia L'Hér.                | ibid.    |
| irpus Jacq.        | 566      | Strabonia DC.                  | 481      |
| pis Cass.          | 114      | Strigia DC.                    | 126      |
| rmum Less.         | 509      | Strobocalyx Blume.             | 21       |
| mia Bertol.        | 630      | Strophopappus DC.              | 75       |
| I. B. K.           | 612      | Struchium P. Br.               | 11       |
| OSCULOSAE Toura.   | 4        | Stylimnus Raf.                 | 450      |
| ONEAE Less.        | 500      | Stylismaus DC.                 | ibid.    |
| ONIDEAE Less.      | 8, 497   | Suprago Gärtn.                 | 128, 129 |
| Gärtn.             | 398      | Suprago DC.                    | 129      |
| irpus Nées.        | 261      | SYNANTHERAE Rich.              | 4        |
| IEAE DC.           | 500      | SYNGENESIA POLYGAMIA Linn. ib. |          |
| Forst.             | 78       | Synchodendron Bojer.           | 92       |
| thus Fras.         | 347      | Synedrella Gärtn.              | 629      |
| eckia Linn.        | 495      |                                |          |
| EAE Less.          | 498, 510 |                                |          |
| m Linn.            | 511      | TAGETAE DC.                    | 499, 638 |
| cline DC.          | 21       | Tagetes Linn.                  | 642      |
| Pers.              | 577      | TAGETINEAE Cass.               | 499, 638 |
| IPHYTUM Neck.      | 4        | TARCHONANTHEAE Less.           | 213, 429 |

|                             |          |                                  |          |
|-----------------------------|----------|----------------------------------|----------|
| <i>Oligogyne</i> DC.        | 669      | <i>Pinguicula</i> Cass.          | 398, 400 |
| <i>Oligolapide</i> Cass.    | 379      | <i>Pinillosia</i> Ossa.          | 400      |
| <i>Oleocarpha</i> DC.       | 692      | <i>Piptodroma</i> Cass.          | 400      |
| <i>Ocellinium</i> DC.       | 133      | <i>Piqueria</i> Cav.             | 400      |
| <i>Ocinia</i> Bertol.       | 104      | <i>Piquerioides</i> DC.          | 400      |
| <i>Ocinia</i> Bertol.       | 478      | <i>Pirarda</i> Adams.            | 400      |
| <i>Oestrada</i> Cass.       | 505      | <i>Pithecosotis</i> Mart.        | 400      |
| <i>Ottalida</i> DC.         | ibid.    | <i>Platodypta</i> DC.            | 400      |
| <i>Oxylobus</i> Moq.        | 115      | <i>Platycarpa</i> DC.            | 400      |
| <i>Oxylobus</i> DC.         | ibid.    | <i>Platyacarpa</i> Less.         | 400      |
| <i>Oxyppus</i> DC.          | 651      | <i>Platypteria</i> H. B. K.      | 400      |
| <i>Oxyura</i> DC.           | 603      | <i>Platypteria</i> DC.           | 400      |
| <i>Oxyura</i> DC.           | 578      | <i>Platocactis</i> DC.           | 400      |
| <i>Pachyderis</i> Cass.     | 360      | <i>Pluchea</i> Cass.             | 400      |
| <i>Pachyderis</i> DC.       | ibid.    | <b>PLUCHINAE</b> Cass.           | 400      |
| <i>Pachyrhina</i> Aubl.     | 14       | <i>Pharidens</i> Neck.           | 400      |
| <i>Pachyrhina</i> Niss.     | ibid.    | <i>Podanthus</i> Lag.            | 400      |
| <i>Palafoxia</i> Lag.       | 114      | <i>Podocoma</i> Cass.            | 400      |
| <i>Palafoxia</i> Cass.      | ibid.    | <i>Pedoneja</i> DC.              | 400      |
| <i>Palafoxia</i> L'Hér.     | 538      | <i>Pellalea</i> H. B. K.         | 400      |
| <i>Palafoxia</i> Cass.      | 48       | <i>Polypteria</i> Less.          | 400      |
| <i>Palafoxia</i> Cass.      | 307      | <i>Pelosa</i> DC.                | 400      |
| <b>PALUDINAE</b> Less.      | 498, 530 | <i>Polyactidium</i> D. C.        | 400      |
| <i>Palafoxia</i> Cass.      | 531      | <i>Polyactis</i> Less.           | 400      |
| <i>Palafoxia</i> Cass.      | 530      | <i>Polyarrhena</i> Cass.         | 400      |
| <i>Palafoxia</i> Cass.      | ibid.    | <i>Polycephalus</i> Forst.       | 400      |
| <i>Palafoxia</i> Linn.      | 531      | <i>Polycephalus</i> DC.          | 400      |
| <i>Palafoxia</i> Ort.       | 549      | <i>Polymania</i> Linn.           | 400      |
| <b>PECTIDAE</b> DC.         | 10, 94   | <i>Polymaniastrum</i> Lam.       | 514, 543 |
| <b>PECTIDAE</b> Less.       | 98       | <i>Polyppus</i> Less.            | 400      |
| <i>Pectidium</i> Less.      | ibid.    | <i>Polypteria</i> Nutt.          | 400      |
| <i>Pectidopsis</i> DC.      | ibid.    | <i>Polypteria</i> Less.          | 400      |
| <i>Pectis</i> Less.         | ibid.    | <b>POROPHYLLAE</b> DC.           | 499, 547 |
| <i>Pectis</i> Linn.         | ibid.    | <i>Porophyllum</i> Vail.         | 400      |
| <i>Pectis</i> Cass.         | ibid.    | <i>Preziosi</i> Cass.            | 133      |
| <i>Pectinellum</i> DC.      | 109      | <i>Priestleya</i> fl. mex.       | 54       |
| <i>Pegolettia</i> Cass.     | 481      | <i>Pronacron</i> Cass.           | 54       |
| <i>Pentameranthes</i> DC.   | 495      | <i>Protopsis</i> Mart.           | 5        |
| <i>Pentanema</i> Cass.      | 474      | <i>Prostia</i> Spreng.           | 5        |
| <i>Paramibus</i> Raf.       | 568      | <i>Psathurocheta</i> DC.         | 609      |
| <i>Parsonia</i> Michx.      | 680      | <i>Pseudo-elephantopus</i> Robr. | 5        |
| <i>Parymenium</i> Schrad.   | 608      | <i>Pseudevax</i> DC.             | 400      |
| <i>Petrobium</i> R. Br.     | 501      | <i>Psadia</i> Jacq.              | 518      |
| <b>PETASITEAE</b> DC.       | 204      | <b>PSIADIEAE</b> DC.             | 512, 517 |
| <i>Petasites</i> Tourn.     | 206      | <i>Psilocarpa</i> DC.            | 517      |
| <i>Phagnalon</i> Cass.      | 396      | <i>Pterigeron</i> DC.            | 517      |
| <i>Phalacraea</i> DC.       | 105      | <i>Pterocaulon</i> Ell.          | 43       |
| <i>Phalacroloma</i> Cass.   | 207      | <i>Pterocaulopsis</i> DC.        | 43       |
| <i>Phalacromes</i> Cass.    | 457      | <i>Pteronia</i> Linn.            | 43       |
| <i>Phania</i> DC.           | 114      | <i>Pteronia</i> Linn. f.         | 43       |
| <i>Phialis</i> Spreng.      | 656      | <i>Pteropappus</i> Less.         | 43       |
| <i>Phyllactis</i> Schrad.   | 534      | <i>Pteropappus</i> DC.           | 43       |
| <i>Philoglossa</i> DC.      | 567      | <i>Pterophora</i> Neck.          | 364      |
| <i>Phyllinema</i> Blum.     | 630      | <i>Pterophorus</i> Cass.         | 43       |
| <i>Phyllocephalum</i> Blum. | 66       | <i>Pterophorus</i> DC.           | 43       |
| <i>Phyllocephalum</i> DC.   | ibid.    | <i>Pterophyton</i> Cass.         | 5-5      |
| <i>Phylloneja</i> DC.       | 325      | <i>Ptilistephium</i> H. B. K.    | 6-8      |
| <i>Phytumopsis</i> Juss.    | 680      | <i>Pycnocephalum</i> DC.         | 43       |
| <i>Pieradenia</i> Hook.     | 665      | <i>Pyrarda</i> Cass.             | 3-3      |
|                             |          | <i>Pyrarda</i> DC.               | 3-3      |



NOMINUM ET SYNONYMORUM.

705

|                            |          |                                   |              |
|----------------------------|----------|-----------------------------------|--------------|
| <i>rum</i> Med.            | 620      | <i>Skirroglyphis</i> DC.          | 533          |
| <i>eta</i> DC.             | 349      | <i>Sobreyra</i> R. et P.          | 638          |
| <i>coma</i> Hook.          | 350      | <i>Sobrya</i> Pers.               | <i>ibid.</i> |
| <i>ria</i> DC.             | 477      | <i>Sogalgina</i> Cass.            | 678          |
| <i>ria</i> Gärtn.          | 478      | <i>Sogaligna</i> Steud.           | <i>ibid.</i> |
|                            |          | <i>Solenogyne</i> Cass.           | 367          |
| <i>ATÆ</i> Tourn.          | 4        | <i>SOLENOGYNEAE</i> DC.           | 212, 365     |
| <i>lla</i> DC.             | 551      | <i>SOLIDAGINEAE</i> DC.           | 212, 328     |
| <i>gua</i> Poepp. et Endl. | 664      | <i>Solidago</i> Linn.             | 330          |
| <i>da</i> Raf.             | 558      | <i>Sommea</i> Bory.               | 3            |
| <i>la</i> DC.              | 559      | <i>Sommerfeltia</i> Less.         | 302          |
| <i>ANIEAE</i> Less.        | 500      | <i>Sparganophorus</i> Vaill.      | 11           |
| <i>erium</i> Desf.         | 463      | <i>Sparganophorus</i> Michx.      | 114          |
| <i>ctina</i> Less.         | 275      | <i>Spermophylla</i> Neck.         | 681          |
| <i>hospermum</i> Reinw.    | 206      | <i>SPHAERANTHEAE</i> DC.          | 212, 368     |
| <i>Adans.</i>              | 575      | <i>Sphaeranthoides</i> Cunn.      | 456          |
| <i>urtia</i> Cass.         | 503      | <i>Sphaeranthus</i> Linn.         | 369          |
| <i>nia</i> DC.             | 345      | <i>Sphenogyne</i> R. Br.          | 681          |
| <i>lra</i> Rottb.          | 90       | <i>SPHENOGYNEAE</i> DC.           | 499, 681     |
| <i>kia</i> Zenk.           | 696      | <i>Spilanthes</i> Jacq.           | 620          |
| <i>sia.</i>                | 183      | <i>Spilanthes</i> Cass.           | 624          |
| <i>i</i> Less.             | 668      | <i>Spilanthus</i> Linn.           | 620          |
| <i>i</i> Lam.              | 658      | <i>Spilanthus</i> Pers.           | 624          |
| <i>ckia</i> Cass.          | 555      | <i>Spiracantha</i> H. B. K.       | 90           |
| <i>ckia</i> Auct.          | 554, 558 | <i>Spixia</i> Schrank.            | 7.           |
| <i>ICKIEAE</i> Cass.       | 498, 554 | <i>Spongotrichum</i> Nées.        | 265          |
| <i>rdia</i> DC.            | 549      | <i>Spongotrichum</i> DC.          | 269          |
|                            |          | <i>Stachyanthus</i> DC.           | 84           |
| <i>a</i> Cass.             | 496      | <i>Stammarium</i> Willd.          | 102          |
| <i>a</i> DC.               | 624      | <i>Stammarium</i> DC.             | <i>ibid.</i> |
| <i>i</i> DC.               | 493      | <i>Starkia</i> Juss.              | 96           |
| <i>alia</i> Gualt.         | 628      | <i>Starkea</i> Willd.             | <i>ibid.</i> |
| <i>themum</i> Cass.        | 367      | <i>Steiractis</i> DC.             | 345          |
| <i>ia</i> Neck.            | 366      | <i>Stemmodontia</i> Cass.         | 538          |
| <i>a</i> DC.               | 262      | <i>Stenaotis</i> Less.            | 298          |
| <i>gyne</i> Cass.          | 479      | <i>Stenaotis</i> Cass.            | 281          |
| <i>tendalia</i> Willd.     | 638      | <i>Stephananthus</i> Lebm.        | 398          |
| <i>carpha</i> Less.        | 608      | <i>Stevia</i> Cav.                | 115          |
| <i>bria</i> Roth.          | 654      | <i>Stilpnopappus</i> Mart.        | 76           |
| <i>iria</i> Moench.        | 496      | <i>Stobæa</i> Spreng.             | 71           |
| <i>carpus</i> Jacq.        | 566      | <i>Stokesia</i> L'Hér.            | <i>ibid.</i> |
| <i>lepis</i> Cass.         | 114      | <i>Strabonia</i> DC.              | 481          |
| <i>permum</i> Less.        | 509      | <i>Strigia</i> DC.                | 126          |
| <i>iania</i> Bertol.       | 630      | <i>Strobocalyx</i> Blume.         | 21           |
| <i>H. B. K.</i>            | 612      | <i>Strophopappus</i> DC.          | 75           |
| <i>LOSCULOSÆ</i> Tourn.    | 4        | <i>Struchium</i> P. Br.           | 11           |
| <i>IONEÆ</i> Less.         | 500      | <i>Stylimnus</i> Raf.             | 450          |
| <i>IONIDEÆ</i> Less.       | 8, 497   | <i>Stylimus</i> DC.               | <i>ibid.</i> |
| <i>is</i> Gärtn.           | 394      | <i>Suprago</i> Gärtn.             | 128, 129     |
| <i>carpus</i> Nées.        | 261      | <i>Suprago</i> DC.                | 129          |
| <i>IEAE</i> DC.            | 500      | <i>SYNANTHERÆ</i> Rich.           | 4            |
| <i>i</i> Forst.            | 78       | <i>SYNGENESIA POLYGAMIA</i> Linn. | <i>ib.</i>   |
| <i>nthus</i> Fras.         | 347      | <i>Synchodendron</i> Bojer.       | 92           |
| <i>beckia</i> Linn.        | 495      | <i>Synedrella</i> Gärtn.          | 629          |
| <i>IEAE</i> Less.          | 498, 510 |                                   |              |
| <i>um</i> Linn.            | 511      | <i>TAGETAE</i> DC.                | 499, 638     |
| <i>ocline</i> DC.          | 21       | <i>Tagetes</i> Linn.              | 642          |
| <i>Pers.</i>               | 577      | <i>TAGETINEAE</i> Cass.           | 499, 638     |
| <i>NIPHYTUM</i> Neck.      | 1        | <i>TARCHONANTHEAE</i> Less.       | 213, 129     |

|                               |          |                             |          |
|-------------------------------|----------|-----------------------------|----------|
| <i>Tarchonanthus</i> Linn.    | 431      | <i>Urolepis</i> DC.         | 131      |
| <i>Tecmarsia</i> DC.          | 93       | <i>Ursinia</i> Gaertn.      | 681      |
| <i>Telekia</i> Baumg.         | 485      | <i>Ursinia</i> Auct.        | 681, 689 |
| <i>Tetramolopium</i> Nees.    | 262      |                             |          |
| <i>Tephrodes</i> DC.          | 24       | <i>Vanillosma</i> Less.     | 18       |
| <i>Tessaria</i> R. et P.      | 456      | <i>Vargasia</i> DC.         | 676      |
| <i>Tetracarpum</i> Moench.    | 654      | <i>Varthemia</i> DC.        | 471      |
| <i>Tetragonotheca</i> Dill.   | 552      | <i>Verbesina</i> Less.      | 612      |
| <i>Tetrantha</i> Poit.        | 503      | <i>Verbesinaria</i> DC.     | 612      |
| <i>Tetranthus</i> Swartz.     | 528      | VERBESINEAE Less.           | 498, 607 |
| <i>Tetradus</i> Cass.         | 666      | <i>Vermifuga</i> R. et Pav. |          |
| <i>Tetrodus</i> DC.           | ibid.    | <i>Vernonia</i> Schreb.     | 15       |
| <i>Tetraotis</i> Blum.        | 637      | <i>Vernonia</i> Less.       | 14, 74   |
| <i>Thelesperma</i> Less.      | 633      | VERNONIACEAE Less.          | 8, 9     |
| <i>Thelythamnos</i> Less.     | 683      | VERNONIEAE Cass.            | 9        |
| <i>Thelythamnos</i> DC.       | 681      | VERNONIEAE DC.              | 10       |
| <i>Therogeron</i> DC.         | 283      | <i>Vicoa</i> Cass.          | 471      |
| <i>Thespis</i> DC.            | 375      | <i>Viguiera</i> H. B. K.    | 579      |
| <i>Thouarsia</i> Vent.        | 318      | <i>Viguiera</i> Less.       | ibid.    |
| <i>Thymophylla</i> Lag.       | 647      | <i>Villanova</i> Ort.       | 511      |
| <i>Tilesia</i> Mey.           | 549      | <i>Virga-aurea</i> Tourn.   | 530      |
| <i>Tithonia</i> Desf.         | 584      | <i>Virgaurea</i> DC.        | ibid.    |
| <i>Tragantes</i> Wallr.       | 176      | <i>Virgilia</i> L'Hér.      | 651      |
| <i>Tragoceras</i> Less.       | 533      | <i>Vittadinia</i> Rich.     | 140      |
| <i>Partragoceros</i> H. B. K. | ibid.    | <i>Viviania</i> Willd.      | 65       |
| <i>rattenickia</i> Pers.      | 680      | <i>Vosacan</i> Adans.       | 585      |
| <i>Trianthema</i> DC.         | 23       |                             |          |
| <i>Trichogonia</i> DC.        | 126      | <i>Wahlenbergia</i> Schum.  | 407      |
| <i>Trichophyllum</i> Nutt.    | 656      | <i>Webbia</i> DC.           | 77       |
| <i>Trichospermum</i> Beauv.   | 531      | <i>Wedelia</i> Jacq.        | 538      |
| <i>Trichospira</i> H. B. K.   | 90       | <i>Wedelia</i> Auct.        | 515, 535 |
| <i>Trichostemma</i> Cass.     | 538      | <i>Wiborgia</i> Roth.       | 677      |
| <i>Trichostephium</i> Cass.   | 540      | <i>Wickstræmia</i> Spreng.  | 140      |
| <i>Tridax</i> Linn.           | 679      | <i>Willdenowa</i> Cav.      | 658      |
| <i>Trigonospermum</i> Less.   | 508      | <i>Wollastonia</i> DC.      | 540      |
| <i>Trilisa</i> Cass.          | 128, 131 | <i>Woodvillea</i> DC.       | 148      |
| <i>Trilisa</i> DC.            | 131      | <i>Wulfia</i> Neck.         | 501      |
| <i>Trimeranthes</i> Cass.     | 496      |                             |          |
| <i>Trimeranthes</i> DC.       | ibid.    | <i>Xanthioides</i> DC.      | 581      |
| <i>Trimorphæa</i> Cass.       | 290      | <i>Xanthisma</i> DC.        | 94       |
| <i>Trimorphæa</i> DC.         | ibid.    | <i>Xanthium</i> Linn.       | 520      |
| <i>Tripolium</i> Nees.        | 253      | <i>Xanthocoma</i> H. B. K.  | 310      |
| <i>Trixidium</i> DC.          | 606      | <i>Xantholepis</i> Willd.   | 77       |
| <i>Trixis</i> Swartz.         | 505      | <i>Xenismia</i> DC.         | 509      |
| <i>Tropidolepis</i> Tausch.   | 216      | <i>Xerolepis</i> Less.      | 607      |
| <i>Tabifera</i> DC.           | 386      | <i>Xerolepis</i> DC.        | ibid.    |
| <i>Tubilium</i> Cass.         | 480      | <i>Ximenesia</i> Cav.       | 677      |
| <i>Tubilium</i> DC.           | 480      | <i>Ximenesia</i> Less.      | ibid.    |
| <i>Turczaninovia</i> DC.      | 257      | <i>Xerothermus</i> DC.      | 311      |
| <i>Turpinia</i> Lallav.       | 77       | <i>Zaluzania</i> Pers.      | 551      |
| <i>Tursenia</i> Cass.         | 398      | <i>Zarabellia</i> Cass.     | 509      |
| TUSSILAGINEAE Less.           | 103, 204 | <i>Zarabellia</i> DC.       | 204      |
| <i>Tussilago</i> Tourn. Cass. | 208      | <i>Zexmenia</i> Lallav.     | 600      |
| <i>Tussilago</i> Linn.        | 204, 208 | <i>Zeyhera</i> Spreng.      | 401      |
|                               |          | <i>Zinnia</i> Linn.         | 524      |
| <i>Ueacea</i> Cass.           | 492      | <i>Zyrphelis</i> Nees.      | 217, 218 |
| <i>Ucucu</i> Adans.           | 629      | <i>Zyrphelis</i> DC.        | 218      |
| <i>Unxia</i> Linn. f.         | 507      |                             |          |

